Number of kinesins engaged in axonal cargo transport: A novel biomarker for neurological disorders

Kumiko Hayashi[1]* and Kazuo Sasaki[2]

[1]Institute for Solid State Physics, The University of Tokyo, Kashiwa, Japan
[2]Department of Applied Physics, Graduate School of Engineering, Tohoku University, Sendai, Japan
*Correspondence to Kumiko Hayashi
hayashi@issp.u-tokyo.ac.jp

ABSTRACT

The motor protein kinesin plays a crucial role in anterograde transport of cargo vesicles in neurons, moving them from the cell body towards the synaptic region. Not only the transport force and velocity, but also the number of kinesin molecules involved in transporting a single cargo vesicle, is pivotal for synapse formation. This collective transport by multiple kinesins ensures stable cargo transport in neurons. Abnormal increases or decreases in the number of engaged kinesin molecules could potentially act as biomarkers for neurodegenerative diseases such as Alzheimer's, Parkinson's, amyotrophic lateral sclerosis (ALS), spastic paraplegia, polydactyly syndrome, and virus transport disorders. Our aim, by employing novel physical measurements to quantify the number of kinesin molecules, is to shed light on the molecular mechanisms of neurodegenerative diseases related to axonal transport.

Introduction

In axonal transport, the motor protein kinesin transports synaptic materials synthesized in the cell body through long axons to the synaptic regions at the terminal. Kinesin is a motor protein that moves in the plus-end direction of microtubules by obtaining energy through ATP hydrolysis (Hirokawa et al., 2009; Vale, 2003). It is known that there are 45 types of kinesin superfamily members in humans. Among them, KIF1A is a kinesin that transports synaptic vesicle precursors, and it is known that mutations in KIF1A can cause excessive transport of synaptic vesicle precursors and transport disorders, leading to abnormalities in synapse formation (Anazawa et al., 2022; Chiba et al., 2019; Niwa et al., 2016). As a result, the mutation in kinesin can lead to severe neurodegenerative diseases (Gabrych et al., 2019; Guedes-Dias and Holzbaur, 2019). While the genetic disease caused by KIF1A is a rare disease (Iqbal et al., 2017; Klebe et al., 2012), a significant number of people are suffering from this newly discovered disease, and a global community to support those with KIF1A-related neurodevelopmental disorders, KIF1A.org, was established in 2016.

Axonal transport mutations caused by KIF1A have been linked to physical deficits, such as a decrease in transport velocity, transport force, or the attachment rate to microtubules (Anazawa et al., 2022; Budaitis et al., 2021; Chiba et al., 2019; Lam et al., 2021). In addition to these phenomena, we have discovered that changes in the number of kinesin molecules transporting cargo can disrupt synapse formation (Hayashi et al., 2018a). It is thought that axonal transport is efficient if the cargo (synaptic vesicle precursors) is transported by the appropriate number of KIF1A molecules, and axonal transport is inefficient if the number of KIF1A molecules decreases or becomes excessive (Hayashi et al., 2018a).

We will discuss a novel measurement method for the number of kinesins, engaging in the transport of synaptic vesicle precursors in the motor neurons of *Caenorhabditis elegans* (*C. elegans*) as an example. Figure 1 shows the results of observing the transport of GFP-labeled synaptic vesicle precursors in the axon using fluorescence microscopy. We can observe the cargo vesicles transported by kinesins, moving in one direction while fluctuating (tracked by particle tracking software). The causes of this fluctuating behavior are thermal fluctuations and collisions with surrounding vesicles and the cytoskeleton.

The details of the physical background of the measurement method are left to the biophysical references (Hasegawa et al., 2019; Hayashi et al., 2018a; Hayashi et al., 2021; Hayashi et al., 2018b), we have successfully identified the parameter χ, a potential biomarker indicative of the quantity of kinesin molecules. Measuring the parameter χ, we found that 1-4 kinesin molecules co-transport the synaptic vesicle precursors, but when there is a deficit in axonal transport, in which binding of kinesin and the cargo vesicle is weak, the number of molecules decreases (Hayashi et al., 2018a).

Simple theoretical model of cargo transport to derive χ

Let's denote the displacement of a cargo vesicle during the time interval Δt as X, and the number of kinesins transporting a single cargo vesicle as N (Fig. 2A). x_i is the displacement of the i-th kinesin ($i = 1, 2, \cdots, N$). Then, the probability distribution function of the displacement X of the cargo vesicle can be written as:

$$P_N(X) = \int_{-\infty}^{+\infty} P(y) \prod_{i=1}^{N} P(x_i) \delta\left(X - \frac{1}{N}\sum_{i=1}^{N} x_i - y\right) \prod_{i=1}^{N} dx_i \, dy$$

By the definition of the delta function

$$\delta(x) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{ikx} dk$$

we have:

$$P_N(X) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{ikX} e^{-iky} P(y) \left(\int_{-\infty}^{+\infty} P(x_1) e^{-ikx_1/N} dx_1 \right)^N dk dy$$

Here, we assume that the cargo vesicle surface (membrane) is flexible and each kinesin moves independently (Fig. 2A). $P(x_i)$ is written as

$$P(x_i) = \frac{1}{\sqrt{2\pi}s} e^{-\frac{1}{2s^2}(x_i-\mu)^2}$$

Here $\mu (= \bar{v}\Delta t)$ and $s^2 (= \sigma^2 \Delta t)$ are the displacement and variance of each kinesin, respectively. $P(y)$ represents the effect of thermal noise acting on the cargo vesicle (Fig. 2A),

$$P(y) = \sqrt{\frac{\beta\kappa}{2\pi}} e^{-\frac{\beta\kappa}{2}y^2}$$

where κ is an effective stiffness between the cargo and kinesin molecules, and $\beta = k_B T$ (k_B:Boltzmann constant, T:Temperature of the environment (temperature in a cell)). Substituting the two equations into $P_N(X)$, we obtain

$$P_N(X) = \frac{1}{2} \left\{ \pi \left(\frac{s^2}{2N} + \frac{1}{2\beta\kappa} \right) \right\}^{-1/2} e^{-\frac{1}{2\left(\frac{s^2}{N} + \frac{1}{\beta\kappa}\right)}(X-\mu)^2}$$

When considering a sufficiently large Δt ($\Delta t \to \infty$) and since $s^2 \propto \Delta t$, $s^2/N \gg 1/\beta\kappa$ (= const), we have:

$$P_N(X) \sim \sqrt{\frac{N}{2\pi s^2}} e^{-\frac{1}{2s^2/N}(X-\mu)^2}$$

At this point, we focus on the mean ($\mu = \bar{v}\Delta t$) and variance ($2D\Delta t = s^2/N$) of the cargo vesicle displacement X, which are measurable quantities ($\bar{v}$ and D) in the experiment, and construct the following quantity:

$$\chi = \frac{\bar{v}}{D} = \frac{\mu}{s^2/N}$$

In Fig. 2B, we provide the simplest force (load)-velocity (F-v) relationship model of kinesin. It should be noted that there exist more sophisticated models of the force-velocity relationship, such as the energy landscape model (Schnitzer et al., 2000) and the three-state model (Sasaki et al., 2018), both of which are derived based on the ATP hydrolysis mechanism of kinesin. The F-v relationship was measured in the *in-vitro* single-molecule experiments (Nishiyama et al., 2002; Schnitzer et al., 2000).

When the viscosity coefficient of the vesicle is denoted by Γ, the average velocity of the cargo vesicle $\bar{v}$ can be written as the intersection between the force-velocity (F-v) relationship and the line representing the Stokes law ($v = F/\Gamma$) as follows:

$$\bar{v} = \frac{V_{\max}}{1 + \frac{\Gamma V_{\max}}{N F_s}}$$

where $V_{\max}$ and F_s is the maximum velocity (the velocity without a cargo vesicle) and the maximum force (the stall force that kinesin can generate against a load F). F_s is known to be of the order of 10 pN from the single-molecule experiments using optical tweezers (Nishiyama et al., 2002; Schnitzer et al., 2000), and $V_{\max}$ is reported to be 4 μm/s as a result of the extreme-value analysis (Naoi et al., 2021). From the result of fluid mechanics, $\Gamma = 6\pi\eta r$ (η:viscosity of the environment, r:radius of the cargo). Then, the expression of χ including the vesicle's viscosity coefficient Γ is:

$$\chi = \frac{V_{\text{max}}}{1 + \frac{\Gamma V_{\text{max}}}{NF_{\text{s}}}} \cdot \frac{N}{\sigma^2}$$

Note that $\mu = \bar{v}\Delta t$ and $s^2 = \sigma^2 \Delta t$. The N dependence of χ is explicitly written as

$$\chi = \frac{c_1 N^2}{N + c_2}$$

where $c_1 = V_{\text{max}}/\sigma^2$ and $c_2 = \Gamma V_{\text{max}}/F_{\text{s}}$. Thus, the number of kinesin molecules N can be related to χ, a measurable quantity in the experiment.

Under the high viscosity inside the cell ($V_{\text{max}} \gg \bar{v}$), where we have:

$$V_{\text{max}} \gg \bar{v} = \frac{F}{\Gamma} \sim \frac{NF_{\text{s}}}{\Gamma}$$

$$\chi \sim \frac{F_{\text{s}}}{\Gamma} \cdot \frac{N^2}{\sigma^2} \propto N^2 \qquad (1)$$

As to the variance σ^2 of a kinesin, the fluorescence observation of cargos in neurons (Hayashi et al., 2021) in Fig. 2C suggests $\sigma^2 \propto 1/\sqrt{\text{FI}}$, where FI is the fluorescence intensity of a cargo vesicle, i.e., $\sqrt{\text{FI}}$ represents the cargo radius when the fluorescence materials are distributed on the cargo vesicle surface uniformly. It indicates $\sigma^2 \propto 1/\Gamma$ because $\Gamma = 6\pi\eta r$. Combine the experimental suggestion with Eq. (1), the following relation is predicted,

$$\chi \sim cF_{\text{s}}N^2 \qquad (2)$$

Here c is the constant which has a unit of energy such as $k_{\text{B}}T$. It is a future research to derive Eq. (2) theoretically.

Experimental results

We show the experimental results to measure χ (Eq. (1)). Fluorescence images of green fluorescence protein (GFP)-labeled synaptic cargo (synaptic vesicle precursors) transported by kinesin, UNC104/KIF1A, in the axon of *C. elegans* were captured using a camera at 100 frames per second as described in Fig. 1B (refer to the reference (Hayashi et al., 2018a) for the detailed experimental procedures). The body movements of *C. elegans* worms were suppressed under anesthesia (Niwa, 2017).

We consider a theoretical model for the motion of a cargo vesicle, transported by kinesins in constant velocity segments (CVSs) (Fig. 1B), which is assumed to be unaffected by other motor proteins based on previous experimental studies (D'Souza et al., 2022; Gennerich and Schild, 2006). This implies that the tug-of-war model, depicting a conflict between the two motors, kinesin and dynein (Gennerich and Schild, 2006; Gross, 2004), was not considered for CVSs. Instead, we support the 'motor coordination model,' in which adaptor proteins that connect motors with cargos deactivate opposing motors (Chen et al., 2019).

For one such segment, we computed χ as a function of Δt, as illustrated in one example. We observed that χ converges to a constant value as Δt becomes large (Fig. 3A). Subsequently, we calculated the χ values for 40 different cargo vesicles and observed the discrete behavior of χ (Fig. 3B, left). We clustered χ based on its discreteness, and then plotted χ against the cluster number N. We subsequently examined the dependency of χ on N, as predicted by Eq. (1) (Fig. 3C).

Kinesins that are not engaged in vesicle transport exist in an autoinhibitory (inactive) state to prevent ATP

wastage. Previous studies revealed that the protein ARL-8 promotes the release of kinesin from this autoinhibitory state (Niwa et al., 2016). In *arl-8* deleted *C. elegans* mutants, the release of kinesin autoinhibition does not proceed adequately, leading to a decrease in transport stability due to a shortage of active kinesin molecules. This causes abnormalities in synapse formation such as mis-locations of synapses (Fig. 3D) (Niwa et al., 2016). We performed the measurement of χ using the same *arl-8* deleted *C. elegans* mutant (Hayashi et al., 2018a). The results are shown in Fig. 3B (right). Consequently, we found a decrease in the number of χ clusters, indicating that χ could detect the reduction in the number of active kinesin molecules.

We also measured χ in mouse neurons (Hayashi et al., 2021; Hayashi et al., 2018b). In the hippocampal neurons of mice, 1-6 kinesin molecules were identified as involved in the transport of synaptic cargos (Fig. 4). When it comes to the transport of endosomes in the superior cervical ganglion (SCG) neurons of mice, we found 1-4 kinesin molecules to be involved.

Regarding dynein, another motor protein, dynein, involved in retrograde transport (Guedes-Dias and Holzbaur, 2019), previous studies have reported on the measurement results of χ (Hasegawa et al., 2019; Hayashi et al., 2021; Peng et al., 2022). The different structures and F-v relationships have been reported for dynein (Brenner et al., 2020; Elshenawy et al., 2019; Gennerich et al., 2007; Hirakawa et al., 2000; Mallik et al., 2004; Toba et al., 2006). Retrograde transport by dynein is also known to be related to neurodegenerative diseases (Guedes-Dias and Holzbaur, 2019; Schmieg et al., 2014; Terenzio et al., 2017).

Conclusion

Similar to a multitude of ants collectively transporting food, the process of axonal transport mediated by kinesin also necessitates the coordinated action of multiple molecules. We introduce a novel physical measurement, termed χ, to provide insights into the mechanisms underlying KIF1A-associated neurological disorders, with particular emphasis on the role of the number of kinesin molecules. We suggest that the physical properties of axonal transport are governed not only by transport force and velocity, but also critically by the number of participating kinesin molecules. Prior research indeed suggests that variations in kinesin count can precipitate abnormalities in synapse formation (Chiba et al., 2019; Niwa et al., 2016). Hence, a comprehensive examination of the linkage between axonal transport aberrations and synaptic formation anomalies represents a crucial focus for future investigations. The potential implication of transport disorders extends to a spectrum of neurodegenerative diseases, including but not limited to Alzheimer's disease, Parkinson's disease, ALS, spastic paraplegia, Short Rib-Polydactyly Syndrome, and viral transport-related disorders. Our intention is to make a meaningful contribution to the demystification of molecular mechanisms underpinning neurological diseases associated with axonal transport, facilitated by our novel approach in physical measurements.

Acknowledgments

This research was conducted with the support of AMED under Grant Number JP18gm5810009, JST under Grant Number JPMJPR1877, the Nakatani Medical and Measurement Technology Promotion Foundation Development Research Grant, the KIF1A.org Mini Grant, and the Grant for Basic Science Research Projects of the Sumitomo Foundation.

Fig. 1. A: Schematics of cargo vesicle transport by kinesin. If the number of kinesins is appropriate, neuronal axonal transport remains stable. However, if the number decreases, it can lead to neurodegenerative diseases. **B:** Fluorescence observation of a cargo vesicle, as reported by (Hayashi et al., 2018a)(left). The image tracks the position of the cargo over time. The inset represents X and Δt, while the highlighted yellow segment represents a constant velocity segment.

Fig. 2. A: Theoretical model of cargo vesicle transport by kinesin. **B:** Simple force-velocity relationship of kinesin. **C:** $D(=s^2/2N\Delta t)$ versus the fluorescence intensity (FI) of cargo vesicles observed in the mouse hippocampal neurons, originally obtained in the supporting material of the reference (Hayashi et al., 2018a).

Fig. 3. Measurement of χ for synaptic vesicle precursor transport by UNC104/KIF1A in *C. elegans* motor neurons (Hayashi et al., 2018a). **A:** Measurement of χ for a cargo vesicle as an example. The thin curves represent χ calculated based on a boot strapping method for the error estimation (Hayashi et al., 2018a). **B:** Measurement of χ for 40 cargo vesicles in the case of the wild type *C. elegans* (left) and the *arl-8* deleted mutant *C. elegans* (right). We found four clusters for the wild type and three clusters for the mutant. **C:** The convergent value, χ^*, of χ as a function of the cluster number N. **D:** Schematic of *en passant* synapses in the motor neuron of *C. elegans*.

Fig. 4. Measurement of χ for synaptic vesicle precursor transport by KIF1A in mouse hippocampal neurons. **A:** Measurement of χ for 131 cargo vesicles. We found six clusters. **B:** The convergent value, χ^*, of χ as a function of the cluster number N.

Reference

Anazawa, Y., Kita, T., Iguchi, R., Hayashi, K., Niwa, S., 2022. De novo mutations in KIF1A-associated neuronal disorder (KAND) dominant-negatively inhibit motor activity and axonal transport of synaptic vesicle precursors. Proc Natl Acad Sci U S A 119, e2113795119.

Brenner, S., Berger, F., Rao, L., Nicholas, M.P., Gennerich, A., 2020. Force production of human cytoplasmic dynein is limited by its processivity. Sci Adv 6, eaaz4295.

Budaitis, B.G., Jariwala, S., Rao, L., Yue, Y., Sept, D., Verhey, K.J., Gennerich, A., 2021. Pathogenic mutations in the kinesin-3 motor KIF1A diminish force generation and movement through allosteric mechanisms. J Cell Biol 220.

Chen, C.W., Peng, Y.F., Yen, Y.C., Bhan, P., Muthaiyan Shanmugam, M., Klopfenstein, D.R., Wagner, O.I., 2019. Insights on UNC-104-dynein/dynactin interactions and their implications on axonal transport in Caenorhabditis elegans. J Neurosci Res 97, 185-201.

Chiba, K., Takahashi, H., Chen, M., Obinata, H., Arai, S., Hashimoto, K., Oda, T., McKenney, R.J., Niwa, S., 2019. Disease-associated mutations hyperactivate KIF1A motility and anterograde axonal transport of synaptic vesicle precursors. Proc Natl Acad Sci U S A 116, 18429-18434.

D'Souza, A.I., Grover, R., Monzon, G.A., Santen, L., Diez, S., 2022. Vesicles driven by dynein and kinesin exhibit directional reversals without external regulators. bioRxiv.

Elshenawy, M.M., Canty, J.T., Oster, L., Ferro, L.S., Zhou, Z., Blanchard, S.C., Yildiz, A., 2019. Cargo adaptors regulate stepping and force generation of mammalian dynein-dynactin. Nat Chem Biol 15, 1093-1101.

Gabrych, D.R., Lau, V.Z., Niwa, S., Silverman, M.A., 2019. Going Too Far Is the Same as Falling Short(dagger): Kinesin-3 Family Members in Hereditary Spastic Paraplegia. Front Cell Neurosci 13, 419.

Gennerich, A., Carter, A.P., Reck-Peterson, S.L., Vale, R.D., 2007. Force-induced bidirectional stepping of cytoplasmic dynein. Cell 131, 952-965.

Gennerich, A., Schild, D., 2006. Finite-particle tracking reveals submicroscopic-size changes of mitochondria during transport in mitral cell dendrites. Phys Biol 3, 45-53.

Gross, S.P., 2004. Hither and yon: a review of bi-directional microtubule-based transport. Phys Biol 1, R1-11.

Guedes-Dias, P., Holzbaur, E.L.F., 2019. Axonal transport: Driving synaptic function. Science 366.

Hasegawa, S., Sagawa, T., Ikeda, K., Okada, Y., Hayashi, K., 2019. Investigation of multiple-dynein transport of melanosomes by non-invasive force measurement using fluctuation unit chi. Sci Rep 9, 5099.

Hayashi, K., Hasegawa, S., Sagawa, T., Tasaki, S., Niwa, S., 2018a. Non-invasive force measurement reveals the number of active kinesins on a synaptic vesicle precursor in axonal transport regulated by ARL-8. Phys Chem Chem Phys 20, 3403-3410.

Hayashi, K., Miyamoto, M.G., Niwa, S., 2021. Effects of dynein inhibitor on the number of motor proteins transporting synaptic cargos. Biophys J 120, 1605-1614.

Hayashi, K., Tsuchizawa, Y., Iwaki, M., Okada, Y., 2018b. Application of the fluctuation theorem for non-invasive force measurement in living neuronal axons. Mol Biol Cell, mbcE18010022.

Hirakawa, E., Higuchi, H., Toyoshima, Y.Y., 2000. Processive movement of single 22S dynein molecules occurs only at low ATP concentrations. Proc Natl Acad Sci U S A 97, 2533-2537.

Hirokawa, N., Noda, Y., Tanaka, Y., Niwa, S., 2009. Kinesin superfamily motor proteins and intracellular transport. Nat Rev Mol Cell Biol 10, 682-696.

Iqbal, Z., Rydning, S.L., Wedding, I.M., Koht, J., Pihlstrom, L., Rengmark, A.H., Henriksen, S.P., Tallaksen, C.M., Toft, M., 2017. Targeted high throughput sequencing in hereditary ataxia and spastic paraplegia. PLoS One 12, e0174667.

Klebe, S., Lossos, A., Azzedine, H., Mundwiller, E., Sheffer, R., Gaussen, M., Marelli, C., Nawara, M., Carpentier, W., Meyer, V., Rastetter, A., Martin, E., Bouteiller, D., Orlando, L., Gyapay, G., El-Hachimi, K.H., Zimmerman, B., Gamliel, M., Misk, A., Lerer, I., Brice, A., Durr, A., Stevanin, G., 2012. KIF1A missense mutations in SPG30, an autosomal recessive spastic paraplegia: distinct phenotypes according to the nature of the mutations. Eur J Hum Genet 20, 645-649.

Lam, A.J., Rao, L., Anazawa, Y., Okada, K., Chiba, K., Dacy, M., Niwa, S., Gennerich, A., Nowakowski, D.W., McKenney, R.J., 2021. A highly conserved 310 helix within the kinesin motor domain is critical for kinesin function and human health. Sci Adv 7.

Mallik, R., Carter, B.C., Lex, S.A., King, S.J., Gross, S.P., 2004. Cytoplasmic dynein functions as a gear in response to load. Nature 427, 649-652.

Naoi, N., Kagawa, Y., Nagino, K., Niwa, S., Hayashi, K., 2021. Extreme-value analysis of intracellular cargo transport by motor proteins. bioRxiv.

Nishiyama, M., Higuchi, H., Yanagida, T., 2002. Chemomechanical coupling of the forward and backward steps of single kinesin molecules. Nat Cell Biol 4, 790-797.

Niwa, S., 2017. Immobilization of Caenorhabditis elegans to Analyze Intracellular Transport in Neurons. J Vis Exp.

Niwa, S., Lipton, D.M., Morikawa, M., Zhao, C., Hirokawa, N., Lu, H., Shen, K., 2016. Autoinhibition of a Neuronal Kinesin UNC-104/KIF1A Regulates the Size and Density of Synapses. Cell Rep 16, 2129-2141.

Peng, C.S., Zhang, Y., Liu, Q., Marti, G.E., Huang, Y.-W.A., Südhof, T.C., Cui, B., S., C., 2022. Nanometer-Resolution Long-term Tracking of Single Cargos Reveals Dynein Motor Mechanisms. bioRxiv.

Sasaki, K., Kaya, M., Higuchi, H., 2018. A Unified Walking Model for Dimeric Motor Proteins. Biophys J 115, 1981-1992.

Schmieg, N., Menendez, G., Schiavo, G., Terenzio, M., 2014. Signalling endosomes in axonal transport: travel updates on the molecular highway. Semin Cell Dev Biol 27, 32-43.

Schnitzer, M.J., Visscher, K., Block, S.M., 2000. Force production by single kinesin motors. Nat Cell Biol 2, 718-723.

Terenzio, M., Schiavo, G., Fainzilber, M., 2017. Compartmentalized Signaling in Neurons: From Cell Biology to Neuroscience. Neuron 96, 667-679.

Toba, S., Watanabe, T.M., Yamaguchi-Okimoto, L., Toyoshima, Y.Y., Higuchi, H., 2006. Overlapping hand-over-hand mechanism of single molecular motility of cytoplasmic dynein. Proc Natl Acad Sci U S A 103, 5741-5745.

Vale, R.D., 2003. The molecular motor toolbox for intracellular transport. Cell 112, 467-480.