# Neural attentional filters and behavioural outcome follow independent individual trajectories over the adult life span

Sarah Tune[1,2] & Jonas Obleser[1,2]

1 – Department of Psychology, University of Lübeck, 23562 Lübeck, Germany
2 – Center of Brain, Behavior, and Metabolism, University of Lübeck, 23562 Lübeck, Germany

*Author correspondence*:
Sarah Tune
Department of Psychology
University of Lübeck
Maria-Goeppert-Str. 9a
23562 Lübeck
Email: sarah.tune@uni-luebeck.de

**Running title:** Modelling trajectories of individual listening behaviour

Number of words in Abstract: 144
Number of words in Introduction: 755
Number of words in Discussion: 1683

**Keywords:** EEG, aging, speech processing, auditory attention, neural speech tracking, longitudinal analysis, latent change score models

**Conflict of Interest:** The authors declare no competing financial interests.

**Author contributions:**
S.T.: Conceptualization, data curation, formal analysis, investigation, visualization, methodology, writing–original draft, writing–review and editing
J.O.: Conceptualization, investigation, methodology, writing–review and editing, supervision, project administration, resources, funding acquisition

## Abstract

Preserved communication abilities promote healthy aging. To this end, the age-typical loss of sensory acuity might in part be compensated for by an individual's preserved attentional neural filtering. Is such a compensatory brain–behaviour link longitudinally stable? Can it predict individual change in listening behaviour? We here show that individual listening behaviour and neural filtering ability follow largely independent developmental trajectories modelling electroencephalographic and behavioural data of N=105 aging individuals (39–82 yrs). First, despite the expected decline in sensory acuity, listening-task performance proved remarkably stable over 2 years. Second, neural filtering and behaviour were correlated only within each separate measurement (T1, T2). Longitudinally, however, our results raise caution on attention-guided neural filtering metrics as predictors of individual trajectories in listening behaviour: Neither neural filtering at T1 nor its T1–T2 change could predict individual two-year behavioural change, under a combination of modelling strategies.

## Introduction

Speech comprehension is an essential aspect of human communication, enabling us to understand and interact with others effectively. Preserved communication is therefore critical to social well-being and healthy aging[1]. Any translational advance aimed at maintaining and restoring successful cognitive aging crucially relies on understanding the factors that explain and predict individual trajectories of listening performance. However, the evidence on these potential factors is astonishingly limited.

As we age, our ability to comprehend speech can decline due to age-related changes in the auditory system (i.e., sensory acuity) and in cognitive resources. Age-related hearing loss reduces the ability to detect and to discriminate speech sounds, especially in noisy environments. However, it has long been recognised that increasing age and hearing loss cannot fully account for the considerable degree of inter-individual differences observed in listening behaviour and its lifespan change[2-4].

Recent research has focused on the neurobiological mechanisms that promote successful speech comprehension by implementing 'neural filters' that segregate behaviourally relevant from irrelevant sounds[5-7]. A growing body of evidence suggests that speech comprehension is neurally supported by an attention-guided filter mechanism arising from primary auditory and perisylvian brain regions: By synchronizing its neural activity with the temporal structure of the speech signal of interest, the brain 'tracks' behaviourally relevant auditory inputs to enable attentive listening[8-11].

In a large age-varying cohort (N=155; 39–80 yrs), we have previously shown how the fidelity of this neural filtering strategy can help explain differences in listening behaviour (i) from individual to individual, and (ii) within individuals from sentence to sentence[12-14]. As participants performed a challenging dual-talker listening task, we recorded their electroencephalogram (EEG). We observed that enhanced neural filtering—defined as stronger neural tracking of attended vs. ignored speech—led to more accurate and overall faster responses. Notably, we observed both neural filtering as well as its link to behaviour to be independent of chronological age and severity of hearing loss[12].

The observation of such brain–behaviour relationships critically advances our understanding of the neurobiological foundation of cognitive functioning. Their translational potential as neural markers predictive of behaviour, however, is often only implicitly assumed but seldomly put to the test[15].

Using auditory cognition as a model system, we here overcome this limitation by testing directly the hitherto unknown longitudinal stability of neural filtering as a neural compensatory mechanism upholding communication success. Going even further, we ask to what extent an individual's attentional neural-filtering ability measured at a given moment is able to predict their future trajectory in listening performance. Only if this is the case, and only if such an association can plausibly be assumed to be causal for future changes in communication ability, neural filtering would be a potential translational target.

We here aim to fill this gap by analysing two-year changes in the sensory, neural, and
behavioural domain in a longitudinal subsample (N=105; 39–82 yrs) of our original
cohort[12] (see Fig. 2A). We apply an advanced combination of cross-sectional and
longitudinal modelling strategies to address the following specific research questions
(see Fig. 1).

First, by focusing on each domain individually, we ask how sensory, neural and
behavioural functioning evolve cross-sectionally across the middle and older adult life
span, and, more importantly, how they change longitudinally across the studied two-
year period. We expect individuals' hearing acuity and behaviour to decrease from T1
to T2, but no systematic longitudinal change in neural filtering.

Second, we test the longitudinal stability of the previously observed age- and
hearing-loss–independent effect of neural filtering on both accuracy and response
speed (Fig. 1A). To this end, we analyse the multivariate direct and indirect relationships
of hearing acuity, neural filtering and listening behaviour within and across timepoints.

Third, leveraging the strengths of latent change score modelling[16,17], we fuse
cross-sectional and longitudinal perspectives to probe the role of neural filtering as a
precursor of behavioural changes in two different ways: we ask (i) whether an
individual's T1 neural filtering strength can predict the observed behavioural
longitudinal change, and (ii) whether two-year changes in neural filtering can explain
concurrent changes in listening behaviour. Here, irrespective of the observed
magnitude and direction of T1–T2 developments, two scenarios are conceivable: Intra-
individual neural and behavioural changes may be either be correlated—lending
support to a compensatory role of neural filtering—or instead follow independent
trajectories[18] (see Fig. 1B, C).

Answering these questions is vital for understanding the neurobiological
mechanisms of successful communication across the lifespan. Answering them will also
critically inform the development of interventions targeted at maintaining or restoring
communication success and therefore concerns basic and applied researchers alike.

**Figure 1. A** Listening behaviour at a given timepoint is shaped by an individuals' sensory and neural functioning. Increased age decreases listening behaviour both directly, and indirectly via age-related hearing loss. Listening behaviour is supported by better neural filtering ability, independently of age and hearing acuity. **B** Conceptual depiction of how two-year changes along the neural (blue) and behavioural (red) domain may be related. Left, Thin coloured lines show individual trajectories across the adult lifespan, thick lines and black arrows highlight two-year changes in a single individual. Right, Across individuals, co-occurring changes in the neural and behavioural domain may be correlated (top) or independent of one another (bottom). **C** Schematic diagram highlighting the key research questions and how they are addressed in the current study using latent change score modelling.

# Results

We studied an age-varying cohort of healthy middle-aged and older adults longitudinally (N=105, 39–82 yrs, median age at T2: 63 yrs). The mean time difference between the two measurement timepoints reported here was 23.2 (sd: 4.0) months. We characterise the multivariate relationship of key measures of sensory, neural, and behavioural functioning to explain and predict individual trajectories of listening performance.

At each of two measurement timepoints, participants underwent audiological assessment followed by EEG recording during which they performed a difficult dual-talker dichotic listening task (see Fig. 2A,B)[12,19,20]. In each trial of the task, participants listened to two temporally aligned but spatially separated five-word sentences. They then had to identify the final word in one of the two sentences from a visual array of four alternatives (Fig. 2C). In 50 % of the trials, a visual spatial-attention cue indicated the side

of target sentence presentation, the other half of trials were preceded by an
uninformative neutral cue.

We extracted individuals' mean accuracy and response speed (calculated as the
inverse of reaction time) across all trials as key readouts of listening behaviour. On the
basis of source-localised 1–8 Hz auditory cortical activity, we further quantified
individuals' neural filtering ability as their differential neural tracking of attended vs.
ignored speech (Fig 2D; see also ref.[12] and Methods for details).

We follow a three-step analysis strategy to address our specific research
questions: First, we provide a largely descriptive overview of the observed average two-
year changes per studied domain. Second, we follow up on this fundamental analysis
with a causal mediation analysis[21] and single-trial mixed-effects model analysis geared
to assess the longitudinal stability of our recently reported effects of age, hearing acuity
and neural filtering on listening task performance. Third and final, we integrate and
extend the first two analysis perspectives in a joint latent change score model (LCSM)[16]
to most directly probe the role of neural filtering ability as a predictor of future attentive
listening ability.

167
168

169
170
**Figure 2. A** Longitudinal cohort of healthy middle-aged and older adults measured twice, two years apart.
Circles represent individual participants at a given measurement time (dark grey: timepoint (T) 1, light grey:
T2, white: drop-outs after T1). Bottom, Age distribution at T1 and T2 across 5-year bins. **B** Left, T1 air
conduction hearing thresholds per individual (thin grey lines) and age group (thick coloured lines). Note
that for didactic purposes, throughout the manuscript, thresholds are expressed as –dB HL to highlight the
decrease in hearing acuity with age (left). Right, Pure-tone average hearing acuity (0.5, 1, 2, and 4kHz across
both ears; higher is better) negatively correlates with age (r=−.43, p=3.73×10⁻⁶). **C** Participants listened to
two sentences presented simultaneously to the left and right ear. In 50 % of trials, a preceding visual cue

indicated the to-be-attended target sentence. Listening behaviour is quantified via the accuracy and speed
in identifying the final word of the target sentence. **D** Left, neural speech tracking as a proxy of an
individual's neural filtering ability. Stimulus envelopes of attended and ignored sentences were
reconstructed from source-localized EEG activity in auditory cortex (see Methods for details) and correlated
with the actual envelopes. Right, Better neural filtering results from stronger neural tracking of attended
compared to ignored speech. We analysed neural filtering derived from the entire sentence presentation
period.
186

### *Listening performance remains stable despite decreased hearing acuity*

In a first analysis (Fig. 3), we characterised how hearing acuity, neural filtering, and
listening performance change across the middle to older adult lifespan. Additionally, we
analysed longitudinal change from timepoint 1 (T1) to timepoint 2 (T2). We used the
same linear mixed-effect models to test cross-sectional effects of age and longitudinal
changes with time. We additionally quantified each measure's test-retest reliability as
their T1–T2 Pearson's correlation.

Note that throughout the manuscript and all analyses, we reversed the sign of
pure-tone average (PTA) values to express them as an index of hearing acuity rather
than hearing loss (i.e., higher values indicating better acuity). Similarly, for more intuitive
interpretation, accuracy is visually presented as mean proportion correct but was logit
transformed for all statistical analyses to satisfy model assumptions.
199

As expected, hearing acuity decreased linearly with increasing age (Fig. 3A, $\beta$=−3.4, SE
(standard error)=.71, p<.001) and on average by 1.2 dB from T1 to T2 ($\beta$=−1.18, SE=.27,
p<.001; $mean_{T1}$: −13.72 dB HL (sd: 7.8); $mean_{T2}$: −14.90 dB HL (8.3)). The effect of age did
not change with time (age x timepoint $\beta$=−.35, SE =.28, p=.21). Assuming constant
individual progression rates, this observed change corresponds to a projected average
decrease in hearing acuity per decade of −6.3 (sd:15.3) dB HL (see Fig. S1). The
magnitude of observed and projected hearing loss progression is well in line with recent
large-sample reports[22-24]. Measurements of hearing acuity showed high test-retest
reliability (r=.94, p<.001), underscoring the high fidelity of our audiological assessment.
209

In line with known deleterious effects of age, both behavioural outcomes (response
speed and accuracy) declined with increasing age, and did so to a similar degree in T1
and T2 (Fig. 3C, D; Speed: $\beta$=−.02, SE=.01, p=.004; Accuracy: $\beta$=−.23, SE=.07, p=.0001;
timepoint x age ps>.12). At the same time, and contrary to our expectations, average
performance levels remained remarkably stable from T1 to T2 (Speed: $\beta$=.004, SE=.01,
p=.44; $mean_{T1}$: .62 $s^{-1}$ (.08); $mean_{T2}$: .62 $s^{-1}$ (.08); Accuracy: $\beta$=.04, SE=.05, p=.36, $mean_{T1}$:
.88 % (.09); $mean_{T2}$: .88. % (.11)). Accuracy and response speed showed moderately high
test-retest reliability (Speed: r=.70, p<.001; Accuracy: r= .72, p<.001).
218

The analysis of change in neural filtering revealed that its strength varied independently
of age at both timepoints (Fig. 3B, $\beta$=.0003, SE=.0005, p=.48; timepoint x age $\beta$=−.0002,
SE=.0006, p=.79), confirming our previously reported T1 results[12]. As shown in Figure 3
(bottom left panel), magnitude and direction of observed longitudinal change are
highly variable across individuals and age groups, and we did not find evidence of any

systematic group-level change from T1 to T2 (β=.001, SE=.001, p=.16). In addition,

individual neural filtering strength correlated only weakly across time (r=.21, p=.03).

We also assessed the reliability of two established neural traits using resting-

state EEG from the same recording sessions: The individual alpha frequency (IAF)[25] and

the slope of 1/f neural noise[26,27]. As expected, both metrics showed high test-retest

reliability (IAF: r=.83, p<.001; 1/f slope: r=.78, p<.001). These findings provide a reference

level on reliability, demonstrating that the weak reliability of the neural filtering metric

is not due primarily to differences in EEG signal quality across sessions.

The temporal instability of neural filtering challenges its status as a potential

trait-like neural marker of attentive listening ability. At the same time, it does not

necessarily preclude the longitudinal stability of the previously observed brain–

behaviour link (see Fig. 1). This is being addressed next.

**Figure 3.** Cross-sectional and longitudinal change along the auditory sensory (**A**), neural (**B**), and
behavioural (**C, D**) domain. For each domain, coloured vectors (colour-coding four age groups for
illustrative purposes, only) in the respective left subpanels show an individual's change from T1 to T2 along
with the cross-sectional trend plus 95% confidence interval (CI) separately for T1 (dark grey) and T2 (light
grey). Top right subpanels: correlation of T1 and T2 as measure of test-retest reliability along with the 45°
line (grey) and individual data points (black circles). Bottom right panels: Mean longitudinal change per age
group (coloured vectors) and grand mean change (grey). Note that accuracy is expressed here as proportion
correct for illustrative purposes, but was analysed logit-transformed or by applying generalized linear
models.

## *Neural filtering reliably supports listening performance independent of age and hearing status*

On the basis of the full T1–T2 dataset, we aimed to replicate our key T1 results and test

their longitudinal stability: We expected an individual's neural filtering ability to impact

their listening outcome independently of age or hearing status[12]. Given the moderately

strong correlation of age and hearing acuity (r=−.43; p<.001; Fig. 2B), we employed

causal mediation analysis to model the direct as well as the hearing-acuity–mediated

effect of age on the behavioural outcome[21]. To formally test the stability of direct and
indirect relationships across time, we used a moderated mediation analysis. In this
analysis, the inclusion of interactions by timepoint tested whether the influence of age,
sensory acuity, and neural filtering varied across time.

Our expectations on the direct relationships were indeed borne out by the data:
higher age was associated with poorer hearing ability ($\beta$=−.43, SE=.09, p<.001) and
listening performance (Speed: $\beta$=−.33, SE=.06, p<.001; Accuracy: $\beta$=−.26, SE=.06,
p<.001). Better hearing ability, on the other hand, boosted accuracy but not response
speed (Accuracy: $\beta$=.30, SE=.1, p=.003; Speed: $\beta$=−.06, SE=.1, p=0.56). These direct
effects remained stable from T1 to T2 (all interactions by timepoint p>.56; all log Bayes
Factors (logBF01)>2.5).

Age also impacted accuracy indirectly: The total effect of age was partially
mediated via its detrimental effect on hearing acuity (Average causal mediation effect
(ACME), $\beta$=−.12, SE=.04, p<.001). We did not find evidence for an analogous indirect
effect on speed (ACME: $\beta$=.008, SE=.03, p=.77). Again, the hearing-acuity mediated
effect of age on accuracy did not change from T1 to T2 as evidenced by moderated
mediation analysis (interaction by timepoint p=.73).

Speaking to the robustness of our previous results, we observed the beneficial
effect of stronger neural filtering fidelity on both measures of listening performance
(Accuracy: $\beta$=.21, SE=, p=.02; Speed: $\beta$=.33, SE=.09, p<.001). Note that the magnitude of
this direct brain–behaviour effect is comparable to that of the direct effect of age.
Alternative models that included indirect, neural filtering-mediated paths from either
age or hearing acuity to behaviour did not reveal any significant mediation effects.

Most importantly, the longitudinal stability of the observed direct brain-
behaviour link was further supported by the absence of any significant changes with
time (interactions by timepoint; ps>.28, logBF01s>2.1).

In our previous T1 analysis (N=155), we had found evidence for the here analysed brain-
behaviour link at two different levels of observation: (i) at the trait level—individuals
with overall stronger neural filtering also performed better overall—and (ii) at the state-
level—stronger neural filtering in a given trial raised the chances of responding
correctly. Aiming at replication of the state-level (i.e., within-participant) relationship,
we ran a single-trial linear mixed-effects model analysis on our longitudinal N=105
sample. This analysis utilised single-trial data of both T1 and T2.

Lending credibility to our previous results, we found that stronger single-trial
neural filtering was associated with higher listening success at both T1 and T2 (logistic
mixed-effects model; within-participant effect of neural filtering on accuracy: odds ratio
(OR)=1.08, SE=.02, p<.001; interaction neural filtering x timepoint: OR=.99, SE=.03,
p=.82; see also Fig. 4C).
294

**Figure 4**. Longitudinal stability of sensory and neural determinants of listening behaviour. Causal mediation analysis of age, hearing acuity, and neural filtering on response speed (**A**) and logit-transformed accuracy (**B**). Graphs next to each path indicate standardised coefficients plus 95% CIs separately for the full dataset (black), T1 (dark grey), and T2 (light grey). We did not find any significant modulation of the observed effects with time (see text for results). **C** Neural filtering strength was found to be predictive of accuracy at the single-trial level at both T1 (top) and T2 (bottom). Grey lines show individual effects, blue thick line shows the group-level fixed effect along with 95 % confidence interval. OR=odds ratio, ** p<.01.

### Accuracy is longitudinally stable but speed and neural filtering increase at T2

Having established the longitudinal stability of the beneficial impact of intact neural filtering on listening performance, we turned to our final, most comprehensive analysis.

In a latent change score model (LCSM), in its bivariate form sometimes termed a parallel process model[16,17], we connected the neural and behavioural domain. This allowed us to more directly probe the potential role of neural filtering as a precursor of behavioural changes. Specifically, we asked: (i) Is an individual's baseline (T1) level of neural filtering ability predictive of their two-year change in behaviour; and (ii) are individual differences in longitudinal dynamics in the behavioural domain associated with those in the neural domain?

As a technical note, it is worth reiterating that in the present data, the highly variable, weakly reliable surface measure of neural tracking was nonetheless robustly connected to the behavioural outcome (see above; see Fig. 4). It is in such scenarios that the LCSM framework comes with particular methodological benefits: By expressing individuals' T1 and T2 levels, as well as their T1–T2 change as latent variables instead of manifest indicators, these types of models circumvent the calculation of notoriously unreliable noisy difference scores. They also avoid potential regression to the mean due to random errors[17,28]. Instead, the measurement errors of both, latent variables and their associated indicators, are explicitly modelled and thus effectively removed from the estimates of individual differences and relationships of interest[16].

Accordingly, for our metrics, we estimated T1 and T2 latent variables of behavioural and neural filtering from two manifest indicators each. These indicators were the average of each metric across the first and second half of the experiment, respectively (see Methods for details). For all measurement models, the standardized factor loadings were significant (all ps<.05; all standardized λ>.55). The assumption of

strict factorial invariance across time could be maintained for all models (all $\Delta\chi^2_{(df=1)}$ < 3.4, all ps > .07).

We then constructed univariate latent change score models (LCSM) to test for significant mean change in each metric from T1 to T2 while adjusting for their respective baseline (T1) level. The univariate models had acceptable (Speed: $\chi^2_{(df=5)}$ =9.7, p=.085, CFI=.988, RMSEA=.094] to excellent fit [Accuracy: $\chi^2_{(df=5)}$=5.1, p=.40, CFI=.999, RMSEA=.016; Neural filtering: $\chi^2_{(df=5)}$=.6, p=.99, CFI=1, RMSEA=0) according to established indices[29].

On average, listening task accuracy remained stable ($b_0$=.053, SE=.045, $\Delta\chi^2_{(df=1)}$ =1.33, p=.25). Response speed, on the other hand, showed a significant mean increase over time ($b_0$=.13, SE=.03, $\Delta\chi^2_{(df=1)}$ =12.79, p<.001; Fig. 5A). Similarly, the model of neural filtering showed a (marginally) significant mean increase ($b_0$=.24, SE=.11, $\Delta\chi^2_{(df=1)}$ =2.98, p=.08; Fig. 5A).

**Figure 5. A** Univariate latent change score models for response speed (left) and neural filtering (right). All paths denoted with Latin letters refer to freely estimated but constrained to be equal parameters of the respective measurement models. Greek letters refer to freely estimated parameters of the structural model. Highlighted in black is the estimated mean longitudinal change from T1 to T2. **B** Latent change score model (LCSM) relating two-year changes in neural filtering strength to changes in response speed. Black arrows indicate paths or covariances of interest. Solid black arrows reflect freely estimated and statistically significant effects, dashed black arrows reflect non-significant effects. All estimates are standardised. Grey arrows show paths that were freely estimated or fixed as part of the structural model but that did not relate to the main research questions. For visual clarity, manifest indicators of the measurement model and all

symbols relating to the estimated mean structure are omitted but are identical to those shown in panel A. ***p<.001, **p<.01, *p<.05, †p=.08. **C** Scatterplots of model-predicted factor scores that refer to the highlighted paths in panel B. Top panel shows that baseline-level neural filtering did not predict two-year change in behavioural functioning, bottom panel shows the absence of a significant change-change correlation.

### *Baseline neural functioning does not predict future change in listening behaviour*

Based on the univariate results, we then connected the two metrics that showed a significant mean change, namely speed and neural filtering, in a bivariate model of change (Fig. 5B; see also Fig. S2 for full model details).

In line with our hypotheses, we modelled the longitudinal impact of T1 neural functioning on the change in speed, as well as the change-change correlation. We also included individuals' age at T1 as a time-invariant covariate to control for its known influence on neural and behavioural functioning at T1, and to test for its association with longitudinal change. The model fit the data well ($\chi^2_{(df=27)}$=25.65, p=.54, CFI=1, RMSEA=0, 95 % CI[0 .07]).

Having ensured factorial invariance and goodness of fit, we can confidently interpret the estimates of individual differences and bivariate relationships that speak to our specific research questions. Crucial to our change-related inquiries, we observed reliable variance (i.e., individual differences) in the longitudinal change in both speed (standardized estimate $\phi$=0.88, SE=.07, $\Delta\chi^2_{(df=5)}$=94.07, p<.01) and neural filtering ($\phi$=.81, SE=.14, $\Delta\chi^2_{(df=4)}$=25.64, p<.001).

Individuals' baseline levels of both speed and neural filtering strength were predictive of their respective longitudinal change: Individuals with relatively strong neural filtering or fast responses at T1 showed a smaller increase from T1 to T2, possibly indicating ceiling effects (Speed: β=−.43, SE=.12, $\Delta\chi^2_{(df=1)}$=8.75, p=.003; Neural filtering: β=−.44, SE=.16, $\Delta\chi^2_{(df=1)}$=4.17, p=.04).

We also observed that participants' age at T1 covaried with the individual degree of change in speed but not with that in neural filtering: The older a participant at T1, the smaller their longitudinal increase in speed (Speed: $\phi$=−.24, SE=.11, $\Delta\chi^2_{(df=1)}$=4.38, p=.037; Neural filtering: $\phi$=−.02, SE=.14, $\Delta\chi^2_{(df=1)}$=.02, p=.89).

Importantly, however, an individual's latent T1 level of neural filtering strength was not predictive of the ensuing latent T1–T2 change in response speed (β=.02, SE=.16, $\Delta\chi^2_{(df=1)}$=.02, p=.90).

### *Neural filtering ability and listening behaviour follow independent developmental trajectories in later adulthood*

Finally, we turn to the last piece in our investigation where we address the question of whether individual differences in the neural and behavioural longitudinal change are connected. In other words: Are the contemporaneous changes along the two studied domains correlated or do they occur largely independently of one another?

Change score modelling revealed that longitudinal change in the neural and the behavioural domain occurred largely independent of one another despite their

systematic relationship within each separate measurement timepoint ($\phi$=.25, SE=.15,
$\Delta\chi^2_{(df=1)}$=2.74, p=.1). In other words, those individuals who showed the largest change
in neural filtering were not necessarily the ones who also changed the most in terms of
their behavioural functioning (see Fig. 5C bottom panel and Fig. S3).

In sum, at the group-level, we observed significant longitudinal increases in
neural filtering strength and listening performance. Importantly, inter-individual
differences in behavioural change could only be predicted by baseline age and baseline
behavioural functioning, and did not correlate with contemporaneous neural changes.

## Discussion

Successfully comprehending speech in noisy environments is a challenging task,
particularly for aging listeners whose hearing ability gradually declines[30]. A much-
researched neural support mechanism here is attention-guided neural 'tracking' of
behaviourally relevant speech signals, as one neural strategy to maintain listening
success[7-10,14,31-33].

However, to date, it is unknown if the fidelity with which an individual
implements this filtering strategy even represents a stable neural trait-like marker of
individual attentive listening ability. Of direct relevant to any future translational efforts
building on neural speech tracking, it is also unknown whether differences in neural
filtering strength observed between aging listeners are predictive at all of how their
attentive listening ability will develop in the future.  We here have addressed these
questions leveraging a new representative prospective cohort sample of healthy
middle-aged to older listeners.

From T1 to T2, individuals' hearing ability worsened as expected[22-24]. Their
listening performance, however, stayed remarkably stable. In addition, an individual's
baseline (T1) neural filtering strength proved to be a strikingly poor indicator of their
future (T2) level of neural functioning. On the other hand, bolstering previous results,
neural filtering reliably supported listening behaviour—both at T1 and T2—and within
each timepoint at two levels of granularity: Individuals with generally stronger neural
tracking of target vs. distractor speech performed the listening task on average more
accurately and faster. They also had higher chances of responding correctly in single
trials with relatively strong neural filtering.

Crucially, momentary states of neural functioning were not predictive of future
behavioural change, and the dynamics of longitudinal change at the neural and
behavioural level appear to follow largely independent trajectories.

### *Neural filtering fidelity as a trait-like neural marker of individual attentive listening ability?*

In recent years, the enhanced representation of behaviourally relevant sounds via their
prioritised neural tracking has been reported in numerous listening studies

investigating different acoustic environments, participant populations, and stages of
auditory processing[12,13,34-39].

This neural signature is commonly interpreted as a neural instantiation of
selective auditory attention in the service of successful speech comprehension. Note
however that its link to behaviour is not always explicitly established (but see refs. [12-
14,40,41]). Given how robustly the current data show the enhanced neural tracking of
attended vs. ignored speech at the group level, the weak reliability of individual neural
filtering strength may come as a surprise. At the same time, stronger neural filtering was
reliably linked to better behavioural performance within both T1 and T2.

How can these two findings be reconciled? Based on the current and previous
results, what may be concluded about the role of neural filtering as a potential neural
marker of individual attentive listening ability?

Previous studies on attention-guided neural speech tracking have not provided
any direct evidence on the temporal stability of neural filtering nor on its relationship
with behaviour. Studies on related neural signatures such as speech-aligned auditory
brainstem responses, or the entrainment of auditory cortical activity to rhythmic (non-
speech) stimulation reported moderate to high reliability[42-45]. However, these studies
have (i) investigated the temporal stability across sessions spaced only days or weeks
apart, (ii) focused on younger normal-hearing populations, or (iii) quantified the neural
encoding of speech or non-speech stimulation that involved less or no attentional
control. These differences render a direct comparison to our approach difficult, but
there is reason to consider a model-derived, latent representation of neural filtering as
employed here the more generalisable metric[46].

Our current results challenge a view of individual neural filtering strength as a
stable neural trait marker of selective auditory attention. Instead, they closely align with
a view of attention-guided neural speech tracking as a form of 'neural entrainment in
the broad sense' that reflects a listener's neural attentional state[10,47,48].

Under this interpretation, the neural metric's poor test–retest reliability is
reconcilable with its stable link to listening behaviour at the different levels of
observation: An individual's ability to exert top-down selective attention to prioritize the
neural encoding of behaviourally relevant information is far from stable but fluctuates
at different time scales[49-52]. This entails that at a longer time scale, here captured by two
distinct measurement timepoints, aging individuals will differ with respect to their
overall level of neural filtering and associated listening behaviour. Their level of neural
and behavioural functioning will differ from other individuals' levels at the same
timepoint but also from their own level at a different timepoint. Moreover, a listener's
behavioural outcome at either timepoint is not only shaped by their broad neural
attentional state with which they enter a communication situation. It is also critically
influenced by short-term fluctuations in neural filtering strength around their current
overall level of neural functioning (see Fig. 4C)[12].

What does this mean for the potential translational value of neural tracking? The
highly dynamic nature gives neural tracking-based metrics value as online neural
indicators of a listener's momentary attentional focus. As such, they could serve as

critical neural read-outs in novel brain-computer interfaces such as neurally-steered
hearing aids[53-55]. At the same time, however, their temporal instability severely limits
their potential as translational targets for diagnosis and therapeutic intervention[45,56,57].
482

### *Individual trajectories of listening behaviour cannot be explained by changes within a single domain*

As a second central query of the current study, we went beyond the establishment of
robust brain–behaviour relationships and directly probed the potency of neural filtering
to predict behavioural change over time[15]. We asked whether individual trajectories of
listening behaviour could be predicted by past levels of neural filtering, or by co-
occurring changes in neural filtering.

Past studies have observed enhanced cortical speech tracking in aging
compared to young adults. This suggests a compensatory role of increased speech-
brain coupling to counteract the deleterious effect of age or of hearing loss on speech
comprehension[41,58-62]. As a corollary of this relationship, typically observed cross-
sectionally, one might expect an individual's neural filtering strength to be connected
not only to present but also to future trajectories of listening behaviour.

Such relationships, if shown to be causal, would provide the strongest evidence
for the role of neural speech tracking as a neural compensatory mechanism supporting
communication success[30]. However, when analysed in our longitudinal sample of aging
listeners, we did not find evidence for a predictive role of neural filtering despite
supporting brain–behaviour links observed within each time point. What are potential
reasons for this absent connection (Fig. 5)?

One obvious explanation, both in statistical and substantive terms, may lie in
the poor reliability of our neural filtering metric as discussed above. Analytically,
however, we were able to mitigate this problem by adopting a modelling approach
which effectively removes the influence of measurement error[28]. Still, individual change
in response speed could only be predicted by an individual's baseline speed and age
but not by baseline neural filtering nor its longitudinal change. These findings call for a
more substantive explanation.

While most desirable from a translational perspective and a core quest in the
cognitive neuroscience of aging, predicting change in cognitive functioning, here
listening behaviour, from baseline or longitudinal change in brain function or structure
is a non-trivial endeavour. Connecting individual trajectories of neural or cognitive
functioning goes beyond the establishment of domain-specific age trends[63,64]. It also
goes beyond the mere extrapolation of (age-independent) brain–behaviour
relationships observed at a given moment[65,66]. Indeed, empirical evidence—and to
some degree also theoretical grounds—for robust brain–behaviour baseline–change or
change–change associations are limited[18].

Most empirical studies reporting such significant cross-domain change-change
correlations have in fact connected behavioural change to alterations in brain structure
rather than brain function[67-72]. Focusing on structural change may be advantageous: not

only can structural feature be quantified more directly and reliably, they also follow
systematic age-dependent trajectories thereby providing clearer causal pathways for
ensuing behavioural change[73,74]. Still, less than half of the studies testing such cross-
domain associations have indeed observed them[18]. From the perspective of theoretical
models of neurobiological and cognitive aging[75-77], the absence of correlated
trajectories of neural and cognitive functioning may indeed be the more expected
result. These models highlight the multifaceted nature of healthy cognitive aging in
which environmental variables, neurobiology, and cognition are dynamically
interrelated[78]. Neural compensatory mechanisms, such as the neural filtering correlate
targeted here, are thought to offset structural decline but are themselves influenced by
a number of factors. This leads to increased inter-individual variability that may
circumvent the emergence of group-level correlated change relationships.

Additionally, it is important to bear in mind that the behavioural outcome of
interest, that is, the speed and accuracy with which an aging individual solves a difficult
listening situation, involves the orchestration of different perceptual and cognitive
processes[79,80]. We here focused on one candidate neurobiological implementation of an
auditory attentional filter to help explain inter-individual differences in listening
behaviour and its lifespan trajectories. Yet, aging individuals may rely on different
alternative neural or cognitive strategies[81]. A complete understanding of inter-
individual differences in listening behaviour in aging adults will therefore depend on a
number of different factors among which the attention-modulated tracking of relevant
speech constitutes one, potentially necessary, but not sufficient neural
correlate[7,10,12,82,83].

Not least, there are a number of methodological choices that might have
constrained the potential fidelity of our study. First, the current study was limited to two
distinct timepoints spaced only two years apart. This limits the ability to model linear as
well as non-linear dynamics of change. Second, it also does not allow the separation of
distinct patterns of change co-occurring at the same time: one continuous, constant
change with age along with a separate process in which relative change is proportional
to the level observed at prior timepoints[84]. Lastly, denser sampling across a longer time
interval would have also increased statistical power to detect correlated change[85]. It
would have also allowed to more directly test hypotheses on causal pathways by which
change in the neural domain should precede change in the behavioural domain.

The conclusions stand, though, that individual trajectories in listening
behaviour cannot be explained by longitudinal change along a single dimension.
Instead, a better understanding of the influences shaping individual listening behaviour
across the adult lifespan will critically rely on uncovering the relative contribution and
age-dependent dynamics of sensory, neural, and cognitive factors.

## Conclusion

The results presented here support the role of attention-guided neural filtering as a
readout of an individual's neural attentional state. At the same time, the state-like nature

of neural tracking-based metrics limits their translational potential as predictors of
longitudinal change in listening behaviour over middle to older adulthood. Our data
caution against explaining audiology-typical listening behaviour solely by changes in
aspects of neural functioning, as listening behaviour and neural filtering ability follow
largely independent developmental trajectories. Our results critically inform
translational efforts aimed at the preservation and restoring of speech comprehension
abilities in aging individuals.

## Materials and Methods

### Data collection

The analysed data are part of a large-scale longitudinal study on the neural and cognitive mechanisms supporting adaptive listening behaviour in a prospective cohort healthy middle-aged and older adults ("The listening challenge: How ageing brains adapt (AUDADAPT)"; https://cordis.europa.eu/project/rcn/197855_en.html). This project encompasses the collection of different demographic, audiological, behavioural, and neurophysiological measures across initially two time points spaced approximately two years apart. The analyses carried out on the data aim at relating adaptive listening behaviour to changes in different neural dynamics. Given the longitudinal nature of the current study, all procedures concerning data collection, as well as EEG recording and analysis are identical to those detailed in our recently published analysis of T1 data using the same experimental paradigm[12].

### Participants and procedure

We here report on a total N = 105 right-handed German native speakers (median age at timepoint 2 (T2) 63 years; age range 39–82 years; 61 females) who underwent audiological, behavioural, and electrophysiological (EEG) assessment at two separate timepoints. On average, the measurement timepoints were spaced 23.2 (± SD 4.0) months apart.

At timepoint 1 (T1), we had screened a total of N=184 participants. Included participants had normal or corrected-to-normal vision, and did not report any neurological, psychiatric, or other disorders. They were also screened for mild cognitive impairment using the German version of the 6-Item Cognitive Impairment Test (6CIT[86] and the MoCA[87]). Only participants with normal hearing or age-adequate mild-to-moderate hearing loss were included (see Fig. 2B for individual audiograms at T1). Handedness was assessed using a translated version of the Edinburgh Handedness Inventory[88]. As a result of the initial screening procedure, 17 participants were excluded prior to EEG recording due to a medical history or non-age-related hearing loss. Three participants dropped out of the study prior to EEG recording and an additional 9 participants were excluded from analyses after EEG recording due to incidental findings after structural MR acquisition (N=3), or due to EEG data quality issues (N=9). Again, all detailed criteria are to be found in ref.[12].

At T2, N=115 participants returned for follow-up measurements. All individuals passed the repeat screening procedures identical to those at T1. Ten participants had to be excluded from the analyses reported here: Three participants had dropped out prior to EEG recording, three participants were excluded due to EEG data quality issues, and four participants because their EEG data had been excluded at T1. This resulted in a final longitudinal sample of N=105 individuals.

Dropout at T2 could not be predicted from participants' T1 age, hearing loss, behavioural performance (accuracy, speed) or neural filtering strength (all ps>.13). This

indicates that compared to the full T1 cohort reported on in previous studies[12,19] our

reduced longitudinal sample was not biased in terms of sensory, cognitive, and neural

functioning.

At each measurement timepoint, participants underwent detailed pure-tone and

speech audiometric measurements, along with an extensive battery of cognitive tests

and personality profiling (see ref.[81] for details). On a separate day, we recorded

participants' electroencephalogram (EEG) during rest (5 min each of eyes-open and

eyes-closed measurements) followed by six blocks of the same dichotic listening task

(see Fig. 2C and ref.[12] for details).

Participants gave written informed consent and received financial

compensation (10€ per hour). Procedures were approved by the ethics committee of

the University of Lübeck and were in accordance with the Declaration of Helsinki.

### *Dichotic listening task*

At each timepoint, participants performed a previously established dichotic listening

task[20] We provide full details on trial structure, stimulus construction, recording and

presentation in our previously published study on the first (N=155) wave of data

collection[12].

In short, in each of 240 trials, participants listened to two competing, dichotically

presented five-word sentences spoken by the same female speaker. They were probed

on the sentence-final noun in one of the two sentences. Two visual cues preceded

auditory presentation. First, a spatial-attention cue either indicated the to-be-probed

ear, thus invoking selective attention, or did not provide any information about the to-

be-probed ear, thus invoking divided attention. Second, but irrelevant to the current

study, a semantic cue specified a general or a specific semantic category for the final

word of both sentences, thus allowing to utilize a semantic prediction. Cue levels were

fully crossed in a 2×2 design and presentation of cue combinations varied on a trial-by-

trial level. All participants listened to the same 240 sentence pairs at each of the two

measurement timepoints. The order of sentence pair presentation was randomised for

each participant and at each timepoint.

To account for differences in hearing acuity within our group of participants, all

stimuli were presented 50 dB above the individual sensation level.

### *EEG recording and analysis*

The approach for EEG recording, preprocessing and subsequent analysis is identical to

the procedures carried out for T1 data collection and analysis[12].

In short, 64-channel EEG data were recorded, cleaned for artifacts using a

custom ICA-based pipeline, down-sampled to 125 Hz, filtered between 1–8Hz, and cut

into single-trials epochs covering the presentation of auditory stimuli. Following source-

localisation via beamforming, we focused on auditory cortical activity to train and test

decoding models of attended and ignored speech using cross-validated regularised

regression. Models were trained on selective-attention trials, only, but then also tested

on divided-attention trials. As results, we obtained single-trial reconstruction accuracy
(Pearson's r) estimates as metrics of the degree of attended and ignored neural speech
tracking, respectively.

We then calculated a neural filtering index across the entire sentence
presentation period. The index quantifies the difference in neural tracking of the to-be-
attended and of the to-be-ignored sentence (Neural filtering index = $(r_{attended} - r_{ignored})/($
$r_{attended} + r_{ignored})$, and thus indexes the strength of neural filtering at the single-trial level.
Positive values indicate successful neural filtering in line with the behavioural goal.

The EEG analyses were carried out in Matlab 2016b using the Fieldtrip toolbox
(v. 2017-04-28), the Human Connectome Project Workbench software (v1.5), FreeSurfer
(v.6.0), and the multivariate temporal response function (mTRF) toolbox (v1.5)[89].

## Behavioural and audiological data analysis

We evaluated participants' behavioural performance in the listening task with respect
to accuracy and response speed. For the binary measure of accuracy, we excluded trials
in which participants failed to answer within the given 4-s response window ('timeouts').
Spatial stream confusions, that is trials in which the sentence-final word of the to-be-
ignored speech stream were selected, and random errors were jointly classified as
incorrect answers. The analysis of response speed, defined as the inverse of reaction
time, was based on correct trials only.

We defined participants' hearing acuity as their pure-tone average (PTA)
composed of (air-conduction) hearing thresholds at the frequencies of .5, 1, 2, and 4 kHz.
Individual PTA values were then averaged across the left and right ear.

## Statistical analysis

For statistical analyses focused on between-participant ('trait-level') effects, behavioural
performance metrics and neural filtering index values were averaged across all trials and
experimental conditions to arrive at one trait-level estimate per participants. This
approach was also motivated by previous results based on T1 data: Here, we had
observed that stronger neural speech tracking led to overall faster and to more trials
with accurate responses irrespective of the specific cue-cue condition[12]. Accuracy was
logit-transformed for statistical analysis but expressed as proportion correct for
illustrative purposes. Similarly, for more intuitive interpretation, we reversed the sign of
PTA values for higher values to correspond to better hearing ability.

All analyses were performed in R (v.4.2.2)[90] using the packages lme4 (v.1.1-31)[91],
mediation (v4.5.0)[92], lavaan (v0.6-12)[93], and OpenMx (v2.21.8)[94].

## Linear mixed-effect modelling

As the first step of our three-part analysis approach, we applied general linear mixed-
effect models to tested for cross-sectional and longitudinal changes in trait-level

sensory acuity, neural filtering, and listening performance. These models included age,
timepoint and their interaction as fixed effect regressors and allowed random
participant-specific intercepts.

In the second step of the analysis, we also aimed at replicating the previously
observed single-trial state-level relationship of neural filtering and accuracy. To this end,
we applied a single generalised linear mixed-effects model (binomial distribution, logit
link function) on single-trial data of both T1 and T2. This model represents an adapted
version of the brain–behaviour model reported in Tune et al. (2021)[12]. In short, we
included all experimental manipulation predictors, as well as age, hearing acuity and
neural filtering metrics. We omitted previously shown to be non-significant higher-
order interactions and additionally included interactions by timepoint to directly test
for any longitudinal change in the effect of neural filtering on behaviour.

To tease apart state-level (i.e., within-participant) and trait-level (i.e., between-
participants) effects, we included two separate neural regressors: For the between-
participant effect regressor, we averaged single-trial neural filtering values per
individual across all trials. By contrast, the within-participant effect of interest was
modelled by the trial-by-trial deviation from the subject-level mean[95]. We included
participant- and item-specific random intercepts as well as random slopes for the effect
of the spatial-attention cue and the probed ear.

We used deviation coding for categorical predictors and z-scored all continuous
regressors. P-values for individual model terms in general linear mixed-effect models
are based on the Satterthwaite approximation for degrees of freedom, and on z-values
and asymptotic Wald tests for the generalised linear mixed-effect model of accuracy[96].
716

### Causal mediation analysis

We performed causal mediation analysis to model the direct as well as the hearing-
acuity–mediated effect of age on accuracy and response speed[21]. Critically, these
models also included a direct (i.e. independent of age and hearing loss) path of neural
filtering on behaviour. To formally test the stability of direct and indirect relationships
across time, we used a moderated mediation analysis[97]. In this analysis, the inclusion of
interactions by timepoint tested whether the influence of age, sensory acuity, and
neural functioning on behaviour varied across time. We z-scored all dependent and
independent variables, and estimated the magnitude of direct and mediated effects
along with percentile-based confidence intervals on the basis of 1,000 replications.

### Bayes Factor calculation

To facilitate interpretation of non-significant effects, we calculated the Bayes factor (BF)
based on the comparison of Bayesian information criterion (BIC)[98]. We report log Bayes
Factors, with a log BF01 of 0 representing equal evidence for and against the null
hypothesis; log BF01s with a positive sign indicating relatively more evidence for the
null hypothesis than the alternative hypothesis, and vice versa. Magnitudes > 1 are

taken as moderate, > 2.3 as strong evidence for either of the alternative or null
hypotheses, respectively.

### *Latent change score modelling*

In the third and final step of our analysis approach, we used structural equation
modelling (SEM) to investigate the role of neural filtering as a predictor of behavioural
change. We used latent change score modelling (LCSM)[16,17] to test (i) whether an
individual's T1 neural filtering strength can be predictive two-year changes in
behaviour, and (ii) whether two-year changes in neural filtering and listening behaviour
are systematically related. All models were specified and fitted with the R-based
package lavaan[93] using maximum likelihood estimation.

To bring all manifest variables onto the same scale while preserving mean
differences over time, we first stacked them across timepoint and then rescaled them
using the proportion of maximum scale ('POMS') method[99,100]. We assessed model fit
using established indices including the $\chi^2$ test, the root mean square error of
approximation (RMSEA), and the comparative fit index (CFI)[29,101]. Likelihood-ratio tests
helped us decide whether (i) constraining individual parameters to be equal
significantly decreased model fit, and (ii) individual parameter estimates were
statistically significant. We report standardized parameter estimates.

In a first step, we specified separate unstructured measurement models of accuracy,
speed, and neural filtering to establish factorial invariance across time using a series of
tests[17,102]. Latent variables representing each metric at T1 and T2 were constructed from
the observed individual means averaged across the first and second half of the
experiment, respectively. Covariances between T1 and T2 latent variables were freely
estimated, residual covariances were set to be equal for the first and second half of the
experiment. The factor loading and mean of the first manifest variable were set to 1 and
0, respectively, to ensure model identification. We sequentially tested for metric (i.e.,
identical factor loadings), strong (i.e., identical means), and strict (i.e., identical residuals)
time invariance using likelihood ratio tests.

Next, for those metrics surviving factorial invariance testing, we constructed
separate (i.e., univariate) change score models to test for group-level mean change
following the tutorial by Kievit et al. (2018). We included a regression path from the T1
latent variable to the latent change variable to test how baseline functioning impacted
the degree of longitudinal change. A likelihood ratio test of nested models (with mean
change being freely estimated vs. constrained to be 0) determined the significance of
group-level mean change.

In the last and final step, we constructed the bivariate LCSM that connected
changes in response speed and neural filtering, which both significantly increased over
time (see Fig. S2 for full model details). We modelled the covariance between the
baseline neural filtering and speed, as well as the covariance between neural and
behavioural change. In addition, we included a regression path from T1 neural filtering

to the T1-T2 change in response speed to directly test the predictive potency of baseline

neural functioning to explain behavioural change.

777            Following the logic of a generalised variance test[103], we tested for significant

778    inter-individual differences in neural and behavioural change. Note that such reliable

779    variance in change is a necessary prerequisite for testing change-change correlations.

780    In short, per domain (neural, behaviour), a likelihood ratio test compared the full model

781    with a restricted model in which the regression paths point to or covariance parameters

782    connected with the latent change variable were fixed to 0.

783            Finally, we asked whether neural and behavioural change covaried significantly.

784    We ran a likelihood ratio test on the full model compared to a restricted model in which

785    the covariance parameter was fixed to 0. For visualisation of results, we re-fit the final

786    model using OpenMx[94] to predict latent factor scores based on maximum likelihood.

### Data and code availability

789    The complete dataset associated with this work including raw data, EEG data analysis

790    results, as well as corresponding code will be publicly available under

791    https://osf.io/28r57/.

## References

1.    Lindenberger, U. Human cognitive aging: Corriger la fortune? *Science* **346,**
572–578 (2014).

2.    Akeroyd, M. A. Are individual differences in speech reception related to
individual differences in cognitive ability? A survey of twenty experimental
studies with normal and hearing-impaired adults. in **47,** S53–71 (2008).

3.    Houtgast, T. & Festen, J. M. On the auditory and cognitive functions that may
explain an individual's elevation of the speech reception threshold in noise.
*Int J Audiol* **47,** 287–295 (2008).

4.    Humes, L. E. *et al.* Central presbycusis: A review and evaluation of the
evidence. *J Am Acad Audiol* **23,** 635–666 (2012).

5.    Cherry, E. C. Some experiments on the recognition of speech, with one and
with two ears. *J. Acoust. Soc. Am.* **25,** 975–979 (1953).

6.    Broadbent, D. E. *Perception and communication*. (Pergamon Press, 1958).
doi:10.1186/s12889-020-10047-2

7.    Obleser, J. & Erb, J. in *The Cognitive Neurosciences* (eds. Poeppel, D., Mangun,
G. S. & Gazzaniga, M. S.) (MIT Press, 2020).
doi:10.7551/mitpress/11442.001.0001

8.    Ding, N. & Simon, J. Z. Emergence of neural encoding of auditory objects
while listening to competing speakers. *Proc. Natl. Acad. Sci. U.S.A.* **109,** 11854–
11859 (2012).

9.    Zion Golumbic, E. M. Z. *et al.* Mechanisms Underlying Selective Neuronal
Tracking of Attended Speech at a '"Cocktail Party"'. *Neuron* **77,** 980–991
(2013).

10.   Obleser, J. & Kayser, C. Neural Entrainment and Attentional Selection in the
Listening Brain. *Trends Cogn. Sci.* **23,** 913–926 (2019).

11.   Lakatos, P., Karmos, G., Mehta, A. D., Ulbert, I. & Schroeder, C. E. Entrainment of
neuronal oscillations as a mechanism of attentional selection. *Science* **320,**
110–113 (2008).

12.   Tune, S., Alavash, M., Fiedler, L. & Obleser, J. Neural attentional-filter
mechanisms of listening success in middle-aged and older individuals. *Nat*
*Commun* 1–14 (2021). doi:10.1038/s41467-021-24771-9

13.   O'Sullivan, J. A. *et al.* Attentional Selection in a Cocktail Party Environment Can
Be Decoded from Single-Trial EEG. *Cereb. Cortex* **25,** 1697–1706 (2014).

14.   Mesgarani, N. & Chang, E. F. Selective cortical representation of attended
speaker in multi-talker speech perception. *Nature* **485,** 233–236 (2012).

15.   Woo, C.-W., Chang, L. J., Lindquist, M. A. & Wager, T. D. Building better
biomarkers: brain models in translational neuroimaging. *Nat. Neurosci.* **20,**
365–377 (2017).

16.   McArdle, J. J. Latent Variable Modeling of Differences and Changes with
Longitudinal Data. *Annu. Rev. Psychol.* **60,** 577–605 (2009).

17.   Kievit, R. A. *et al.* Developmental cognitive neuroscience using latent change
score models: A tutorial and applications. *Dev Cogn Neurosci* **33,** 99–117
(2018).

18.   Oschwald, J. *et al.* Brain structure and cognitive ability in healthy aging: A
review on longitudinal correlated change. *Rev Neurosci* **31,** 1–57 (2019).

19.   Alavash, M., Tune, S. & Obleser, J. Dynamic large-scale connectivity of intrinsic
cortical oscillations supports adaptive listening in challenging conditions.
*PLoS Biol* **19,** e3001410 (2021).

20.   Alavash, M., Tune, S. & Obleser, J. Modular reconfiguration of an auditory
control brain network supports adaptive listening behavior. *Proc. Natl. Acad.*
*Sci. U.S.A.* **116,** 660–669 (2019).

21.   Imai, K., Keele, L. & Tingley, D. A general approach to causal mediation
analysis. *Psychol. Methods* **15,** 309–334 (2010).
22.   Linssen, A. M., Van Boxtel, M. P. J., Joore, M. A. & Anteunis, L. J. C. Predictors of
hearing acuity: Cross-sectional and longitudinal analysis. *J. Gerontol. A Biol. Sci.*
*Med. Sci.* **69,** 759–765 (2014).
23.   Rigters, S. C., Van Der Schroeff, M. P., Papageorgiou, G., Baatenburg De Jong,
R. J. & Goedegebure, A. Progression of hearing loss in the aging population:
Repeated auditory measurements in the Rotterdam study. *Audiol Neurootol*
**23,** 290–297 (2019).
24.   Wiley, T. L., Chappell, R., Carmichael, L., Nondahl, D. M. & Cruickshanks, K. J.
Changes in hearing thresholds over 10 years in older adults. *J Am Acad Audiol*
**19,** 281–92– quiz 371 (2008).
25.   Corcoran, A. W., Alday, P. M., Schlesewsky, M. & Bornkessel-Schlesewsky, I.
Toward a reliable, automated method of individual alpha frequency (IAF)
quantification. *Psychophysiology* **57,** e13064–21 (2018).
26.   Donoghue, T. *et al.* Parameterizing neural power spectra into periodic and
aperiodic components. *Nat. Neurosci.* 1–24 (2020). doi:10.1038/s41593-020-
00744-x
27.   Voytek, B. *et al.* Age-related changes in 1/f neural electrophysiological noise. *J.*
*Neurosci.* **35,** 13257–13265 (2015).
28.   McArdle, J. J. & Nesselroade, J. R. Using multivariate data to structure
developmental change. *Life-span developmental psychology: Methodological*
*contributions* 223–267 (1994).
29.   Raykov, T. & Marcoulides, G. A. On multilevel model reliability estimation from
the perspective of structural equation modeling. *Structural Equation Modeling:*
*A Multidisciplinary Journal* **13,** 130–141 (2006).
30.   Peelle, J. E. & Wingfield, A. The Neural Consequences of Age-Related Hearing
Loss. *Trends Neurosci.* **39,** 486–497 (2016).
31.   Power, A. J., Foxe, J. J., Forde, E.-J., Reilly, R. B. & Lalor, E. C. At what time is the
cocktail party? A late locus of selective attention to natural speech. *Eur. J.*
*Neurosci.* **35,** 1497–1503 (2012).
32.   O'Sullivan, J. *et al.* Hierarchical Encoding of Attended Auditory Objects in
Multi-talker Speech Perception. *Neuron* **104,** 1195–1204 (2019).
33.   Gross, J. *et al.* Speech Rhythms and Multiplexed Oscillatory Sensory Coding in
the Human Brain. *PLoS Biol* **11,** e1001752–14 (2013).
34.   Fiedler, L., Wöstmann, M., Herbst, S. K. & Obleser, J. Late cortical tracking of
ignored speech facilitates neural selectivity in acoustically challenging
conditions. *NeuroImage* **186,** 33–42 (2019).
35.   Kaufman, M. & Golumbic, E. Z. Listening to two speakers: Capacity and
tradeoffs in neural speech tracking during Selective and Distributed
Attention. *NeuroImage* **270,** 119984 (2023).
36.   Kraus, F., Tune, S., Ruhe, A., Obleser, J. & Wöstmann, M. Unilateral Acoustic
Degradation Delays Attentional Separation of Competing Speech. *Trends in*
*Hearing* **25,** 23312165211013242 (2021).
37.   Brodbeck, C., Jiao, A., Hong, L. E. & Simon, J. Z. Neural speech restoration at
the cocktail party: Auditory cortex recovers masked speech of both attended
and ignored speakers. *PLoS Biol* **18,** e3000883–22 (2020).
38.   Forte, A. E., Etard, O. & Reichenbach, T. The human auditory brainstem
response to running speech reveals a subcortical mechanism for selective
attention. *eLife* **6,** (2017).
39.   Fuglsang, S. A., Märcher-Rørsted, J., Dau, T. & Hjortkjær, J. Effects of
Sensorineural Hearing Loss on Cortical Synchronization to Competing Speech
during Selective Attention. *J. Neurosci.* **40,** 2562–2572 (2020).

40.   Orf, M., Wöstmann, M., Hannemann, R. & Obleser, J. Target enhancement but
not distractor suppression in auditory neural tracking during continuous
speech. *ISCIENCE* **26,** 106849 (2023).
41.   Schmitt, R., Meyer, M. & Giroud, N. Better speech-in-noise comprehension is
associated with enhanced neural speech tracking in older adults with hearing
impairment. *CORTEX* **151,** 133–146 (2022).
42.   Song, J. H., Nicol, T. & Kraus, N. Test-retest reliability of the speech-evoked
auditory brainstem response. *Clin Neurophysiol* **122,** 346–355 (2011).
43.   Easwar, V., Scollie, S., Aiken, S. & Purcell, D. Test-Retest Variability in the
Characteristics of Envelope Following Responses Evoked by Speech Stimuli.
*Ear and Hearing* **41,** 150–164 (2020).
44.   Cabral-Calderin, Y. & Henry, M. J. Reliability of Neural Entrainment in the
Human Auditory System. *J. Neurosci.* **42,** 894–908 (2022).
45.   Panela, R. A., Copelli, F. & Herrmann, B. Reliability and generalizability of
neural speech tracking in younger and older adults. *Nature Communications*
2023.07.26.550679 (2023). doi:10.1101/2023.07.26.550679
46.   Hertzog, C. & Schaie, K. W. Stability and change in adult intelligence: 2.
Simultaneous analysis of longitudinal means and covariance structures.
*Psychol Aging* **3,** 122–130 (1988).
47.   Lakatos, P. *et al.* Global dynamics of selective attention and its lapses in
primary auditory cortex. *Nat. Neurosci.* **19,** 1707–1717 (2016).
48.   Lakatos, P., Gross, J. & Thut, G. A New Unifying Account of the Roles of
Neuronal Entrainment. *Curr. Biol.* **29,** R890–R905 (2019).
49.   Schroeder, C. E. & Lakatos, P. Low-frequency neuronal oscillations as
instruments of sensory selection. *Trends Neurosci.* **32,** 9–18 (2009).
50.   Schroeder, C. E., Wilson, D. A., Radman, T., Scharfman, H. & Lakatos, P.
Dynamics of Active Sensing and perceptual selection. *Current Opinion in
Neurobiology* **20,** 172–176 (2010).
51.   Helfrich, R. F. *et al.* Neural Mechanisms of Sustained Attention Are Rhythmic.
*Neuron* **99,** 854–865.e5 (2018).
52.   Buschman, T. J. & Kastner, S. From Behavior to Neural Dynamics: An Integrated
Theory of Attention. *Neuron* **88,** 127–144 (2015).
53.   Van Eyndhoven, S., Francart, T. & Bertrand, A. EEG-informed attended speaker
extraction from recorded speech mixtures with application in neuro-steered
hearing prostheses. *IEEE Transactions on Biomedical Engineering* **64,** 1045–
1056 (2016).
54.   Ceolini, E. *et al.* Brain-informed speech separation (BISS) for enhancement of
target speaker in multitalker speech perception. 1–12 (2020).
doi:10.1016/j.neuroimage.2020.117282
55.   O'Sullivan, J. *et al.* Neural decoding of attentional selection in multi-speaker
environments without access to clean sources. *J Neural Eng* **14,** 056001 (2017).
56.   Gillis, M., Vanthornhout, J., Simon, J. Z., Francart, T. & Brodbeck, C. Neural
Markers of Speech Comprehension: Measuring EEG Tracking of Linguistic
Speech Representations, Controlling the Speech Acoustics. *J. Neurosci.* **41,**
10316–10329 (2021).
57.   Di Liberto, G. M., Hjortkjær, J. & Mesgarani, N. Editorial: Neural Tracking:
Closing the Gap Between Neurophysiology and Translational Medicine. *Front.
Neurosci.* **16,** 872600 (2022).
58.   Gillis, M., Decruy, L., Vanthornhout, J. & Francart, T. Hearing loss is associated
with delayed neural responses to continuous speech. *Eur. J. Neurosci.* **55,**
1671–1690 (2022).

59. Presacco, A., Simon, J. Z. & Anderson, S. Speech-in-noise representation in the aging midbrain and cortex: Effects of hearing loss. *PLoS ONE* **14,** e0213899–26 (2019).

60. Presacco, A., Simon, J. Z. & Anderson, S. Evidence of degraded representation of speech in noise, in the aging midbrain and cortex. *J. Neurophysiol.* **116,** 2346–2355 (2016).

61. Decruy, L., Vanthornhout, J. & Francart, T. Evidence for enhanced neural tracking of the speech envelope underlying age-related speech-in-noise difficulties. *J. Neurophysiol.* **122,** 601–615 (2019).

62. Decruy, L., Vanthornhout, J. & Francart, T. Hearing impairment is associated with enhanced neural tracking of the speech envelope. *Hearing Research* **393,** 107961 (2020).

63. Lindenberger, U., Oertzen, von, T., Ghisletta, P. & Hertzog, C. Cross-Sectional Age Variance Extraction: What's Change Got To Do With It? *Psychol Aging* **26,** 34–47 (2011).

64. Tucker-Drob, E. M. *et al.* A strong dependency between changes in fluid and crystallized abilities in human cognitive aging. *Sci Adv* **8,** eabj2422 (2022).

65. Boker, S. M. & Martin, M. A Conversation between Theory, Methods, and Data. *Multivariate Behav Res* **0,** 1–14 (2018).

66. Raz, N. & Lindenberger, U. Only time will tell: Cross-sectional studies offer no solution to the age-brain-cognition triangle: Comment on salthouse (2011). *Psychological Bulletin* **137,** 790–795 (2011).

67. Raz, N. Regional brain changes in aging healthy adults: general trends, individual differences and modifiers. *Cereb. Cortex* **15,** 1676–1689 (2005).

68. McArdle, J. J. *et al.* Structural modeling of dynamic changes in memory and brain structure using longitudinal data from the normative aging study. *J Gerontol B Psychol Sci Soc Sci* **59,** P294–304 (2004).

69. Bender, A. R., Prindle, J. J., Brandmaier, A. M. & Raz, N. White matter and memory in healthy adults: Coupled changes over two years. *NeuroImage* **131,** 193–204 (2016).

70. Ritchie, S. J. *et al.* Coupled changes in brain white matter microstructure and fluid intelligence in later life. *J. Neurosci.* **35,** 8672–8682 (2015).

71. Lövdén, M. *et al.* Changes in perceptual speed and white matter microstructure in the corticospinal tract are associated in very old age. *NeuroImage* **102,** 520–530 (2014).

72. Persson, N. *et al.* Regional brain shrinkage and change in cognitive performance over two years: The bidirectional influences of the brain and cognitive reserve factors. *NeuroImage* **126,** 15–26 (2016).

73. Bennett, I. J. & Madden, D. J. Disconnected aging: cerebral white matter integrity and age-related differences in cognition. *Neuroscience* **276,** 187–205 (2014).

74. Grady, C. The cognitive neuroscience of ageing. *Nat. Rev. Neurosci.* **13,** 491–505 (2012).

75. Reuter-Lorenz, P. A. & Park, D. C. How Does it STAC Up? Revisiting the Scaffolding Theory of Aging and Cognition. *Neuropsychol Rev* **24,** 355–370 (2014).

76. Cabeza, R. *et al.* Maintenance, reserve and compensation: the cognitive neuroscience of healthy ageing. *Nat. Rev. Neurosci.* **19,** 1–10 (2018).

77. Stern, Y., Barnes, C. A., Grady, C., Jones, R. N. & Raz, N. Brain reserve, cognitive reserve, compensation, and maintenance: operationalization, validity, and mechanisms of cognitive resilience. *Neurobiology of Aging* **83,** 124–129 (2019).

78. Freund, J. *et al.* Emergence of individuality in genetically identical mice. *Science* **340,** 756–759 (2013).

79. Anderson, S., White-Schwoch, T., Parbery-Clark, A. & Kraus, N. A dynamic auditory-cognitive system supports speech-in-noise perception in older adults. *Hearing Research* **300,** 18–32 (2013).

80. Peelle, J. E. Language and aging. *The Oxford Handbook of Neurolinguistics* (de Zubicaray and Schiller, eds), pp. 295–316. Oxford University Press.

81. Tune, S., Wöstmann, M. & Obleser, J. Probing the limits of alpha power lateralisation as a neural marker of selective attention in middle-aged and older listeners. *Eur. J. Neurosci.* **48,** 2537–2550 (2018).

82. Gillis, M., Van Canneyt, J., Francart, T. & Vanthornhout, J. Neural tracking as a diagnostic tool to assess the auditory pathway. *Hearing Research* 108607 (2022).

83. Strauß, A., Wöstmann, M. & Obleser, J. Cortical alpha oscillations as a tool for auditory selective inhibition. *Front. Hum. Neurosci.* **8,** 1–7 (2014).

84. Jacobucci, R., Serang, S. & Grimm, K. J. A Short Note on Interpretation in the Dual Change Score Model. *Structural Equation Modeling: A Multidisciplinary Journal* **0,** 1–7 (2019).

85. Rast, P. & Hofer, S. M. Longitudinal design considerations to optimize power to detect variances and covariances among rates of change: simulation results based on actual longitudinal studies. *Psychol. Methods* **19,** 133 (2014).

86. Jefferies, K. & Gale, T. M. in *Cognitive Screening Instruments* 209–218 (Springer London, 2012). doi:10.1007/978-1-4471-2452-8_11

87. Nasreddine, Z. S. *et al.* The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc* **53,** 695–699 (2005).

88. Oldfield, R. C. The assessment and analysis of handedness: The Edinburgh inventory. **9,** 97–113 (1971).

89. Crosse, M. J., Di Liberto, G. M., Bednar, A. & Lalor, E. C. The Multivariate Temporal Response Function (mTRF) Toolbox: A MATLAB Toolbox for Relating Neural Signals to Continuous Stimuli. *Front. Hum. Neurosci.* **10,** 604 (2016).

90. R Core Team. R: A language and environment for statistical computing. (2019). doi:https://www.R-project.org/

91. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using lme4. *J. Stat. Soft.* **67,** 1–48 (2015).

92. Tingley, D., Yamamoto, T., Keele, L. & Imai, K. Mediation: R package for causal mediation analysis. *J. Stat. Soft.* **59,** 1–38 (2014).

93. Rosseel, Y. lavaan: An R package for structural equation modeling. *J. Stat. Soft.* **48,** 1–36 (2012).

94. Neale, M. C. *et al.* OpenMx 2.0: Extended Structural Equation and Statistical Modeling. *Psychometrika* **81,** 535–549 (2016).

95. Bell, A., Fairbrother, M. & Jones, K. Fixed and random effects models: making an informed choice. *Qual Quant* **53,** 1051–1074 (2018).

96. Luke, S. G. Evaluating significance in linear mixed-effects models in R. *Behav Res* **49,** 1494–1502 (2017).

97. Preacher, K. J., Rucker, D. D. & Hayes, A. F. Addressing Moderated Mediation Hypotheses: Theory, Methods, and Prescriptions. *Multivariate Behav Res* **42,** 185–227 (2007).

98. Wagenmakers, E.-J. A practical solution to the pervasive problems of p values. *Psychonomic Bulletin & Review* **14,** 779–804 (2007).

99. Moeller, J. A word on standardization in longitudinal studies: don't. *Front. Psychology* **6,** 1389 (2015).

100. Little, T. D. *Longitudinal structural equation modeling.* (Guilford press, 2013).

101. Kline, R. B. *Principles and practice of structural equation modeling.* (2023).

102. Polk, S. E. *et al*. Change in Latent Gray-Matter Structural Integrity Is Associated With Change in Cardiovascular Fitness in Older Adults Who Engage in At-Home Aerobic Exercise. *Front. Hum. Neurosci.* **16,** 852737 (2022).

103. Brandmaier, A. M. Precision, Reliability, and Effect Size of Slope Variance in Latent Growth Curve Models: Implications for Statistical Power Analysis. 1–16 (2018). doi:10.3389/fpsyg.2018.00294

## Acknowledgments

Research was funded by the European Research Council (grant no. ERC-CoG-2014-646696 "Audadapt" awarded to J.O.).