Interpreting SNP heritability in admixed populations

Jinguo Huang[1,2], Nicole Kleman[3], Saonli Basu[4], Mark D. Shriver[2], and
Arslan A. Zaidi[†3,5]

[1]Bioinformatics and Genomics, Huck Institutes of the Life Sciences, Pennsylvania State University

[2]Department of Anthropology, Pennsylvania State University

[3]Department of Genetics, Cell Biology, and Development, University of Minnesota

[4]Department of Biostatistics, University of Minnesota

[5]Institute of Health Informatics, University of Minnesota

[†]Correspondence to A.A.Z (aazaidi@umn.edu)

August 2, 2024

Abstract

SNP heritability (h^2_{snp}) is defined as the proportion of phenotypic variance explained by genotyped SNPs and is believed to be a lower bound of heritability (h^2), being equal to it if all causal variants are known. Despite the simple intuition behind h^2_{snp}, its interpretation and equivalence to h^2 is unclear, particularly in the presence of population structure and assortative mating. It is well known that population structure can lead to inflation in $\hat{h}^2_{snp}$ estimates because of confounding due to linkage disequilibrium (LD) or shared environment. Here we use analytical theory and simulations to demonstrate that h^2_{snp} estimates can be biased in admixed populations, even in the absence of confounding and even if all causal variants are known. This is because admixture generates LD, which contributes to the genetic variance, and therefore to heritability. Genome-wide restricted maximum likelihood (GREML) does not capture this contribution leading to under- or over-estimates of h^2_{snp} relative to h^2, depending on the genetic architecture. In contrast, Haseman-Elston (HE) regression exaggerates the LD contribution leading to biases in the opposite direction. For the same reason, GREML and HE estimates of local ancestry heritability (h^2_γ) are also biased. We describe this bias in $\hat{h}^2_{snp}$ and $\hat{h}^2_\gamma$ as a function of admixture history and the genetic architecture of the trait and show that it can be recovered under some conditions. We clarify the interpretation of $\hat{h}^2_{snp}$ in admixed populations and discuss its implication for genome-wide association studies and polygenic prediction.

Introduction

The ability to estimate (narrow-sense) heritability (h^2) from unrelated individuals was a major advance in genetics. Traditionally, h^2 was estimated from family-based studies in which the phenotypic resemblance between relatives could be modeled as a function of their expected genetic relatedness [1]. However, this approach was limited to analysis of closely related individuals where pedigree information is available and the realized genetic relatedness is not too different from expectation [2]. With the advent of genome-wide association studies (GWAS), we hoped that many of the variants underlying this heritability would be uncovered. However, when genome-wide significant SNPs explained a much smaller fraction of the phenotypic variance, it became important to explain the missing heritability – were family-based estimates inflated or were GWAS just underpowered, limited by variant discovery?

Yang *et al.* (2010) [3] made the key insight that one could estimate the portion of h^2 tagged by genotyped SNPs, regardless of whether or not they were genome-wide significant, by exploiting the subtle variation in the realized genetic relatedness among apparently unrelated individuals [3–5]. This quantity came to be known colloquially as 'SNP heritability' (h^2_{snp}) and it is believed to be equal to h^2 if all causal variants are included among genotyped SNPs [3]. Indeed, estimates of h^2_{snp} explain a much larger fraction of trait heritability than GWAS SNPs [3], approaching family-based estimates of h^2 when whole genome sequence data, which captures rare variants, are used [6]. This has made it clear that GWAS have yet to uncover more variants with increasing sample size. Now, h^2_{snp} has become an important aspect of the design of genetic studies and is often used to define the power of variant discovery in GWAS and the upper limit of polygenic prediction accuracy.

Despite the utility and simple intuition of h^2_{snp}, there is much confusion about its interpretation and equivalence to h^2, particularly in the presence of population structure and assortative mating [7–12]. But much of the discussion of heritability in structured populations has focused on biases in $\hat{h}^2_{snp}$ – the estimator – due to confounding effects of shared environment and linkage disequilibrium (LD) with other variants [7, 9–11, 13]. There is comparatively little discussion, at least in human genetics, on the fact that LD due to population structure also contributes to genetic variance, and therefore, is a component of heritability [1] (but see [14–16] for a rigorous discussion). We think this is at least partly due to the fact that most studies are carried out in cohorts with primarily European ancestry, where the degree of population structure is minimal and large effects of LD can be ignored. However, that is not the case for diverse, multi-ethnic cohorts, which have historically been underrepresented in genetic studies, but thanks to a concerted effort in the field, are now becoming increasingly common [17–23]. The complex structure in these cohorts also brings unique methodological challenges and it is imperative that we understand whether existing methods, which have largely been evaluated in more homogeneous groups, generalize to more diverse cohorts.

Our goal in this paper is to study the behavior of $\hat{h}^2_{snp}$ in admixed populations. What is its interpretation in the ideal situation where causal variants are known? Is it an unbiased estimate of h^2? To answer these questions, we derived a general expression for the genetic variance in admixed populations, decomposing it in terms of the contribution of population structure, which influences both the genotypic variance at individual loci and the LD across loci. We used theory and simulations to show that $\hat{h}^2_{snp}$

67 estimated with genome-wide restricted maximum likelihood (GREML) [3, 5] and Haseman-Elston (HE)

68 regression [24] – two widely used approaches – can be biased in admixed and other structured popula-

69 tions, even in the absence of confounding and when all causal variants are known. We explain this in

70 terms of the discrepancy between the model assumed in $\hat{h}^2_{snp}$ estimation and the generative model from

71 which the genetic architecture of the trait in the population may have been sampled. We describe the

72 bias in $\hat{h}^2_{snp}$ as a function of admixture history and genetic architecture and discuss its implications for

73 GWAS and polygenic prediction accuracy.

Model

74

Genetic architecture

75

76 We begin by describing a generative model for the phenotype. Let $y = g + e$, where y is the phenotypic

77 value of an individual, g is the genotypic value, and e is random error. We assume additive effects such

78 that $g = \sum_{i=1}^{m} \beta_i x_i$ where β_i is the effect size of the i^{th} biallelic locus and $x_i \in \{0, 1, 2\}$ is the number

79 of copies of the trait-increasing allele. Importantly, the effect sizes are fixed quantities and differences in

80 genetic values among individuals are due to random variation in genotypes. Note, that this is different

81 from the model assumed by GREML where genotypes are fixed and effect sizes are random [14].

82 We denote the mean, variance, and covariance with $\mathbb{E}(.)$, $\mathbb{V}(.)$, and $\mathbb{C}(.)$, respectively, where the

83 expectation is measured over random draws from the population rather than random realizations of the

84 evolutionary process. We can express the additive genetic variance of a quantitative trait as follows:

$$V_g = \mathbb{V}(\sum_{i=1}^{m} \beta_i x_i) = \sum_{i=1}^{m} \beta_i^2 \, \mathbb{V}(x_i) + \sum_{j \neq i} \beta_i \beta_j \, \mathbb{C}(x_i, x_j)$$

85 Here the first term represents the contribution of individual loci (genic variance) and the second term

86 is the contribution of linkage disequilibrium (LD contribution). We make the assumption that loci are

87 unlinked and therefore, the LD contribution is entirely due to population structure. We describe the

88 behavior of V_g in a population that is a mixture of two previously isolated populations A and B that

89 diverged from a common ancestor. To do this, we denote θ as the fraction of the genome of an individual

90 with ancestry from population A. Thus, $\theta = 1$ if the individual is from population A, 0 if they are from

91 population B, and $\theta \in (0, 1)$ if they are admixed. Then, V_g can be expressed in terms of ancestry as

92 (Appendix):

$$V_g = 2 \, \mathbb{E}(\theta) \sum_{i=1}^{m} \beta_i^2 \, f_{\mathrm{i}}^{\mathrm{A}}(1 - f_{\mathrm{i}}^{\mathrm{A}}) + 2\{1 - \mathbb{E}(\theta)\} \sum_{i=1}^{m} \beta_i^2 \, f_{\mathrm{i}}^{\mathrm{B}}(1 - f_{\mathrm{i}}^{\mathrm{B}}) \tag{1.1}$$

$$+ 2 \, \mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} \sum_{i=1}^{m} \beta_i^2 (f_{\mathrm{i}}^{\mathrm{A}} - f_{\mathrm{i}}^{\mathrm{B}})^2 \tag{1.2}$$

$$+ 2 \, \mathbb{V}(\theta) \sum_{i=1}^{m} \beta_i^2 (f_{\mathrm{i}}^{\mathrm{A}} - f_{\mathrm{i}}^{\mathrm{B}})^2 \tag{1.3}$$

$$+ 4 \, \mathbb{V}(\theta) \sum_{i \neq j} \beta_i \beta_j (f_{\mathrm{i}}^{\mathrm{A}} - f_{\mathrm{i}}^{\mathrm{B}})(f_{\mathrm{j}}^{\mathrm{A}} - f_{\mathrm{j}}^{\mathrm{B}}) \tag{1.4}$$

93 where f_i^A and f_i^B are the allele frequencies in populations A and B, and $\mathbb{E}(\theta)$ and $\mathbb{V}(\theta)$ are the mean

94 and variance of individual ancestry. The sum of the first three terms represents the genic variance and

95 the last term represents the LD contribution.

Demographic history

97 From Eq. 1, it is clear that, conditional on the genetic architecture in the source populations (β, f^A, f^B),

98 V_g is a function of the mean, $\mathbb{E}(\theta)$, and variance, $\mathbb{V}(\theta)$, of individual ancestry in the admixed population.

99 We consider two demographic models that affect $\mathbb{E}(\theta)$ and $\mathbb{V}(\theta)$ in qualitatively different ways. In the

100 first model, the source populations meet once t generations ago (we refer to this as $t = 0$) in proportions

101 p and $1 - p$, after which there is no subsequent admixture (Fig. 1A). In the second model, there is

102 continued gene flow in every generation from one of the source populations such that the mean overall

103 amount of ancestry from population A is the same as in the first model (Fig. 1A). For brevity, we refer

104 to these as the hybrid-isolation (HI) and continuous gene flow (CGF) models, respectively, following

105 Pfaff *et al.* (2001) [25]. $\mathbb{V}(\theta)$ is also affected by ancestry-based assortative mating, where individuals are

106 more likely to partner with others of similar ancestry. We refer to this simply as assortative mating for

107 brevity and model this following Zaitlen *et al.* (2017) using a parameter $P \in (0, 1)$, which represents the

108 correlation of the ancestry of individuals across mating pairs in the population [26].

109 Under these conditions, the behavior of $\mathbb{E}(\theta)$ and $\mathbb{V}(\theta)$ has been described previously [26, 27] (Fig. 1B

110 and C). Briefly, in the HI model, $\mathbb{E}(\theta)$ remains constant at p in the generations after admixture as there

111 is no subsequent gene flow. $\mathbb{V}(\theta)$ is at its maximum at $t = 0$ when each individuals carries chromosomes

112 either from population A or B, but not both. This genome-wide correlation in ancestry breaks down

113 in subsequent generations as a function of mating, independent assortment, and recombination, leading

114 to a decay in $\mathbb{V}(\theta)$, the rate depending on the strength of assortative mating (Fig. 1C). In the CGF

115 model, both $\mathbb{E}(\theta)$ and $\mathbb{V}(\theta)$ increase with time as new chromosomes are introduced from the source

116 populations. But while $\mathbb{E}(\theta)$ continues to increase monotonically, $\mathbb{V}(\theta)$ will plateau and decrease due to

117 the countervailing effects of independent assortment and recombination which redistribute ancestry in

118 the population, reaching equilibrium at zero if there is no more gene flow and the population is mating

119 randomly. $\mathbb{V}(\theta)$ provides an intuitive and quantitative measure of the degree of population structure

120 (along the axis of ancestry) in admixed populations.

Results

Genetic variance in admixed populations

123 To understand the expectation of genetic variance in admixed populations, it is first worth discussing

124 its behavior in the source populations. In Eq. 1, the first term represents the within-population com-

125 ponent (V_{gw}) and the last three terms altogether represent the component of genetic variance between

126 populations A and B (V_{gb}). Note that $V_{gb} = \frac{(\bar{g_A} - \bar{g_B})^2}{2}$ is positive only if there is a difference in the mean

127 genotypic values (Fig. 2). This variance increases with genetic divergence since the expected values

128 of both $(f_i^A - f_i^B)^2$ and $(f_i^A - f_i^B)(f_j^A - f_j^B)$ are functions of F_{ST}. While $\beta_i^2 (f_i^A - f_i^B)^2$ is expected to

Figure 1: The behavior of mean and variance of individual ancestry as a function of admixture history. (A) Shows the demographic models under which simulations were carried out. Admixture might occur once (Hybrid Isolation, HI, left column) or continuously (Continuous Gene Flow, CGF, right column). (B) The mean individual ancestry, $\mathbb{E}(\theta)$ remains constant over time in the HI model and increases in the CGF model with continued gene flow. (C) The variance in individual ancestry, $\mathbb{V}(\theta)$ is maximum at $t = 0$ in the HI model, decaying subsequently. $\mathbb{V}(\theta)$ increases with gene flow in the CGF model and will subsequently decrease with time. P measures the strength of assortative mating, which slows the decay of $\mathbb{V}(\theta)$. P=0.6 is missing for simulations run for 50 and 100 generations and $\theta \in \{0.1, 0.2\}$ due to the difficulty in finding mate pairs (Methods).

129 increase monotonically with increasing divergence, $\beta_i\beta_j(f_i^A - f_i^B)(f_j^A - f_j^B)$ is expected to be zero under

130 neutrality because the direction of frequency change will be uncorrelated across loci. In this case, the LD

131 contribution, i.e., (1.4), is expected to be zero and $V_{gb} = (1.1) + (1.2) + (1.3)$. However, this is true only

132 in expectation over the evolutionary process and the realized LD contribution may be non-zero even for

133 neutral traits.

Figure 2: Decomposing genetic variance in a two-population system. The plot illustrates the expected distribution of genetic values in two populations under different selective pressures and the terms on the right list the total (V_g) and between-population genetic variance (V_{gb}) expected over the evolutionary process. For neutrally evolving traits (top row), we expect there to be an absolute difference in the mean genetic values ($|\bar{g_A} - \bar{g_B}|$) that is proportional to F_{ST}. For traits under divergent selection (middle), $|\bar{g_A} - \bar{g_B}|$ is expected to be greater than that expected under genetic drift. For traits under stabilizing selection, $|\bar{g_A} - \bar{g_B}|$ will be less than that expected under genetic drift, and zero in the extreme case.

For traits under selection, the LD contribution is expected to be greater or less than zero, depending

on the type of selection. Under divergent selection, trait-increasing alleles will be systematically more

frequent in one population over the other, inducing positive LD across loci [28, 29], increasing the

LD contribution, i.e., term (1.4). Stabilizing selection, on the other hand, induces negative LD [30,

31]. In the extreme case, the mean genetic values of the two populations are exactly equal and $V_{gb} =$

$(1.2) + (1.3) + (1.4) = 0$. For this to be true, (1.4) has to be negative and equal to (1.2) + (1.3), which

are both positive, and the total genetic variance is reduced to the within-population variance, i.e., term

(1.1) (Fig. 2). This is relevant because, as we show in the following sections, the behavior of the genetic

variance in admixed populations depends on the magnitude of V_{gb} between the source populations.

We illustrate this by tracking the genetic variance in admixed populations for two traits, both with

the same mean F_{ST} at causal loci but with different LD contributions (term 1.4): one where the LD

145 contribution is positive (Trait 1) and the other where it is negative (Trait 2). Thus, traits 1 and 2

146 can be thought of as examples of phenotypes under divergent and stabilizing selection, respectively, and

147 we refer to them as such from hereon. To simulate the genetic variance of such traits, we drew the

148 allele frequencies (f^A and f^B) in populations A and B for 1,000 causal loci with $F_{ST} \sim 0.2$ using the

149 Balding-Nichols model [32]. We drew their effects (β) from $\mathcal{N}(0, \frac{1}{2m\bar{f}(1-\bar{f})})$ where $\bar{f}$ is the mean allele

150 frequency between the two populations, m is the number of loci. To simulate positive and negative

151 LD, we permuted the effect signs across variants 100 times and selected the combinations that gave the

152 most positive and negative LD contribution to represent the genetic architecture of traits that might

153 be under directional (Trait 1) and stabilizing (Trait 2) selection, respectively (Methods). We simulated

154 the genotypes of 10,000 individuals under the HI and CGF models for $t \in \{10, 20, 50, 100\}$ generations

155 post-admixture and calculated genetic values for both traits using $g = \sum_{i=1}^{m} \beta_i x_i$, where $m = 1,000$

156 (Method). The observed genetic variance at any time can then be calculated simply as the variance in

157 genetic values, i.e. $V_g = \mathbb{V}(g)$.

158 In the HI model, $\mathbb{E}(\theta)$ does not change (Fig. 1B) so terms (1.1) and (1.2) are constant through time.

159 Terms (1.3) and (1.4) decay towards zero as the variance in ancestry goes to zero and V_g ultimately

160 converges to (1.1) + (1.2) (Fig. 3). This equilibrium value is equal to the $\mathbb{E}(V_g|\theta)$ (Appendix) and the

161 rate of convergence depends on the strength of assortative mating, which slows the rate at which $\mathbb{V}(\theta)$

162 decays. V_g approaches equilibrium from a higher value for traits under divergent selection and lower value

163 for traits under stabilizing selection because of positive and negative LD contributions, respectively, at

164 $t = 0$ (Fig. 3). In the CGF model, V_g increases initially for both traits with increasing gene flow (Fig.

165 3). This might seem counter-intuitive at first because gene flow increases admixture LD, which leads

166 to more negative values of the LD contribution for traits under stabilizing selection (Fig. S1). But this

167 is outweighed by positive contributions from the genic variance – terms (1.1) + (1.2) + (1.3) – all of

168 which initially increase with gene flow (Fig. S1). After a certain point, the increase in V_g slows down as

169 any increase in $\mathbb{V}(\theta)$ due to gene flow is counterbalanced by recombination and independent assortment.

170 Ultimately, V_g will decrease if there is no more gene flow, reaching the same equilibrium value as in the

171 HI model, i.e., $\mathbb{E}(V_g|\theta) = (1.1) + (1.2)$. Because the loci are unlinked, we refer to the sum (1.3) + (1.4)

172 as the contribution of population structure.

GREML estimation

174 In their original paper, Yang *et al.* (2010) defined h_{snp}^2 as the variance explained by genotyped SNPs and

175 not as heritability [3]. This is because h^2 is the genetic variance explained by causal variants, which are

176 unknown. Genotyped SNPs may not overlap with or tag all causal variants and thus, h_{snp}^2 is understood

177 to be a lower bound of h^2, both being equal if causal variants are known [3]. Our goal is to demonstrate

178 that this may not be true in structured populations and quantify the bias in $\hat{h}_{snp}^2$, even in the ideal

179 situation when causal variants are known.

180 We used GREML, implemented in GCTA [3, 5], to estimate the genetic variance for our simulated

181 traits. GCTA assumes the following model: $\boldsymbol{y} = \boldsymbol{Zu} + \boldsymbol{\epsilon}$ where $\boldsymbol{Z}$ is an $n \times m$ standardized genotype

182 matrix such that the genotype of the k^{th} individual at the i^{th} locus is $z_{ik} = \frac{x_{ik} - 2f_i}{\sqrt{2f_i(1-f_i)}}$, f_i being the

Figure 3: Genetic variance in admixed populations under the (A) HI and (B) CGF models. Dotted lines represent the expected genetic variance based on Eq. (1) and solid lines represent results of simulations averaged over ten replicates. Red and blue lines represent traits under divergent and stabilizing selection, respectively. $P = 0.6$ is missing for simulations run for 50 and 100 generations and $\theta \in \{0.1, 0.2\}$ due to the difficulty in finding mate pairs (Methods)

183 allele frequency. The SNP effects corresponding to the scaled genotypes are assumed to be random and
184 independent such that $\boldsymbol{u} \sim \mathcal{N}(0, \boldsymbol{I}\frac{\sigma_u^2}{m})$ and $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \boldsymbol{I}\sigma_\epsilon^2)$ is random environmental error. Then, the
185 phenotypic variance can be decomposed as:

$$\mathbb{V}(\boldsymbol{y}) = \mathbb{V}(\boldsymbol{Z}\boldsymbol{u}) + \mathbb{V}(e)$$
$$= \frac{\boldsymbol{Z}\boldsymbol{Z}'}{m}\sigma_u^2 + \sigma_\epsilon^2$$

186 where $\frac{\boldsymbol{Z}\boldsymbol{Z}'}{m}$ is the genetic relationship matrix (GRM), the variance components σ_u^2 and σ_ϵ^2 are estimated
187 using restricted maximum likelihood, and $\hat{h}_{snp}^2$ is calculated as $\frac{\hat{\sigma}_u^2}{\hat{\sigma}_u^2 + \hat{\sigma}_\epsilon^2}$. We are interested in asking
188 whether $\hat{\sigma}_u^2$ is an unbiased estimate of V_g. To answer this, we constructed the GRM with causal variants
189 and estimated $\hat{\sigma}_u^2$ using GCTA [3, 4].

190   GCTA under- and over-estimates the genetic variance in admixed populations for traits under diver-
191 gent (Trait 1) and stabilizing selection (Trait 2), respectively, when there is population structure, i.e.,
192 when $\mathbb{V}(\theta) > 0$ (Fig. 4A). One reason for this bias is that the GREML model assumes that the effects
193 are independent, and therefore the LD contribution is zero. This, as discussed in the previous section, is
194 not true for traits under divergent or stabilizing selection between the source populations, and only true
195 for neutral traits in expectation. Because of this, $\hat{\sigma}_u^2$ does not capture the LD contribution, i.e. term
196 (1.4) (Fig. 4A). But $\hat{\sigma}_u^2$ can be biased even if the LD contribution is zero if the genotypes are scaled with
197 $\sqrt{2f_i(1 - f_i)}$ – the standard practice – where f_i is the frequency of the allele in the population. This
198 scaling assumes that $\mathbb{V}(x_i) = 2f_i(1 - f_i)$, which is true only if the population were mating randomly.
199 In an admixed population $\mathbb{V}(x_i) = 2f_i(1 - f_i) + 2\mathbb{V}(\theta)(f_i^A - f_i^B)^2$, where f_i, f_i^A, and f_i^B correspond
200 to frequency in the admixed population, and source populations, A and B, respectively (Appendix).
201 Alternatively, if the genotypes are scaled, $\mathbb{V}(z_i) = 1 + 2\mathbb{V}(\theta)F_{st}^{(i)}$ where $F_{st}^{(i)}$ is the F_{st} at the i^{th} locus.
202 We show that this assumption biases $\hat{\sigma}_u^2$ downwards by a factor of $2\mathbb{V}(\theta)(f_i^A - f_i^B)^2$ (or $2\mathbb{V}(\theta)F_{st}^{(i)}$ if
203 genotypes are scaled) – term (1.3) (Fig. 4B, Appendix). Thus, with the standard scaling, $\hat{\sigma}_u^2$ gives a
204 biased estimate in the presence of population structure, even of the genic variance.

205   The overall bias in $\hat{\sigma}_u^2$ is determined by the relative magnitude and direction of terms (1.3) and
206 (1.4), both of which are functions of $\mathbb{V}(\theta)$, and therefore, of the degree of structure in the population.
207 The contribution of term (1.3) will be modest, even in highly structured populations (Fig. S1) and
208 therefore, the overall bias is largely driven by the LD contribution. If there is no more gene flow, $\mathbb{V}(\theta)$
209 will ultimately go to zero and V_g will converge towards $\hat{\sigma}_u^2$. Thus, $\hat{\sigma}_u^2$ is more accurately interpreted
210 as the genetic variance expected if the LD contribution were zero and if the population were mating
211 randomly. In other words, $\mathbb{E}(\hat{\sigma}_u^2) = (1.1) + (1.2) \neq V_g$ (Fig. 4B).

212   In principle, we can recover the missing components of V_g by scaling the genotypes appropriately.
213 For example, we can recover term (1.3) by scaling the genotype at each variant i by its sample variance,
214 i.e., $z_{ik} = \frac{x_{ik} - 2f_i}{\sqrt{\mathbb{V}(x_i)}}$ (Fig. 4C) (Appendix). We can also recover term (1.4) by scaling the genotypes with
215 the covariance between SNPs, i.e., the LD matrix, as previously proposed [33, 34] (Methods). In matrix
216 form, the 'LD-scaled' genotypes can be written as $\boldsymbol{Z} = (\boldsymbol{X} - 2\boldsymbol{P})\boldsymbol{U}^{-1}$ where $\boldsymbol{P}$ is an $n \times m$ matrix such
217 that all elements of the i^{th} column contain the frequency of the i^{th} SNP and $\boldsymbol{U}$ is the (upper triangular)
218 square root matrix of the LD matrix, i.e., $\boldsymbol{\Sigma} = \boldsymbol{U}'\boldsymbol{U}$ [33]. GREML recovers the LD contribution under

Figure 4: The behavior of GREML estimates of the genetic variance ($\hat{\sigma}_u^2$) in admixed populations under the HI (left column) and CGF (right column) models either without (A-D) or with (E-H) individual ancestry as a fixed effect. The solid lines represent estimates from simulated data averaged across ten replicates with red and blue colors representing estimates for traits under divergent and stabilizing selection, respectively. P indicates the strength of assortative mating. The shaded area represents the 95% confidence bands generated by bootstrapping (sampling with replacement 100 times) the point estimate reported by GCTA. The dotted lines either represent the expected variance in the population based on Eq. 1 (A & B) or the expected estimate for three different ways of scaling genotypes (B-D & F-H). (A-B & E-F) show the behavior of $\hat{\sigma}_u^2$ for the default scaling, (C, G) shows $\hat{\sigma}_u^2$ when the genotype at a locus is scaled by its sample variance ($\mathbb{V}(x)$ scaled), and (D, H) when it is scaled by the sample covariance (LD scaled).

219 this scaling, resulting in unbiased estimates of V_g for both traits (Fig. 4D, Appendix).

220 In practice, however, the LD contribution may not be fully recoverable for two reasons. One, the

221 LD-scaled GRM requires computing the inverse of $\mathbf{\Sigma}$ or $\mathbf{U}$ which may not exist, especially if the number

222 of markers is greater than the sample size – the case for most human genetic studies. Second, it is

223 common to include individual ancestry or principal components of the GRM as fixed effects in the model

224 to account for inflation in heritability estimates due to shared environment. This should also have the

225 effect of removing the components of genetic variance along the ancestry axes, the residual variance being

226 equal to $\mathbb{E}\{\mathbb{V}(g|\theta)\} = (1.1) + (1.2) - (1.3)$ (Appendix). Indeed, this is what we observe in Fig. 4H. Thus,

227 if ancestry is included as a fixed effect, we expect V_g to be underestimated in the presence of population

228 structure, regardless of genetic architecture.

HE estimation

230 Haseman-Elston (HE) regression also assumes a random-effects model but uses a method-of-moments

231 approach, as opposed to GREML, which maximizes the likelihood to estimate V_g. Previous work has

232 shown that as long as all causal variants are included in the GRM calculation, the HE estimator will

233 not be biased, even if they are in LD with each other [35]. We show that in the presence of positive

234 and negative LD between causal loci, as exemplified by traits under divergent and stabilizing selection,

235 respectively, the HE estimates of V_g are biased upwards and downwards, respectively (Fig. 5A-B). To

236 understand this discrepancy and the source of bias in our simulations, recall that HE estimates V_g from

237 the regression of the (pairwise) phenotypic covariance between individuals on their genotypic covariance

238 [24]. More specifically, if we denote $Y_{kl} = y_k y_l$ as the product of the (centered) phenotypes of k^{th} and

239 l^{th} individuals, and ψ_{kl} as the k^{th} and l^{th} entry of the GRM, then the HE estimator can be written as:

$$
\begin{aligned}
\hat{V}_g &= \frac{Cov(Y_{kl}, \psi_{kl})}{Var(\psi_{kl})} \\[6pt]
&= \frac{\mathbb{E}(y_k y_l \sum_{w=1}^{M} z_{wk} z_{wl})}{\mathbb{E}(\sum_{i=1}^{M} z_{ik} z_{il} \sum_{w=1}^{M} z_{wk} z_{wl})} \\[6pt]
&= \frac{\mathbb{E}\{(g_k + e_k)(g_l + e_l) \sum_{w=1}^{M} z_{wk} z_{wl}\}}{\mathbb{E}(\sum_{i=1}^{M} z_{ik} z_{il} \sum_{w=1}^{M} z_{wk} z_{wl})} \\[6pt]
&= \frac{\mathbb{E}(g_k g_l \sum_{w=1}^{M} z_{wk} z_{wl})}{\mathbb{E}(\sum_{i=1}^{M} z_{ik} z_{il} \sum_{j=1}^{M} z_{wk} z_{wl})} \\[6pt]
&= \frac{\mathbb{E}(\sum_{i=1}^{M} \sum_{j=1}^{M} u_i u_j z_{ik} z_{jl} \sum_{w=1}^{M} z_{wk} z_{wl})}{\mathbb{E}(\sum_{i=1}^{M} z_{ik} z_{il} \sum_{w=1}^{M} z_{wk} z_{wl})} \\[6pt]
&= \frac{\mathbb{E}(\sum_{i=1}^{M} \sum_{j=1}^{M} u_i u_j \sum_{w=1}^{M} z_{ik} z_{jl} z_{wk} z_{wl})}{\mathbb{E}(\sum_{i=1}^{M} \sum_{w=1}^{M} z_{ik} z_{il} z_{wk} z_{wl})} \\[6pt]
&= \frac{\mathbb{E}(\sum_{i=1}^{M} u_i^2 \sum_{w=1}^{M} z_{ik} z_{jl} z_{wk} z_{wl})}{\mathbb{E}(\sum_{i=1}^{M} \sum_{w=1}^{M} z_{ik} z_{il} z_{wk} z_{wl})} + \frac{\mathbb{E}(\sum_{i=1}^{M} \sum_{j \neq i} u_i u_j \sum_{w=1}^{M} z_{ik} z_{jl} z_{wk} z_{wl})}{\mathbb{E}(\sum_{i=1}^{M} \sum_{w=1}^{M} z_{ik} z_{il} z_{wk} z_{wl})}
\end{aligned}
\tag{2}
$$

240 Where the first and second terms represent the genic and LD components, respectively, of the esti-

241 mate. Population structure induces correlations between the alleles at a given locus as well as across

242 loci (i.e., LD). But the LD may not be directional, i.e., trait-increasing alleles may be as likely to be

243 co-inherited with each other as they are to trait-decreasing alleles, and vice versa – implicit under the

244 standard random-effects model. Thus, in the absence of directional LD, the second term is zero and the
245 first term is unaffected as long as all causal variants are included in the GRM, because the increase in
246 the numerator due to population structure is proportional to the denominator [35]. Directional LD does
247 not affect the first term but exaggerates the contribution from the second term, i.e., the LD component
248 (see Appendix section A3.2). Consequently, HE regression over- and under-estimates V_g for traits with
249 positive and negative LD, respectively. Note that this bias is in the opposite direction of the bias observed
250 with GREML, which fails to capture the LD contribution. Scaling the genotype at a locus by its LD with
251 other loci, as discussed in the previous section, corrects for the bias in HE regression regardless of genetic
252 architecture, yielding estimates consistent with GREML (Fig. 5C). Thus, GREML and HE regression
253 are guaranteed to yield the same estimates only if the underlying model specifying the distribution of
254 effects is consistent with the true architecture of the trait.

255   The practice of including individual ancestry as a covariate in HE regression to account for shared
256 environment [11] reduces the bias from exaggerated LD contributions (Fig. 5D-F). But, as with GREML,
257 this also removes any genetic variance that may exist along the ancestry axis, yielding underestimates
258 of V_g, regardless of genetic architecture.

259 Local ancestry heritability

260 A related quantity of interest in admixed populations is local ancestry heritability (h_γ^2), which is defined
261 as the proportion of phenotypic variance that can be explained by local ancestry. Zaitlen *et al.* (2014)
262 [36] showed that this quantity is related to, and can be used to estimate, h^2 in admixed populations.
263 The advantage of this approach is that local ancestry segments shared between individuals are identical
264 by descent and are therefore, more likely to tag causal variants compared to array markers, allowing
265 one to potentially capture the contributions of rare variants [36]. Here, we show that in the presence of
266 population structure, (i) the relationship between h_γ^2 and h^2 is not straightforward and (ii) $\hat{h}_\gamma^2$ may be a
267 biased estimate of local ancestry heritability under the random effects model for the same reasons that
268 $\hat{h}_{snp}^2$ is biased.

269   We define local ancestry $\gamma_i \in \{0, 1, 2\}$ as the number of alleles at locus i that trace their ancestry
270 to population A. Thus, ancestry at the i^{th} locus in individual k is a binomial random variable with
271 $\mathbb{E}(\gamma_{ik}) = 2\theta_k$, θ_k being the ancestry of the k^{th} individual. Similar to genetic value, the 'ancestry value'
272 of an individual can be defined as $\sum_{i=1}^m \phi_i \gamma_i$, where $\phi_i = \beta_i(f_i^A - f_i^B)$ is the effect size of local ancestry
273 (Appendix). Then, the genetic variance due to local ancestry can be expressed as (Appendix):

Figure 5: Genetic variance ($\hat{V}_g$) estimated with HE regression in admixed populations under the HI (left column) and CGF (right column) models either without (A-C) or with (D-F) adjustment for individual ancestry. The solid lines represent estimates from simulated data averaged across ten replicates with red and blue colors representing estimates for traits under divergent and stabilizing selection, respectively. P indicates the strength of assortative mating. (A & D) show the behavior of $\hat{V}_g$ for the default scaling, (B, E) shows $\hat{V}_g$ when the genotype at a locus is scaled by its sample variance ($\mathbb{V}(x)$ scaled), and (C, F) when it is scaled by the sample covariance (LD scaled). The dotted lines in A-E represent the expected V_g in the population based on Eq. 1 and in F, represent the expected V_g after removing any genetic variance along the ancestry axis. The shaded areas represent the 95% bootstrapped confidence bands of the estimate.

$$V_\gamma = \mathbb{V}\left(\sum_{i=1}^m \phi_i \gamma_i\right) = \sum_{i=1}^m \phi_i^2\, \mathbb{V}(\gamma_i) + \sum_{i=1}^m \sum_{j\neq i} \phi_i \phi_j\, \mathbb{C}(\gamma_i, \gamma_j)$$

$$= 2\,\mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\}\sum_{i=1}^m \phi_i^2 + 2\,\mathbb{V}(\theta)\sum_{i=1}^m \phi_i^2 + 4\,\mathbb{V}(\theta)\sum_{i=1}^m \sum_{j\neq i} \phi_i\phi_j$$

$$= 2\,\mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\}\sum_{i=1}^m \beta_i^2(f_{\mathrm{i}}^{\mathrm{A}} - f_{\mathrm{i}}^{\mathrm{B}})^2$$

$$+ 2\,\mathbb{V}(\theta)\sum_{i=1}^m \beta_i^2(f_{\mathrm{i}}^{\mathrm{A}} - f_{\mathrm{i}}^{\mathrm{B}})^2$$

$$+ 4\,\mathbb{V}(\theta)\sum_{i=1}^m \sum_{j\neq i} \beta_i\beta_j(f_{\mathrm{i}}^{\mathrm{A}} - f_{\mathrm{i}}^{\mathrm{B}})(f_{\mathrm{j}}^{\mathrm{A}} - f_{\mathrm{j}}^{\mathrm{B}})$$

and heritability explained by local ancestry is simply the ratio of V_γ and the phenotypic variance. Note
that $V_\gamma = (1.2) + (1.3) + (1.4)$ and therefore its behavior is similar to V_g in that the terms (1.3) and (1.4)
decay towards zero as $\mathbb{V}(\theta) \to 0$, and V_γ converges to (1.2) (Fig. S2). Additionally, the dependence of
V_γ on both $\mathbb{E}(\theta)$ and $\mathbb{V}(\theta)$ precludes a straightforward derivation between local ancestry heritability and
h^2.

GREML estimation of $\hat{h}_\gamma^2$ is similar to that of $\hat{h}_{snp}^2$, the key difference being that the former involves
constructing the GRM using local ancestry instead of genotypes [36]. The following model is assumed:
$\boldsymbol{y} = \boldsymbol{W}\boldsymbol{v} + \boldsymbol{\xi}$ where $\boldsymbol{W}$ is an $n \times m$ standardized local ancestry matrix, $\boldsymbol{v} \sim \mathcal{N}(0, \boldsymbol{I}\frac{\sigma_v^2}{m})$ are local
ancestry effects, and $\boldsymbol{\xi} \sim \mathcal{N}(0, \boldsymbol{I}\sigma_\xi^2)$. Note that σ_ξ^2 captures both environmental noise as well as any
genetic variance independent of local ancestry. The phenotypic variance is decomposed as $\mathbb{V}(\boldsymbol{y}) =$
$\mathbb{V}(\boldsymbol{W}\boldsymbol{v}) + \mathbb{V}(\boldsymbol{\xi}) = \frac{\boldsymbol{W}\boldsymbol{W}'}{m}\sigma_v^2 + \sigma_\xi^2$ where $\frac{\boldsymbol{W}\boldsymbol{W}'}{m}$ is the local ancestry GRM and σ_v^2 is the parameter of
interest, which is believed to be equal to V_γ – the genetic variance due to local ancestry.

We show that, in the presence of population structure, i.e., when $\mathbb{V}(\theta) > 0$, GREML $\hat{\sigma}_v^2$ is biased
downwards relative to V_γ for traits under divergent selection and upwards for traits under stabilizing
selection because it does not capture the contribution of LD (Fig. 6A). But there is another source of
bias in $\hat{\sigma}_v^2$, which tends to be inflated in the presence of population structure if individual ancestry is
not included as a covariate, even with respect to the expectation of V_γ under equilibrium (seen more
clearly in Fig. 6B-C). We suspect this inflation is because of strong correlations between local ancestry
– local ancestry disequilibrium – across loci that inflates $\hat{\sigma}_v^2$ in a way that is not adequately corrected
even when all causal variants are included in the model [4, 10]. Scaling local ancestry by its covariance
removes this bias and recovers the contribution of LD (Fig. 6D) presumably because this accounts for
the correlation in genotypes across loci. Including individual ancestry as a fixed effect also corrects for
the inflation in $\hat{\sigma}_v^2$ (Fig. 6E-H). But as with $\hat{\sigma}_u^2$, this practice will underestimate the genetic variance
due to local ancestry in the presence of population structure because it removes the variance along the
ancestry axis (Fig. 6E-H).

Based on the above, GREML $\hat{h}_\gamma^2$ and corresponding estimates of h^2 are more accurately interpreted as
the heritability due to local ancestry and heritability, respectively, expected in the absence of population
structure. We believe $\hat{h}_\gamma^2$ is still useful in that, because it should capture the effects of rare variants, it
can be used to estimate the upper bound of $\hat{h}_{snp}^2$.

In a previous paper, we suggested that local ancestry heritability could potentially be used to estimate the genetic variance between populations [37]. Our results suggest this is not possible for two reasons. First, the GREML estimator of local ancestry heritability, as we show in this section is biased and does not capture the LD contribution. But even if we were able to recover the LD component, our decomposition shows that local ancestry is equal to the genetic variance between populations (V_{gb}) only when $\mathbb{E}(\theta) = 0.5$ and $\mathbb{V}(\theta) = \mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} = 0.25$, which is only possible at $t = 0$ in the HI model. After admixture, $\mathbb{V}(\theta)$ decays and the equivalence between V_γ and V_{gb} is lost, making it impossible to estimate the latter from admixed populations, especially for traits under divergent or stabilizing selection, even if the environment is randomly distributed with respect to ancestry. We note that this conclusion was recently reached independently by Schraiber and Edge (2023) [38].

Figure 6: The behavior of GREML estimates of the variance due to local ancestry ($\hat{\sigma}_v^2$) in admixed populations under the HI (left column) and CGF (right column) models either without (A-D) or with (E-H) individual ancestry included as a fixed effect. The solid lines represent estimates from simulated data averaged across ten replicates with red and blue colors representing estimates for traits under divergent and stabilizing selection, respectively. P indicates the strength of assortative mating. The dotted lines either represent the expected variance in the population (A & B) or the expected estimate for three different ways of scaling local ancestry (B-D & F-H). (A-B & E-F) show the behavior of $\hat{\sigma}_v^2$ for the default scaling, (C, G) shows $\hat{\sigma}_v^2$ when local ancestry is scaled by the sample variance, and (D, H) when it is scaled by the sample covariance. Shaded regions represent the 95% confidence bands. Some runs in (D & H) failed to converge as seen by the missing segments of the solid lines because the expected variance in such cases was too small.

How much does LD contribute to V_g in practice?

In the previous sections, we showed theoretically that $\hat{h}^2_{snp}$ may be biased in admixed populations even if the causal variants are known and in the absence of confounding by shared environment. GREML fails to capture the LD contribution whereas HE regression overestimates it. The extent to which $\hat{h}^2_{snp}$ is biased because of this reason in practice is ultimately an empirical question, which is difficult to answer because the true genetic architecture – the LD contribution in particular – is unknown. In this section, we develop some intuition for this contribution among individuals with mixed African and European ancestry using a combination of simulations and empirical data analysis.

First, we simulated a neutral trait using genotype data from the African Americans (ASW) from the 1,000 Genomes Project (1KGP) [39]. To do this, we sampled $m \in \{10, 100, 1,000\}$ causal loci from a set of common (MAF > 0.01), LD pruned variants and assigned them effects such that $\beta_i \sim \mathcal{N}\left(0, \frac{1}{\sqrt{m\,\mathbb{V}(x_i)}}\right)$, i.e., the expected *genic* variance is $\mathbb{E}\{\sum_{i=1}^{m} \beta_i^2 Var(x_i)\} = 1$ (Methods). We computed the genic and LD contributions and repeated this process 1,000 times where each replicate can be thought of as an independent realization of the genetic architecture of a neutrally evolving trait. We show that the LD contribution may be zero in expectation but can be substantial for a given trait (up to 50% of the genic variance, Fig. S4), even in the absence of selection.

Second, we estimated the LD contribution of genome-wide significant SNPs for 26 quantitative traits from the GWAS catalog [40]. To do this, we decomposed the variance explained in ASW into the four components in Equation 1 using allele frequencies (f^A and f^B) from the YRI and CEU and the mean ($\mathbb{E}(\theta) \approx 0.77$) and variance ($\mathbb{V}(\theta) \approx 0.02$) of individual ancestry from ASW (Methods). We show that for skin pigmentation – a trait under strong divergent selection – the LD contribution, i.e. term (1.4), is positive and accounts for $\approx 40 - 50\%$ of the total variance explained. This is because of large allele frequency differences between Africans and Europeans that are correlated across skin pigmentation loci, consistent with strong polygenic selection favoring alleles for darker pigmentation in regions with high UV exposure and vice versa [37, 41–44]. But for most other traits, LD contributes relatively little, explaining a modest, but non-negligible proportion of the genetic variance in height, LDL and HDL cholestrol, mean corpuscular hemoglobin (MCH), neutrophil count (NEU), and white blood cell count (WBC) (Fig. 7). Because we selected independent associations for this exercise (Methods), the LD contribution is driven entirely due to population structure in ASW. The contribution of population structure to the genic variance, i.e., term (1.3) is also small even for traits like skin pigmentation and neutrophil count with large effect alleles that are highly diverged in frequency between Africans and Europeans [42, 43, 45–47]. Overall, this suggests that population structure contributes relatively little, as least to the variance explained by GWAS SNPs.

Discussion

Despite the growing size of GWAS and discovery of thousands of variants for hundreds of traits [40], the heritability explained by GWAS SNPs remains a fraction of twin-based heritability estimates. Yang *et al.* (2010) introduced the concept of SNP heritability (h^2_{snp}) that does not depend on the discovery of

Figure 7: Decomposing the genetic variance explained by GWAS SNPs in the 1000 Genomes ASW (African Americans from Southwest). We calculated the four variance components listed in Eq. 1, their values shown on the y-axis as a fraction of the total variance explained (shown as percentage at the bottom). The LD contribution, which can be positive or negative, is shown in yellow. The number of variants used to calculate variance components for each trait is also shown at the bottom.

causal variants but assumes that they are numerous and are more or less uniformly distributed across the genome (the infinitesimal model), their contributions to the genetic variance 'tagged' by genotyped SNPs [3]. h_{snp}^2 is now routinely estimated in most genomic studies and at least for some traits (e.g. height and BMI), these estimates now approach twin-based heritability [6]. But despite the widespread use of $\hat{h}_{snp}^2$, its interpretation remains unclear, particularly in the presence of admixture and population structure. It is generally accepted that $\hat{h}_{snp}^2$ can be biased in structured populations because of confounding effects of unobserved environmental factors and LD between causal variants [4, 7, 9–11, 48]. But $\hat{h}_{snp}^2$ may be biased even in the absence of confounding because of misspecification of the underlying random-effects model, i.e., if the model does not represent the genetic architecture from which the trait is sampled [14–16, 49, 50].

Under the standard GREML model, SNP effects are assumed to be uncorrelated and the total genetic variance can be represented as the sum of the variance explained by individual loci, i.e. the genic variance [14–16]. In admixed populations, there is substantial LD, which can contribute to the genetic variance, and can persist for a number of generations, despite recombination, due to continued gene flow and/or ancestry-based assortative mating. GREML does not capture this LD contribution [12, 15], and therefore, may lead to biased estimates of h_{snp}^2. The LD contribution can be negative for traits under stabilizing selection, and positive for traits under divergent selection between the source populations, leading to over- or under-estimates, respectively. Thus, GREML estimates of h_{snp}^2, assuming genotypes are scaled properly (see below), is better interpreted as the proportion of phenotypic variance explained by the

369 *genic* variance. Estimates of local ancestry heritability ($\hat{h}_\gamma^2$) [36, 51] should be interpreted similarly.

370 We show that with GREML, $\hat{h}_{snp}^2$ can be biased even when the LD contribution is zero if the geno-
371 types are scaled by $\sqrt{2f(1-f)}$ – the standard approach, which implicitly assumes a randomly mating
372 population. In the presence of population structure, the variance in genotypes can be higher and $\hat{h}_{snp}^2$
373 does not capture this additional variance, which we show can be recovered by scaling genotypes by the
374 SNP variance ($\sqrt{Var(x)}$). In principle, the LD contribution can also be recovered by scaling genotypes
375 by the SNP covariance, i.e., the LD matrix, as previously suggested [33, 34]. But this approach is limited
376 to situations where the sample size is much larger than the number of markers.

377 We also investigated the behavior of another widely used approach to estimate h_{snp}^2 – Haseman-
378 Elston regression. We show that $\hat{h}_{snp}^2$ estimated with HE regression is also biased, but for different
379 reasons and in the opposite direction of the bias observed with GREML. HE regression exaggerates
380 the LD contribution, leading to over- and under-estimates of h_{snp}^2 for traits where the causal loci are
381 in positive and negative LD, respectively. Approaches that correct for population structure [35] should
382 remove this source of bias but would also remove any genetic variance in the trait along the ancestry axis,
383 including the LD contribution. This results in underestimates of h_{snp}^2, regardless of trait architecture.

384 One limitation of this paper is that we have focused on random-effects estimators of h_{snp}^2 because of
385 their widespread use. Estimators of h_{snp}^2 can be broadly grouped into random- and fixed effect estimators
386 based on how they treat SNP effects [35]. Fixed effect estimators make fewer distributional assumptions
387 but they are not as widely used because they require conditional estimates of all variants – a high-
388 dimensional problem where the number of markers is often far larger than the sample size [52]. This is
389 one reason why random effect estimators, such as GREML, are popular – because they reduce the number
390 of parameters that need to be estimated by assuming that the effects are drawn from some distribution
391 where the variance is the only parameter of interest. Fixed effects estimators, in principle, should be able
392 to capture the LD contribution but this is not obvious in practice since the simulations used to evaluate
393 the accuracy of such estimators still assume uncorrelated effects [35, 52, 53]. Further research is needed
394 to clarify the interpretation of the different estimators of h_{snp}^2 in structured populations under a range
395 of genetic architectures.

396 Does the LD contribution to the genetic variance have practical implications? The answer to this
397 depends on the context in which SNP heritability is used. $\hat{h}_{snp}^2$ can be useful in quantifying the power
398 to detect variants in GWAS where the quantity of interest is the genic variance. But $\hat{h}_{snp}^2$ can lead to
399 misleading conclusions if used to measure the extent to which genetic variation contributes to phenotypic
400 variation, in predicting the response to selection, or in defining the upper limit of polygenic prediction
401 accuracy [2] – applications where the LD contribution is important.

402 Ultimately, the discrepancy between $\hat{h}_{snp}^2$ and h^2 in practice is an empirical question, the answer to
403 which depends on the degree of population structure (which we can measure) and the genetic architecture
404 of the trait (which we do not know *a priori*). We show that for most traits, the contribution of population
405 structure to the variance explained by GWAS SNPs is modest among African Americans. Thus, if we
406 assume that the genetic architecture of GWAS SNPs represents that of all causal variants, then despite
407 incorrect assumptions, the discrepancy between $\hat{h}_{snp}^2$ and h^2 should be fairly modest. But this assumption
408 is unrealistic given that GWAS SNPs are common variants that in most cases cumulatively explain a

409 fraction of trait heritability. What is the LD contribution of the rest of the genome, particularly rare
410 variants? This is not obvious and will become clearer in the near future through large sequence-based
411 studies [54]. While these are underway, theoretical studies are needed to understand how different
412 selection regimes influence the directional LD between causal variants – clearly an important aspect of
413 the genetic architecture of complex traits.

Methods

Simulating genetic architecture

We first drew the allele frequency (f^0) of 1,000 biallelic causal loci in the ancestor of populations A and
B from a uniform distribution, $U(0.001, 0.999)$. Then, we simulated their frequency in populations A and
B (f^A and f^B) under the Balding-Nichols model [32], such that $f^A, f^B \sim Beta(\frac{f^0(1-F)}{F}, \frac{(1-f^0)(1-F)}{F})$
where $F = 0.2$ is the inbreeding coefficient. We implemented this using code adapted from Lin *et al.*
(2021) [55]. To avoid drawing extremely rare alleles, we continued to draw f^A and f^B until we had 1,000
loci with $f^A, f^B \in (0.01, 0.99)$.

We generated the effect size (β) of each locus by sampling from $\mathcal{N}(0, \frac{1}{2m\bar{f}(1-\bar{f})})$, where m is the
number of loci and $\bar{f}$ is the mean allele frequency across populations A and B. Thus, rare variants have
larger effects than common variants and the total genetic variance sums to 1. Given these effects, we
simulated two different traits, one with a large difference in means between populations A and B (Trait
1) and the other with roughly no difference (Trait 2). This was achieved by permuting the signs of the
effects 100 times to get a distribution of V_{gb} – the genetic variance between populations. This has the
effect of varying the LD contribution without changing the F_{ST} at causal loci. We selected the maximum
and minimum of V_{gb} to represent Traits 1 and 2.

Simulating admixture

We simulated the genotypes, local ancestry, and phenotype for 10,000 admixed individuals per generation
under the hybrid isolation (HI) and continuous gene flow (CGF) models by adapting the code from Zaitlen
et al. (2017) [26]. We denote the ancestry of a randomly selected individual k with θ, the fraction of their
genome from population A. At $t = 0$ under the HI model, we set θ to 1 for individuals from population A
and 0 if they were from population B such that $\mathbb{E}(\theta) = p \in \{0.1, 0.2, 0.5\}$ with no further gene flow from
either source population. In the CGF model, population B receives a constant amount q from population
A in every generation starting at $t = 0$. The mean overall proportion of ancestry in the population is
kept the same as the HI model by setting $q = 1 - (1-p)^{\frac{1}{t}}$ where t is the number of generations of gene
flow from A. In every generation, we simulated ancestry-based assortative mating by selecting mates
such that the correlation between their ancestries is $P \in \{0, 0.3, 0.6, 0.9\}$ in every generation. We do this
by repeatedly permuting individuals with respect to each other until P falls within ± 0.01 of the desired
value. It becomes difficult to meet this criterion when $\mathbb{V}(\theta)$ is small (Fig.1C). To overcome this, we
relaxed the threshold up to 0.04 for some conditions, i.e., when $\theta \in \{0.1, 0.2\}$ and $t \geq 50$. We generated
expected variance in individual ancestry using the expression in Zaitlen *et al.* (2017) [26]. At time t

since admixture, $\mathbb{V}(\theta_t) = \mathbb{V}(\theta_{t-1})\frac{(1+P)}{2}$ under the HI model where P measures the strength of assortative

mating, i.e, the correlation between the ancestry between individuals in a mating pair. Under the CGF

model, $\mathbb{V}(\theta_t) = q(1-q)\,\mathbb{E}(\theta_{t-1})^2 + q(1-q)\{1 - 2\,\mathbb{E}(\theta_{t-1})\} + (1-q)\,\mathbb{V}(\theta_{t-1})\frac{(1+P)}{2}$ (Appendix).

We sampled the local ancestry at each i^{th} locus as $\gamma_i = \gamma_{if} + \gamma_{im}$ where $\gamma_{im} \sim Bin(1, \theta_m), \gamma_{if} \sim$

$Bin(1, \theta_f)$ and θ_m and θ_f represent the ancestry of the maternal and paternal chromosome, respectively.

The global ancestry of the individual is then calculated as $\theta_k = \sum_{i=1}^{m} \frac{\gamma_{im} + \gamma_{if}}{2m}$, where m is the number of

loci. We sample the genotypes x_{im} and x_{if} from a binomial distribution conditioning on local ancestry.

For example, the genotype on the maternal chromosome is $x_{im} \sim Bin(1, f_i^{A})$ if $\gamma_{im} = 1$ and $x_{im} \sim$

$Bin(1, f_i^{B})$ if $\gamma_{im} = 0$ where f_i^{A} and f_i^{B} represent the allele frequency in populations A and B, respectively.

Then, the genotype can be obtained as the sum of the maternal and paternal genotypes: $x_i = x_{im} + x_{ip}$.

We calculate the genetic value of each individual as $g = \sum_{i=1}^{m} \beta_i x_i$ and the genetic variance as $\mathbb{V}(g)$.

456 Heritability estimation with GREML

We used the *--reml* and *--reml-no-constrain* flags in GCTA [5] to estimate σ_u^2 and σ_v^2, the genetic variance

due to genotypes and local ancestry, respectively. We could not run GCTA without noise in the genetic

values so we simulated individual phenotypes with a heritability of $h^2 = 0.8$ by adding random noise

$e \sim \mathcal{N}(0, V_g\frac{1-h^2}{h^2})$. We computed three different GRMs, which correspond to different transformations

of the genotypes: (i) standard, (i) Variance or $V(x)$ scaled, and (ii) LD-scaled.

For the standard GRM, the genotypes at the i^{th} SNP are standardized such that $z_i = \frac{x_i - 2f_i}{\sqrt{2f_i(1-f_i)}}$.

For the variance scaled GRM, we computed $z_i = \frac{x_i - 2f_i}{\sqrt{\mathbb{V}(x_i)}}$ where $\mathbb{V}(x_i)$ is the sample variance of the

genotypes at the i^{th} SNP. The LD-scaled GRM conceptually corresponds to standardizing the genotypes

by the SNP covariance, rather than its variance. Let $\boldsymbol{X}$ represent the $n \times m$ *unstandardized* matrix of

genotypes and $\boldsymbol{P}$ represent an $n \times m$ matrix where the i^{th} column contains the allele frequency of that

SNP. Let $\boldsymbol{U}$ be the upper triangular 'square root' matrix of the $m \times m$ SNP covariance matrix $\boldsymbol{\Sigma}$ such

that $\boldsymbol{\Sigma} = \boldsymbol{U}'\boldsymbol{U}$. Then, the standardized genotypes are computed as $\boldsymbol{Z} = (\boldsymbol{X} - 2\boldsymbol{P})\boldsymbol{U}^{-1}$ and the GRM

becomes $(\boldsymbol{X} - 2\boldsymbol{P})\boldsymbol{\Sigma}^{-1}(\boldsymbol{X} - 2\boldsymbol{P})'$ [33]. Similarly, the three GRMs for local ancestry were computed by

scaling local ancestry with (i) $\sqrt{2\bar{\gamma}_i(1 - \bar{\gamma}_i)}$ where we denote $\bar{\gamma}_i$ as the mean local ancestry at the i^{th}

SNP, or with the (ii) variance, or (iii) covariance of local ancestry, respectively. We estimated σ_u^2 and

σ_v^2 with and without individual ancestry as a fixed effect to correct for any confounding due to genetic

stratification. This was done by using the *--qcovar* flag.

474 Heritability estimation with HE regression

475   Haseman-Elston regression with and without ancestry correction was implemented using custom scripts

476   in R [56]. To estimate V_g without ancestry correction, we first computed the cross-product of the centered

477   phenotypes ($\boldsymbol{y}$), resulting in an $n \times n$ matrix $\boldsymbol{y}\boldsymbol{y}'$. We stacked the upper-triangular matrix of $\boldsymbol{y}\boldsymbol{y}'$ into a

478   vector and regressed it on the corresponding elements of the GRM ($\boldsymbol{\psi}$), taking the slope as an estimate

479   of V_g:

$$\hat{V}_g = \frac{\sum_{k=1} \sum_{l<k} y_k y_l \psi_{kl}}{\sum_k \sum_{l<k} \psi_{kl}^2}$$

480   To correct for individual ancestry, we followed the approach of Min et al. (2022) [35]. To do this, we
481   first regressed out the effect of individual ancestry ($\boldsymbol{\theta}$) on phenotype. The regression coefficient can be
482   expressed as $\boldsymbol{\theta}(\boldsymbol{\theta}'\boldsymbol{\theta})^{-1}\boldsymbol{\theta}'$ and the residuals as $\boldsymbol{y}* = (\boldsymbol{I} - \boldsymbol{\theta}(\boldsymbol{\theta}'\boldsymbol{\theta})^{-1}\boldsymbol{\theta}')\boldsymbol{y}$. Then, we fit the following model:

$$\mathbb{E}(\boldsymbol{y} * \boldsymbol{y}*') = V_g \boldsymbol{\psi} + V_e \boldsymbol{I} + \delta \boldsymbol{\theta}\boldsymbol{\theta}'$$

where $\boldsymbol{\theta}\boldsymbol{\theta}'$ represents the cross-product of individual ancestry, δ represents its corresponding regression
coefficient, and V_g represents the parameter of interest, i.e., the genetic variance and V_e, the residual
variance.

To demonstrate that the bias in HE estimates arises because of a bias in the estimate of LD contri-
bution, not the genic variance, we carried out a simple simulation where half of the individuals in the
population derive their ancestry from population A and the rest from population B. This is equivalent to
the meta-population under the HI model at $t = 0$ where $\mathbb{E}(\theta) = 0.5$. We simulated genotypes for $1,000$
individuals for $m = 100$ loci where the allele frequencies in populations A and B were set to $f_A = 0.1$
and $f_B = 0.8$, respectively. We standardized the genotypes at each locus i using the square-root of the
sample variance and assigned effect sizes such that the total genetic variance explained by all loci is equal
to 1, i.e., the effect of the scaled genotype at the i^{th} locus is $u_i = \frac{1}{\sqrt{m}}$. This is equivalent to the effect
size of the unscaled genotypes being $\beta_i = \frac{1}{\sqrt{m \mathbb{V}(x_i)}}$ where $\mathbb{V}(x_i)$ is the sample variance at the i^{th} locus.
We introduced randomness in the direction of the effect by assigning a negative or positive sign to each
locus uniformly at random 100 times to generate 100 traits with the same genic variance but varying LD
contributions. Then, for each trait we computed the two terms in Eq. 2, which should converge to the
genic variance and LD contributions, which represent the genic and LD components to the HE regression
estimate. Fig. S5 shows that in the presence of directional LD, the overall bias is in the HE regression
estimate is due to an exaggerated estimate of the LD contribution.

Estimating variance explained by GWAS SNPs

To decompose the variance explained by GWAS SNPs in African Americans, we needed four quantities:
(i) effect sizes of GWAS SNPs, (ii) their allele frequencies in Africans and Europeans, and (iii) the mean
and variance of global ancestry in African Americans (Equation 1).

We retrieved the summary statistics of 26 traits from GWAS catalog [40]. Full list of traits and the
source papers [44, 57–64] are listed in Table S1. To maximize the number of variants discovered, we chose
summary statistics from studies that were conducted in both European and multi-ancestry samples and
that reported the following information: effect allele, effect size, p-value, and genomic position. For birth
weight, we downloaded the data from the Early Growth Genetics (EGG) consortium website [61] since the
version reported on the GWAS catalog is incomplete. For skin pigmentation, we chose summary statistics

from the UKB [65] released by the Neale Lab (http://www.nealelab.is/uk-biobank) and processed by Ju

and Mathieson [44] to represent effect sizes estimated among individuals of European ancestry. We

also selected summary statistics from Lona-Durazo *et al.* (2019) where effect sizes were meta-analyzed

across four admixed cohorts [57]. Lona-Durazo *et al.* provide summary statistics separately with and

without conditioning on rs1426654 and rs35397 – two large effect variants in *SLC24A5* and *SLC45A2*.

We used the 'conditioned' effect sizes and added in the effects of rs1426654 and rs35397 to estimate

genetic variance.

We selected independent hits for each trait by pruning and thresholding with PLINK v1.90b6.21 [66]

in two steps as in Ju *et al.* (2020) [44]. We used the genotype data of GBR from the 1000 genome

project [39] as the LD reference panel. We kept only SNPs (indels were removed) that passed the

genome-wide significant threshold (*--clump-p1 5e-8*) with a pairwise LD cutoff of 0.05 (*--clump-r2 0.05*)

and a physical distance threshold of 250Kb (*--clump-kb 250*) for clumping. Second, we applied a second

round of clumping (*--clump-kb 100*) to remove SNPs within 100kb.

When GWAS was carried out separately in different ancestry cohorts in the same study, we used

inverse-variance weighting to meta-analyze effect sizes for variants that were genome-wide significant

(p-value $< 5 \times 10^{-8}$) in at least one cohort. This allowed us to maximize the discovery of variants such

as the Duffy null allele that are absent among individuals of European ancestry but polymorphic in other

populations [47].

We used allele frequencies from the 1000 Genomes CEU and YRI to represent the allele frequencies

of GWAS SNPs in Europeans and Africans, respectively, making sure that the alleles reported in the

summary statistics matched the alleles reported in the 1000 Genomes. We estimated the global ances-

try of ASW individuals (N = 74) with CEU and YRI individuals from 1000 genome (phase 3) using

ADMIXTURE 1.3.0 [67] with k=2 and used it to calculate the mean (proportion of African ancestry =

0.767) and variance (0.018) of global ancestry in ASW. With the effect sizes, allele frequencies, and the

mean and variance in ancestry, we calculated the four components of genetic variance using Equation 1

and expressed them as a fraction of the total genetic variance.

Initially, the multi-ancestry summary statistics for a few traits (NEU, WBC, MON, MCH, BAS)

yielded values > 1 for the proportion of variance explained. This is likely because, despite LD pruning,

some of the variants in the model are not independent and tag large effect variants under divergent

selection such as the Duffy null allele, leading to an inflated contribution of LD. We checked this by

calculating the pairwise contribution , i.e., $\beta_i \beta_j (f_i^A - f_i^B)(f_j^A - f_j^B)$, of all SNPs in the model and show

long-range positive LD between variants on chromosome 1 for NEU, WBC, and MON, especially with the

Duffy null allele (Fig. S6A-C). A similar pattern was observed on chromosome 16 for MCH, confirming

our suspicion. This also suggests that for certain traits, pruning and thresholding approaches are not

guaranteed to yield independent hits. To get around this problem, we retained only one association with

the lowest p-value, each from chromosome 1 (rs2814778 for NEU, WBC, and MON) and chromosome 16

(rs13331259 for MCH) (Fig. S6D). For BAS, we observed that the variance explained was driven by a

rare variant (rs188411703, MAF = 0.0024) of large effect ($\beta = -2.27$). We believe this effect estimate

to be inflated and therefore, we removed it from our calculation.

As a sanity check, we independently estimated the genetic variance as the variance in polygenic

scores, calculated using --*score* flag in PLINK, [66] in ASW individuals. We compared the first estimate of the genetic variance to the second (Fig. S7) to confirm two things: (i) the allele frequencies, and mean and variance in ancestry are estimated correctly, and (ii) the variants are more or less independent in that they do not absorb the effects of other variants in the model. We show that the two estimates of the genetic variance are strongly correlated ($r \sim 0.85$, Fig. S7). The 95% confidence intervals were calculated by sampling individuals with replacement 10,000 times.

Code availability

We carried out all analyses in R version 4.2.3 [56], PLINK v1.90b6.21 and PLINK 2.0 [66, 68], and GCTA version 1.94.1 [5]. All code is freely available on https://github.com/jinguohuang/admix_heritability.git.

Acknowledgements

We thank Iain Mathieson and Doc Edge for helpful comments on the manuscript. This study was funded by National Institute of General Medical Sciences award R00GM137076 to A.A.Z. The content of this paper is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1. Lynch, M. & Walsh, B. *Genetics and analysis of quantitative traits* 1–980 (Sinauer Associates, Inc, 1998).

2. Visscher, P. M., Hill, W. G. & Wray, N. R. Heritability in the genomics era–concepts and misconceptions. *Nature reviews. Genetics* **9,** 255–66 (4 2008).

3. Yang, J., Benyamin, B., *et al.* Common SNPs explain a large proportion of the heritability for human height. *Nature Genetics* **42,** 565–569 (7 2010).

4. Yang, J., Manolio, T. A., *et al.* Genome partitioning of genetic variation for complex traits using common SNPs. *Nature Genetics 2011 43:6* **43,** 519–525 (6 2011).

5. Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide complex trait analysis. *American journal of human genetics* **88,** 76–82 (1 2011).

6. Wainschtein, P., Jain, D., *et al.* Assessing the contribution of rare variants to complex trait heritability from whole-genome sequence data. *Nature Genetics 2022 54:3* **54,** 263–273 (3 2022).

7. Browning, S. R. & Browning, B. L. Population structure can inflate SNP-based heritability estimates. *American Journal of Human Genetics* **89,** 191–193 (1 2011).

8. Goddard, M. E., Lee, S. H., Yang, J., Wray, N. R. & Visscher, P. M. Response to browning and browning. *American Journal of Human Genetics* **89,** 193–195 (1 2011).

9. Kumar, S. K., Feldman, M. W., Rehkopf, D. H. & Tuljapurkar, S. Limitations of GCTA as a solution to the missing heritability problem. *Proceedings of the National Academy of Sciences of the United States of America* **113,** E61–E70 (1 2016).

10. Yang, J., Leec, S. H., Wraya, N. R., Goddardd, M. E. & Visscher, P. M. GCTA-GREML accounts for linkage disequilibrium when estimating genetic variance from genome-wide SNPs. *Proceedings of the National Academy of Sciences* **113,** E4579–E4580 (32 2016).

11. Lin, Z., Seal, S. & Basu, S. Estimating SNP heritability in presence of population substructure in biobank-scale datasets. *Genetics* **220** (4 2022).

12. Border, R., O'Rourke, S., *et al.* Assortative mating biases marker-based heritability estimators. *Nature Communications 2022 13:1* **13,** 1–10 (1 2022).

13. Visscher, P. M., Yang, J. & Goddard, M. E. A Commentary on 'Common SNPs Explain a Large Proportion of the Heritability for Human Height' by Yang et al. (2010). *Twin Research and Human Genetics* **13,** 517–524 (6 2010).

14. De los Campos, G., Sorensen, D. & Gianola, D. Genomic Heritability: What Is It? *PLOS Genetics* **11,** e1005048 (5 2015).

15. Rawlik, K., Canela-Xandri, O., Oolliams, J. W. & Tenesa, A. SNP heritability: What are we estimating? *bioRxiv,* 2020.09.15.276121 (2020).

599 16. Lara, L. A. C., Pocrnic, I., de P. Oliveira, T., Gaynor, R. C. & Gorjanc, G. Temporal and genomic
600   analysis of additive genetic variance in breeding programmes. *Heredity 2021 128:1* **128,** 21–32 (1
601   2021).

602 17. Wojcik, G. L., Graff, M., *et al.* Genetic analyses of diverse populations improves discovery for
603   complex traits. *Nature* **570,** 514–518 (7762 2019).

604 18. Ben-Eghan, C., Sun, R., *et al.* Don't ignore genetic data from minority populations. *Nature 2021*
605   *585:7824* **585,** 184–186 (7824 2020).

606 19. Verma, A., Damrauer, S. M., *et al.* The Penn Medicine BioBank: Towards a Genomics-Enabled
607   Learning Healthcare System to Accelerate Precision Medicine in a Diverse Population. *Journal of*
608   *Personalized Medicine* **12,** 1974 (12 2022).

609 20. Fatumo, S., Chikowore, T., *et al.* A roadmap to increase diversity in genomic studies. *Nature*
610   *Medicine 2022 28:2* **28,** 243–250 (2 2022).

611 21. The All of Us Research Program Investigators. The "All of Us" Research Program. *New England*
612   *Journal of Medicine* **381,** 668–676 (7 2019).

613 22. Sohail, M., Chong, A. Y., *et al.* Nationwide genomic biobank in Mexico unravels demographic
614   history and complex trait architecture from 6,057 individuals. *bioRxiv,* 2022.07.11.499652 (2022).

615 23. Johnson, R., Ding, Y., *et al.* The UCLA ATLAS Community Health Initiative: Promoting precision
616   health research in a diverse biobank. *Cell Genomics* **3,** 100243 (1 2023).

617 24. Haseman, J. K. & Elston, R. C. The investigation of linkage between a quantitative trait and a
618   marker locus. *Behavior genetics* **2,** 3–19 (1 1972).

619 25. Pfaff, C. L., Parra, E. J., *et al.* Population structure in admixed populations: effect of admixture
620   dynamics on the pattern of linkage disequilibrium. *American journal of human genetics* **68,** 198–207
621   (1 2001).

622 26. Zaitlen, N., Huntsman, S., *et al.* The Effects of Migration and Assortative Mating on Admixture
623   Linkage Disequilibrium. *Genetics* **205,** 375–383 (1 2017).

624 27. Verdu, P. & Rosenberg, N. A. A General Mechanistic Model for Admixture Histories of Hybrid
625   Populations. *Genetics* **189,** 1413–1426 (4 2011).

626 28. Corre, V. L. & Kremer, A. Genetic Variability at Neutral Markers, Quantitative Trait Loci and
627   Trait in a Subdivided Population Under Selection. *Genetics* **164,** 1205–1219 (3 2003).

628 29. Berg, J. J. & Coop, G. A Population Genetic Signal of Polygenic Adaptation. *PLoS Genetics* **10,**
629   e1004412 (8 2014).

630 30. Bulmer, M. G. The Effect of Selection on Genetic Variability. *The American Naturalist* **105,** 201–
631   211 (943 1971).

632 31. Yair, S. & Coop, G. Population differentiation of polygenic score predictions under stabilizing
633   selection. *Philosophical Transactions of the Royal Society B* **377** (1852 2022).

32. Balding, D. J. & Nichols, R. A. A method for quantifying differentiation between populations at multi-allelic loci and its implications for investigating identity and paternity. *Genetica* **96**, 3–12 (1-2 1995).

33. Mathew, B., Léon, J. & Sillanpää, M. J. A novel linkage-disequilibrium corrected genomic relationship matrix for SNP-heritability estimation and genomic prediction. *Heredity 2017 120:4* **120**, 356–368 (4 2017).

34. Ma, R. & Dicker, L. H. The Mahalanobis kernel for heritability estimation in genome-wide association studies: fixed-effects and random-effects methods (2019).

35. Min, A., Thompson, E. & Basu, S. Comparing heritability estimators under alternative structures of linkage disequilibrium. *G3 Genes|Genomes|Genetics* **12** (8 2022).

36. Zaitlen, N., Pasaniuc, B., *et al.* Leveraging population admixture to characterize the heritability of complex traits. *Nature Genetics* **46**, 1356–1362 (12 2014).

37. Zaidi, A. A., Mattern, B. C., *et al.* Investigating the case of human nose shape and climate adaptation. *PLoS Genetics* **13**, 2017 (3 2017).

38. Schraiber, J. G. & Edge, M. D. Heritability within groups is uninformative about differences among groups: cases from behavioral, evolutionary, and statistical genetics. *bioRxiv,* 2023.11.06.565864 (2023).

39. Auton, A., Abecasis, G. R., *et al.* A global reference for human genetic variation. *Nature 2015 526:7571* **526**, 68–74 (7571 2015).

40. Sollis, E., Mosaku, A., *et al.* The NHGRI-EBI GWAS Catalog: knowledgebase and deposition resource. *Nucleic Acids Research* **51**, D977–D985 (D1 2023).

41. Jablonski, N. G. the Evolution of Human Skin and Skin Color. *Annual Review of Anthropology* **33**, 585–623 (1 2004).

42. Lamason, R. L., Mohideen, M. A. P., *et al.* SLC24A5, a putative cation exchanger, affects pigmentation in zebrafish and humans. *Science* **310**, 1782–1786 (5755 2005).

43. Beleza, S., Johnson, N. A., *et al.* Genetic architecture of skin and eye color in an African-European admixed population. *PLoS genetics* **9** (ed Spritz, R. A.) e1003372 (3 2013).

44. Ju, D. & Mathieson, I. The evolution of skin pigmentation-associated variation in West Eurasia. *Proceedings of the National Academy of Sciences of the United States of America* **118**, e2009227118 (1 2020).

45. Nalls, M. A., Wilson, J. G., *et al.* Admixture Mapping of White Cell Count: Genetic Locus Responsible for Lower White Blood Cell Count in the Health ABC and Jackson Heart Studies. *The American Journal of Human Genetics* **82**, 81–87 (1 2008).

46. Reich, D., Nalls, M. A., *et al.* Reduced Neutrophil Count in People of African Descent Is Due To a Regulatory Variant in the Duffy Antigen Receptor for Chemokines Gene. *PLoS Genetics* **5** (ed Visscher, P. M.) e1000360 (1 2009).

47. McManus, K. F., Taravella, A. M., *et al.* Population genetic analysis of the DARC locus (Duffy)
reveals adaptation from standing variation associated with malaria resistance in humans. *PLOS*
Genetics **13,** e1006560 (3 2017).

48. Kumar, S. K., Feldman, M. W., Rehkop, D. H. & Tuljapurkar, S. Reply to Yang et al.: GCTA
produces unreliable heritability estimates. *Proceedings of the National Academy of Sciences* **113,**
E4581–E4581 (32 2016).

49. Speed, D., Hemani, G., Johnson, M. R. & Balding, D. J. Improved heritability estimation from
genome-wide SNPs. *American journal of human genetics* **91,** 1011–21 (6 2012).

50. Speed, D., Cai, N., Johnson, M. R., Nejentsev, S. & Balding, D. J. Reevaluation of SNP heritability
in complex human traits. *Nature Genetics* **49,** 986–992 (7 2017).

51. Chan, T. F., Rui, X., *et al.* Estimating heritability explained by local ancestry and evaluating strat-
ification bias in admixture mapping from summary statistics. *bioRxiv,* 2023.04.10.536252 (2023).

52. Schwartzman, A., Schork, A. J., Zablocki, R. & Thompson, W. K. A simple, consistent estimator
of SNP heritability from genome-wide association studies. *https://doi.org/10.1214/19-AOAS1291*
13, 2509–2538 (4 2019).

53. Hou, K., Ding, Y., *et al.* Causal effects on complex traits are similar for common variants across
segments of different continental ancestries within admixed individuals. *Nature Genetics 2023 55:4*
55, 549–558 (4 2023).

54. Backman, J. D., Li, A. H., *et al.* Exome sequencing and analysis of 454,787 UK Biobank participants.
Nature 2021 599:7886 **599,** 628–634 (7886 2021).

55. Lin, M., Park, D. S., Zaitlen, N. A., Henn, B. M. & Gignoux, C. R. Admixed Populations Improve
Power for Variant Discovery and Portability in Genome-Wide Association Studies. *Frontiers in*
Genetics **12,** 673167 (2021).

56. R Core Team. *R: A Language and Environment for Statistical Computing* R Foundation for Sta-
tistical Computing (Vienna, Austria, 2023).

57. Lona-Durazo, F., Hernandez-Pacheco, N., *et al.* Meta-analysis of GWA studies provides new insights
on the genetic architecture of skin pigmentation in recently admixed populations. *BMC Genetics*
20, 1–16 (1 2019).

58. Yengo, L., Vedantam, S., *et al.* A saturated map of common genetic variants associated with human
height. *Nature* **610,** 704–712 (7933 2022).

59. Hoffmann, T. J., Choquet, H., *et al.* A large multiethnic genome-wide association study of adult
body mass index identifies novel loci. *Genetics* **210,** 499–515 (2 2018).

60. Pulit, S. L., Stoneman, C., *et al.* Meta-Analysis of genome-wide association studies for body fat
distribution in 694 649 individuals of European ancestry. *Human Molecular Genetics* **28,** 166–174
(1 2019).

61. Warrington, N. M., Beaumont, R. N., *et al.* Maternal and fetal genetic effects on birth weight and
their relevance to cardio-metabolic risk factors. *Nature Genetics 2019 51:5* **51,** 804–814 (5 2019).

62.   Surendran, P., Feofanova, E. V., *et al.* Discovery of rare variants associated with blood pressure
regulation through meta-analysis of 1.3 million individuals. *Nature Genetics* **52,** 1314–1332 (12
2020).

63.   Graham, S. E., Clarke, S. L., *et al.* The power of genetic diversity in genome-wide association studies
of lipids. *Nature* **600,** 675–679 (7890 2021).

64.   Chen, M. H., Raffield, L. M., *et al.* Trans-ethnic and Ancestry-Specific Blood-Cell Genetics in
746,667 Individuals from 5 Global Populations. *Cell* **182,** 1198–1213.e14 (5 2020).

65.   Bycroft, C., Freeman, C., *et al.* The UK Biobank resource with deep phenotyping and genomic
data. *Nature* **562,** 203–209 (7726 2018).

66.   Chang, C. C., Chow, C. C., *et al.* Second-generation PLINK: rising to the challenge of larger and
richer datasets. *GigaScience* **4,** s13742–015–0047–8 (1 2015).

67.   Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in unrelated
individuals. *Genome research* **19,** 1655–64 (9 2009).

68.   Purcell, S., Neale, B., *et al.* PLINK: a tool set for whole-genome association and population-based
linkage analyses. *American journal of human genetics* **81,** 559–75 (3 2007).

69.   Chen, G. B. Estimating heritability of complex traits from genome-wide association studies using
IBS-based Haseman-Elston regression. *Frontiers in Genetics* **5,** 72296 (APR 2014).

 # Appendix

 ## A1   Variance in ancestry

726 We denote variance and covariance with $\mathbb{V}(.)$ and $\mathbb{C}(.)$ and used the expressions in [26] to generate the

727 expected value for the variance in ancestry, i.e., $\mathbb{V}(\theta)$. This is straightforward for the HI model, where at

728 time t $\mathbb{V}(\theta_t) = \mathbb{V}(\theta_{t-1})\frac{(1+P_{t-1})}{2}$. $P_t = Cor(\theta_m, \theta_f)$ measures the strength of assortative mating, i.e, the

729 correlation between the ancestry across mating pairs (θ_m, θ_f) at time t. For simplicity, we assumed this to

730 be constant in every generation, i.e. $P_t = P_{t-1} = P$ following [26]. Since our notation slightly differs from

731 [26], we re-derived the expression for $V(\theta_t)$ for the CGF model where population B receives a constant

732 amount q of gene flow from population A in every generation. Note, that $\mathbb{E}(\theta_t) = q + (1-q)\mathbb{E}(\theta_{t-1})$.

733 Then,

$$
\begin{aligned}
\mathbb{V}(\theta_t) =& \mathbb{E}(\theta_t^2) - \mathbb{E}(\theta_t)^2 \\
=& q + (1-q)\mathbb{E}\left[\left(\frac{\theta_{t-1}^m + \theta_{t-1}^f}{2}\right)\left(\frac{\theta_{t-1}^m + \theta_{t-1}^f}{2}\right)\right] - \{q + (1-q)\mathbb{E}(\theta_{t-1})\}^2 \\
=& q + \frac{(1-q)}{4}\{2\mathbb{E}(\theta_{t-1}^2) + 2\mathbb{E}(\theta_{t-1}^m\theta_{t-1}^f)\} - \{q^2 + 2q(1-q)\mathbb{E}(\theta_{t-1}) + (1-q)^2\mathbb{E}(\theta_{t-1})^2\} \\
=& q + \frac{1-q}{2}\mathbb{E}(\theta_{t-1}^2) + \frac{1-q}{2}\mathbb{E}(\theta_{t-1}^m\theta_{t-1}^f) - q^2 - 2q(1-q)\mathbb{E}(\theta_{t-1}) - (1-q)^2\mathbb{E}(\theta_{t-1})^2 \\
=& q(1-q) + \frac{1-q}{2}\{\mathbb{V}(\theta_{t-1}) + \mathbb{E}(\theta_{t-1})^2\} + \frac{1-q}{2}\{\mathbb{C}(\theta_{t-1}^m, \theta_{t-1}^f) + \mathbb{E}(\theta_{t-1})^2\} - 2q(1-q)\mathbb{E}(\theta_{t-1}) - \mathbb{E}(\theta_{t-1})^2 \\
=& q(1-q) + \frac{1-q}{2}\mathbb{V}(\theta_{t-1}) + \frac{1-q}{2}\mathbb{E}(\theta_{t-1})^2 + \frac{1-q}{2}P_{t-1}\mathbb{V}(\theta_{t-1}) + \frac{1-q}{2}\mathbb{E}(\theta_{t-1})^2 - 2q(1-q)\mathbb{E}(\theta_{t-1}) - \mathbb{E}(\theta_{t-1})^2 \\
=& q(1-q) + \frac{1-q}{2}\mathbb{V}(\theta_{t-1})\{1 + P_{t-1}\} + (1-q)\mathbb{E}(\theta_{t-1})^2 - 2q(1-q)\mathbb{E}(\theta_{t-1}) - (1-q)^2\mathbb{E}(\theta_{t-1})^2 \\
=& q(1-q)\mathbb{E}(\theta_{t-1})^2 + q(1-q)\{1 - 2\mathbb{E}(\theta_{t-1})\} + \frac{1-q}{2}\mathbb{V}(\theta_{t-1})\{1 + P_{t-1}\}
\end{aligned}
$$

 ## A2   Genetic variance

735 Let $y = g + e$, where y is the phenotypic value of an individual, g is the genotypic value, and e is random

736 error. We assume additive effects such that $g = \sum_{i=1}^{m}\beta_i x_i$ where β_i is the effect size of the i^{th} biallelic

737 locus and $x_i \in \{0, 1, 2\}$ is the number of copies of the trait-increasing allele. Then, the genetic variance

738 V_g is:

$$
V_g = \mathbb{V}(\sum_{i=1}^{m}\beta_i x_i) = \sum_{i=1}^{m}\beta_i^2\,\mathbb{V}(x_i) + \sum_{j\neq i}\beta_i\beta_j\,\mathbb{C}(x_i, x_j)
$$

739 In the following sections, we decompose $\mathbb{V}(x_i)$ and $\mathbb{C}(x_i, x_j)$ further as functions of ancestry.

 ### A2.1 $\mathbb{V}(x_i)$

741 We first derive $\mathbb{V}(x_i)$ as a function of ancestry (θ) using the law of total variance:

$$\mathbb{V}(x_i) = \mathbb{E}_{\theta}\{\mathbb{V}(x_i|\theta)\} + \mathbb{V}\{\mathbb{E}_{\theta}(x_i|\theta)\}$$

742   where $\mathbb{E}_{\theta}$ represents the expectation taken over θ.

743   **A2.1.1** $\mathbb{E}_{\theta}\{\mathbb{V}(x_i|\theta)\}$

744   We derive $\mathbb{V}(x_i|\theta)$ by further conditioning on the local ancestry at each locus.

$$\mathbb{V}(x_i|\theta) = \mathbb{E}_{\gamma}\{\mathbb{V}(x_i|\gamma, \theta)\} + \mathbb{V}\{\mathbb{E}_{\gamma}(x_i|\gamma, \theta)\}$$

745   where $\mathbb{E}_{\gamma}$ represents expectation taken over local ancestry. Since we are interested in the variance at
746   a single locus, we will ignore the subscript i and denote the frequency of the trait-increasing allele in
747   populations A and B with f^{A} and f^{B}, respectively.

$$
\begin{aligned}
\mathbb{E}_{\gamma}\{\mathbb{V}(x_i|\gamma, \theta)\} &= \mathbb{V}(x_i|\gamma = 0, \theta)\,\mathbb{P}(\gamma = 0|\theta) + \mathbb{V}(x_i|\gamma = 1, \theta)\,\mathbb{P}(\gamma = 1|\theta) + \mathbb{V}(x_i|\gamma = 2, \theta)\,\mathbb{P}(\gamma = 2|\theta) \\
&= 2\,f^{\mathrm{B}}(1 - f^{\mathrm{B}})(1 - \theta)^2 + \{f^{\mathrm{A}}(1 - f^{\mathrm{A}}) + f^{\mathrm{B}}(1 - f^{\mathrm{B}})\}2\theta(1 - \theta) + 2\,f^{\mathrm{A}}(1 - f^{\mathrm{A}})\theta^2 \\
&= (2\,f^{\mathrm{B}} - 2f^{A^2})(1 - 2\theta + \theta^2) + (f^{\mathrm{A}} - f^{A^2} + f^{\mathrm{B}} - f^{B^2})(2\theta - 2\theta^2) + (2\,f^{\mathrm{A}} - 2f^{A^2})\theta^2 \\
&= 2\,f^{\mathrm{B}} - 2\theta\,f^{\mathrm{B}} - 2f^{B^2} + 2\theta f^{B^2} + 2\theta\,f^{\mathrm{A}} - 2\theta f^{A^2} \\
&= 2\,f^{\mathrm{B}}(1 - \theta) - 2f^{B^2}(1 - \theta) + 2\theta\,f^{\mathrm{A}}(1 - f^{\mathrm{A}}) \\
&= 2\,f^{\mathrm{B}}(1 - f^{\mathrm{B}})(1 - \theta) + 2\theta\,f^{\mathrm{A}}(1 - f^{\mathrm{A}})
\end{aligned}
$$

748   To derive $\mathbb{V}\{\mathbb{E}_{\gamma}(x|\gamma, \theta)\}$, note that

$$
\begin{aligned}
\mathbb{E}_{\gamma}(x|\gamma, \theta) &= \mathbb{E}_{\gamma}\{\mathbb{E}(x|\theta)\} \\
&= \mathbb{E}(x|\gamma = 0, \theta)\,\mathbb{P}(\gamma = 0|\theta) + \mathbb{E}(x|\gamma = 1, \theta)\,\mathbb{P}(\gamma = 1|\theta) + \mathbb{E}(x|\gamma = 2, \theta)\,\mathbb{P}(\gamma = 2|\theta) \\
&= 2\theta\,f^{\mathrm{A}} + 2(1 - \theta)\,f^{\mathrm{B}}
\end{aligned}
$$

749   And,

$$
\begin{aligned}
\mathbb{V}\{\mathbb{E}_{\gamma}(x|\gamma, \theta)\} &= [\mathbb{E}(x|\gamma = 0, \theta) - \mathbb{E}(x|\theta)]^2\,\mathbb{P}(\gamma = 0|\theta) \\
&\quad + [\mathbb{E}(x|\gamma = 1, \theta) - \mathbb{E}(x|\theta)]^2\,\mathbb{P}(\gamma = 1|\theta) \\
&\quad + [\mathbb{E}(x|\gamma = 2, \theta) - \mathbb{E}(x|\theta)]^2\,\mathbb{P}(\gamma = 2|\theta) \\
&= \theta^2\left[2\,f^{\mathrm{A}} - \{2\theta\,f^{\mathrm{A}} + 2(1 - \theta)\,f^{\mathrm{B}}\}\right]^2 \\
&\quad + 2\theta(1 - \theta)\left[f^{\mathrm{A}} + f^{\mathrm{B}} - \{2\theta\,f^{\mathrm{A}} + 2(1 - \theta)\,f^{\mathrm{B}}\}\right]^2 \\
&\quad + (1 - \theta)^2\left[2\,f^{\mathrm{B}} - \{2\theta\,f^{\mathrm{A}} + 2(1 - \theta)\,f^{\mathrm{B}}\}\right]^2 \\
&= 2\theta(1 - \theta)(f^{\mathrm{A}} - f^{\mathrm{B}})^2
\end{aligned}
$$

750 Putting this together,

$$
\begin{aligned}
\mathbb{E}_{\theta}\{\mathbb{V}(x_i|\theta)\} &= \mathbb{E}_{\theta}\{2\,f^{\mathrm{B}}(1-f^{\mathrm{B}})(1-\theta) + 2\theta\,f^{\mathrm{A}}(1-f^{\mathrm{A}}) + 2\theta(1-\theta)(f^{\mathrm{A}}-f^{\mathrm{B}})^2\} \\
&= 2\,f^{\mathrm{B}}(1-f^{\mathrm{B}})\{1-\mathbb{E}_{\theta}(\theta)\} + 2\,\mathbb{E}_{\theta}(\theta)\,f^{\mathrm{A}}(1-f^{\mathrm{A}}) + 2\,\mathbb{E}_{\theta}(\theta-\theta^2)(f^{\mathrm{A}}-f^{\mathrm{B}})^2 \\
&= 2\,f^{\mathrm{B}}(1-f^{\mathrm{B}})\{1-\mathbb{E}_{\theta}(\theta)\} + 2\,\mathbb{E}_{\theta}(\theta)\,f^{\mathrm{A}}(1-f^{\mathrm{A}}) + 2\{\mathbb{E}_{\theta}(\theta)-\mathbb{E}_{\theta}(\theta^2)\}(f^{\mathrm{A}}-f^{\mathrm{B}})^2 \\
&= 2\,f^{\mathrm{B}}(1-f^{\mathrm{B}})\{1-\mathbb{E}_{\theta}(\theta)\} + 2\,\mathbb{E}_{\theta}(\theta)\,f^{\mathrm{A}}(1-f^{\mathrm{A}}) + 2\{\mathbb{E}_{\theta}(\theta)-\mathbb{V}(\theta)-\mathbb{E}_{\theta}(\theta)^2\}(f^{\mathrm{A}}-f^{\mathrm{B}})^2 \\
&= 2\,f^{\mathrm{B}}(1-f^{\mathrm{B}})\{1-\mathbb{E}_{\theta}(\theta)\} + 2\,\mathbb{E}_{\theta}(\theta)\,f^{\mathrm{A}}(1-f^{\mathrm{A}}) + 2\,\mathbb{E}_{\theta}(\theta)(1-\mathbb{E}_{\theta}(\theta))(f^{\mathrm{A}}-f^{\mathrm{B}})^2 - 2\,\mathbb{V}(\theta)(f^{\mathrm{A}}-f^{\mathrm{B}})^2
\end{aligned}
$$

751 **A2.1.2** $\mathbb{V}\{\mathbb{E}_{\theta}(x_i|\theta)\}$

752 Recall from the previous section that $\mathbb{E}_{\theta}(x_i|\theta) = 2\theta\,f^{\mathrm{A}} + 2(1-\theta)\,f^{\mathrm{B}}$. Then,

$$
\begin{aligned}
\mathbb{V}\{\mathbb{E}_{\theta}(x_i|\theta)\} &= \mathbb{V}\{2\theta\,f^{\mathrm{A}} + 2(1-\theta)\,f^{\mathrm{B}}\} \\
&= 4\,\mathbb{V}(\theta)f^{A^2} + 4\,\mathbb{V}(1-\theta)f^{B^2} + 2\,\mathbb{C}(2\theta\,f^{\mathrm{A}}, 2(1-\theta\,f^{\mathrm{B}}) \\
&= 4\,\mathbb{V}(\theta)f^{A^2} + 4\,\mathbb{V}(1-\theta)f^{B^2} - 8\,f^{\mathrm{A}}\,f^{\mathrm{B}}\,\mathbb{V}(\theta) \\
&= 4\,\mathbb{V}(\theta)(f^{\mathrm{A}}-f^{\mathrm{B}})^2
\end{aligned}
$$

753 We are now ready to express $\mathbb{V}(x_i)$:

$$
\begin{aligned}
\mathbb{V}(x_i) &= 2\,f^{\mathrm{B}}(1-f^{\mathrm{B}})\{1-\mathbb{E}_{\theta}(\theta)\} + 2\,\mathbb{E}_{\theta}(\theta)\,f^{\mathrm{A}}(1-f^{\mathrm{A}}) + 2\,\mathbb{E}_{\theta}(\theta)(1-\mathbb{E}_{\theta}(\theta))(f^{\mathrm{A}}-f^{\mathrm{B}})^2 \\
&\quad - 2\,\mathbb{V}(\theta)(f^{\mathrm{A}}-f^{\mathrm{B}})^2 + 4\,\mathbb{V}(\theta)(f^{\mathrm{A}}-f^{\mathrm{B}})^2 \\
&= 2\,\mathbb{E}_{\theta}(\theta)\,f_i^{\mathrm{A}}(1-f_i^{\mathrm{A}}) + 2\{1-\mathbb{E}_{\theta}(\theta)\}\,f_i^{\mathrm{B}}(1-f_i^{\mathrm{B}}) \\
&\quad + 2\,\mathbb{E}_{\theta}(\theta)\{1-\mathbb{E}_{\theta}(\theta)\}(f_i^{\mathrm{A}}-f_i^{\mathrm{B}})^2 - 2\,\mathbb{V}(\theta)(f_i^{\mathrm{A}}-f_i^{\mathrm{B}})^2
\end{aligned}
$$

754 Note, that we can also express $V(x_i)$ as:

$$
\mathbb{V}(x_i) = 2f_i(1-f_i) + 2\,\mathbb{V}(\theta)(f_i^{A}-f_i^{B})^2
$$

755 where the second term is the contribution of population structure to the genetic variance at locus i.

756 **A2.2** $\mathbb{C}(x_i, x_j)$

757 We can derive $\mathbb{C}(x_i, x_j)$ using the law of total covariance:

$$\mathbb{C}(x_i, x_j) = \mathop{\mathbb{E}}_{\theta}\{\mathbb{C}(x_i, x_j | \theta)\} + \mathbb{C}\{\mathop{\mathbb{E}}_{\theta}(x_i|\theta), \mathop{\mathbb{E}}_{\theta}(x_j|\theta)\}$$

$$=0 + \mathbb{C}\{2\,f_i^{A}\,\theta + 2\,f_i^{B}(1-\theta), 2\,f_j^{A}\,\theta + 2\,f_j^{B}(1-\theta)\}$$

$$= \mathbb{C}(2\,f_i^{A}\,\theta, 2\,f_j^{A}\,\theta) + \mathbb{C}(2\,f_i^{A}\,\theta, 2\,f_j^{B}(1-\theta)) +$$

$$\mathbb{C}(2\,f_i^{B}(1-\theta), 2\,f_j^{A}\,\theta) + \mathbb{C}(2\,f_i^{B}(1-\theta), 2\,f_j^{B}(1-\theta))$$

$$= 4\,\mathbb{V}(\theta)(f_i^{A} - f_i^{B})(f_j^{A} - f_j^{B})$$

758 $\mathbb{E}_{\theta}\{\mathbb{C}(x_i, x_j | \theta)\} = 0$ because we assume that the loci are unlinked and therefore, x_i and x_j are condi-

759 tionally independent. Putting this all together, we get the genetic variance in admixed populations as

760 presented in the main text:

$$V_g = \sum_{i=1}^{m} \beta_i^2\,\mathbb{V}(x_i) + \sum_{j\neq i} \beta_i \beta_j\,\mathbb{C}(x_i, x_j)$$

$$= \sum_{i=1}^{m} \beta_i^2\,2\,\mathop{\mathbb{E}}_{\theta}(\theta)\,f_i^{A}(1 - f_i^{A}) + \sum_{i=1}^{m} \beta_i^2\,2\{1 - \mathop{\mathbb{E}}_{\theta}(\theta)\}\,f_i^{B}(1 - f_i^{B})$$

$$+ \sum_{i=1}^{m} \beta_i^2\,2\,\mathop{\mathbb{E}}_{\theta}(\theta)\{1 - \mathop{\mathbb{E}}_{\theta}(\theta)\}(f_i^{A} - f_i^{B})^2 +$$

$$+ \sum_{i=1}^{m} \beta_i^2\,2\,\mathbb{V}(\theta)(f_i^{A} - f_i^{B})^2]$$

$$+ \sum_{j\neq i} \beta_i \beta_j\,4\,\mathbb{V}(\theta)(f_i^{A} - f_i^{B})(f_j^{A} - f_j^{B})$$

761 The only difference being that in the main text we use $\mathbb{E}$ instead of $\mathbb{E}_{\theta}$ for simplicity. With two 'unad-

762 mixed' source populations with equal number of individuals, $\mathbb{E}(\theta) = 0.5$ and $\mathbb{V}(\theta) = \mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} = 0.25$

763 and V_g reduces to:

$$V_g = \mathbb{V}\left(\sum_{i=1}^{m} \beta_i x_i\right) = \sum_{i=1}^{m} \beta_i^2\,\mathbb{V}(x_i) + \sum_{j\neq i} \beta_i \beta_j\,\mathbb{C}(x_i, x_j)$$

$$= \sum_{i=1}^{m} \beta_i^2\left[f_i^{A}(1 - f_i^{A}) + f_i^{B}(1 - f_i^{B})\right]$$

$$+ \sum_{i=1}^{m} \beta_i^2(f_i^{A} - f_i^{B})^2$$

$$+ \sum_{i\neq j} \beta_i \beta_j(f_i^{A} - f_i^{B})(f_j^{A} - f_j^{B})$$

764 A3 The effect of genotype scale on $\hat{V}_g$

765 In the main text, we showed that both GREML and Haseman-Elston regression estimates of V_g depend

766 on how the genotypes are scaled. We provide an explanation of this behavior using the Haseman-Elston

767 (HE) regression estimator, which is asymptotically equivalent to the GREML estimator if effects are

768 uncorrelated [69] but which, unlike GREML, has a closed-form solution.

769 A3.1   No directional LD

770 A3.1.1   Scaling by $2f_i(1-f_i)$

771 First, let's assume a genetic architecture where all loci contribute equally to the genetic variance and there
772 is no LD contribution. With the standard scaling, the genotype at a given locus i is $z_i = \frac{x_i - 2f_i}{2f_i(1-f_i)}$ where
773 f_i is the frequency of the allele in the population. Under the random-effects model, this is equivalent
774 to saying that the unscaled effects are: $\beta_i \sim \mathcal{N}(0, \frac{\sigma_u^2}{2mf_i(1-f_i)})$, σ_u^2 being the parameter of interest. In a
775 panmictic population,

$$
\begin{aligned}
V_g &= \mathbb{V}\left(\sum_{i=1}^m \beta_i x_i\right) = \sum_{i=1}^m \beta_i^2\, \mathbb{V}(x_i) \\
&= \sum_{i=1}^m \frac{\sigma_u^2}{2mf_i(1-f_i)} 2f_i(1-f_i) \\
&= \sigma_u^2
\end{aligned}
$$

776 In an admixed population,

$$
\begin{aligned}
V_g &= \sum_{i=1}^m \beta_i^2 \{2f_i(1-f_i) + 2\,\mathbb{V}(\theta)(f_i^A - f_i^B)^2\} \\
&= \sum_{i=1}^m \frac{\sigma_u^2}{2mf_i(1-f_i)} \{2f_i(1-f_i) + 2\,\mathbb{V}(\theta)(f_i^A - f_i^B)^2\} \\
&= \frac{\sigma_u^2}{m} \sum_{i=1}^m \{1 + \mathbb{V}(\theta)\frac{(f_i^A - f_i^B)^2}{f_i(1-f_i)}\} \\
&= \sigma_u^2 + \underbrace{\mathbb{V}(\theta)\frac{\sigma_u^2}{m}\sum_{i=1}^m \frac{(f_i^A - f_i^B)^2}{f_i(1-f_i)}}_{\substack{\text{contribution of population structure}\\\text{to the genic variance}}}
\end{aligned}
$$

777 The HE estimator of V_g is based on the regression of products of (centered) phenotypes $y_k y_l$ for all pairs
778 of individuals $k \neq l$ on the corresponding entries of the GRM (ψ) where $\psi_{kl} = \frac{\sum_{i=1}^m z_{ik} z_{il}}{m}$ and z_{ik} is the
779 centered and scaled genotype of individual k for locus i:

$$
\begin{aligned}
\hat{V}_g &= \frac{\mathbb{C}(y_k y_l, \psi_{kl})}{\mathbb{V}(\psi_{kl})} \\
&= \frac{\mathbb{E}_{kl}(y_k y_l \psi_{kl}) - \mathbb{E}_{kl}(y_k y_l)\,\mathbb{E}_{kl}(\psi_{kl})}{\mathbb{E}_{kl}(\psi_{kl}^2) - \mathbb{E}(\psi_{kl})^2} \\
&= \frac{\mathbb{E}_{kl}(y_k y_l \psi_{kl})}{\mathbb{E}_{kl}(\psi_{kl}^2)}
\end{aligned}
$$

780 Where $\mathbb{E}_{kl}$ represents the expectation over all $k \times l$ pairwise comparisons between individuals. It is
781 simpler to express the HE estimator in terms of the scaled effects $u_i \sim \mathcal{N}(0, \frac{\sigma_u^2}{m})$.

$$\hat{V}_g = \frac{\mathbb{E}_{kl}(y_k y_l \psi_{kl})}{\mathbb{E}_{kl}(\psi_{kl}^2)} = \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^m u_i z_{ik} \sum_{i=1}^m u_i z_{il} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)}$$

$$= \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^m u_i^2 z_{ik} z_{il} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)} + \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^m \sum_{j \neq i} u_i u_j z_{ik} z_{jl} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)}$$

$$= \frac{\mathbb{E}(u_i^2)\,\mathbb{E}_{kl}\left(\sum_{i=1}^m z_{ik} z_{il} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)} + \frac{\mathbb{E}(u_i u_j)\,\mathbb{E}_{kl}\left(\sum_{i=1}^m \sum_{j \neq i} z_{ik} z_{jl} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)}$$

$$= \frac{\mathbb{E}(u_i^2)\,\mathbb{E}_{kl}(m\psi_{kl}^2)}{\mathbb{E}_{kl}(\psi_{kl}^2)} + 0 = \sigma_u^2$$

Where $\mathbb{E}(u_i)$ and $\mathbb{E}(u_i u_j)$ represent expectations over random realizations of effect sizes. Thus, the last

line follows from our assumption that the effect sizes are independent in expectation, i.e., $\mathbb{E}(u_i u_j) = 0$.

Note, that the estimate is still biased since it does not capture the contribution of population structure.

A3.1.2 Scaling by $\mathbb{V}(x_i)$

Next, we consider the case where the genotypes are standardized instead by the sample variance, i.e.,

$z_{kl} = \frac{x_{ik} - 2f_i}{\sqrt{\mathbb{V}(x_i)}}$ such that $\mathbb{V}(z_i) = 1$. We can derive $\mathbb{E}(u_i^2)$ corresponding to this scaling by noting that the

genetic variance is invariant under linear transformations of the genotype [14]:

$$\sum_{i=1}^m \beta_i^2 \, \mathbb{V}(x_i) = \sum_{i=1}^m u_i^2 \, \mathbb{V}(z_i)$$

$$m\,\mathbb{E}(u_i^2) = \sigma_u^2 + \mathbb{V}(\theta)\frac{\sigma_u^2}{m}\sum_{i=1}^m \frac{(f_i^A - f_i^B)^2}{f_i(1 - f_i)}$$

$$\mathbb{E}(u_i^2) = \frac{\sigma_u^2}{m} + \mathbb{V}(\theta)\frac{\sigma_u^2}{m^2}\sum_{i=1}^m \frac{(f_i^A - f_i^B)^2}{f_i(1 - f_i)}$$

789 Then, the HE estimator becomes:

$$\hat{V}_g = m\,\mathbb{E}(u_i^2)$$

$$= m\left(\frac{\sigma_u^2}{m} + \mathbb{V}(\theta)\frac{\sigma_u^2}{m^2}\sum_{i=1}^m \frac{(f_i^A - f_i^B)^2}{f_i(1 - f_i)}\right)$$

$$= \sigma_u^2 + \mathbb{V}(\theta)\frac{\sigma_u^2}{m}\sum_{i=1}^m \frac{(f_i^A - f_i^B)^2}{f_i(1 - f_i)}$$

Which provides an unbiased estimate of the genic variance. It's important to note that even though we

assumed effect sizes under a random-effect model, the above result holds under a fixed-effect model as

long as there is no directional LD. We discuss the implications of directional LD in the following section.

A3.2 Directional LD

Under the standard random-effect model, the effect sizes are assumed to be independent *in expectation*.

We discussed in the main text how certain processes (e.g. selection and assortative mating) can induce

directional LD across causal loci. But directional LD might arise even for neutral traits and under the

random-effects model for any given realization of effects. This can lead to biases both HE and GREML

estimates of V_g, though the direction and reason for bias is different for the two methods. GREML does
not have a closed-form solution so the exact estimand is difficult to derive. Instead, we develop some
intuition based on HE regression.

A3.2.1   Scaling by $\mathbb{V}(x_i)$

To do this, let $\boldsymbol{u}' = [u_1, u_2, ..., u_m]$ represent the vector of a given realization of (fixed) effects correspond-
ing to the standardized genotypes such that each locus contributes equally to σ_u^2, the genic variance, i.e.,
$u_i^2 = \frac{\sigma_u^2}{m}$. Let there be positive LD across loci such that all cross-product terms are $u_i u_j = \frac{\sigma_u^2}{m}$. Then,
the genetic variance explained by all loci is:

$$
\begin{aligned}
V_g &= \sum_{i=1}^{m} u_i^2 \, \mathbb{V}(z_i) + \sum_{j \neq i} u_i u_j \, \mathbb{C}(z_i, z_j) \\
&= \sum_{i=1}^{m} u_i^2 + \sum_{j \neq i} u_i u_j \, \mathbb{C}(z_i, z_j) \\
&= \sigma_u^2 + \frac{\sigma_u^2}{m} \sum_{j \neq i} \mathbb{C}(z_i, z_j)
\end{aligned}
\tag{3}
$$

806 where $\mathbb{C}(z_i, z_j)$ is the LD between the i^{th} and j^{th} loci that ranges from 0 (no LD) to 1 (perfect LD).
807 Thus, the LD contribution to V_g ranges from 0 to $(m-1)\sigma_m^2$. In comparison, the HE estimator is:

$$
\begin{aligned}
\hat{V}_g &= \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^{m} u_i^2 z_{ik} z_{il} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)} + \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^{m} \sum_{j \neq i} u_i u_j z_{ik} z_{jl} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)} \\
&= \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^{m} \frac{\sigma_u^2}{m} z_{ik} z_{il} \sum_{w=1}^{m} z_{wk} z_{wl}/m\right)}{\mathbb{E}_{kl}(\sum_{i=1}^{m} z_{ik} z_{il}/m \sum_{w=1}^{m} z_{wk} z_{wl}/m)} + \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^{m} \sum_{j \neq i} \frac{\sigma_u^2}{m} z_{ik} z_{jl} \sum_{w=1}^{m} z_{wk} z_{wl}/m\right)}{\mathbb{E}_{kl}(\sum_{i=1}^{m} z_{ik} z_{il}/m \sum_{w=1}^{m} z_{wk} z_{wl}/m)} \\
&= \sigma_u^2 + \sigma_u^2 \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^{m} \sum_{j \neq i} z_{ik} z_{jl} z_{wk} z_{wl}\right)}{\mathbb{E}_{kl}(\sum_{i=1}^{m} z_{ik} z_{il} \sum_{w=1}^{m} z_{wk} z_{wl})}
\end{aligned}
\tag{4}
$$

This shows that the bias due to directional LD in the HE estimate of V_g does not come from the genic,
but the LD component. When there is no LD, e.g. if the population has reached equilibrium after
generations of random mating, this component goes to zero and both the estimate and V_g converge to
the same value – the genic variance. The LD component is maximum when the i^{th} and j^{th} loci are
in perfect LD. In this case, i and j are exchangeable and the second term of the estimator reduces
to $(m-1)\sigma_m^2$. Thus, HE regression should give an unbiased estimate of V_g, even in the presence of
directional LD, but only when LD is perfect. For any other value $0 < C(z_i, z_j) < 1$, the estimate is
biased (Fig. A1). An interpretable, analytical derivation of the second term in Eq. 4 is complicated but
we illustrate the bias with simulations below.
For unlinked markers, $\mathbb{C}(z_i, z_j)$ is a function of $4\mathbb{V}(\theta)(f_i^A - f_i^B)(f_j^A - f_j^B)$ (see A2.1). Perfect LD arises
when (i) both $f_i^A - f_i^B = 1$ and $f_j^A - f_j^B = 1$ and (ii) $\mathbb{V}(\theta)$ is maximum, which occurs at the time of
admixture when source populations mix equally, i.e, $\mathbb{E}(\theta) = 0.5$. To generate a range of LD, we simulated
an admixed population ($N = 1,000$) with equal number of individuals from populations A and B. Thus,

821 $4\mathbb{V}(\theta) = 4\mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} = 1$. We simulated genotypes for each individual at 50 'causal' loci where

822 the difference between the frequencies in the source populations, $f_i^A - f_i^B \in [0,1]$ with the condition

823 that $\frac{f_i^A + f_i^B}{2} = 0.5$. We assigned each locus the same effect size (on the variance-standardized scale) of

824 $+1/\sqrt{m}$ summing up to a genic variance of 1. The positive sign ensures positive LD across loci, i.e,

825 all off-diagonal elements of $\boldsymbol{uu}'$ are set to $1/m$. For each simulation, we computed the expected and

826 estimated LD component using the second terms in Eqs. 3 and 4, respectively, and averaged the results

827 over 100 replications.

Figure A1: The behavior of the LD contribution (y-axis) to the genetic variance (red) and the Haseman-Elston regression estimate (blue) as a function of LD, i.e. $\mathbb{C}(z_i, z_j)$ (x-axis). Each point represents the contribution calculated from a random draw of genotypes, given $\mathbb{C}(z_i, z_j) \propto (f_i^A - f_i^B)(f_j^A - f_j^B)$. The red line represents the expected LD contribution and the black dashed line represents the contribution expected in the case of perfect LD.

828 ## A3.2.2   Scaling by LD

829 In the main text, we showed that standardizing the genotypes at a locus by its covariance with other loci

830 accounts for the bias for GREML and HE estimators. More specifically, the 'LD-scaled' genotypes can

831 be written as $\boldsymbol{Z} = (\boldsymbol{X} - 2\boldsymbol{P})\boldsymbol{U}^{-1}$ where $\boldsymbol{P}$ is an $n \times m$ matrix such that all elements of the i^{th} column

832 contain the frequency of the i^{th} SNP and $\boldsymbol{U}$ is the (upper triangular) square root of the LD matrix, i.e,

833 $\boldsymbol{\Sigma} = \boldsymbol{U}'\boldsymbol{U}$. Under this scheme, the standardized genotypes are uncorrelated and therefore, the second

834 term in Eqs. 3 and 4 are zero. This reduces the estimator to the first term, representing the sum of

835 squares of effect sizes, i.e. $\boldsymbol{u}'\boldsymbol{u} = \sum_{i=1}^{m} u_i^2$. The effect sizes corresponding to the LD scaled genotypes

836 are $\boldsymbol{u} = \boldsymbol{U}\boldsymbol{\beta}$ and the sum of squares is:

$$\boldsymbol{u'u} = (\boldsymbol{U\beta})'(\boldsymbol{U\beta}) = \boldsymbol{\beta'U'U\beta} = \boldsymbol{\beta\Sigma\beta} = \sum_{i=1}^{m}\sum_{j=1}^{m}\beta_i\beta_j\,\mathbb{C}(x_i, x_j)$$

Which captures both the genic and LD contributions and therefore, provides an unbiased estimate of V_g.

A4 Genetic variance after correction for individual ancestry

In the main text, we stated that including individual ancestry as a fixed effect in GREML can lead to an underestimate of V_g in the presence of population structure. Mixed effect models deal with fixed effects (ancestry in our case) by projecting them out of the phenotypes, and estimating the residual variance. This is conceptually equivalent to measuring the residual variance of the regression between phenotype and ancestry. As a result, any variance in the phenotype that is explained by ancestry is removed. To understand this quantitatively, it is helpful to decompose V_g into components of variance explained by and variance orthogonal to ancestry:

$$\mathbb{V}(g) = \underbrace{\mathbb{V}\{\mathbb{E}_{\theta}(g|\theta)\}}_{\substack{\text{variance along} \\ \text{ancestry axis}}} + \underbrace{\mathbb{E}_{\theta}\{\mathbb{V}(g|\theta)\}}_{\substack{\text{variance orthogonal} \\ \text{to ancestry axis}}}$$

We can express the residual variance as:

$$
\begin{aligned}
\mathbb{E}_{\theta}\{\mathbb{V}(g|\theta)\} &= \mathbb{E}_{\theta}\{\mathbb{V}(\sum_{i=1}^{M}\beta_i^2 x_i|\theta)\} \\
&= \mathbb{E}_{\theta}\{\sum_{i=1}^{M}\beta_i^2\,\mathbb{V}(x_i|\theta)\} + \mathbb{E}_{\theta}\{\sum_{i\neq j}\beta_i\beta_j\,\mathbb{C}(x_i, x_j|\theta) \\
&= \sum_{i=1}^{M}\beta_i^2\,\mathbb{E}_{\theta}\{\mathbb{V}(x_i|\theta)\} + 0 \\
&= 2\,\mathbb{E}_{\theta}(\theta)\sum_{i=1}^{M}\beta^2\,f_i^{\mathrm{A}}(1 - f_i^{\mathrm{A}}) + 2\{1 - \mathbb{E}_{\theta}(\theta)\}\sum_{i=1}^{M}\beta^2\,f_i^{\mathrm{B}}(1 - f_i^{\mathrm{B}}) \\
&\quad + 2\,\mathbb{E}_{\theta}(\theta)\sum_{i=1}^{M}\beta^2\{1 - \mathbb{E}_{\theta}(\theta)\}(f_i^{\mathrm{A}} - f_i^{\mathrm{B}})^2 - 2\,\mathbb{V}(\theta)\sum_{i=1}^{M}\beta^2(f_i^{\mathrm{A}} - f_i^{\mathrm{B}})^2
\end{aligned}
$$

Note, that this represents the following components of V_g from the main text: (1.1) + (1.2) - (1.3). Note, that (1.3), which is subtracted out, is always positive and depends on $\mathbb{V}(\theta)$. Thus, the residual genetic variance will be underestimated, regardless of trait architecture, in the presence of population structure, i.e. when $\mathbb{V}(\theta) > 0$.

A5 Effect size of local ancestry

We define local ancestry $\gamma_i \in \{0, 1, 2\}$ as the number of alleles at locus i that trace their ancestry to population A. Thus, the local ancestry at locus i in individual k is a Binomial random variable with

854 $\mathbb{E}(\gamma_{i,k}) = 2\theta_k$. We define the ancestry value of an individual as the weighted sum of their local ancestry:

855 $\sum_{i=1}^{m} \phi_i \gamma_i$ where $\phi_i = \beta_i(f_i^B - f_i^A)$.

856   To show this, note that $\phi = \mathbb{E}(y|\gamma = 1) - \mathbb{E}(y|\gamma = 0)$ where $\mathbb{E}(y|\gamma = 1) = \int_{-\infty}^{\infty} yh(y|\gamma = 1)$ and h is

857   a density function. Our goal is to express ϕ in terms of β, which is equal to $\mathbb{E}(y|x = 1) - \mathbb{E}(y|x = 0)$.

858   Furthermore, $\mathbb{E}(y|x = 1) = \int_{-\infty}^{\infty} yh(y|x = 1)$. We can express $h(y|\gamma)$ in terms of $h(y|x)$ as follows:

$$
\begin{aligned}
h(y|\gamma = 1) = \; & h(y|x = 0)\,\mathbb{P}(x = 0|\gamma = 1) + h(y|x = 1)\,\mathbb{P}(x = 1|\gamma = 1) + h(y|x = 2)\,\mathbb{P}(x = 2|\gamma = 1) \\
= \; & h(y|x = 0)2(1 - f^A)(1 - f^A) + h(y|x = 1)\{f^A(1 - f^B) + f^B(1 - f^A)\} + h(y|x = 2)2\,f^A\,f^B
\end{aligned}
$$

$$
\begin{aligned}
\mathbb{E}(y|\gamma = 1) = & \int_{-\infty}^{\infty} yh(y|\gamma = 1)dy \\
= & (1 - f^A)(1 - f^B) \int_{-\infty}^{\infty} yh(y|x = 0)dy \\
& + \{f^A(1 - f^B) + f^B(1 - f^A)\} \int_{-\infty}^{\infty} yh(y|x = 1)dy \\
& + f^A f^B \int_{-\infty}^{\infty} yh(y|x = 2)dy \\
= & (1 - f^A)(1 - f^B)\,\mathbb{E}(y|x = 0) + \{f^A(1 - f^B) + f^B(1 - f^A)\}\,\mathbb{E}(y|x = 1) + f^A f^B\,\mathbb{E}(y|x = 2) \\
= & 0 + \{f^A(1 - f^B) + f^B(1 - f^A)\}\beta + f^A f^B\,2\beta \\
= & \beta\,f^A + \beta\,f^B
\end{aligned}
$$

859   Similary, $\mathbb{E}(y|\gamma = 0) = 2\beta\,f^B$ and $\phi = \mathbb{E}(y|\gamma = 1) - \mathbb{E}(y|\gamma = 0) = \beta(f^B - f^A)$

860   # A6   Genetic variance due to local ancestry

$$
\begin{aligned}
V_\gamma = \; & \mathbb{V}\left(\sum_{i=1}^{m} \phi_i \gamma_i\right) \\
= & \sum_{i=1}^{m} \phi_i^2\,\mathbb{V}(\gamma_i) + \sum_{i=1}^{m} \sum_{j \neq i}^{m} \phi_i \phi_j\,\mathbb{C}(\gamma_i, \gamma_j)
\end{aligned}
\tag{5}
$$

861   We use the law of total variance and covariance to derive $\mathbb{V}(\gamma_i)$ and $\mathbb{C}(\gamma_i, \gamma_j)$:

$$
\begin{aligned}
\mathbb{V}(\gamma_i) = & \mathbb{E}\{\mathbb{V}(\gamma_i|\theta)\} + \mathbb{V}\{\mathbb{E}(\gamma_i|\theta)\} \\
= & \mathbb{E}\{2\theta(1 - \theta)\} + \mathbb{V}(2\theta) \\
= & 2\,\mathbb{E}(\theta) - 2\,\mathbb{E}(\theta^2) + 4\,\mathbb{V}(\theta) \\
= & 2\,\mathbb{E}(\theta) - 2\,\mathbb{V}(\theta) - 2\,\mathbb{E}(\theta)^2 + 4\,\mathbb{V}(\theta) \\
= & 2\,\mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} + 2\,\mathbb{V}(\theta)
\end{aligned}
$$

$$
\begin{aligned}
\mathbb{C}(\gamma_i, \gamma_j) = & \mathbb{E}\{\mathbb{C}(\gamma_i, \gamma_j|\theta)\} + \mathbb{C}\{\mathbb{E}(\gamma_i, \gamma_j|\theta)\} \\
= & 0 + \mathbb{C}(2\theta, 2\theta) = 4\,\mathbb{V}(\theta)
\end{aligned}
$$

$$
\begin{aligned}
V_\gamma = &\, 2\,\mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} \sum_{i=1}^{m} \phi_i^2 + 2\,\mathbb{V}(\theta) \sum_{i=1}^{m} \phi_i^2 + 4\,\mathbb{V}(\theta) \sum_{i=1}^{m} \sum_{j \neq i} \phi_i \phi_j \\
= &\, 2\,\mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} \sum_{i=1}^{m} \beta_i^2 (f_{\mathrm{i}}^{\mathrm{B}} - f_{\mathrm{i}}^{\mathrm{A}})^2 \\
&+ 2\,\mathbb{V}(\theta) \sum_{i=1}^{m} \beta_i^2 (f_{\mathrm{i}}^{\mathrm{B}} - f_{\mathrm{i}}^{\mathrm{A}})^2 \\
&+ 4\,\mathbb{V}(\theta) \sum_{i=1}^{m} \sum_{j \neq i} \beta_i \beta_j (f_{\mathrm{i}}^{\mathrm{B}} - f_{\mathrm{i}}^{\mathrm{A}})(f_{\mathrm{j}}^{\mathrm{B}} - f_{\mathrm{j}}^{\mathrm{A}})
\end{aligned}
$$

Figure S1: The behavior of the four components of genetic variance in admixed populations under the (A) HI and (B) CGF models. We assume that the mean ancestry proportion in the population is 0.5. The solid lines represent values observed in simulations averaged across ten replicates and the dotted lines represent the expected values based on Eq. 1 of the main text. The red and blue lines represent values for traits 1 and 2, respectively. P indicates the strength of assortative mating. P=0.6 is missing for simulations run for 50 and 100 generations and $\theta \in \{0.1, 0.2\}$ due to the difficulty in finding mate pairs (Methods).

Figure S2: The behavior of the genetic variance due to local ancestry in admixed populations under the (A) HI and (B) CGF models. The solid lines represent values observed in simulations averaged across ten replicates and the dotted lines represent the expected values based on Eq. 1 of the main text. The red and blue lines represent values for traits 1 and 2, respectively. P indicates the strength of assortative mating. P=0.6 is missing for simulations run for 50 and 100 generations and $\theta \in \{0.1, 0.2\}$ due to the difficulty in finding mate pairs (Methods).

Figure S3: The behavior of GREML estimates of SNP heritability ($\hat{h}^2_{snp}$) in admixed populations under the HI (left column) and CGF (right column) models either without (A-C) or with (D-F) individual ancestry as a fixed effect. The solid lines represent $\hat{h}^2_{snp}$ averaged across ten replicates, with red and blue colors representing estimates for traits under divergent and stabilizing selection, respectively. (A, D) show the behavior of $\hat{h}^2_{snp}$ for the default scaling, (B, E) shows $\hat{h}^2_{snp}$ when the genotype at a locus is scaled by its sample variance ($\mathbb{V}(x)$ scaled), and (C, F) when it is scaled by the sample covariance (LD scaled). The shaded area represents the 95% confidence bands generated by bootstrapping (sampling with replacement 100 times) the point estimate reported by GCTA. The black dotted lines represent the expected heritability value given the simulation settings ($h^2 = 0.8$). P indicates the strength of assortative mating

Figure S4: Distribution of the total genetic variance (left), genic variance (middle), and LD component (right) for a neutral trait simulated by drawing effects for 10, 100, or 1,000 causal variants in ASW. The total genetic variance is the sum of the genic and LD components.

Figure S5: The effect of directional LD on Haseman-Elston estimate of genetic variance (V_g). Each individual point is an independent simulation where the effects were drawn from a normal distribution and applied to genotypes from an admixed population (Methods). The solid red line shows the $y = x$ line and the color of each point represents the contribution of LD to V_g.

Figure S6: The LD contribution to the variance explained by variant pairs for (A) neutrophil counts (NEU), (B) white blood count (WBC), (C) monocyte count (MON), and (D) mean corpuscular hemoglobin (MCH). Only chromosomes where we suspected there was a disproportionate contribution to the variance explained are shown.

Figure S7: Expected variance explained estimated using Equation 1 (x-axis) vs the observed variance in polygenic scores (y-axis) in ASW are strongly correlated. Confidence intervals were generated by non-parametric bootstrap (Methods).