Interpreting SNP heritability in admixed populations

Jinguo Huang[*1,2], Nicole Kleman[*3], Saonli Basu[4], Mark D. Shriver[2], and

Arslan A. Zaidi[†3,5]

[1]Bioinformatics and Genomics, Huck Institutes of the Life Sciences, Pennsylvania State University

[2]Department of Anthropology, Pennsylvania State University

[3]Department of Genetics, Cell Biology, and Development, University of Minnesota

[4]Department of Biostatistics, University of Minnesota

[5]Institute of Health Informatics, University of Minnesota

[*]Contributed equally

[†]Correspondence to A.A.Z (aazaidi@umn.edu)

March 11, 2025

Abstract

SNP heritability (h^2_{snp}) is defined as the proportion of phenotypic variance explained by genotyped SNPs and is believed to be a lower bound of heritability (h^2), being equal to it if all causal variants are genotyped. Despite the simple intuition behind h^2_{snp}, its interpretation and equivalence to h^2 is unclear, particularly in the presence of admixture and assortative mating. Here we use analytical theory and simulations to describe the behavior of h^2 and three widely used random-effect estimators of h^2_{snp} – Genome-wide restricted maximum likelihood (GREML), Haseman-Elston (HE) regression, and LD score regression (LDSC) – in admixed populations. We show that h^2_{snp} estimates can be biased in admixed populations, even if all causal variants are genotyped and in the absence of confounding due to shared environment. This is largely because admixture generates directional LD, which contributes to the genetic variance, and therefore to heritability. Random-effect estimators of h^2_{snp}, because they assume that SNP effects are independent, do not capture the contribution, which can be positive or negative depending on the genetic architecture, leading to under- or over-estimates of h^2_{snp} relative to h^2. For the same reason, estimates of local ancestry heritability ($\hat{h}^2_\gamma$) are also biased in the presence of directional LD. We describe this bias in $\hat{h}^2_{snp}$ and $\hat{h}^2_\gamma$ as a function of admixture history and the genetic architecture of the trait, clarifying their interpretation and implication for genome-wide association studies and polygenic prediction in admixed populations.

Introduction

The ability to estimate (narrow-sense) heritability (h^2) from unrelated individuals was a major advance in genetics. Traditionally, h^2 was estimated from family-based studies in which the phenotypic resemblance between relatives could be modeled as a function of their expected genetic relatedness [1]. However, this approach was limited to analysis of closely related individuals where pedigree information is available and the realized genetic relatedness is not too different from expectation [2]. With the advent of genome-wide association studies (GWAS), we hoped that many of the variants underlying this heritability would be uncovered. However, when genome-wide significant single nucleotide polymorphisms (SNPs) explained a much smaller fraction of the phenotypic variance, it became important to explain the missing heritability – were family-based estimates inflated or were GWAS just underpowered, limited by variant discovery?

Yang *et al.* (2010) [3] made the key insight that one could estimate the portion of h^2 tagged by genotyped SNPs, regardless of whether or not they were genome-wide significant, by exploiting the subtle variation in the realized genetic relatedness among apparently unrelated individuals [3–5]. This quantity came to be known colloquially as 'SNP heritability' (h^2_{snp}) and it is believed to be equal to h^2 if all causal variants are included among genotyped SNPs [3]. Indeed, estimates of h^2_{snp} explain a much larger fraction of trait heritability than GWAS SNPs [3], approaching family-based estimates of h^2 when whole genome sequence data, which captures rare variants, are used [6]. This has made it clear that GWAS have yet to uncover more variants with increasing sample size. Now, h^2_{snp} has become an important aspect of the design of genetic studies and is often used to define the power of variant discovery in GWAS and the upper limit of polygenic prediction accuracy.

Despite the utility and simple intuition of h^2_{snp}, there is much confusion about its interpretation and equivalence to h^2, particularly in the presence of population structure and assortative mating [7–12]. But much of the discussion of heritability in structured populations has focused on biases in $\hat{h}^2_{snp}$ – the estimator – due to confounding effects of shared environment and linkage disequilibrium (LD) with other variants [7, 9–11, 13]. There is comparatively little discussion, at least in human genetics, on the fact that LD due to population structure also contributes to genetic variance, and therefore, is a component of heritability [1, 14, 15]. We think this is at least partly due to the fact that most studies are carried out in cohorts with primarily European ancestry, where the degree of population structure is minimal and large effects of LD can be ignored. However, that is not the case for diverse, multi-ethnic cohorts, which have historically been underrepresented in genetic studies, but thanks to a concerted effort in the field, are now becoming increasingly common [16–22]. The complex structure in these cohorts also brings unique methodological challenges and it is imperative that we understand whether existing methods, which have largely been evaluated in more homogeneous groups, generalize to more diverse cohorts.

Our goal in this paper is to study the behavior of h^2 and $\hat{h}^2_{snp}$ in admixed populations. How should we interpret $\hat{h}^2_{snp}$ in the ideal situation where causal variants are directly genotyped? Is it an unbiased estimate of h^2? To answer these questions, we derived a general expression for the genetic variance in admixed populations, decomposing it in terms of the contribution of population structure, which influences both the genotypic variance at individual loci and the LD across loci. We used theory and simulations to show that $\hat{h}^2_{snp}$ estimated with genome-wide restricted maximum likelihood (GREML)

68 [3, 5], linkage disequilibrium score regression (LDSC) [23], and Haseman-Elston (HE) regression [24] –

69 three widely used approaches – can be biased in admixed and other structured populations, even in the

70 absence of confounding and when all causal variants are genotyped. We explain this in terms of the

71 discrepancy between the model assumed in $\hat{h}^2_{snp}$ estimation and the generative model from which the

72 genetic architecture of the trait in the population may have been sampled. We describe the bias in $\hat{h}^2_{snp}$

73 as a function of admixture history and genetic architecture and discuss its implications for GWAS and

74 polygenic prediction accuracy.

Model

Genetic architecture

77 We begin by describing a generative model for the phenotype. Let $y = g + e$, where y is the phenotypic

78 value of an individual, g is the genotypic value, and e is random error. We assume additive effects such

79 that $g = \sum_{i=1}^{m} \beta_i x_i$ where β_i is the effect size of the i^{th} biallelic locus (out of a total m causal loci) and

80 $x_i \in \{0, 1, 2\}$ is the number of copies of the trait-increasing allele. Importantly, the effect sizes are fixed

81 quantities and differences in genetic values among individuals are due to random variation in genotypes.

82 Note, that this is different from the random-effects model assumed by many heritability estimators where

83 genotypes are fixed and effect sizes are random [25].

84 We denote the mean, variance, and covariance with $\mathbb{E}(.)$, $\mathbb{V}(.)$, and $\mathbb{C}(.)$, respectively, where the

85 expectation is measured over random draws from the population rather than random realizations of the

86 evolutionary process. We can express the additive genetic variance of a quantitative trait as follows:

$$V_g = \mathbb{V}(\sum_{i=1}^{m} \beta_i x_i) = \sum_{i=1}^{m} \beta_i^2 \, \mathbb{V}(x_i) + \sum_{j \neq i} \beta_i \beta_j \, \mathbb{C}(x_i, x_j)$$

87 Here, the first term represents the contribution of individual loci (genic variance) and the second term

88 is the contribution of linkage disequilibrium (LD contribution). We make the assumption that loci are

89 unlinked and therefore, the LD contribution is entirely due to population structure. We describe the

90 behavior of V_g in a population that is a mixture of two previously isolated populations A and B that

91 diverged from a common ancestor. To do this, we denote θ as the fraction of the genome of an individual

92 with ancestry from population A. Thus, $\theta = 1$ if the individual is from population A, 0 if they are from

93 population B, and $\theta \in (0, 1)$ if they are admixed. Then, V_g can be expressed in terms of ancestry as

94 (Appendix):

$$V_g = 2\,\mathbb{E}(\theta)\sum_{i=1}^{m}\beta_i^2\,f_i^{\mathrm{A}}(1-f_i^{\mathrm{A}}) + 2\{1-\mathbb{E}(\theta)\}\sum_{i=1}^{m}\beta_i^2\,f_i^{\mathrm{B}}(1-f_i^{\mathrm{B}}) \tag{1.1}$$

$$+ 2\,\mathbb{E}(\theta)\{1-\mathbb{E}(\theta)\}\sum_{i=1}^{m}\beta_i^2(f_i^{\mathrm{A}}-f_i^{\mathrm{B}})^2 \tag{1.2}$$

$$+ 2\,\mathbb{V}(\theta)\sum_{i=1}^{m}\beta_i^2(f_i^{\mathrm{A}}-f_i^{\mathrm{B}})^2 \tag{1.3}$$

$$+ 4\,\mathbb{V}(\theta)\sum_{i\neq j}\beta_i\beta_j(f_i^{\mathrm{A}}-f_i^{\mathrm{B}})(f_j^{\mathrm{A}}-f_j^{\mathrm{B}}) \tag{1.4}$$

95 where f_i^{A} and f_i^{B} are the allele frequencies in populations A and B, and $\mathbb{E}(\theta)$ and $\mathbb{V}(\theta)$ are the mean
96 and variance of individual ancestry. The sum of the first three terms represents the genic variance and
97 the last term represents the LD contribution.

Demographic history

From Eq. 1, it is clear that, conditional on the genetic architecture in the source populations (β, f^A, f^B),
V_g is a function of the mean, $\mathbb{E}(\theta)$, and variance, $\mathbb{V}(\theta)$, of individual ancestry in the admixed population.
We consider two demographic models that affect $\mathbb{E}(\theta)$ and $\mathbb{V}(\theta)$ in qualitatively different ways. In the
first model, the source populations meet once t generations ago (we refer to this as $t=0$) in proportions
p and $1-p$, after which there is no subsequent admixture (Fig. 1A). In the second model, there is
continued gene flow in every generation from one of the source populations such that the mean overall
amount of ancestry from population A is the same as in the first model (Fig. 1A). For brevity, we refer
to these as the hybrid-isolation (HI) and continuous gene flow (CGF) models, respectively, following
Pfaff *et al.* (2001) [26]. $\mathbb{V}(\theta)$ is also affected by ancestry-based assortative mating, where individuals are
more likely to partner with others of similar ancestry. We refer to this simply as assortative mating for
brevity and model this following Zaitlen *et al.* (2017) using a parameter $P \in (0,1)$, which represents the
correlation of the ancestry of individuals across mating pairs in the population [27].

Under these conditions, the behavior of $\mathbb{E}(\theta)$ and $\mathbb{V}(\theta)$ has been described previously [27, 28] (Fig. 1B
and C). Briefly, in the HI model, $\mathbb{E}(\theta)$ remains constant at p in the generations after admixture as there
is no subsequent gene flow. $\mathbb{V}(\theta)$ is at its maximum at $t=0$ when each individuals carries chromosomes
either from population A or B, but not both. This genome-wide correlation in ancestry breaks down
in subsequent generations as a function of mating, independent assortment, and recombination, leading
to a decay in $\mathbb{V}(\theta)$, the rate depending on the strength of assortative mating (Fig. 1C). In the CGF
model, both $\mathbb{E}(\theta)$ and $\mathbb{V}(\theta)$ increase with time as new chromosomes are introduced from the source
populations. But while $\mathbb{E}(\theta)$ continues to increase monotonically, $\mathbb{V}(\theta)$ will plateau and decrease due
to the countervailing effects of independent assortment and recombination which redistribute ancestry
in the population, approaching zero at equilibrium if there is no more gene flow and the population
is mating randomly. $\mathbb{V}(\theta)$ provides an intuitive and quantitative measure of the degree of population
structure (along the axis of ancestry) in admixed populations.

Figure 1: The behavior of mean and variance of individual ancestry as a function of admixture history. (A) Shows the demographic models under which simulations were carried out. Admixture might occur once (Hybrid Isolation, HI, left column) or continuously (Continuous Gene Flow, CGF, right column). (B) The mean individual ancestry, $\mathbb{E}(\theta)$ remains constant over time in the HI model and increases in the CGF model with continued gene flow. (C) The variance in individual ancestry, $\mathbb{V}(\theta)$ is maximum at $t = 0$ in the HI model, decaying subsequently. $\mathbb{V}(\theta)$ increases with gene flow in the CGF model and will subsequently decrease with time. P measures the strength of assortative mating, which slows the decay of $\mathbb{V}(\theta)$. P=0.6 is missing for simulations run for 50 and 100 generations and $\theta \in \{0.1, 0.2\}$ due to the difficulty in finding mate pairs (Methods).

123 Results

124 Genetic variance in admixed populations

To understand the expectation of genetic variance in admixed populations, it is first worth discussing

its behavior in the source populations. In Eq. 1, the first term represents the within-population com-

ponent (V_{gw}) and the last three terms altogether represent the component of genetic variance between

populations A and B (V_{gb}). Note that $V_{gb} = \frac{(\bar{g}_A - \bar{g}_B)^2}{2}$ is positive only if there is a difference in the

mean genotypic values (Fig. 2). This variance increases with increasing F_{ST}, i.e., genetic divergence,

between the two populations. While $\beta_i^2 (f_i^A - f_i^B)^2$ is expected to increase monotonically with F_{ST},

$\beta_i \beta_j (f_i^A - f_i^B)(f_j^A - f_j^B)$ is expected to be zero under neutrality because the direction of frequency change

will be uncorrelated across loci. In this case, the LD contribution, i.e., (1.4), is expected to be zero and

$V_{gb} = (1.1) + (1.2) + (1.3)$. However, this is true only in expectation over the evolutionary process and

the realized LD contribution may be non-zero even for neutral traits.

Figure 2: Decomposing genetic variance in a two-population system. The plot illustrates the expected distribution of genetic values in two populations under different selective pressures and the terms on the right list the total (V_g) and between-population genetic variance (V_{gb}) expected over the evolutionary process. For neutrally evolving traits (top row), we expect there to be an absolute difference in the mean genetic values ($|\bar{g}_A - \bar{g}_B|$) that is proportional to F_{ST}. For traits under divergent selection (middle), $|\bar{g}_A - \bar{g}_B|$ is expected to be greater than that expected under genetic drift. For traits under stabilizing selection, $|\bar{g}_A - \bar{g}_B|$ will be less than that expected under genetic drift, and zero in the extreme case.

135 For traits under selection, the LD contribution is expected to be greater or less than zero, depending

136 on the type of selection. Under divergent selection, trait-increasing alleles will be systematically more

137 frequent in one population over the other, inducing positive LD across loci [29, 30], increasing the

138 LD contribution, i.e., term (1.4). Stabilizing selection, on the other hand, induces negative LD [31,

139 32]. In the extreme case, the mean genetic values of the two populations are exactly equal and $V_{gb} =$

140 $(1.2) + (1.3) + (1.4) = 0$. For this to be true, (1.4) has to be negative and equal to $(1.2) + (1.3)$, which

141 are both positive, and the total genetic variance is reduced to the within-population variance, i.e., term

142 (1.1) (Fig. 2). This is relevant because, as we show in the following sections, the behavior of the genetic

143 variance in admixed populations depends on the magnitude of V_{gb} between the source populations.

144   We illustrate this by tracking the genetic variance in admixed populations for two traits, both with

145 the same mean F_{ST} at causal loci but with different LD contributions (term 1.4): one where the LD

146 contribution is positive (Trait 1) and the other where it is negative (Trait 2). Thus, traits 1 and 2 can

147 be thought of as examples of phenotypes under divergent and stabilizing selection, respectively, and we

148 refer to them as such from hereon. To simulate the genetic variance of such traits, we drew the allele

149 frequencies (f^A and f^B) in populations A and B for 1,000 causal loci with $F_{ST} \approx 0.2$ using the Balding-

150 Nichols model [33]. We assigned effect sizes such that each locus contributes equally to the genetic

151 variance, i.e., $\beta = \frac{1}{\sqrt{2m\bar{f}(1-\bar{f})}}$ where $\bar{f}$ is the mean allele frequency between the two populations, and m

152 is the number of loci. To simulate positive and negative LD, we permuted the effect signs across variants

153 100 times and selected the combinations that gave the most positive and negative LD contribution to

154 represent the genetic architecture of traits that might be under directional (Trait 1) and stabilizing (Trait

155 2) selection, respectively (Methods). We simulated the genotypes of 10,000 individuals under the HI and

156 CGF models for $t \in \{10, 20, 50, 100\}$ generations post-admixture and calculated genetic values for both

157 traits using $g = \sum_{i=1}^{m} \beta_i x_i$, where $m = 1,000$ (Method). The observed genetic variance at any time can

158 then be calculated simply as the variance in genetic values, i.e. $V_g = \mathbb{V}(g)$.

159   In the HI model, $\mathbb{E}(\theta)$ does not change (Fig. 1B) so terms (1.1) and (1.2) are constant through time.

160 Terms (1.3) and (1.4) decay towards zero as the variance in ancestry goes to zero and V_g ultimately

161 converges to $(1.1) + (1.2)$ (Fig. 3). This equilibrium value is equal to the $\mathbb{E}(V_g|\theta)$ (Appendix) and the

162 rate of convergence depends on the strength of assortative mating, which slows the rate at which $\mathbb{V}(\theta)$

163 decays. V_g approaches equilibrium from a higher value for traits under divergent selection and lower value

164 for traits under stabilizing selection because of positive and negative LD contributions, respectively, at

165 $t = 0$ (Fig. 3). In the CGF model, V_g increases initially for both traits with increasing gene flow (Fig.

166 3). This might seem counter-intuitive at first because gene flow increases admixture LD, which leads

167 to more negative values of the LD contribution for traits under stabilizing selection (Fig. S1). But this

168 is outweighed by positive contributions from the genic variance – terms $(1.1) + (1.2) + (1.3)$ – all of

169 which initially increase with gene flow (Fig. S1). After a certain point, the increase in V_g slows down as

170 any increase in $\mathbb{V}(\theta)$ due to gene flow is counterbalanced by recombination and independent assortment.

171 Ultimately, V_g will decrease if there is no more gene flow, reaching the same equilibrium value as in the

172 HI model, i.e., $\mathbb{E}(V_g|\theta) = (1.1) + (1.2)$. Because the loci are unlinked, we refer to the sum $(1.3) + (1.4)$

173 as the contribution of population structure.

Figure 3: Genetic variance in admixed populations under the (A) HI and (B) CGF models. Dotted lines represent the expected genetic variance based on Eq. (1) and solid lines represent results of simulations averaged over ten replicates. Red and blue lines represent traits under divergent and stabilizing selection, respectively. $P = 0.6$ is missing for simulations run for 50 and 100 generations and $\theta \in \{0.1, 0.2\}$ due to the difficulty in finding mate pairs (Methods).

GREML estimation

In their original paper, Yang *et al.* (2010) defined h_{snp}^2 as the variance explained by genotyped SNPs and not as heritability [3]. This is because h^2 is the genetic variance explained by causal variants, which are unknown. Genotyped SNPs may not overlap with or tag all causal variants and thus, h_{snp}^2 is understood to be a lower bound of h^2, both being equal if causal variants are directly genotyped [3]. Our goal is to demonstrate (i) that this may not be true in structured populations, (ii) quantify the bias in $\hat{h}_{snp}^2$, and (iii) understand the source and behavior of this bias even in the ideal situation when causal variants are genotyped.

We first used GREML, implemented in GCTA [3, 5], to estimate the genetic variance for our simulated traits. GCTA assumes the following model: $\boldsymbol{y} = \boldsymbol{Zu} + \boldsymbol{\epsilon}$ where $\boldsymbol{Z}$ is an $n \times m$ standardized genotype matrix such that the genotype of the k^{th} individual at the i^{th} locus is $z_{ik} = \frac{x_{ik} - 2f_i}{\sqrt{2f_i(1-f_i)}}$, f_i being the allele frequency. The SNP effects corresponding to the scaled genotypes are assumed to be random and independent such that $\boldsymbol{u} \sim \mathcal{N}(0, \boldsymbol{I}\frac{\sigma_u^2}{m})$ and $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \boldsymbol{I}\sigma_\epsilon^2)$ is random environmental error. Then, the phenotypic variance can be decomposed as:

$$\mathbb{V}(\boldsymbol{y}) = \mathbb{V}(\boldsymbol{Zu}) + \mathbb{V}(e)$$
$$= \frac{\boldsymbol{ZZ}'}{m} \sigma_u^2 + \boldsymbol{I}\sigma_\epsilon^2$$

where $\frac{\boldsymbol{ZZ}'}{m}$ is the genetic relationship matrix (GRM), the variance components σ_u^2 and σ_ϵ^2 are estimated using restricted maximum likelihood, and $\hat{h}_{snp}^2$ is calculated as $\frac{\hat{\sigma}_u^2}{\hat{\sigma}_u^2 + \hat{\sigma}_\epsilon^2}$. We are interested in asking whether $\hat{\sigma}_u^2$ is an unbiased estimate of V_g. To answer this, we constructed the GRM with causal variants and estimated $\hat{\sigma}_u^2$ using GCTA [3, 4].

GCTA under- and over-estimates the genetic variance in admixed populations for traits under divergent (Trait 1) and stabilizing selection (Trait 2), respectively, when there is population structure, i.e., when $\mathbb{V}(\theta) > 0$ (Fig. 4A). One reason for this bias is that the GREML model assumes that the effects are independent, and therefore the LD contribution is zero. This, as discussed in the previous section, is not true for traits under divergent or stabilizing selection between the source populations, and only true for neutral traits in expectation. Because of this, $\hat{\sigma}_u^2$ does not capture the LD contribution, i.e. term (1.4) (Fig. 4A). But $\hat{\sigma}_u^2$ can be biased even if the LD contribution is zero if the genotypes are scaled with $\sqrt{2f_i(1-f_i)}$ – the standard practice – where f_i is the frequency of the allele in the population. This scaling assumes that $\mathbb{V}(x_i) = 2f_i(1-f_i)$, which is true only if the population were mating randomly. In an admixed population $\mathbb{V}(x_i) = 2f_i(1-f_i) + 2\mathbb{V}(\theta)(f_i^A - f_i^B)^2$, where f_i, f_i^A, and f_i^B correspond to frequency in the admixed population, and source populations, A and B, respectively (Appendix). Alternatively, if the genotypes are scaled, $\mathbb{V}(z_i) = 1 + 2\mathbb{V}(\theta)F_{st}^{(i)}$ where $F_{st}^{(i)}$ is the F_{st} at the i^{th} locus. We show that this assumption biases $\hat{\sigma}_u^2$ downwards by a factor of $2\mathbb{V}(\theta)(f_i^A - f_i^B)^2$ (or $2\mathbb{V}(\theta)F_{st}^{(i)}$ if genotypes are scaled) – term (1.3) (Fig. 4B, Appendix). Thus, with the standard scaling, $\hat{\sigma}_u^2$ gives a biased estimate in the presence of population structure, even of the genic variance.

The overall bias in $\hat{\sigma}_u^2$ is determined by the relative magnitude and direction of terms (1.3) and (1.4), both of which are functions of $\mathbb{V}(\theta)$, and therefore, of the degree of structure in the population.

Figure 4: The behavior of GREML estimates of the genetic variance ($\hat{\sigma}_u^2$) in admixed populations under the HI (left column) and CGF (right column) models either without (A-D) or with (E-H) individual ancestry as a fixed effect. The solid lines represent estimates from simulated data averaged across ten replicates with red and blue colors representing estimates for traits under divergent and stabilizing selection, respectively. P indicates the strength of assortative mating. The shaded area represents the 95% confidence bands generated by bootstrapping (sampling with replacement 100 times) the point estimate reported by GCTA. The dotted lines either represent the expected variance in the population based on Eq. 1 (A & B) or the expected estimate for three different ways of scaling genotypes (B-D & F-H). (A-B & E-F) show the behavior of $\hat{\sigma}_u^2$ for the default scaling, (C, G) shows $\hat{\sigma}_u^2$ when the genotype at a locus is scaled by its sample variance ($\mathbb{V}(x)$ scaled), and (D, H) when it is scaled by the sample covariance (LD scaled).

209 The contribution of term (1.3) will be modest, even in highly structured populations (Fig. S1) and

210 therefore, the overall bias is largely driven by the LD contribution. If there is no more gene flow, $\mathbb{V}(\theta)$

211 will ultimately go to zero and V_g will converge towards $\hat{\sigma}_u^2$. Thus, $\hat{\sigma}_u^2$ is more accurately interpreted as

212 the genic variance or the genetic variance expected if the LD contribution were zero and if the population

213 were mating randomly. In other words, $\mathbb{E}(\hat{\sigma}_u^2) = (1.1) + (1.2) \neq V_g$ (Fig. 4B).

214 In principle, we can recover the missing components of V_g by scaling the genotypes appropriately.

215 For example, we can recover term (1.3) by scaling the genotype at each variant i by its sample variance,

216 i.e., $z_{ik} = \frac{x_{ik} - 2f_i}{\sqrt{\mathbb{V}(x_i)}}$ (Fig. 4C) (Appendix). We can also recover term (1.4) by scaling the genotypes with

217 the covariance between SNPs, i.e., the LD matrix, as previously proposed [34, 35] (Methods). In matrix

218 form, the 'LD-scaled' genotypes can be written as $\boldsymbol{Z} = (\boldsymbol{X} - 2\boldsymbol{P})\boldsymbol{U}^{-1}$ where $\boldsymbol{P}$ is an $n \times m$ matrix such

219 that all elements of the i^{th} column contain the frequency of the i^{th} SNP and $\boldsymbol{U}$ is the (upper triangular)

220 square root matrix of the LD matrix, i.e., $\boldsymbol{\Sigma} = \boldsymbol{U}'\boldsymbol{U}$ [34]. GREML recovers the LD contribution under

221 this scaling, resulting in unbiased estimates of V_g for both traits (Fig. 4D, Appendix).

222 In practice, however, the LD contribution may not be fully recoverable for two reasons. One, the

223 LD-scaled GRM requires computing the inverse of $\boldsymbol{\Sigma}$ or $\boldsymbol{U}$ which may not exist, especially if the number

224 of markers is greater than the sample size – the case for most human genetic studies. Second, it is

225 common to include individual ancestry or principal components of the GRM as fixed effects in the model

226 to account for inflation in heritability estimates due to shared environment. This also has the effect of

227 removing the components of genetic variance along the ancestry axes, the residual variance being equal

228 to $\mathbb{E}\{\mathbb{V}(g|\theta)\} = (1.1) + (1.2) - (1.3)$ (Appendix). Indeed, this is what we observe in Fig. 4H.

HE estimation

230 Haseman-Elston (HE) regression also assumes a random-effects model but uses a method-of-moments

231 approach, as opposed to GREML, which maximizes the likelihood to estimate V_g. Previous work has

232 shown that as long as all causal variants are included in the GRM calculation, the HE estimator will

233 not be biased, even if they are in LD with each other [36]. We show that in the presence of positive

234 and negative LD between causal loci, as exemplified by traits under divergent and stabilizing selection,

235 respectively, the HE estimates of V_g *are* biased upwards and downwards, respectively (Fig. 5A-B). To

236 understand this discrepancy and the source of bias in our simulations, recall that HE estimates V_g from

237 the regression of the (pairwise) phenotypic covariance between individuals on their genotypic covariance

238 [24]. More specifically, if we denote $Y_{kl} = y_k y_l$ as the product of the (centered) phenotypes of k^{th} and

239 l^{th} individuals, and ψ_{kl} as the k^{th} and l^{th} entry of the GRM, then the HE estimator can be written as:

$$\hat{V}_g^{he} = \frac{Cov(Y_{kl}, \psi_{kl})}{Var(\psi_{kl})} = \frac{\mathbb{E}(y_k y_l \sum_{w=1}^M z_{wk} z_{wl})}{\mathbb{E}(\sum_{i=1}^M z_{ik} z_{il} \sum_{w=1}^M z_{wk} z_{wl})}$$

$$= \frac{\mathbb{E}\{(g_k + e_k)(g_l + e_l) \sum_{w=1}^M z_{wk} z_{wl}\}}{\mathbb{E}(\sum_{i=1}^M z_{ik} z_{il} \sum_{w=1}^M z_{wk} z_{wl})} = \frac{\mathbb{E}(g_k g_l \sum_{w=1}^M z_{wk} z_{wl})}{\mathbb{E}(\sum_{i=1}^M z_{ik} z_{il} \sum_{j=1}^M z_{wk} z_{wl})}$$

$$= \frac{\mathbb{E}(\sum_{i=1}^M \sum_{j=1}^M u_i u_j z_{ik} z_{jl} \sum_{w=1}^M z_{wk} z_{wl})}{\mathbb{E}(\sum_{i=1}^M z_{ik} z_{il} \sum_{w=1}^M z_{wk} z_{wl})} = \frac{\mathbb{E}(\sum_{i=1}^M \sum_{j=1}^M u_i u_j \sum_{w=1}^M z_{ik} z_{jl} z_{wk} z_{wl})}{\mathbb{E}(\sum_{i=1}^M \sum_{w=1}^M z_{ik} z_{il} z_{wk} z_{wl})}$$

$$= \underbrace{\frac{\mathbb{E}(\sum_{i=1}^M u_i^2 \sum_{w=1}^M z_{ik} z_{jl} z_{wk} z_{wl})}{\mathbb{E}(\sum_{i=1}^M \sum_{w=1}^M z_{ik} z_{il} z_{wk} z_{wl})}}_{\text{genic component}} + \underbrace{\frac{\mathbb{E}(\sum_{i=1}^M \sum_{j \neq i} u_i u_j \sum_{w=1}^M z_{ik} z_{jl} z_{wk} z_{wl})}{\mathbb{E}(\sum_{i=1}^M \sum_{w=1}^M z_{ik} z_{il} z_{wk} z_{wl})}}_{\text{directional LD}} \qquad (2)$$

240 Where the first and second terms represent the genic and LD components, respectively, of the esti-
241 mate. Population structure induces correlations between the alleles at a given locus as well as across
242 loci (i.e., LD). But the LD may not be directional, i.e., trait-increasing alleles may be as likely to be
243 co-inherited with each other as they are to trait-decreasing alleles, and vice versa – implicit under the
244 standard random-effects model. Thus, in the absence of directional LD, the second term is zero and the
245 first term is unaffected as long as all causal variants are included in the GRM, because the increase in
246 the numerator due to population structure is proportional to the denominator [36]. Directional LD does
247 not affect the first term but exaggerates the contribution from the second term, i.e., the LD component
248 (see Appendix section A4.2). Consequently, HE regression over- and under-estimates V_g for traits with
249 positive and negative LD, respectively. Note that this bias is in the opposite direction of the bias observed
250 with GREML, which fails to capture the LD contribution. Scaling the genotype at a locus by its LD with
251 other loci, as discussed in the previous section, corrects for the bias in HE regression regardless of genetic
252 architecture, yielding estimates consistent with GREML (Fig. 5C). Thus, GREML and HE regression
253 are guaranteed to yield the same estimates only if the underlying model specifying the distribution of
254 effects is consistent with the true architecture of the trait.

255 As with GREML, it is common to account for ancestry in HE regression to correct for any inflation
256 in heritability estimates due to shared environment [11, 37]. We show that this removes the bias from
257 exaggerated LD contributions but also removes any genetic variation along the ancestry axis (Fig. 5D-
258 F, Methods). Thus, ancestry-corrected estimates of V_g should be consistent between GREML and HE
259 regression and should be interpreted as the sum of the contributions of individual loci, i.e., the genic
260 variance.

LD score regression

261

262 LD score regression (LDSC) is different from GREML and HE regression in that it uses GWAS summary
263 statistics instead of individual-level data [23]. Intuitively, LDSC estimates V_g (or h_{snp}^2 if the phenotype
264 is scaled to unit variance) from the slope of a regression of marginal association statistics from GWAS
265 (χ_k^2) on 'LD scores', defined as the sum of squared correlations between a given k^{th} variant and all other
266 variants, i.e. $l_k = \sum_{i=1}^m r_{ik}^2$. To understand the effect of directional LD on LDSC, note that the estimated
267 marginal effect ($\hat{\tau}_k$) of the k^{th} marker is a function of its true marginal effect (τ_k) and some estimation
268 error (ε_k): $\hat{\tau}_k = \tau_k + \varepsilon_k$ assuming for simplicity that there is no residual stratification in the upstream

Figure 5: Genetic variance ($\hat{V}_g$) estimated with HE regression in admixed populations under the HI (left column) and CGF (right column) models either without (A-C) or with (D-F) adjustment for individual ancestry. The solid lines represent estimates from simulated data averaged across ten replicates with red and blue colors representing estimates for traits under divergent and stabilizing selection, respectively. P indicates the strength of assortative mating. (A & D) show the behavior of $\hat{V}_g$ for the default scaling, (B, E) shows $\hat{V}_g$ when the genotype at a locus is scaled by its sample variance ($\mathbb{V}(x)$ scaled), and (C, F) when it is scaled by the sample covariance (LD scaled). The dotted lines in A-E represent the expected V_g in the population based on Eq. 1 and in F, represent the expected V_g after removing any genetic variance along the ancestry axis. The shaded areas represent the 95% bootstrapped confidence bands of the estimate.

269 GWAS. Then, $\mathbb{E}(\hat{\tau}_k) = \tau_k$ and $\mathbb{E}(\hat{\tau}_k^2) = \tau_k^2 + \mathbb{E}(\varepsilon^2) = \tau_k^2 + SE^2(\hat{\tau}_k)$, where $SE(.)$ is the standard error.

270 Furthermore, $\tau_k = \sum_{i=1}^m r_{ik} u_i$, where r_{ik} is the genotypic correlation between the k^{th} marker and i^{th}

271 causal variant [23, 38].

$$\mathbb{E}(\chi_k^2) = \frac{\mathbb{E}(\hat{\tau}_k^2)}{SE^2(\hat{\tau}_k)} = \frac{\tau_k^2 + SE^2(\hat{\tau}_k)}{SE^2(\hat{\tau}_k)}$$

272 If we assume a highly polygenic architecture – as LDSC does – such that the effect of any individual

273 causal variant is small, then $SE^2(\hat{\tau}_k) \approx \frac{1}{n\,\mathbb{V}(z_k)} = 1/n$ and $\mathbb{E}(\chi_k^2) \approx n\tau_k^2 + 1$. We can decompose the

274 LDSC regression slope as follows:

$$
\begin{aligned}
\mathbb{E}(\hat{\beta}_{ldsc}) &= \frac{\mathbb{C}(\chi_k^2, l_k)}{\mathbb{V}(l_k)} \approx \frac{\mathbb{C}(n\tau_k^2 + 1, l_k)}{\mathbb{V}(l_k)} = \frac{\mathbb{C}\{n\left(\sum_{i=1}^m r_{ik} u_i\right)^2, l_k\}}{\mathbb{V}(l_k)} \\
&\approx \frac{n\,\mathbb{C}\left(\sum_{i=1}^m r_{ik}^2 u_i^2 + 2\sum_{i=1}^m \sum_{j<i}^m r_{ik} r_{jk} u_i u_j, l_k\right)}{\mathbb{V}(l_k)} \\
&\approx \underbrace{\frac{n\,\mathbb{C}\left(\sum_{i=1}^m r_{ik}^2 u_i^2, l_k\right)}{\mathbb{V}(l_k)}}_{\text{genic component}} + \underbrace{\frac{n\,\mathbb{C}\left(2\sum_{i=1}^m \sum_{j<i} r_{ik} r_{jk} u_i u_j, l_k\right)}{\mathbb{V}(l_k)}}_{\text{directional LD}}
\end{aligned}
$$

275 Thus, the LDSC slope can also be decomposed into contributions of individual loci (first term) and

276 directional LD (second term). As with GREML and HE regression, LDSC assumes random independent

277 effects where $u_i \sim \mathcal{N}(0, \frac{\sigma_u^2}{m})$ such that the second term is zero over variants i and j, and the slope reduces

278 to:

$$\mathbb{E}(\hat{\beta}_{ldsc}) = \frac{n\,\mathbb{C}\left(\sum_{i=1}^m r_{ik}^2 u_i^2, l_k\right)}{\mathbb{V}(l_k)} = \frac{n\sigma_u^2}{m} \frac{\mathbb{C}\left(\sum_{i=1}^m r_{ik}^2, l_k\right)}{\mathbb{V}(l_k)} = \frac{n\sigma_u^2}{m}$$

279 from which $\hat{V}_g$ can be derived, i.e., $\hat{V}_g = \hat{\sigma}_u^2 = \frac{\hat{\beta}_{ldsc} m}{n}$.

280 In practice, LD scores are computed in a user-defined window around each variant. If the causal

281 variants are far enough apart, i.e., not in physical linkage – implicit in our generative model – such that

282 there is effectively only a single variant per window, or if there is no directional LD within a window,

283 the LDSC estimate of V_g should reflect the sum of the contributions of individual loci, i.e., the genic

284 variance, and therefore, be equivalent to (ancestry-adjusted) GREML and HE regression. As a result,

285 LDSC estimates of V_g, and therefore, of h_{snp}^2, will also be biased downwards and upwards in the presence

286 of positive and negative LD, respectively, across causal variants.

287 To illustrate this, we simulated genotype data on chromosome 2 ($M = 88,112$ SNPs) of individuals

288 of mixed African and European ancestry ($N = 5,000$) under the HI and CGF models using haplotypes

289 from the 1000 Genomes YRI and CEU [39] (Methods). We selected $p \in \{0.01, 0.05, 0.1\}$ proportion of

290 variants uniformly at random to be causal and assigned them effects $\beta_i = \frac{1}{\sqrt{pM2\bar{f}_i(1-\bar{f}_i)}}$, where $\bar{f}_i$ is

291 the mean frequency between YRI and CEU. To simulate directional LD, we assigned '+' or '−' signs

292 to causal effects uniformly at random 1,000 times and selected the combination that gave the largest

293 and smallest V_{gb} between YRI and CEU to represent traits under divergent and stabilizing selection,

294 respectively. We carried out GWAS for the unscaled genetic values in the admixed population at every

generation under the HI and CGF models with 20 genetic principal components (PCs) as covariates. We also projected out 20 PCs from the genotypes in computing LD scores to account for admixture LD with cov-LDSC [40]. Because we used the unscaled genetic values as our GWAS phenotype, the LDSC slope gives an estimate of V_g, the behavior of which is shown in Fig. 6 as a function of genetic architecture and admixture history.

Figure 6: Behavior of LDSC estimates of (A) V_g and (B) $\mathbb{E}(\chi^2)$ under the HI and CGF admixture models for traits under divergent (red) and stabilizing (blue) selection as a function of time since admixture (x-axis). In (A) the dashed and solid lines indicate the true (simulated) and estimated V_g, respectively, while the dotted horizontal line shows the expected genic variance. In (B) the solid line shows the mean χ^2 of the GWAS summary statistics while the dashed line shows the LDSC intercept (λ_{ldsc}). Shaded area indicates the 95% CI across ten replicates. Results here shown only for $p = 0.01$. Results for $p \in \{0.05, 0.1\}$ are shown in Fig. S6.

Under the HI model, the LDSC $\hat{V}_g$ is also consistent with an estimate of the genic variance, in line with theoretical expectation. As a result, LDSC under- and over-estimates V_g, and therefore, h^2_{snp} for traits under divergent and stabilizing selection, respectively (Fig. 6A). With time and random mating, V_g will converge to the genic variance, and therefore, to $\hat{V}_g$. The pattern is more complex for the CGF model, which shows an inflation in $\hat{V}_g$ in the first few generations of admixture, regardless of genetic architecture (Fig. 6A). We believe this is due to inflation in the GWAS summary statistics that is not fully captured by the LDSC intercept (λ_{ldsc}). This can be seen in Fig. 6B, which shows that the mean χ^2 is larger in the first few generations of the CGF model compared with that in the HI model

308 even though the simulated effects are the same. We confirm this by observing over-dispersion in GWAS

309 summary statistics especially in the first few generations of the CGF model (Fig. S7). Importantly, the

310 LDSC intercepts are similar under the HI and CGF models, suggesting that at least some of the inflation

311 in summary statistics is not captured by λ_{ldsc} under the CGF model. Because we performed GWAS

312 on genetic values directly, the inflation in test statistics is not because of environmental stratification.

313 Instead, we believe it arises from subtle patterns of admixture LD in the first few generations of the

314 CGF model when the minor (CEU) ancestry is rare in the sample. As such, rare variants tagging CEU

315 ancestry segments would be susceptible to subtle biases that are difficult to correct by including PCs as

316 linear covariates in the GWAS (Fig. S7E). The residual inflation, because it might be correlated with

317 admixture LD, is absorbed by the LDSC slope, leading to an over-estimate in V_g, regardless of genetic

318 architecture. Consistent with this, the bias appears to decrease with time under the CGF model (Fig. 6)

319 as the proportion of CEU ancestry increases in the population. As the bias decreases, the LDSC estimate

320 converges to the genic variance. Importantly, this estimate is also consistent with (ancestry-adjusted)

321 estimates from GREML and HE regression. We confirm this by applying GREML and HE regression to

322 data from the same set of simulations used for LDSC (Fig. S8).

323   In summary, LDSC estimates of V_g and h_{snp}^2 are susceptible to confounding due to population struc-

324 ture in admixed populations if the minor ancestry is represented by a small proportion of individuals in

325 the GWAS sample. But as long as this is not the case, LDSC estimates of V_g and h_{snp}^2 should also be

326 interpreted as the sum of the contributions of individual loci, and therefore should be consistent with

327 GREML and HE regression. This estimate may or may not be equal to h^2, depending on the genetic

328 architecture.

Local ancestry heritability

330 A related quantity of interest in admixed populations is local ancestry heritability (h_γ^2), which is defined

331 as the proportion of phenotypic variance that can be explained by local ancestry. Zaitlen *et al.* (2014)

332 [41] showed that this quantity is related to, and can be used to estimate, h^2 in admixed populations.

333 The advantage of this approach is that local ancestry segments shared between individuals are identical

334 by descent and are therefore, more likely to tag causal variants compared to array markers, allowing

335 one to potentially capture the contributions of rare variants [41]. Here, we show that in the presence of

336 population structure, (i) the relationship between h_γ^2 and h^2 is not straightforward and (ii) $\hat{h}_\gamma^2$ may be a

337 biased estimate of local ancestry heritability under the random effects model for the same reasons that

338 $\hat{h}_{snp}^2$ is biased.

339   We define local ancestry $\gamma_i \in \{0, 1, 2\}$ as the number of alleles at locus i that trace their ancestry

340 to population A. Thus, ancestry at the i^{th} locus in individual k is a binomial random variable with

341 $\mathbb{E}(\gamma_{ik}) = 2\theta_k$, θ_k being the ancestry of the k^{th} individual. Similar to genetic value, the 'ancestry value'

342 of an individual can be defined as $\sum_{i=1}^{m} \phi_i \gamma_i$, where $\phi_i = \beta_i(f_i^A - f_i^B)$ is the effect size of local ancestry

343 (Appendix). Then, the genetic variance due to local ancestry can be expressed as (Appendix):

$$V_\gamma = \mathbb{V}\left(\sum_{i=1}^{m} \phi_i \gamma_i\right) = \sum_{i=1}^{m} \phi_i^2 \, \mathbb{V}(\gamma_i) + \sum_{i=1}^{m} \sum_{j \neq i} \phi_i \phi_j \, \mathbb{C}(\gamma_i, \gamma_j)$$

$$= 2\,\mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} \sum_{i=1}^{m} \phi_i^2 + 2\,\mathbb{V}(\theta) \sum_{i=1}^{m} \phi_i^2 + 4\,\mathbb{V}(\theta) \sum_{i=1}^{m} \sum_{j \neq i} \phi_i \phi_j$$

$$= 2\,\mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} \sum_{i=1}^{m} \beta_i^2 (f_{\mathrm{i}}^{\mathrm{A}} - f_{\mathrm{i}}^{\mathrm{B}})^2$$

$$+ 2\,\mathbb{V}(\theta) \sum_{i=1}^{m} \beta_i^2 (f_{\mathrm{i}}^{\mathrm{A}} - f_{\mathrm{i}}^{\mathrm{B}})^2$$

$$+ 4\,\mathbb{V}(\theta) \sum_{i=1}^{m} \sum_{j \neq i} \beta_i \beta_j (f_{\mathrm{i}}^{\mathrm{A}} - f_{\mathrm{i}}^{\mathrm{B}})(f_{\mathrm{j}}^{\mathrm{A}} - f_{\mathrm{j}}^{\mathrm{B}})$$

and heritability explained by local ancestry is simply the ratio of V_γ and the phenotypic variance. Note

that $V_\gamma = (1.2) + (1.3) + (1.4)$ and therefore its behavior is similar to V_g in that the terms (1.3) and (1.4)

decay towards zero as $\mathbb{V}(\theta) \to 0$, and V_γ converges to (1.2) (Fig. S2). Additionally, the dependence of

V_γ on both $\mathbb{E}(\theta)$ and $\mathbb{V}(\theta)$ precludes a straightforward derivation between local ancestry heritability and

h^2.

GREML estimation of $\hat{h}_\gamma^2$ is similar to that of $\hat{h}_{snp}^2$, the key difference being that the former involves

constructing the GRM using local ancestry instead of genotypes [41]. The following model is assumed:

$y = Wv + \xi$ where W is an $n \times m$ standardized local ancestry matrix, $v \sim \mathcal{N}(0, I\frac{\sigma_v^2}{m})$ are local

ancestry effects, and $\xi \sim \mathcal{N}(0, I\sigma_\xi^2)$. Note that σ_ξ^2 captures both environmental noise as well as any

genetic variance independent of local ancestry. The phenotypic variance is decomposed as $\mathbb{V}(y) =$

$\mathbb{V}(Wv) + \mathbb{V}(\xi) = \frac{WW'}{m}\sigma_v^2 + \sigma_\xi^2$ where $\frac{WW'}{m}$ is the local ancestry GRM and σ_v^2 is the parameter of

interest, which is believed to be equal to V_γ – the genetic variance due to local ancestry.

We show that, in the presence of population structure, i.e., when $\mathbb{V}(\theta) > 0$, GREML $\hat{\sigma}_v^2$ is biased

downwards relative to V_γ for traits under divergent selection and upwards for traits under stabilizing

selection because it does not capture the contribution of LD (Fig. 7A). But there is another source of

bias in $\hat{\sigma}_v^2$, which tends to be inflated in the presence of population structure if individual ancestry is

not included as a covariate, even with respect to the expectation of V_γ under equilibrium (seen more

clearly in Fig. 7B-C). We suspect this inflation is because of strong correlations between local ancestry

– local ancestry disequilibrium – across loci that inflates $\hat{\sigma}_v^2$ in a way that is not adequately corrected

even when all causal variants are included in the model [4, 10]. Scaling local ancestry by its covariance

removes this bias and recovers the contribution of LD (Fig. 7D) presumably because this accounts for

the correlation in genotypes across loci. Including individual ancestry as a fixed effect also corrects for

the inflation in $\hat{\sigma}_v^2$ (Fig. 7E-H). But as with $\hat{\sigma}_u^2$, this practice will underestimate the genetic variance

due to local ancestry in the presence of population structure because it removes the variance along the

ancestry axis (Fig. 7E-H).

Based on the above, GREML $\hat{h}_\gamma^2$ and corresponding estimates of h^2 are more accurately interpreted as

the heritability due to local ancestry and heritability, respectively, expected in the absence of population

structure. We believe $\hat{h}_\gamma^2$ is still useful in that, because it should capture the effects of rare variants, it

can be used to estimate the upper bound of $\hat{h}_{snp}^2$.

In a previous paper, we suggested that local ancestry heritability could potentially be used to estimate
the genetic variance between populations [42]. Our results suggest this is not possible for two reasons.
First, the GREML estimator of local ancestry heritability, as we show in this section is biased and
does not capture the LD contribution. But even if we were able to recover the LD component, our
decomposition shows that local ancestry is equal to the genetic variance between populations (V_{gb}) only
when $\mathbb{E}(\theta) = 0.5$ and $\mathbb{V}(\theta) = \mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} = 0.25$, which is only possible at $t = 0$ in the HI model. After
admixture, $\mathbb{V}(\theta)$ decays and the equivalence between V_γ and V_{gb} is lost, making it impossible to estimate
the latter from admixed populations, especially for traits under divergent or stabilizing selection, even
if the environment is randomly distributed with respect to ancestry. We note that this conclusion was
recently reached independently by Schraiber and Edge (2023) [43].

Figure 7: The behavior of GREML estimates of the variance due to local ancestry ($\hat{\sigma}_v^2$) in admixed populations under the HI (left column) and CGF (right column) models either without (A-D) or with (E-H) individual ancestry included as a fixed effect. The solid lines represent estimates from simulated data averaged across ten replicates with red and blue colors representing estimates for traits under divergent and stabilizing selection, respectively. P indicates the strength of assortative mating. The dotted lines either represent the expected variance in the population (A & B) or the expected estimate for three different ways of scaling local ancestry (B-D & F-H). (A-B & E-F) show the behavior of $\hat{\sigma}_v^2$ for the default scaling, (C, G) shows $\hat{\sigma}_v^2$ when local ancestry is scaled by the sample variance, and (D, H) when it is scaled by the sample covariance. Shaded regions represent the 95% confidence bands. Some runs in (D & H) failed to converge as seen by the missing segments of the solid lines because the expected variance in such cases was too small.

How much does LD contribute to V_g in practice?

384 In the previous sections, we showed theoretically that $\hat{h}^2_{snp}$ may be biased in admixed populations even in

385 the absence of confounding by shared environment and even if the causal variants are directly genotyped.

386 All three estimators fail to capture the LD contribution. The extent to which $\hat{h}^2_{snp}$ is biased because

387 of this reason in practice is ultimately an empirical question, which is difficult to answer because the

388 true genetic architecture – the LD contribution in particular – is unknown. In this section, we develop

389 some intuition for this contribution among individuals with mixed African and European ancestry using

390 a combination of simulations and empirical data analysis.

391 First, we simulated a neutral trait using genotype data from the African Americans (ASW) from

392 the 1,000 Genomes Project [39]. To do this, we sampled $m \in \{10, 100, 1,000\}$ causal loci from a set of

393 common (MAF > 0.01), LD pruned variants and assigned them effects such that $\beta_i \sim \mathcal{N}\left(0, \frac{1}{\sqrt{m \, \mathbb{V}(x_i)}}\right)$,

394 i.e., the expected *genic* variance is $\mathbb{E}\{\sum_{i=1}^{m} \beta_i^2 Var(x_i)\} = 1$ (Methods). We computed the genic and

395 LD contributions and repeated this process 1,000 times where each replicate can be thought of as an

396 independent realization of the genetic architecture of a neutrally evolving trait. We show that the LD

397 contribution may be zero in expectation but can be substantial for a given trait (up to $\approx 50\%$ of the genic

398 variance, Fig. S4), even in the absence of selection.

399 Second, we estimated the LD contribution of genome-wide significant SNPs for 26 quantitative traits

400 among admixed Americans with primarily African and European ancestry. To do this, we decomposed

401 the variance explained by GWAS SNPs into the four components in Equation 1 using previously published

402 effect sizes, allele frequencies (f^A and f^B) from the 1000 Genomes YRI and CEU, and the mean ($\mathbb{E}(\theta) \approx$

403 0.77) and variance ($\mathbb{V}(\theta) \approx 0.02$) of individual ancestry from ASW (Methods). We show that for skin

404 pigmentation – a trait under strong divergent selection – the LD contribution, i.e. term (1.4), is positive

405 and accounts for $\approx 40\%$ of the total genetic variance. This is because of large allele frequency differences

406 between Africans and Europeans that are correlated across skin pigmentation loci, consistent with strong

407 polygenic selection favoring alleles for darker pigmentation in regions with high UV exposure and vice

408 versa [42, 44–47]. But for most other traits, LD contributes relatively little, explaining a modest, but

409 non-negligible proportion of the genetic variance. For example, GWAS SNPs for height, LDL and HDL

410 cholestrol, and mean corpuscular hemoglobin (MCH) exhibit positive LD, whereas SNPs for blood count

411 – particularly neutrophil (NEU) and white blood cell count (WBC) – loci exhibit negative LD (Fig.

412 8). Because we selected independent associations for this analysis (Methods), the LD contribution is

413 driven largely due to population structure in ASW. The contribution of population structure to the

414 genic variance, i.e., term (1.3) is small even for traits like skin pigmentation and neutrophil count with

415 large effect alleles that are highly diverged in frequency between Africans and Europeans [45, 46, 48–50].

416 Overall, this suggests a modest contribution of population structure to the genetic variance explained

417 by GWAS SNPs for most traits. Nevertheless, this contribution is non-negligible and may be higher for

418 other traits or for variants that do not reach genome-wide significance. We recommend that, in addition

419 to considering the contribution of population structure in overall heritability estimation, researchers

420 also account for it when reporting the total variance explained by GWAS SNPs in admixed and other

421 structured populations. This can be done with Equation 1 instead of the standard $2 \sum_{i=1}^{m} \hat{\beta}_i^2 f_i (1 - f_i)$

422 formula used in most studies.

Figure 8: Decomposing the genetic variance explained by GWAS SNPs in the 1000 Genomes ASW (African Americans from Southwest). We calculated the four variance components listed in Eq. 1, their values shown on the y-axis as a proportion of total genetic variance and confidence intervals from a parametric bootstrap (Methods). The LD contribution, which can be positive or negative, is shown in red, and the components of the genic variance are shown in shades of blue. The number of variants used to calculate variance components for each trait is shown at the bottom.

423 Discussion

424 Despite the growing size of GWAS and discovery of thousands of variants for hundreds of traits [51], the
425 heritability explained by GWAS SNPs remains a fraction of twin-based heritability estimates. Yang *et*
426 *al.* (2010) introduced the concept of SNP heritability (h^2_{snp}) that does not depend on the discovery of
427 causal variants but assumes that they are numerous and are more or less uniformly distributed across the
428 genome (the infinitesimal model), their contributions to the genetic variance 'tagged' by genotyped SNPs
429 [3]. h^2_{snp} is now routinely estimated in most genomic studies and at least for some traits (e.g. height and
430 BMI), these estimates now approach twin-based heritability [6]. But despite the widespread use of $\hat{h}^2_{snp}$,
431 its interpretation remains unclear, particularly in the presence of admixture and population structure.
432 It is generally accepted that $\hat{h}^2_{snp}$ can be biased in structured populations because of confounding effects
433 of unobserved environmental factors and LD between causal variants [4, 7, 9–11, 52]. But $\hat{h}^2_{snp}$ may be
434 biased even in the absence of confounding because of misspecification of the underlying random-effects
435 model, i.e., if the model does not represent the genetic architecture from which the trait is sampled [14,
436 15, 25, 53, 54].
437   Under the standard random-effects model, SNP effects are assumed to be uncorrelated and the total
438 genetic variance can be represented as the sum of the variance explained by individual loci, i.e. the

439 genic variance [14, 15, 25]. In admixed populations, there is substantial LD, which can contribute to the

440 genetic variance, and can persist for a number of generations, despite recombination, due to continued

441 gene flow and/or ancestry-based assortative mating. GREML and HE regression do not capture this

442 LD contribution, and therefore, may lead to biased estimates of h^2_{snp}. The LD contribution can be

443 negative for traits under stabilizing selection, and positive for traits under divergent selection between

444 the source populations, leading to over- or under-estimates, respectively. We show that in admixed

445 and other structured populations, GREML and HE regression estimates can be biased even when the

446 LD contribution is zero if the genotypes are scaled by $\sqrt{2f(1-f)}$ – the standard approach, which

447 implicitly assumes a randomly mating population. In the presence of population structure, the variance

448 in genotypes can be higher and $\hat{h}^2_{snp}$ does not capture this additional variance, which we show can be

449 recovered by scaling genotypes by the SNP variance ($\sqrt{Var(x)}$). Thus, as long as the genotypes are

450 scaled properly GREML and HE estimates of h^2_{snp} should be interpreted as the proportion of phenotypic

451 variance explained by the *genic* variance. Estimates of local ancestry heritability ($\hat{h}^2_\gamma$) [41, 55] should be

452 interpreted similarly.

453 The behavior of LD Score regression, which is different from GREML and HE in that it uses variant

454 summary statistics as input as opposed to individual genotypes, depends on the magnitude and pattern

455 of inflation in the GWAS and the extent to which it is captured by the LDSC intercept. We show that

456 under the HI model, where all the admixture occurs at once, the LDSC estimate of h^2_{snp} is equivalent

457 to GREML and HE regression in that it measures the sum of the individual contribution of each locus,

458 i.e., the genic variance. However, we note that under some population structures, e.g. under the CGF

459 model where the amount of gene flow in the first few generations is small, inflation in summary statistics

460 due to population structure may not uniformly affect all variants and can be correlated with local levels

461 of LD. This can lead to inflated estimates of h^2_{snp} that are not interpretable. As such, we recommend

462 caution when using LDSC in estimating heritability from GWAS in admixed and other structured cohorts,

463 especially when the ancestry composition of the cohort is highly unbalanced, i.e., when certain ancestries

464 are represented by a small number of individuals. We suggest that researchers remove any individuals

465 who may represent a very small fraction of the GWAS sample (in terms of ancestry) as well as report

466 SNP heritability estimates from all three approaches so that any inconsistencies can be evaluated for

467 potential sources of bias.

468 Does the bias in SNP heritability estimates due to the missing LD contribution have practical impli-

469 cations? The answer to this depends on the context in which SNP heritability is used. In the absence of

470 residual bias due to stratification (see previous paragraph), $\hat{h}^2_{snp}$ is an interpretable quantity: the sum

471 of the proportion of phenotypic variance explained by individual loci. As such, $\hat{h}^2_{snp}$ can be useful in

472 quantifying the power to detect variants in GWAS where the quantity of interest is the genic variance.

473 But because it does not capture the LD contribution to h^2, $\hat{h}^2_{snp}$ may not be appropriate in measuring

474 the extent to which genetic variation contributes to phenotypic variation, in predicting the response to

475 selection, or in defining the upper limit of polygenic prediction accuracy [2], especially for traits where

476 the genetic risk varies as a function of ancestry (e.g. health disparities).

477 One limitation of this paper is that we have focused on random-effects estimators of h^2_{snp} because of

478 their widespread use. Estimators of h^2_{snp} can be broadly grouped into random- and fixed effect estimators

based on how they treat SNP effects [36]. Fixed effect estimators make fewer distributional assumptions
but they are not as widely used because they require conditional estimates of all variants – a high-
dimensional problem where the number of markers is often far larger than the sample size [56]. This is
one reason why random effect estimators, such as GREML, are popular – because they reduce the number
of parameters that need to be estimated by assuming that the effects are drawn from some distribution
where the variance is the only parameter of interest. Fixed effects estimators, in principle, should be able
to capture the LD contribution but this is not obvious in practice since the simulations used to evaluate
the accuracy of such estimators still assume uncorrelated effects [36, 56, 57]. Further research is needed
to clarify the interpretation of the different estimators of h^2_{snp} in structured populations under a range
of genetic architectures.

Ultimately, the discrepancy between $\hat{h}^2_{snp}$ and h^2 in practice is an empirical question, the answer to
which depends on the degree of population structure (which we can measure) and the genetic architecture
of the trait (which we do not know *a priori*). We show that for most traits, the contribution of population
structure to the variance explained by GWAS SNPs is modest among African Americans. Thus, if we
assume that the genetic architecture of GWAS SNPs represents that of all causal variants, then despite
incorrect assumptions, the discrepancy between $\hat{h}^2_{snp}$ and h^2 should be fairly small. But this assumption
is unrealistic given that GWAS SNPs are common variants that in most cases cumulatively explain a
fraction of trait heritability. What is the LD contribution of the rest of the genome, particularly rare
variants? This is not obvious and will become clearer in the near future through large sequence-based
studies [58]. While these are underway, theoretical studies are needed to understand how different
selection regimes influence the directional LD between causal variants – clearly an important aspect of
the genetic architecture of complex traits.

Methods

Simulating genetic architecture

We first drew the allele frequency (f^0) of 1,000 biallelic causal loci in the ancestor of populations A and
B from a uniform distribution, $U(0.001, 0.999)$. Then, we simulated their frequency in populations A and
B (f^A and f^B) under the Balding-Nichols model [33], such that $f^A, f^B \sim Beta(\frac{f^0(1-F)}{F}, \frac{(1-f^0)(1-F)}{F})$
where $F = 0.2$ is the inbreeding coefficient. We implemented this using code adapted from Lin *et al.*
(2021) [59]. To avoid drawing extremely rare alleles, we continued to draw f^A and f^B until we had 1,000
loci with $f^A, f^B \in (0.01, 0.99)$.

We generated the effect size of the i^{th} locus by setting $\beta_i = \frac{1}{\sqrt{2m\bar{f}_i(1-\bar{f}_i)}}$, where m is the number
of loci and $\bar{f}$ is the mean allele frequency across populations A and B. Thus, rare variants have larger
effects than common variants and the total genetic variance sums to 1. Given these effects, we simulated
two different traits, one with a large difference in means between populations A and B (Trait 1) and
the other with roughly no difference (Trait 2). This was achieved by permuting the signs of the effects
100 times to get a distribution of V_{gb} – the genetic variance between populations. This has the effect of
varying the LD contribution without changing the F_{ST} at causal loci. We selected the maximum and

516 minimum of V_{gb} to represent Traits 1 and 2.

Simulating admixture

518 We simulated the genotypes, local ancestry, and phenotype for 10,000 admixed individuals per generation

519 under the hybrid isolation (HI) and continuous gene flow (CGF) models by adapting the code from Zaitlen

520 *et al.* (2017) [27]. We denote the ancestry of a randomly selected individual k with θ, the fraction of their

521 genome from population A. At $t = 0$ under the HI model, we set θ to 1 for individuals from population A

522 and 0 if they were from population B such that $\mathbb{E}(\theta) = p \in \{0.1, 0.2, 0.5\}$ with no further gene flow from

523 either source population. In the CGF model, population B receives a constant amount q from population

524 A in every generation starting at $t = 0$. The mean overall proportion of ancestry in the population is

525 kept the same as the HI model by setting $q = 1 - (1 - p)^{\frac{1}{t}}$ where t is the number of generations of gene

526 flow from A. In every generation, we simulated ancestry-based assortative mating by selecting mates

527 such that the correlation between their ancestries is $P \in \{0, 0.3, 0.6, 0.9\}$ in every generation. We do this

528 by repeatedly permuting individuals with respect to each other until P falls within ± 0.01 of the desired

529 value. It becomes difficult to meet this criterion when $\mathbb{V}(\theta)$ is small (Fig.1C). To overcome this, we

530 relaxed the threshold up to 0.04 for some conditions, i.e., when $\theta \in \{0.1, 0.2\}$ and $t \geq 50$. We generated

531 expected variance in individual ancestry using the expression in Zaitlen *et al.* (2017) [27]. At time t

532 since admixture, $\mathbb{V}(\theta_t) = \mathbb{V}(\theta_{t-1})\frac{(1+P)}{2}$ under the HI model where P measures the strength of assortative

533 mating, i.e, the correlation between the ancestry between individuals in a mating pair. Under the CGF

534 model, $\mathbb{V}(\theta_t) = q(1-q)\mathbb{E}(\theta_{t-1})^2 + q(1-q)\{1 - 2\mathbb{E}(\theta_{t-1})\} + (1-q)\mathbb{V}(\theta_{t-1})\frac{(1+P)}{2}$ (Appendix).

535 We sampled the local ancestry at each i^{th} locus as $\gamma_i = \gamma_{if} + \gamma_{im}$ where $\gamma_{im} \sim Bin(1, \theta_m), \gamma_{if} \sim$

536 $Bin(1, \theta_f)$ and θ_m and θ_f represent the ancestry of the maternal and paternal chromosome, respectively.

537 The global ancestry of the individual is then calculated as $\theta_k = \sum_{i=1}^{m} \frac{\gamma_{im} + \gamma_{if}}{2m}$, where m is the number of

538 loci. We sample the genotypes x_{im} and x_{if} from a binomial distribution conditioning on local ancestry.

539 For example, the genotype on the maternal chromosome is $x_{im} \sim Bin(1, f_i^A)$ if $\gamma_{im} = 1$ and $x_{im} \sim$

540 $Bin(1, f_i^B)$ if $\gamma_{im} = 0$ where f_i^A and f_i^B represent the allele frequency in populations A and B, respectively.

541 Then, the genotype can be obtained as the sum of the maternal and paternal genotypes: $x_i = x_{im} + x_{ip}$.

542 We calculate the genetic value of each individual as $g = \sum_{i=1}^{m} \beta_i x_i$ and the genetic variance as $\mathbb{V}(g)$.

543 For LDSC estimation, we simulated phased genotypes on chromosome 2 ($M = 88,112$ HapMap 3

544 SNPs) for $N = 5,000$ individuals for 20 generations of admixture under the HI and CGF models from

545 CEU and YRI haplotypes from the 1000 Genomes Project ([39]) using Haptools [60] implemented in

546 Admix-kit [61]. We used recombination maps for the hg38 build of the Human genome downloadable

547 from the Beagle (v5.5) website [62]. We sampled $p \in \{0.01, 0.05, 0.1\}$ of the variants to be causal

548 and assigned them effects such that $\beta_i = \frac{1}{2pM\bar{f}(1-\bar{f})}$ where $\bar{f} = \frac{f_{CEU} + f_{YRI}}{2}$ with the condition that

549 $0.01 < \bar{f} < 0.99$. We permuted the signs of the effects 1,000 times and selected the combination that

550 gave the smallest and largest value of genetic variance between CEU and YRI (V_{gb}) to represent traits

551 under stabilizing and divergent selection, respectively.

Heritability estimation with GREML

We used the *--reml* and *--reml-no-constrain* flags in GCTA [5] to estimate σ_u^2 and σ_v^2, the genetic variance due to genotypes and local ancestry, respectively. We could not run GCTA without noise in the genetic values so we simulated individual phenotypes with a heritability of $h^2 = 0.8$ by adding random noise $e \sim \mathcal{N}(0, V_g \frac{1-h^2}{h^2})$. We computed three different GRMs, which correspond to different transformations of the genotypes: (i) standard, (i) Variance or $V(x)$ scaled, and (ii) LD-scaled.

For the standard GRM, the genotypes at the i^{th} SNP are standardized such that $z_i = \frac{x_i - 2f_i}{\sqrt{2f_i(1-f_i)}}$. For the variance scaled GRM, we computed $z_i = \frac{x_i - 2f_i}{\sqrt{\mathbb{V}(x_i)}}$ where $\mathbb{V}(x_i)$ is the sample variance of the genotypes at the i^{th} SNP. The LD-scaled GRM conceptually corresponds to standardizing the genotypes by the SNP covariance, rather than its variance. Let $\boldsymbol{X}$ represent the $n \times m$ *unstandardized* matrix of genotypes and $\boldsymbol{P}$ represent an $n \times m$ matrix where the i^{th} column contains the allele frequency of that SNP. Let $\boldsymbol{U}$ be the upper triangular 'square root' matrix of the $m \times m$ SNP covariance matrix $\boldsymbol{\Sigma}$ such that $\boldsymbol{\Sigma} = \boldsymbol{U}'\boldsymbol{U}$. Then, the standardized genotypes are computed as $\boldsymbol{Z} = (\boldsymbol{X} - 2\boldsymbol{P})\boldsymbol{U}^{-1}$ and the GRM becomes $(\boldsymbol{X} - 2\boldsymbol{P})\boldsymbol{\Sigma}^{-1}(\boldsymbol{X} - 2\boldsymbol{P})'$ [34]. Similarly, the three GRMs for local ancestry were computed by scaling local ancestry with (i) $\sqrt{2\bar{\gamma}_i(1 - \bar{\gamma}_i)}$ where we denote $\bar{\gamma}_i$ as the mean local ancestry at the i^{th} SNP, or with the (ii) variance, or (iii) covariance of local ancestry, respectively. We estimated σ_u^2 and σ_v^2 with and without individual ancestry as a fixed effect to correct for any confounding due to genetic stratification. This was done by using the *--qcovar* flag.

Heritability estimation with HE regression

Haseman-Elston regression without ancestry correction was implemented using GCTA [4]. To demonstrate that the bias in unadjusted HE estimates arises because of a bias in the estimate of LD contribution, not the genic variance, we carried out a simple simulation where half of the individuals in the population derive their ancestry from population A and the rest from population B. This is equivalent to the meta-population under the HI model at $t = 0$ where $\mathbb{E}(\theta) = 0.5$. We simulated genotypes for $1,000$ individuals for $m = 100$ loci where the allele frequencies in populations A and B were set to $f_A = 0.1$ and $f_B = 0.8$, respectively. We standardized the genotypes at each locus i using the square-root of the sample variance and assigned effect sizes such that the total genetic variance explained by all loci is equal to 1, i.e., the effect of the scaled genotype at the i^{th} locus is $u_i = \frac{1}{\sqrt{m}}$. This is equivalent to the effect size of the unscaled genotypes being $\beta_i = \frac{1}{\sqrt{m \mathbb{V}(x_i)}}$ where $\mathbb{V}(x_i)$ is the sample variance at the i^{th} locus. We introduced randomness in the direction of the effect by assigning a negative or positive sign to each locus uniformly at random 100 times to generate 100 traits with the same genic variance but varying LD contributions. Then, for each trait we computed the two terms in Eq. 2, which should converge to the genic variance and LD contributions, which represent the genic and LD components to the HE regression estimate. Fig. S5 shows that in the presence of directional LD, the overall bias is in the HE regression estimate is due to an exaggerated estimate of the LD contribution.

To correct for individual ancestry, we followed the moment-matching approach of Ge et al. (2017) implemented in MMHE [37]. In this approach, conceptually, individual ancestry is first projected out from both the phenotype and genotypes. Then, the cross-product of the residual phenotypes is regressed

590 on the corresponding entries of the GRM (computed from the residual genotypes) to estimate V_g. In
591 practice, a more efficient algorithm is used.

Heritability estimation with LDSC

593 We used estimate LD score regression heritability with cov-LDSC [40]. We used summary statistics from a
594 GWAS of the genetic values with 20 PCs as covariates. We used cov-LDSC to compute ancestry-adjusted
595 LD scores (for 20 PCs) and to estimate the LDSC slope and intercept (λ_{ldsc}) for each combination of
596 the parameter space, i.e., (i) demographic model (HI or CGF), (ii) generations since admixture (2
597 - 20), (iii) genetic architecture (divergent or stabilizing), (iv) proportion of variants that are causal
598 $p \in \{0.01, 0.05, 0.1\}$, and (v) replicate (1-10). Because we used the genetic values directly in the GWAS,
599 the LDSC slope yields an estimate of the genetic variance ($\hat{V}_g^{ldsc}$). For consistency, we also carried out
600 GREML and HE regression estimation on the same data using GCTA [5] (Fig. S8).
601   To explore the inflation in GWAS test statistics observed under the CGF model (Fig. 6, Fig. S6),
602 compared the GWAS effect sizes of all variants with their expected effects computed from their LD with
603 the causal variants. Specifically, the expected effect of the j^{th} SNP is $\mathbb{E}(\hat{\beta}_j) = \sum_{i=1}^{pM} \beta_i r_{ij}$ where β_i is the
604 true (simulated) effect of the i^{th} causal variant and r_{ij} is the genotypic correlation between them. We
605 compared the expected and estimated GWAS effects for a single replicate of genotypes simulated under
606 the HI and CGF models for generations 2 and 20 since admixture. These results are shown in Fig. S7.

Estimating variance explained by GWAS SNPs

608 We retrieved the summary statistics of 26 traits from GWAS catalog [51]. Full list of traits and the
609 source papers [47, 63–70] are listed in Table S1. To maximize the number of variants discovered, we
610 chose summary statistics from studies that were conducted in both European and multi-ancestry samples
611 and that reported the following information: effect allele, effect size, p-value, and genomic position.
612 For height, we used the conditional and joint (COJO) effect sizes reported in Yengo *et al.* (2022)
613 [64]. For birth weight, we downloaded the data from the Early Growth Genetics (EGG) consortium
614 website [67] since the version reported on the GWAS catalog is incomplete. For skin pigmentation,
615 we chose summary statistics from the UKB [71] released by the Neale Lab (http://www.nealelab.is/uk-
616 biobank) and processed by Ju and Mathieson [47] to represent effect sizes estimated among individuals
617 of European ancestry. We also selected summary statistics from Lona-Durazo *et al.* (2019) where
618 effect sizes were meta-analyzed across four admixed cohorts [63]. Lona-Durazo *et al.* provide summary
619 statistics separately with and without conditioning on rs1426654 and rs35397 – two large effect variants
620 in *SLC24A5* and *SLC45A2*. We used the 'conditioned' effect sizes and added in the effects of rs1426654
621 and rs35397 to estimate genetic variance.
622   We selected independent hits for each trait by pruning and thresholding with PLINK v1.90b6.21 [72]
623 in two steps as in Ju *et al.* (2020) [47]. We used the genotype data of GBR from the 1000 genome
624 project [39] as the LD reference panel. We kept only SNPs (indels were removed) that passed the
625 genome-wide significant threshold (*--clump-p1 5e-8*) with a pairwise LD cutoff of 0.05 (*--clump-r2 0.05*)
626 and a physical distance threshold of 250kb (*--clump-kb 250*) for clumping. Second, we applied a second

627 round of clumping (*--clump-kb 100*) to remove SNPs within 100kb.

628   When GWAS was carried out separately in different ancestry cohorts in the same study, we used
629 inverse-variance weighting to meta-analyze effect sizes for variants that were genome-wide significant
630 (p-value $< 5 \times 10^{-8}$) in at least one cohort. This allowed us to maximize the discovery of variants such
631 as the Duffy null allele that are absent among individuals of European ancestry but polymorphic in other
632 populations [50].

633   We used allele frequencies from the 1000 Genomes CEU and YRI to represent the allele frequencies
634 of GWAS SNPs in Europeans and Africans, respectively, making sure that the alleles reported in the
635 summary statistics matched the alleles reported in the 1000 Genomes. We estimated the global ancestry
636 of ASW individuals (N = 74) with CEU and YRI individuals from 1000 genome (phase 3) using ADMIX-
637 TURE 1.3.0 [73] with k=2 and used it to calculate the mean (proportion of African ancestry = 0.767)
638 and variance (0.018) of global ancestry in ASW. With the effect sizes, allele frequencies, and the mean
639 and variance in ancestry, we calculated the four components of genetic variance using Equation 1. The
640 confidence intervals were calculated using a parametric bootstrap by randomly sampling the estimated
641 effect of each variant 100 times from a standard normal distribution with a mean and standard deviation
642 equal to the effect size size and standard error reported by the original GWAS.

643   Initially, the multi-ancestry summary statistics for a few traits (NEU, WBC, MON, MCH, BAS)
644 yielded values > 1 for the proportion of variance explained. This is likely because, despite LD pruning,
645 some of the variants in the model are not independent and tag large effect variants under divergent
646 selection such as the Duffy null allele, leading to an inflated contribution of LD. We checked this by
647 calculating the pairwise contribution , i.e., $\beta_i \beta_j (f_i^A - f_i^B)(f_j^A - f_j^B)$, of all SNPs in the model and show
648 long-range positive LD between variants on chromosome 1 for NEU, WBC, and MON, especially with the
649 Duffy null allele (Fig. S9A-C). A similar pattern was observed on chromosome 16 for MCH, confirming
650 our suspicion. This also suggests that for certain traits, pruning and thresholding approaches are not
651 guaranteed to yield independent hits. To get around this problem, we retained only one association with
652 the lowest p-value, each from chromosome 1 (rs2814778 for NEU, WBC, and MON) and chromosome 16
653 (rs13331259 for MCH) (Fig. S9D). For BAS, we observed that the variance explained was driven by a
654 rare variant (rs188411703, MAF = 0.0024) of large effect ($\beta = -2.27$). We believe this effect estimate
655 to be inflated and therefore, we removed it from our calculation.

656   As a sanity check to ensure transferability of GWAS effect sizes, we re-estimated the effects using
657 genotypic and phenotypic data from admixed individuals from the All of Us Research Program (AoU) [20].
658 To restrict the analysis to individuals with primarily mixed African and European ancestry, we focused
659 on 61,461 individuals who self-identified as Black or African American, excluding those of Hispanic or
660 Latino ethnicity. Of those 61,461 individuals, 48,587 individuals had trait and covariate data and were
661 used as our final cohort. Phenotypes were matched from the GWAS catalog to AoU, with codes found in
662 Table S1 [20, 51]. Of the 26 traits, 21 were available in AoU, all except skin pigmentation, birth weight,
663 pulse pressure, red blood cell count distribution width, and mean platelet volume. When necessary, we
664 converted all observations to the most frequent unit of measurement for each trait (e.g., for height, the
665 most common unit of measure was centimeters). We visually removed any outliers that were extreme
666 or where the unit value was misspecified (Table S1). The GWAS variants were extracted from vcf files

called by the AoU from whole genome sequencing (WGS) data. We re-estimated the effect sizes for
GWAS variants for the 21 traits using the *--glm* flag in PLINK [72] with age, age^2, sex assigned at birth,
and the first 16 genetic PCs included as covariates. All phenotypes and covariates were standardized
to unit variance. We show that effect sizes re-estimated in AoU are highly correlated with the original
GWAS effects with the exception of a few outliers (Fig S10).

Code availability

We carried out all analyses in R version 4.2.3 [74], PLINK v1.90b6.21 and PLINK 2.0 [72, 75], and GCTA
version 1.94.1 [5]. All code is freely available on https://github.com/zaidilab/admix_heritability.git.

Data availability

Summary statistics for each trait can be found in Table S1. This study used data from the All of
Us Research Program's Controlled Tier Dataset v7, available to authorized users on the Researcher
Workbench [20].

Acknowledgements

We thank Iain Mathieson and Doc Edge for helpful comments on the manuscript. This study was funded
by National Institute of General Medical Sciences award R00GM137076 to AAZ and T32GM132063 to
NK. The content of this paper is solely the responsibility of the authors and does not necessarily represent
the official views of the National Institutes of Health. The All of Us Research Program would not be
possible without the partnership of its participants.

References

1. Lynch, M. & Walsh, B. *Genetics and analysis of quantitative traits* 1–980 (Sinauer Associates, Inc, 1998).

2. Visscher, P. M., Hill, W. G. & Wray, N. R. Heritability in the genomics era–concepts and misconceptions. *Nature reviews. Genetics* **9,** 255–66 (4 2008).

3. Yang, J., Benyamin, B., *et al.* Common SNPs explain a large proportion of the heritability for human height. *Nature Genetics* **42,** 565–569 (7 2010).

4. Yang, J., Manolio, T. A., *et al.* Genome partitioning of genetic variation for complex traits using common SNPs. *Nature Genetics 2011 43:6* **43,** 519–525 (6 2011).

5. Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide complex trait analysis. *American journal of human genetics* **88,** 76–82 (1 2011).

6. Wainschtein, P., Jain, D., *et al.* Assessing the contribution of rare variants to complex trait heritability from whole-genome sequence data. *Nature Genetics 2022 54:3* **54,** 263–273 (3 2022).

7. Browning, S. R. & Browning, B. L. Population structure can inflate SNP-based heritability estimates. *American Journal of Human Genetics* **89,** 191–193 (1 2011).

8. Goddard, M. E., Lee, S. H., Yang, J., Wray, N. R. & Visscher, P. M. Response to browning and browning. *American Journal of Human Genetics* **89,** 193–195 (1 2011).

9. Kumar, S. K., Feldman, M. W., Rehkopf, D. H. & Tuljapurkar, S. Limitations of GCTA as a solution to the missing heritability problem. *Proceedings of the National Academy of Sciences of the United States of America* **113,** E61–E70 (1 2016).

10. Yang, J., Leec, S. H., Wraya, N. R., Goddardd, M. E. & Visscher, P. M. GCTA-GREML accounts for linkage disequilibrium when estimating genetic variance from genome-wide SNPs. *Proceedings of the National Academy of Sciences* **113,** E4579–E4580 (32 2016).

11. Lin, Z., Seal, S. & Basu, S. Estimating SNP heritability in presence of population substructure in biobank-scale datasets. *Genetics* **220** (4 2022).

12. Border, R., O'Rourke, S., *et al.* Assortative mating biases marker-based heritability estimators. *Nature Communications 2022 13:1* **13,** 1–10 (1 2022).

13. Visscher, P. M., Yang, J. & Goddard, M. E. A Commentary on 'Common SNPs Explain a Large Proportion of the Heritability for Human Height' by Yang et al. (2010). *Twin Research and Human Genetics* **13,** 517–524 (6 2010).

14. Rawlik, K., Canela-Xandri, O., Oolliams, J. W. & Tenesa, A. SNP heritability: What are we estimating? *bioRxiv,* 2020.09.15.276121 (2020).

15. Lara, L. A. C., Pocrnic, I., de P. Oliveira, T., Gaynor, R. C. & Gorjanc, G. Temporal and genomic analysis of additive genetic variance in breeding programmes. *Heredity 2021 128:1* **128,** 21–32 (1 2021).

16. Wojcik, G. L., Graff, M., *et al.* Genetic analyses of diverse populations improves discovery for complex traits. *Nature* **570,** 514–518 (7762 2019).

17. Ben-Eghan, C., Sun, R., *et al.* Don't ignore genetic data from minority populations. *Nature 2021 585:7824* **585,** 184–186 (7824 2020).

18. Verma, A., Damrauer, S. M., *et al.* The Penn Medicine BioBank: Towards a Genomics-Enabled Learning Healthcare System to Accelerate Precision Medicine in a Diverse Population. *Journal of Personalized Medicine* **12,** 1974 (12 2022).

19. Fatumo, S., Chikowore, T., *et al.* A roadmap to increase diversity in genomic studies. *Nature Medicine 2022 28:2* **28,** 243–250 (2 2022).

20. The All of Us Research Program Investigators. The "All of Us" Research Program. *New England Journal of Medicine* **381,** 668–676 (7 2019).

21. Sohail, M., Chong, A. Y., *et al.* Nationwide genomic biobank in Mexico unravels demographic history and complex trait architecture from 6,057 individuals. *bioRxiv,* 2022.07.11.499652 (2022).

22. Johnson, R., Ding, Y., *et al.* The UCLA ATLAS Community Health Initiative: Promoting precision health research in a diverse biobank. *Cell Genomics* **3,** 100243 (1 2023).

23. Bulik-Sullivan, B. K., Loh, P.-R., *et al.* LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nature Genetics* **47,** 291–295 (3 2015).

24. Haseman, J. K. & Elston, R. C. The investigation of linkage between a quantitative trait and a marker locus. *Behavior genetics* **2,** 3–19 (1 1972).

25. De los Campos, G., Sorensen, D. & Gianola, D. Genomic Heritability: What Is It? *PLOS Genetics* **11,** e1005048 (5 2015).

26. Pfaff, C. L., Parra, E. J., *et al.* Population structure in admixed populations: effect of admixture dynamics on the pattern of linkage disequilibrium. *American journal of human genetics* **68,** 198–207 (1 2001).

27. Zaitlen, N., Huntsman, S., *et al.* The Effects of Migration and Assortative Mating on Admixture Linkage Disequilibrium. *Genetics* **205,** 375–383 (1 2017).

28. Verdu, P. & Rosenberg, N. A. A General Mechanistic Model for Admixture Histories of Hybrid Populations. *Genetics* **189,** 1413–1426 (4 2011).

29. Corre, V. L. & Kremer, A. Genetic Variability at Neutral Markers, Quantitative Trait Loci and Trait in a Subdivided Population Under Selection. *Genetics* **164,** 1205–1219 (3 2003).

30. Berg, J. J. & Coop, G. A Population Genetic Signal of Polygenic Adaptation. *PLoS Genetics* **10,** e1004412 (8 2014).

31. Bulmer, M. G. The Effect of Selection on Genetic Variability. *The American Naturalist* **105,** 201–211 (943 1971).

32. Yair, S. & Coop, G. Population differentiation of polygenic score predictions under stabilizing selection. *Philosophical Transactions of the Royal Society B* **377** (1852 2022).

33. Balding, D. J. & Nichols, R. A. A method for quantifying differentiation between populations at multi-allelic loci and its implications for investigating identity and paternity. *Genetica* **96,** 3–12 (1-2 1995).

34. Mathew, B., Léon, J. & Sillanpää, M. J. A novel linkage-disequilibrium corrected genomic relationship matrix for SNP-heritability estimation and genomic prediction. *Heredity 2017 120:4* **120,** 356–368 (4 2017).

35. Ma, R. & Dicker, L. H. The Mahalanobis kernel for heritability estimation in genome-wide association studies: fixed-effects and random-effects methods. *arXiv* (2019).

36. Min, A., Thompson, E. & Basu, S. Comparing heritability estimators under alternative structures of linkage disequilibrium. *G3 Genes|Genomes|Genetics* **12** (8 2022).

37. Ge, T., Chen, C. Y., Neale, B. M., Sabuncu, M. R. & Smoller, J. W. Phenome-wide heritability analysis of the UK Biobank. *PLOS Genetics* **13,** e1006711 (4 2017).

38. Pirinen, M. *GWAS 8: Heritability and mixed models* https://www.mv.helsinki.fi/home/mjxpirin/GWAS_course/material/GWAS8.html.

39. Auton, A., Abecasis, G. R., *et al.* A global reference for human genetic variation. *Nature 2015 526:7571* **526,** 68–74 (7571 2015).

40. Luo, Y., Li, X., *et al.* Estimating heritability and its enrichment in tissue-specific gene sets in admixed populations. *Human Molecular Genetics* **30,** 1521–1534 (16 2021).

41. Zaitlen, N., Pasaniuc, B., *et al.* Leveraging population admixture to characterize the heritability of complex traits. *Nature Genetics* **46,** 1356–1362 (12 2014).

42. Zaidi, A. A., Mattern, B. C., *et al.* Investigating the case of human nose shape and climate adaptation. *PLoS Genetics* **13,** 2017 (3 2017).

43. Schraiber, J. G. & Edge, M. D. Heritability within groups is uninformative about differences among groups: cases from behavioral, evolutionary, and statistical genetics. *bioRxiv,* 2023.11.06.565864 (2023).

44. Jablonski, N. G. the Evolution of Human Skin and Skin Color. *Annual Review of Anthropology* **33,** 585–623 (1 2004).

45. Lamason, R. L., Mohideen, M. A. P., *et al.* SLC24A5, a putative cation exchanger, affects pigmentation in zebrafish and humans. *Science* **310,** 1782–1786 (5755 2005).

46. Beleza, S., Johnson, N. A., *et al.* Genetic architecture of skin and eye color in an African-European admixed population. *PLoS genetics* **9** (ed Spritz, R. A.) e1003372 (3 2013).

47. Ju, D. & Mathieson, I. The evolution of skin pigmentation-associated variation in West Eurasia. *Proceedings of the National Academy of Sciences of the United States of America* **118,** e2009227118 (1 2020).

48. Nalls, M. A., Wilson, J. G., *et al.* Admixture Mapping of White Cell Count: Genetic Locus Responsible for Lower White Blood Cell Count in the Health ABC and Jackson Heart Studies. *The American Journal of Human Genetics* **82,** 81–87 (1 2008).

49. Reich, D., Nalls, M. A., *et al.* Reduced Neutrophil Count in People of African Descent Is Due To
a Regulatory Variant in the Duffy Antigen Receptor for Chemokines Gene. *PLoS Genetics* **5** (ed
Visscher, P. M.) e1000360 (1 2009).

50. McManus, K. F., Taravella, A. M., *et al.* Population genetic analysis of the DARC locus (Duffy)
reveals adaptation from standing variation associated with malaria resistance in humans. *PLOS*
Genetics **13,** e1006560 (3 2017).

51. Sollis, E., Mosaku, A., *et al.* The NHGRI-EBI GWAS Catalog: knowledgebase and deposition re-
source. *Nucleic Acids Research* **51,** D977–D985 (D1 2023).

52. Kumar, S. K., Feldman, M. W., Rehkop, D. H. & Tuljapurkar, S. Reply to Yang et al.: GCTA
produces unreliable heritability estimates. *Proceedings of the National Academy of Sciences* **113,**
E4581–E4581 (32 2016).

53. Speed, D., Hemani, G., Johnson, M. R. & Balding, D. J. Improved heritability estimation from
genome-wide SNPs. *American journal of human genetics* **91,** 1011–21 (6 2012).

54. Speed, D., Cai, N., Johnson, M. R., Nejentsev, S. & Balding, D. J. Reevaluation of SNP heritability
in complex human traits. *Nature Genetics* **49,** 986–992 (7 2017).

55. Chan, T. F., Rui, X., *et al.* Estimating heritability explained by local ancestry and evaluating strat-
ification bias in admixture mapping from summary statistics. *bioRxiv,* 2023.04.10.536252 (2023).

56. Schwartzman, A., Schork, A. J., Zablocki, R. & Thompson, W. K. A simple, consistent estimator
of SNP heritability from genome-wide association studies. *https://doi.org/10.1214/19-AOAS1291*
13, 2509–2538 (4 2019).

57. Hou, K., Ding, Y., *et al.* Causal effects on complex traits are similar for common variants across
segments of different continental ancestries within admixed individuals. *Nature Genetics 2023 55:4*
55, 549–558 (4 2023).

58. Backman, J. D., Li, A. H., *et al.* Exome sequencing and analysis of 454,787 UK Biobank participants.
Nature 2021 599:7886 **599,** 628–634 (7886 2021).

59. Lin, M., Park, D. S., Zaitlen, N. A., Henn, B. M. & Gignoux, C. R. Admixed Populations Improve
Power for Variant Discovery and Portability in Genome-Wide Association Studies. *Frontiers in*
Genetics **12,** 673167 (2021).

60. Massarat, A. R., Lamkin, M., *et al.* Haptools: a toolkit for admixture and haplotype analysis.
Bioinformatics **39** (3 2023).

61. Hou, K., Gogarten, S., *et al.* Admix-kit: an integrated toolkit and pipeline for genetic analyses of
admixed populations. *Bioinformatics* **40** (4 2024).

62. Browning, B. L., Zhou, Y. & Browning, S. R. A One-Penny Imputed Genome from Next-Generation
Reference Panels. *The American Journal of Human Genetics* **103,** 338–348 (2018).

63. Lona-Durazo, F., Hernandez-Pacheco, N., *et al.* Meta-analysis of GWA studies provides new insights
on the genetic architecture of skin pigmentation in recently admixed populations. *BMC Genetics*
20, 1–16 (1 2019).

64.   Yengo, L., Vedantam, S., *et al.* A saturated map of common genetic variants associated with human
height. *Nature* **610,** 704–712 (7933 2022).

65.   Hoffmann, T. J., Choquet, H., *et al.* A large multiethnic genome-wide association study of adult
body mass index identifies novel loci. *Genetics* **210,** 499–515 (2 2018).

66.   Pulit, S. L., Stoneman, C., *et al.* Meta-Analysis of genome-wide association studies for body fat
distribution in 694 649 individuals of European ancestry. *Human Molecular Genetics* **28,** 166–174
(1 2019).

67.   Warrington, N. M., Beaumont, R. N., *et al.* Maternal and fetal genetic effects on birth weight and
their relevance to cardio-metabolic risk factors. *Nature Genetics 2019 51:5* **51,** 804–814 (5 2019).

68.   Surendran, P., Feofanova, E. V., *et al.* Discovery of rare variants associated with blood pressure
regulation through meta-analysis of 1.3 million individuals. *Nature Genetics* **52,** 1314–1332 (12
2020).

69.   Graham, S. E., Clarke, S. L., *et al.* The power of genetic diversity in genome-wide association studies
of lipids. *Nature* **600,** 675–679 (7890 2021).

70.   Chen, M. H., Raffield, L. M., *et al.* Trans-ethnic and Ancestry-Specific Blood-Cell Genetics in
746,667 Individuals from 5 Global Populations. *Cell* **182,** 1198–1213.e14 (5 2020).

71.   Bycroft, C., Freeman, C., *et al.* The UK Biobank resource with deep phenotyping and genomic
data. *Nature* **562,** 203–209 (7726 2018).

72.   Chang, C. C., Chow, C. C., *et al.* Second-generation PLINK: rising to the challenge of larger and
richer datasets. *GigaScience* **4,** s13742–015–0047–8 (1 2015).

73.   Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in unrelated
individuals. *Genome research* **19,** 1655–64 (9 2009).

74.   R Core Team. *R: A Language and Environment for Statistical Computing* R Foundation for Sta-
tistical Computing (Vienna, Austria, 2023).

75.   Purcell, S., Neale, B., *et al.* PLINK: a tool set for whole-genome association and population-based
linkage analyses. *American journal of human genetics* **81,** 559–75 (3 2007).

 # Appendix

 ## A1 Variance in ancestry

We denote variance and covariance with $\mathbb{V}(.)$ and $\mathbb{C}(.)$ and used the expressions in [27] to generate the

expected value for the variance in ancestry, i.e., $\mathbb{V}(\theta)$. This is straightforward for the HI model, where at

time t $\mathbb{V}(\theta_t) = \mathbb{V}(\theta_{t-1})\frac{(1+P_{t-1})}{2}$. $P_t = Cor(\theta_m, \theta_f)$ measures the strength of assortative mating, i.e, the

correlation between the ancestry across mating pairs (θ_m, θ_f) at time t. For simplicity, we assumed this to

be constant in every generation, i.e. $P_t = P_{t-1} = P$ following [27]. Since our notation slightly differs from

[27], we re-derived the expression for $V(\theta_t)$ for the CGF model where population B receives a constant

amount q of gene flow from population A in every generation. Note, that $\mathbb{E}(\theta_t) = q + (1-q)\mathbb{E}(\theta_{t-1})$.

Then,

$$
\begin{aligned}
\mathbb{V}(\theta_t) = {}& \mathbb{E}(\theta_t^2) - \mathbb{E}(\theta_t)^2 \\
= {}& q + (1-q)\,\mathbb{E}\left[\left(\frac{\theta_{t-1}^m + \theta_{t-1}^f}{2}\right)\left(\frac{\theta_{t-1}^m + \theta_{t-1}^f}{2}\right)\right] - \{q + (1-q)\,\mathbb{E}(\theta_{t-1})\}^2 \\
= {}& q + \frac{(1-q)}{4}\{2\,\mathbb{E}(\theta_{t-1}^2) + 2\,\mathbb{E}(\theta_{t-1}^m\theta_{t-1}^f)\} - \{q^2 + 2q(1-q)\,\mathbb{E}(\theta_{t-1}) + (1-q)^2\,\mathbb{E}(\theta_{t-1})^2\} \\
= {}& q + \frac{1-q}{2}\,\mathbb{E}(\theta_{t-1}^2) + \frac{1-q}{2}\,\mathbb{E}(\theta_{t-1}^m\theta_{t-1}^f) - q^2 - 2q(1-q)\,\mathbb{E}(\theta_{t-1}) - (1-q)^2\,\mathbb{E}(\theta_{t-1})^2 \\
= {}& q(1-q) + \frac{1-q}{2}\{\mathbb{V}(\theta_{t-1}) + \mathbb{E}(\theta_{t-1})^2\} + \frac{1-q}{2}\{\mathbb{C}(\theta_{t-1}^m, \theta_{t-1}^f) + \mathbb{E}(\theta_{t-1})^2\} - 2q(1-q)\,\mathbb{E}(\theta_{t-1}) - \mathbb{E}(\theta_{t-1})^2 \\
= {}& q(1-q) + \frac{1-q}{2}\,\mathbb{V}(\theta_{t-1}) + \frac{1-q}{2}\,\mathbb{E}(\theta_{t-1})^2 + \frac{1-q}{2}P_{t-1}\,\mathbb{V}(\theta_{t-1}) + \frac{1-q}{2}\,\mathbb{E}(\theta_{t-1})^2 - 2q(1-q)\,\mathbb{E}(\theta_{t-1}) - \mathbb{E}(\theta_{t-1})^2 \\
= {}& q(1-q) + \frac{1-q}{2}\,\mathbb{V}(\theta_{t-1})\{1 + P_{t-1}\} + (1-q)\,\mathbb{E}(\theta_{t-1})^2 - 2q(1-q)\,\mathbb{E}(\theta_{t-1}) - (1-q)^2\,\mathbb{E}(\theta_{t-1})^2 \\
= {}& q(1-q)\,\mathbb{E}(\theta_{t-1})^2 + q(1-q)\{1 - 2\,\mathbb{E}(\theta_{t-1})\} + \frac{1-q}{2}\,\mathbb{V}(\theta_{t-1})\{1 + P_{t-1}\}
\end{aligned}
$$

 ## A2 Genetic variance

867 Let $y = g + e$, where y is the phenotypic value of an individual, g is the genotypic value, and e is random

868 error. We assume additive effects such that $g = \sum_{i=1}^m \beta_i x_i$ where β_i is the effect size of the i^{th} biallelic

869 locus and $x_i \in \{0, 1, 2\}$ is the number of copies of the trait-increasing allele. Then, the genetic variance

870 V_g is:

$$
V_g = \mathbb{V}(\sum_{i=1}^m \beta_i x_i) = \sum_{i=1}^m \beta_i^2\,\mathbb{V}(x_i) + \sum_{j\neq i} \beta_i\beta_j\,\mathbb{C}(x_i, x_j)
$$

871 In the following sections, we decompose $\mathbb{V}(x_i)$ and $\mathbb{C}(x_i, x_j)$ further as functions of ancestry.

 ### A2.1 $\mathbb{V}(x_i)$

873 We first derive $\mathbb{V}(x_i)$ as a function of ancestry (θ) using the law of total variance:

$$\mathbb{V}(x_i) = \mathbb{E}_\theta\{\mathbb{V}(x_i|\theta)\} + \mathbb{V}_\theta\{\mathbb{E}(x_i|\theta)\}$$

874 where $\mathbb{E}_\theta$ represents the expectation taken over θ.

875 A2.1.1 $\quad \mathbb{E}_\theta\{\mathbb{V}(x_i|\theta)\}$

876 We derive $\mathbb{V}(x_i|\theta)$ by further conditioning on the local ancestry at each locus.

$$\mathbb{V}(x_i|\theta) = \mathbb{E}_\gamma\{\mathbb{V}(x_i|\gamma,\theta)\} + \mathbb{V}\{\mathbb{E}_\gamma(x_i|\gamma,\theta)\}$$

877 where $\mathbb{E}_\gamma$ represents expectation taken over local ancestry. Since we are interested in the variance at
878 a single locus, we will ignore the subscript i and denote the frequency of the trait-increasing allele in
879 populations A and B with f^A and f^B, respectively.

$$
\begin{aligned}
\mathbb{E}_\gamma\{\mathbb{V}(x_i|\gamma,\theta)\} &= \mathbb{V}(x_i|\gamma=0,\theta)\,\mathbb{P}(\gamma=0|\theta) + \mathbb{V}(x_i|\gamma=1,\theta)\,\mathbb{P}(\gamma=1|\theta) + \mathbb{V}(x_i|\gamma=2,\theta)\,\mathbb{P}(\gamma=2|\theta) \\
&= 2\,f^B(1-f^B)(1-\theta)^2 + \{f^A(1-f^A) + f^B(1-f^B)\}2\theta(1-\theta) + 2\,f^A(1-f^A)\theta^2 \\
&= (2\,f^B - 2f^{A^2})(1-2\theta+\theta^2) + (f^A - f^{A^2} + f^B - f^{B^2})(2\theta-2\theta^2) + (2\,f^A - 2f^{A^2})\theta^2 \\
&= 2\,f^B - 2\theta\,f^B - 2f^{B^2} + 2\theta f^{B^2} + 2\theta\,f^A - 2\theta f^{A^2} \\
&= 2\,f^B(1-\theta) - 2f^{B^2}(1-\theta) + 2\theta\,f^A(1-f^A) \\
&= 2\,f^B(1-f^B)(1-\theta) + 2\theta\,f^A(1-f^A)
\end{aligned}
$$

880 To derive $\mathbb{V}\{\mathbb{E}_\gamma(x|\gamma,\theta)\}$, note that

$$
\begin{aligned}
\mathbb{E}_\gamma(x|\gamma,\theta) &= \mathbb{E}_\gamma\{\mathbb{E}(x|\theta)\} \\
&= \mathbb{E}(x|\gamma=0,\theta)\,\mathbb{P}(\gamma=0|\theta) + \mathbb{E}(x|\gamma=1,\theta)\,\mathbb{P}(\gamma=1|\theta) + \mathbb{E}(x|\gamma=2,\theta)\,\mathbb{P}(\gamma=2|\theta) \\
&= 2\theta\,f^A + 2(1-\theta)\,f^B
\end{aligned}
$$

881 And,

$$
\begin{aligned}
\mathbb{V}\{\mathbb{E}_\gamma(x|\gamma,\theta)\} &= [\mathbb{E}(x|\gamma=0,\theta) - \mathbb{E}(x|\theta)]^2\,\mathbb{P}(\gamma=0|\theta) \\
&\quad + [\mathbb{E}(x|\gamma=1,\theta) - \mathbb{E}(x|\theta)]^2\,\mathbb{P}(\gamma=1|\theta) \\
&\quad + [\mathbb{E}(x|\gamma=2,\theta) - \mathbb{E}(x|\theta)]^2\,\mathbb{P}(\gamma=2|\theta) \\
&= \theta^2\left[2\,f^A - \{2\theta\,f^A + 2(1-\theta)\,f^B\}\right]^2 \\
&\quad + 2\theta(1-\theta)\left[f^A + f^B - \{2\theta\,f^A + 2(1-\theta)\,f^B\}\right]^2 \\
&\quad + (1-\theta)^2\left[2\,f^B - \{2\theta\,f^A + 2(1-\theta)\,f^B\}\right]^2 \\
&= 2\theta(1-\theta)(f^A - f^B)^2
\end{aligned}
$$

882 Putting this together,

$$\begin{aligned}
\mathbb{E}_{\theta}\{\mathbb{V}(x_i|\theta)\} =& \mathbb{E}_{\theta}\{2\,f^{\mathrm{B}}(1-f^{\mathrm{B}})(1-\theta) + 2\theta\,f^{\mathrm{A}}(1-f^{\mathrm{A}}) + 2\theta(1-\theta)(f^{\mathrm{A}}-f^{\mathrm{B}})^2\} \\
=& 2\,f^{\mathrm{B}}(1-f^{\mathrm{B}})\{1-\mathbb{E}_{\theta}(\theta)\} + 2\,\mathbb{E}_{\theta}(\theta)\,f^{\mathrm{A}}(1-f^{\mathrm{A}}) + 2\,\mathbb{E}_{\theta}(\theta-\theta^2)(f^{\mathrm{A}}-f^{\mathrm{B}})^2 \\
=& 2\,f^{\mathrm{B}}(1-f^{\mathrm{B}})\{1-\mathbb{E}_{\theta}(\theta)\} + 2\,\mathbb{E}_{\theta}(\theta)\,f^{\mathrm{A}}(1-f^{\mathrm{A}}) + 2\{\mathbb{E}_{\theta}(\theta)-\mathbb{E}_{\theta}(\theta^2)\}(f^{\mathrm{A}}-f^{\mathrm{B}})^2 \\
=& 2\,f^{\mathrm{B}}(1-f^{\mathrm{B}})\{1-\mathbb{E}_{\theta}(\theta)\} + 2\,\mathbb{E}_{\theta}(\theta)\,f^{\mathrm{A}}(1-f^{\mathrm{A}}) + 2\{\mathbb{E}_{\theta}(\theta)-\mathbb{V}(\theta)-\mathbb{E}_{\theta}(\theta)^2\}(f^{\mathrm{A}}-f^{\mathrm{B}})^2 \\
=& 2\,f^{\mathrm{B}}(1-f^{\mathrm{B}})\{1-\mathbb{E}_{\theta}(\theta)\} + 2\,\mathbb{E}_{\theta}(\theta)\,f^{\mathrm{A}}(1-f^{\mathrm{A}}) + 2\,\mathbb{E}_{\theta}(\theta)(1-\mathbb{E}_{\theta}(\theta))(f^{\mathrm{A}}-f^{\mathrm{B}})^2 - 2\,\mathbb{V}(\theta)(f^{\mathrm{A}}-f^{\mathrm{B}})^2
\end{aligned}$$

883 **A2.1.2** $\mathbb{V}\{\mathbb{E}_{\theta}(x_i|\theta)\}$

884 Recall from the previous section that $\mathbb{E}_{\theta}(x_i|\theta) = 2\theta\,f^{\mathrm{A}} + 2(1-\theta)\,f^{\mathrm{B}}$. Then,

$$\begin{aligned}
\mathbb{V}\{\mathbb{E}_{\theta}(x_i|\theta)\} =& \mathbb{V}\{2\theta\,f^{\mathrm{A}} + 2(1-\theta)\,f^{\mathrm{B}}\} \\
=& 4\,\mathbb{V}(\theta)f^{A^2} + 4\,\mathbb{V}(1-\theta)f^{B^2} + 2\,\mathbb{C}(2\theta\,f^{\mathrm{A}}, 2(1-\theta\,f^{\mathrm{B}}) \\
=& 4\,\mathbb{V}(\theta)f^{A^2} + 4\,\mathbb{V}(1-\theta)f^{B^2} - 8\,f^{\mathrm{A}}\,f^{\mathrm{B}}\,\mathbb{V}(\theta) \\
=& 4\,\mathbb{V}(\theta)(f^{\mathrm{A}}-f^{\mathrm{B}})^2
\end{aligned}$$

885 We are now ready to express $\mathbb{V}(x_i)$:

$$\begin{aligned}
\mathbb{V}(x_i) =& 2\,f^{\mathrm{B}}(1-f^{\mathrm{B}})\{1-\mathbb{E}_{\theta}(\theta)\} + 2\,\mathbb{E}_{\theta}(\theta)\,f^{\mathrm{A}}(1-f^{\mathrm{A}}) + 2\,\mathbb{E}_{\theta}(\theta)(1-\mathbb{E}_{\theta}(\theta))(f^{\mathrm{A}}-f^{\mathrm{B}})^2 \\
& - 2\,\mathbb{V}(\theta)(f^{\mathrm{A}}-f^{\mathrm{B}})^2 + 4\,\mathbb{V}(\theta)(f^{\mathrm{A}}-f^{\mathrm{B}})^2 \\
=& 2\,\mathbb{E}_{\theta}(\theta)\,f_i^{\mathrm{A}}(1-f_i^{\mathrm{A}}) + 2\{1-\mathbb{E}_{\theta}(\theta)\}\,f_i^{\mathrm{B}}(1-f_i^{\mathrm{B}}) \\
& + 2\,\mathbb{E}_{\theta}(\theta)\{1-\mathbb{E}_{\theta}(\theta)\}(f_i^{\mathrm{A}}-f_i^{\mathrm{B}})^2 - 2\,\mathbb{V}(\theta)(f_i^{\mathrm{A}}-f_i^{\mathrm{B}})^2
\end{aligned}$$

886 Note, that we can also express $V(x_i)$ as:

$$\mathbb{V}(x_i) = 2f_i(1-f_i) + 2\,\mathbb{V}(\theta)(f_i^A - f_i^B)^2$$

887 where the second term is the contribution of population structure to the genetic variance at locus i.

888 **A2.2** $\mathbb{C}(x_i, x_j)$

889 We can derive $\mathbb{C}(x_i, x_j)$ using the law of total covariance:

$$\mathbb{C}(x_i, x_j) = \underset{\theta}{\mathbb{E}}\{\mathbb{C}(x_i, x_j|\theta)\} + \underset{\theta}{\mathbb{C}}\{\underset{\theta}{\mathbb{E}}(x_i|\theta), \underset{\theta}{\mathbb{E}}(x_j|\theta)\}$$

$$=0 + \mathbb{C}\{2\,f_i^A\,\theta + 2\,f_i^B(1-\theta), 2\,f_j^A\,\theta + 2\,f_j^B(1-\theta)\}$$

$$=\mathbb{C}(2\,f_i^A\,\theta, 2\,f_j^A\,\theta) + \mathbb{C}(2\,f_i^A\,\theta, 2\,f_j^B(1-\theta)) +$$

$$\mathbb{C}(2\,f_i^B(1-\theta), 2\,f_j^A\,\theta) + \mathbb{C}(2\,f_i^B(1-\theta), 2\,f_j^B(1-\theta))$$

$$=4\,\mathbb{V}(\theta)(f_i^A - f_i^B)(f_j^A - f_j^B)$$

890 $\mathbb{E}_\theta\{\mathbb{C}(x_i, x_j|\theta)\} = 0$ because we assume that the loci are unlinked and therefore, x_i and x_j are condi-
891 tionally independent. Putting this all together, we get the genetic variance in admixed populations as
892 presented in the main text:

$$V_g = \sum_{i=1}^{m} \beta_i^2 \,\mathbb{V}(x_i) + \sum_{j \neq i} \beta_i \beta_j \,\mathbb{C}(x_i, x_j)$$

$$= \sum_{i=1}^{m} \beta_i^2 2\,\underset{\theta}{\mathbb{E}}(\theta)\, f_i^A(1 - f_i^A) + \sum_{i=1}^{m} \beta_i^2 2\{1 - \underset{\theta}{\mathbb{E}}(\theta)\}\, f_i^B(1 - f_i^B)$$

$$+ \sum_{i=1}^{m} \beta_i^2 2\,\underset{\theta}{\mathbb{E}}(\theta)\{1 - \underset{\theta}{\mathbb{E}}(\theta)\}(f_i^A - f_i^B)^2 +$$

$$+ \sum_{i=1}^{m} \beta_i^2 2\,\mathbb{V}(\theta)(f_i^A - f_i^B)^2]$$

$$+ \sum_{j \neq i} \beta_i \beta_j 4\,\mathbb{V}(\theta)(f_i^A - f_i^B)(f_j^A - f_j^B)$$

893 The only difference being that in the main text we use $\mathbb{E}$ instead of $\mathbb{E}_\theta$ for simplicity. With two 'unad-
894 mixed' source populations with equal number of individuals, $\mathbb{E}(\theta) = 0.5$ and $\mathbb{V}(\theta) = \mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} = 0.25$
895 and V_g reduces to:

$$V_g = \mathbb{V}\left(\sum_{i=1}^{m} \beta_i x_i\right) = \sum_{i=1}^{m} \beta_i^2 \,\mathbb{V}(x_i) + \sum_{j \neq i} \beta_i \beta_j \,\mathbb{C}(x_i, x_j)$$

$$= \sum_{i=1}^{m} \beta_i^2 \left[f_i^A(1 - f_i^A) + f_i^B(1 - f_i^B)\right]$$

$$+ \sum_{i=1}^{m} \beta_i^2 (f_i^A - f_i^B)^2$$

$$+ \sum_{i \neq j} \beta_i \beta_j (f_i^A - f_i^B)(f_j^A - f_j^B)$$

896 A3 Genetic variance after correction for individual ancestry

897 It can also be helpful to decompose V_g into components of variance explained by and variance orthogonal
898 to ancestry:

$$\mathbb{V}(g) = \underbrace{\mathbb{V}\{\mathbb{E}_{\theta}(g|\theta)\}}_{\substack{\text{variance along} \\ \text{ancestry axis}}} + \underbrace{\mathbb{E}_{\theta}\{\mathbb{V}(g|\theta)\}}_{\substack{\text{variance orthogonal} \\ \text{to ancestry axis}}}$$

899 We can express the residual variance as:

$$
\begin{aligned}
\mathbb{E}_{\theta}\{\mathbb{V}(g|\theta)\} =& \mathbb{E}_{\theta}\{\mathbb{V}(\sum_{i=1}^{M} \beta_i^2 x_i|\theta)\} \\
=& \mathbb{E}_{\theta}\{\sum_{i=1}^{M} \beta_i^2 \, \mathbb{V}(x_i|\theta)\} + \mathbb{E}_{\theta}\{\sum_{i \neq j} \beta_i \beta_j \, \mathbb{C}(x_i, x_j|\theta) \\
=& \sum_{i=1}^{M} \beta_i^2 \, \mathbb{E}_{\theta}\{\mathbb{V}(x_i|\theta)\} + 0 \\
=& 2 \, \mathbb{E}_{\theta}(\theta) \sum_{i=1}^{M} \beta^2 \, f_{i}^{A}(1 - f_{i}^{A}) + 2\{1 - \mathbb{E}_{\theta}(\theta)\} \sum_{i=1}^{M} \beta^2 \, f_{i}^{B}(1 - f_{i}^{B}) \\
& + 2 \, \mathbb{E}_{\theta}(\theta) \sum_{i=1}^{M} \beta^2 \{1 - \mathbb{E}_{\theta}(\theta)\}(f_{i}^{A} - f_{i}^{B})^2 - 2 \, \mathbb{V}(\theta) \sum_{i=1}^{M} \beta^2 (f_{i}^{A} - f_{i}^{B})^2
\end{aligned}
$$

900 Note, that this represents the following components of V_g: (1.1) + (1.2) - (1.3).

A4 Haseman-Elston regression

902 The Haseman-Elston (HE) estimator of V_g is based on the regression of products of (centered) phenotypes
903 $y_k y_l$ for all pairs of individuals $k \neq l$ on the corresponding entries of the GRM (ψ) where $\psi_{kl} = \frac{\sum_{i=1}^{m} z_{ik} z_{il}}{m}$
904 and z_{ik} is the centered and scaled genotype of individual k for locus i. In this and the following sections,
905 we denote the effect sizes corresponding to the scaled genotypes as u_i and show the impact of alternative
906 scaling schemes of the genotype on the HE estimator.

$$
\begin{aligned}
\hat{V}_g =& \frac{\mathbb{C}(y_k y_l, \psi_{kl})}{\mathbb{V}(\psi_{kl})} \\
=& \frac{\mathbb{E}_{kl}(y_k y_l \psi_{kl}) - \mathbb{E}_{kl}(y_k y_l) \, \mathbb{E}_{kl}(\psi_{kl})}{\mathbb{E}_{kl}(\psi_{kl}^2) - \mathbb{E}(\psi_{kl})^2} \\
=& \frac{\mathbb{E}_{kl}(y_k y_l \psi_{kl})}{\mathbb{E}_{kl}(\psi_{kl}^2)} = \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^{m} u_i z_{ik} \sum_{i=1}^{m} u_i z_{il} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)} \\
=& \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^{m} u_i^2 z_{ik} z_{il} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)} + \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^{m} \sum_{j \neq i} u_i u_j z_{ik} z_{jl} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)}
\end{aligned}
$$

907 Where $\mathbb{E}_{kl}$ represents the expectation over all $k \times l$ pairwise comparisons between individuals and the
908 first and second terms represent the genic and LD components of the estimator.

909 A4.1   No directional LD

910 First, let's assume a genetic architecture where the effect sizes are random and there is no LD contribu-
911 tion, i.e., $\mathbb{E}_{ij}(u_i u_j) = 0$. We can further simplify the estimator as follows:

$$
\begin{aligned}
\hat{V}_g =& \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^m u_i^2 z_{ik} z_{il} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)} + \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^m \sum_{j\neq i} u_i u_j z_{ik} z_{jl} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)}\\
=& \frac{\mathbb{E}(u_i^2)\,\mathbb{E}_{kl}\left(\sum_{i=1}^m z_{ik} z_{il} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)} + \frac{\mathbb{E}(u_i u_j)\,\mathbb{E}_{kl}\left(\sum_{i=1}^m \sum_{j\neq i} z_{ik} z_{jl} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)}\\
=& \frac{\mathbb{E}_i(u_i^2)\,\mathbb{E}_{kl}(m\psi_{kl}^2)}{\mathbb{E}_{kl}(\psi_{kl}^2)} + 0 = m\,\mathbb{E}_{ij}(u_i^2)
\end{aligned}
$$

912 Thus, in the absence of directional LD, the estimator is a function of $\mathbb{E}(u_i^2)$.

913 A4.1.1   Scaling by $2f_i(1-f_i)$

914 With the standard scaling, the genotype at a given locus i is $z_i = \frac{x_i - 2f_i}{\sqrt{2f_i(1-f_i)}}$ where f_i is the frequency
915 of the allele in the population. Under the random-effects model, this is equivalent to saying that the
916 unscaled effects are $\beta_i \sim \mathcal{N}(0, \frac{\sigma_u^2}{2mf_i(1-f_i)})$ and the scaled effects are $u_i \sim \mathcal{N}(0, \frac{\sigma_u^2}{m})$. In a panmictic
917 population, $V_g = \sum_{i=1}^m \beta_i^2 \mathbb{V}(x_i) = \sum_{i=1}^m \frac{\sigma_u^2}{2mf_i(1-f_i)} 2f_i(1-f_i) = \sigma_u^2$. Thus, in a panmictic population,
918 $\mathbb{E}(\hat{V}_g) = \sigma_u^2 = V_g$, i.e., the standard scaling yields unbiased estimates of the genetic variance. In an
919 admixed population,

$$
\begin{aligned}
V_g =& \sum_{i=1}^m \beta_i^2 \{2f_i(1-f_i) + 2\,\mathbb{V}(\theta)(f_i^A - f_i^B)^2\}\\
=& \sum_{i=1}^m \frac{\sigma_u^2}{2mf_i(1-f_i)}\{2f_i(1-f_i) + 2\,\mathbb{V}(\theta)(f_i^A - f_i^B)^2\}\\
=& \frac{\sigma_u^2}{m}\sum_{i=1}^m \{1 + \mathbb{V}(\theta)\frac{(f_{\mathrm{i}}^A - f_{\mathrm{i}}^B)^2}{f_i(1-f_i)}\}\\
=& \sigma_u^2 + \underbrace{\mathbb{V}(\theta)\frac{\sigma_u^2}{m}\sum_{i=1}^m \frac{(f_{\mathrm{i}}^A - f_{\mathrm{i}}^B)^2}{f_i(1-f_i)}}_{\substack{\text{contribution of population structure}\\\text{to the genic variance}}}
\end{aligned}
$$

920 Thus, with the standard scaling, HE regression does not capture the contribution of population structure
921 and therefore, gives a biased estimate of V_g.

922 A4.1.2   Scaling by $\mathbb{V}(x_i)$

923 Next, we consider the case where the genotypes are standardized instead by the sample variance, i.e.,
924 $z_{kl} = \frac{x_{ik} - 2f_i}{\sqrt{\mathbb{V}(x_i)}}$ such that $\mathbb{V}(z_i) = 1$. We can derive $\mathbb{E}(u_i^2)$ corresponding to this scaling by noting that the
925 genetic variance is invariant under linear transformations of the genotype [25]:

$$\sum_{i=1}^{m} \beta_i^2 \, \mathbb{V}(x_i) = \sum_{i=1}^{m} u_i^2 \, \mathbb{V}(z_i)$$

$$m \, \mathbb{E}(u_i^2) = \sigma_u^2 + \mathbb{V}(\theta) \frac{\sigma_u^2}{m} \sum_{i=1}^{m} \frac{(f_i^{\mathrm{A}} - f_i^{\mathrm{B}})^2}{f_i(1 - f_i)}$$

$$\mathbb{E}(u_i^2) = \frac{\sigma_u^2}{m} + \mathbb{V}(\theta) \frac{\sigma_u^2}{m^2} \sum_{i=1}^{m} \frac{(f_i^{\mathrm{A}} - f_i^{\mathrm{B}})^2}{f_i(1 - f_i)}$$

Then, the HE estimator becomes:

$$
\begin{aligned}
\hat{V}_g &= m \, \mathbb{E}(u_i^2) \\
&= m \left(\frac{\sigma_u^2}{m} + \mathbb{V}(\theta) \frac{\sigma_u^2}{m^2} \sum_{i=1}^{m} \frac{(f_i^{\mathrm{A}} - f_i^{\mathrm{B}})^2}{f_i(1 - f_i)} \right) \\
&= \sigma_u^2 + \mathbb{V}(\theta) \frac{\sigma_u^2}{m} \sum_{i=1}^{m} \frac{(f_i^{\mathrm{A}} - f_i^{\mathrm{B}})^2}{f_i(1 - f_i)}
\end{aligned}
$$

Which provides an unbiased estimate of the genic variance. It's important to note that even though we assumed effect sizes under a random-effect model, the above result holds under a fixed-effect model as long as there is no directional LD. We discuss the implications of directional LD in the following section.

A4.2 Directional LD

Under the standard random-effect model, the effect sizes are assumed to be independent *in expectation*. We discussed in the main text how certain processes (e.g. selection and assortative mating) can induce directional LD across causal loci. But directional LD might arise even for neutral traits and under the random-effects model for any given realization of effects. This can lead to biases in both HE and GREML estimates of V_g, though the direction and reason for bias is different for the two methods. GREML does not have a closed-form solution so the exact estimand is difficult to derive. Here, we develop some intuition for HE regression.

A4.2.1 Scaling by $\mathbb{V}(x_i)$

To do this, let $\boldsymbol{u}' = [u_1, u_2, ..., u_m]$ represent the vector of a given realization of (fixed) effects corresponding to the standardized genotypes such that each locus contributes equally to σ_u^2, the genic variance, i.e., $u_i^2 = \frac{\sigma_u^2}{m}$. Let there be positive LD across loci such that all cross-product terms are $u_i u_j = \frac{\sigma_u^2}{m}$. Then, the genetic variance explained by all loci is:

$$
\begin{aligned}
V_g &= \sum_{i=1}^{m} u_i^2 \, \mathbb{V}(z_i) + \sum_{j \neq i} u_i u_j \, \mathbb{C}(z_i, z_j) \\
&= \sum_{i=1}^{m} u_i^2 + \sum_{j \neq i} u_i u_j \, \mathbb{C}(z_i, z_j) \\
&= \sigma_u^2 + \frac{\sigma_u^2}{m} \sum_{j \neq i} \mathbb{C}(z_i, z_j)
\end{aligned}
\tag{3}
$$

943 where $\mathbb{C}(z_i, z_j)$ is the LD between the i^{th} and j^{th} loci that ranges from 0 (no LD) to 1 (perfect LD).

944 Thus, the LD contribution to V_g ranges from 0 to $(m-1)\sigma_m^2$. In comparison, the HE estimator is:

$$
\begin{aligned}
\hat{V}_g &= \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^m u_i^2 z_{ik} z_{il} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)} + \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^m \sum_{j \neq i} u_i u_j z_{ik} z_{jl} \psi_{kl}\right)}{\mathbb{E}_{kl}(\psi_{kl}^2)} \\
&= \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^m \frac{\sigma_u^2}{m} z_{ik} z_{il} \sum_{w=1}^m z_{wk} z_{wl}/m\right)}{\mathbb{E}_{kl}(\sum_{i=1}^m z_{ik} z_{il}/m \sum_{w=1}^m z_{wk} z_{wl}/m)} + \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^m \sum_{j \neq i} \frac{\sigma_u^2}{m} z_{ik} z_{jl} \sum_{w=1}^m z_{wk} z_{wl}/m\right)}{\mathbb{E}_{kl}(\sum_{i=1}^m z_{ik} z_{il}/m \sum_{w=1}^m z_{wk} z_{wl}/m)} \\
&= \sigma_u^2 + \sigma_u^2 \frac{\mathbb{E}_{kl}\left(\sum_{i=1}^m \sum_{j \neq i} z_{ik} z_{jl} z_{wk} z_{wl}\right)}{\mathbb{E}_{kl}(\sum_{i=1}^m z_{ik} z_{il} \sum_{w=1}^m z_{wk} z_{wl})}
\end{aligned}
\tag{4}
$$

This shows that the bias due to directional LD in the HE estimate of V_g does not come from the genic,

but the LD component. When there is no LD, e.g. if the population has reached equilibrium after

generations of random mating, this component goes to zero and both the estimate and V_g converge to

the same value – the genic variance. The LD component is maximum when the i^{th} and j^{th} loci are

in perfect LD. In this case, i and j are exchangeable and the second term of the estimator reduces

to $(m-1)\sigma_m^2$. Thus, HE regression should give an unbiased estimate of V_g, even in the presence of

directional LD, but only when LD is perfect. For any other value $0 < C(z_i, z_j) < 1$, the estimate is

biased (Fig. A1). An interpretable, analytical derivation of the second term in Eq. 4 is complicated but

we illustrate the bias with simulations below.

For unlinked markers, $\mathbb{C}(z_i, z_j)$ is a function of $4\mathbb{V}(\theta)(f_i^A - f_i^B)(f_j^A - f_j^B)$ (see A2.1). Perfect LD arises

when (i) both $f_i^A - f_i^B = 1$ and $f_j^A - f_j^B = 1$ and (ii) $\mathbb{V}(\theta)$ is maximum, which occurs at the time of

admixture when source populations mix equally, i.e, $\mathbb{E}(\theta) = 0.5$. To generate a range of LD, we simulated

an admixed population ($N = 1,000$) with equal number of individuals from populations A and B. Thus,

$4\mathbb{V}(\theta) = 4\mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} = 1$. We simulated genotypes for each individual at 50 'causal' loci where

the difference between the frequencies in the source populations, $f_i^A - f_i^B \in [0, 1]$ with the condition

that $\frac{f_i^A + f_i^B}{2} = 0.5$. We assigned each locus the same effect size (on the variance-standardized scale) of

$+1/\sqrt{m}$ summing up to a genic variance of 1. The positive sign ensures positive LD across loci, i.e,

all off-diagonal elements of $\boldsymbol{uu'}$ are set to $1/m$. For each simulation, we computed the expected and

estimated LD component using the second terms in Eqs. 3 and 4, respectively, and averaged the results

over 100 replications.

A4.2.2   Scaling by LD

In the main text, we showed that standardizing the genotypes at a locus by its covariance with other loci

accounts for the bias for GREML and HE estimators. More specifically, the 'LD-scaled' genotypes can

be written as $\boldsymbol{Z} = (\boldsymbol{X} - 2\boldsymbol{P})\boldsymbol{U}^{-1}$ where $\boldsymbol{P}$ is an $n \times m$ matrix such that all elements of the i^{th} column

contain the frequency of the i^{th} SNP and $\boldsymbol{U}$ is the (upper triangular) square root of the LD matrix, i.e,

$\boldsymbol{\Sigma} = \boldsymbol{U'U}$. Under this scheme, the standardized genotypes are uncorrelated and therefore, the second

term in Eqs. 3 and 4 are zero. This reduces the estimator to the first term, representing the sum of

squares of effect sizes, i.e. $\boldsymbol{u'u} = \sum_{i=1}^m u_i^2$. The effect sizes corresponding to the LD scaled genotypes

Figure A1: The behavior of the LD contribution (y-axis) to the genetic variance (red) and the Haseman-Elston regression estimate (blue) as a function of LD, i.e. $\mathbb{C}(z_i, z_j)$ (x-axis). Each point represents the contribution calculated from a random draw of genotypes, given $\mathbb{C}(z_i, z_j) \propto (f_i^A - f_i^B)(f_j^A - f_j^B)$. The red line represents the expected LD contribution and the black dashed line represents the contribution expected in the case of perfect LD.

973 are $\boldsymbol{u} = \boldsymbol{U\beta}$ and the sum of squares is:

$$\boldsymbol{u}'\boldsymbol{u} = (\boldsymbol{U\beta})'\,(\boldsymbol{U\beta}) = \boldsymbol{\beta}'\boldsymbol{U}'\boldsymbol{U\beta} = \boldsymbol{\beta\Sigma\beta} = \sum_{i=1}^{m}\sum_{j=1}^{m}\beta_i\beta_j\,\mathbb{C}(x_i, x_j)$$

974 Which captures both the genic and LD contributions and therefore, provides an unbiased estimate of V_g.

A4.3 Genetic variance after correction for individual ancestry

976 In GREML it is common to include ancestry or principal components of the GRM to correct for popu-
977 lation structure, which can lead to inflated estimates of heritability [4, 7]. In HE regression, a moment-
978 matching approach (MMHE) can be used to account for covariates [37, 59]. To explain this, we use
979 matrix notation. Recall the generative model $\boldsymbol{y} = \boldsymbol{g} + \boldsymbol{e} = \boldsymbol{Zu} + \boldsymbol{e}$ where $\boldsymbol{y}$ is an $n \times 1$ (centered)
980 phenotype vector, $\boldsymbol{Z}$ is $n \times m$ matrix of standardized genotypes, and $\boldsymbol{u}$ is an $m \times 1$ vector of effect sizes.
981 In the absence of covariates, HE regression estimates $\hat{\sigma}_u^2$ and $\hat{\sigma}_e^2$ by regressing the empirical covariance
982 of the phenotypes onto the GRM: $\text{vec}(\boldsymbol{yy}') = \text{vec}(\boldsymbol{Zuu}'\boldsymbol{Z}') + \text{vec}(\boldsymbol{ee}')$ where $\text{vec}(.)$ is the vectorization
983 operator. If the effects are independent, this simplifies to $\sigma_u^2\,\text{vec}(\boldsymbol{\psi}) + \sigma_e^2\,\text{vec}(\boldsymbol{I})$. Then $\hat{\sigma}_u^2$ and $\hat{\sigma}_e^2$ can be
984 obtained from ordinary least squares (OLS).
985 Conceptually, the MMHE approach adjusts for covariates (e.g. sex, age, and ancestry) by projecting them
986 out of both the phenotypes and genotypes [37]. We denote the projection matrix as $\boldsymbol{P} = \boldsymbol{C}(\boldsymbol{C}'\boldsymbol{C})^{-1}\boldsymbol{C}'$
987 where $\boldsymbol{C}$ is the $n \times q$ design matrix of covariates. In our case, $\boldsymbol{C} = \begin{bmatrix} \boldsymbol{1} & \boldsymbol{\theta} \end{bmatrix}$ where $\boldsymbol{\theta}$ is the $n \times 1$ vector
988 of individual ancestry. Then, let $\dot{\boldsymbol{y}} = \boldsymbol{My}$ and $\dot{\boldsymbol{Z}} = \boldsymbol{MZ}$ can be thought of as the residuals of a
989 regression of $\boldsymbol{y}$ and $\boldsymbol{Z}$, respectively, on $\boldsymbol{\theta}$ and $\dot{\boldsymbol{\psi}} = \dot{\boldsymbol{Z}}\dot{\boldsymbol{Z}}'$ as the corresponding kinship matrix. Then, the
990 ancestry-adjusted estimate of σ_u^2 can be obtained by regressing $\text{vec}(\dot{\boldsymbol{y}}\dot{\boldsymbol{y}}')$ on $\text{vec}(\dot{\boldsymbol{\psi}}')$:

$$\begin{aligned}
\text{vec}(\dot{\boldsymbol{y}}\dot{\boldsymbol{y}}') &= \text{vec}(\dot{\boldsymbol{g}}\dot{\boldsymbol{g}}') + \text{vec}(\dot{\boldsymbol{e}}\dot{\boldsymbol{e}}') \\
&= \text{vec}((\boldsymbol{MZu})(\boldsymbol{MZu})') + \sigma_e^2\boldsymbol{I} \\
&= \text{vec}(\boldsymbol{MZuu}'\boldsymbol{Z}'\boldsymbol{M}) + \sigma_e^2\boldsymbol{I} \\
&= \text{vec}(\boldsymbol{MZ}\sigma_u^2\boldsymbol{Z}'\boldsymbol{M}) + \sigma_e^2\boldsymbol{I} \\
&= \text{vec}(\dot{\boldsymbol{Z}}\sigma_u^2\dot{\boldsymbol{Z}}') + \sigma_e^2\boldsymbol{I} \\
&= \sigma_u^2\,\text{vec}(\dot{\boldsymbol{\psi}}) + \sigma_e^2\boldsymbol{I}_{(n-q)}
\end{aligned}$$

991 The simplification in line 3 follows if we assume that (i) the off-diagonal terms of $\boldsymbol{uu}'$ are 0, i.e., the
992 effects are independent, and (ii) all loci contribute equally to the genetic variance. The OLS estimates
993 of σ_u^2 and σ_e^2 can be obtained by solving the following linear system [11, 37]:

$$\begin{bmatrix} \text{tr}(\boldsymbol{M\psi M\psi}) & \text{tr}(\boldsymbol{M\psi}) \\ \text{tr}(\boldsymbol{\psi}) & n-q \end{bmatrix} \begin{bmatrix} \sigma_u^2 \\ \sigma_e^2 \end{bmatrix} \begin{bmatrix} \boldsymbol{y}'\boldsymbol{M\psi My} \\ \boldsymbol{y}'\boldsymbol{My} \end{bmatrix}$$

994 where $\text{tr}(.)$ represents the trace of the matrix. In practice, we use the more efficient algorithm in Ge *et*
995 *al.* (2017) [37], which we used to estimate $\hat{\sigma}_u^2$.

 A5 LD Score regression

 LD score regression (LDSC) estimates h_{snp}^2 under a random-effects model from GWAS summary statistics

 using method-of-moments. Here, we provide intuition for how LDSC can be biased in the presence of

 directional LD. We define χ_k^2 as the marginal association statistic of the k^{th} marker and $l_k = \sum_{w=1}^{W} r_{wk}^2$

 as its LD score. Thus, LDSC estimates h_{snp}^2 from the regression slope of χ_k^2 on l_k over all k markers.

 To understand this, let's denote the estimated marginal effect of the k^{th} marker as a function of its

 true marginal effect and some estimation error: $\hat{\tau}_k = \tau_k + \varepsilon_k$ assuming for simplicity that there is no

 residual bias due to stratification in the upstream GWAS. Then, $\mathbb{E}(\hat{\tau}_k) = \tau_k$ and $\mathbb{E}(\hat{\tau}_k^2) = \tau_k^2 + \mathbb{E}(\varepsilon^2) =$

 $\tau_k^2 + SE^2(\hat{\tau}_k)$. Furthermore, $\tau_k = \sum_{i=1}^{m} r_{ik} u_i$, where r_{ik} is the LD between the k^{th} marker and the i^{th}

 causal locus. If the individual causal effects are small, then $SE^2(\hat{\tau}_k) \approx 1/n$ and

$$\mathbb{E}(\chi_k^2) = \frac{\mathbb{E}(\hat{\tau}_k^2)}{SE^2(\hat{\tau}_k)} = \frac{\tau_k^2 + SE^2(\hat{\tau}_k)}{SE^2(\hat{\tau}_k)} = \frac{\tau_k^2 + 1/n}{1/n} = n\tau_k^2 + 1$$

 The expected slope of LD Score regression is:

$$\hat{\beta}_{\chi^2,l} = \frac{\mathbb{C}(\chi_k^2, l_k)}{\mathbb{V}(l_k)} = \frac{\mathbb{C}(n\tau_k^2 + 1, l_k)}{\mathbb{V}(l_k)} = \frac{\mathbb{C}\left((\sum_{i=1}^{m} r_{ik} u_i)^2, l_k\right)}{\mathbb{V}(l_k)}$$

$$= \frac{n\,\mathbb{C}\left(\sum_{i=1}^{m} r_{ik}^2 u_i^2 + 2\sum_{i=1}^{m}\sum_{j<i}^{m} r_{ik} r_{jk} u_i u_j, \sum_{i=1}^{m} r_{ik}^2\right)}{\mathbb{V}(l_k)}$$

$$= \frac{n\,\mathbb{C}\left(\sum_{i=1}^{m} r_{ik}^2 u_i^2, \sum_{i=1}^{m} r_{ik}^2\right) + n\,\mathbb{C}\left(2\sum_{i=1}^{m}\sum_{j<i}^{m} r_{ik} r_{jk} u_i u_j, \sum_{i=1}^{m} r_{ik}^2\right)}{\mathbb{V}(l_k)}$$

 As with GREML and HE regression, LDSC assumes a polygenic, random-effects model where $u_i \sim$

 $\mathcal{N}(0, \frac{\sigma_u^2}{m})$. As such, the cross-product term inside the brackets is zero over variants i and j, and the slope

 reduces to:

$$\hat{\beta}_{\chi^2,l} = \frac{n\,\mathbb{C}\left(\sum_{i=1}^{m} r_{ik}^2 u_i^2, \sum_{i=1}^{m} r_{ik}^2\right)}{\mathbb{V}(l_k)} = \frac{n\sigma_u^2}{m}$$

 from which $\hat{V}_g$ and therefore, $\hat{h}_{snp}^2$ can be derived. To understand the behavior of LDSC under directional

 LD, consider a simple generative model with m causal variants, each with the same scaled (fixed) effect u

 in positive LD such that $V_g = \sum_{i=1}^{m} u_i^2\,\mathbb{V}(z_i) + 2\sum_{i=1}^{m}\sum_{j<i} \mathbb{C}(z_i, z_j) = \sum_{i=1}^{m} u_i^2 + 2\sum_{i=1}^{m}\sum_{j<i} u_i u_j r_{ij} =$

 $mu^2 + 2u^2 \sum_{i=1}^{m}\sum_{j<i} r_{ij}$. We further assume that causal variants are unlinked, that any LD between

 them is due to population structure, and that each causal variant is tagged by W markers. The marginal

 effect of a marker is a function of its LD with the causal loci, i.e., $\tau_k = \sum_{i=1}^{m} r_{ik} u_i = u \sum_{i=1}^{m} r_{ik}$.

 However, if, as per standard practice, individual ancestry was included as a covariate in the GWAS,

 then the marginal effect of the marker will only absorb the effect of the causal variant nearby but not

 of variants in LD due to population structure. Thus, if we note r_{kc} as the LD between the k^{th} marker

 and the causal variant in its vicinity, $\tau_k = u r_{kc}$, $\tau_k^2 = r_{kc}^2 u^2$, and $\mathbb{E}(\chi_k^2) = n r_{kc}^2 u^2 + 1$. In practice, LDSC

 regresses χ_k^2 on $l_k = \sum_{w=1}^{W} r_{kw}^2$:

$$LDSC = \frac{\mathbb{C}(\chi_k^2, l_k)}{\mathbb{V}(l_k)} = \frac{\mathbb{C}(\chi_k^2, \sum_w^W r_{kw}^2)}{\mathbb{V}(l_k)} = \frac{\sum_w^W \mathbb{C}(\chi_k^2, r_{kw}^2)}{\mathbb{V}(l_k)}$$
$$= \frac{\mathbb{C}(\chi_k^2, r_{kc}^2)}{\mathbb{V}(l_k)} + \frac{\sum_{w \neq c} \mathbb{C}(\chi_k^2, r_{kw}^2)}{\mathbb{V}(l_k)} = \frac{nu^2 r_{kc}^2}{\mathbb{V}(l_k)}$$

A6 Effect size of local ancestry

We define local ancestry $\gamma_i \in \{0, 1, 2\}$ as the number of alleles at locus i that trace their ancestry to population A. Thus, the local ancestry at locus i in individual k is a Binomial random variable with $\mathbb{E}(\gamma_{i,k}) = 2\theta_k$. We define the ancestry value of an individual as the weighted sum of their local ancestry: $\sum_{i=1}^m \phi_i \gamma_i$ where $\phi_i = \beta_i(f_i^B - f_i^A)$.

To show this, note that $\phi = \mathbb{E}(y|\gamma = 1) - \mathbb{E}(y|\gamma = 0)$ where $\mathbb{E}(y|\gamma = 1) = \int_{-\infty}^{\infty} yh(y|\gamma = 1)$ and h is a density function. Our goal is to express ϕ in terms of β, which is equal to $\mathbb{E}(y|x = 1) - \mathbb{E}(y|x = 0)$. Furthermore, $\mathbb{E}(y|x = 1) = \int_{-\infty}^{\infty} yh(y|x = 1)$. We can express $h(y|\gamma)$ in terms of $h(y|x)$ as follows:

$$h(y|\gamma = 1) = h(y|x = 0)\,\mathbb{P}(x = 0|\gamma = 1) + h(y|x = 1)\,\mathbb{P}(x = 1|\gamma = 1) + h(y|x = 2)\,\mathbb{P}(x = 2|\gamma = 1)$$
$$= h(y|x = 0)2(1 - f^A)(1 - f^A) + h(y|x = 1)\{f^A(1 - f^B) + f^B(1 - f^A)\} + h(y|x = 2)2\,f^A\,f^B$$

$$\mathbb{E}(y|\gamma = 1) = \int_{-\infty}^{\infty} yh(y|\gamma = 1)dy$$
$$= (1 - f^A)(1 - f^B)\int_{-\infty}^{\infty} yh(y|x = 0)dy$$
$$\quad + \{f^A(1 - f^B) + f^B(1 - f^A)\}\int_{-\infty}^{\infty} yh(y|x = 1)dy$$
$$\quad + f^A f^B \int_{-\infty}^{\infty} yh(y|x = 2)dy$$
$$= (1 - f^A)(1 - f^B)\,\mathbb{E}(y|x = 0) + \{f^A(1 - f^B) + f^B(1 - f^A)\}\,\mathbb{E}(y|x = 1) + f^A f^B\,\mathbb{E}(y|x = 2)$$
$$= 0 + \{f^A(1 - f^B) + f^B(1 - f^A)\}\beta + f^A f^B\,2\beta$$
$$= \beta\,f^A + \beta\,f^B$$

Similary, $\mathbb{E}(y|\gamma = 0) = 2\beta\,f^B$ and $\phi = \mathbb{E}(y|\gamma = 1) - \mathbb{E}(y|\gamma = 0) = \beta(f^B - f^A)$

A7 Genetic variance due to local ancestry

$$V_\gamma = \mathbb{V}(\sum_{i=1}^m \phi_i \gamma_i)$$
$$= \sum_{i=1}^m \phi_i^2\,\mathbb{V}(\gamma_i) + \sum_{i=1}^m \sum_{j \neq i}^m \phi_i \phi_j\,\mathbb{C}(\gamma_i, \gamma_j) \tag{5}$$

We use the law of total variance and covariance to derive $\mathbb{V}(\gamma_i)$ and $\mathbb{C}(\gamma_i, \gamma_j)$:

$$\mathbb{V}(\gamma_i) = \mathbb{E}\{\mathbb{V}(\gamma_i|\theta)\} + \mathbb{V}\{\mathbb{E}(\gamma_i|\theta)\}$$

$$= \mathbb{E}\{2\theta(1-\theta)\} + \mathbb{V}(2\theta)$$

$$= 2\mathbb{E}(\theta) - 2\mathbb{E}(\theta^2) + 4\mathbb{V}(\theta)$$

$$= 2\mathbb{E}(\theta) - 2\mathbb{V}(\theta) - 2\mathbb{E}(\theta)^2 + 4\mathbb{V}(\theta)$$

$$= 2\mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} + 2\mathbb{V}(\theta)$$

$$\mathbb{C}(\gamma_i, \gamma_j) = \mathbb{E}\{\mathbb{C}(\gamma_i, \gamma_j|\theta)\} + \mathbb{C}\{\mathbb{E}(\gamma_i, \gamma_j|\theta)\}$$

$$= 0 + \mathbb{C}(2\theta, 2\theta) = 4\mathbb{V}(\theta)$$

$$V_\gamma = 2\mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} \sum_{i=1}^{m} \phi_i^2 + 2\mathbb{V}(\theta) \sum_{i=1}^{m} \phi_i^2 + 4\mathbb{V}(\theta) \sum_{i=1}^{m} \sum_{j \neq i} \phi_i \phi_j$$

$$= 2\mathbb{E}(\theta)\{1 - \mathbb{E}(\theta)\} \sum_{i=1}^{m} \beta_i^2 (f_i^{\mathrm{B}} - f_i^{\mathrm{A}})^2$$

$$+ 2\mathbb{V}(\theta) \sum_{i=1}^{m} \beta_i^2 (f_i^{\mathrm{B}} - f_i^{\mathrm{A}})^2$$

$$+ 4\mathbb{V}(\theta) \sum_{i=1}^{m} \sum_{j \neq i} \beta_i \beta_j (f_i^{\mathrm{B}} - f_i^{\mathrm{A}})(f_j^{\mathrm{B}} - f_j^{\mathrm{A}})$$

 Supplement

Figure S1: The behavior of the four components of genetic variance in admixed populations under the (A) HI and (B) CGF models. We assume that the mean ancestry proportion in the population is 0.5. The solid lines represent values observed in simulations averaged across ten replicates and the dotted lines represent the expected values based on Eq. 1 of the main text. The red and blue lines represent values for traits 1 and 2, respectively. P indicates the strength of assortative mating. P=0.6 is missing for simulations run for 50 and 100 generations and $\theta \in \{0.1, 0.2\}$ due to the difficulty in finding mate pairs (Methods).

Figure S2: The behavior of the genetic variance due to local ancestry in admixed populations under the (A) HI and (B) CGF models. The solid lines represent values observed in simulations averaged across ten replicates and the dotted lines represent the expected values based on Eq. 1 of the main text. The red and blue lines represent values for traits 1 and 2, respectively. P indicates the strength of assortative mating. P=0.6 is missing for simulations run for 50 and 100 generations and $\theta \in \{0.1, 0.2\}$ due to the difficulty in finding mate pairs (Methods).

Figure S3: The behavior of GREML estimates of SNP heritability ($\hat{h}^2_{snp}$) in admixed populations under the HI (left column) and CGF (right column) models either without (A-C) or with (D-F) individual ancestry as a fixed effect. The solid lines represent $\hat{h}^2_{snp}$ averaged across ten replicates, with red and blue colors representing estimates for traits under divergent and stabilizing selection, respectively. (A, D) show the behavior of $\hat{h}^2_{snp}$ for the default scaling, (B, E) shows $\hat{h}^2_{snp}$ when the genotype at a locus is scaled by its sample variance ($\mathbb{V}(x)$ scaled), and (C, F) when it is scaled by the sample covariance (LD scaled). The shaded area represents the 95% confidence bands generated by bootstrapping (sampling with replacement 100 times) the point estimate reported by GCTA. The black dotted lines represent the expected heritability value given the simulation settings ($h^2 = 0.8$). P indicates the strength of assortative mating

Figure S4: Distribution of the total genetic variance (left), genic variance (middle), and LD component (right) for a neutral trait simulated by drawing effects for 10, 100, or 1,000 causal variants in ASW. The total genetic variance is the sum of the genic and LD components.

Figure S5: The effect of directional LD on Haseman-Elston estimate of genetic variance (V_g). Each individual point is an independent simulation where the effects were drawn from a normal distribution and applied to genotypes from an admixed population (Methods). The solid red line shows the $y = x$ line and the color of each point represents the contribution of LD to V_g.

Figure S6: Behavior of LDSC estimates of V_g under the HI and CGF admixture models as a function of time since admixture (x-axis) for traits under directional (blue) and stabilizing selection (red) and with $p \in \{0.01, 0.05, 0.1\}$ proportion of causal variants. The dashed lines indicate (A) the simulated V_g whereas the solid lines show the estimated value. Dotted horizontal lines indicate the expected genic variance. Shaded ribbons indicate the 95% CI across ten replicates.

Figure S7: Expected vs Observed effect sizes of 87,564 variants on chromosome 2 for a GWAS carried out in 5,000 admixed individuals simulated under the HI and CGF models (panel columns) for generations 2 and 20 (panel rows) since admixture. Variants are grouped into bins and each bin is colored by the (A) mean χ^2, (B) mean LD score, (C) no. of variants, (D) mean MAF, (E) mean CEU frequency, and (F) mean YRI frequency across variants in that bin. The $\mathbb{E}(\hat{\beta}_j)$ of the j^{th} SNP was computed as $\sum_{i=1}^m r_{ij}\beta_i$ where β_i is the simulated effect of the i^{th} causal variant and r_{ij} is the genotypic correlation between them. The plot shows overdispersion in $\hat{\beta}$ due to population structure, especially in the 2nd generation of the CGF model. This appears to be driven by incoming variants that are common in CEU (E) but rare or absent in YRI (F).

Figure S8: (A) GREML and (B) HE estimates of V_g (y-axis) on simulated genotype data of 5,000 admixed individuals for chromosome 2 under the HI and CGF models as a function of time since admixture (x-axis) and trait architecture (color). Results are also shown for varying proportion of causal variants (p) and whether or not ancestry (20 PCs) were included as covariates in the model. GREML and HE estimates are consistent with analytical expectations and simulations of unlinked markers.

Figure S9: The LD contribution to the variance explained by variant pairs for (A) neutrophil counts (NEU), (B) white blood count (WBC), (C) monocyte count (MON), and (D) mean corpuscular hemoglobin (MCH). Only chromosomes where we suspected there was a disproportionate contribution to the variance explained are shown.

Figure S10: Effect sizes estimated in the AoU cohort compared to effect sizes from original GWAS summary statistics (Methods). Each β is colored by the amount of variance explained by that variant with low variance explained shown in blue and higher variance explained shown in red. Variance explained is computed as $\beta^2 2f(1-f)$, where f is the minor allele frequency in the African American AoU cohort. The $y = x$ line is shown as a dotted black line and the gray lines are 95% confidence intervals for each β in the re-estimated cohort.