Resistance and resiliency of soil microbiomes under climate change

Julia A. Boyle[1], Bridget K. Murphy[2,3], Ingo Ensminger[1,2,3], John R. Stinchcombe[1,4]*, Megan. E.

Frederickson[1]*

(*) Equal contribution

1. Department of Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks St.,

Toronto, ON, Canada, M5S3B2

2. Department of Biology, University of Toronto, Mississauga, ON, Canada, L5L1C8

3. Department of Cell and Systems Biology, University of Toronto, Toronto, ON, Canada,

M5S1A1

4. Koffler Scientific Reserve, University of Toronto, Toronto, ON, Canada, M5S3B2

Corresponding author information: Julia Boyle, julia.boyle@mail.utoronto.ca, Department of
Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks St., Toronto, ON,
Canada, M5S3B2

Keywords: resilience; resistance; climate change; fungi; bacteria; legacy effects; symbiosis; soil

microbiome

Abstract

Soil microbiomes play key roles in plant productivity and nutrient cycling, and we need

to understand whether and how they will withstand the effects of global climate change. We

exposed *in situ* soil microbial communities to multiple rounds of heat, drought, or both

treatments, and profiled microbial communities with 16S rRNA and ITS amplicon sequencing

during and after these climatic changes, and then tested how domain and symbiotic lifestyle

affected response. Fungal community composition strongly shifted due to drought and its legacy.

In contrast, bacterial community composition resisted change during the experiment, but still was

affected by the legacy of drought. We identified fungal and bacterial taxa with differential

abundance due to heat and drought and found that taxa affected during climate events are not

necessarily the taxa affected in recovery periods, showing the complexity and importance of

legacy effects. Additionally, we found evidence that symbiotic groups of microbes important to

plant performance respond in diverse ways to climate treatments and their legacy, suggesting

plants may be impacted by past climatic events like drought and warming even if they do not

experience the event themselves.

38
Introduction

Anthropogenic climate change is increasing both global temperature and the frequency

and severity of weather events like drought (Thuiller 2007). Soil microbial communities are

among the largest and most diverse ecosystems on earth (Singh et al. 2009) and their response to

climate change will affect elemental cycling, greenhouse gas production, plant-soil interactions,

and ecosystem services and agriculture (Dubey et al. 2019; Jansson & Hofmockel 2020; ASM

Report 2022). Here, we examine the effect of repeated heat and drought events on a soil

microbiome across time, characterizing how fungal and bacterial communities resist, and recover

from, heat and drought events.

Resistance in a microbial community represents how quickly and by what magnitude a

community responds to shifting environmental conditions, whereas recovery and resilience can

be measured by whether community composition reflects historical conditions (Bissett et al.

2013; Hartmann et al. 2013; Martiny et al. 2017). Microbial communities can resist more or less

depending on the type and duration of environmental change (Schimel et al. 2007; Zhou et al.

2020), with evidence of microbes being more resistant to short-term than long-term changes (Pec

et al. 2021). After an environmental shift, microbial communities are often assumed to have high

resilience due to short microbial generation times (Allison and Martiny 2008), but empirical

evidence is mixed, as discussed below. Soil ecosystems exhibit differing microbial resistance and

resilience to environmental change (Waldrop & Firestone 2006). Ecosystems characterized by

natural fluctuations, such as northern temperate environments that experience large differences

between winter low and summer high temperatures, provide an interesting opportunity to test

microbial resistance and resilience because microbes are expected to be adapted to both

temperature extremes.

Microbial communities display highly variable responses to warming and drought.

Microbial biomass can increase (Castro et al. 2010) or decrease (Allison & Treseder 2008) in

response to heat, with a decrease in microbial biomass accompanying a loss of overall soil

carbon (Melillo et al. 2017). Heat can increase species richness (Allison & Treseder 2008) or

decrease it (Sheik et al. 2011), yet a meta-analysis of studies found no overall trend of warming

or altered precipitation on microbial alpha diversity (Zhou et al. 2020). Instead, both heat and

drought often shift microbial community composition (Allison & Treseder 2008; Castro et al.

2010; Zhou et al. 2020; Pec et al. 2021). Drought and heat combined can have more of an effect

on microbes than drought alone (Bérard et al. 2011), highlighting the importance of studying

multiple climatic factors and their interactions; yet multifactor studies remain rare (Zhou et al.

2020). Furthermore, studies measuring the persistence of these effects, i.e., a community's

resilience, show mixed results; some find no persisting legacy effects from climate events on soil

community composition (Rousk et al. 2013; Dacal et al. 2022), while others find lasting legacy

effects (Averill et al. 2016; Martiny et al. 2017; de Nijs et al. 2019). More empirical evidence on

how resilient microbial communities are to climate events is needed (Jansson & Hofmockel

2020) and we need to investigate factors that underlie microbial resilience itself, given that

microbes are a major contributor to global nutrient cycling of carbon, phosphorus, and nitrogen

(ASM Report 2022).

Differences in resistance and resilience are possible among component members of the

microbiome, due to the vast taxonomic and functional diversity of soil microbes (Singh et al.

2009). For example, bacteria are generally fast-growing, with short generation times and the

ability to horizontally transfer potentially adaptive genes (Ochman et al. 2000), while fungi show

a myriad of life history strategies, many of which require multiple stages (Andrews 1992) and

thus longer generation times. In the soil food web, bacteria are considered a 'fast' energy channel

that cycles nutrients rapidly, while fungi are considered a 'slow' energy channel that cycles

nutrients slowly (Rooney et al. 2006; Bardgett & Caruso 2020). Consistent with this

classification, turnover times of soil fungi are estimated to be an order of magnitude longer than

those of soil bacteria (Rousk & Bååth 2011). Slow fungal turnover times mean that it could take

longer to observe a shift in fungal than in bacterial community composition, but that the effect of

climate stressors on fungi may persist. We therefore hypothesized that fungal communities may

be more resistant but less resilient than bacterial communities to climatic perturbations.

Microbial lifestyle (e.g., symbiosis propensity and type) could have implications for how

well a microbe can resist or recover from climatic events. In the soil, microbes can live

symbiotically with (and generally derive a benefit from) host organisms like plants, animals, and

macroscopic fungi (Berendsen et al. 2012), with the outcome of symbiosis ranging from negative

to positive for the host. For example, a polyphyletic group of soil bacteria known as rhizobia can

facultatively live inside specialized legume root structures called nodules, where the rhizobia

provide biologically available nitrogen for the plant in return for carbon (Beringer et al. 1979).

Rhizobia genomes are large and contain symbiotic plasmids required for nodule formation

(MacLean et al. 2007), and this extra genetic material could hamper growth rates and select for

genome streamlining under warming (Hessen et al. 2010; Sabath et al. 2013; Simonsen 2021).

The bacterial plant pathogen *Agrobacterium tumefaciens* also carries large plasmids that contain

the tumor-inducing genes required for its lifestyle (White & Winans 2007), which could slow

replication times. In addition, plants have a ubiquitous, often mutualistic, symbiosis with

mycorrhizal fungi wherein mycorrhizae provide mineral nutrients to plants in return for carbon

and improve plant tolerance of water stress by extending the root system (Kakouridis et al.

2022). However, mycorrhizal fungi are obligately symbiotic (Smith & Read 1997), and thus the

fate of a mycorrhizal fungus under climate stress may depend on the performance of their plant

partner and vice versa, to positive or negative effect (Stachowicz 2001; Kiers et al. 2010). In

sum, there are clear pathways through which the life history and symbiotic nature of a microbe

could underlie facets of the microbe's resistance or resilience to climate events.

We used a factorial experiment to examine the response of soil microbial communities to

the interacting climate factors of heat and drought, with an emphasis on measuring resiliency and

legacy effects. We identify microbial groups and lifestyles particularly altered under climate

change, and discuss why they may be differentially affected.

117
Methods

Site description and sampling

We conducted this study at the Koffler Scientific Reserve (KSR, www.ksr.utoronto.ca) in

Ontario, Canada (latitude 44°01'48"N, longitude 79°32'01"W) in an old field environment

(Figure 1A). The average annual precipitation is 767 mm, with the average high temperature

reaching 27°C in July and the average low temperature reaching -11°C in January (Koffler

Scientific Reserve 2023).

The temperature-free-air-controlled enhancement experiment at KSR was established

following Kimball et al. (2008). Plots are 3m in diameter, with six plots having six 100W

infrared heaters each (Mor Electric Heating Association) and six plots having equivalent dummy

infrared heaters. In 2017, we excavated twelve plots 30 cm deep and filled them with a 1:1:1

mixture of local soil, peat moss, and sand; then, we planted white spruce seedlings (*Picea*

glauca, provided by Natural Resources Canada, Laurentian Forestry Centre, Quebec QC) in the

plots. A top layer of commercially supplied cedar bark mulch was added in 2020 and 2021, and

weeding occurred biweekly to keep plant cover constant. Starting in 2020, we applied climate

treatments to plots: control, heat, drought, or heat and drought (Figure 1B). The heaters were

activated in the heated plots to increase leaf temperature by 5°C day and night, while the drought

plots had concave-up (U) slotted rainout structures that covered ~50% of the plot surface area

and drained rain out and away from all plots. To control for the effects of the structures

themselves, non-drought plots had the same rainout structures except in the concave-down

position (∩), allowing all water to pass off the structure and into the plot. There were two

periods of active treatment (Table S1): rainout structures were present July-November 2020, and

June-October 2021, for a total of 8 months; heaters were active in heated plots August-December

2020, and August 2021-January 2022, for a total of 9 months. Heat and drought treatments were

set up in this way to study the physiological and molecular responses of different white spruce

genotypes to climatic stressors through the growing season and into early winter.

On June 11th 2021, September 14th 2021, and June 5th 2022, we sampled from the first

10 cm of topsoil from three locations ~50 cm apart in each plot. We collected the June 2021 and

2022 samples when there were no active climate treatments being applied (plots in a recovery

period) to test microbial resilience, and the September 2021 samples when climate treatments

were being actively imposed to test resistance.

Sequencing and QIIME2

We extracted soil microbial DNA using the QIAGEN DNeasy PowerSoil Pro Kit

and sequenced samples at Genome Quebec (Montréal, Canada) using Illumina MiSeq PE 250bp

16S rRNA amplicon sequencing on the conserved hypervariable V4 region (primer pair 515F-

806R) and ITS region (primer pair ITS1FP2-58A2RP3). Samples from 2021 and 2022 were

sequenced separately. We used Quantitative Insights Into Microbial Ecology 2 (QIIME2)

v.2022.2 (Bolyen et al. 2019) to trim sequences and then denoised sequences with DADA2

(Callahan et al. 2016). Using QIIME2, we removed ASVs that had fewer than 10 reads across all

samples, and assigned taxonomy using the 'sklearn' feature classifier (Pedregosa et al. 2011); we

used Greengenes 16S V4 region reference for bacteria (McDonald et al. 2012), and UNITE

(Nilsson et al. 2019) version 9.0 with dynamic clustering of global and 97% singletons for fungi.

We then filtered out reads assigned as cyanobacteria and mitochondria to remove plant and

animal DNA. Finally, for each of the bacterial and fungal datasets we constructed a phylogeny

using QIIME2's MAFFT (Katoh & Standley 2013) and FastTree (Price et al. 2010) functions to

obtain a rooted tree.

Statistical Analysis

We performed analyses in R v4.2.0 (R Core Team 2022), with the 'tidyverse' (Wickham

et al. 2019), 'phyloseq' (McMurdie & Holmes 2013), and 'microbiome' (Lahti & Shetty 2019)

packages. For statistical tests, we merged the subsamples of plots such that each plot had one set

of reads for each time point and domain, then rarefied samples to the minimum read number for

that sampling time point (Table S2). We analyzed the data from each time point separately unless

otherwise stated. All models followed the same general structure, with heat treatment, drought

treatment, and the heat $\times$ drought interaction as predictors. We tested for differences in Shannon

diversity using the rarefied datasets and linear models, then used type 3 ANOVAs from the 'car'

package (Fox & Weisburg 2019) on the linear models to calculate F values. We used weighted

Unifrac principal coordinates analysis (PCoA) on the rarefied relative abundance data. Next, we

tested for differences in rarefied relative community composition using adonis2 permutational

analysis of variance with the Bray-Curtis method and 9999 permutations ('vegan' package,

Oksanen et al. 2022); and results did not change based on the order of factors. To detect

differentially abundant taxa we used the analysis of compositions of microbiomes with bias

correction ('ANCOMBC') package (Lin & Peddada 2020). We performed the analysis at the

genus level with Holm's method p-value adjustment.

To assess treatment effects in a group of known symbiotic bacteria, we subsetted the

rarefied datasets to include only rhizobia genera (Lajudie et al. 2019) and tested for differences

in Shannon diversity and community composition as described above. To assess treatment

effects in symbiotic fungi, we first classified fungi into guilds using FUNGuild annotation

(Nguyen et al. 2016) and kept classifications only when 'Probable' or 'Highly Probable.' We

concatenated classifications across years and relativized each guild's raw read number to sample

read number, then used repeated measures ANOVAs with time included as a factor and plot as

the error term. The response variables were the relative abundance of guilds of interest: parasites,

pathogens, endophytes, and mycorrhizae; response variables were natural log transformed as

needed to improve the normality of error distributions.

<u>Results</u>

Fungal communities

Active treatment and recovery periods

During active treatment, fungal community composition was not resistant; it was strongly

shaped by drought ($F_{1,8}$=1.24, p<0.05; Appendix S1: Table S4), but not heat or heat × drought.

Shannon diversity was not significantly affected by any active experimental treatment (Appendix

S1: Table S3).

Fungal communities were not resilient, as they were slow to recover after experiencing

climate treatments. In the first recovery period, there was reduced Shannon diversity in the

treatments with drought and heat × drought legacies ($F_{1,8}$>7.19, p<0.05 for all effects; Figure 2A,

Appendix S1: Table S3). Then, in the second recovery period, the legacy of drought also

predicted fungal community composition ($F_{1,8}$=2.35, p<0.05; Figure 2B, Appendix S1: Table

S4).

Taxonomic and guild responses

Alongside whole community shifts, we identified differentially abundant genera due to

both heat and drought. The full list of differentially abundant fungal genera is shown in Figure 4.

Notably, Saccharomycetales (yeasts) showed strikingly higher relative abundance in drought

plots (Figure 3), driven by the *Babjevia* genus (~3x log fold increase in drought, Holm's adjusted

$p<0.05$; Figure 4). We also saw interactions of seasonality and climate treatments; for example,

Pezizales visually increased in relative abundance in our September samples (Figure 3), but

drought significantly dampened this increase compared to control plots, leading Pezizales to

have one of the largest negative log fold changes among fungi (Holm's adjusted $p<0.05$; Figure

4).

Our analysis of symbiotic fungal guilds showed the mycorrhizal guild significantly

decreased in relative abundance due to heat ($F_{1,8}=6.15$, $p<0.05$; Appendix S1: Table S5) and had a

non-significant trend of lower abundance in drought. Most of the mycorrhizal reads were

ectomycorrhizal, which is not surprising given that the spruce saplings in the plots are

ectomycorrhizal, and mycorrhizal relative abundance was much greater during September

samples compared to June samples ($F_{2,22}=13.2$, $p<0.001$; Appendix S1: Table S5). Our differential

abundance analysis found two ectomycorrhizal genera, *Wilcoxina* and *Suillus* (Dahlberg &

Finlay 1999; Yu et al. 2001), that associate with Pinaceae tree saplings both had decreased

relative abundance due to legacy of drought, but differing responses during active drought, with

Wilcoxina being decreased while *Suillus* was increased (Holm's adjusted $p<0.05$; Figure 4).

Additionally, our guild analysis showed the relative abundance of parasitic fungal guilds

(parasitic on other fungi, lichen, plants, or insects) was increased by drought ($F_{1,8}=13.0$, $p<0.01$;

Figure 5, Appendix S1: Table S5). Pathogens and endophytes were not significantly affected by

the climate treatments (Appendix S1: Table S5).

Bacterial communities

Active treatment and recovery periods

Bacteria communities were more resistant to the climate events than fungi, with no

significant shifts due to active treatments (Appendix S1: Table S6). Shannon diversity was also

not affected during active treatment (Appendix S1: Table S7).

Bacteria diversity and composition were also very resilient to climate treatments,

especially heat. The first recovery period suggested no significant legacy effects from treatment.

In the second recovery period, only the legacy of drought increased bacterial Shannon diversity

($F_{1,8}=9.30$, $p<0.05$; Figure 2C, Appendix S1: Table S7).

Taxonomic and guild responses

Despite whole bacterial communities having strong resistance to climate treatments, we

identified differentially abundant bacterial genera due to heat and drought during active

treatment and recovery periods (Holm's adjusted $p<0.05$; Figure 4), though fewer genera were

found to be significantly affected compared to fungi. Drought and legacy of drought not only

increased the abundances of some rhizobia (*Burkholderia*, *Cupriavidus*, *Methylobacterium*,

Rhizobium, Phyllobacterium), but also increased other plant growth-promoting rhizobacteria in

the soil, for example *Variovorax* spp., and *Pseudoxanthomonas* spp., which can help plants

tolerate abiotic stresses such as drought (Kang et al. 2014; Kaushal 2019; Singh et al. 2021). In

contrast, a potentially pathogenic genus *Agrobacterium* decreased in abundance due to drought,

and a human pathogen, *Legionella*, significantly increased due to the legacy of heat.

The rhizobia guild's Shannon diversity significantly increased due to active heat

($F_{1,8}=9.78$, p<0.05, Appendix S1: Table S8) and active drought ($F_{1,8}=5.29$, p=0.05, Appendix S1:

Table S8), and remained significantly increased due to legacy of drought in the second recovery

period ($F_{1,8}$=5.94, p<0.05, Appendix S1: Table S8, Figure 5). While rhizobia communities

visually clustered into treatment groups in PCoAs (Figure 5), only the first recovery period was

significantly predicted by the heat × drought interaction ($F_{1,8}$=1.78, p<0.05, Appendix S1: Table

S9); community composition was not significantly affected by heat and drought factors in the

second recovery period or the active treatment (Appendix S1: Table S9).

Discussion

Our results suggest that climate change, and especially drought, will have more persistent

effects on soil fungi than on soil bacteria. In fungal communities, drought shaped community

composition, decreased diversity, and altered the relative abundance of symbiotic fungi

important to plant communities. In contrast, bacterial communities were largely resistant to

climate treatments, although drought tended to increase diversity and increased the relative

abundance of some bacteria beneficial to plants. We discuss the implications of the differences

between these members of the microbial community, and the effects of climatic events on

symbiotic microbes below.

268
Fungi are less resistant and resilient

Microbial communities experienced legacy effects from the climate treatments, with the

effect of drought being stronger than the effect of heat. The study site regularly experiences

extreme annual swings in temperature, potentially conditioning microbes to resist and recover

faster from heat (Hawkes & Keitt 2015). In contrast to some studies (de Vries et al. 2018; Pec et

al. 2021), even relatively short-term climate treatments had strong immediate and lasting impacts

on fungal communities; droughted fungal communities did not resemble control communities

during or after treatment. Moreover, fungal community resilience was lower than bacterial

resilience to drought. Our prediction that fungi would be more resistant but less resilient to

climate change than bacteria was therefore only partially supported; their more complex life

history strategies and longer generation times may contribute to fungi's lack of resilience under

climate stress, but do not prevent immediate shifts in relative abundance or presence. In contrast

to our prediction, the bacterial community composition in our plots was very resistant to climate

treatments, despite literature predicting high turnover rates (Rousk & Bååth 2011). These results

are consistent with climate conditions affecting slow-growing microbes more, and fungal

communities having a higher proportion of very slow-growing microbes compared to bacteria,

thus showing greater change. While some literature suggests slow-growing microbes have lower

relative fitness under more extreme climate conditions (Sabath et al. 2013), slow-growing

microbes may also benefit from climate change relative to faster-growing microbes, especially if

heat and drought make resources scarce or reduce the cost of slower growth (Konstantinidis &

Tiedje 2004; Abreu et al. 2023).

The life histories of bacteria and fungi likely affected their resistance and resilience to

climatic shifts, although differences in dispersal could have also played a role. It is possible that

climate treatments may have influenced local bacterial populations, but dispersal was high

enough among bacterial metacommunities to mask these effects. If bacteria could disperse easily

between our plots, their community compositions would converge more readily and the strength

of signal would be smaller (Gianuca et al. 2017), while if fungi cannot disperse as easily, shifts

in composition would be more evident. Actual rates and distances of microbial dispersal are not

well described, but there is evidence that fungi are more dispersal-limited than bacteria (Zhang et

al. 2021) meaning fungi may be less able to escape a poor environment. While we cannot rule

out the role of dispersal, dispersal did not mask the significant effects of treatment on rhizobia

and other specific bacterial genera. While surprising, highly resistant bacterial communities are

not uncommon; for example, Waring & Hawkes (2018) also observed no direct compositional

changes to altered precipitation. Despite no significant changes in community composition, the

bacteria in our plots likely experienced physiological changes and costs (Schimel et al. 2007) as

well as turnover in allele frequencies and gene functions that we could not detect with 16S rRNA

sequencing, which may have then contributed to drought's legacy effects on bacterial diversity.

The community-wide shifts in composition and diversity we detected are likely a product

of both the direct effects of heat and drought on microbes and the indirect effects of climate

treatment on microbes mediated through species interactions, such that microbes not directly

affected by climate may still have increased or decreased in relative abundance. Microbes that

interact strongly with other microbes may have a linked response if any of their network is

disturbed by climate (Bardgett & Caruso 2020). Indirect effects of climate may also be mediated

through plant-microbe or animal-microbe interactions, where stressed organisms may alter the

soil environment in some way, for example plants may secrete different root exudates

(Karlowsky et al. 2018; Bakhshandeh et al. 2019), or, stressed organisms may themselves be an

altered environment for symbionts due to factors like growth or immune responses (Nejat &

Mantri 2017). While examining differential abundance trends over time, it became apparent that

microbes affected during active treatment were not necessarily the ones affected in recovery

periods (Figure 4). This observation demonstrates the complexity of legacy effects and

resiliency, and suggests a strong role of persisting direct and indirect effects of climate on

microbial communities.

Fungal communities showed low resistance and resilience, indicating that the ecosystem

services they provide may change under future climate conditions. Fungi are major decomposers

of organic material, and therefore unlock a large supply of nutrients for higher soil trophic levels

(Boddy & Watkinson 1995). Mycorrhizal fungi in particular contribute to nutrient cycling and

mineral weathering (Courty et al. 2010), as well as carbon sequestration (Clemmenson et al.

2013). Despite their importance to global elemental cycling and other ecosystem processes, most

research on fungal stress responses focuses solely on yeasts (Branco et al 2022). In this

experiment, yeasts were common in drought plots and did not represent the average fungal

response to treatments. Although yeasts thrived under drought, overall fungal diversity was

lower after drought and mycorrhizal abundance was reduced after heat, suggesting the breadth of

ecosystem services provided by fungal communities will be decreased under future climatic

conditions (Mori et al. 2016). Additionally, low resilience in soil communities means that plants

that recruit into a population following a climatic shift may interact with an altered pool of

microbes even if they do not experience the climate event themselves. Soil microbiomes affect

plant performance and fitness (Berendsen et al. 2012; Rubin et al. 2018; Lu et al. 2018;

Fitzpatrick et al. 2019) through mechanisms such as nutrient provisioning (Beringer et al. 1979)

and altering plant gene expression profiles (Lu et al. 2018). For plants, associations with

microbes can result in improved stress tolerance (Gehring et al. 2017; Allsup et al. 2023) or even

broadened geographic range size (Afkhami et al. 2014), which can be adaptive under climate

change. Therefore, soil microbes modulate how plants respond to climate events (Classen et al.

2015; Rudgers et al. 2020) and soil legacy effects may have far-reaching ecological and

environmental consequences.

343
Climate events on symbiotic microbes

Free-living and symbiotic microbes may respond differently to climate change, for

example if living inside host cells or tissue buffers microbes against changes to the abiotic

environment. Plant symbionts sometimes responded differently to climate treatments than the

average response of bacterial and fungal communities. We do not attribute these results to a

shifting plant community, because it was held constant throughout the experiment, however

results may be influenced by shifts in plant traits under different climate treatments.

The rhizobia guild showed a similar post-drought response compared to the overall

bacteria community with an increase in diversity; however, because we also saw significantly

higher diversity when heat and drought were being actively applied, this indicates rhizobia

positively responded more to active treatment than the overall community. Since rhizobia fix

biological nitrogen through legume symbiosis (Beringer et al. 1979), their high abundance bodes

well for global nitrogen cycles under climate change. This was unexpected, as we predicted

rhizobia would be a worse soil competitor under climate treatments due to their relatively large

genomes and thus longer replication times. However, if slower-growing microbes are more

sensitive to changes in climate, as we saw with our domain comparison, then this rhizobia result

is consistent. At higher temperatures, slow-growing marine bacteria cells retain more normal

membrane functions and have a lower mortality burden than fast growers (Schaum & Collins

2014; Abreu et al. 2023), which may apply to slow-growing soil bacteria too. Given our results,

we might predict the likelihood of symbiosis between specific plant-rhizobia partners to differ in

future climate conditions. A change in the availability of rhizobia partners matters because

rhizobia-plant interactions vary in specificity (Chen et al. 2021), which could lead to shifts in

plant communities and legume ranges (Harrison et al. 2018). The increased abundance and

diversity of rhizobia could also shape evolution in the rhizobia-legume system through increased

bacterial competition (Burghardt & diCenzo 2023).

While beneficial bacteria increased with our climate treatments, mutualistic fungi were

negatively affected. Mycorrhizae, mainly composed of ectomycorrhizae, had significantly lower

relative abundance in heated plots, suggesting mycorrhizae were less resistant to heat than the

fungal community overall (which showed no major responses to heat). We also observed

significant decreases in the abundance of ectomycorrhizal genera in the bulk soil due to active

drought and its legacy, and possibly as a result of this decreased abundance, drought has been

shown to reduce ectomycorrhizal root colonization (Erlandson et al. 2022). A recent study by Fu

et al. (2022) found that drought lowered the diversity of field arbuscular mycorrhizae, and they

suggested this shift was mediated by plant communities. In our experiment, because the

ectomycorrhizae obligately associated with the spruce saplings in our plots, it is likely there were

negative indirect climate effects on the mycorrhizae mediated through changes in the physiology

of the spruce, indirect microbe-microbe interactions, as well as the direct effects of climate

treatment. Ectomycorrhizal fungi have a narrow climatic tolerance and thus are predicted to be

disproportionately affected by climate change (Baldrian et al. 2023), which our results support.

Mutualistic and parasitic fungi both derive nutrients from hosts, however while heat

decreased mycorrhizae, drought increased parasites in recovery periods. If parasitic fungal

species become more prevalent in post-drought conditions this could increase the likelihood of

soil dwelling organisms hosting a parasite. Drought and heat may be especially problematic in

temperate forests, where most trees are ectomycorrhizal (Baldrian et al. 2023), since trees may

simultaneously lack their main mutualist partners and experience increased parasite pressure.

Organisms hosting parasites may have reduced growth and fitness, leading to reduced ecosystem

productivity. Additionally, parasitized hosts may experience more negative effects from climate

events than non-parasitized hosts (Baldrian et al. 2023). In positive feedback, pathogens and

parasitic microbes are predicted to increase in heated environments in part due to increased

susceptibility of stressed hosts (Pritchard 2011; Altizer et al. 2013; Baker et al. 2018). While we

saw limited evidence of heat differentially affecting parasites, our results did suggest drought

may also play a role in moderating parasite abundances under future climate conditions.

396
Conclusions

Soil microbiomes are tightly linked to global nutrient and chemical cycling, and there is a

rich literature describing how microbes buffer plants against the abiotic stresses of climate

change (e.g. Yang et al. 2009; Lau & Lennon 2012; Rudgers et al. 2020; Marro et al. 2022;

Allsup et al. 2023). Given the soil microbiome's important role in ecosystems, it is important to

understand how microbial populations respond to climate events and the factors underlying

variation in resistance and resilience. We found that fungi were less resistant and less resilient to

drought and its legacy compared to bacteria, while heat had limited overall effect on both

domains. Symbiosis generated variation in resistance and resilience within the domains, but did

not have a universal effect; for example notable mutualistic bacteria performed well under

climate stress while mutualistic fungi did not. Our results suggest that slower-growing microbes

will be more affected by climate change. Low resistance and resilience of soil fungal

communities will have far-reaching consequences for ecosystems under climate change.

Acknowledgments

We thank KSR staff throughout the years, and Chris Carlson for his help collecting and

transporting soil. This work was funded by the Government of Canada through Genome Canada

and Ontario Genomics (OGI-211), NSERC Discovery grants to IE, JRS, and MEF, and the

Canada Foundation for Innovation. We also acknowledge the support by the Forest and

Environmental Genomics group of the Canadian Forest Service Quebec who provided white

spruce seedlings and technical support during planting, and Genome Quebec for their sequencing

facilities.

<u>Conflict of Interest Statement</u>

The authors declare no conflicts of interest.

<u>Literature cited</u>

Abreu, C. I., M. Dal Bello, C. Bunse, J. Pinhassi, and J. Gore. 2023. Warmer temperatures
favor slower-growing bacteria in natural marine communities. Science Advances 9.

Afkhami, M. E., P. J. McIntyre, and S. Y. Strauss. 2014. Mutualist-mediated effects on
species' range limits across large geographic scales. Ecology Letters 17:1265–1273.

Allison, S. D., and J. B. H. Martiny. 2008. Resistance, resilience, and redundancy in
microbial communities. Proceedings of the National Academy of Sciences of the
United States of America 105:11512–11519.

Allison, S. D., and K. K. Treseder. 2008. Warming and drying suppress microbial activity
and carbon cycling in boreal forest soils. Global Change Biology 14:2898–2909.

Allsup, C. M., I. George, and R. A. Lankau. 2023. Shifting microbial communities can
enhance tree tolerance to changing climates. Science 380:835–840.

Altizer, S., R. S. Ostfeld, P. T. J. Johnson, S. Kutz, and C. D. Harvell. 2013. Climate
Change and Infectious Diseases: From Evidence to a Predictive Framework. Science
341:514–519.

American Society for Microbiology Report. 2022. Microbes and Climate Change – Science,
People & Impacts: Report on an American Academy of Microbiology Virtual
Colloquium held on Nov. 5, 2021. Washington (DC).

Andrews, J. 1992. Fungal Life-History Strategies. Pages 119–145 *in* J. Dighton, J. White,
and P. Oudemans, editors. The Fungal Community: Its Organization and Role in the
Ecosystem. Second edition. CRC Press.

Averill, C., B. G. Waring, and C. V. Hawkes. 2016. Historical precipitation predictably
alters the shape and magnitude of microbial functional response to soil moisture.
Global Change Biology 22:1957–1964.

Baker, D. M., C. J. Freeman, J. C. Y. Wong, M. L. Fogel, and N. Knowlton. 2018. Climate
change promotes parasitism in a coral symbiosis. The ISME Journal 12:921–930.

Bakhshandeh, S., P. E. Corneo, L. Yin, and F. A. Dijkstra. 2019. Drought and heat stress
reduce yield and alter carbon rhizodeposition of different wheat genotypes. Journal of
Agronomy and Crop Science 205:157–167.

Baldrian, P., R. López-Mondéjar, and P. Kohout. 2023. Forest microbiome and global
change. Nature Reviews Microbiology 21:487–501.

Bardgett, R. D., and T. Caruso. 2020. Soil microbial community responses to climate
extremes: resistance, resilience and transitions to alternative states. Philosophical
Transactions of the Royal Society B: Biological Sciences 375:20190112.

Bérard, A., T. Bouchet, G. Sévenier, A. L. Pablo, and R. Gros. 2011. Resilience of soil
microbial communities impacted by severe drought and high temperature in the
context of Mediterranean heat waves. European Journal of Soil Biology 47:333–342.

Berendsen, R. L., C. M. J. Pieterse, and P. A. H. M. Bakker. 2012. The rhizosphere
microbiome and plant health. Trends in Plant Science 17:478–486.

Beringer, J. E., N. Brewin, A. W. B. Johnston, H. M. Schulman, and D. A. Hopwood. 1979.
The Rhizobium-legume symbiosis. Proceedings of the Royal Society of London. Series
B. Biological Sciences 204:219–233.

Bissett, A., M. V. Brown, S. D. Siciliano, and P. H. Thrall. 2013. Microbial community
responses to anthropogenically induced environmental change: towards a systems
approach. Ecology Letters 16:128–139.

Boddy, L., and S. C. Watkinson. 1995. Wood decomposition, higher fungi, and their role in
nutrient redistribution. Canadian Journal of Botany 73:1377–1383.

Bolyen, E., J. R. Rideout, M. R. Dillon, N. A. Bokulich, C. C. Abnet, G. A. Al-Ghalith, H.
Alexander, E. J. Alm, M. Arumugam, F. Asnicar, Y. Bai, J. E. Bisanz, K. Bittinger, A.
Brejnrod, C. J. Brislawn, C. T. Brown, B. J. Callahan, A. M. Caraballo-Rodríguez, J.

Chase, E. K. Cope, R. Da Silva, C. Diener, P. C. Dorrestein, G. M. Douglas, D. M. Durall, C. Duvallet, C. F. Edwardson, M. Ernst, M. Estaki, J. Fouquier, J. M. Gauglitz, S. M. Gibbons, D. L. Gibson, A. Gonzalez, K. Gorlick, J. Guo, B. Hillmann, S. Holmes, H. Holste, C. Huttenhower, G. A. Huttley, S. Janssen, A. K. Jarmusch, L. Jiang, B. D. Kaehler, K. Bin Kang, C. R. Keefe, P. Keim, S. T. Kelley, D. Knights, I. Koester, T. Kosciolek, J. Kreps, M. G. I. Langille, J. Lee, R. Ley, Y. X. Liu, E. Loftfield, C. Lozupone, M. Maher, C. Marotz, B. D. Martin, D. McDonald, L. J. McIver, A. V. Melnik, J. L. Metcalf, S. C. Morgan, J. T. Morton, A. T. Naimey, J. A. Navas-Molina, L. F. Nothias, S. B. Orchanian, T. Pearson, S. L. Peoples, D. Petras, M. L. Preuss, E. Pruesse, L. B. Rasmussen, A. Rivers, M. S. Robeson, P. Rosenthal, N. Segata, M. Shaffer, A. Shiffer, R. Sinha, S. J. Song, J. R. Spear, A. D. Swafford, L. R. Thompson, P. J. Torres, P. Trinh, A. Tripathi, P. J. Turnbaugh, S. Ul-Hasan, J. J. J. van der Hooft, F. Vargas, Y. Vázquez-Baeza, E. Vogtmann, M. von Hippel, W. Walters, Y. Wan, M. Wang, J. Warren, K. C. Weber, C. H. D. Williamson, A. D. Willis, Z. Z. Xu, J. R. Zaneveld, Y. Zhang, Q. Zhu, R. Knight, and J. G. Caporaso. 2019. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nature Biotechnology 2019 37:8 37:852–857.

Burghardt, L. T., and G. C. diCenzo. 2023. The evolutionary ecology of rhizobia: multiple facets of competition before, during, and after symbiosis with legumes. Current Opinion in Microbiology 72:102281.

Callahan, B. J., P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, and S. P. Holmes. 2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nature Methods 13:581–583.

Castro, H. F., A. T. Classen, E. E. Austin, R. J. Norby, and C. W. Schadt. 2010. Soil microbial community responses to multiple experimental climate change drivers. Applied and Environmental Microbiology 76:999–1007.

Chen, W. F., E. T. Wang, Z. J. Ji, and J. J. Zhang. 2021. Recent development and new insight of diversification and symbiosis specificity of legume rhizobia: mechanism and application. Journal of Applied Microbiology 131:553–563.

Clemmensen, K. E., A. Bahr, O. Ovaskainen, A. Dahlberg, A. Ekblad, H. Wallander, J. Stenlid, R. D. Finlay, D. A. Wardle, and B. D. Lindahl. 2013. Roots and Associated Fungi Drive Long-Term Carbon Sequestration in Boreal Forest. Science 339:1615–1618.

Courty, P.-E., M. Buée, A. G. Diedhiou, P. Frey-Klett, F. Le Tacon, F. Rineau, M.-P. Turpault, S. Uroz, and J. Garbaye. 2010. The role of ectomycorrhizal communities in

forest ecosystem processes: New perspectives and emerging concepts. Soil Biology
and Biochemistry 42:679–698.

Dacal, M., P. García-Palacios, S. Asensio, J. Wang, B. K. Singh, and F. T. Maestre. 2022.
Climate change legacies contrastingly affect the resistance and resilience of soil
microbial communities and multifunctionality to extreme drought. Functional Ecology
36:908–920.

Dahlberg, A., and R. D. Finlay. 1999. Suillus. Pages 33–64 Ectomycorrhizal Fungi Key
Genera in Profile. Springer Berlin Heidelberg, Berlin, Heidelberg.

Dubey, A., M. A. Malla, F. Khan, K. Chowdhary, S. Yadav, A. Kumar, S. Sharma, P. K.
Khare, and M. L. Khan. 2019. Soil microbiome: a key player for conservation of soil
health under changing climate. Biodiversity and Conservation 28:8 28:2405–2429.

Erlandson, S. R., R. Margis, A. Ramirez, N. Nguyen, L. A. Lofgren, H. Liao, R. Vilgalys, P.
G. Kennedy, and K. G. Peay. 2022. Transcriptional acclimation and spatial
differentiation characterize drought response by the ectomycorrhizal fungus *Suillus
pungens*. New Phytologist 234:1910–1913.

Fitzpatrick, C. R., Z. Mustafa, and J. Viliunas. 2019. Soil microbes alter plant fitness under
competition and drought. Journal of Evolutionary Biology 32:438–450.

Fox J, Weisberg S. 2019. An R Companion to Applied Regression, Third edition. Sage,
Thousand Oaks CA.

Fu, W., B. Chen, M. C. Rillig, J. Jansa, W. Ma, C. Xu, W. Luo, H. Wu, Z. Hao, H. Wu, A.
Zhao, Q. Yu, and X. Han. 2022. Community response of arbuscular mycorrhizal fungi
to extreme drought in a cold-temperate grassland. New Phytologist 234:2003–2017.

Gehring, C. A., C. M. Sthultz, L. Flores-Rentería, A. V. Whipple, and T. G. Whitham.
2017. Tree genetics defines fungal partner communities that may confer drought
tolerance. Proceedings of the National Academy of Sciences 114:11169–11174.

Gianuca, A. T., S. A. J. Declerck, P. Lemmens, and L. De Meester. 2017. Effects of
dispersal and environmental heterogeneity on the replacement and nestedness
components of β-diversity. Ecology 98:525–533.

Harrison, T. L., A. K. Simonsen, J. R. Stinchcombe, and M. E. Frederickson. 2018. More
partners, more ranges: generalist legumes spread more easily around the globe.
Biology Letters 14.

Hartmann, M., P. A. Niklaus, S. Zimmermann, S. Schmutz, J. Kremer, K. Abarenkov, P.
Lüscher, F. Widmer, and B. Frey. 2013. Resistance and resilience of the forest soil
microbiome to logging-associated compaction. The ISME Journal 8:226–244.

Hawkes, C. V., and T. H. Keitt. 2015. Resilience vs. historical contingency in microbial
responses to environmental change. Ecology Letters 18:612–625.

Hessen, D. O., P. D. Jeyasingh, M. Neiman, and L. J. Weider. 2010. Genome streamlining
and the elemental costs of growth. Trends in Ecology & Evolution 25:75–80.

Jansson, J. K., and K. S. Hofmockel. 2020. Soil microbiomes and climate change. Nature
Reviews Microbiology 18:35–46.

Kakouridis, A., J. A. Hagen, M. P. Kan, S. Mambelli, L. J. Feldman, D. J. Herman, P. K.
Weber, J. Pett-Ridge, and M. K. Firestone. 2022. Routes to roots: direct evidence of
water transport by arbuscular mycorrhizal fungi to host plants. New Phytologist
236:210–221.

Kang, S. M., A. L. Khan, M. Waqas, Y. H. You, J. H. Kim, J. G. Kim, M. Hamayun, and I.
J. Lee. 2014. Plant growth-promoting rhizobacteria reduce adverse effects of salinity
and osmotic stress by regulating phytohormones and antioxidants in *Cucumis sativus*.
Journal of Plant Interactions 9:673–682.

Karlowsky, S., A. Augusti, J. Ingrisch, M. K. U. Akanda, M. Bahn, and G. Gleixner. 2018.
Drought-Induced Accumulation of Root Exudates Supports Post-drought Recovery of
Microbes in Mountain Grassland. Frontiers in Plant Science 9.

Katoh, K., and D. M. Standley. 2013. MAFFT Multiple Sequence Alignment Software
Version 7: Improvements in Performance and Usability. Molecular Biology and
Evolution 30:772–780.

Kaushal, M. 2019. Portraying Rhizobacterial Mechanisms in Drought Tolerance. Pages
195–216 PGPR Amelioration in Sustainable Agriculture. Elsevier.

Kiers, E.T., T. M. Palmer, A. R. Ives, J. F. Bruno, and J. L. Bronstein. 2010. Mutualisms in
a changing world: an evolutionary perspective. Ecology Letters 13:1459–1474.

Kimball, B. A., M. M. Conley, S. Wang, X. Lin, C. Luo, J. Morgan, and D. Smith. 2008.
Infrared heater arrays for warming ecosystem field plots. Global Change Biology
14:309–320.

Koffler Scientific Reserve. 2023. Koffler Scientific Reserve Quick Facts & Weather.

Konstantinidis, K. T., and J. M. Tiedje. 2004. Trends between gene content and genome
571   size in prokaryotic species with larger genomes. Proceedings of the National Academy
572   of Sciences 101:3160–3165.

573   Lahti, L., and S. Shetty. 2019. microbiome R package.

574   Lau, J. A., and J. T. Lennon. 2012. Rapid responses of soil microorganisms improve plant
575   fitness in novel environments. Proceedings of the National Academy of Sciences of the
576   United States of America 109:14058–14062.

577   de Lajudie, P. M., M. Andrews, J. Ardley, B. Eardly, E. Jumas-Bilak, N. Kuzmanović, F.
578   Lassalle, K. Lindström, R. Mhamdi, E. Martínez-Romero, L. Moulin, S. A. Mousavi,
579   X. Nesme, A. Peix, J. Puławska, E. Steenkamp, T. Stępkowski, C.-F. Tian, P. Vinuesa,
580   G. Wei, A. Willems, J. Zilli, and P. Young. 2019. Minimal standards for the
581   description of new genera and species of rhizobia and agrobacteria. International
582   Journal of Systematic and Evolutionary Microbiology 69:1852–1863.

583   Lin, H., and S. Das Peddada. 2020. Analysis of compositions of microbiomes with bias
584   correction. Nature Communications 11:1–11.

585   Lu, T., M. Ke, M. Lavoie, Y. Jin, X. Fan, Z. Zhang, Z. Fu, L. Sun, M. Gillings, J. Peñuelas,
586   H. Qian, and Y. G. Zhu. 2018. Rhizosphere microorganisms can influence the timing
587   of plant flowering. Microbiome 6:1–12.

588   MacLean, A. M., T. M. Finan, and M. J. Sadowsky. 2007. Genomes of the Symbiotic
589   Nitrogen-Fixing Bacteria of Legumes. Plant Physiology 144:615–622.

590   Marro, N., G. Grilli, F. Soteras, M. Caccia, S. Longo, N. Cofré, V. Borda, M. Burni, M.
591   Janoušková, and C. Urcelay. 2022. The effects of arbuscular mycorrhizal fungal
592   species and taxonomic groups on stressed and unstressed plants: a global meta-
593   analysis. New Phytologist 235:320–332.

594   Martiny, J. B., A. C. Martiny, C. Weihe, Y. Lu, R. Berlemont, E. L. Brodie, M. L. Goulden,
595   K. K. Treseder, and S. D. Allison. 2017. Microbial legacies alter decomposition in
596   response to simulated global change. The ISME Journal 11:490–499.

597   McDonald, D., M. N. Price, J. Goodrich, E. P. Nawrocki, T. Z. Desantis, A. Probst, G. L.
598   Andersen, R. Knight, and P. Hugenholtz. 2011. An improved Greengenes taxonomy
599   with explicit ranks for ecological and evolutionary analyses of bacteria and archaea.
600   The ISME Journal 6:610–618.

601   McMurdie, P. J., and S. Holmes. 2013. Phyloseq: An R Package for Reproducible
602   Interactive Analysis and Graphics of Microbiome Census Data. PLoS ONE 8.

Melillo, J. M., S. D. Frey, K. M. DeAngelis, W. J. Werner, M. J. Bernard, F. P. Bowles, G.
Pold, M. A. Knorr, and A. S. Grandy. 2017. Long-term pattern and magnitude of soil
carbon feedback to the climate system in a warming world. Science 358:101–105.

Mori, A. S., F. Isbell, S. Fujii, K. Makoto, S. Matsuoka, and T. Osono. 2016. Low
multifunctional redundancy of soil fungal diversity at multiple scales. Ecology Letters
19:249–259.

Nejat, N., and N. Mantri. 2017. Plant Immune System: Crosstalk Between Responses to
Biotic and Abiotic Stresses the Missing Link in Understanding Plant Defence. Current
Issues in Molecular Biology:1–16.

Nguyen, N. H., Z. Song, S. T. Bates, S. Branco, L. Tedersoo, J. Menke, J. S. Schilling, and
P. G. Kennedy. 2016. FUNGuild: An open annotation tool for parsing fungal
community datasets by ecological guild. Fungal Ecology 20:241–248.

de Nijs, E. A., L. C. Hicks, A. Leizeaga, A. Tietema, and J. Rousk. 2019. Soil microbial
moisture dependences and responses to drying–rewetting: The legacy of 18 years
drought. Global Change Biology 25:1005–1015.

Nilsson, R. H., K. H. Larsson, A. F. S. Taylor, J. Bengtsson-Palme, T. S. Jeppesen, D.
Schigel, P. Kennedy, K. Picard, F. O. Glöckner, L. Tedersoo, I. Saar, U. Kõljalg, and
K. Abarenkov. 2019. The UNITE database for molecular identification of fungi:
handling dark taxa and parallel taxonomic classifications. Nucleic Acids Research
47:D259–D264.

Ochman, H., J. G. Lawrence, and E. A. Grolsman. 2000. Lateral gene transfer and the
nature of bacterial innovation. Nature 405:299–304.

Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, O'Hara R, Solymos P,
Stevens M, Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D,
Carvalho G, Chirico M, De Caceres M, Durand S, Evangelista H, FitzJohn R, Friendly
M, Furneaux B, Hannigan G, Hill M, Lahti L, McGlinn D, Ouellette M, Ribeiro Cunha
E, Smith T, Stier A, Ter Braak C, Weedon J. 2022. vegan: Community Ecology
Package. R package version 2.6-2.

Pec, G. J., L. T. A. Van Diepen, M. Knorr, A. S. Grandy, J. M. Melillo, K. M. Deangelis, J.
L. Blanchard, S. D. Frey, C. : Pec, L. T. A. Van Diepen, M. Knorr, A. S. Grandy, J. M.
Melillo, K. M. Deangelis, and J. L. Blanchard. 2021. Fungal community response to
long-term soil warming with potential implications for soil carbon dynamics.

Pedregosa, F., G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P.
Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A. Passos, D. Cournapeau, M.

Brucher, M. Perrot, and E. Duchesnay. 2011. Scikit-learn: Machine Learning in Python . Journal of Machine Learning Research 12:2825–2830.

Price, M. N., P. S. Dehal, and A. P. Arkin. 2010. FastTree 2 – Approximately Maximum-Likelihood Trees for Large Alignments. PLOS ONE 5:e9490.

Pritchard, S. G. 2011. Soil organisms and global climate change. Plant Pathology 60:82–99.

R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Rooney, N., K. McCann, G. Gellner, and J. C. Moore. 2006. Structural asymmetry and the stability of diverse food webs. Nature 442:265–269.

Rousk, J., and E. Bååth. 2011. Growth of saprotrophic fungi and bacteria in soil. FEMS Microbiology Ecology 78:17–30.

Rousk, J., A. R. Smith, and D. L. Jones. 2013. Investigating the long-term legacy of drought and warming on the soil microbial community across five European shrubland ecosystems. Global Change Biology 19:3872–3884.

Rubin, R. L., G. W. Koch, A. Martinez, R. L. Mau, M. A. Bowker, and B. A. Hungate. 2018. Developing climate-smart restoration: Can plant microbiomes be hardened against heat waves? Ecological Applications 28:1594–1605.

Rudgers, J. A., M. E. Afkhami, L. Bell-Dereske, Y. A. Chung, K. M. Crawford, S. N. Kivlin, M. A. Mann, and M. A. Nuntildez. 2020. Climate Disruption of Plant-Microbe Interactions. Annual Review of Ecology, Evolution, and Systematics 51:561–586.

Sabath, N., E. Ferrada, A. Barve, and A. Wagner. 2013. Growth Temperature and Genome Size in Bacteria Are Negatively Correlated, Suggesting Genomic Streamlining During Thermal Adaptation. Genome Biology and Evolution 5:966–977.

Schaum, C. E., and S. Collins. 2014. Plasticity predicts evolution in a marine alga. Proceedings of the Royal Society B: Biological Sciences 281:20141486.

Schimel, J., T. C. Balser, and M. Wallenstein. 2007. Microbial Stress-Response Physiology And Its Implications For Ecosystem Function. Ecology 88:1386–1394.

Sheik, C. S., W. H. Beasley, M. S. Elshahed, X. Zhou, Y. Luo, and L. R. Krumholz. 2011. Effect of warming and drought on grassland microbial communities. ISME Journal 5:1692–1700.

Simonsen, A. K. 2021. Environmental stress leads to genome streamlining in a widely
distributed species of soil bacteria. The ISME Journal 16:423–434.

Singh, B. K., C. D. Campbell, S. J. Sorenson, and J. Zhou. 2009. Soil genomics. Nature
Reviews Microbiology 7:756–756.

Singh, S., V. Kumar, S. Datta, P. C. Ramamurthy, and J. Singh. 2021. Molecular
mechanism and signaling pathways interplay between plant hormones during plant-
microbe crosstalk. Microbial Management of Plant Stresses: Current Trends,
Application and Challenges:93–105.

Stachowicz, J. 2001. Mutualism, Facilitation, and the Structure of Ecological Communities:
Positive interactions play a critical, but underappreciated, role in ecological
communities by reducing physical or biotic stresses in existing habitats and by creating
new habitats on which many species depend. BioScience 51:235–246.

Smith, S.E., Read, D.J.1997. Mycorrhizal symbiosis. London: Academic Press.

T Classen, A. E., M. K. Sundqvist, J. A. Henning, G. S. Newman, J. A. M Moore, M. A.
Cregger, L. C. Moorhead, C. M. Patterson, M. K. Sundqvist, J. A. Henning, G. S.
Newman, J. A. M Moore, M. A. Cregger, L. C. Moorhead, and C. M. Patterson. 2015.
Direct and indirect effects of climate change on soil microbial and soil microbial-plant
interactions: What lies ahead? Ecosphere 6:1–21.

Thuiller, W. 2007. Climate change and the ecologist. Nature 448:550–552.

de Vries, F. T., R. I. Griffiths, M. Bailey, H. Craig, M. Girlanda, H. S. Gweon, S. Hallin, A.
Kaisermann, A. M. Keith, M. Kretzschmar, P. Lemanceau, E. Lumini, K. E. Mason, A.
Oliver, N. Ostle, J. I. Prosser, C. Thion, B. Thomson, and R. D. Bardgett. 2018. Soil
bacterial networks are less stable under drought than fungal networks. Nature
Communications 9:3033.

Waldrop, M. P., and M. K. Firestone. 2006. Response of Microbial Community
Composition and Function to Soil Climate Change. Microbial Ecology 52:716–724.

Waring, B., and C. V Hawkes. 2018. Ecological mechanisms underlying soil bacterial
responses to rainfall along a steep natural precipitation gradient. FEMS Microbiology
Ecology 94.

White, C. E., and S. C. Winans. 2007. Cell–cell communication in the plant pathogen
Agrobacterium tumefaciens. Philosophical Transactions of the Royal Society B:
Biological Sciences 362:1135–1148.

Wickham, H., M. Averick, J. Bryan, W. Chang, L. McGowan, R. François, G. Grolemund,
A. Hayes, L. Henry, J. Hester, M. Kuhn, T. Pedersen, E. Miller, S. Bache, K. Müller, J.
Ooms, D. Robinson, D. Seidel, V. Spinu, K. Takahashi, D. Vaughan, C. Wilke, K.
Woo, and H. Yutani. 2019. Welcome to the Tidyverse. Journal of Open Source
Software 4:1686.

Yang, J., J. W. Kloepper, and C.-M. Ryu. 2009. Rhizosphere bacteria help plants tolerate
abiotic stress. Trends in Plant Science 14:1–4.

Yu, T. E., K. N. Egger, and L. R. Peterson. 2001. Ectendomycorrhizal associations -
Characteristics and functions. Mycorrhiza 11:167–177.

Zhang, G., G. Wei, F. Wei, Z. Chen, M. He, S. Jiao, Y. Wang, L. Dong, and S. Chen. 2021.
Dispersal Limitation Plays Stronger Role in the Community Assembly of Fungi
Relative to Bacteria in Rhizosphere Across the Arable Area of Medicinal Plant.
Frontiers in Microbiology 12.

Zhou, Z., C. Wang, and Y. Luo. 2020. Meta-analysis of the impacts of global change factors
on soil microbial diversity and functionality. Nature Communications 11:3072.

Figure captions

Figure 1. Experimental design at Koffler Scientific Reserve. (a) The temperature-free-air-

controlled enhancement plots showing the infrared heaters and rainout structures, as well as the

water drainage system. Photo by Julia Boyle. (b) Schematic diagram of the experimental plots

(numbered) and their climate treatments. Dots represent sampled sites within plots.

Figure 2. Community structure of fungal and bacterial soil microbes across climate conditions

and time. Boxplots of Shannon diversity across time in fungal (a) and bacterial (c) microbial

communities. Weighted Unifrac PCoA plots of fungal (b) and bacterial (d) microbial

communities. When significant, main effects of heat, drought, and their interaction are included

with panels.

Figure 3. Community composition over time for each subsample in each treatment. (a)-(c) Fungi

relative abundances at the order level. (d)-(f) Bacteria relative abundance at the order level.

Orders were aggregated and included when they were at least 1% of the compositional

abundance, and found in at least 30% of samples. Plot order within treatment groups remains

consistent, with 2-3 samples per plot.

Figure 4. Differentially abundant microbe genera due to drought (a) and heat (b) across all

sampling times, compared to control plots. Taxa shown are significant ($p<0.05$) with Holm's

method p-value adjustment, grouped by domain, and ordered by descending average log fold

change.

Figure 5. Symbiotic microbes under climate treatments. (a) Rhizobia Shannon diversity. (b)

Rhizobia community weighted Unifrac PCoAs. (c) and (d) Fungal parasite and mycorrhizae

relative abundance across treatment and time. When significant, main effects of heat, drought,

and their interaction are included with panels.

743
744   **<u>Figures</u>**
745
746

747
748   Figure 1
749
750
751

752
753   Figure 2

754
755   Figure 3
756
757
758

759
760   Figure 4

761
762   Figure 5

(a)

(b)

12	11	10	9	8	7

1	2	3	4	5	6

Climate treatment

Control   **Drought**   **Heat**   **Heat and Drought**

⊙ = Sample site

Fungal Order

Acrospermales	Fungi_ord_Incertae_sedis	Pleosporales
Agaricales	Glomerellales	Pleurotheciales
Agaricostilbales	Helotiales	Polyporales
Auriculariales	Hypocreales	Rozellomycota_ord_Incertae_sedis
Boletales	Leucosporidiales	Saccharomycetales
Branch06	Microbotryales	Sakaguchiales
Cantharellales	Microbotryomycetes_ord_Incertae_sedis	Sebacinales
Capnodiales	Microascales	Sordariales
Chaetosphaeriales	Mortierellales	Sporidiobolales
Chaetothyriales	Myrmecridiales	Thelephorales
Coniochaetales	Onygenales	Trechisporales
Cystofilobasidiales	Orbiliales	Tremellales
Eurotiales	Other	Xylariales
Filobasidiales	Pezizales	

Bacterial Order

[Chthoniobacterales]	Armatimonadales	Clostridiales	iii1-15	Oceanospirillales	S0208
[Entotheonellales]	B07_WMSP1	Cytophagales	IS-44	OPB54	SBR1031
[Marinicellales]	B97	Desulfuromonadales	JG30-KF-CM45	Opitutales	SC-I-84
[Pedosphaerales]	Bacillales	DS-100	LD1-PA13	Other	Sediment-1
[Roseiflexales]	BD7-3	DS-18	Legionellales	Phycisphaerales	Solibacterales
[Saprospirales]	Bdellovibrionales	Ellin329	Methylacidiphilales	Pirellulales	Solirubrobacterales
0319-7L14	BPC015	Ellin5290	Methylophilales	PK29	Sphaerobacterales
11-24	Burkholderiales	Ellin6067	Micrococcales	PK329	Sphingobacterales
258ds10	C114	Ellin6513	MIZ46	Planctomycetales	Sphingomonadales
32-20	Caldilineales	Enterobacteriales	MND1	Pseudomonadales	Spirobacillales
A31	Caulobacterales	FAC88	MVS-107	RB41	Sva0725
A89	CCM11a	Flavobacteriales	MVS-40	Rhizobiales	Syntrophobacterales
Acidimicrobiales	CCU21	Gaiellales	Myxococcales	Rhodobacterales	Thiotrichales
Actinomycetales	CFB-26	GCA004	N1423WL	Rhodocyclales	Unknown
agg27	Chlamydiales	Gemmatales	NB1-j	Rhodospirillales	Verrucomicrobiales
AKIW781	Chloroflexales	Gemmatimonadales	Nitrososphaerales	Rickettsiales	wb1_H11
AKYG1722	Chromatiales	H39	Nitrospirales	Rubrobacterales	WCHB1-50
AKYG885	Chthonomonadales	Herpetosiphonales	Nostocales	S-70	WD2101
Alteromonadales	CL500-15	HTCC2188		S-BQ2-57	Xanthomonadales

(a) Rhizobia diversity

First recovery period | Active treatment | Second recovery period

Shannon diversity index

Active treatment: Drought p=0.05, Heat p<0.05
Second recovery period: Drought p<0.05

Treatment: Control, Drought, Heat, Heat and drought

(b) Rhizobia communities

First recovery period — Axis 1 [79.3%], Axis 2 [9.1%], Heat x Drought p<0.05
Active treatment — Axis 1 [72%], Axis 2 [16.2%]
Second recovery period — Axis 1 [84.9%], Axis 2 [11%]

Treatment: Control, Drought, Heat, Heat and drought

(c) Fungal parasites across treatment and recovery Drought p<0.01, Month p<0.001

First recovery period | Active treatment | Second recovery period

Parasite relative abundance

Treatment

(d) Mycorrhizae across treatment and recovery Heat p<0.05, Month p<0.05

First recovery period | Active treatment | Second recovery period

Mycorrhizal relative abundance

Treatment