1   **TITLE:** Allergic inflammation triggers dyslipidemia via IgG signalling

AUTHORS: Nieves Fernández-Gallego[1,2#], Raquel Castillo-González[1,3#], Lucía Moreno-Serna[2],

Antonio J. García-Cívico[4,5], Elisa Sánchez-Martínez[2], Celia López-Sanz[2], Ana Luiza Fontes[6],

Lígia L. Pimentel[6], Ana Gradillas[5], David Obeso[4,5], Marta Ramírez-Huesca[1], Ignacio Ruiz-

Fernández[1], Emilio Nuñez-Borque[7], Yolanda R. Carrasco[8], Borja Ibáñez[9,10,11], Pilar Martín[1,11],

Carlos Blanco[12], Coral Barbas[5], Domingo Barber[4], Luis M. Rodríguez-Alcalá[6], Alma Villaseñor[4,5],

Vanesa Esteban[7,13], Francisco Sánchez-Madrid[1,2,11], Rodrigo Jiménez-Saiz[2,8,14,15]*

AFFILIATIONS:

[1]Vascular Pathophysiology Area, Centro Nacional de Investigaciones Cardiovasculares (CNIC),

28029 Madrid, Spain

[2]Department of Immunology, Instituto de Investigación Sanitaria Hospital Universitario de La

Princesa (IIS-Princesa), Universidad Autónoma de Madrid (UAM), 28006 Madrid, Spain

[3]Department of Immunology, Ophthalmology and Ear, Nose and Throat (ENT), Universidad

Complutense, 28040 Madrid, Spain.

[4]Department of Basic Medical Sciences, Faculty of Medicine, Instituto de Medicina Molecular

Aplicada (IMMA), Universidad San Pablo-CEU, CEU Universities, 28668 Madrid, Spain

[5]Centro de Metabolómica y Bioanálisis (CEMBIO), Faculty of Pharmacy, Universidad San

Pablo-CEU, CEU Universities, 28668 Madrid, Spain

[6]Universidade Católica Portuguesa, CBQF - Centro de Biotecnologia e Química Fina –

Laboratório Associado, Escola Superior de Biotecnologia, 4169-005 Porto, Portugal

[7]Department of Allergy and Immunology, Instituto de Investigación Sanitaria Fundación Jiménez

Díaz (IIS-FJD), Universidad Autónoma de Madrid (UAM), 28040 Madrid, Spain

[8]Department of Immunology and Oncology, Centro Nacional de Biotecnología (CNB)-CSIC,

28049 Madrid, Spain

[9]Myocardial Pathophysiology Area, Centro Nacional de Investigaciones Cardiovasculares

(CNIC), Madrid, Spain

[10]Department of Cardiology, Instituto de Investigación Sanitaria Fundación Jiménez Díaz (IIS-

FJD), 28040 Madrid, Spain

[11]CIBER de Enfermedades Cardiovasculares (CIBERCV), Instituto de Salud Carlos III, 28029

Madrid, Spain

[12]Department of Allergy, Instituto de Investigación Sanitaria Hospital Universitario de La

Princesa (IIS-Princesa), 28006 Madrid, Spain

[13]Faculty of Medicine and Biomedicine, Universidad Alfonso X El Sabio, 28037 Madrid, Spain

[14]Department of Medicine, McMaster Immunology Research Centre (MIRC), Schroeder Allergy

and Immunology Research Institute (SAIRI), McMaster University, Hamilton, ON, L8N 3Z5,

Canada

[15]Faculty of Experimental Sciences, Universidad Francisco de Vitoria (UFV), 28223 Madrid,

Spain

#First co-authors.

CONTACT INFORMATON:

*Correspondence to: Rodrigo Jiménez-Saiz, PhD. Department of Immunology, Instituto de

Investigación Sanitaria Hospital Universitario de La Princesa (IIS-Princesa), Diego de León 62,

Madrid 28006, Spain.

Email address: rodrigo.jimenez@ufv.es / carlosrodrigo.jimenez.externo@salud.madrid.org

SUMMARY

Allergic diseases begin early in life and are often chronic, thus creating an inflammatory environment
that may lead to metabolic disorders, although the underlying mechanisms remain incompletely
understood. Here, we show that allergic inflammation induces diet-independent dyslipidemia in a
mouse model of allergy and atherosclerosis. Using untargeted lipidomics in mouse plasma, we found
that allergic inflammation induces a unique lipid signature that extends beyond acute and late
inflammation and that is characterized by triglyceride (TG) changes in circulation. Alterations in blood
TGs following an allergic reaction are independent of T-cell-driven late phase inflammation. On the
contrary, the humoral component is sufficient to induce a TG increase and a unique lipid profile
through the IgG-mediated alternative pathway of anaphylaxis. Lastly, we demonstrated blood TG
changes in patients after undergoing an allergic reaction. Overall, this study reveals the importance
of IgG-mediated allergic inflammation insofar as it regulates lipid metabolism, which may contribute
to atherosclerosis and, ultimately, to cardiovascular events.

INTRODUCTION

Type 2 immunity provides protection against parasites and non-microbial noxious substances and
regulates several homeostatic processes[1,2]. However, when it is directed toward innocuous antigens
from food, pollen, animal dander, *etc*. it causes allergic diseases[3]. These begin early in life and are
often chronic, thus creating an inflammatory ambient that may precede or exacerbate other
pathologies[4]. In this regard, allergy has been associated with a higher risk of atherosclerosis[5,6], which

accounts for over 80% of deaths caused by cardiovascular disease, the major cause of mortality
worldwide[7].

Atherosclerosis is a disease characterized by the chronic inflammation of blood vessels[8]. From an
immunological perspective, atherosclerosis is predominantly driven by type 1 immunity, which is
opposed to the hallmark type 2 identity of allergic disease[9]. Intriguingly, rather than counteracting
atherosclerosis, allergy seems to exacerbate this condition[10]. Atherosclerosis begins early in life,
typically associated with dyslipidemia, progresses slowly, and can remain silent until clinical
manifestations arise in the form of ischemic heart disease and/or stroke[11,12].

The pathology of allergic disease is heterogeneous and can generally be divided into two phases
that are initiated by allergen exposure in a time-dependent manner. Upon allergen exposure, acute
allergic reactions take place within minutes, or even seconds, via IgE (classical pathway)[13] or,
although less characterized in humans, via IgG (alternative pathway)[14,15]. Then, late phase
inflammation develops hours, or even days, after allergen encounter, and it is mainly orchestrated
by CD4 Th2 cells[16].

Recent studies in human allergic patients point towards metabolic alterations linked to allergic
disease[17-20]. Also, a recent transcriptomic study in the blood of healthy donors found an association
between triglyceride (TG) levels and canonical genes of type 2 immunity[21], which underscores the
existence of an interplay between allergic disease and lipid metabolism. Yet the relationship of
allergic disease and its different phases to metabolic changes and, particularly, to dyslipidemia, the
main risk factor of atherosclerosis, remains poorly understood.

Here, we investigated the impact of allergic inflammation in dyslipidemia, and the effect of the diet,
in a murine model of allergy and atherosclerosis. Furthermore, we ascertained the contribution of
acute and late phase inflammation to allergic dyslipidemia using cell-depleting strategies and passive
sensitization models and delved into the role of the classical and alternative anaphylaxis pathways.
Moreover, untargeted lipidomics were applied to define the lipid fingerprint of allergic inflammation.
Lastly, we evaluated changes in serum TG levels in patients undergoing allergic reactions. Overall,
this study reveals the importance of humoral allergic inflammation in lipid metabolism, which may
contribute to atherosclerosis and, ultimately, to cardiovascular events.

RESULTS

Allergic reactivity causes a diet-independent increase in blood triglycerides. To investigate the
clinical association between allergic disease and atherosclerosis, we applied a well-established
model of allergy and anaphylaxis[22,23] in LDL receptor-deficient mice (LDLr KO) (**Fig. 1a**). This mouse
strain is prone to atherosclerosis, particularly when fed a high cholesterol (HC) diet[24]. Following

sensitization, LDLr KO mice developed high serum levels of allergen-specific IgE and IgG1 (**Fig. 1b**),
and acute allergic responses upon allergen challenge, measured as changes in core temperature
(**Fig. 1c**). Also, allergic mice exhibited late phase inflammation, characterized by eosinophilia in the
peritoneal cavity (site of challenge) (**Fig. 1d; Fig. S1**). Given that the blood lipid profile can inform of
cardiovascular risk, we collected fasting blood samples at different time points of allergic pathology
including baseline, after sensitization, and post-allergen challenge (**Fig. 1a**).

107
Fig. 1 Allergic sensitization and clinical responses in atherosclerosis-prone mice. a Protocol
for allergen sensitization, challenge, and sample collection in LDLr KO mice fed a high cholesterol
(HC) or regular (RE) diet. **b** Serum levels of peanut (PN)-specific IgE and IgG1 after sensitization of
HC diet-fed naive (n=14) and allergic (n=15) mice. **c** Acute allergic response following allergen
challenge measured as hypothermia in HC diet-fed naive (n=16) and allergic mice (n=22). **d** Late
phase inflammation in the peritoneal lavage at 3 days post-allergen challenge quantified in terms of
total cell number and eosinophilia in HC diet-fed naive (n=15) or allergic (n=18) mice. Data for **b** and
d are presented as mean ± s.e.m. In **c** and **d,** each dot represents an individual mouse. Statistically
significant differences in **d** were calculated with the Mann-Whitney test (****p≤0.0001). Pool of 3
independent experiments.
118

We did not observe any significant differences in the lipid profile of allergic and naive mice except
for TGs, which were increased in the allergic group, but only after allergen challenge (**Fig. 2a**).

Fig. 2 Serum triglyceride (TG) levels increase following allergic inflammation. a, b Serum levels
of total cholesterol, free cholesterol, LDL:HDL ratio and TGs determined at baseline (Base), after
sensitization (Sens) and 3 days post-allergen challenge (PC) in naive and allergic mice fed a high-
cholesterol (HC) diet **(a)** or a regular (RE) diet **(b)**. The data shown in **a** include 12 naive and 12
allergic mice; the data shown in **b** include 15 naive and 17 allergic mice. Each dot represents an
individual mouse. Data are presented as mean ± s.e.m. Statistically significant differences between
groups for each time point were calculated with the Mann Whitney test (*p≤0.05; ***p≤0.0005). Pool
of 3 independent experiments.
129

The fact that elevated TG levels have been recently identified as a risk factor of subclinical
atherosclerosis in the PESA study (an observational, longitudinal and prospective study with 4,000
patients), even in patients with normal LDL:HDL[25], prompted us to investigate these changes further.
We questioned whether the systemic TG increase observed in allergic mice following allergen
exposure was partly due to the HC diet used to study atherosclerosis or was inherent to allergic
inflammation. To address this question, we followed a similar experimental design (**Fig. 1a**), but mice
were fed a regular diet (RE) rather than a HC one. Also, we collected blood samples after the first

challenge, to assess lipid changes in the incipient stages of atherosclerosis. In agreement with the previous data, we did not observe differences in the lipid profile of allergic and naive mice at different phases of allergic pathology, except for TGs, which were again increased in the allergic group after allergen challenge (**Fig. 2b**). Altogether, these results indicate that allergic inflammation induces a systemic increase in blood TGs that is diet independent.

Allergic inflammation causes a unique lipid signature in circulation. The systemic increase in serum TG levels following an allergic reaction was suggestive of more profound changes in the lipid profile. Thus, we performed a comprehensive, kinetic blood analysis at different times post-allergen (or PBS) exposure in allergic and naive mice under HC or RE diets, via untargeted lipidomics (**Fig. 3a**). The plasma samples, collected at 1-, 3-, and 7-days post-allergen or PBS challenge, were treated for lipid extraction and were analysed using liquid chromatography coupled to mass spectrometry (LC-MS). After curating the data with the respective quality assurance using the blanks, the presence in samples and quality controls (QCs), and according to the coefficient of variance of the QCs (<30%), we obtained 1,374 features including both positive and negative ionization modes. Then, we assessed for the major separations between groups based on the overall lipid profile via principal component analysis (PCA) models (**Fig. 3b**). As expected, the group on a RE diet differentiated well from those on a HC diet at all time points. Interestingly, the lipid profile of naive and PBS-challenged allergic mice on a HC diet was comparable and overlapping across the study, which indicates that quiescent allergy, or atopy *per se*, does not alter the blood lipid profile. Intriguingly, upon allergen challenge, the lipid profile of HC-diet-fed allergic mice became increasingly different, as compared to that of naive mice or unchallenged allergic mice on a HC diet. This effect became most obvious 7 days after inducing allergic inflammation, at which point the lipid profile of allergen-exposed mice on a HC diet was clearly distinct from that of the rest of the groups (**Fig. 3b**).

Fig. 3 Principal component analysis (PCA) models of the blood lipid profile at different phases of allergic pathology. a Protocol for allergen sensitization, challenge, and kinetic of blood sample collection in LDLr KO mice fed a high cholesterol (HC) or regular (RE) diet for lipidomics. **b** Unsupervised PCA model of liquid chromatography coupled to mass spectrometry (LC-MS) in negative and positive ionization mode of plasma samples from allergic or naive mice at different time points (1-, 3-, and 7-days post-allergen challenge, PC) of allergic pathology. The analysis includes allergen-challenged allergic mice on a RE diet (n=5, black color); non-challenged allergic mice on a HC diet (n=5, blue color); allergen-challenged naive mice on a HC diet (n=5, white color); and allergen-challenged allergic mice on a HC diet (n=6, yellow color). All data were unit variance (UV) scaled. The X and Y axis indicate the percentage of variability explained by each component. R^2 represents the total variability explained by the model to differentiate groups.

189
190
Fig. 4 Allergic inflammation induces a unique blood lipid profile of triglycerides (TGs). a
Heatmap of the 25 specific TGs differentially expressed (p-value<0.05) between allergic mice with

quiescent allergy (non-challenged, blue color, n=5) and active allergy (allergen-challenged, yellow
color, n=6), fed a high cholesterol diet, at 7 days post-allergen challenge. The figure shows the
clustering results in the form of a dendrogram. Relative lipid abundance is represented along a color
gradient from blue (diminished in active allergy) to red (increased in active allergy). **b** Analysis of the
lipid metabolic networks of differentially expressed TGs in allergic mice with active allergy according
to their chain length, desaturation, and level of significance with LINEX. The intensity of the color
indicates the extent of TG upregulation; the sphere size indicates the level of significance of the
change according to the -log10 value FDR; the association between TGs according to their chain
length and level of desaturation are indicated by a blue and an orange line, respectively. TG 52:2
(TG 16:0_18:1_18:1); TG 56:6 (TG 16:0_18:1_22:5); FDR, false discovery rate.
203

In view of the unique lipid fingerprint exhibited by allergic mice 7 days after following allergen

exposure, we decided to compare it to the lipid profile of mice with quiescent allergy (sensitized but

not exposed to allergen), to better understand the impact of allergic inflammation on the lipid profile.

We detected 392 features that were significantly different in this comparison (adjusted p-value, p-

false discovery rate (FDR)< 0.05) including those from positive and negative ionization mode. Thus,

from these, we were able to identify 171 lipid species **(Fig. S2; Table S1)**. Among them, we detected

25 specific TGs, all of which were increased during allergic inflammation **(Fig. 4a)**. These TGs were

further analyzed using lipid metabolic networks according to their chain length, desaturation, and

level of significance (**Fig. 4b**). The analysis showed a cluster of TGs that have been associated with

cardiovascular disease (TG 54:2; TG 52:2; TG 50:2; TG 48:2)[26,27]. In addition to these, but not

clustered together, TG 56:6, TG 56:4 and TG 53:3 have been also linked to cardiovascular disease[26]

or risk factors of it (*i.e.*, hypertension, diabetes)[28]. Collectively, these results show that allergic

inflammation yields a unique lipid signature, characterized by TG changes, that extends beyond the

acute phase of the pathology.

Blood changes in triglyceride levels are independent of T-cell driven allergic inflammation.
Allergic inflammation is characterized by an acute phase, which happens within minutes, or seconds,

following allergen exposure, and by a late phase, that occurs hours or even days after the reaction[16].

The fact that the unique blood lipid signature triggered by allergic inflammation was detected days

after allergen exposure made us postulate that late phase inflammation was driving it. CD4 Th2 cells

orchestrate late phase inflammation, hence we decided to deplete CD4 T cells prior allergen

challenge to block it (**Fig. 5a**). We confirmed by flow cytometry that the antibody regime followed

was able to deplete CD4 T cells (**Fig. 5b; Fig. S3**), as previously reported[22,29]. CD4 T-cell depletion

began 2 weeks after sensitization (1 week before allergen challenge) to avoid interfering with the

generation of allergen-specific IgE/IgG1, both of which contribute to the acute phase. Anti-CD4-

depleted and isotype-treated allergic mice showed comparable levels of allergen-specific IgE and

IgG1 (**Fig. 5c**). Likewise, both groups of allergic mice underwent anaphylaxis following allergen
challenge (**Fig. 5d**), thus confirming that CD4 T-cell depletion had neither interfered with sensitization
nor with acute allergic responses. The analysis of the blood lipid profile revealed that the TG peak
previously observed during allergic inflammation remained intact despite CD4 T-cell depletion (**Fig.**
5e). These data suggest that T-cell-driven late phase inflammation is not involved in the TG changes
triggered by allergic inflammation.

Fig. 5 Blood changes in triglyceride (TG) levels during allergic inflammation are independent
of CD4 T cells. a Protocol for allergen sensitization, CD4 T cell depletion, allergen challenge, and
sample collection in LDLr KO mice fed a high cholesterol diet. **b** Flow cytometry analysis of CD4 T
cell percentage in the spleen of anti-CD4-depleted and isotype-treated allergic mice. **c** Serum levels
of peanut (PN)-specific IgE and IgG1 after sensitization. **d** Acute allergic responses following
allergen challenge. **e** Serum TG levels determined at baseline (Base), after sensitization (Sens) and
1- and 3- days post-allergen challenge (PC) in naive (n=8), and anti-CD4-depleted- (n=8) and

isotype-treated (n=7) allergic mice. Data for **b, c** and **e** are presented as mean ± s.e.m. In **b** and **d,**
each dot represents an individual mouse. Statistically significant differences in **e** were calculated
with the Kruskal-Wallis test (****p≤0.005**). Pool of 2 independent experiments.
247

Blood changes in TG levels during allergic inflammation are driven by IgG signalling. The
seemingly dispensable contribution of late phase allergic inflammation to dyslipidemia raised the
possibility that this was triggered by the acute allergic response, which is strictly dependent on IgE
and IgG1. To test this hypothesis, we devised a model of passive sensitization, which is based on
the transfer of sera from allergic mice into naive recipients[22]. With this approach, only the humoral
allergic machinery is transferred to naive mice, thus allowing for intact acute allergic responses and
precluding Th2-driven late phase inflammation (**Fig. 6a**). Given that acute allergic responses can
operate through the classical (IgE-mediated) and/or the alternative pathway (IgG-mediated), we
examined the contribution of each pathway to blood TG levels.

Unlike other antibodies, IgE is irreversibly denatured at 57°C due to a heat-labile domain, which is
not present in other antibody isotypes[30,31]. We exploited this biochemical property to generate allergic
sera without functional IgE. Following the heat-treatment of allergic sera, we observed a decrease
in allergen-specific IgE levels as compared to the native allergic sera, while IgG1 levels remain
unchanged (**Fig. S4a**). To ascertain if the remnants of denatured IgE were functional, we used a cell
activation assay based on bone marrow-derived mast cells[32]. Mast cells sensitized with native
allergic sera showed a dose-dependent activation following allergen stimulation by means of CD63
expression detected by flow cytometry (**Fig. 6b; Fig S4b**). On the other hand, mast cells sensitized
with heat-denatured allergic or naive sera did not increase CD63 expression after allergen
stimulation, which was comparable to that of unstimulated cells (**Fig. 6b; Fig S4b**). These data
demonstrate that the heat-treatment rendered allergen-specific IgE completely ineffective.

We also leveraged on the ability of anaphylactic IgE to remain bound, for months, to mast cells via
its high affinity receptor (FcεRI), which contrasts with the rapid clearance of circulatory IgG from the
circulation in a matter of days[33]. Thus, resting the mice for 2 weeks after passive sensitization with
allergic serum allows for blood IgG clearance and restricts anaphylaxis to the classical pathway[22].
Therefore, adjusting the sensitization time prior challenge as well as the type of sera used for it
(native and heat-denatured), allowed us to induce anaphylaxis via the classical and/or alternative
pathway (**Fig. 6a**). Upon challenge, all the allergic mice developed anaphylaxis to a similar level as
determined by hypothermia (**Fig. 6c**), regardless of the sensitization sera used (native or heat-
treated) and of the resting period prior challenge (1 day or 2 weeks). In line with the previous
experiments, we did not observe changes between groups in the lipid profile following allergen
challenge except for TGs (**Fig. 6d, Fig. S4c**). Interestingly, the group of allergic mice that underwent

anaphylaxis through the classical pathway (IgE-mediated) did not exhibit a TG peak. However, the
groups of mice that experienced anaphylaxis through the alternative pathway (IgG) or a combination
of both, showed a significant increase in blood TGs (**Fig. 6d**). Furthermore, we determined the lipid
profile of plasma from these mice at 7-days post-allergen challenge via LC-MS analysis, as
abovementioned. We identified 1,198 features that were grouped into 36 lipid classes for heatmap
analysis (**Fig. 6e**). Remarkably, the naive and IgE groups were clustered together, while the IgE+IgG
and IgG groups exhibited a profile that was more alike. The IgG group had a distinct lipid profile
characterized by high peak intensities in TGs, as well as in compounds belonging to lipid classes
such as sphingosine, N-acyl ethanolamines, lysiphosphatidylethanolamines, hexosylceramides,
monogalactosyldiacylglycerols, lysophosphatidylcholine, sulfonolipids, sphingomyelins,
phosphatidylcholine, ceramides and diglycerides. In line with this, the univariate analysis showed
that 456 lipids changed significantly (p-value<0.05) when comparing the IgG *vs* control group, while
210 lipids did it in the IgE *vs* control group comparison (**Table S2**). Of note, 645 lipids were
differentially expressed when comparing the IgG *vs* IgE group. These differences were attenuated
when comparing the IgE+IgG group *vs* the IgE (239 lipids) or IgG (325 lipids) groups, due to their
complementary profiles (**Table S2**). Altogether, these data indicate that allergic reactions are
sufficient to induce a blood TG increase and a unique lipid signature that is mostly triggered by the
IgG-mediated alternative pathway of anaphylaxis.
297

Fig. 6 Blood changes in triglyceride (TG) levels and in the lipid profile are driven by the IgG-mediated alternative pathway of anaphylaxis. a Protocol for passive sensitization with native and heated sera to induce anaphylaxis via de classical and/or the alternative pathway. **b** Activation assay with mast cells sensitized with naive sera, native allergic sera, and heat-treated allergic sera (representative dot plots of 4 independent experiments). **c** Acute allergic responses following allergen challenge of mice passively sensitized with naive sera (n=17) or allergic sera (native or

heated) to induce anaphylaxis via the classical (IgE; n=19) or alternative pathway (IgG; n=19), or
both (IgE+IgG; n=18). **d** Serum TG levels measured at 1- and 7-days post-allergen challenge (PC)
in passively sensitized mice. **e** Heatmap of the lipid species differentially expressed (p-value<0.05)
at 7 days PC in different groups of passively sensitized mice (a pool of 5 individual samples from
each group was analyzed in triplicate). The figure shows the clustering results in the form of a
dendrogram. Relative lipid abundance is represented along a color gradient from blue (diminished)
to red (increased). Data for **d** are presented as mean ± s.e.m. In **c** and **d** and each dot represents
an individual mouse from a pool of 3 independent experiments. Statistically significant differences in
d were calculated with a two-way-ANOVA followed by Bonferroni's multiple comparisons test
(*p≤0.05; ***p≤0.0005; ****p<0.0001). AHexCer, acylhexosylceramides; ASG, acylsterylgycoside;
BMP, bis(monoacylglycero)phosphates; CAR, acyl carnitines; CE, cholesteryl esters; Cer,
ceramides; DG, diacylglycerides; DGDG, digalactosyldiacylglycerol; Hex2Cer, dihexosylceramides;
Hex3Cer, trihexosylceramides; HexCer, hexosylceramides; LPC, lysophosphatidylcholines; LPE,
lysophosphatidylethanolamines; MG, monoacylglycerides; MGDG, mongalactosyldiacylglycerols;
NAE, N-acyl ethanolamines; NAGly, N-acyl glycines; NAGlySer, N-acyl glycyl serines; NAOrn, N-
acyl ornithines; NAPhe, N-acyl phenylalanines; NATau, N-acyl taurines; PC, phosphatidylcholines;
PE, phosphatidylethanolamines; PI, phosphatidylinositols; PS, phosphatidylserines; SE, steryl
esters; SG, sterylglycoside; SHexCer, sulfatides; SL, sulfonolipids; SM, sphingomyelins; SPB,
sphingoid bases; ST, sterols; VAE, vitamin A fatty acid esters.
324
325
Allergic patients show increased blood triglyceride levels after an allergic reaction.
Considering the murine data, we decided to explore whether allergic reactions were able to induce

blood TG changes in humans. Many common allergens are environmental and ubiquitous[34,35], which

makes it difficult to control allergen exposure. In addition, subclinical exposures to the allergen might

have an immunological impact too complex to be evaluated and controlled. Notwithstanding, we

collected blood samples from 59 patients during the onset of an allergic reaction, and another one

≥14 days later (**Fig. 7a**) to measure if allergic inflammation had induced changes in blood TG levels.

Most of the patients presented hypersensitivity reactions of grade I and II severity[36], triggered by

drugs or foods (**Table 1**). Given that some anti-inflammatory drugs may alter the blood TG levels[37],

the initial sample was taken prior treatment of the reaction, and the second sample was taken at

least 14 days later to allow for drug clearance. We also evaluated in different preclinical models of

anaphylaxis that blood TG levels did not change shortly after the allergic reaction (data not shown)

and remain unchanged for up to 24 h after it (**Fig. 4e**).

346 **Table 1. Allergic reactions**

Allergic patients	
Number	59
Age (years, SD)	38.7 ± 19.2
Sex (♀)	61%
Trigger of the reaction	
Drugs	69%
Foods	29%
Others	2%
Severity	
Level 1	49%
Level 2	41%
Level 3	10%
Symptoms	
Skin	92%
Mucosal	51%
Airways	36%
Gastrointestinal	20%
Neurologic	12%
Cardiovascular	10%

Of note, the serum TG levels of allergic patients increased significantly following an allergic reaction
(**Fig. 7b**). These data support our previous findings in mice and confirm that allergic reactions alter
blood TG levels in humans as well.

Fig. 7 Allergic patients show increased blood triglyceride (TG) levels days after undergoing
an allergic reaction. a Protocol followed for serum collection from 59 allergic patients. **b** Serum TG
levels measured at the onset of the allergic reaction and ≥14 days later. Data presented as mean ±
s.e.m. Each dot represents an individual patient. Statistically significant differences were calculated
with the Wilcoxon rank-sum test.
357
358

DISCUSSION

Recent studies in humans indicate that allergic disease may cause metabolic changes. For example, it has been reported that children with food allergy have lower serum levels of sphingolipids and ceramides[17], and severe respiratory allergy has been associated with higher plasma levels of fatty acids[19]. Along the same line, the metabolomic profiling of serum from food allergic children, at the time of diagnosis and allergy resolution, revealed that the lipid profile is closely related to allergy outgrowth[18]. Also, it has been reported that patients with allergic rhinitis, asthma and atopic dermatitis have increased risk of dyslipidemia[38-40]. While these studies evince the existence of an interplay between allergic disease and lipid metabolism, the mechanisms involved are poorly understood, particularly the contribution of the different phases of allergic inflammation to dyslipidemia.

Using a murine model of allergy and atherosclerosis, we observed that allergic mice exhibited dyslipidemia, which was characterized by increased levels of serum TGs. Intriguingly, this serum TG increase detected in allergic animals was not inherent to allergic disease, but specific to allergic inflammation. Hence, quiescent allergy or atopy in and of itself did not alter serum TG levels, but an allergic reaction was required for this to happen. Also, these changes in serum TGs appear to occur regardless of the diet because the blood was collected under fasting conditions in mice fed atherosclerosis-prone diets and regular diets. Further studies should determine whether the lipids released during allergic inflammation, and particularly TGs, are due to *de novo* synthesis in the liver, local release from adipocytes, and/or to changes in the lipid metabolism of immune cells.

Several studies in humans have established a long-lasting association between elevated serum TG levels and cardiovascular disease[41-44], although it is still unclear whether TGs promote disease via direct or indirect mechanisms. A recent study combined large-scale human genetic analysis with experimental evidence. It showed that TG-lowering alleles involved in hepatic production of TG-rich lipoproteins (*TM6SF2* and *PNPLA3*) and peripheral lipolysis (*LPL* and *ANGPTL4*) were associated with lower risk of cardiovascular disease[45]. Also, elevated serum TG levels have been recently identified as a risk factor of subclinical atherosclerosis in the PESA study, even when the LDL:HDL ratio was normal[25].

A number of studies both in rodents[46,47] and humans[48,49] have explored metabolic changes following an allergic reaction, but focusing on the acute phase, to unveil biomarkers or mediators of anaphylaxis. In other words, these studies analyzed metabolic changes in circulation minutes or hours after the allergic reaction was triggered. Our comprehensive, untargeted lipidomic analysis in plasma was performed days (1, 3, and 7) after allergen challenge, well beyond the acute phase reaction was over. Strikingly, these analyses revealed a unique lipid fingerprint in allergic mice following an allergic reaction, particularly at 7 days post allergen exposure, as compared to

unchallenged allergic mice. Their profile was characterized by an increase in 25 specific TGs, some of which have been positively associated with risk of cardiovascular disease. For example, the lipidomic analysis of 685 plasma samples of the prospective population-based Bruneck study found that TG 56:6, TG 52:2, TG 50:2, and particularly TG 54:2, were significantly associated with cardiovascular disease risk[26]. Furthermore, using 5,662 serum lipid profiles from PROMIS (a case-control study of first-ever acute myocardial infarction), Harshfield *et al*. identified lipids associated with the genetic polymorphism rs662799 in the *APOA5−APOC3* locus, which is linked to coronary artery disease. Of these, TG 53:3 and TG 56:4 were positively associated with risk factors of cardiovascular disease including hypertension and diabetes, respectively[28]. Also, TG 38:0 has been identified as a serum biomarker of Wilson´s disease, in a cohort of 34 patients that included 34 relatives of these patients and 64 healthy individuals as controls. Wilson´s disease is a genetic pathology characterized by copper accumulation and dysregulated lipid metabolism[50]. Although the relationship between Wilson´s disease and atherosclerosis has not been defined, it has been reported that Wilson´s disease patients are at a higher risk of suffering cardiovascular events[51]. On the other hand, Fernandez *et al*. performed a plasma lipidome using 427 samples from the prospective population-based Malmö Diet and Cancer (MDC) study. They determined that TG 48:2 was negatively associated with cardiovascular disease[27]. While further validation of these TG species is required in well-characterized cohorts of allergic patients, it is noteworthy that a substantial number of the TG species identified in the murine model have been associated with risk of cardiovascular disease in humans.

To better understand the interplay between allergic disease and lipid metabolism, we took different approaches to ascertain the contribution of cellular and humoral inflammation to serum TG changes. Given that the TG peak in serum was observed between 3- and 7-days post-allergen exposure, and the lipid profile was unique 7 days after the allergic reaction, we considered that these changes were associated with late phase inflammation, which peaks 3 days after challenge in this model[52]. Although different players such as type 2 innate lymphoid cells can contribute to late phase inflammation[53], CD4 Th2 cells orchestrate it[16]. Surprisingly, the impairment of CD4-T-cell-driven late-phase inflammation did not affect the increase in serum TG changes, thus pointing towards the acute phase response. Indeed, using models of passive sensitization, where there is no type 2 adaptive immune response established, but exclusively the humoral components transferred, we demonstrated that the acute phase alone was sufficient to recapitulate the increase in serum TGs. Furthermore, we delved into the role of the classical (IgE-mediated) and alternative (IgG-mediated) pathways of anaphylaxis to dyslipidemia. Strikingly, the classical pathway of anaphylaxis appeared to be dispensable for the changes in serum TGs, as these were only observed when the alternative

pathway or both were active. Also, the alternative pathway was sufficient to induce a unique lipid
signature, clearly different from the one produced by the classical pathway, which was similar to that
of the control group. The identification of lipid species that are specifically upregulated days after the
onset of an IgG-mediated reaction may be helpful to appraise its actual contribution to human
anaphylaxis, which still lacks specific biomarkers[14].

From a broader perspective, these data display a novel role for humoral responses, in particular
those of the IgG isotype insofar as they can regulate the lipid metabolism in response to allergens.
However, it is plausible to consider that this mechanism may operate in other diseases, not
necessarily type 2 driven, where strong IgG responses are chronically induced[54]. For example,
autoimmune diseases, such systemic lupus erythematosus and rheumatoid arthritis, are associated
with dyslipidemia and high serum TG levels, even in patients naive to corticosteroids[55-57].

Assessing the precise contribution of acute *vs* late allergic inflammation to serum TG levels in
humans is almost an unsurmountable challenge. Even so, we attained to show in a group of 59
patients that an allergic reaction induced an increase in serum TG levels days after the reaction. A
recent transcriptomic study in blood of 3,229 individuals from the BIOS (Biobank-based Integrative
Omics Study) consortium found an association between TG levels and canonical genes of type 2
immunity[21]. Although information pertaining to allergic disease of these patients was not available,
they found that TG levels were associated with a downregulation of the hallmark Th2 cytokine, *IL4,*
two IgE-receptor genes (*FCER1A* and *MS4A2*) as well as genes related to allergic mediators or its
metabolism (*HDC, HRH4, CPA3, HPGDS, CYP11A1, PTGER3*)[21]. In this context, one may speculate
that the increase of TGs observed following an allergic reaction is the output of a regulatory
mechanism that prevents immunopathology by reducing the expression of key molecules of the
allergic machinery. Yet, the chronic use of this putative mechanism of control may be at the cost of
a higher risk of developing cardiovascular events.

EXPERIMENTAL PROCEDURES

Mice. Age-, sex-, and strain-matched controls were used in all the experiments. B6;129S7-
Ldlrtm1Her/J mice were crossed with B6.SJL-Ptprc[a]Pepc[b]/BoyCrl mice and selected by genotype
(LDLr KO) at CNIC´s Animal facility. Mice were maintained in biohazard specific pathogen-free
conditions on a 12-hour light-dark cycle with a chow diet (RE; LASQCdiet Rod18-A, Altromin) or a
HC diet (SSNIFF, S9167-E010) and water *ad libitum*. All procedures were approved by the
Environmental Council of the Community of Madrid (Madrid, Spain) with PROEX references 45.2/20
and 9.7/23.

Allergy model. Male mice were sensitized with 4 mg of PN butter (~1 mg of protein; Capitán Mani) and 5 µg of cholera toxin (C8052, Sigma) in 0.5 mL of PBS administered intragastrically, weekly for 4 weeks. Following sensitization, mice were challenged intraperitoneally with 2.5 mg of crude peanut extract (CPE; XPF171D3A25; Stallergens Greer), or 88 µg of CPE per gram of body weight, and core temperature was assessed rectally with a dual thermometer (620-2049, VWR). Serum or plasma samples were collected at different times of allergic pathology under 12-16 hours of fasting and stored at -80°C for further analyses. In some instances, at 3- or 7-days post-allergen challenge, peritoneal lavage was performed with ice-cold 10 mM EDTA in PBS and spleens were processed to assess eosinophilia and CD4 T-cell depletion by flow cytometry[23].

Antigen-specific Ig ELISA. ELISAs were performed in a 96-well flat bottom polystyrene plate (3590, Corning). Plates were washed with 0.05% Tween-20 in PBS (washing buffer), and 10% FBS/1% BSA in washing buffer was used as blocking solution. Plates were developed with 50 µl TMB (421101, BioLegend) and stopped using 50 µl 1.5 N H_2SO_4. Plates were read at 450 nm using a Thermo Scientific Multiskan FC. No-sample controls were included in all assays. The average optical density of the no-sample control wells was subtracted from all samples prior to plotting.

To determine the <u>serum levels of PN-specific IgE</u>, plates were coated with 50 µl of 2 µg/ml anti-mouse IgE (clone R35-72M; 553413, BD) in PBS, sealed with aluminum foil, and incubated overnight at 4°C. The next day, plates were washed and blocked for 2 hours at room temperature (RT). Plates were washed and incubated with 50 µl of serum diluted to the indicated concentrations in blocking solution overnight at 4°C. A standard curve was included of purified mouse IgE k Isotype control (557079, BD Biosciences) from 120 ng/ml to 0.9 ng/ml diluted in PBS. The next day, plates were washed and incubated with 50 µl of biotinylated-CPE at 150 ng/ml while standards were incubated with 50 µl of blocking solution, for 90 minutes at RT. Plates were washed and the samples were incubated with 50 µl of streptavidin-HRP diluted in blocking solution and standards with 50 µl of 0.175 µg/ml rat anti-mouse IgE-HRP (1130-05, Southern Biotech) diluted in blocking solution for 1 hour at RT. Lastly, plates were washed, developed, stopped, and read.

To determine the <u>serum levels of PN-specific IgG1</u>, plates were coated with 10 µg/ml CPE in 50 µl of carbonate bicarbonate buffer, sealed with aluminum foil, and incubated overnight at 4°C. The next day, plates were washed and blocked for 2 hours at RT. Plates were washed and incubated with 50 µl of serum diluted to the indicated concentrations in blocking solution overnight at 4°C. The next day, plates were washed and incubated with 50 µl of 0.5 µg/ml goat anti-mouse IgG1-HRP (A10551, Thermo Fisher Scientific) in blocking solution for 2 hours at RT. Ultimately, plates were washed, developed, stopped, and read.

493 Antibody treatments and passive sensitization. CD4 T-cell depletion was accomplished via 2
intraperitoneal (i.p.) administrations of 200 µg of anti-CD4 (clone GK1.5; BE0003, BioXCell) one
week and one day before challenge[22]. IgG from rat serum (I4131, Sigma-Aldrich) was used as
isotype control. Cell depletion was confirmed by flow cytometry.

For passive sensitization, a pool of sera collected from PN-allergic or non-allergic mice under fasting
conditions was administered to male naive LDLr KO mice via i.p. injection. For IgG-passive
sensitization, part of the PN-allergic sera pool was heated at 57°C for 4 hours to accomplish IgE
denaturation[30,31]. Then, 700 ul of heat-treated sera was transferred to naive recipients the previous
day to challenge. The irreversible denaturation and deactivation of IgE was confirmed by ELISA and
mast cell activation test[32]. In the rest of the groups (naive, IgE, and IgE+IgG), the naive or native
allergic sera, respectively, were serially transferred as follows: 100 µL; 200 µL; and 400 µL with an
interval of 48 hours between each administration. Then, the mice were challenged either the day
after the last administration, to induce IgE- and IgG-mediated anaphylaxis, or 10 days later to allow
for IgG clearance and IgE-mediated anaphylaxis[58].

Flow cytometry. Staining was conducted in FACS buffer (2.5 mM EDTA, 0.5% BSA, in PBS) in a
96 well V- or U-bottom plate (Corning, 353077) with combinations of the antibodies described in
Table S3 and the dilutions they were used at, based on 50 µl final staining volume. In all assays,
cells were incubated with anti-FcγRII/III (clone 93; 101302, BioLegend) before incubation with
fluorochrome-conjugated antibodies. Dead cells were excluded by fixable viability dye eFluor780
(65-0865-18, eBioscience) and gated on singlets. On average, a minimum of 50,000 live and singlet
cells were analyzed. Fluorescence minus one (FMO) and unstained controls were used for gating.
Data were acquired on a LSR II or Fortessa (BD) and analyzed using FlowJo (FlowJo LLC, Ashland,
USA).

Blood lipid profile. Murine serum samples were analyzed with the automated Dimension® RxL
Max® clinical chemistry system from Siemens at the Comparative Medicine Unit of CNIC. The assay
was performed using a Flex ® reagent cartridge. The analysis of the lipid profile included reagents
from Siemens (HDL, DF48B; Cholesterol DF27; DF131, LDL; DF55A, Lipase; DF69A, TGs),
Spinreact (41035, Free Cholesterol), and Wako (436-91995, 436-91995; NEFA).

Lipidomics. Data acquisition and analysis was performed at CEMBIO and CBQF. Common
reagents include reverse-osmosed ultrapure water from Millipore; LC-MS grade methanol (MeOH),
acetonitrile (ACN), isopropanol (IPA) and sodium hydroxide from Fisher Scientific, Honeywell or
VWR; HPLC grade methyl tert-butyl ether (MTBE), chloroform, ammonium fluoride (NH4F) (ACS
reagent, ≥ 98%) and ammonium formate from Sigma-Aldrich or VWR. Analytical grade ammonia

solution (28%, GPR RECTAPUR®) and acetic acid glacial (AnalaR® NORMAPUR®) were obtained
from VWR Chemicals.

Lipidomics at <u>CEMBIO.</u> Lipid extraction was performed as reported[59]. In brief, 50 µl of plasma from
63 mice were thawed on ice for approximately 1 hour and vortexed for 2 min. Subsequently, 800 µl
of a previously prepared mixture consisting of MeOH:MTBE:Chloroform (1.33:1:1, v/v/v) and 25 µl
light Splash Lipidomics (Mass Spec Standard, 330707, Avanti Polar Lipids, Inc.) used as the internal
standard were added to each sample at RT for deproteinization and lipid extraction. Then, samples
were vortex-mixed for 5 min at RT and centrifuged for 15 min at 15°C and 16,100 x g. Finally,
supernatants were transferred to chromatography vials with inserts and were directly injected into
the LC-MS system. In addition, QC samples were set in parallel to the sample preparation. The QC
was prepared by pooling equal volumes (30 µl) of each supernatant sample from the study. The QC
was split into 5 vials, and these were analyzed throughout the analysis process.
Data were acquired using an Agilent 1290 Infinity II UHPLC system coupled to an Agilent 6545
quadrupole time-of-flight (QTOF) mass spectrometer. The Agilent 1290 Infinity II Multisampler
system, equipped with a multi-wash option, was used to uptake 1 and 2 µl of extracted samples in
positive and negative ionization modes, respectively. The multisampler was maintained at 15°C to
preserve lipids in a stable environment and avoid precipitation. An Agilent InfinityLab Poroshell 120
ECsingle bondC18 (3.0 × 100 mm, 2.7 µm) (Agilent Technologies) column and a compatible guard
column (Agilent InfinityLab Poroshell 120 ECsingle bondC18, 3.0 × 5 mm, 2.7 µm) were used and
maintained at 50°C. The chromatography gradient started at 70% of B at 0 – 1 min, 86% at 3.5 – 10
min, 100% B at 11–17 min. The starting conditions were recovered by min 17, followed by a 2 min
re-equilibration time; the total running time was 19 min. The mobile phases used for both positive
and negative ionization modes consisted of (A) 10 mM ammonium acetate, 0.2 mM NH4F in 9:1
water/MeOH and (B) 10 mM ammonium acetate, 0.2 mM NH4F in 2:3:5 ACN/MeOH/IPA. The flow
rate was held constant, set at 0.6 ml/min. The multi-wash strategy consisted of a mixture of
MeOH:IPA (50:50, v/v) with the wash time set at 15 s, and aqueous phase:organic phase (30:70,
v/v) mixture to assist in the starting conditions.
The Agilent 6545 QTOF mass spectrometer equipped with a dual AJS ESI ion source was set with
the following parameters: 150 V fragmentor, 65 V skimmer, 3500 V capillary voltage, 750 V octopole
radio frequency voltage, 10 L/min nebulizer gas flow, 200°C gas temperature, 50 psi nebulizer gas
pressure, 12 L/min sheath gas flow, and 300°C sheath gas temperature. Data were collected in
centroid in mode positive and negative ESI modes in separate runs, operated in full scan mode from
50 to 1700 m/z with a scan rate of 3 spectra/s. A solution consisting of two reference mass

compounds was used throughout the whole analysis: purine ($C_5H_4N_4$) at *m/z* 121.0509 for the

positive and *m/z* 119.0363 for the negative ionization modes; and HP-0921 ($C_{18}H_{18}O_6N_3P_3F_{24}$) at *m/z*

922.0098 for the positive and *m/z* 980.0163 (HP-0921+acetate) for the negative ionization modes.

These masses were continuously infused into the system through an Agilent 1260 Iso Pump at a 1

mL/min (split ratio 1:100) to provide a constant mass correction. Ten Iterative-MS/MS runs were

performed for both ion modes at the end of the analytical run. They were operated with an MS and

MS/MS scan rates of 3 spectra/s, 40–1700 *m/z* mass window, a narrow ($\sim$ 1.3 amu) MS/MS isolation

width, 3 precursors per cycle, 5,000 counts, and 0.001% of MS/MS threshold. Five iterative-MS/MS

runs were set with a collision energy of 20 eV, and the subsequent five runs were performed at 40

eV. References masses and contaminants detected in blank samples were excluded from the

analysis to avoid inclusion in the iterative-MS/MS.

The data collected after the LC-MS analyses in both positive and negative ion modes were

reprocessed with the Agilent MassHunter Profinder B.10.0.2 software. The datasets were extracted

using the Batch Recursive Feature Extraction (RFE) workflow integrated into the software. This

workflow comprises two steps: the Batch Molecular Feature Extraction (MFE) and the Batch Find by

Ion Feature extraction (FbI). The MFE algorithm consists in removing unwanted information,

including the background noise, and then creating a list of possible components that represent the

full range of TOF mass spectral data features, which are the sum of coeluting ions that are related

by charge-state envelope, isotope pattern, and/or the presence of different adducts and dimers.

Additionally, the MFE is intended to detect coeluting adducts of the same feature, selecting the

following adducts: $[M+H]^+$, $[M+Na]^+$, $[M+K]^+$, $[M+NH4]^+$ and $[M+C2H6N2+H]^+$ in LC-MS positive

ionization; $[M-H]^-$, $[M+CH3COOH-H]^-$, and $[M+Cl]^-$ in LC-MS negative ion mode. The neutral loss

(NL) of water is also considered for both ion modes. The algorithm then aligns the molecular features

across the study samples using the mass and retention time to build a single spectrum for each

compound group. The next step involves FbI, using the median values derived from the MFE process

to perform a targeted extraction to improve the reliability of finding and reporting features from

complex datasets used for differential analysis.

After data pre-processing, quality assurance (QA) consisted of raw data filtration by keeping all

features that were present after blank subtraction, were detected in >50% of QCs and >75% in at

least one sample group and had Relative Standard Deviation (RSD) <30% in the QCs. The rest of

the signals were excluded from the analyses. Finally, 1,106 and 268 chemical entities were obtained

that passed LC-MS quality control in positive and negative ionization, respectively. Missing values

were replaced using the k-nearest neighbors (kNN) algorithm[60] using an in-house script developed

in Matlab® (v.R2018b, MathWorks®). The quality of the analyses was tested using PCA models[61].

The lipid annotation process was carried out in three steps: the first one was an initial tentative
identification of lipid features, based on the MS1 data, using our online tool CEU Mass Mediator
(CMM) (http://ceumass.eps.uspceu.es/mediator/)[62]. This tool for mass-based compound annotation
comprises the information available in different databases (KEGG, HMDB, LIPID MAPS, Metlin,
MINE and an in-house library). This stage started with the tentative assignment based on (i) accurate
mass (maximum mass error tolerance 20 ppm); (ii) retention time; (iii) isotopic pattern distribution;
(iv) the possibility of cation and anion formation; and (v) adduct formation pattern.
Secondly, to increase the level of confidence annotation, the raw LC-MS/MS data obtained was
imported to the Lipid Annotator software (Agilent Technologies Inc.) and the MS-DIAL software[63], to
build a fragmentation-based (MS/MS) library comprising the *m/z* of all the precursors identified as
lipids by the software, together with their corresponding RT. The Lipid Annotator method[64] was set
as follows: ion species $[M+H]^+$, $[M+Na]^+$, and $[M+NH4]^+$ for positive; and $[M-H]^-$, and $[M+CH3COOH-$
$H]^-$ for negative ionization mode. Then, for both ion modes, the Q-Score was set at $\geq$ 50; all the lipid
classes were selected, the mass deviation was established as $\leq$ 20 ppm, fragment score threshold
was fixed as $\geq$ 30, and the total score was set at $\geq$ 60. For the case of MS-DIAL software, analytical
parameters were also set as described: ion species $[M+H]^+$, $[M-H2O+H]^+$, $[M+Na]^+$, $[M+K]^+$,
$[M+NH4]^+$ and $[M+C2H6N2+H]^+$ for positive; and $[M-H]^-$, $[M-H2O-H]^-$, $[M+Cl]^-$, $[M+HCOOH-H]^-$ and
$[M+CH3COOH-H]^-$ for negative mode. Adduct formation with formic acid were observed
experimentally in this study even though the mobile phases used for LC are lacking this compound.
Formic acid presence was considered due to trace amount contamination levels of formate in acetate
salts of LC-MS grade, or MeOH oxidation during ESI. The search was fixed to be performed across
a mass range from 50 to 1500 Da (for both MS1 and MS/MS levels), the mass deviation accepted
was $\leq$ 0.01 Da and $\leq$ 0.025 for MS1 and MS/MS levels respectively, and the identification score cut
off was also set at $\geq$ 60.
Finally, a manual MS/MS spectral interpretation was carried out using the software Agilent
MassHunter (version 10.0), matching the retention time and MS/MS fragmentation to the available
spectral data included in MetFrag and Lipid Maps[65]. Different forms of cations and anions were
targeted for the same molecule to ensure its annotations. The interpretation of each obtained
spectrum led to the same ID in terms of belonging to the same lipid category and in terms of fatty
acids composition, which together give a robust and complete identification. The nomenclature used
in this article for the lipid species reported follows the latest update of the shorthand annotation.[66]
Lipid metabolic networks according to their chain length, saturation, and level of significance were
performed by version 1 LINEX webapp.

<u>Lipidomics at CBQF.</u> Lipid extraction from 20 plasma samples was performed as reported[67] with
some modifications. Briefly, tubes were added with 1.5 mL of MeOH and 5 mL of MTBE. Samples
were homogenized for 1 min followed by 15 min of sonication in ultrasound. This procedure was
repeated once again and then the tubes let under rotation (70 rpm) overnight (18 h). Afterward, tubes
were added with 1.25 mL of HPLC-water, homogenized for 1 min and centrifuged (1250 x g, 10 min,
room temperature). About 3 mL of the upper layer (MTBE containing the extracted lipids) were
recovered to a new tube previously weighted. Then, 3 mL of MTBE were added to extraction tubes
and these homogenized and centrifuged at the same above-mentioned conditions. Upper layer was
recovered, and the MTBE extract let to evaporate completely, under nitrogen, for further
determination of lipid extract mass.

Plasma lipid extracts were pooled according to their group, dissolved in IPA:ACN (9:1, v/v) at 0.2
mg/mL and analyzed in triplicate on an UHPLC instrument (Elute; Bruker), equipped with an Acquity
UPLC BEH C18 (17 μm) pre-column (Waters), an Intensity Solo 2 C18 (100 x 2.1 mm) column
(Bruker), and coupled with an UHR–QTOF detector (Impact II; Bruker). The injection method was
based on conditions previously reported with some modifications[68]. Thus, mobile phases consisted
of ACN:upH2O (6:4, v/v) (Phase A) and IPA:ACN (9:1, v/v) (Phase B), each one added with 0.1%
(v/v) formic acid and 10 mM ammonium formate. Phase B gradient flow was set as follows: 0.0 min:
40%, 2.0 min: 43%, 2.1 min: 50%, 12.0 min: 54%, 12.1 min: 70%, 18.0 min: 99%, 20.0 min: 99%,
20.1 min: 40%, and 22 min: 40%. Flow rate was set at 0.4 mL/min and column temperature at 55
°C. The injection volume was 3 μL in positive ionization mode and 5 μL in negative ionization mode.
For MS analysis, the following parameters were applied: end plate offset voltage 500 V, capillary
voltage 4500 V (positive ionization) or 3000 V (negative ionization), nebulizing gas pressure 35 psi,
drying gas flow 8 L/min, drying gas temperature 325 °C, quadrupole ion energy 3 eV (positive
ionization) or 5 eV (negative ionization), collision energy 10 eV (positive ionization) or 5 eV (negative
ionization). Acquisition was performed in an auto MS/MS scan mode over a mass range of m/z 50-
1500. For both ionization modes, an external mass calibration was performed with a solution of
IPA:upH2O (1:1, v/v) added with 0.2% (v/v) formic acid and 0.6% (v/v) NaOH 1M, continuously
injected at 180 μL/h.

The files resulted from LC-ESI-QTOF analysis were converted into abf format through Reifycs
Analysis Base File Converter (Reifycs Inc.). These converted files were used for lipid composition
analysis in MS-Dial 5.1.230517 (Riken). The alignment results were exported and identified
compounds were grouped by lipid class, according to LIPID MAPS database nomenclature. Peak
intensities of compounds belonging to same lipid class were totalized and used as data for heatmap
analysis in ClustVis, a web tool available for visualizing clustering of multivariate data[69]. The MSDial

software was able to detect 2,718 different ions among the serum samples, being identified 1,236
compounds by the software. From these data was excluded, among the compounds with attributed
equal identification, the ones that revealed considerably low signal, leading to a total of 1,198
compounds to assess further statistical analysis. For heatmap analysis, compounds were grouped
into 36 lipid classes.

TG measurements in human sera. Human TGs were measured using the Cobas c 501 analyzer
(Roche Diagnostic). For this purpose, serum samples (BD Vacutainer tubes) were collected from
each patient under two conditions: one during the onset of a hypersensitivity reaction and the other
at least 14 days after the reaction. If the allergic reaction occurred accidentally, the first samples
were collected in the emergency departments, whereas if it was due to a controlled challenge test,
the sera were obtained in the allergy units.

Study population included 118 paired sera from 59 patients presenting hypersensitivity reactions and
recruited from three Spanish hospitals (Hospital Universitario Fundación Jiménez Díaz, Hospital
Central de la Cruz Roja, Hospital Infantil Universitario Niño Jesús). Each patient's sex, age, reaction
trigger, clinical signs and symptoms were recorded. In addition, the severity of the reaction was
determined according to the criteria established by Brown[36]. The study was approved by the Ethics
Committee (CEIm FJD, PIC057-19), and authors adhered to the declaration of Helsinki. All patients
were included after giving informed consent by the donors or their parents. Inclusion criteria included
acceptance to participate in the study and an objective diagnosis of the reaction. Exclusion criteria
were the presence of a blood-borne disease, any psychiatric illness or psychological pathology that
prevented acceptance for the study or receiving any treatment prior to obtaining the acute phase
sample, as these interfere with TG levels.

Statistics. Independent experiments were performed with multiple biological replicates to ensure
reproducibility and statistical power to detect meaningful differences. GraphPad Prism v9 was used
for all statistical analyses. Quantitative variables are expressed as means ± standard error of the
mean (s.e.m.). Normal distribution was determined by D'Agostino and Pearson test, Shapiro–Wilk
test, Kolmogorov–Smirnov test, and Anderson-Darling test. A two-way analysis of variance (ANOVA)
or mixed-effects test were used to evaluate differences between more than two groups with a
parametric distribution, and Bonferroni´s multiple comparison test was applied. Data with
nonparametric distribution were evaluated with the Kruskal–Wallis test for more than two groups,
and the Mann–Whitney or Wilcoxon signed-rank test for two independent groups. A p-value<0.05
was considered significant.

For lipidomic data, multivariate analysis was performed using SIMCA v.16.0 (Sartorius Stedim Data
Analytics). The PCA models were used to evaluate data quality and find patterns in the experimental
samples. The models were unit variance scaled and evaluated using R^2 and Q^2 parametersm, which
are the classification and prediction capacity, respectively. Complementary, univariate analysis was
performed in MATLAB (v.R2018b, MathWorks®) or with the Lipidmaps' statistical tool to obtain the
significance of each feature in the study. The focus was set on the comparison between two groups
at day 7. For this statistical analysis, T-test student or Mann-Whitney U test was applied for each
metabolite, depending on the normality of the data, which were evaluated by the Kolmogorow-
Smirnow-Lilliefors test and the Levene test, respectively. The FDR was performed by using the
Benjamini-Hochberg correction for these p-values, and statistical significance was set at 95% level
(pFDR< 0.05 for the adjusted p-value)[70]. The MetaboAnalyst online tool (v. 5.0) was used to produce
heat maps with hierarchical clustering. Euclidean distance measure and Ward's clustering method
were chosen as the clustering parameters. For lipid maps, LINEX – LipidNetworkExplorer online tool
was used (https://exbio.wzw.tum.de/linex). GraphPad Prism v9 was used to represent plots with
metabolomics data.

AUTHOR CONTRIBUTIONS. N.F.-G, R.C.-G. and R.J.-S. performed the *in vivo* experiments and
analyzed the data. L.M.-S. did ELISAs. A.J.G.-C., A.G., D.O., L.M-S, A.L.-F. and L.L.-P. performed
lipidomics, and L.M.R.-A., A.V., D.B., and C.Barbas provided supervision and aided in lipidomics
data processing. E.S.-M. and C.L.-S. conducted mast cell cultures and activation assays. M.R.-H.
and I.R.-F. helped with experiments. E.N.-B., V.E., and C.Blanco provided samples from allergic
patients and/or measured their lipid profile. Y.R.C., B.I., P.M. and F.S.-M. provided expertise in
immunology and/or atherosclerosis throughout the project. R.J.-S conceptualized and oversaw the
project, designed experiments, and wrote the manuscript. All authors discussed the results, assisted
in the preparation of the manuscript, and approved its publication.

COMPETING INTERESTS: The authors declare no competing interests.

ACKNOWLEDGEMENTS. R.J.-S.´s laboratory has received support from the Severo Ochoa
Program (AEI/SEV-2017- 0712; Spain), FSE/FEDER through the Instituto de Salud Carlos III (ISCIII;
CP20/00043; PI22/00236; Spain), The Nutricia Research Foundation (NRF-2021-13; The
Netherlands), New Frontiers in Research Fund (NFRFE-2019-00083; Canada) and SEAIC
(BECA20A9; Spain). F.S.-M. reports grants from Ministerio de Ciencia e Innovación (MICIN;
PID2020-120412RB-I00; Spain), and Comunidad de Madrid (INTEGRAMUNE, P2022/BMD7209;
Spain). V.E. obtained support from the ISCIII (PI21/00158; Spain). N.F.-G. and I.R-F. received a

Formación de Profesorado Universitario grant (FPU16/03953 and FPU20/05176, respectively) from Ministerio de Universidades (Spain). N.F.-G. and L.M.-S. are supported by the INVESTIGO Program of the Comunidad de Madrid and Ministerio de Trabajo y Economía Social, Servicio Público de Empleo (SEPE), respectively, which is funded by "Plan de Recuperación, Transformación y Resiliencia" and "NextGenerationEU" of the European Union (09-PIN1-00015.6/2022 and 2022-C23.I01.P03.S0020-0000031). R.C.-G. is supported by Ayudas para contratos Juan de la Cierva-formación 2021 (FJC2021-047282-I) from MICIN (Spain). P.M. is supported by MICIN-ISCIII-Fondo de Investigación Sanitaria (PI22/01759; PMPTA22/00090-BIOCARDIOTOX) and Comunidad de Madrid (P2022/BMD-7209-INTEGRAMUNE-CM; Spain). The CNIC is supported by the ISCIII, the MICIN, and the Pro CNIC Foundation. CNIC is a Severo Ochoa Center of Excellence (grant CEX2020-001041-S funded by MICIN/AEI/10.13039/501100011033). The R.J.-S.´s lab recognizes Ana Cayuela, Ibon Redondo, Pedro L. Majano, Francisco Vega, María Ángeles Vallejo, Melissa Gordon, María Jesús Andrés and all the members of the R.J.-S. and F.S.-M. labs for technical help and/or scientific input.

REFERENCES

Molofsky, A. B. & Locksley, R. M. The ins and outs of innate and adaptive type 2 immunity. *Immunity* **56**, 704-722 (2023). https://doi.org:10.1016/j.immuni.2023.03.014

Gause, W. C., Rothlin, C. & Loke, P. n. Heterogeneity in the initiation, development and function of type 2 immunity. *Nature Reviews Immunology* **20**, 603-614 (2020). https://doi.org:10.1038/s41577-020-0301-x

Kopp, E. B., Agaronyan, K., Licona-Limón, I., Nish, S. A. & Medzhitov, R. Modes of type 2 immune response initiation. *Immunity* **56**, 687-694 (2023). https://doi.org:10.1016/j.immuni.2023.03.015

Akdis, C. A. Does the epithelial barrier hypothesis explain the increase in allergy, autoimmunity and other chronic conditions? *Nature Reviews Immunology* **21**, 739-751 (2021). https://doi.org:10.1038/s41577-021-00538-7

Knoflach, M. *et al.* Allergic Rhinitis, Asthma, and Atherosclerosis in the Bruneck and ARMY Studies. *Archives of Internal Medicine* **165**, 2521-2526 (2005). https://doi.org:10.1001/archinte.165.21.2521

Wilson, J. M. *et al.* IgE to the Mammalian Oligosaccharide Galactose-α-1,3-Galactose Is Associated With Increased Atheroma Volume and Plaques With Unstable Characteristics—Brief Report. *Arteriosclerosis, Thrombosis, and Vascular Biology* **38**, 1665-1669 (2018). https://doi.org:10.1161/atvbaha.118.311222

Roth, G. A. *et al.* Global Burden of Cardiovascular Diseases and Risk Factors, 1990–2019. *Journal of the American College of Cardiology* **76**, 2982-3021 (2020). https://doi.org:10.1016/j.jacc.2020.11.010

Libby, P. *et al.* Atherosclerosis. *Nature Reviews Disease Primers* **5**, 56 (2019). https://doi.org:10.1038/s41572-019-0106-z

9 Björkegren, J. L. M. & Lusis, A. J. Atherosclerosis: Recent developments. *Cell* **185**,
1630-1645 (2022). https://doi.org:10.1016/j.cell.2022.04.004
10 Fernández-Gallego, N. *et al.* The impact of type 2 immunity and allergic diseases in
atherosclerosis. *Allergy* **77**, 3249-3266 (2022). https://doi.org:10.1111/all.15426
11 Wang, H. *et al.* Global, regional, and national life expectancy, all-cause mortality, and
cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for
the Global Burden of Disease Study 2015. *The Lancet* **388**, 1459-1544 (2016).
https://doi.org:10.1016/s0140-6736(16)31012-1
12 Townsend, N. *et al.* Epidemiology of cardiovascular disease in Europe. *Nature Reviews*
Cardiology **19**, 133-143 (2021). https://doi.org:10.1038/s41569-021-00607-3
13 Galli, S. J. & Tsai, M. IgE and mast cells in allergic disease. *Nature Medicine* **18**, 693-
704 (2012). https://doi.org:10.1038/nm.2755
14 Finkelman, F. D., Khodoun, M. V. & Strait, R. Human IgE-independent systemic
anaphylaxis. *Journal of Allergy and Clinical Immunology* **137**, 1674-1680 (2016).
https://doi.org:10.1016/j.jaci.2016.02.015
15 Jönsson, F. *et al.* An IgG-induced neutrophil activation pathway contributes to human
drug-induced anaphylaxis. *Science Translational Medicine* **11**, eaat1479 (2019).
https://doi.org:10.1126/scitranslmed.aat1479
16 Galli, S. J., Tsai, M. & Piliponsky, A. M. The development of allergic inflammation.
Nature **454**, 445-454 (2008). https://doi.org:10.1038/nature07204
17 Crestani, E. *et al.* Untargeted metabolomic profiling identifies disease-specific signatures
in food allergy and asthma. *Journal of Allergy and Clinical Immunology* **145**, 897-906
(2020). https://doi.org:10.1016/j.jaci.2019.10.014
18 Jang, H. *et al.* Metabolomic profiling revealed altered lipid metabolite levels in childhood
food allergy. *Journal of Allergy and Clinical Immunology* **149**, 1722-1731.e1729 (2022).
https://doi.org:10.1016/j.jaci.2021.10.034
19 Obeso, D. *et al.* Multi-omics analysis points to altered platelet functions in severe food-
associated respiratory allergy. *Allergy* **73**, 2137-2149 (2018).
https://doi.org:10.1111/all.13563
20 Ellul, S. *et al.* Plasma metabolomic profiles associated with infant food allergy with
further consideration of other early life factors. *Prostaglandins, Leukotrienes and*
Essential Fatty Acids **159**, 102099 (2020). https://doi.org:10.1016/j.plefa.2020.102099
21 Dekkers, K. F. *et al.* Lipid-induced transcriptomic changes in blood link to lipid
metabolism and allergic response. *Nature Communications* **14**, 544 (2023).
https://doi.org:10.1038/s41467-022-35663-x
22 Jiménez-Saiz, R. *et al.* Lifelong memory responses perpetuate humoral T H 2 immunity
and anaphylaxis in food allergy. *Journal of Allergy and Clinical Immunology* **140**, 1604-
1615.e1605 (2017). https://doi.org:10.1016/j.jaci.2017.01.018
23 Chu, D. K. *et al.* Indigenous enteric eosinophils control DCs to initiate a primary Th2
immune response in vivo. *Journal of Experimental Medicine* **211**, 1657-1672 (2014).
https://doi.org:10.1084/jem.20131800
24 Ishibashi, S. *et al.* Hypercholesterolemia in low density lipoprotein receptor knockout
mice and its reversal by adenovirus-mediated gene delivery. *J. Clin. Invest.* **92**, 883-893
(1993). https://doi.org:10.1172/jci116663
25 Raposeiras-Roubin, S. *et al.* Triglycerides and Residual Atherosclerotic Risk. *Journal of*
the American College of Cardiology **77**, 3031-3041 (2021).
https://doi.org:10.1016/j.jacc.2021.04.059
26 Stegemann, C. *et al.* Lipidomics Profiling and Risk of Cardiovascular Disease in the
Prospective Population-Based Bruneck Study. *Circulation* **129**, 1821-1831 (2014).
https://doi.org:10.1161/circulationaha.113.002500

27 Kiechl, S. *et al.* Plasma Lipid Composition and Risk of Developing Cardiovascular
Disease. *PLoS ONE* **8**, e71846 (2013). https://doi.org:10.1371/journal.pone.0071846
28 Harshfield, E. L. *et al.* An Unbiased Lipid Phenotyping Approach To Study the Genetic
Determinants of Lipids and Their Association with Coronary Heart Disease Risk Factors.
Journal of Proteome Research **18**, 2397-2410 (2019).
https://doi.org:10.1021/acs.jproteome.8b00786
29 Bruton, K. *et al.* Interrupting reactivation of immunologic memory diverts the allergic
response and prevents anaphylaxis. *Journal of Allergy and Clinical Immunology* **147**,
1381-1392 (2021). https://doi.org:10.1016/j.jaci.2020.11.042
30 Ishizaka, K., Ishizaka, T. & Menzel, A. E. Physicochemical properties of reaginic
antibody. VI. Effect of heat on gamma-E-, gamma-G- and gamma-A-antibodies in the
sera of ragweed sensitive patients. *J Immunol* **99**, 610-618 (1967).
31 Dorrington, K. J. & Bennich, H. Thermally induced structural changes in immunoglobulin
E. *J Biol Chem* **248**, 8378-8384 (1973).
32 López-Sanz, C., Sánchez-Martínez, E. & Jiménez-Saiz, R. Protocol to desensitize
human and murine mast cells after polyclonal IgE sensitization. *STAR Protocols* **3**,
101755 (2022). https://doi.org:10.1016/j.xpro.2022.101755
33 Kubo, S., Nakayama, T., Matsuoka, K., Yonekawa, H. & Karasuyama, H. Long Term
Maintenance of IgE-Mediated Memory in Mast Cells in the Absence of Detectable Serum
IgE. *The Journal of Immunology* **170**, 775-780 (2003).
https://doi.org:10.4049/jimmunol.170.2.775
34 Martín-López, L. *et al.* Environmental exposure of Der p 23 in household dust samples.
Clinical & Experimental Allergy **51**, 1645-1647 (2021). https://doi.org:10.1111/cea.14004
35 Brough, H. A., Mills, E. N. C., Richards, K., Lack, G. & Johnson, P. E. Mass
spectrometry confirmation that clinically important peanut protein allergens are present
in household dust. *Allergy* **75**, 709-712 (2019). https://doi.org:10.1111/all.14070
36 Brown, S. G. A. Clinical features and severity grading of anaphylaxis. *Journal of Allergy*
and Clinical Immunology **114**, 371-376 (2004). https://doi.org:10.1016/j.jaci.2004.04.029
37 Ettinger, W. H. & Hazzard, W. R. Prednisone increases very low density lipoprotein and
high density lipoprotein in healthy men. *Metabolism* **37**, 1055-1058 (1988).
https://doi.org:10.1016/0026-0495(88)90067-4
38 Sheha, D. S., El-Korashi, L. A., AbdAllah, A. M., El-Begermy, M. M. & Elmahdi, A. R.
Dyslipidemia among allergic rhinitis patients: Frequency and risk factors. *World Allergy*
Organization Journal **14**, 100523 (2021). https://doi.org:10.1016/j.waojou.2021.100523
39 La Mantia, I., Andaloro, C., Albanese, P. G. & Varricchio, A. Blood lipid levels related to
allergic rhinitis: a significant association. *EuroMediterranean Biomed J* **12**, 144-147
(2017).
40 Rhee, T.-M., Choi, E.-K., Han, K.-D., Lee, S.-R. & Oh, S. Impact of the Combinations of
Allergic Diseases on Myocardial Infarction and Mortality. *The Journal of Allergy and*
Clinical Immunology: In Practice **9**, 872-880.e874 (2021).
https://doi.org:10.1016/j.jaip.2020.09.008
41 Sarwar, N. *et al.* Triglycerides and the Risk of Coronary Heart Disease. *Circulation* **115**,
450-458 (2007). https://doi.org:10.1161/circulationaha.106.637793
42 Austin, M. A., Hokanson, J. E. & Edwards, K. L. Hypertriglyceridemia as a
Cardiovascular Risk Factor. *The American Journal of Cardiology* **81**, 7B-12B (1998).
https://doi.org:10.1016/s0002-9149(98)00031-9
43 Nordestgaard, B. G. & Varbo, A. Triglycerides and cardiovascular disease. *The Lancet*
384, 626-635 (2014). https://doi.org:10.1016/s0140-6736(14)61177-6
44 Klempfner, R. *et al.* Elevated Triglyceride Level Is Independently Associated With
Increased All-Cause Mortality in Patients With Established Coronary Heart Disease.

Circulation: Cardiovascular Quality and Outcomes **9**, 100-108 (2016). https://doi.org/10.1161/circoutcomes.115.002104

Liu, D. J. *et al.* Exome-wide association study of plasma lipids in >300,000 individuals. *Nature Genetics* **49**, 1758-1766 (2017). https://doi.org:10.1038/ng.3977

Xu, Y., Guo, N., Dou, D., Ran, X. & Liu, C. Metabolomics analysis of anaphylactoid reaction reveals its mechanism in a rat model. *Asian Pacific Journal of Allergy and Immunology* **35**, 224-232 (2018). https://doi.org:10.12932/ap0845

Sun, S. *et al.* Integrating Widely Targeted Lipidomics and Transcriptomics Unravels Aberrant Lipid Metabolism and Identifies Potential Biomarkers of Food Allergies in Rats. *Molecular Nutrition & Food Research* **67**, e2200365 (2023). https://doi.org:10.1002/mnfr.202200365

Perales-Chorda, C. *et al.* Characterization of anaphylaxis reveals different metabolic changes depending on severity and triggers. *Clinical & Experimental Allergy* **51**, 1295-1309 (2021). https://doi.org:10.1111/cea.13991

Nuñez-Borque, E. *et al.* Proteomic profile of extracellular vesicles in anaphylaxis and their role in vascular permeability. *Allergy* **76**, 2276-2279 (2021). https://doi.org:10.1111/all.14792

Zhi, Y. *et al.* HR-MS Based Untargeted Lipidomics Reveals Characteristic Lipid Signatures of Wilson's Disease. *Front Pharmacol* **12**, 754185 (2021). https://doi.org:10.3389/fphar.2021.754185

Grandis, D. J. *et al.* Wilson's Disease and Cardiac Myopathy. *The American Journal of Cardiology* **120**, 2056-2060 (2017). https://doi.org:10.1016/j.amjcard.2017.08.025

Sun, J. *et al.* Impact of CD40 Ligand, B Cells, and Mast Cells in Peanut-Induced Anaphylactic Responses. *The Journal of Immunology* **179**, 6696-6703 (2007). https://doi.org:10.4049/jimmunol.179.10.6696

Bartemes, K. R. & Kita, H. Roles of innate lymphoid cells (ILCs) in allergic diseases: The 10-year anniversary for ILC2s. *Journal of Allergy and Clinical Immunology* **147**, 1531-1547 (2021). https://doi.org:10.1016/j.jaci.2021.03.015

Toms, T. E. Dyslipidaemia in Rheumatological Autoimmune Diseases. *The Open Cardiovascular Medicine Journal* **5**, 64-75 (2011). https://doi.org:10.2174/1874192401105010064

van Halm, V. P. *et al.* Lipids and inflammation: serial measurements of the lipid profile of blood donors who later developed rheumatoid arthritis. *Annals of the Rheumatic Diseases* **66**, 184-188 (2006). https://doi.org:10.1136/ard.2006.051672

Rodríguez-Carrio, J. *et al.* High triglycerides and low high-density lipoprotein cholesterol lipid profile in rheumatoid arthritis: A potential link among inflammation, oxidative status, and dysfunctional high-density lipoprotein. *Journal of Clinical Lipidology* **11**, 1043-1054.e1042 (2017). https://doi.org:10.1016/j.jacl.2017.05.009

Nuttall, S. L. Cardiovascular risk in systemic lupus erythematosus--evidence of increased oxidative stress and dyslipidaemia. *Rheumatology* **42**, 758-762 (2003). https://doi.org:10.1093/rheumatology/keg212

Vieira, P. & Rajewsky, K. The half-lives of serum immunoglobulins in adult mice. *European journal of immunology* **18**, 313-316 (1988). https://doi.org:10.1002/eji.1830180221

Gonzalez-Riano, C., Gradillas, A. & Barbas, C. Exploiting the formation of adducts in mobile phases with ammonium fluoride for the enhancement of annotation in liquid chromatography-high resolution mass spectrometry based lipidomics. *Journal of Chromatography Open* **1**, 100018 (2021). https://doi.org:10.1016/j.jcoa.2021.100018

Armitage, E. G., Godzien, J., Alonso-Herranz, V., López-Gonzálvez, Á. & Barbas, C. Missing value imputation strategies for metabolomics data. *Electrophoresis* **36**, 3050-3060 (2015). https://doi.org:10.1002/elps.201500352

61 Kellogg, J. J., Kvalheim, O. M. & Cech, N. B. Composite score analysis for unsupervised
comparison and network visualization of metabolomics data. *Analytica Chimica Acta*
1095, 38-47 (2020). https://doi.org:10.1016/j.aca.2019.10.029
62 Gil de la Fuente, A. *et al.* Knowledge-based metabolite annotation tool: CEU Mass
Mediator. *Journal of Pharmaceutical and Biomedical Analysis* **154**, 138-149 (2018).
https://doi.org:10.1016/j.jpba.2018.02.046
63 Tsugawa, H. *et al.* MS-DIAL 4: accelerating lipidomics using an MS/MS, CCS, and
retention time atlas. *bioRxiv*, 2020.2002.2011.944900 (2020).
https://doi.org:10.1101/2020.02.11.944900
64 Koelmel, J. P. *et al.* Lipid Annotator: Towards Accurate Annotation in Non-Targeted
Liquid Chromatography High-Resolution Tandem Mass Spectrometry (LC-HRMS/MS)
Lipidomics Using a Rapid and User-Friendly Software. *Metabolites* **10**, 101 (2020).
https://doi.org:10.3390/metabo10030101
65 Han, X. *Lipidomics: Comprehensive mass spectrometry of lipids*. (John Wiley & Sons,
2016).
66 Liebisch, G. *et al.* Update on LIPID MAPS classification, nomenclature, and shorthand
notation for MS-derived lipid structures. *Journal of lipid research* **61**, 1539-1555 (2020).
https://doi.org:10.1194/jlr.S120001025
67 Matyash, V., Liebisch, G., Kurzchalia, T. V., Shevchenko, A. & Schwudke, D. Lipid
extraction by methyl-tert-butyl ether for high-throughput lipidomics. *Journal of lipid*
research **49**, 1137-1146 (2008). https://doi.org:10.1194/jlr.D700041-JLR200
68 Teixeira, F. S. *et al.* Differential Lipid Accumulation on HepG2 Cells Triggered by
Palmitic and Linoleic Fatty Acids Exposure. *Molecules* **28** (2023).
https://doi.org:10.3390/molecules28052367
69 Metsalu, T. & Vilo, J. ClustVis: a web tool for visualizing clustering of multivariate data
using Principal Component Analysis and heatmap. *Nucleic Acids Research* **43**, W566-
W570 (2015). https://doi.org:10.1093/nar/gkv468
70 Benjamini, Y., Krieger, A. M. & Yekutieli, D. Adaptive linear step-up procedures that
control the false discovery rate. *Biometrika* **93**, 491-507 (2006).
https://doi.org:10.1093/biomet/93.3.491

961 **SUPPLEMENTARY FIGURE**

964 **Fig. S1 Assessment of eosinophilia by flow cytometry.** Cytometric identification of eosinophils in
965 the peritoneal lavage of naive and allergic mice at 3 days post-allergen challenge, based on F4/80
966 and Siglec-F expression gated on live, singlets and CD45+ cells. Representative plots of 4
967 independent experiments. FMO, fluorescence minus one control.

Quiescent allergy
Active allergy

Fig. S2 Allergic inflammation induces a unique blood lipid profile. Heatmap of the 171 identified
lipid species differentially expressed (p-value < 0.05) between allergic mice with quiescent allergy
(non-challenged, blue color, n=5) and active allergy (allergen-challenged, yellow color, n=6) 7 days
after challenge. The figure shows the clustering results in the form of a dendrogram. Relative lipid
abundance is represented along a gradient color from blue (diminished in active allergy) to red
(increased in active allergy). DG; diglycerides; Cer, ceramides; CAR, carnitines; LPE,
lysophosphatidylethanolamine; LPC, lysophosphatidylcholine; PC, phosphatidylcholine; CE,
cholesteryl ester; PI, phosphoinositide; LPI, lysophoinositide; FA, fatty acid; HerCer,
hexceramide; SM, sphingomyelin; PS, glycerophosphoserines; PE, glycerophosphoethanolamines;
PE-O-, oxidate glycerophosphoethanolamines; PC-O-, oxidate phosphatidylcholine; FAHFA, fatty
acid esters of hydroxy fatty acids; TG, triglycerides.

1032
1033 **Fig. S3 Assessment of CD4 T-cell depletion by flow cytometry.** Cytometric identification of CD4
1034 T cells in the spleen of naive and allergic mice 7 days after α-CD4- (clone GK1.5) or isotype-
1035 treatment based on CD4 expression (clone RM4-5) gated on live, singlets and CD45+ cells.
1036 Representative plots of 3 independent experiments.

1037
1038
1039
1040

1041

Fig. S4. Characterization of allergic sera for passive sensitization and measurement of serum lipids in circulation following allergen challenge. a Levels of peanut (PN)-specific IgE and IgG1 of naive and allergic sera (native of heated) used for passive sensitization. **b** Mast cell activation assays with cells sensitized with naive or allergic sera (native or heated) and challenged with different concentrations of crude peanut extract (CPE) (pooled data from 3 independent experiments). **c** Assessment of serum levels of total cholesterol, free cholesterol and LDL:HDL ratio at 1- and 7- days post-allergen challenge (PC) in passively sensitized mice with naive (n=17) or allergic sera (native or heated) to induce anaphylaxis via the classical (IgE; n=19), the alternative pathway (IgG; n=19), or both (IgE+IgG; n=18).

1051