Short Title: Engineering *Synechococcus* sp. PCC 11901

Title: **Engineering the highly productive cyanobacterium *Synechococcus* sp. PCC 11901**

Authors: Angelo J. Victoria[1,2*], Tiago Toscano Selão[3*], José Ángel Moreno-Cabezuelo[4],

Lauren A. Mills[4], Grant A. R. Gale[1,2], David J. Lea-Smith[4], Alistair J. McCormick[†,1,2]

[1] Institute of Molecular Plant Sciences, School of Biological Sciences, University of

Edinburgh, EH9 3BF, UK.

[2] Centre for Engineering Biology, School of Biological Sciences, University of Edinburgh,

EH9 3BF, UK.

[3] Department of Chemical and Environmental Engineering, University of Nottingham,

Nottingham NG7 2RD, UK.

[4] School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich,

NR4 7TJ, UK.

Corresponding author: Alistair McCormick, School of Biological Sciences, University of

Edinburgh, +44 (0)1316505316, alistair.mccormick@ed.ac.uk

On sentence summary: Genetic parts were characterised in *Synechococcus* sp. PCC 11901,

including a tightly regulated inducible promoter system, efficient CRISPRi and a novel

markerless Cas12a genome editing approach.

Author Contributions: AJV and TTS designed the study with input from AM and DL-S. AJV,

TTS, JAMC and LAM performed experiments. AM and AJV prepared the manuscript with

input from GARG, TTS, and DL-S.

ABSTRACT

Synechococcus sp. PCC 11901 (PCC 11901) is a fast-growing marine cyanobacterial strain that has a capacity for sustained biomass accumulation to very high cell densities, comparable to that achieved by commercially relevant heterotrophic organisms. However, genetic tools to engineer PCC 11901 for biotechnology applications are limited. Here we describe a suite of tools based on the CyanoGate MoClo system to unlock the engineering potential of PCC 11901. First, we characterised neutral sites suitable for stable genomic integration that do not affect growth even at high cell densities. Second, we tested a suite of constitutive promoters, terminators, and inducible promoters including a 2,4-diacetylphloroglucinol (DAPG)-inducible PhlF repressor system, which has not previously been demonstrated in cyanobacteria and showed tight regulation and a 228-fold dynamic range of induction. Lastly, we developed a DAPG-inducible dCas9-based CRISPR interference (CRISPRi) system and a modular method to generate markerless mutants using CRISPR-Cas12a. Based on our findings, PCC 11901 is highly responsive to CRISPRi-based repression and showed high efficiencies for single insertion (31-81%) and multiplex double insertion (25%) genome editing with Cas12a. We envision that these tools will lay the foundations for the adoption of PCC 11901 as a robust model strain for engineering biology and green biotechnology.

INTRODUCTION

Climate change has necessitated a global shift towards more sustainable production practices and the building of a bioeconomy centred on Net Zero Carbon policies (*Net Zero Strategy*, 2021). Cyanobacteria are an attractive alternative to heterotrophic microbial bioproduction chassis, such as *Escherichia coli* and yeast, due to their capacity for biology-based carbon capture and utilisation and potential for the production of a wide array of useful chemicals (Zhang et al., 2017; Daneshvar et al., 2022). Unicellular model cyanobacterial strains, such as *Synechocystis* sp. PCC 6803 (hereafter PCC 6803) and *Synechococcus elongatus* PCC 7942, have been investigated as biorefineries for bulk commodity products such as biofuels (Kopka et al., 2017), bioplastics (Khetkorn et al., 2016), natural colourants (Puzorjov et al., 2022) and terpenoids (Rautela and Kumar, 2022), while filamentous strains such as *Arthrospira platensis* have been developed for the production of food and high-value therapeutic antibodies (Jester et al., 2022; Saveria et al., 2022). Nevertheless, slow growth rates and low biomass productivity remain key bottlenecks that limit the economic competitiveness and commercial expansion of cyanobacterial biotechnology (Lea-Smith et al., 2021).

Several fast-growing cyanobacterial strains have been reported that show strong potential for

overcoming yield challenges (for recent review see Selão, 2022), including fresh water strains

Synechococcus elongatus UTEX 2973 (hereafter UTEX 2973) (Yu et al., 2015),

Synechococcus elongatus PCC 11801 (hereafter PCC 11801) (Jaiswal et al., 2018),

Synechococcus elongatus PCC 11802 (Jaiswal et al., 2020), and marine strains

Synechococcus sp. PCC 7002 (hereafter PCC 7002) (Batterton and Van Baalen, 1971) and

Synechococcus sp. PCC 11901 (hereafter PCC 11901) (Włodarczyk et al., 2020). Marine

cyanobacteria are of particular interest as they can utilise sea/brackish water, circumventing

the need for freshwater resources (Hitchcock et al., 2020). In contrast to other fast-growing

strains, PCC 11901 has an additional capacity for sustained growth to high densities (up to 30

g/L dry cell weight, DCW), similar to the maximum biomass accumulation observed for

batch-cultured *E. coli* cultures (24.7 g/L DCW) (Soini et al., 2008). Furthermore, PCC 11901

can tolerate high light intensities (>900 µmol photons m^{-2} s^{-1}), temperatures (up to 43°C), and

salinities over 2-fold higher than sea water (Cho et al., 2023). PCC 11901 is amenable to

natural transformation and its genome is fully-sequenced (Włodarczyk et al., 2020). PCC

11901 has also been engineered to produce free fatty acids yielding >6 mM (1.5 g L^{-1}), which

is comparable to that achieved by similarly engineered heterotrophic organisms (Xiao et al.,

2018). The recent isolation of a cobalamin-independent strain of PCC 11901 (*Synechococcus*

sp. UTEX 3154) that does not require the addition of vitamin B12 will help reduce scale-up

costs (Mills et al., 2022). Together, these qualities make PCC 11901 an attractive model

species for fundamental research and biotechnology applications.

In recent years, standardised molecular tools have been developed to help progress

synthetic/engineering biology in cyanobacteria, such as the CyanoGate Modular Cloning

system (Vasudevan et al., 2019) and the Start-Stop Assembly method (Taylor et al., 2021).

Additional advances include the development of self-replicating vectors for

extrachromosomal expression (Taton et al., 2014; Opel et al., 2022), recombineering (Jones

et al., 2021), genetic circuits (Taton et al., 2017; Zhang et al., 2022), CRISPR interference

(CRISPRi) (Yao et al., 2020), and CRISPR-based gene editing tools to reprogram

metabolism (Baldanta et al., 2022; Cengic et al., 2022; Wang et al., 2023). Strategies for

biocontainment of mutant cyanobacterial strains have also been developed, for example,

using inducible kill-switches (Zhou et al., 2019). Nevertheless, such tools are often not

readily transferrable between different species, and to date very few engineering approaches

have been characterised in PCC 11901. Notably, Mills et al. (2022) recently reported that

PCC 11901 was not compatible with the negative selection markers *sacB* and *codA* used for

generating markerless mutants (i.e. a genome-modified mutant that lacks a selective

antibiotic resistance cassette) that have key advantages for biotechnology applications (Lea-

Smith et al., 2016).

In this study we investigated a broad set of available genetic parts and developed several

novel tools for engineering PCC 11901. We tested five putative neutral integration sites to

serve as loci for introducing heterologous DNA and explored the amenability of PCC 11901

to transconjugation. We then characterised the functionality of a suite of known and new

genetic parts, including the 2,4-diacetylphloroglucinol (DAPG)-inducible PhlF repressor

system. Building on this, we developed an inducible CRISPRi gene repression system and a

novel hybrid plasmid approach to successfully generate markerless mutants using CRISPR-

Cas12a. Together these tools should fast-track the further development of PCC 11901 as a

commercially viable chassis strain for cyanobacterial biotechnology.

RESULTS AND DISCUSSION

Identification of robust neutral sites in PCC 11901 suitable for high-density growth

PCC 11901 is naturally transformable but exhibits partial resistance to kanamycin

(Włodarczyk et al., 2020). We first tested the susceptibility of PCC 11901 to five common

selective antibiotics used to generate marked mutants (**Figure 1A**). We observed that PCC

11901 exhibited partial resistance to kanamycin at 50 µg mL^{-1} and gentamicin at 10 µg mL^{-1}.

In contrast, PC11901 was fully susceptible to spectinomycin, erythromycin, and

chloramphenicol at all concentrations tested. Previous work has reported that many

filamentous and unicellular species have innate resistances to specific antibiotics, including

aminoglycosides such as kanamycin and gentamicin (Dias et al., 2015). Aminoglycoside

resistance has been attributed to native redox-active compounds such as glutathione in

Synechocystis sp. PCC 6803 (Cameron and Pakrasi, 2011), and the presence of native

resistance genes such as the kanamycin aminoglycoside acetyltransferase homolog recently

described in *Arthrospira platensis* (NIES39_D01030, GenBank ID: BAI89523.1) (Jester et

al., 2022; Shiraishi and Nishida, 2022), which is a member of the Gcn5-related N-

acetyltransferase (GNAT) superfamily (Favrot et al., 2016). A BLAST analysis of the PCC

11901 genome yielded 13 genes belonging to the GNAT family. The closest homolog to

BAI89523.1 (FEK30_08270, GenBank ID: QCS49435.1) shared a 33% peptide sequence

identity with similar conserved domains and a coenzyme A binding pocket motif, and thus

may account for the partial resistance observed (**Supplementary Figure S1**).

Neutral sites are genomic loci that allow for stable integration of heterologous genes into the

genome with no or minimal phenotypic impact. Several neutral sites have been identified in

model cyanobacterial species, such as PCC 6803 and PCC 7002 (Pinto et al., 2015; Ruffing et

al., 2016; Vogel et al, 2017). However, no neutral sites have yet been characterised in PCC

11901, specifically at the high densities achievable in this strain. We identified five putative

neutral site loci in PCC 11901 based on previously identified neutral integration sites in other

cyanobacteria and analysis of the PCC 11901 genome. *desB* (FEK_04840) encodes for a

putative fatty acid desaturase, which in PCC 7002 is involved in modulating membrane

fluidity at temperatures below 22ºC but does not impact growth at 30ºC (Sakamoto et al.,

1997; Ruffing et al., 2016). *glgA1* (FEK_14880) encodes for one of two putative glycogen

synthase isoforms in PCC 11901, which has previously been characterised as a neutral site in

PCC 6803 and PCC 11801 (Sengupta et al., 2020; Mittermair et al., 2021). The loci for *mrr*

(FEK30_09380) and *aquI* (FEK30_10065) encode for a putative Type IV restriction

endonuclease and a Type II site-specific deoxyribonuclease, respectively, and were selected

based on the hypothesis that several endonuclease genes may be redundant for immunity and

not essential for growth. Studies in other cyanobacterial strains, such as

Thermosynechococcus elongatus BP-1 and PCC 6803, have also reported improved

transformation efficiencies in nuclease mutants (Kufryk et al., 2002; Iwai et al., 2004).

Lastly, we identified an intergenic region of 185 bp between two convergent predicted open

reading frames encoding hypothetical proteins (FEK30_11550 and FEK30_11555), which we

hypothesised would not contain important regulatory elements. We called this locus neutral

site 1 (NS1). Such regions have previously been used successfully as neutral sites in PCC

6803 and other bacterial chassis strains, including *E. coli, Bacillus subtilis* and *Pseudomonas*

putida (Bernhards et al., 2022).

To establish and validate the concentrations at which selection can be performed, we chose

the putative neutral site *desB*, previously tested in PCC 11901 (Mills et al., 2022), and

introduced antibiotic resistance cassettes (AbRs) for each antibiotic via natural transformation

(**Figure 1B**). Following three days of growth, we observed hundreds of transformant colonies

for each AbR at concentrations of 100 µg mL^{-1} kanamycin, 25 µg mL^{-1} spectinomycin, 50 µg

mL^{-1} gentamicin, 1.25 µg mL^{-1} erythromycin, and 5 µg mL^{-1} chloramphenicol, respectively.

Based on the robust susceptibility of PCC 11901 to spectinomycin and previous success with

using spectinomycin for selection in model strains (Vasudevan et al., 2019), we next

assembled integrative vectors carrying a spectinomycin resistance cassette (SpR) flanked by

homologous regions for each of the five target putative neutral sites to facilitate homologous

recombination (HR), and transformed these into wild-type PCC 11901 to produce the mutants

Δmrr::SpR, $\Delta aquI$::SpR, $\Delta desB$::SpR, $\Delta glgA1$::SpR, and $\Delta NS1$::SpR (**Figure 1C**).

Remarkably, we observed full segregation for each of the five mutants following a single re-

streak from the transformation plates containing 25 µg mL^{-1} spectinomycin (**Supplementary**

Figure S2). The segregated mutants were then subjected to a comparative growth analysis to

assess the suitability of each putative neutral site. We observed that mrr and $aquI$ were the

best performing neutral sites, with Δmrr::SpR and $\Delta aquI$::SpR reaching optical densities

similar to wild-type (i.e. $OD_{750} > 100$). In contrast, $\Delta desB$::SpR and $\Delta NS1$::SpR grew

similarly to wild-type up to $OD_{750} \sim 50$, but then growth rates declined, suggesting that these

neutral sites should only be used at lower cell densities. The growth rate of $\Delta glgA1$::SpR

declined from $OD_{750} \sim 20$, indicating that $glgA2$ is not able to compensate for the loss of

$glgA1$ as observed in other strains (Sengupta et al., 2020; Mittermair et al., 2021). Notably,

all five mutants grew to higher densities compared to wild-type PCC 6803 and PCC 7002.

Together, our results show that endonuclease encoding genes are promising targets for

identifying additional robust neutral sites in PCC 11901.

PCC 11901 is amenable to conjugation

Delivery of heterologous DNA into genetically tractable cyanobacteria can also be achieved

through conjugal transfer or triparental mating using an *E. coli* 'helper' strain with

appropriate transmissible plasmids to enable conjugation of mobilizable plasmid vectors,

such as RSF1010-based vectors (Gale et al., 2019). As genome integration and segregation

are not required, conjugation has proved to be a powerful tool for rapidly testing genetic parts

and libraries in several cyanobacterial strains (Bishé et al., 2019; Puzorjov et al., 2021). We

sought to determine whether PCC 11901 is amenable to conjugation by introducing the

empty RSF1010-based CyanoGate acceptor vectors pPMQAK1-T and pPMQAK1-T-eYFP,

previously assembled by Vasudevan et al., (2019), which both contain a kanamycin

resistance cassette (KmR) (**Figure 2A**). We initially obtained colonies on selective media

containing kanamycin (**Figure 2B**), but subsequently found that a portion of re-streaked

colonies exhibited a yellowing or chlorotic phenotype and were unable to survive successive

rounds of streaking (**Figure 2C, D**). We hypothesised that the initial colonies were likely

false-positives due to the native kanamycin resistance of PCC 11901 (**Figure 1A,**
Supplementary Figure S1). Thus, we constructed new acceptor vectors pPMQSK1-1 and
pPMQSK1-T (for level 1 and level T in CyanoGate) that carried SpR, and assembled
pPMQSK1-1 vectors carrying either an eYFP (pPMQSK1-1-eYFP) or *Francisella novicida*
derived Cas12a (also known as *Fn*Cpf1) expression cassette (pPMQSK1-1-Cas12a).
Following conjugal transfer of pPMQSK1-1-based vectors into PCC 11901 we observed no
chlorotic phenotypes in any transconjugant strains after re-streaking (**Figure 2D**),
demonstrating that spectinomycin allows for robust selection of transconjugants.

We next performed a comparative growth analysis to explore the potential impact of
RSF1010-based vectors on high density growth in PCC 11901. We included a stably
integrated Δ*mrr*::eYFP mutant for comparison between transconjugants and transformants
(pC1.493, **Supplementary Table S2**). The growth of transconjugants with empty acceptor
vectors pPMQAK1-T or pPMQSK1-1 were not significantly different to wild-type PCC
11901 and Δ*mrr*::eYFP (**Figure 2E**). However, growth was reduced in transconjugants with
pPMQAK1-T-eYFP, pPMQSK1-1-eYFP or pPMQSK1-1-Cas12a, suggesting that the
expression of genes from RSF1010-based vectors could represent a metabolic burden for
PCC 11901, at least at high cell densities (Meyer, 2009). A comparison of eYFP fluorescence
levels between Δ*mrr*::eYFP and the transconjugants with pPMQAK1-T-eYFP or pPMQSK1-
1-eYFP showed that eYFP expression was $50 \pm 1\%$ lower in both transconjugants compared
to the stable integration mutant (**Figure 2F-G**). Overall, our results show for the first time
that conjugation is feasible in PCC 11901. However, for this species the speed of
transformation and segregation for genome integration appears more rapid than using
conjugation. Furthermore, protein expression levels in stable mutants exceeded
transconjugants with RSF1010-based vectors (at least for eYFP), suggesting that stable
integration may be the favoured engineering choice in PCC 11901. Nevertheless, the capacity
to maintain self-replicating vectors in PCC 11901 could still greatly facilitate the
characterisation of genetic parts, an important aspect of strain engineering, and facilitate
transient gene expression.

Characterisation of constitutive promoters and transcriptional terminators in PCC
11901
Constitutive promoters provide a stable level of gene expression and are essential parts in the
engineering toolbox of any chassis strain. Thus, we characterised a suite of 12 constitutive

promoters derived from Vasudevan et al. (2019) to provide a promoter library with varying

strengths for use in engineering PCC 11901. Included were constitutive promoters from PCC

6803 (P_{cpc560} and P_{psbA2L}) and synthetic promoters (P_{J23119}, P_{J23115}, P_{J23113}, P_{J23111}, P_{J23110},

P_{J23103}, P_{J23101}, P_{V02}, P_{V07} and P_{trc10}). Promoter strengths were assessed as expression cassettes

driving eYFP through stable genomic integration at the *mrr* neutral site, or on the self-

replicating RSF1010-based vector pPMQAK1-T. We were successful in assembling and

characterising all 12 constitutive promoters following conjugation using pPMQAK1-T

(**Figure 3A**). We were also able to generate and characterise 10 segregated integrative

transformants at the *mrr* neutral site (**Figure 3B**). Unfortunately, we were not able to generate

transformant colonies for P_{J23119} and P_{trc10} despite multiple attempts. Overall, the three

strongest constitutive promoters were P_{cpc560}, P_{J23119}, P_{psbA2L}, which is consistent with findings

in other cyanobacterial species (Li et al., 2018; Vasudevan et al., 2019; Sengupta et al.,

2020). The PCC 6803 *cpc* operon promoter (P_{cpc560}) was used to drive high expression levels

of heterologous genes in several model species, including PCC 6803 and PCC 7002. The

P_{psbA2L} promoter drives expression of the photosystem II reaction centre subunit D1. The high

light levels used to grow PCC 11901 may support the strong levels of expression seen with

P_{psbA2L}, similar to previous work in PCC 6803 and UTEX 2973 under high light conditions

(Lindberg et al., 2010; Sakurai et al., 2012; Li et al., 2018). Although the observed trend in

promoter strength showed a strong correlation ($R^2 = 0.95$) between those integrated at *mrr*

and on pPMQAK1-T (**Figure 3C**), we found that genomic integration resulted in a 32%

mean increase in eYFP expression for each promoter over those from RSF1010-based

vectors. Previous work in PCC 6803 has observed contrasting results, with promoters on

RSF1010-based vectors achieving a 3-fold higher level of expression compared to neutral site

integrations (Ng et al., 2015). The latter finding was attributed to an increased plasmid copy-

number for the RSF1010-based vector compared to the native chromosome. Thus, in PCC

11901 the copy number ratio may be lower for RSF1010-based vectors relative to the native

chromosomes. The chromosomal copy number for PCC 11901 remains unclear and should be

a focus for further work.

We next characterised the efficiencies of a suite of 10 transcriptional terminators in PCC

11901 based on those previously characterised in PCC 6803, UTEX 2973 and *E. coli* by Gale

et al. (2021). Cyanobacteria rely on the mechanism of rho-independent transcription

termination or intrinsic termination, which is defined by the formation of a hairpin loop on

the terminator sequence that leads to dissociation of the RNA polymerase and release of the

mRNA transcript (Wilson and von Hippel, 1995). Our suite of intrinsic terminators included

six native terminators from *E. coli* (T$_{ECK120010850}$, T$_{ECK120033736}$, T$_{ECK120029600}$, T$_{bba\_B0011}$,

T$_{bba\_B0061}$, and T$_{rrnB}$), two synthetic terminators derived from *E. coli* sequences (T$_{L3S2P21}$ and

T$_{L3S2P11}$), and two native terminators from PCC 6803 from photosystem II subunit D1 (T$_{psbA2}$)

and photosystem I subunit B (T$_{psaB}$) (Chen et al., 2013; Liu & Pakrasi, 2018). Of these, six

had termination efficiency (TE) values of >95%, with highest and lowest TE values observed

for T$_{ECK120029600}$ (99.8%) and T$_{bba\_B0061}$ (47.8%), respectively (**Figure 3D**). Our results were

relatively consistent with those from PCC 6803 and UTEX 2973, supporting our previous

observation that these transcriptional terminators perform similarly across different

cyanobacterial chassis (Gale et al., 2021).

Inducible gene expression systems in PCC 11901

Chemically inducible promoters can be used to modulate gene expression in response to an

external stimulus and are powerful tools for fundamental research and advanced engineering

approaches (e.g. gene circuit assembly) (Meyer et al., 2019). We first tested the L-rhamnose

inducible promoter P$_{rhaBAD}$ and its cognate transcription factor, RhaS, which has previously

been characterised in PCC 6803 (Kelly et al., 2018; Behle et al., 2020). To evaluate this

system in PCC 11901, we initially investigated if L-rhamnose impacted the growth of wild-

type PCC 11901 over a 72 h growth period. We found that PCC 11901 growth was not

affected by concentrations of up to 20 mM L-rhamnose, which was twice the highest

concentration tested by Behle et al. (2020), suggesting that L-rhamnose is not toxic to PCC

11901 (**Supplementary Figure S3A**). We then constructed an RSF1010-based reporter

system RhaS/P$_{rhaBAD}$-*eYFP*-T$_{rrnB}$ in the pPMQAK1-T vector, using the medium strength

constitutive promoter P$_{J23101}$ to drive RhaS expression and eYFP as the fluorescent reporter

(**Figure 4A**). Following successful conjugal transfer into PCC 11901, cultures were induced

with increasing concentrations of L-rhamnose (0-20 mM) and eYFP fluorescence was

measured after 24 h (**Figure 4B**). In the uninduced state (0 mM L-rhamnose), a low level of

eYFP fluorescence was detected (305 ± 9 AU, 6.4% of the maximum expression level),

demonstrating that the P$_{rhaBAD}$ promoter was leaky. Upon induction, we found that the

promoter achieved maximum expression (4,776 ± 185 AU) with 10 mM L-rhamnose, giving

a 15-fold dynamic range. Leaky expression with the P$_{rhaBAD}$ promoter has been reported

previously in PCC 6803 (Liu et al., 2020). Those authors successfully increased the control of

expression by replacing the promoter ribosome binding site with a theophylline riboswitch,

although their system required two small molecules for induction.

RNA riboswitches are structured noncoding RNA domains that can regulate gene expression

and exert translational control in bacteria by binding small molecules (Kavita and Breaker,

2023). We tested the engineered theophylline-inducible riboswitch E* fused to the *trc*

promoter (P_{trc}) first demonstrated in PCC 7942 (Nakahira et al., 2013). We observed that

growth of wild-type PCC 11901 was not affected at concentrations up to 2 mM theophylline

but was reduced at 5 mM (**Supplementary Figure S3B**). This is consistent with reports in

other cyanobacteria of the negative impact of theophylline on growth at concentrations above

2 mM (Ma et al., 2014). We then assembled an RSF1010-based reporter harbouring P_{trcE*}-

eYFP-T_{rrnB} in the pPMQAK1-T vector for conjugal transfer into PCC1901 (**Figure 4C**). In

the absence of theophylline, an eYFP fluorescence signal above background levels was still

detected in conjugant strains (336 ± 163.38 AU, 4.8% of the maximum expression level),

indicating that P_{trcE*} was leaky in PCC 11901, similar to P_{rhaBAD} (**Figure 4D**). Maximum

eYFP expression (8,555 ± 315 AU) was achieved with 2 mM theophylline (giving a 25-fold

dynamic range), but expression levels decreased at 5 mM theophylline, likely due to growth

inhibition. The observed leaky expression with P_{trcE*} was consistent with reports in several

other cyanobacterial species (Taton et al., 2017; Chi et al., 2019).

We sought to identify an inducible system for PCC 11901 that has tighter regulation than the

P_{rhaBAD} and P_{trcE*} promoters, so we next evaluated the 2,4-diacetylphloroglucinol (DAPG)-

inducible promoter P_{phlF} and its cognate transcription factor PhlF, which has tight regulation

in *E. coli* but has not yet been characterised in cyanobacteria (Meyer et al., 2019). In contrast

to the RhaS/P_{rhaBAD} system, the PhlF transcription factor functions as a repressor to the P_{phlF}

promoter, which undergoes a conformational change upon binding to DAPG and releases the

promoter leading to transcription, similar to the TetR family of repressors (Abbas et al.,

2002). We assembled two pPMQAK1-T vectors with i) a no TF control harbouring the

expression cassette PhlF/P_{phlF}-*eYFP*-T_{rrnB} to test the functionality of the P_{phlF} promoter in the

absence of the PhlF repressor transcription factor and DAPG, and ii) the reporter system

phlF/P_{phlF}-*eYFP*-T_{rrnB}, where the medium strength constitutive promoter PJ$_{23101}$ was used to

drive PhlF expression (**Figure 4E**). Following conjugation into PCC 11901, we observed

robust levels of YFP expression for the P_{phlF} promoter in the no TF control (94% of P_{cpc560})

(**Figure 2F** and **4F**). For the reporter system, increasing levels of eYFP fluorescence was

observed with increasing DAPG concentrations, with maximum expression (6,440 ± 279 AU)

achieved upon induction with 10 µM DAPG, similar to that for the no TF control. We

observed near background levels for eYFP fluorescence (28 ± 27 AU) in the uninduced state

(0 µM DAPG), indicating very tight repression of P$_{phlF}$ by PhlF. We found that DAPG had no

impact on the growth of PCC 11901 up to 10 µM, but that growth rates were reduced at 25

µM DAPG, indicating partial toxicity at higher DAPG concentrations (**Supplementary**

Figure S3C). The latter result was not unexpected, as DAPG has been shown to have broad

spectrum activity against bacteria and fungi, particularly those pathogenic to plants (Keel,

1992; Julian et al., 2021). To the best of our knowledge this is the first report of the PhlF/P$_{phlF}$

system being successfully utilised in a cyanobacterial strain. Overall, it provided a wide, 228-

fold dynamic range of induction in PCC 11901 and showed tight repression in the absence of

DAPG.

Gene repression using CRISPRi

CRISPRi is now well established as a powerful tool to explore gene function and pathways in

a variety of model cyanobacterial species (Liu et al., 2020; Yao et al., 2020; Dallo et al.,

2023). However, the number of characterised induction systems for CRISPRi remains

relatively   low   for   cyanobacteria.   Many   publications   utilise   the   TetR-based

anhydrotetracycline (aTc)-inducible promoters (Huang and Lindblad, 2013), but partial

leakiness (i.e. transcriptional repression in the absence of the inducer molecule) and the

degree of repression of the available systems can limit effective application, depending on the

strain used. Furthermore, aTc is photosensitive and degrades rapidly in UV or blue light (Zess

et al., 2016), which we believed would be problematic given the light intensities used to

culture PCC 11901 for high density growth and exposure to sunlight for outdoor growth.

To alleviate the potential challenge of aTc instability under high light, we sought to develop a

robust and tightly regulated inducible dCas9 CRISPRi approach in PCC 11901 by testing our

three inducible promoter systems. Using the CyanoGate system, we assembled the dCas9

expression cassettes *rhaS*/P$_{rhaBAD}$-*dCas9*-T$_{rrnB}$, P$_{trcE*}$-*dCas9*-T$_{rrnB}$ and *phlF*/P$_{phlF}$-*dCas9*-T$_{rrnB}$

in level 1 position 1 (L1P1). In level 1 position 2 (L1P2), we assembled four different single-

guide RNA (sgRNA) expression cassettes (P$_{trc10}$_TSS-sgRNA-sgRNA scaffold) targeting eYFP

at four different sequence locations (**Figure 5A**) (Vasudevan et al., 2019). As a control we

also assembled a constitutive expression cassette with a low strength synthetic promoter

(P$_{J23113}$-*dCas9*-T$_{rrnB}$), as previous work has indicated that strong expression of large proteins,

such as dCas9, can produce a metabolic burden that negatively impacts growth (Depardieu

and Bikard, 2020). L1P1 and L1P2 vectors were then assembled together into the level T

acceptor pUC19-T pCAT.015 vector for subsequent integration into the *desB* neutral site of

the Δ*mrr*::eYFP strain.

We initially tested the functionality of dCas9 in PCC 11901 using the weak constitutive

promoter P$_{J23113}$. Following transformation and segregation, we found that only strains

carrying both dCas9 and an sgRNA showed a reduction in eYFP fluorescence, which was

similar for all four sgRNAs used (52 ± 3.9%) (**Figure 5B**). No significant impacts on growth

were observed between strains (**Supplementary Figure S4A**). Although this demonstrated

that CRISPRi-dCas9 was functional in PCC 11901, the reductions in eYFP fluorescence were

unimpressive, which we attributed to the low strength of the promoter P$_{J23113.}$ Our results are

in line with those in PCC 7002 and *E. coli* where weaker promoters driving dCas9 expression

resulted in lower repression of the target gene (Gordon et al., 2016; Fontana et al., 2018).

We then tested the three inducible CRISPRi-dCas9 systems by inducing transformed cultures

with 10 mM L-rhamnose, 2 mM theophylline, or 10 µM DAPG, respectively, and measured

the change in eYFP fluorescence. Leakiness of dCas9 expression was investigated by

measuring eYFP fluorescence in the absence of the respective inducer molecules. For the

RhaS/P$_{rhaBAD}$ CRISPRi-dCas9 system we observed a 45-58% reduction in eYFP fluorescence

in the absence of L-rhamnose (**Figure 5C**), indicative of the P$_{rhaBAD}$ promoter leakiness

observed previously (**Figure 4B**). P$_{rhaBAD}$ appeared able to constitutively express dCas9 at

similar levels to that of the weak constitutive promoter P$_{J23113}$. Overall, our results were

consistent with the leakiness reported for P$_{rhaBAD}$ when used to express ddCas12a in PCC

6803 (Liu et al., 2020). Induction with L-rhamnose resulted in a 98 ± 1% decrease in eYFP

fluorescence for all sgRNAs in the presence of dCas9. The leakiness of the theophylline-

inducible P$_{trcE*}$ was also apparent when used to regulate dCas9, showing a 27% reduction in

eYFP fluorescence in the absence of theophylline (**Figure 5D**). Induction of dCas9

expression with theophylline resulted in a similar robust decrease in eYFP fluorescence (98 ±

1%). In contrast, cultures transformed with the DAPG-inducible PhlF/P$_{phlF}$ CRISPRi-dCas9

system showed no reductions in eYFP fluorescence in the absence of DAPG, while the

addition of DAPG resulted in an average 97 ± 1% decrease in eYFP fluorescence for cultures

with sgRNAs and dCas9 (**Figure 5E**).

Growth analyses showed that expressing dCas9 under the control of the RhaS/P$_{rhaBAD}$, P$_{trcE*}$

or PhlF/P$_{phlF}$ system with their respective inducer molecules did not impair growth

(**Supplementary Figure S4B-D**). Overall, our results show that the PhlF/P$_{phlF}$ system tightly

regulated dCas9 expression with no measurable leakiness, while induction with DAPG

provided robust transcriptional repression. Notably, similar levels of repression were

observed for all four sgRNAs for each promoter tested. This was unexpected, as previous

work in Vasudevan et al. (2019) showed different levels of repression for the same four

sgRNAs in PCC 6803, and may indicate that PCC 11901 is more amenable to CRISPRi

repression.

We next tested the PhlF/P$_{phlF}$ CRISPRi-dCas9 system on two endogenous gene targets in

PCC 11901 to characterise its capacity for affecting native metabolism. Our gene targets were

cpcB and *nblA*, which are involved in the synthesis and degradation of phycobiliproteins,

respectively, and have been targeted previously in CRISPR-based studies in PCC 7002 and

UTEX 2973 (Gordon et al., 2016; Ungerer and Pakrasi, 2016; Wendt et al., 2016). The *cpc*

operon is involved in the synthesis of the phycocyanin rods of the phycobilisome complex

(**Figure 5F**), an assemblage of proteins that bind to photosystem I or II and increase the

efficiency of light harvesting (Puzorjov and McCormick, 2020; Domínguez-Martín et al.,

2022). We designed an 18 bp sgRNA targeting *cpcB* (**Supplementary Table S3**), the first

gene of the *cpc* operon, and checked to ensure that no off-target sites were present in the

genome of PCC 11901 (Bae et al., 2014). Following induction with DAPG, we found that the

CRISPRi strain expressing dCas9 and the sgRNA targeting *cpcB* had turned a yellow-green

colour after 24 h of induction, similar to the olive colour observed in phycobilisome-deficient

PCC 6803 mutants (**Figure 5G**) (Lea-Smith et al., 2014; Vasudevan et al., 2019), and its

growth was reduced compared to the controls (**Figure 5H**). A significant decrease in the

absorption peak at 625 nm was also observed in phycobilisome extracts from the latter strain

(**Figure 5I**), demonstrating a robust reduction in phycobilisome abundance. SDS-PAGE

analysis of phycobilisome extracts showed a decrease in both *cpc* and *apc* peptides in the

induced (but not uninduced) CRISPRi strain, indicating that CRISPRi repression of *cpcB*

resulted in a reduction in overall phycobilisome abundance in PCC 11901 (**Supplementary**

Figure S5A). The apparent co-repression of allophycocyanin was unexpected, as previous

cpc mutants in PCC 6803 showed no reduction of allophycocyanin (Lea-Smith et al., 2014;

Liberton et al., 2017). However, previous attempts to knockout *cpcB* in PCC 7002 (the

closest relative species to PCC 11901) failed , suggesting that *cpcB* is essential in PCC 7002

(Gordon et al., 2016). Our results indicate that phycocyanin and allophycocyanin may be co-

regulated in PCC 11901, which may also be the case in PCC 7002. Measurement of

chlorophyll content in the CRISPRi strains showed no significant differences

(**Supplementary Figure S5B**), suggesting that the abundance of other components of the

light reactions were not affected by the repression of *cpcB*. Finally, we tested if removal of

DAPG could derepress *cpcB* and restore the phycobilisome pool. Washing the cells in fresh

media to remove DAPG resulted in increased growth after 24 h and restoration of

phycobilisome abundance to wild-type levels after 48 h (**Supplementary Figure S5C** and

S5D). Thus, our results indicate that ~24 h is sufficient for the turnover and degradation of

the dCas9 pool in PCC 11901.

We next targeted the *nblA* gene, which plays a key role in phycobilisome degradation under

nitrogen deficient conditions (**Figure 5J, Supplementary Table S3**) (Collier and Grossman,

1994). *nblA* is a common target to test CRISPR-Cas functionality in cyanobacteria as

disruption of *nblA* leads to an easily detectable phenotype under nitrogen-limiting conditions;

wild-type strains grown in media lacking nitrate exhibit chlorosis or bleaching characteristic

of phycobilisome degradation, while *nblA* mutants show a non-bleaching phenotype and

remain blue-green (Wendt et al., 2016; Baldanta et al., 2022; Cengic et al., 2022). Following

induction with DAPG in nitrogen-depleted medium, we found that the CRISPRi strain

expressing dCas9 and the sgRNA targeting *nblA* remained blue-green, whereas all the

controls turned chlorotic (**Figure 5K**). All cultures grew very slowly due to the lack of

nitrogen (**Figure 5L**), which is not unexpected as nitrogen plays a crucial role in the fast-

growing phenotype of PCC 11901 (Włodarczyk et al., 2020). Chlorophyll content was

reduced overall (**Supplementary Figure S5E**), indicating that all cultures were stressed.

Nevertheless, a strong absorption peak at 625 nm was maintained in phycobilisome extracts

from the functional CRISPRi strain, while all other controls showed a much-reduced peak

indicative of phycobilisome degradation (**Figure 5M**).

Overall, the PhlF/P$_{phlF}$ inducible CRISPRi system shows tight regulation of dCas9 activity

and robust repression of heterologous and native gene expression. Furthermore, we have

shown that removal of the inducer DAPG can reversibly attenuate gene repression and related

physiological effects rapidly in PCC 11901. This system should provide a powerful tool for

studying gene function in PCC 11901 and enable metabolic engineering approaches to exploit

the yield potential of PCC 11901 (Li et al., 2016; Yao et al., 2020; Miao et al., 2023).

Markerless gene editing using CRISPR-Cas12a and a 'double HR' approach

The generation of markerless mutants has key advantages for biotechnology as strains can be
repeatedly genetically manipulated and the absence of genes encoding antibiotic resistance
proteins avoids the possibility of antibiotic resistant organisms being released into the
environment (Lea-Smith et al., 2016). However, recent efforts to use the common negative
selection markers *sacB* and *codA* to generate markerless mutants proved unsuccessful in PCC
11901 (Mills et al., 2022). To overcome this, we attempted to use a CRE-lox recombination
system approach recently demonstrated in PCC 7002 (Jones et al., 2021). This system first
involved generation of marked mutants via insertion of an AbR flanked by two LoxP sites,
lox71 (5'-ATAACTTCGTATAATGTATGCTATACGAACGGTA-3') and lox66 (5'-
TACCGTTCGTATAATGTATGCTATACGAAGTTAT-3') into a target site. A plasmid
vector encoding CRE and a second AbR was then introduced into marked mutants and
integrated into an essential locus (*rbcLXS* or *psbEFJL*) with CRE under control of the native
promoter. As these genes are essential, the strains were maintained in a partially segregated
state under antibiotic selection. Subsequent expression of CRE resulted in excision of the
lox71-AbR-lox66 cassette from the genome. Growth of these mutants on plates lacking the
second antibiotic resulted in loss of chromosomes containing the CRE/AbR insertion and
generation of markerless mutants. We attempted to replicate this system in PCC 11901.
Marked, segregated mutants targeting five different loci (*ctaDIEI*, *ctacII*, *acs*, *ldhA*, *sdhA*)
were successfully generated using a gentamicin resistance cassette (GmR) or KmR flanked
by lox66 and lox71 (**Supplementary Figure S6)**. We then generated two plasmids that
allowed recombination of CRE and SpR into the *rbcLXS* or *psbEFJL* locus. However, despite
repeated attempts to transform these plasmids into each of the five markerless mutants, we
were unable to obtain spectinomycin resistant colonies. Moreover, no colonies were obtained
when the plasmids were introduced into wild-type cells. Our inability to generate markerless
mutants in PCC 11901 using this approach suggests that CRE is either toxic or that the
generation of partially segregated mutants in key essential genes is extremely challenging.
495
In conjunction, alternate CRISPR-Cas gene editing approaches for generating markerless
mutants were trialled. These allow for more efficient engineering in cyanobacteria without
the requirement for positive selection markers. As the double stranded breaks generated by
Cas proteins are lethal to most microbes (including cyanobacteria) due to their general lack of
a non-homologous end joining pathway (Su et al., 2016), selection is based on the uptake of
supplied repair template sequences to replace the sgRNA target locus by HR (Behler et al.,
2018). Due to the documented cytotoxicity of Cas9 in cyanobacteria (Li et al., 2016;

Racharaks et al., 2021), much CRISPR-Cas work to date has focused on reportedly non-toxic

Cas12a (*Fn*Cpf1), which has led to several successful examples of gene editing in a variety of

cyanobacteria (Lin et al., 2021; Baldanta et al., 2022). Recently, an RNA riboswitch-

inducible dCas9 system was demonstrated in PCC 6803 that could overcome the toxicity of

Cas9 (Cengic et al., 2022).

Here, we sought to build a novel CyanoGate-compatible inducible CRISPR-Cas12a system

that allows for the generation of markerless mutants and multiplex editing at high

efficiencies. Our system is comprised of two vectors i) an RSF1010-based (pPMQSK1-1)

self-replicating 'editing vector' carrying Cas12a (pC1.509, **Supplementary Table S1**), and

ii) a pUC19-T 'hybrid suicide vector' designed to deliver an sgRNA expression cassette into

the editing vector for stable expression and a repair template with homology to the sgRNA

target locus on the genome by two separate HR events, which we called the 'double HR'

approach (**Figure 6A**). We chose the DAPG-inducible PhlF/P$_{phlF}$ promoter system to drive

expression of Cas12a, which we have previously shown to be tightly regulated in PCC 11901

(**Figure 4** and **Figure 5**), and conjugated the editing vector into PCC 11901 to generate an

'editing strain'. To assist with assembling the hybrid suicide vector, we constructed the

CyanoGate-compatible sgRNA assembly acceptor vector pCA0.421 based on the CRATES

system from Liao et al., (2019), which enables one-pot assembly of single or multiplexed

sgRNA arrays for targeted gene editing (**Supplementary Table S1**). We hypothesised that

with this approach new hybrid suicide vectors carrying one or more sgRNAs and up to six

repair templates could be rapidly generated and iteratively transformed into the editing strain

to generate markerless mutants with single or multiple chromosomal alterations.

As a proof-of-concept, we used the editing strain (carrying the editing vector with SpR) to

perform a markerless knock-in of an eYFP expression cassette into the *mrr* neutral site

(**Figure 6B**). A hybrid suicide vector pCT.590 (**Supplementary Table S2**) was assembled

and transformed into PCC 11901 in the absence or presence of DAPG, and the transformation

mix was then plated onto gentamicin-supplemented agar or gentamicin- and DAPG-

supplemented agar, respectively. In the absence of DAPG, no edits were observed in the

colonies that grew, indicating that Cas12a expression was tightly repressed. However, with

DAPG we observed that 25% of colonies showed insertions of the eYFP expression cassette

and eYFP fluorescence. This demonstrated that in the Δ*mrr*::eYFP colonies i) HR had

occurred between the hybrid vector and the editing vector to allow for growth on gentamicin,

and ii) HR had occurred between the hybrid vector and the genome. In the remaining 75% of

colonies, only i) had occurred, which indicated that HR with the editing vector was more

prominent than with the genome. We confirmed that double HR had occurred with colonies

expressing eYFP by PCR of the editing vector, which showed the presence of the sgRNA

cassette and that SpR had been replaced by GmR (**Supplementary Figure S9**). Notably, the

absence of a WT band in the Δ*mrr*::eYFP colonies indicated that these strains were fully

segregated.

To further improve the efficiency of editing, we performed transformation in the absence of

DAPG for 4 h, then added DAPG for 12 h before plating onto gentamicin- and DAPG-

supplemented agar to induce Cas12a expression. We observed a significant increase in the

efficiency of double HR (81%) (**Figure 6B**), which demonstrated the benefit of increasing the

time period for genomic HR to occur prior to the induction of Cas12a expression. This

approach was used in all subsequent gene editing experiments. We next assembled the hybrid

suicide vector pCT.621 to test the *aquI* neutral site as a second target locus for insertion of an

mCherry expression cassette (**Supplementary Table S2**). Here we observed a 56%

efficiency in generating fully segregated Δ*aquI*::mCherry lines (**Figure 6C**), suggesting that

the double HR approach was still robust at different genomic loci.

We next investigated if we could generate the double insertion mutant Δ*mrr*::eYFP/

Δ*aquI*::mCherry through iterative or multiplex genome editing. For the former, we assembled

a hybrid suicide vector pCT.622 for transformation into Δ*mrr*::eYFP, which carries an editing

vector with GmR (**Supplementary Table S2**). Following transformation of Δ*mrr*::eYFP, we

observed mCherry expression cassette insertions in *aquI* in 31% of colonies (**Figure 6D**).

Fluorescence measurements confirmed expression of both eYFP and mCherry, and thus

successful generation of Δ*mrr*::eYFP/ Δ*aquI*::mCherry double mutants through iterative gene

editing. For the latter multiplex approach, we assembled the suicide vector pCT.623 carrying

an sgRNA array and homology repair templates designed to insert expression cassettes for

eYFP and mCherry into *mrr* and *aquI*, respectively. Following transformation of the editing

strain, we screened colonies for single and double insertion mutants by PCR. We observed

insertion efficiencies of 56% and 25% for the eYFP and mCherry expression cassettes,

respectively, while 25% of the colonies contained both cassettes at their expected loci

(**Figure 6E**). Notably, we did not find examples of an mCherry insertion in the absence of

eYFP, suggesting that the first sgRNA in the array targeting the *mrr* locus may have been

more efficient or is more abundantly expressed.

Overall, our results support PCC 11901 as a highly amenable strain for CRISPR-Cas12a-

based editing. To our knowledge, this is the first report of iterative CRISPR-Cas12a gene

editing and multiplex editing using an sgRNA array in a cyanobacterial strain. The iterative

editing scheme was designed to allow as many sites to be targeted as desired, by cycling

through SpR and GmR cassettes in the hybrid suicide vector, enabling more complex editing

schemes to occur. The two editing schemes (iterative and multiplex) could also be further

combined, allowing further flexibility in strain design. Additional efficiency improvements

could be made by establishing the mechanism by which 'escaper colonies' survive on

selective media but avoid editing, a common phenomenon observed in bacteria and

cyanobacteria (Vento et al., 2019; Cengic et al., 2022). As we were able to achieve iterative

gene editing, escape was not due to mutations in Cas12a and thus may have been due to point

mutations in the sgRNA or the sgRNA genomic target site. A possible strategy to improve the

efficiency of edits could be to reduce the expression levels of Cas12a and/or the sgRNA(s),

an approach that that has been shown to increase editing efficiencies in *E. coli* and *Klebsiella*

spp. (Collias et al., 2023).

Finally, we sought to cure the editing strain of the self-replicating editing vector. Various

methods have been employed to cure plasmids from bacteria, with the standard approach

being repeated subculture of mutants in antibiotic-free medium and screening for

spontaneous plasmid loss (Bishé et al., 2019). However, RSF1010-based vectors appear to

persist in cyanobacteria for long periods, even in the absence of antibiotic selection (Nagy et

al., 2021; Puzorjov et al., 2022). Previous CRISPR-Cas work in *E. coli* and *P. putida* utilised

a 'self-targeting' sgRNA for efficient removal of RSF1010-based vectors (Lauritsen et al.,

2017). Here, we generated a hybrid suicide vector containing an sgRNA that targeted the

editing vector pC1.509. We transformed cells in the presence of DAPG and plated the

transformation culture onto agar without antibiotics. We then screened colonies for the

absence of the editing vector by PCR and observed a curing efficiency of 50% (**Figure 6F**).

We further verified the loss of the editing vector by patching the cured colonies onto

gentamicin-supplemented agar, which resulted in no growth indicating sensitivity to the

antibiotic. Thus, we demonstrated that PCC 11901 mutants generated by CRISPR-Cas12a

can be cured of the editing vector to produce fully markerless mutant strains containing no
scars or AbR cassettes.

CONCLUSION

Here we have investigated the amenability of the fast-growing marine cyanobacterium PCC
11901 to engineering and assembled a comprehensive suite of tools compatible with the
CyanoGate MoClo platform to enable future work in this strain. We hope that the tools
described will accelerate the wider adoption of this next-generation cyanobacterial chassis
strain. The fast-growing and highly productive phenotype of PCC 11901 offers much in terms
of advancing our fundamental understanding of the genetic basis and regulation of these
processes (Ungerer et al., 2018). Furthermore, PCC 11901 shows promise for applied work
aiming to develop commercially viable green biotechnology chassis for renewable
biomanufacture and biomaterials production (Goodchild-Michelman et al., 2023),
sequestration of CO_2 emissions in hard-to-abate sectors (e.g. capture of CO_2 from point-
source flue gases) (Zhang et al., 2017), and sustainable space exploration (Santomartino et
al., 2023).

MATERIALS AND METHODS

Cyanobacterial culture conditions

Synechococcus sp. PCC 11901 was cultured in AD7 or MAD liquid medium (Włodarczyk et
al., 2020), or on 1.5% (w/v) agar plates as described in (Włodarczyk et al., 2020). Cultures
were grown in an Algaetron AG 230 incubator (Photon Systems Instruments) at 30°C, 2%
(v/v) CO_2 under continuous warm white LED light (150 µmol photons m^{-2} s^{-1}) and shaking at
120 rpm. Agar plates were incubated under identical conditions, without shaking.

Plasmid vector assembly

Level 0, 1, and T plasmid vectors were assembled using the CyanoGate MoClo kit
(Vasudevan et al., 2019). Native PCC 11901 genetic parts were amplified from genomic
DNA using Q5 High-Fidelity DNA Polymerase (New England Biolabs). Where necessary,
native genetic parts were domesticated (i.e. sites for Type IIS restriction endonucleases BsaI
and BpiI were removed) using specific primers. Alternatively, parts were synthesized as
Gblocks DNA fragments (Integrated DNA Technologies) and cloned directly into an
appropriate level 0 acceptor (Engler et al., 2014) (see **Supplementary Table S1** and
Supplementary Data S1 for new vectors, and **Supplementary Table S2** for all plasmids

assembled in this study). Vectors were transformed into One Shot TOP10 chemically
competent *Escherichia coli* (Thermo Fisher Scientific) cells as per the manufacturer's
instructions. Transformed cultures were grown at 37°C on 1.5% (w/v) LB agar or in liquid
LB medium shaking at 125 rpm with appropriate antibiotic selection.

sgRNA selection and CRISPR-Cas assemblies
sgRNAs were designed by selecting 18-22 bp sequences adjacent to the protospacer adjacent
motif (PAM) sequence 5'-NGG-3' for *Streptococcus pyogenes* dCas9 or 5'-TTTV-3' for
Francisella novicida Cas12a. Candidate sgRNAs were checked for potential off-target sites
in the PCC 11901 genome using Cas-OFFinder (Bae et al., 2014). The sgRNAs for dCas9
were made by annealing complementary oligonucleotides carrying the required overhangs
and BsaI recognition sites (**Supplementary Table S3**), and were assembled into the level 1
position 2 (L1P2) acceptor vector pICH47742 together with the P$_{trc10\_TSS}$ promoter (pC0.220)
and the sgRNA scaffold (pC0.122) as described in Vasudevan et al., (2019). The sgRNAs for
Cas12a were also made by annealing complementary oligonucleotides carrying overhangs
and BpiI recognition sites as described in Liao et al. (2019) (see **Supplementary Figure S3**
and **S7**) and were assembled into the new acceptor vector pC0.421 (**Supplementary Table**
S1).

Natural transformation of PCC 11901
Purified plasmid (1 μg) was added to 1 mL of wild-type PCC 11901 culture at exponential
growth phase (OD$_{750}$ = 0.8) and incubated for 12 h at 30°C under continuous warm white
LED light (150 μmol photons m^{-2} s^{-1}) and shaking at 120 rpm in an Infors Multitron Pro
incubator (Infors HT). The cultures were then plated onto AD7 agar plates supplemented with
appropriate antibiotics (25 μg mL^{-1} spectinomycin, 50 μg mL^{-1} gentamicin, 100 μg mL^{-1}
carbenicillin or 100 μg mL^{-1} kanamycin). Plates were then sealed with 3M Micropore® tape
to allow for gas exchange and incubated at 30°C, 2% (v/v) CO$_2$ under warm white LED light
(150 μmol photons m^{-2} s^{-1}). Colonies typically appeared after 2-4 days.

Transconjugation of PCC 11901
Genetic modification by conjugal transfer was performed using an approach adapted from
Gale et al. (2019). Overnight cultures of *E. coli* strain HB101 harbouring vectors pRK2013
(ATCC 37159) and pRL528 (the helper strain) and a TOP10 strain harbouring an RSF1010-
derived level T vector were each washed three times with LB medium to remove antibiotics.

The *E. coli* cultures were then combined (450 μL each) and incubated for 1 h at room

temperature (RT). PCC 11901 cultures were grown to OD_{750} ~1.0 and washed three times

with fresh AD7 medium. The combined *E. coli* culture was added to 900 μL of PCC 11901

culture and the mixture incubated at 30°C under warm white LED light (150 μmol photons

m^{-2} s^{-1}) for 4 h, without shaking. The mix was centrifuged at 4,000 *g* and the cell pellet was

plated onto 0.45 μm Metricel® membrane filter discs (Pall Corporation) laid on top of non-

selective AD7 agar. After 24 h of incubation, the membranes were transferred to AD7 agar

supplemented with appropriate antibiotics (25 μg mL^{-1} spectinomycin, or 100 μg mL^{-1}

kanamycin) and incubated as above. Conjugant colonies typically appeared six days post

membrane transfer.

Comparative growth assays

Growth curve experiments were performed by inoculating a seed culture containing 30 mL of

MAD medium with single colonies of cyanobacteria picked from agar plates and grown as

described above to OD_{750} ~1.0. The seed cultures were then used to prepare triplicate 15 mL

starter cultures adjusted to OD_{750} ~0.2 and aliquoted into Corning® 25 cm^2 cell culture flasks

with canted necks and vented caps (Corning). To facilitate gas exchange and prevent

foaming, 0.5 μL of Antifoam 204 (Sigma Aldrich) was added to the cultures. Cultures were

grown at 30°C, 2% (v/v) CO_2, shaking at 150 rpm under 150 μmol photons m^{-2} s^{-1} for the

first 24 h, which was then increased to 750 μmol photons m^{-2} s^{-1}. Optical density was

measured every 24 h using a WPA Biowave II UV-Vis spectrophotometer (Biochrom) for

eight days.

eYFP quantifications

Mutant PCC 11901 strains were grown in 6-well culture plates (Starlab CytoOne) and

incubated in an Algaetron® AG 230 incubator under the same culturing conditions as

described above. OD_{750} and eYFP fluorescence of cultures were measured using a FLUOstar

OMEGA microplate reader (BMG Labtech). Fluorescence of eYFP for individual cells

(10,000 cells per culture) was measured by flow cytometry using a BD® LSR II Fortessa

flow cytometer (Becton Dickinson). Cells were gated using forward and side scatter, and

median eYFP fluorescence was calculated from excitation/emission wavelengths 488

nm/515–545 nm (Kelly et al., 2018), and reported after 24 h of growth unless otherwise

stated.

Terminator efficiency calculations

The efficiency of terminator sequences (terminator efficiency (TE)) was calculated by

assembling terminator sequences into the pDUOTK1-L1 vector as described in Gale et al.,

(2021).

Measurement of chlorophyll content

Cultures were diluted to $OD_{750} = 1.0$ and centrifuged at 17,000 g for 2 min. The resulting

pellet was resuspended in 100% (v/v) methanol and shaken at 2400 rpm for 1 h in the dark at

RT using an IKA-VIBRAX-VXR bead beater. The homogenates were then centrifuged at

17,000 g for 10 min and the absorbance of the supernatant was measured at 652, 665 and

750 nm. The mean concentration of chlorophyll a was calculated from triplicates as described

in Porra et al., (1989).

Extraction of phycobiliproteins and analysis

Phycobiliproteins (PBS) were extracted and quantified using absorbance spectroscopy as

described previously (Zavřel et al., 2018). Briefly, PCC 11901 cells were pelleted by

centrifugation at 15,000 g for 5 min, washed in phosphate buffered saline three times and

freeze-dried overnight. Dried samples were lysed with 0.5 mm glass beads (BioSpec

Products) on a TissueLyser II homogeniser (Qiagen) for 15 s at RT. 1 mL of pre-cooled

phosphate buffered saline at 4°C was added to each tube and mixed for 5 s on the

homogeniser. Samples were then incubated on ice for 60 min to efficiently extract soluble

proteins and prevent protein degradation. Following centrifugation for 5 min at 4°C and

15,000 g, the aqueous blue liquid layer containing PBS was transferred to sterile 1.5 mL

microcentrifuge tubes and frozen for future use. Samples were measured from 550-750 nm on

a Biochrom WPA Biowave II Spectrophotometer. For SDS-PAGE analysis, samples were

run on a Bolt 12% Bis-Tris Plus Mini protein gel (Invitrogen) at 150 V for 1 h. A pre-stained

protein standard (Proteintech) was used as a ladder. The gels were then stained with 1% (w/v)

Coomassie Brilliant Blue solution (Bio-Rad) and destained with a methanol: acetic acid:

distilled water (50%: 10%: 40%) solution.

ACKNOWLEDGEMENT

We would like to thank the Frankenberg-Dinkel Lab (RPTU Kaiserslautern-Landau) for

kindly providing their facilities for several experiments while hosting A.J.V.

DATA AVAILABILITY

Plasmids in **Supplementary Table S1** are available from Addgene (https://www.addgene.org/Alistair_McCormick), Addgene IDs: 203934-203943, 205441.

FUNDING

A.J.V. was funded by a postgraduate research scholarship from the Darwin Trust of Edinburgh. T.T.S. acknowledges funding from the Green Chemicals Beacon of Excellence, University of Nottingham. J.A.M.C. acknowledges funding support from a FEBS short term fellowship and University of Cordoba fellowship. L.A.M. acknowledges funding support from the BBSRC Norwich Research Park Doctoral Training Partnership program (grant number BB/S507404/1). D.J. and L.-S. acknowledge funding from BBSRC grant [BB/S020365/1]. A.J.M acknowledges funding from the UK Biotechnology and Biological Sciences Research Council (BBSRC) grants [BB/S020128/1] and [BB/W003538/1].

COMPETING INTERESTS

The authors declare that no competing interests exist.

SUPPLEMENTARY DATA

Supplementary Table S1. Table of all new CyanoGate-compatible parts generated in this work.

Supplementary Table S2. All plasmids made and used in this study. See 'Supplementary Table S2 - Plasmids (all).xlsx'.

Supplementary Table S3. Primer and sgRNA oligonucleotides used in this study. See 'Supplementary Table S3 - Primer and sgRNA oligonucleotides.xlsx'.

Supplementary Data S1. Sequence maps (.gb files) for plasmid vectors in Supplementary Table S1. See 'Supplementary Data S1.zip'.

Supplementary Figure S1. Analysis of putative Gcn5-related N-acetyltransferase (GNAT) family genes in PCC 11901.

Supplementary Figure S2. Representative PCR-based segregation analysis of PCC 11901 transformants targeting neutral sites *glgA1* and *aquI*.

Supplementary Figure S3. Growth of PCC 11901 with varying doses of small molecule inducers.

Supplementary Figure S4. Growth analysis of CRISPRi-dCas9 strains targeting eYFP.

Supplementary Figure S5. Analysis of CRISPRi-dCAs9 strains targeting *cpcB* and *nblA*.

Supplementary Figure S6. Attempted generation of markerless mutants using the CRE-Lox

system.

Supplementary Figure S7. CRISPR-Cas12a double HR editing approach – sgRNA, repair

template and hybrid suicide vector assembly.

Supplementary Figure S8. Pipeline for iterative CRISPR-Cas12a editing using the double

HR approach.

Supplementary Figure S9. Confirmation of recombination between the hybrid suicide

vector and editing vector.

REFERENCES

Abbas A, Morrissey JP, Marquez PC, Sheehan MM, Delany IR, O'Gara F (2002)
Characterization of interactions between the transcriptional repressor PhlF and its
binding site at the phlA promoter in Pseudomonas fluorescens F113. J Bacteriol **184**:
3008–3016

Bae S, Park J, Kim J-S (2014) Cas-OFFinder: a fast and versatile algorithm that searches
for potential off-target sites of Cas9 RNA-guided endonucleases. Bioinformatics **30**:
1473–1475

Baldanta S, Guevara G, Navarro-Llorens JM (2022) SEVA-Cpf1, a CRISPR-Cas12a
vector for genome editing in cyanobacteria. Microb Cell Factories **21**: 103

Batterton JC, Van Baalen C (1971) Growth responses of blue-green algae to sodium
chloride concentration. Archiv Mikrobiol **76**: 151–165

Behle A, Saake P, Germann AT, Dienst D, Axmann IM (2020) Comparative dose–
response analysis of inducible promoters in cyanobacteria. ACS Synth Biol **9**: 843–855

Behler J, Vijay D, Hess WR, Akhtar MK (2018) CRISPR-based technologies for metabolic
engineering in cyanobacteria. Trends Biotechnol **36**: 996–1010

BEIS (2021) 'Net Zero Strategy: Build Back Greener'. Department for Business, Energy, and
International Strategy. Available at: https://www.gov.uk/government/publications/net-
zero-strategy (Accessed: 13 September 2022).

Bernhards CB, Liem AT, Berk KL, Roth PA, Gibbons HS, Lux MW (2022) Putative
phenotypically neutral genomic insertion points in prokaryotes. ACS Synth Biol **11**:
1681–1685

Bishé B, Taton A, Golden JW (2019) Modification of RSF1010-based broad-host-range
plasmids for improved conjugation and cyanobacterial bioprospecting. iScience **20**:
216–228

Cameron JC, Pakrasi HB (2011) Glutathione facilitates antibiotic resistance and
photosystem I stability during exposure to gentamicin in cyanobacteria. Appl Environ
Microbiol **77**: 3547–3550

Cengic I, Cañadas IC, Minton NP, Hudson EP (2022) Inducible CRISPR/Cas9 allows for
multiplexed and rapidly segregated single-target genome editing in Synechocystis sp.
PCC 6803. ACS Synth Biol **11**: 3100–3113

Chen Y-J, Liu P, Nielsen AAK, Brophy JAN, Clancy K, Peterson T, Voigt CA (2013)
Characterization of 582 natural and synthetic terminators and quantification of their
design constraints. Nat Methods **10**: 659–664

Chi X, Zhang S, Sun H, Duan Y, Qiao C, Luan G, Lu X (2019) Adopting a theophylline-
responsive riboswitch for flexible regulation and understanding of glycogen metabolism
in synechococcus elongatus PCC 7942. Front Microbiol 10:

Cho BA, Moreno-Cabezuelo JÁ, Mills LA, del Río Chanona EA, Lea-Smith DJ, Zhang
D (2023) Integrated experimental and photo-mechanistic modelling of biomass and
optical density production of fast versus slow growing model cyanobacteria. Algal Res
70: 102997

Collias D, Vialetto E, Yu J, Co K, Almási É d H, Rüttiger A-S, Achmedov T, Strowig T,
Beisel CL (2023) Systematically attenuating DNA targeting enables CRISPR-driven
editing in bacteria. Nat Commun **14**: 680

Collier JL, Grossman AR (1994) A small polypeptide triggers complete degradation of
light-harvesting phycobiliproteins in nutrient-deprived cyanobacteria. EMBO J **13**:
1039–1047

Dallo T, Krishnakumar R, Kolker SD, Ruffing AM (2023) High-Density Guide RNA
Tiling and Machine Learning for Designing CRISPR Interference in Synechococcus sp.
PCC 7002. ACS Synth Biol. doi: 10.1021/acssynbio.2c00653

Daneshvar E, Wicker RJ, Show P-L, Bhatnagar A (2022) Biologically-mediated carbon
capture and utilization by microalgae towards sustainable CO_2 biofixation and biomass
valorization – A review. Chem Eng J **427**: 130884

Depardieu F, Bikard D (2020) Gene silencing with CRISPRi in bacteria and optimization of
dCas9 expression levels. Methods **172**: 61–75

Dias E, Oliveira M, Jones-Dias D, Vasconcelos V, Ferreira E, Manageiro V, Caniça M
(2015) Assessing the antibiotic susceptibility of freshwater Cyanobacteria spp. Front
Microbiol **6**: 799

Domínguez-Martín MA, Sauer PV, Kirst H, Sutter M, Bína D, Greber BJ, Nogales E,
Polívka T, Kerfeld CA (2022) Structures of a phycobilisome in light-harvesting and
photoprotected states. Nature **609**: 835–845

Engler C, Youles M, Gruetzner R, Ehnert T-M, Werner S, Jones JDG, Patron NJ,
Marillonnet S (2014) A golden gate modular cloning toolbox for plants. ACS Synth
Biol **3**: 839–843

Favrot L, Blanchard JS, Vergnolle O (2016) Bacterial GCN5-Related N-
Acetyltransferases: From Resistance to Regulation. Biochemistry **55**: 989–1002

Fontana J, Dong C, Ham JY, Zalatan JG, Carothers JM (2018) Regulated expression of
sgRNAs tunes CRISPRi in E. coli. Biotechnology Journal **13**: 1800069

Gale GAR, Schiavon Osorio AA, Mills LA, Wang B, Lea-Smith DJ, McCormick AJ
(2019) Emerging species and genome editing tools: future prospects in cyanobacterial
synthetic biology. Microorganisms **7**: 409

Gale GAR, Wang B, McCormick AJ (2021) Evaluation and comparison of the efficiency of
transcription terminators in different cyanobacterial species. Front Microbiol 11:

Goodchild-Michelman IM, Church GM, Schubert MG, Tang T-C (2023) Light and
carbon: Synthetic biology toward new cyanobacteria-based living biomaterials. Mater
Today Bio **19**: 100583

Gordon GC, Korosh TC, Cameron JC, Markley AL, Begemann MB, Pfleger BF (2016)
CRISPR interference as a titratable, trans-acting regulatory tool for metabolic
engineering in the cyanobacterium Synechococcus sp. strain PCC 7002. Metab Eng **38**:
170–179

Hitchcock A, Hunter CN, Canniffe DP (2020) Progress and challenges in engineering
cyanobacteria as chassis for light-driven biotechnology. Microb Biotechnol **13**: 363–367

Huang H-H, Lindblad P (2013) Wide-dynamic-range promoters engineered for
cyanobacteria. J Biol Eng **7**: 10

Iwai M, Katoh H, Katayama M, Ikeuchi M (2004) Improved genetic transformation of the
thermophilic cyanobacterium, thermosynechococcus elongatus BP-1. Plant Cell Physiol
45: 171–175

Jaiswal D, Sengupta A, Sengupta S, Madhu S, Pakrasi HB, Wangikar PP (2020) A novel
cyanobacterium Synechococcus elongatus PCC11802 has distinct genomic and
metabolomic characteristics compared to its neighbor PCC 11801. Sci Rep **10**: 191

Jaiswal D, Sengupta A, Sohoni S, Sengupta S, Phadnavis AG, Pakrasi HB, Wangikar
PP (2018) Genome features and biochemical characteristics of a robust, fast growing
and naturally transformable cyanobacterium Synechococcus elongatus PCC11801
isolated from India. Sci Rep **8**: 16632

Jester BW, Zhao H, Gewe M, Adame T, Perruzza L, Bolick DT, Agosti J, Khuong N,
Kuestner R, Gamble C, et al (2022) Development of spirulina for the manufacture and
oral delivery of protein therapeutics. Nat Biotechnol **40**: 956–964

Jones CM, Parrish S, Nielsen DR (2021) Exploiting polyploidy for markerless and plasmid-
free genome engineering in cyanobacteria. ACS Synth Biol **10**: 2371–2382

Kavita K, Breaker RR (2023) Discovering riboswitches: the past and the future. Trends
Biochem Sci **48**: 119–141

Kelly CL, Taylor GM, Hitchcock A, Torres-Méndez A, Heap JT (2018) A rhamnose-
inducible system for precise and temporal control of gene expression in cyanobacteria.
ACS Synth Biol **7**: 1056–1066

Khetkorn W, Incharoensakdi A, Lindblad P, Jantaro S (2016) Enhancement of poly-3-
hydroxybutyrate production in Synechocystis sp. PCC 6803 by overexpression of its
native biosynthetic genes. Bioresour Technol **214**: 761–768

Kopka J, Schmidt S, Dethloff F, Pade N, Berendt S, Schottkowski M, Martin N,
Dühring U, Kuchmina E, Enke H, et al (2017) Systems analysis of ethanol production
in the genetically engineered cyanobacterium Synechococcus sp. PCC 7002. Biotechnol
Biofuels **10**: 56

Kufryk GI, Sachet M, Schmetterer G, Vermaas WFJ (2002) Transformation of the
cyanobacterium Synechocystis sp. PCC 6803 as a tool for genetic mapping: optimization
of efficiency. FEMS Microbiol Lett **206**: 215–219

Lauritsen I, Porse A, Sommer MOA, Nørholm MHH (2017) A versatile one-step
CRISPR-Cas9 based approach to plasmid-curing. Microb Cell Factories **16**: 135

Lea-Smith DJ, Bombelli P, Dennis JS, Scott SA, Smith AG, Howe CJ (2014)
Phycobilisome-deficient strains of Synechocystis sp. PCC 6803 have reduced size and
require carbon-limiting conditions to exhibit enhanced productivity. Plant Physiol **165**:
705–714

Lea-Smith DJ, Summerfield TC, Ducat DC, Lu X, McCormick AJ, Purton S (2021)
Editorial: Exploring the growing role of cyanobacteria in industrial biotechnology and
sustainability. Front Microbiol 12:

Lea-Smith DJ, Vasudevan R, Howe CJ (2016) Generation of marked and markerless
mutants in model cyanobacterial species. J Vis Exp 54001

Li H, Shen CR, Huang C-H, Sung L-Y, Wu M-Y, Hu Y-C (2016) CRISPR-Cas9 for the
genome engineering of cyanobacteria and succinate production. Metab Eng **38**: 293–302

Li S, Sun T, Xu C, Chen L, Zhang W (2018) Development and optimization of genetic
toolboxes for a fast-growing cyanobacterium Synechococcus elongatus UTEX 2973.
Metab Eng **48**: 163–174

Liberton M, Chrisler WB, Nicora CD, Moore RJ, Smith RD, Koppenaal DW, Pakrasi
HB, Jacobs JM (2017) Phycobilisome truncation causes widespread proteome changes
in Synechocystis sp. PCC 6803. PLoS One **12**: e0173251

Lin P-C, Zhang F, Pakrasi HB (2021) Enhanced limonene production in a fast-growing cyanobacterium through combinatorial metabolic engineering. Metab Eng Commun **12**: e00164

Lindberg P, Park S, Melis A (2010) Engineering a platform for photosynthetic isoprene production in cyanobacteria, using Synechocystis as the model organism. Metab Eng **12**: 70–79

Liu D, Johnson VM, Pakrasi HB (2020) A reversibly induced CRISPRi system targeting photosystem II in the cyanobacterium Synechocystis sp. PCC 6803. ACS Synth Biol **9**: 1441–1449

Liu D, Pakrasi HB (2018) Exploring native genetic elements as plug-in tools for synthetic biology in the cyanobacterium Synechocystis sp. PCC 6803. Microb Cell Factories **17**: 48

Ma AT, Schmidt CM, Golden JW (2014) Regulation of gene expression in diverse cyanobacterial species by using theophylline-responsive riboswitches. Appl Environ Microbiol **80**: 6704–6713

Meyer AJ, Segall-Shapiro TH, Glassey E, Zhang J, Voigt CA (2019) Escherichia coli "Marionette" strains with 12 highly optimized small-molecule sensors. Nat Chem Biol **15**: 196–204

Meyer R (2009) Replication and conjugative mobilization of broad host-range IncQ plasmids. Plasmid **62**: 57–70

Miao R, Jahn M, Shabestary K, Hudson EP (2023) CRISPR interference screens reveal tradeoffs between growth rate and robustness in Synechocystis sp. PCC 6803 across trophic conditions. 2023.02.13.528328

Mills LA, Moreno-Cabezuelo JÁ, Włodarczyk A, Victoria AJ, Mejías R, Nenninger A, Moxon S, Bombelli P, Selão TT, McCormick AJ, et al (2022) Development of a biotechnology platform for the fast-growing cyanobacterium Synechococcus sp. PCC 11901. Biomolecules **12**: 872

Mittermair S, Lakatos G, Nicoletti C, Ranglová K, Manoel JC, Grivalský T, Kozhan DM, Masojídek J, Richter J (2021) Impact of glgA1, glgA2 or glgC overexpression on growth and glycogen production in Synechocystis sp. PCC 6803. J Biotechnol **340**: 47–56

Nakahira Y, Ogawa A, Asano H, Oyama T, Tozawa Y (2013) Theophylline-dependent riboswitch as a novel genetic tool for strict regulation of protein expression in cyanobacterium synechococcus elongatus PCC 7942. Plant Cell Physiol **54**: 1724–1735

Ng AH, Berla BM, Pakrasi HB (2015) Fine-tuning of photoautotrophic protein production by combining promoters and neutral sites in the cyanobacterium Synechocystis sp. strain PCC 6803. Appl Environ Microbiol **81**: 6857–6863

Opel F, Siebert NA, Klatt S, Tüllinghoff A, Hantke JG, Toepel J, Bühler B, Nürnberg DJ, Klähn S (2022) Generation of synthetic shuttle vectors enabling modular genetic engineering of cyanobacteria. ACS Synth Biol **11**: 1758–1771

Pinto F, Pacheco CC, Oliveira P, Montagud A, Landels A, Couto N, Wright PC, Urchueguía JF, Tamagnini P (2015) Improving a Synechocystis-based photoautotrophic chassis through systematic genome mapping and validation of neutral sites. DNA Res **22**: 425–437

Porra RJ, Thompson WA, Kriedemann PE (1989) Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. Biochim Biophys Acta - Bioenerg **975**: 384–394

Puzorjov A, Dunn KE, McCormick AJ (2021) Production of thermostable phycocyanin in a mesophilic cyanobacterium. Metab Eng Commun **13**: e00175

Puzorjov A, McCormick AJ (2020) Phycobiliproteins from extreme environments and their potential applications. J Exp Bot **71**: 3827–3842

Puzorjov A, Mert Unal S, Wear MA, McCormick AJ (2022) Pilot scale production, extraction and purification of a thermostable phycocyanin from Synechocystis sp. PCC 6803. Bioresour Technol **345**: 126459

Racharaks R, Arnold W, Peccia J (2021) Development of CRISPR-Cas9 knock-in tools for free fatty acid production using the fast-growing cyanobacterial strain Synechococcus elongatus UTEX 2973. J Microbiol Methods **189**: 106315

Rautela A, Kumar S (2022) Engineering plant family TPS into cyanobacterial host for terpenoids production. Plant Cell Rep **41**: 1791–1803

Ruffing AM, Jensen TJ, Strickland LM (2016) Genetic tools for advancement of Synechococcus sp. PCC 7002 as a cyanobacterial chassis. Microb Cell Factories **15**: 190

Sakamoto T, Shen G, Higashi S, Murata N, Bryant DA (1997) Alteration of low-temperature susceptibility of the cyanobacterium Synechococcus sp. PCC 7002 by genetic manipulation of membrane lipid unsaturation. Arch Microbiol **169**: 20–28

Sakurai I, Stazic D, Eisenhut M, Vuorio E, Steglich C, Hess WR, Aro E-M (2012) Positive regulation of psbA gene expression by cis-encoded antisense RNAs in Synechocystis sp. PCC 6803. Plant Physiol **160**: 1000–1010

Santomartino R, Averesch NJH, Bhuiyan M, Cockell CS, Colangelo J, Gumulya Y, Lehner B, Lopez-Ayala I, McMahon S, Mohanty A, et al (2023) Toward sustainable space exploration: a roadmap for harnessing the power of microorganisms. Nat Commun **14**: 1391

Saveria T, Parthiban C, Seilie AM, Brady C, Martinez A, Manocha R, Afreen E, Zhao H, Krzeszowski A, Ferrara J, et al (2022) Needle-free, spirulina-produced Plasmodium falciparum circumsporozoite vaccination provides sterile protection against pre-erythrocytic malaria in mice. NPJ Vaccines **7**: 1–11

Selão TT (2022) Exploring cyanobacterial diversity for sustainable biotechnology. J Exp Bot **73**: 3057–3071

Sengupta A, Pritam P, Jaiswal D, Bandyopadhyay A, Pakrasi HB, Wangikar PP (2020a) Photosynthetic co-production of succinate and ethylene in a fast-growing cyanobacterium, Synechococcus elongatus PCC 11801. Metabolites **10**: 250

Sengupta S, Jaiswal D, Sengupta A, Shah S, Gadagkar S, Wangikar PP (2020b) Metabolic engineering of a fast-growing cyanobacterium Synechococcus elongatus PCC 11801 for photoautotrophic production of succinic acid. Biotechnol Biofuels **13**: 89

Shiraishi H, Nishida H (2022) Complete genome sequence of the edible filamentous cyanobacterium arthrospira platensis nies-39, based on long-read sequencing. Microbiol Resour Announc **12**: e01139-22

Soini J, Ukkonen K, Neubauer P (2008) High cell density media for Escherichia coli are generally designed for aerobic cultivations – consequences for large-scale bioprocesses and shake flask cultures. Microb Cell Factories **7**: 26

Su T, Liu F, Gu P, Jin H, Chang Y, Wang Q, Liang Q, Qi Q (2016) A CRISPR-Cas9 assisted non-homologous end-joining strategy for one-step engineering of bacterial genome. Sci Rep **6**: 37895

Taton A, Ma AT, Ota M, Golden SS, Golden JW (2017) NOT gate genetic circuits to control gene expression in cyanobacteria. ACS Synth Biol **6**: 2175–2182

Taton A, Unglaub F, Wright NE, Zeng WY, Paz-Yepes J, Brahamsha B, Palenik B, Peterson TC, Haerizadeh F, Golden SS, et al (2014) Broad-host-range vector system for synthetic biology and biotechnology in cyanobacteria. Nucleic Acids Res **42**: e136

Taylor GM, Hitchcock A, Heap JT (2021) Combinatorial assembly platform enabling engineering of genetically stable metabolic pathways in cyanobacteria. Nucleic Acids Res **49**: e123

Ungerer J, Pakrasi HB (2016) Cpf1 is a versatile tool for CRISPR genome editing across diverse species of cyanobacteria. Sci Rep **6**: 39681

Ungerer J, Wendt KE, Hendry JI, Maranas CD, Pakrasi HB (2018) Comparative genomics reveals the molecular determinants of rapid growth of the cyanobacterium Synechococcus elongatus UTEX 2973. Proc Natl Acad Sci USA **115**: E11761–E11770

Vasudevan R, Gale GAR, Schiavon AA, Puzorjov A, Malin J, Gillespie MD, Vavitsas K, Zulkower V, Wang B, Howe CJ, et al (2019) CyanoGate: a modular cloning suite for engineering cyanobacteria based on the plant MoClo syntax. Plant Physiol **180**: 39–55

Vento JM, Crook N, Beisel CL (2019) Barriers to genome editing with CRISPR in bacteria. J Ind Microbiol Biotechnol **46**: 1327–1341

Vogel AIM, Lale R, Hohmann-Marriott MF (2017) Streamlining recombination-mediated genetic engineering by validating three neutral integration sites in Synechococcus sp. PCC 7002. J Biol Eng **11**: 19

Wang S-Y, Li X, Wang S-G, Xia P-F (2023) Base editing for reprogramming cyanobacterium Synechococcus elongatus. Metab Eng **75**: 91–99

Wendt KE, Ungerer J, Cobb RE, Zhao H, Pakrasi HB (2016) CRISPR/Cas9 mediated targeted mutagenesis of the fast growing cyanobacterium Synechococcus elongatus UTEX 2973. Microb Cell Factories **15**: 115

Wilson KS, von Hippel PH (1995) Transcription termination at intrinsic terminators: the role of the RNA hairpin. Proc Natl Acad Sci USA **92**: 8793–8797

Włodarczyk A, Selão TT, Norling B, Nixon PJ (2020) Newly discovered Synechococcus sp. PCC 11901 is a robust cyanobacterial strain for high biomass production. Commun Biol **3**: 1–14

Xiao K, Yue X-H, Chen W-C, Zhou X-R, Wang L, Xu L, Huang F-H, Wan X (2018) Metabolic engineering for enhanced medium chain omega hydroxy fatty acid production in Escherichia coli. Front Microbiol **9**: 139

Yao L, Shabestary K, Björk SM, Asplund-Samuelsson J, Joensson HN, Jahn M, Hudson EP (2020) Pooled CRISPRi screening of the cyanobacterium Synechocystis sp PCC 6803 for enhanced industrial phenotypes. Nat Commun **11**: 1666

Yu J, Liberton M, Cliften PF, Head RD, Jacobs JM, Smith RD, Koppenaal DW, Brand JJ, Pakrasi HB (2015) Synechococcus elongatus UTEX 2973, a fast growing cyanobacterial chassis for biosynthesis using light and CO2. Sci Rep **5**: 8132

Zavřel T, Chmelík D, Sinetova MA, Červený J (2018) Spectrophotometric determination of phycobiliprotein content in cyanobacterium Synechocystis. J Vis Exp 58076

Zess EK, Begemann MB, Pfleger BF (2016) Construction of new synthetic biology tools for the control of gene expression in the cyanobacterium Synechococcus sp. strain PCC 7002. Biotechnol Bioeng **113**: 424–432

Zetsche B, Gootenberg JS, Abudayyeh OO, Slaymaker IM, Makarova KS, Essletzbichler P, Volz SE, Joung J, van der Oost J, Regev A, et al (2015) Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system. Cell **163**: 759–771

Zhang A, Carroll AL, Atsumi S (2017) Carbon recycling by cyanobacteria: improving CO2 fixation through chemical production. FEMS Microbiol Lett **364**: fnx165

Zhou Y, Sun T, Chen Z, Song X, Chen L, Zhang W (2019) Development of a new biocontainment strategy in model cyanobacterium Synechococcus strains. ACS Synth Biol **8**: 2576–2584

FIGURE LEGENDS

Figure 1. Antibiotic susceptibility and characterisation of putative neutral integration sites in PCC 11901. **(A)** Susceptibility of wild-type PCC 11901 to increasing concentrations of common antibiotics. PCC 11901 wild-type cultures were inoculated at $OD_{750} = 0.2$ and grown in MAD medium as described in the Materials and Methods for 48 h. **(B)** The diagram illustrates the transformation strategy used to introduce antibiotic resistance (AbR) cassettes into each putative neutral site via homologous recombination. An integrative pUC19 plasmid vector was assembled using the CyanoGate MoClo system (Vasudevan et al., 2019) and 1 µg of each plasmid was transformed into wild-type (WT) PCC 11901 (see **Supplementary Table S2** for plasmid vectors). Colony images and numbers of mutants transformed with different AbR cassettes integrated into the *desB* neutral site. Colony counts were estimated by dividing the plate into nine sectors and taking the average colony counts of three sectors. Based on these results, we recommend concentrations of 100 µg mL^{-1} kanamycin, 25 µg mL^{-1} spectinomycin, 50 µg mL^{-1} gentamicin, 1.25 µg mL^{-1} erythromycin, and 5 µg mL^{-1} chloramphenicol for selection using the respective AbR cassettes. **(C)** Growth analysis of five putative neutral site mutants transformed with a spectinomycin resistance cassette (SpR) and grown in MAD medium as described in the Materials and Methods. Lowercase letters indicating significant difference ($P < 0.05$) are shown, as determined by ANOVA followed by Tukey's honestly significant difference tests. Error bars show the mean ±SEM of three biological replicates. Abbreviations: Cm, chloramphenicol; DF, Down Flank; Em, erythromycin; Gm, gentamicin; Km, kanamycin; Sp, spectinomycin; UF, Up Flank.

Figure 2. Transconjugation in PCC 11901. **(A)** Illustration showing conjugal transfer of a self-replicating RSF1010-based vector from an *E. coli* 'cargo strain' into PCC 11901. A transmissible helper plasmid is transferred from an *E. coli* helper strain to the cargo *E. coli* strain, which in turn aids the transfer of the mobilizable RSF1010 vector into the recipient PCC 11901. This series of steps is facilitated by the formation of conjugation pili where genetic material is transferred. **(B)** Representative image of PCC 11901 colony growth on membrane filters following selection on AD7 agar supplemented with 100 µg mL^{-1} kanamycin **(C)** pPMQAK1-T-based conjugants showed a typical dark green phenotype or a pale phenotype (red box) after streaking colonies from membranes filters onto agar media containing kanamycin. Pale colonies did not survive re-streaking. **(D)** Survival rates of transconjugants harbouring RSF1010-based vectors selected for with kanamycin (pPMQAK1-T and pPMQAK1-T-eYFP) or spectinomycin (pPMQSK1-1, pPMQSK1-1-

eYFP and pPMQSK1-1-Cas12a). **(E)** Growth comparison of transconjugant and transformant
strains. **(F)** Fluorescence of transconjugant and transformant strains expressing eYFP
measured at 24 h. **(G)** Growth comparison for transformant and transconjugant strains
expressing eYFP. For (E), (F) and (G) lowercase letters indicating significant difference ($P <$
0.05) are shown, as determined by ANOVA followed by Tukey's honestly significant
difference tests. Error bars show the mean ±SEM of three biological replicates.

1102

Figure 3. Characterisation of constitutive promoters and transcriptional terminators in PCC
11901. **(A)** Expression levels of eYFP driven by constitutive promoters integrated into the
mrr neutral site, or **(B)** on the self-replicating pPMQAK1-T vector. **(C)** Correlation analysis
of the expression levels of constitutive promoters integrated into the *mrr* neutral site or on a
pPMQAK1-T vector. **(D)** Termination efficiency (TE) values for terminator sequences
calculated as in Gale et al. (2021). Error bars represent the mean ±SEM of three biological
replicates, each calculated from 10,000 individual cells. Abbreviations: P_{const}, constitutive
promoter; T_{term}, transcriptional terminator.

1111

Figure 4. Evaluation of three inducible promoter systems in PCC 11901**.** Overview of the
genetic components and dose-response expression levels of eYFP 24 h after induction with
increasing concentrations of their respective substrate for the **(A, B)** L-rhamnose-inducible
promoter RhaS/P_{rhaBAD} fluorescent reporter system, **(C, D)** theophylline-inducible promoter
P_{trcE*} fluorescent reporter system and **(E, F)** DAPG-inducible PhlF/P_{phlF} fluorescent reporter
system. The eYFP expression level for the constitutive P_{cpc560} promoter (pPMQAK1-T-eYFP,
Figure 2F) is included for comparison. Error bars show the mean ±SEM of three biological
replicates.

1120

Figure 5. Inducible CRISPR interference (CRISPRi) for conditional knockdown of gene
expression in PCC 11901. **(A)** Overview of the approaches used to test dCas9 functionality
by targeting an eYFP expression cassette integrated into the *mrr* neutral site (bottom
schematic) using sgRNAs targeting four different sites neighbouring a dCas9 protospacer
adjacent motif sequence 5'-NGG-3' in the eYFP open reading frame (Vasudevan et al.,
2019). **(B-E)** eYFP fluorescence of plasmid vectors carrying sgRNAs with and without a
constitutively expressed (P_{J23113}) or inducible (RhaS/P_{rhaBAD}, PhlF/P_{phlF} and P_{trcE*}) dCas9 and
a strain carrying no sgRNA as a control. **(F)** Illustration of the *cpc* operon and the sgRNA
target site (red bar, see **Supplementary Table S3** for sequence) in *cpcB* (c-phycocyanin

1130 subunit β, FEK30_15275). **(G)** The *cpcB* CRISPRi strains with and without inducer after 24

1131 h of growth in MAD medium. **(H)** Growth of the *cpcB* CRISPRi strains with and without

1132 inducer over 48 h. **(I)** Absorbance spectra of PBS extracts from strains in (G). **(J)** Illustration

1133 of the sgRNA target site in *nblA* (nonbleaching A, FEK30_13550). **(K)** The *nblA* CRISPRi

1134 strains with and without inducer after 24 h of growth in MAD medium without nitrogen. **(L)**

1135 Growth of the *nblA* CRISPRi strains with and without inducer over 48 h. **(M)** Absorbance

1136 spectra of PBS extracts from strains in (K). Error bars show the mean ±SEM of three

1137 biological replicates.

1138

1139 **Figure 6.** Genome editing of PCC 11901 with CRISPR-Cas12a using a double homologous

1140 recombination approach. **(A)** The editing strategy relies on an RSF1010-based editing vector

1141 (pC1.509, **Supplementary Figure S1**) and a pUC19 suicide vector. Transformation of the

1142 pUC19 suicide vector into an 'editing strain' carrying pC1.509 results in (1) HR with

1143 pC1.509 to deliver the sgRNA(s) and GmR (components in blue) and (2) HR with the target

1144 genomic locus to deliver a template for homology directed repair (components in purple).

1145 Subsequent DAPG-induction of Cas12a results in cleavage of the target locus (in grey),

1146 leaving only edited copies of the genome (in purple) intact (for a detailed methodology and

1147 protocols see **Supplementary Figure S7** and **S8**). **(B,C)** Demonstration of single genome

1148 editing through markerless insertion of eYFP into the *mrr* locus and mCherry into the *aquI*

1149 locus, respectively. (**D**) Iterative genome editing through markerless insertion of mCherry

1150 into the *aquI* locus of the Δ*mrr*::eYFP strain. (**E**) Multiplexed genome editing through

1151 simultaneous insertion of eYFP and mCherry into the *mrr* and *aquI* locus, respectively. (F)

1152 Demonstration of curing of the RSF1010 editing vector using a self-targeting sgRNA.

1153

Figure 1. Antibiotic susceptibility and characterisation of putative neutral integration sites in PCC 11901. **(A)** Susceptibility of wild-type PCC 11901 to increasing concentrations of common antibiotics. PCC 11901 wild-type cultures were inoculated at $OD_{750} = 0.2$ and grown in MAD medium as described in the Materials and Methods for 48 h. **(B)** The diagram illustrates the transformation strategy used to introduce antibiotic resistance (AbR) cassettes into each putative neutral site via homologous recombination. An integrative pUC19 plasmid vector was assembled using the CyanoGate MoClo system (Vasudevan et al., 2019) and 1 μg of each plasmid was transformed into wild-type (WT) PCC 11901 (see **Supplementary Table S2** for plasmid vectors). Colony images and numbers of mutants transformed with different AbR cassettes integrated into the *desB* neutral site. Colony counts were estimated by dividing the plate into nine sectors and taking the average colony counts of three sectors. Based on these results, we recommend concentrations of 100 μg mL^{-1} kanamycin, 25 μg mL^{-1} spectinomycin, 50 μg mL^{-1} gentamicin, 1.25 μg mL^{-1} erythromycin, and 5 μg mL^{-1} chloramphenicol for selection using the respective AbR cassettes. **(C)** Growth analysis of five putative neutral site mutants transformed with a spectinomycin resistance cassette (SpR) and grown in MAD medium as described in the Materials and Methods. Lowercase letters indicating significant difference ($P < 0.05$) are shown, as determined by ANOVA followed by Tukey's honestly significant difference tests. Error bars show the mean ±SEM of three biological replicates. Abbreviations: Cm, chloramphenicol; DF, Down Flank; Em, erythromycin; Gm, gentamicin; Km, kanamycin; Sp, spectinomycin; UF, Up Flank.

Figure 2. Transconjugation in PCC 11901. **(A)** Illustration showing conjugal transfer of a self-replicating RSF1010-based vector from an *E. coli* 'cargo strain' into PCC 11901. A transmissible helper plasmid is transferred from an *E. coli* helper strain to the cargo *E. coli* strain, which in turn aids the transfer of the mobilizable RSF1010 vector into the recipient PCC 11901. This series of steps is facilitated by the formation of conjugation pili where genetic material is transferred. **(B)** Representative image of PCC 11901 colony growth on membrane filters following selection on AD7 agar supplemented with 100 μg mL^{-1} kanamycin **(C)** pPMQAK1-T-based conjugants showed a typical dark green phenotype or a pale phenotype (red box) after streaking colonies from membranes filters onto agar media containing kanamycin. Pale colonies did not survive re-streaking. **(D)** Survival rates of transconjugants harbouring RSF1010-based vectors selected for with kanamycin (pPMQAK1-T and pPMQAK1-T-eYFP) or spectinomycin (pPMQSK1-1, pPMQSK1-1-eYFP and pPMQSK1-1-Cas12a). **(E)** Growth comparison of transconjugant and transformant strains. **(F)** Fluorescence of transconjugant and transformant strains expressing eYFP measured at 24 h. **(G)** Growth comparison for transformant and transconjugant strains expressing eYFP. For (E), (F) and (G) lowercase letters indicating significant difference (P <

1194   0.05) are shown, as determined by ANOVA followed by Tukey's honestly significant

1195   difference tests. Error bars show the mean ±SEM of three biological replicates.

1196

Figure 3. Characterisation of constitutive promoters and transcriptional terminators in PCC 11901. **(A)** Expression levels of eYFP driven by constitutive promoters integrated into the *mrr* neutral site, or **(B)** on the self-replicating pPMQAK1-T vector. **(C)** Correlation analysis of the expression levels of constitutive promoters integrated into the *mrr* neutral site or on a pPMQAK1-T vector. **(D)** Termination efficiency (TE) values for terminator sequences calculated as in Gale et al. (2021). Error bars represent the mean ±SEM of three biological replicates, each calculated from 10,000 individual cells. Abbreviations: P_{const}, constitutive promoter; T_{term}, transcriptional terminator.

Figure 4. Evaluation of three inducible promoter systems in PCC 11901. Overview of the genetic components and dose-response expression levels of eYFP 24 h after induction with increasing concentrations of their respective substrate for the **(A, B)** L-rhamnose-inducible promoter RhaS/P_{rhaBAD} fluorescent reporter system, **(C, D)** theophylline-inducible promoter P_{trcE*} fluorescent reporter system and **(E, F)** DAPG-inducible PhlF/P_{phlF} fluorescent reporter system. The eYFP expression level for the constitutive P_{cpc560} promoter (pPMQAK1-T-eYFP, **Figure 2F**) is included for comparison. Error bars show the mean ±SEM of three biological replicates.

1218

Figure 5. Inducible CRISPR interference (CRISPRi) for conditional knockdown of gene expression in PCC 11901. **(A)** Overview of the approaches used to test dCas9 functionality

1221 by targeting an eYFP expression cassette integrated into the *mrr* neutral site (bottom

1222 schematic) using sgRNAs targeting four different sites neighbouring a dCas9 protospacer

1223 adjacent motif sequence 5'-NGG-3' in the eYFP open reading frame (Vasudevan et al.,

1224 2019). **(B-E)** eYFP fluorescence of plasmid vectors carrying sgRNAs with and without a

1225 constitutively expressed (P_{J23113}) or inducible (RhaS/P_{rhaBAD}, PhlF/P_{phlF} and P_{trcE*}) dCas9 and

1226 a strain carrying no sgRNA as a control. **(F)** Illustration of the *cpc* operon and the sgRNA

1227 target site (red bar, see **Supplementary Table S3** for sequence) in *cpcB* (c-phycocyanin

1228 subunit β, FEK30_15275). **(G)** The *cpcB* CRISPRi strains with and without inducer after 24

1229 h of growth in MAD medium. **(H)** Growth of the *cpcB* CRISPRi strains with and without

1230 inducer over 48 h. **(I)** Absorbance spectra of PBS extracts from strains in (G). **(J)** Illustration

1231 of the sgRNA target site in *nblA* (nonbleaching A, FEK30_13550). **(K)** The *nblA* CRISPRi

1232 strains with and without inducer after 24 h of growth in MAD medium without nitrogen. **(L)**

1233 Growth of the *nblA* CRISPRi strains with and without inducer over 48 h. **(M)** Absorbance

1234 spectra of PBS extracts from strains in (K). Error bars show the mean ±SEM of three

1235 biological replicates.

1236

1237

Figure 6. Genome editing of PCC 11901 with CRISPR-Cas12a using a double homologous recombination approach. **(A)** The editing strategy relies on an RSF1010-based editing vector (pC1.509, **Supplementary Figure S1**) and a pUC19 suicide vector. Transformation of the pUC19 suicide vector into an 'editing strain' carrying pC1.509 results in (1) HR with pC1.509 to deliver the sgRNA(s) and GmR (components in blue) and (2) HR with the target genomic locus to deliver a template for homology directed repair (components in purple). Subsequent DAPG-induction of Cas12a results in cleavage of the target locus (in grey), leaving only edited copies of the genome (in purple) intact (for a detailed methodology and protocols see **Supplementary Figure S7** and **S8**). **(B,C)** Demonstration of single genome editing through markerless insertion of eYFP into the *mrr* locus and mCherry into the *aquI* locus, respectively. **(D)** Iterative genome editing through markerless insertion of mCherry into the *aquI* locus of the Δ*mrr*::eYFP strain. **(E)** Multiplexed genome editing through simultaneous insertion of eYFP and mCherry into the *mrr* and *aquI* locus, respectively. (F) Demonstration of curing of the RSF1010 editing vector using a self-targeting sgRNA.

1252 **Supplementary Table S1.** Table of all new CyanoGate-compatible parts generated in this work.

1253

No.	Vector ID	Part name	Level	5' overhang	3' overhang	Backbone	Domesticated	Selection	Notes	References
Acceptors										
Level T										
1	pCAT.482	pPMQSK1-T	T	TGCC	GGGA	pPMQSK	yes	SpR/KmR	Modified BioBrick vector, RSF1010-derived ori.	Huang et al., 2010
Level 1										
1	pCA1.339	pPMQSK1-1	1	GGAG	GGGA	pPMQSK	yes	SpR/KmR	Modified BioBrick vector, RSF1010-derived ori.	Huang et al., 2010
Level 0										
1	pCA0.421	sgRNA acceptor	0	AATG	AGGT	pICH41258	no	SpR	Acceptor of sgRNA(s) for CRISPR-Cas12a gene editing	this study
Insertion Sites										
1	pC0.378	*desB* Down Flank	0	GCTT	CGCT	pICH41276	no	SpR	*Synechococcus* sp. PCC 11901 sequence downstream of FEK30_04840.	Mills et al., 2022 Vogel et al., 2017
2	pC0.385	*desB* Up Flank	0	GGAG	AATG	pICH41295	yes	SpR	*Synechococcus* sp. PCC 11901 sequence upstream of FEK30_04840.	Mills et al., 2022 Vogel et al., 2017
3	pC0.386	*glgA1* Up Flank	0	GGAG	AATG	pICH41295	no	SpR	*Synechococcus* sp. PCC 11901 sequence upstream of FEK30_14880.	Sengupta et al., 2020, Mittermair et al., 2021
4	pC0.387	*glgA1* Down Flank	0	GCTT	CGCT	pICH41276	no	SpR	*Synechococcus* sp. PCC 11901 sequence downstream of FEK30_14880.	Sengupta et al., 2020, Mittermair et al., 2021
5	pC0.388	*mrr* Up Flank	0	GGAG	AATG	pICH41295	yes	SpR	*Synechococcus* sp. PCC 11901 sequence upstream of FEK30_09380.	this study
6	pC0.389	*mrr* Down Flank	0	GCTT	CGCT	pICH41276	yes	SpR	*Synechococcus* sp. PCC 11901 sequence downstream of FEK30_09380.	this study
7	pC0.419	*aquI* Down Flank	0	GCTT	CGCT	pICH41276	no	SpR	*Synechococcus* sp. PCC 11901 sequence downstream of FEK30_10065.	this study
8	pC0.420	*aquI* Up Flank	0	GGAG	AATG	pICH41295	no	SpR	*Synechococcus* sp. PCC 11901 sequence upstream of FEK30_10065.	this study
9	pC0.425	RSF1010 Up Flank	0	GGAG	AATG	pICH41295	no	SpR	RSF1010 upstream homology sequence	this study

#	ID	Name		5'	3'	Backbone		Res	Description	Reference
10	pC0.426	RSF1010 Down Flank	0	GCTT	CGCT	pICH41276	no	SpR	RSF1010 downstream homology sequence	this study
11	pC0.437	NS1 Up Flank	0	GGAG	AATG	pICH41295	no	SpR	*Synechococcus* sp. PCC 11901 sequence upstream of an intergenic region of 185bp between FEK30_11550 and FEK30_11555)	Bernhards et al., 2022
12	pC0.438	NS1 Down Flank	0	GCTT	CGCT	pICH41276	yes	SpR	*Synechococcus* sp. PCC 11901 sequence downstream of an intergenic region of 185bp between FEK30_11550 and FEK30_11555)	Bernhards et al., 2022

Promoters

#	ID	Name		5'	3'	Backbone		Res	Description	Reference
1	pC0 422	P_{rhaBAD}	0	GGAG	AATG	pICH41295	no	SpR	L-rhamnose inducible promoter	Kelly et al., 2018
2	pC0.423	P_{phlF}	0	GGAG	AATG	pICH41295	no	SpR	2,4-diacetylphloroglucinol inducible promoter from *Pseudomonas fluorescens*	Meyer et al., 2019
3	pC0.394	P_{trcE*}	0	GGAG	AATG	pSB4K5	no	SpR	theophylline responsive riboswitch E*	Nakahira et al., 2014

Parts

#	ID	Name		5'	3'	Backbone		Res	Description	Reference
1	pC0.393	Ab[R]Gent	0	AGGT	GCCT	pICH41264	no	SpR/ GmR	Gentamicin resistance cassette	this study
2	pC0.391	Cas12a	0			pICH41308	no	SpR	Cas12a/Cpf1 from Francisella novicida	Zetsche et al., 2015

Other

#	ID	Name		5'	3'	Backbone		Res	Description	Reference
1	pC1.509	Cas12a editing vector	1			pPMQSK1-1	no	SpR	Editing vector for CRISPR-Cas12a gene editing (see Figure 6 for details)	this study
2	pC1.530	Curing vector (GmR) for editing strain	1			pPMQSK1-1	no	SpR	Curing vector with sgRNA and GmR for removing Cas12a editing vector (see Figure 6x for details)	this study

Additional references

Zetsche, B., Gootenberg, J. S., Abudayyeh, O. O., Slaymaker, I. M., Makarova, K. S., Essletzbichler, P., Volz, S. E., Joung, J., van der Oost, J., Regev, A., Koonin, E. V., & Zhang, F. (2015). Cpf1 Is a Single RNA-Guided Endonuclease of a Class 2 CRISPR-Cas System. *Cell*, *163*(3), 759–771. https://doi.org/10.1016/j.cell.2015.09.038

1254   **Supplementary Table S2.** All plasmids made and used in this study. See 'Supplementary

1255   Table S2 - Plasmids (all).xlsx'. RSF1010 vectors used for promoter and terminator

1256   characterisations in Figure 3B and 3C, respectively, have been published previously

1257   (Vasudevan et al., 2019; Gale et al., 2021).

1258

1259   **Supplementary Table S3.** Primer and sgRNA oligonucleotides used in this study. See

1260   'Supplementary Table S3 - Primer and sgRNA oligonucleotides.xlsx'.

1261

1262   **Supplementary Data S1.** Sequence maps (.gb files) for plasmid vectors in Supplementary

1263   Table S1. See 'Supplementary Data S1.zip'.

1264

1265

A

1266

1267 **B**

1268 Aligned_sequences: 2
1: BAI89523.1

1269 # 2: QCS49435.1
Matrix: EBLOSUM62

1270 # Gap_penalty: 10.0
Extend_penalty: 0.5

1271 # Length: 219
Identity: 46/219 (21.0%)

1272 # Similarity: 68/219 (31.1%)
Gaps: 87/219 (39.7%)

1273 # Score: 100.0

1274 CLUSTAL O(1.2.4) multiple sequence alignment

1275 BAI89523.1   MRIINLSPNNQNHIHQAATLLV-AEFRENWPNAWPTYERGLAEVMESFGDDRVNLVAVDE 59

QCS49435.1   MNKISDCSNILFSVDKARVDLVQLQALFNATAFWA-RERSLADLETAIA-YSDPVVTVWD 58

1276   *. *. . *   :.:* . ** : *   *   **.**:: ::.   :*:* :

1277

BAI89523.1   NDNLLGWIGGIS--QYQGHVWELHPLVVKSDYQGLGIGRKLVANLEDYVRSQGGLTLWLG 117

1278 QCS49435.1   GDRLIGFTRGTSDGVFRATVWD---VVIHPDYQGLGLGRKLVETLISHPRMCRVERVYLM 115

1279   .*.*:*: * *   ::. **:   :*:: ******:***** .* .: *   ::*

1280

BAI89523.1   TDDENNLTSLSGVELYPHFLENIANIKNHGRHPYEFYQKCGFVIMGVVPDANGIGKPDIL 177

1281 QCS49435.1   TTH--------------------------QQTFYERIGFKENATTTM---------- 136

1282   * .   **:: **   ...

1283 BAI89523.1   MAKSLRIDNHPKSN-------------   191

1284 QCS49435.1   ---VLHNHEHPISVITCAVEETAPLPA   160

              *: .:** *

1285

Supplementary Figure S1. Analysis of putative Gcn5-related N-acetyltransferase (GNAT)

family genes in PCC 11901. **(A)** Phylogenetic tree of 13 GNAT family proteins in PCC

11901 and the aminoglycoside acetyltransferase from *Arthrospira platensis* (BAI89523.1)

(built by Clustal Omega (EMBL-EBI)). **(B)** Sequence alignment and scores between

BAI89523.1 and a putative aminoglycoside acetyltransferase from PCC 11901

(QCS49435.1). The conserved coenzyme A binding pocket is highlighted in yellow. The

alignment was performed in EMBOSS Needle.

1293

1294

Supplementary Figure S2. Representative PCR-based segregation analysis of PCC 11901

transformants targeting neutral sites *glgA1* and *aquI*. Electrophoresis gel profiles are shown

for four independent transformant colonies after one round of re-streaking on selective agar.

Full segregation for each transformant is demonstrated by the presence of amplicons showing

insertion of the SpR cassette at the *glgA1* (1,618 bp) and *aquI* (1,832 bp) loci and the absence

of a respective wild-type (WT) amplicon (1,902 bp for *glgA1* and 1,541 bp for *aquI*).

Amplicons from each transformation plasmid (+) are included as positive controls.

Supplementary Figure S3. Growth of PCC 11901 with varying doses of small molecule inducers. Growth of wild-type PCC 11901 in increasing concentrations of **(A)** L-rhamnose, **(B)** theophylline, and **(C)** DAPG over 72 h. Error bars represent the mean ±SEM of three biological replicates.

Supplementary Figure S4. Growth analysis of CRISPRi-dCas9 strains targeting eYFP. **(A)** Growth of P_{J23113} CRISPRi strains in MAD medium. **(B)** Growth of RhaS/P_{rhaBAD} CRISPRi strains in the presence or absence of 10 mM L-rhamnose. **(C)** Growth of P_{trcE*} CRISPRi strains in the presence or absence of 2 mM theophylline. **(D)** Growth of PhlF/P_{phlF} CRISPRi strains in the presence or absence of 10 μM DAPG. Error bars represent mean ±SEM of three biological replicates.

Supplementary Figure S5. Analysis of CRISPRi-dCAs9 strains targeting *cpcB* and *nblA*. **(A)** Coomassie-stained SDS-PAGE gel of PBS extracts from the *cpcB* CRISPRi strain carrying both sgRNA and dCas9 in the absence (uninduced) or presence (induced) of 10 μM DAPG. Arrows indicate the bands for *apcB* (18.7 kdA), *cpcB* (18.1 kdA), *cpcA* (17.6 kdA) and *apcA* (17.3 kdA). **(B)** Chlorophyll contents of the *cpcB* CRISPRi strains. **(C)** Growth of the *cpcB* CRISPRi strain carrying both sgRNA and dCas9 in comparison to wild-type PCC 11901 following removal of DAPG. **(D)** Absorbance spectra of PBS extracts following removal of DAPG at 48 h. **(E)** Chlorophyll contents of *nblA* CRISPRi strains. Error bars show the mean ±SEM of three biological replicates for B, C and E.

1329

Supplementary Figure S6. Attempted generation of markerless mutants using the CRE-Lox system. Schematic representations of locus location in the PCC 11901 genome (top) and the

wild-type (middle) and marked knockout (KO, bottom) profiles expected in the **(A)** *ctaDIEI*,

(C) *ctaCII*, **(E)** *acs*, **(G)** *ldhA* and **(I)** *sdhA* strains following amplification with primers

flanking the deleted sequence. Regions deleted in the mutant strains are shaded in black.

Amplification of genomic DNA in: **(B)** Δ*ctaDIEI* KmR marked KO (lane 1); Δ*ctaDIEI* GmR

marked KO (lanes 2-5); WT (lane 6); **(D)** negative control (lane 1); WT (lane 2); Δ*ctaCII*

GmR marked KO (lanes 3-4); **(F)** WT (lane 1); negative control (lane 2); Δ*acs* GmR marked

KO (lane 3); **(H)** WT (lane 1); negative control (lane 2); Δ*ldhA* GmR marked KO (lane 3);

(J) negative control (lane 1); WT (lane 2); Δ*sdhA* KmR marked KO (lane 3). Δ*ctaDIEI* GmR

marked   KOs   were   sequenced   to   confirm   the   correct   KO   profile.

Assembly of sgRNA plasmid vectors (adapted from Liao et al., 2019)

1. To anneal oligonucleotides for making dsDNA parts for assembly (A), add 15 µL of the forward and reverse oligonucleotides into a PCR tube and incubate in a thermal cycler using the following step-down program: 95°C for 5 min, then decrease by 1°C incrementally and hold for 30 seconds for 79 cycles (until 25°C).

2. Level 0 assembly (B): In a PCR tube, add 2 µL of T4 ligation buffer, 2 µL of ATP (10 mM), 1 µL of each annealed repeat-spacer subunit (from 1.), 50 ng of the sgRNA acceptor vector (pCA0.421, Supplementary Table S1), 0.5 µL of T4 ligase, 1.5 µL of BpiI, and water to a total volume of 20 µL. Mix by pipetting and spin down briefly. Incubate the tube in a thermal cycler using the following program: 30 cycles of alternating digestion and ligation (37°C for 5 min, 16°C for 5 min) followed by a final digestion step (60°C for 5 min).

3. In a 1.5 mL microcentrifuge tube, transform 10 µL of the assembly reaction into 30 µl of competent TOP10 *E. coli* cells using the heat shock method. Add 500 µL LB to the cells and shake at 37°C for 1 h. Plate the cells in LB gar supplemented with 100 ug/mL spectinomycin and 40 ug/mL X-gal, and incubate overnight at 37 °C.

4. Screen the white colonies by colony PCR and Sanger sequencing.

Supplementary Figure S7. CRISPR-Cas12a double HR editing approach – sgRNA, repair template and hybrid suicide vector assembly. **(A)** sgRNA were assembled by annealing oligonucleotides incorporating the desired 18-22bp sgRNA (yellow), direct repeat (grey),

unique 4 bp overhangs (red) and BpiI sites (underlined) for **(B)** assembly into the pC0.421

sgRNA acceptor vector. Overhangs match those from the CRATES system and allow for

assembly of an array of between one and seven sgRNAs (Liao et al., 2019). **(C)** MoClo

assembly of level 0 parts into level 1 vectors and then a level T hybrid suicide vector

incorporating the sgRNA(s) in position 1 and homology repair template(s) in position 2 up to

position 7 using the available Plant MoClo level 1 acceptor vectors (Engler et al. 2014).

Using this system between one and six genomic loci can be targeted simultaneously. A

protocol for assembly of the sgRNA plasmid vectors is included.

1368
1369

Protocol for transformation with hybrid suicide vectors

1. Grow the editing strain from OD_{750} = 0.2 to the initial stages of exponential growth (OD_{750} = ~0.8).

2. Centrifuge 1 mL of the editing strain for 10 min at 3,500 g and wash the cell pellet with fresh culture media three
times to remove antibiotics.

3. Dilute the washed editing strain to OD_{750} = 0.2 and add 1.5 µg of purified hybrid suicide vector to 1 mL of the editing
strain.

4. Incubate for 4 h at 30°C in 2% (v/v) CO_2 under warm white LED light (150 µmol photons m^{-2} s^{-1}) before adding 10 µM
DAPG and continuing incubation overnight.

5. Plate the transformation mixture on to agar plates supplemented with appropriate antibiotics (25 µg mL^{-1}
spectinomycin, 50 µg mL^{-1} gentamicin) and 10 µM DAPG and incubate as described in step 4.

6. Screen for mutant colonies using colony PCR.

1381

Supplementary Figure S8. Pipeline for iterative CRISPR-Cas12a editing using the double

HR approach. In the first round of editing the hybrid suicide vector recombines with the

editing vector to introduce an sgRNA (or sgRNA array) and replaces the SpR with GmR for

subsequent selection on gentamicin-supplemented agar plates (also see **Figure 6A**).

Subsequent rounds of editing involved cycling between hybrid suicide vectors that exchange

SpR and GmR for delivery of new sgRNA(s) to the editing vector. Curing the edited strain of

the editing vector to generate a fully markerless strain is done using a hybrid suicide vector

(e.g. pC1.530, **Supplementary Table S1**), which delivers a self-targeting sgRNA to the

editing vector. A protocol for transformation with the hybrid suicide vector and induction

with DAPG is included.

1392

1393

1394 **Supplementary Figure S9.** Confirmation of recombination between the hybrid suicide

1395 vector and editing vector. Example colony PCR of a transformant showing replacement of

1396 SpR by GmR

1397

Parsed Citations

Abbas A, Morrissey JP, Marquez PC, Sheehan MM, Delany IR, O'Gara F (2002) Characterization of interactions between the transcriptional repressor PhlF and its binding site at the phlA promoter in Pseudomonas fluorescens F113. J Bacteriol 184: 3008–3016
Google Scholar: Author Only Title Only Author and Title

Bae S, Park J, Kim J-S (2014) Cas-OFFinder: a fast and versatile algorithm that searches for potential off-target sites of Cas9 RNA-guided endonucleases. Bioinformatics 30: 1473–1475
Google Scholar: Author Only Title Only Author and Title

Baldanta S, Guevara G, Navarro-Llorens JM (2022) SEVA-Cpf1, a CRISPR-Cas12a vector for genome editing in cyanobacteria. Microb Cell Factories 21: 103
Google Scholar: Author Only Title Only Author and Title

Batterton JC, Van Baalen C (1971) Growth responses of blue-green algae to sodium chloride concentration. Archiv Mikrobiol 76: 151–165
Google Scholar: Author Only Title Only Author and Title

Behle A, Saake P, Germann AT, Dienst D, Axmann IM (2020) Comparative dose–response analysis of inducible promoters in cyanobacteria. ACS Synth Biol 9: 843–855
Google Scholar: Author Only Title Only Author and Title

Behler J, Vijay D, Hess WR, Akhtar MK (2018) CRISPR-based technologies for metabolic engineering in cyanobacteria. Trends Biotechnol 36: 996–1010
Google Scholar: Author Only Title Only Author and Title

BEIS (2021) 'Net Zero Strategy: Build Back Greener'. Department for Business, Energy, and International Strategy. Available at: https://www.gov.uk/government/publications/net-zero-strategy (Accessed: 13 September 2022).
Google Scholar: Author Only Title Only Author and Title

Bernhards CB, Liem AT, Berk KL, Roth PA, Gibbons HS, Lux MW (2022) Putative phenotypically neutral genomic insertion points in prokaryotes. ACS Synth Biol 11: 1681–1685
Google Scholar: Author Only Title Only Author and Title

Bishé B, Taton A, Golden JW (2019) Modification of RSF1010-based broad-host-range plasmids for improved conjugation and cyanobacterial bioprospecting. iScience 20: 216–228
Google Scholar: Author Only Title Only Author and Title

Cameron JC, Pakrasi HB (2011) Glutathione facilitates antibiotic resistance and photosystem I stability during exposure to gentamicin in cyanobacteria. Appl Environ Microbiol 77: 3547–3550
Google Scholar: Author Only Title Only Author and Title

Cengic I, Cañadas IC, Minton NP, Hudson EP (2022) Inducible CRISPR/Cas9 allows for multiplexed and rapidly segregated single-target genome editing in Synechocystis sp. PCC 6803. ACS Synth Biol 11: 3100–3113
Google Scholar: Author Only Title Only Author and Title

Chen Y-J, Liu P, Nielsen AAK, Brophy JAN, Clancy K, Peterson T, Voigt CA (2013) Characterization of 582 natural and synthetic terminators and quantification of their design constraints. Nat Methods 10: 659–664
Google Scholar: Author Only Title Only Author and Title

Chi X, Zhang S, Sun H, Duan Y, Qiao C, Luan G, Lu X (2019) Adopting a theophylline-responsive riboswitch for flexible regulation and understanding of glycogen metabolism in synechococcus elongatus PCC 7942. Front Microbiol 10:
Google Scholar: Author Only Title Only Author and Title

Cho BA, Moreno-Cabezuelo JÁ, Mills LA, del Río Chanona EA, Lea-Smith DJ, Zhang D (2023) Integrated experimental and photo-mechanistic modelling of biomass and optical density production of fast versus slow growing model cyanobacteria. Algal Res 70: 102997
Google Scholar: Author Only Title Only Author and Title

Collias D, Vialetto E, Yu J, Co K, Almási É d H, Rüttiger A-S, Achmedov T, Strowig T, Beisel CL (2023) Systematically attenuating DNA targeting enables CRISPR-driven editing in bacteria. Nat Commun 14: 680
Google Scholar: Author Only Title Only Author and Title

Collier JL, Grossman AR (1994) A small polypeptide triggers complete degradation of light-harvesting phycobiliproteins in nutrient-deprived cyanobacteria. EMBO J 13: 1039–1047
Google Scholar: Author Only Title Only Author and Title

Dallo T, Krishnakumar R, Kolker SD, Ruffing AM (2023) High-Density Guide RNA Tiling and Machine Learning for Designing CRISPR Interference in Synechococcus sp. PCC 7002. ACS Synth Biol. doi: 10.1021/acssynbio.2c00653

Google Scholar: Author Only Title Only Author and Title

Daneshvar E, Wicker RJ, Show P-L, Bhatnagar A (2022) Biologically-mediated carbon capture and utilization by microalgae towards sustainable CO2 biofixation and biomass valorization – A review. Chem Eng J 427: 130884

Google Scholar: Author Only Title Only Author and Title

Depardieu F, Bikard D (2020) Gene silencing with CRISPRi in bacteria and optimization of dCas9 expression levels. Methods 172: 61–75

Google Scholar: Author Only Title Only Author and Title

Dias E, Oliveira M, Jones-Dias D, Vasconcelos V, Ferreira E, Manageiro V, Caniça M (2015) Assessing the antibiotic susceptibility of freshwater Cyanobacteria spp. Front Microbiol 6: 799

Google Scholar: Author Only Title Only Author and Title

Domínguez-Martín MA, Sauer PV, Kirst H, Sutter M, Bína D, Greber BJ, Nogales E, Polívka T, Kerfeld CA (2022) Structures of a phycobilisome in light-harvesting and photoprotected states. Nature 609: 835–845

Google Scholar: Author Only Title Only Author and Title

Engler C, Youles M, Gruetzner R, Ehnert T-M, Werner S, Jones JDG, Patron NJ, Marillonnet S (2014) A golden gate modular cloning toolbox for plants. ACS Synth Biol 3: 839–843

Google Scholar: Author Only Title Only Author and Title

Favrot L, Blanchard JS, Vergnolle O (2016) Bacterial GCN5-Related N-Acetyltransferases: From Resistance to Regulation. Biochemistry 55: 989–1002

Google Scholar: Author Only Title Only Author and Title

Fontana J, Dong C, Ham JY, Zalatan JG, Carothers JM (2018) Regulated expression of sgRNAs tunes CRISPRi in E. coli. Biotechnology Journal 13: 1800069

Google Scholar: Author Only Title Only Author and Title

Gale GAR, Schiavon Osorio AA, Mills LA, Wang B, Lea-Smith DJ, McCormick AJ (2019) Emerging species and genome editing tools: future prospects in cyanobacterial synthetic biology. Microorganisms 7: 409

Google Scholar: Author Only Title Only Author and Title

Gale GAR, Wang B, McCormick AJ (2021) Evaluation and comparison of the efficiency of transcription terminators in different cyanobacterial species. Front Microbiol 11:

Google Scholar: Author Only Title Only Author and Title

Goodchild-Michelman IM, Church GM, Schubert MG, Tang T-C (2023) Light and carbon: Synthetic biology toward new cyanobacteria-based living biomaterials. Mater Today Bio 19: 100583

Google Scholar: Author Only Title Only Author and Title

Gordon GC, Korosh TC, Cameron JC, Markley AL, Begemann MB, Pfleger BF (2016) CRISPR interference as a titratable, trans-acting regulatory tool for metabolic engineering in the cyanobacterium Synechococcus sp. strain PCC 7002. Metab Eng 38: 170–179

Google Scholar: Author Only Title Only Author and Title

Hitchcock A, Hunter CN, Canniffe DP (2020) Progress and challenges in engineering cyanobacteria as chassis for light-driven biotechnology. Microb Biotechnol 13: 363–367

Google Scholar: Author Only Title Only Author and Title

Huang H-H, Lindblad P (2013) Wide-dynamic-range promoters engineered for cyanobacteria. J Biol Eng 7: 10

Google Scholar: Author Only Title Only Author and Title

Iwai M, Katoh H, Katayama M, Ikeuchi M (2004) Improved genetic transformation of the thermophilic cyanobacterium, thermosynechococcus elongatus BP-1. Plant Cell Physiol 45: 171–175

Google Scholar: Author Only Title Only Author and Title

Jaiswal D, Sengupta A, Sengupta S, Madhu S, Pakrasi HB, Wangikar PP (2020) A novel cyanobacterium Synechococcus elongatus PCC11802 has distinct genomic and metabolomic characteristics compared to its neighbor PCC 11801. Sci Rep 10: 191

Google Scholar: Author Only Title Only Author and Title

Jaiswal D, Sengupta A, Sohoni S, Sengupta S, Phadnavis AG, Pakrasi HB, Wangikar PP (2018) Genome features and biochemical characteristics of a robust, fast growing and naturally transformable cyanobacterium Synechococcus elongatus PCC11801 isolated from India. Sci Rep 8: 16632

Google Scholar: Author Only Title Only Author and Title

Jester BW, Zhao H, Gewe M, Adame T, Perruzza L, Bolick DT, Agosti J, Khuong N, Kuestner R, Gamble C, et al (2022) Development of spirulina for the manufacture and oral delivery of protein therapeutics. Nat Biotechnol 40: 956–964

Google Scholar: Author Only Title Only Author and Title

Jones CM, Parrish S, Nielsen DR (2021) Exploiting polyploidy for markerless and plasmid-free genome engineering in cyanobacteria. ACS Synth Biol 10: 2371–2382
Google Scholar: Author Only Title Only Author and Title

Kavita K, Breaker RR (2023) Discovering riboswitches: the past and the future. Trends Biochem Sci 48: 119–141
Google Scholar: Author Only Title Only Author and Title

Kelly CL, Taylor GM, Hitchcock A, Torres-Méndez A, Heap JT (2018) A rhamnose-inducible system for precise and temporal control of gene expression in cyanobacteria. ACS Synth Biol 7: 1056–1066
Google Scholar: Author Only Title Only Author and Title

Khetkorn W, Incharoensakdi A, Lindblad P, Jantaro S (2016) Enhancement of poly-3-hydroxybutyrate production in Synechocystis sp. PCC 6803 by overexpression of its native biosynthetic genes. Bioresour Technol 214: 761–768
Google Scholar: Author Only Title Only Author and Title

Kopka J, Schmidt S, Dethloff F, Pade N, Berendt S, Schottkowski M, Martin N, Dühring U, Kuchmina E, Enke H, et al (2017) Systems analysis of ethanol production in the genetically engineered cyanobacterium Synechococcus sp. PCC 7002. Biotechnol Biofuels 10: 56
Google Scholar: Author Only Title Only Author and Title

Kufryk GI, Sachet M, Schmetterer G, Vermaas WFJ (2002) Transformation of the cyanobacterium Synechocystis sp. PCC 6803 as a tool for genetic mapping: optimization of efficiency. FEMS Microbiol Lett 206: 215–219
Google Scholar: Author Only Title Only Author and Title

Lauritsen I, Porse A, Sommer MOA, Nørholm MHH (2017) A versatile one-step CRISPR-Cas9 based approach to plasmid-curing. Microb Cell Factories 16: 135
Google Scholar: Author Only Title Only Author and Title

Lea-Smith DJ, Bombelli P, Dennis JS, Scott SA, Smith AG, Howe CJ (2014) Phycobilisome-deficient strains of Synechocystis sp. PCC 6803 have reduced size and require carbon-limiting conditions to exhibit enhanced productivity. Plant Physiol 165: 705–714
Google Scholar: Author Only Title Only Author and Title

Lea-Smith DJ, Summerfield TC, Ducat DC, Lu X, McCormick AJ, Purton S (2021) Editorial: Exploring the growing role of cyanobacteria in industrial biotechnology and sustainability. Front Microbiol 12:
Google Scholar: Author Only Title Only Author and Title

Lea-Smith DJ, Vasudevan R, Howe CJ (2016) Generation of marked and markerless mutants in model cyanobacterial species. J Vis Exp 54001
Google Scholar: Author Only Title Only Author and Title

Li H, Shen CR, Huang C-H, Sung L-Y, Wu M-Y, Hu Y-C (2016) CRISPR-Cas9 for the genome engineering of cyanobacteria and succinate production. Metab Eng 38: 293–302
Google Scholar: Author Only Title Only Author and Title

Li S, Sun T, Xu C, Chen L, Zhang W (2018) Development and optimization of genetic toolboxes for a fast-growing cyanobacterium Synechococcus elongatus UTEX 2973. Metab Eng 48: 163–174
Google Scholar: Author Only Title Only Author and Title

Liberton M, Chrisler WB, Nicora CD, Moore RJ, Smith RD, Koppenaal DW, Pakrasi HB, Jacobs JM (2017) Phycobilisome truncation causes widespread proteome changes in Synechocystis sp. PCC 6803. PLoS One 12: e0173251
Google Scholar: Author Only Title Only Author and Title

Lin P-C, Zhang F, Pakrasi HB (2021) Enhanced limonene production in a fast-growing cyanobacterium through combinatorial metabolic engineering. Metab Eng Commun 12: e00164
Google Scholar: Author Only Title Only Author and Title

Lindberg P, Park S, Melis A (2010) Engineering a platform for photosynthetic isoprene production in cyanobacteria, using Synechocystis as the model organism. Metab Eng 12: 70–79
Google Scholar: Author Only Title Only Author and Title

Liu D, Johnson VM, Pakrasi HB (2020) A reversibly induced CRISPRi system targeting photosystem II in the cyanobacterium Synechocystis sp. PCC 6803. ACS Synth Biol 9: 1441–1449
Google Scholar: Author Only Title Only Author and Title

Liu D, Pakrasi HB (2018) Exploring native genetic elements as plug-in tools for synthetic biology in the cyanobacterium Synechocystis sp. PCC 6803. Microb Cell Factories 17: 48
Google Scholar: Author Only Title Only Author and Title

Ma AT, Schmidt CM, Golden JW (2014) Regulation of gene expression in diverse cyanobacterial species by using theophylline-responsive riboswitches. Appl Environ Microbiol 80: 6704–6713

Google Scholar: Author Only Title Only Author and Title

Meyer AJ, Segall-Shapiro TH, Glassey E, Zhang J, Voigt CA (2019) Escherichia coli "Marionette" strains with 12 highly optimized small-molecule sensors. Nat Chem Biol 15: 196–204
Google Scholar: Author Only Title Only Author and Title

Meyer R (2009) Replication and conjugative mobilization of broad host-range IncQ plasmids. Plasmid 62: 57–70
Google Scholar: Author Only Title Only Author and Title

Miao R, Jahn M, Shabestary K, Hudson EP (2023) CRISPR interference screens reveal tradeoffs between growth rate and robustness in Synechocystis sp. PCC 6803 across trophic conditions. 2023.02.13.528328
Google Scholar: Author Only Title Only Author and Title

Mills LA, Moreno-Cabezuelo JÁ, Wlodarczyk A, Victoria AJ, Mejías R, Nenninger A, Moxon S, Bombelli P, Selão TT, McCormick AJ, et al (2022) Development of a biotechnology platform for the fast-growing cyanobacterium Synechococcus sp. PCC 11901. Biomolecules 12: 872
Google Scholar: Author Only Title Only Author and Title

Mittermair S, Lakatos G, Nicoletti C, Ranglová K, Manoel JC, Grivalský T, Kozhan DM, Masojídek J, Richter J (2021) Impact of glgA1, glgA2 or glgC overexpression on growth and glycogen production in Synechocystis sp. PCC 6803. J Biotechnol 340: 47–56
Google Scholar: Author Only Title Only Author and Title

Nakahira Y, Ogawa A, Asano H, Oyama T, Tozawa Y (2013) Theophylline-dependent riboswitch as a novel genetic tool for strict regulation of protein expression in cyanobacterium synechococcus elongatus PCC 7942. Plant Cell Physiol 54: 1724–1735
Google Scholar: Author Only Title Only Author and Title

Ng AH, Berla BM, Pakrasi HB (2015) Fine-tuning of photoautotrophic protein production by combining promoters and neutral sites in the cyanobacterium Synechocystis sp. strain PCC 6803. Appl Environ Microbiol 81: 6857–6863
Google Scholar: Author Only Title Only Author and Title

Opel F, Siebert NA, Klatt S, Tüllinghoff A, Hantke JG, Toepel J, Bühler B, Nürnberg DJ, Klähn S (2022) Generation of synthetic shuttle vectors enabling modular genetic engineering of cyanobacteria. ACS Synth Biol 11: 1758–1771
Google Scholar: Author Only Title Only Author and Title

Pinto F, Pacheco CC, Oliveira P, Montagud A, Landels A, Couto N, Wright PC, Urchueguía JF, Tamagnini P (2015) Improving a Synechocystis-based photoautotrophic chassis through systematic genome mapping and validation of neutral sites. DNA Res 22: 425–437
Google Scholar: Author Only Title Only Author and Title

Porra RJ, Thompson WA, Kriedemann PE (1989) Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. Biochim Biophys Acta - Bioenerg 975: 384–394
Google Scholar: Author Only Title Only Author and Title

Puzorjov A, Dunn KE, McCormick AJ (2021) Production of thermostable phycocyanin in a mesophilic cyanobacterium. Metab Eng Commun 13: e00175
Google Scholar: Author Only Title Only Author and Title

Puzorjov A, McCormick AJ (2020) Phycobiliproteins from extreme environments and their potential applications. J Exp Bot 71: 3827–3842
Google Scholar: Author Only Title Only Author and Title

Puzorjov A, Mert Unal S, Wear MA, McCormick AJ (2022) Pilot scale production, extraction and purification of a thermostable phycocyanin from Synechocystis sp. PCC 6803. Bioresour Technol 345: 126459
Google Scholar: Author Only Title Only Author and Title

Racharaks R, Arnold W, Peccia J (2021) Development of CRISPR-Cas9 knock-in tools for free fatty acid production using the fast-growing cyanobacterial strain Synechococcus elongatus UTEX 2973. J Microbiol Methods 189: 106315
Google Scholar: Author Only Title Only Author and Title

Rautela A, Kumar S (2022) Engineering plant family TPS into cyanobacterial host for terpenoids production. Plant Cell Rep 41: 1791–1803
Google Scholar: Author Only Title Only Author and Title

Ruffing AM, Jensen TJ, Strickland LM (2016) Genetic tools for advancement of Synechococcus sp. PCC 7002 as a cyanobacterial chassis. Microb Cell Factories 15: 190
Google Scholar: Author Only Title Only Author and Title

Sakamoto T, Shen G, Higashi S, Murata N, Bryant DA (1997) Alteration of low-temperature susceptibility of the cyanobacterium Synechococcus sp. PCC 7002 by genetic manipulation of membrane lipid unsaturation. Arch Microbiol 169: 20–28
Google Scholar: Author Only Title Only Author and Title

Sakurai I, Stazic D, Eisenhut M, Vuorio E, Steglich C, Hess WR, Aro E-M (2012) Positive regulation of psbA gene expression by cis-encoded antisense RNAs in Synechocystis sp. PCC 6803. Plant Physiol 160: 1000–1010

Google Scholar: Author Only Title Only Author and Title

Santomartino R, Averesch NJH, Bhuiyan M, Cockell CS, Colangelo J, Gumulya Y, Lehner B, Lopez-Ayala I, McMahon S, Mohanty A, et al (2023) Toward sustainable space exploration: a roadmap for harnessing the power of microorganisms. Nat Commun 14: 1391

Google Scholar: Author Only Title Only Author and Title

Saveria T, Parthiban C, Seilie AM, Brady C, Martinez A, Manocha R, Afreen E, Zhao H, Krzeszowski A, Ferrara J, et al (2022) Needle-free, spirulina-produced Plasmodium falciparum circumsporozoite vaccination provides sterile protection against pre-erythrocytic malaria in mice. NPJ Vaccines 7: 1–11

Google Scholar: Author Only Title Only Author and Title

Selão TT (2022) Exploring cyanobacterial diversity for sustainable biotechnology. J Exp Bot 73: 3057–3071

Google Scholar: Author Only Title Only Author and Title

Sengupta A, Pritam P, Jaiswal D, Bandyopadhyay A, Pakrasi HB, Wangikar PP (2020a) Photosynthetic co-production of succinate and ethylene in a fast-growing cyanobacterium, Synechococcus elongatus PCC 11801. Metabolites 10: 250

Google Scholar: Author Only Title Only Author and Title

Sengupta S, Jaiswal D, Sengupta A, Shah S, Gadagkar S, Wangikar PP (2020b) Metabolic engineering of a fast-growing cyanobacterium Synechococcus elongatus PCC 11801 for photoautotrophic production of succinic acid. Biotechnol Biofuels 13: 89

Google Scholar: Author Only Title Only Author and Title

Shiraishi H, Nishida H (2022) Complete genome sequence of the edible filamentous cyanobacterium arthrospira platensis nies-39, based on long-read sequencing. Microbiol Resour Announc 12: e01139-22

Google Scholar: Author Only Title Only Author and Title

Soini J, Ukkonen K, Neubauer P (2008) High cell density media for Escherichia coli are generally designed for aerobic cultivations – consequences for large-scale bioprocesses and shake flask cultures. Microb Cell Factories 7: 26

Google Scholar: Author Only Title Only Author and Title

Su T, Liu F, Gu P, Jin H, Chang Y, Wang Q, Liang Q, Qi Q (2016) A CRISPR-Cas9 assisted non-homologous end-joining strategy for one-step engineering of bacterial genome. Sci Rep 6: 37895

Google Scholar: Author Only Title Only Author and Title

Taton A, Ma AT, Ota M, Golden SS, Golden JW (2017) NOT gate genetic circuits to control gene expression in cyanobacteria. ACS Synth Biol 6: 2175–2182

Google Scholar: Author Only Title Only Author and Title

Taton A, Unglaub F, Wright NE, Zeng WY, Paz-Yepes J, Brahamsha B, Palenik B, Peterson TC, Haerizadeh F, Golden SS, et al (2014) Broad-host-range vector system for synthetic biology and biotechnology in cyanobacteria. Nucleic Acids Res 42: e136

Google Scholar: Author Only Title Only Author and Title

Taylor GM, Hitchcock A, Heap JT (2021) Combinatorial assembly platform enabling engineering of genetically stable metabolic pathways in cyanobacteria. Nucleic Acids Res 49: e123

Google Scholar: Author Only Title Only Author and Title

Ungerer J, Pakrasi HB (2016) Cpf1 is a versatile tool for CRISPR genome editing across diverse species of cyanobacteria. Sci Rep 6: 39681

Google Scholar: Author Only Title Only Author and Title

Ungerer J, Wendt KE, Hendry JI, Maranas CD, Pakrasi HB (2018) Comparative genomics reveals the molecular determinants of rapid growth of the cyanobacterium Synechococcus elongatus UTEX 2973. Proc Natl Acad Sci USA 115: E11761–E11770

Google Scholar: Author Only Title Only Author and Title

Vasudevan R, Gale GAR, Schiavon AA, Puzorjov A, Malin J, Gillespie MD, Vavitsas K, Zulkower V, Wang B, Howe CJ, et al (2019) CyanoGate: a modular cloning suite for engineering cyanobacteria based on the plant MoClo syntax. Plant Physiol 180: 39–55

Google Scholar: Author Only Title Only Author and Title

Vento JM, Crook N, Beisel CL (2019) Barriers to genome editing with CRISPR in bacteria. J Ind Microbiol Biotechnol 46: 1327–1341

Google Scholar: Author Only Title Only Author and Title

Vogel AIM, Lale R, Hohmann-Marriott MF (2017) Streamlining recombination-mediated genetic engineering by validating three neutral integration sites in Synechococcus sp. PCC 7002. J Biol Eng 11: 19

Google Scholar: Author Only Title Only Author and Title

Wang S-Y, Li X, Wang S-G, Xia P-F (2023) Base editing for reprogramming cyanobacterium Synechococcus elongatus. Metab Eng 75: 91–99

Google Scholar: Author Only Title Only Author and Title

Wendt KE, Ungerer J, Cobb RE, Zhao H, Pakrasi HB (2016) CRISPR/Cas9 mediated targeted mutagenesis of the fast growing cyanobacterium Synechococcus elongatus UTEX 2973. Microb Cell Factories 15: 115

Google Scholar: Author Only Title Only Author and Title

Wilson KS, von Hippel PH (1995) Transcription termination at intrinsic terminators: the role of the RNA hairpin. Proc Natl Acad Sci USA 92: 8793–8797

Google Scholar: Author Only Title Only Author and Title

Włodarczyk A, Selão TT, Norling B, Nixon PJ (2020) Newly discovered Synechococcus sp. PCC 11901 is a robust cyanobacterial strain for high biomass production. Commun Biol 3: 1–14

Google Scholar: Author Only Title Only Author and Title

Xiao K, Yue X-H, Chen W-C, Zhou X-R, Wang L, Xu L, Huang F-H, Wan X (2018) Metabolic engineering for enhanced medium chain omega hydroxy fatty acid production in Escherichia coli. Front Microbiol 9: 139

Google Scholar: Author Only Title Only Author and Title

Yao L, Shabestary K, Björk SM, Asplund-Samuelsson J, Joensson HN, Jahn M, Hudson EP (2020) Pooled CRISPRi screening of the cyanobacterium Synechocystis sp PCC 6803 for enhanced industrial phenotypes. Nat Commun 11: 1666

Google Scholar: Author Only Title Only Author and Title

Yu J, Liberton M, Cliften PF, Head RD, Jacobs JM, Smith RD, Koppenaal DW, Brand JJ, Pakrasi HB (2015) Synechococcus elongatus UTEX 2973, a fast growing cyanobacterial chassis for biosynthesis using light and CO2. Sci Rep 5: 8132

Google Scholar: Author Only Title Only Author and Title

Zavřel T, Chmelík D, Sinetova MA, Červený J (2018) Spectrophotometric determination of phycobiliprotein content in cyanobacterium Synechocystis. J Vis Exp 58076

Google Scholar: Author Only Title Only Author and Title

Zess EK, Begemann MB, Pfleger BF (2016) Construction of new synthetic biology tools for the control of gene expression in the cyanobacterium Synechococcus sp. strain PCC 7002. Biotechnol Bioeng 113: 424–432

Google Scholar: Author Only Title Only Author and Title

Zetsche B, Gootenberg JS, Abudayyeh OO, Slaymaker IM, Makarova KS, Essletzbichler P, Volz SE, Joung J, van der Oost J, Regev A, et al (2015) Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system. Cell 163: 759–771

Google Scholar: Author Only Title Only Author and Title

Zhang A, Carroll AL, Atsumi S (2017) Carbon recycling by cyanobacteria: improving CO2 fixation through chemical production. FEMS Microbiol Lett 364: fnx165

Google Scholar: Author Only Title Only Author and Title

Zhou Y, Sun T, Chen Z, Song X, Chen L, Zhang W (2019) Development of a new biocontainment strategy in model cyanobacterium Synechococcus strains. ACS Synth Biol 8: 2576–2584

Google Scholar: Author Only Title Only Author and Title