Intrinsic control of DRG sensory neuron diversification by Pten

Alejandra Fernandez[1‡], Nick Sarn[2], Charis Eng[2,3,4], Kevin M. Wright[1,‡]

[1]Vollum Institute, Oregon Health & Science University, Portland, OR, 97239

[2]Genomic Medicine Institute, Lerner Research Institute, Cleveland Clinic, Cleveland, OH 44195

[3]Center for Personalized Genetic Healthcare, Medical Specialties Institute, Cleveland Clinic, Cleveland, OH 44195

[4]Department of Genetics and Genome Sciences, Case Western Reserve University School of Medicine, Cleveland, OH 44106

[‡]Corresponding Authors: fernaale@ohsu.edu; wrighke@ohsu.edu

Summary

Phosphatase and tensin homolog (PTEN) modulates intracellular survival and differentiation signaling pathways downstream of neurotrophin receptors in the developing peripheral nervous system (PNS). Although well-studied in the context of brain development, our understanding of the *in vivo* role of PTEN in the PNS is limited to models of neuropathic pain and nerve injury. Here, we assessed how alterations in PTEN signaling affects the development of peripheral somatosensory circuits. We found that sensory neurons within the dorsal root ganglia (DRG) in *Pten* heterozygous (*Pten^Het*) mice exhibit defects in neuronal subtype diversification. Abnormal DRG differentiation in *Pten^Het* mice arises early in development, with subsets of neurons

expressing both progenitor and neuronal markers. DRGs in *Pten^{Het}* mice show dysregulation of both mTOR and GSK-3β signaling pathways downstream of PTEN. Finally, we show that mice with an autism-associated mutation in *Pten* (*Pten^{Y68H/+}*) show abnormal DRG development. Thus, we have discovered a crucial role for PTEN signaling in the intrinsic diversification of primary sensory neuron populations in the DRG during development.

Running Title

PTEN regulates DRG diversification

Keywords:

PTEN, somatosensory neuron, neuronal differentiation, diversification, DRG, Autism Spectrum Disorder

Introduction

The development of neural circuits depends on tightly regulated signaling pathways that guide the differentiation and diversification of progenitors to establish populations of mature neuronal subtypes [1-7]. Many neurodevelopmental disorders are thought to arise from alterations in neuronal progenitor maintenance and disruptions in the balance between proliferation and differentiation in the central nervous system (CNS) [8]. In contrast, little attention has been given to how altered development of the peripheral nervous system (PNS) contributes to neurodevelopmental disorders.

Neurons in the dorsal root ganglia (DRG) play a critical role in relaying sensory information to the CNS. During embryonic development, neural crest cells (NCCs) delaminate from the dorsal neural tube, migrate laterally, and coalesce to form the nascent DRG. These NCCs are bi-potential progenitors that possess the ability to self-renew and generate both neurons and glia [9]. Through asymmetric divisions, some DRG progenitors exit the cell cycle to initially give rise to transcriptionally diverse populations of sensory neurons, while a subset remains in a proliferative state to subsequently generate satellite glia [10]. There are at least 14 distinct neuronal subtypes in the DRG, each one conveying a specific sensory modality [1, 4, 11, 12]. Neuronal subtype diversity within DRGs is initially established during embryogenesis in overlapping developmental waves of neurogenesis. As postmitotic cells acquire a neuronal fate, they begin to express specific tropomyosin receptor kinase (*Trk*) receptors [13, 14]. In mice, large and medium diameter neurons expressing the neurotrophin receptors TrkC or TrkB are generated first, between embryonic day 9.5 (E9.5) and E11.5, and ultimately differentiate into proprioceptors and

mechanoreceptors, respectively [7, 14, 15]. The small diameter neurons that give rise

to nociceptors initially express TrkA and are generated between E10.5 and E13.5 [11,

15]. Neurotrophin signaling through Trk receptors is required for multiple aspects of

DRG neuron development, including neuronal survival, axonal outgrowth, target

innervation, and neuronal maturation [13, 16].

While the intrinsic molecular pathways downstream of Trk receptors are shared

by most DRG neuron populations, how these signals are integrated to regulate fate-

specification in sensory neurons *in vivo* is poorly understood [16-18]. The

Phosphatidylinositol 3 kinase/protein kinase B/mammalian Target of Rapamycin

(PI3K/AKT/mTOR) pathway is one of the main survival and differentiation signaling

pathways initiated downstream of Trk receptors in sensory neurons [13, 19-21].

Phosphatase and tensin homolog (PTEN) is the primary negative regulator of this

pathway, counteracting PI3K activity [22-24]. Dysfunctional or impaired PTEN activity

leads to increased AKT and mTOR activation [25]. In humans, germline heterozygous

mutations in *PTEN* or other components of the PI3K/AKT/mTOR pathway have been

linked to cancer and neurodevelopmental disorders, including autism spectrum

disorders (ASD) [26-30]. Moreover, PTEN has been previously shown to have

neurodevelopmental roles in mouse models [31]. In the developing CNS, PTEN

promotes cell cycle exit and the transition to post-mitotic neuronal fates [32, 33].

Heterozygous mutations of *Pten* during development lead to increased cell proliferation

and increased production of glial cells [34, 35]. PTEN is also involved in neuronal

differentiation [36]. In post-mitotic neurons, PTEN plays a role in axon outgrowth,

guidance and branching patterns, and overall neuronal growth [37-40]. When *Pten* is

conditionally deleted in neurons, they have hypertrophic cell somas and dendrites in the

cerebellum, cortex, and hippocampus [41-43]. At synapses, PTEN regulates dendritic

spine density and connectivity [42, 44-46]. Within the PNS, PTEN is involved in growth

cone collapse, restricts axon regeneration, and regulates myelin thickness [39, 47, 48].

However, despite its well-known function regulating signaling pathways downstream of

Trk receptor activation, how alterations in PTEN activity affect neuronal differentiation

and population diversity in DRG neurons *in vivo* remains unclear. Moreover, whether

heterozygosity of *Pten*, which more accurately models neurodevelopmental disorders, is

sufficient to elicit phenotypes in the PNS remains unknown.

In this study, we examined DRG development in mice lacking a single copy of

wildtype *Pten (Pten^{Het})*. We find that adult *Pten^{Het}* mice have a normal number of DRG

neurons, but that population diversification is dramatically altered. These alterations are

cell-intrinsic and arise during early stages of neuronal differentiation, with nascent DRG

neurons remaining in an intermediate state, expressing both progenitor and neuronal

markers simultaneously. We find that both the mTOR and GSK-3β pathways

downstream of Akt are dysregulated in *Pten^{Het}* DRG neurons. Finally, we show that

mice harboring a heterozygous ASD-associated mutation that shows predominant

nuclear localization (*Pten$^{Y68H/+}$*) also show alterations in DRG diversification [49-51].

Taken together, these results show that tight regulation and proper dosage of PTEN

activity is critical for the proper development and diversification of somatosensory

neurons *in vivo,* and identify a key contribution of *Pten*-mediated signaling to the

development of primary sensory neurons in the context of syndromic ASD.

Results

DRG population diversity is altered in *Pten^{Het}* mice

Recent studies have identified altered development and function of primary DRG neurons in several mouse models of syndromic ASD, including *Mecp2^{-/y}*, *Gabrb3^{+/-}*, *Shank3^{+/-}* and *Fmr1^{-/y}* mutants [52-55]. Additionally, single copy mutations in several genes in the PI3K/Akt/mTOR pathway are associated with ASD, and this pathway plays a critical role downstream of Trk receptors [16, 56, 57]. While several studies have focused on the role of PTEN in DRG neurons in the adult or in the context of axon regeneration, how the loss of *Pten* affects DRG development has not been studied *in vivo*. To test the hypothesis that disruptions in this pathway affect DRG development, we examined how global loss of a single *Pten* allele (*Pten^{Het}*) affects somatosensory neuron development. We collected hindlimb-innervating DRGs (lumbar level 3-4; L3-L4, **Figure 1A**) from P21 mice and compared the number of neurons in *Pten^{Het}* mutants and wild type (WT; *Pten^{+/+}*) littermates using Islet1/2 (ISL1) as a neuronal marker. There was no difference in the total number of ISL1$^+$ neurons in the DRG of adult *Pten^{Het}* mutants compared to WT littermate controls (**Figure 1B**). In contrast, when we examined DRGs using subtype-specific markers (**Figure 1C**), we identified alterations in neuronal diversification in *Pten^{Het}* mutants. The number of TrkA$^+$ peptidergic nociceptors was decreased by 27% in *Pten^{Het}* mutants relative to control littermates (**Figure 1D**), whereas IB4$^+$ non-peptidergic nociceptors were unchanged (**Figure 1E**). In contrast, the Tyrosine Hydroxylase positive (TH+) C-low-threshold mechanoreceptor (C-LTMR) population was increased by 156% (**Figure 1F**). There was no significant change in the

number of TrkB$^+$ Aδ-LTMRs or Calbindin$^+$ Aβ-rapidly adapting LTMRs (RA-LTMRs

(**Figure 1G, H**). There was also a 56% increase in the number of Parvalbumin positive

(PV$^+$) neurons, which includes proprioceptors and populations of low-threshold

mechanoreceptors (LTMRs) (**Figure 1I**). Together, these data show that a

physiologically optimal expression level of *Pten* is critical for DRG neuron diversification.

Control of DRG population diversification by *Pten* is neuron-autonomous

Our examination of *Pten^{Het}* mice identified a role for PTEN in DRG differentiation

and population diversification. To test whether this reflected a neuron-autonomous role

for PTEN, we asked whether conditional deletion of *Pten* from primary sensory neurons

would be sufficient to recapitulate the phenotypes in constitutive *Pten^{Het}* mice. We

generated transgenic mice carrying: 1: a *Cre* driver for DRG neurons (*Isl1^{Cre}*); 2: a

Pten^{flox} allele. We quantified population subtypes in DRGs from *Isl1^{Cre};Pten$^{fl/+}$* animals

and *Isl1^{Cre};Pten$^{+/+}$* littermate controls at P21. In line with our results in constitutive

Pten^{Het} animals, we found that single allele deletion of *Pten* from DRG neurons resulted

in altered subtype diversification. There was a 42% decrease in the number of TrkA$^+$

peptidergic nociceptors, and no change in the number of IB4$^+$ non-peptidergic

nociceptors (**Figure S1A, C, G**). Moreover, we observed a 99% increase in the number

of TH$^+$ c-LTMRs in *Pten^{Het}* mutants (**Figure S1B, G**). The number of TrkB+ Aδ-LTMRs

and Calbindin+ Aβ RA-LTMR was not affected in mutants, while PV$^+$ neurons were

increased by 58% (**Figure S1D, E, F, G**). These data show that PTEN signaling

functions within DRG neurons to control their specification and population diversity.

***Pten* is essential for regulating neuronal subtype specification during early**

development

To resolve when heterozygous deletion of *Pten* disrupts adult DRG population

diversity, we next investigated whether these defects arose early in development or

whether they emerged in adult animals. During embryonic development, large/medium

diameter neurons expressing TrkC and/or TrkB are generated first, followed by TrkA

expressing small diameter neurons [15]. Together, these neurons ultimately give rise to

the full complement of genetically and physiologically distinct subtypes of

somatosensory neurons (**Figure 2A**). When we examined the proportions of TrkA, TrkB,

and TrkC positive neurons between E10.5 and E14.5, we found a significant decrease

in the number of TrkA$^+$ neurons in *Pten^{Het}* mice compared to WT littermates at all ages

examined (**Figure 2B**). In contrast, the proportion of TrkC$^+$ neurons was significantly

increased between E11.5-E14.5 (**Figure 2D**). In line with our results in adult mice,

which did not detect any changes in the proportion of neurons from the TrkB lineage, we

saw no change in the number of TrkB$^+$ neurons between E11.5-E14.5 (**Figure 2C**).

Therefore, PTEN plays a critical role in the early differentiation of TrkA$^+$ and TrkC$^+$ DRG

neurons.

To better define the role of PTEN during DRG neuron development, we assessed

PTEN's cellular localization by immunostaining DRGs of WT mice between E9.5 and

E14.5. At E9.5, when the nascent DRG is primarily comprised of mitotic precursors,

PTEN levels are relatively low (**Figure 3A**). PTEN levels sharply increased at E10.5,

when DRG neurons are exiting the cell cycle, and remained elevated through E14.5

(**Figure 3A**) [58, 59]. We next triple-labeled developing DRGs with PTEN and ISL1 to

mark newly differentiated neurons, as well as SOX10 to label DRG precursors and glial

cells. At all ages examined, PTEN levels were significantly higher in ISL1$^+$ neurons

compared to SOX10$^+$ cells (**Figure 3B, C**), suggesting that PTEN plays a role in newly

generated DRG neurons after they exit the cell cycle.

DRG neuron proliferation and apoptosis are normal in the DRG of *Pten*^{Het} mice

The final number of DRG neurons in the adult organism are regulated by a

balance of progenitor proliferation and developmental cell death. We therefore used

EdU birthdating to examine whether abnormal progenitor proliferation contributed to the

altered number of TrkA$^+$ and TrkC$^+$ neurons observed in *Pten*^{Het} mutants. We

administered EdU to pregnant dams at a range of timepoints from E9.5 to E14.5 to mark

cells undergoing terminal mitosis at the time of injection. We harvested embryos 2 hours

after injection and quantified the proportion of Sox10$^+$ progenitors that were EdU

positive in DRG sections [60]. There was no difference in the number of EdU$^+$/SOX10$^+$

cells at any of the time points examined (**Figure 4A**), suggesting that progenitor

proliferation is normal in *Pten*^{Het} mutants.

Target-derived neurotrophins binding to Trk receptors activate PI3K/AKT

signaling, which regulates pro-survival pathways to ensure the proper number of DRG

neurons [6, 61, 62]. Since PTEN functions as a negative regulator of this pathway [63],

we expected that we would see increased neuronal survival in mice lacking a single

Pten allele. Somewhat surprisingly, when we examined activated Caspase-3 in DRGs at

a range of timepoints that correspond to the window during which most developmental

cell death occurs (E10.5-E14.5), we saw no difference between *Pten*^{Het} mice and WT

littermates (**Figure 4B**). Taken together, these data indicate that proliferation and

developmental cell death are largely normal in *Pten^Het* mice, and are unlikely to

significantly contribute to the alterations in DRG subtype diversification.

Pten heterozygous deletion alters the specification of DRG neurons

Since alterations in DRG subtype populations in *Pten^Het* mice do not appear to

arise from abnormal proliferation, we carefully examined the transition of neural

progenitors to postmitotic neurons. As progenitors in the DRG exit the cell cycle and

differentiate into neurons, SOX10 is downregulated and ISL1 levels increase, resulting

in largely mutually exclusive expression [9, 64-66]. When we quantified the proportion

of SOX10$^+$ progenitors and ISL1$^+$ neurons in WT and *Pten^Het* animals, we observed that

proportion of SOX10$^+$ progenitors was slightly elevated and the number of ISL1$^+$

neurons were slightly reduced in *Pten^Het* mice at E12.5 and E13.5, and were not

statistically different at any other timepoints (**Figure 5A, B**). However, we observed that

while SOX10 and ISL1 are mostly mutually exclusive in WT animals as previously seen

[67], there was a significant increase in the proportion of cells co-expressing both

progenitor (SOX10) and neuronal (ISL1) markers at every timepoint examined in *Pten^Het*

animals (**Figure 5A, C**). These results suggest that altering PTEN levels results in

abnormal neuronal differentiation in DRG sensory neurons.

Intracellular signaling pathways downstream of *Pten* are altered in developing

sensory neurons in *Pten^Het* mutants

We next sought to determine whether specific signaling pathways downstream of

PTEN were altered in *Pten* heterozygous mutants. PTEN negatively regulates

PI3K/AKT-mediated signaling, which activates both the TSC/mTOR/S6 pathway and the

GSK-3β/β-catenin pathway [34, 63, 68] (**Figure 6A**). Since *Pten* haploinsufficiency has

different effects on TrkA, TrkB, and TrkC populations within developing DRGs, we

examined key readouts downstream of each pathway in developing DRGs by

immunostaining for pS6 and using a β-catenin reporter *TCF/Lef:H2B-GFP* mouse

crossed onto a *Pten^Het* background (**Figure 6**) [69]. We examined levels of pS6 in

cryosections of *Pten^Het* and WT littermates at E11.5, and observed that the number of

pS6$^+$/ISL1$^+$ neurons in DRGs of *Pten^Het* embryos was increased (**Figure 6B**). When we

co-labeled neurons with antibodies to pS6 and specific Trk receptors, we found that

both TrkA$^+$ (**Figure 6C**) and TrkC$^+$ (**Figure 6E**) neurons showed a significant increase in

the proportion of pS6$^+$ neurons. Surprisingly, the proportion of TrkB$^+$ neurons that were

pS6$^+$ was unchanged in *Pten^Het* embryos (**Figure 6D**).

We next used GFP fluorescence in the *TCF/Lef:H2B-GFP* mouse as a readout of

activation of the GSK-3β/β-catenin signaling pathway [34, 69]. The number of GFP$^+$

cells that showed activation of β-catenin signaling in E11.5 DRGs was significantly

increased in ISL1$^+$ neurons (**Figure 6F**). Examination of population specific markers

showed no significant change in the number of TrkA$^+$/GFP$^+$ and TrkB$^+$/GFP$^+$ neurons,

and a 46% increase in the number of GFP+ neurons within the TrkC$^+$ population

(**Figure 6G-I**).

DRG neuron diversification is abnormal in mice with an ASD-associated mutation

in *Pten*.

Given the prevalence of *PTEN* mutations within the ASD population and the high

incidence of sensory phenotypes in ASD [26, 70], we decided to determine if DRG

subtype diversification was impacted by mutations observed in individuals with ASD.

The ASD-associated Y68H mutation (*Pten*$^{Y68H/+}$) results in reduced protein stability and

nuclear predominant PTEN subcellular localization in CNS neurons [51, 71]. Similar to

our results in *Pten*^{Het} mice, *Pten*$^{Y68H/+}$ mice had no change in the total number of DRG

neurons, but did have significant alterations in the proportions of specific subtypes

(**Figure 7**). Quantification revealed a 50% decrease in the TrkA$^+$ peptidergic nociceptive

population, a 79% increase in TH$^+$ C-LTMRs and a 96% increase in PV$^+$ neurons in

Pten$^{Y68H/+}$ mice when compared to WT littermates (**Figure 7A, C**). There were no

changes in the number of IB4$^+$ non-peptidergic nociceptors, TrkB$^+$ Aδ-LTMRs, or

Calbindin$^+$ Aβ RA-LTMRs. These numbers are in concordance with our observations in

Pten^{Het} mice, and suggest that signaling pathways downstream of cytoplasmic PTEN

plays a predominant role in regulating DRG subtype diversification.

Discussion

Pten functions as the primary regulator of PI3K/AKT signaling downstream of

neurotrophin receptor activation, however little is known about its role in cell-fate

determination and differentiation pathways. Here, we identified a critical developmental

role for *Pten* in modulating cell-fate determination in TrkA and TrkC lineages of primary

sensory neurons in the DRG.

DRG development and the role of *Pten*

Primary sensory neurons within DRGs differentiate from neural crest progenitors that delaminate from the dorsal neural tube [7]. Given that there are at least 14 molecularly and functionally distinct somatosensory neuron subtypes in adult DRGs, a central question of ongoing investigation is how this neuronal diversity arises during development. Recent work has begun to trace the developmental trajectories and gene expression patterns of each developing population of DRG neurons [3, 4, 72]. As soon as immature DRG neurons exit the cell cycle, they progress through hierarchical steps of specification driven by specific gene expression programs, following specific chronological sequences [1, 3, 4, 73]. The diversity of sensory modalities emerges early in embryonic development through two overlapping waves of sensory neurogenesis that occur between E9.5 and E13 in mice [1, 3, 7, 15, 73]. The first wave (E9.5-E11.5) gives rise to myelinated, medium- and large-diameter neurons which will eventually differentiate into mechanoreceptors, proprioceptors, and a small population of nociceptors [15]. Mechanoreceptors express RET and MAFA and/or TrkB [74], proprioceptors express tropomyosin receptor kinase C (TrkC) and RUNX3 [72], and the small Aδ-nociceptor population expresses TrkA [15, 75, 76]. A second wave of neurogenesis (E10.5-E12.5) generates unmyelinated small-diameter nociceptive neurons [7, 77]. These neurons co-express TrkA and Runx1 and later diversify into pain, itch, thermosensation as well as gentle and, presumably, affective touch [77-79].

The identity of each DRG subtype relies on their expression of specific transmembrane receptors and transcription factors [4, 12]. Recent work suggests that

counteracting gene-regulatory networks operate within immature postmitotic DRG

neurons to establish the transcriptional programs defining neuron subtype-specific fate

choice [3]. Extrinsic cues are also necessary for DRG cell-fate determination and

transcriptional control of subtype identity [4]. Our understanding of how these intrinsic

and extrinsic signaling pathways are integrated and the cellular mechanisms underlying

the hierarchical fate-decisions leading to molecular specificity in each DRG neuronal

subpopulation remain incomplete. Our data provide evidence that precise regulation of

Pten signaling plays a critical role in neuronal diversification in DRG neurons. Loss of a

single *Pten* allele (*Pten^Het^*) or incorporation of an ASD-associated mutation in *Pten*

(*Pten^Y68H/+^*) leads to significant alterations in the numbers of TrkC- and TrkA-lineage

derived populations. These defects are evident as early as E10.5, as developing DRG

neurons are acquiring specific fates and persist into adulthood, where the resulting

mature neuronal subtypes are affected. Moreover, we show that these defects are

modulated intrinsically within the DRG neuron population (*Isl1^Cre^;Pten^Flox/+^*). These

results raise a host of new questions, including: 1. *How is Pten signaling contributing to*

cell-fate decisions during development? 2. *Are neurons in Pten^Het^ and Pten^Y68H/+^ mice*

only failing to express specific markers, or are these neurons switching fates? 3. *Does*

altered Pten signaling lead to neurons that are either unspecified or stalled in

intermediate states? Further studies using single cell RNA sequencing will help address

these questions, and provide additional insight into how reduced *Pten* signaling affects

DRG neuron development and function.

***Pten* regulates neuronal specification broadly throughout the developing nervous**

system

The current model of neuronal specification in the DRG suggests that nascent

neurons are initially unspecified and express a broad array of transcription factors, and

progressively acquire molecular specificity through a mechanism of hierarchical binary

fate decisions [3, 4]. In other neuronal circuits, *Pten* plays a role in neuronal

specification, controlling progenitor cell proliferation to maintain ratios of populations in

several lineages [80, 81]. In the retina, *Pten* regulates cell division and is required to

produce the full complement of rod photoreceptors and amacrine cells [82]. Moreover,

in cortical GABAergic neurons, conditional deletion of *Pten* results in a preferential loss

of somatostatin (SST$^+$) interneurons, leading to an altered ratio of PV$^+$/SST$^+$

interneurons [36]. Whether this results from changes in transcriptional control of cell-fate

determination remains unknown. However, in both retina and developing brain, AKT

signaling was found to be increased. Moreover, conditional deletion of the Tuberous

Sclerosis Complex (TSC) gene, *Tsc1*, which functions downstream of PTEN and inhibits

mTOR, causes ectopic PV expression and altered physiological properties in SST$^+$

cortical interneurons [83]. While these data demonstrate a role for TSC/mTOR signaling

events in mediating interneuron differentiation, additional studies have also identified a

contribution of the GSK-3β/β-Catenin pathway to cellular organization in the cortex [34].

One particularly relevant study showed overactivation of the GSK-3β/β-Catenin and not

the mTOR pathway in the developing cortex of *Pten* heterozygous conditional mutants.

GSK-3β/β-Catenin overactivation resulted in altered neuronal differentiation and

excessive gliogenesis in *Pten*^{Het} mice [34]. In contrast, homozygous mutants showed

overactivation of both the GSK-3β and the mTOR pathway, and lowering mTOR activity

was not sufficient to rescue the phenotypes in *Pten^Het* mice [34]. More recently, a

transcriptomic analysis revealed that, during development, cellular populations within

the cerebral cortex have different requirements of PTEN, as heterozygous deletion of

Pten affects downstream signaling differently in astrocytes and neurons [84]. In our

current study, we found that both mTOR and GSK-3β/β-Catenin pathways are

hyperactive in TrkC$^+$ DRG neurons in *Pten$^{+/-}$* mutants, whereas only pS6 was

significantly elevated in TrkA$^+$ DRG neurons. This could reflect differences in

downstream signaling in the different populations, or could reflect a lag between β-

catenin activity and reporter expression in newly generated TrkA$^+$ neurons at E11.5 [80].

Along with the different effects of diminished *Pten* signaling in each population

subtype derived from the TrkA and TrkC lineages, we saw no changes in population

diversification in neurons derived from the TrkB lineage. This particularity of the TrkB

lineage may result from a difference in the intrinsic neuronal response to neurotrophins

during development. Brain-derived neurotrophin factor (BDNF) signaling through TrkB is

qualitatively different than signaling through TrkA and TrkC, and can be regulated by co-

factors, modulators, and transactivators including G protein-coupled receptors (GPCRs)

and cyclic adenosine monophosphate (cAMP) [85, 86]. Therefore, it is possible that

other pathways, along with the PI3K/AKT, are engaged during population diversification

within the TrkB lineage, or at least, to modulate the expression of the surface proteins

used to identify each population. In this case, either neuronal subtypes derived from the

TrkB lineage are unaffected in *Pten^Het* mutants, or the phenotypes in these populations

are undetectable in our assays.

Pten regulates gene expression in DRG neurons

A number of studies have examined the consequence of suppressing Pten

activity in adult DRGs in the context of axon regeneration following injury [87-91]. While

these experiments show that loss of PTEN can promote axon regeneration, the majority

of these studies have not examined how this affects the maintenance of DRG neuron

identity. However, a recent study showed that inducible deletion of *Pten* from adult

DRGs under non-injury conditions results in pathological itch and elevated responses to

pruritogens and thermal stimulation [92]. This abnormal sensory response is due in part

to ectopic expression of *TRPV1* and *MrgprA3*, particularly within the IB4$^+$ population.

Simultaneous conditional deletion of *Pten* and *TrpV1* restored itch and thermal

responses back to baseline. These results suggest that in addition to the role we have

identified for PTEN in regulating DRG subtype differentiation during development,

sustained PTEN levels are also required in a subset of adult DRG neurons to maintain

their proper molecular identity and function. This prior study only examined the

expression of a subset of itch-related genes, so it is not clear what other molecular

changes occur in the absence of *Pten* in adult neurons. It is also unknown whether

deleting a single allele of *Pten* from adult DRGs is sufficient to alter neuron identity

and/or function, as this previous study did not report results from heterozygous *Pten^{cKO}*

mice [92].

Tuberous Sclerosis Complex 2 (*Tsc2*) functions to negatively modulate mTOR

activity downstream of Pten. Conditional deletion of *Tsc2* from developing nociceptors

leads to defective innervation of both central and peripheral targets and reduced heat

sensitivity and injury-induced cold hypersensitivity [93, 94]. While the overall number of

neurons was normal in $Tsc2^{cKO}$ mice, there were significant alterations to neuronal

subtype populations and gene expression profiles in the DRG. Similar to our results in

$Pten^{Het}$ and $Pten^{Y68H/+}$ mice, they reported a significant decrease in TrkA$^+$ neurons in

$Tsc2^{cKO}$ mice. They also observed an increase in IB4$^+$ neurons, which we did not see in

our $Pten$ mutant mice. This difference may be due to the timing of $Tsc2$ deletion

(perinatal), or due to the loss of both $Tsc2$ alleles, compared to our analysis of $Pten$

heterozygous mice. Among the genes that showed altered expression following $Tsc2$

deletion, ion channels implicated in DRG neuron function were particularly affected.

Whether this reflects abnormal differentiation of neuronal subtypes, or abnormal

maintenance of adult subtype identity is unclear.

Implications of altered $Pten$ signaling during DRG development in autism

spectrum disorders (ASD)

In addition to the core features of ASD, which include social and communication

deficits and repetitive/restrictive interests, it is estimated that up to 90 percent of

individuals with ASD experience some form of atypical sensory sensitivity [70]. Recent

work has begun to characterize how mutations in ASD-susceptibility genes affect

sensory circuit development and/or function [52, 54, 95-100]. Rats with mutations in

$Mecp2$ show abnormal mechanical and thermal sensitivity due to abnormal sensory

neuron development and function during postnatal development [55]. Moreover,

deletion of $Mecp2$ or $Gabrb3$ in somatosensory neurons during development results in

tactile deficits, defects in social interactions, and anxiety-like behaviors, all of which are

associated with ASD [52]. Pharmacologically reducing tactile overactivity in *Mecp2* and

Shank3 mutants reduced anxiety-like behaviors and some social deficits, further

establishing the link between PNS-specific sensory defects and ASD phenotypes [54].

Finally, a recent study has emphasized the link between the developmental timing of

somatosensory dysfunction and emergence of ASD-like behaviors [53]. Our data

showing that primary sensory deficits in $Pten^{Het}$ mutants arise early during development

suggest that these defects may contribute to later behavioral abnormalities. Notably, we

see the same defects in DRG neuron diversification in both $Pten^{Het}$ and $Pten^{Y68H/+}$ mice.

These results show that loss of a single allele of Pten is sufficient to cause defects in

DRG diversification, and that this arises predominantly due to the cytosolic signaling

function of PTEN. This is supported by the observation that both TSC/mTOR and GSK-

3β/β-Catenin signaling pathways are dysregulated in $Pten^{Het}$ DRGs. PTEN also has

several functions within the nucleus, and whether these play a role in regulating DRG

development remains to be determined. In future studies, the complex cellular functions

of PTEN could be genetically dissected using mouse models with mutations affecting

distinct functional aspects of PTEN. Furthermore, parsing out how the developmental

defects in primary sensory neurons contribute to the wider range of ASD-like

phenotypes in $Pten^{Het}$ mutants could be done using conditional deletion of a single copy

of *Pten* specifically in DRG neurons.

Acknowledgements

We would like to thank the members of the Wright laboratory for their assistance and

discussion throughout the course of this study. Special thanks to Dr. Matthew Pomaville

for his insight and help with experimental troubleshooting. We also thank Dr. Martin

Riccomagno for his helpful suggestions on the manuscript. This work was supported by

NIH grant R01NS091027 (K.M.W.), OHSU Fellowship for Diversity in Research (A.F.),

Collins Medical Trust (A.F.) and NIH grant K01NS116168 (A.F.). Confocal microscopy

and analysis was performed in the OHSU Advanced Light Microscopy Core, supported

by the NIH grant P30NS061800. C.E. is the Sondra J and Stephen R Hardis Chair of

Cancer Genomic Medicine at the Cleveland Clinic and would like to thank the Ambrose

Monell Foundation.

Materials and Methods

Mouse lines. All experiments were performed in accordance to OHSU IACUC Protocol

#IS00000539. Animals were housed in standard conditions and cared for by the

Department of Comparative Medicine at Oregon Health & Science University. Mice

were maintained in a 12-hour light/dark cycle, and provided food and water *ad libitum*.

For experiments involving embryonic timepoints, noon on the date that a vaginal plug

was detected was considered E0.5. For experiments involving constitutive *Pten*

heterozygous mice, male *Pten^{Het}* breeders (*B6;129P2-Pten^{tm1Mak}/Mmjax* [101]) were

crossed with CD1 females to generate experimental mice on a mixed genetic

background. For conditional deletion experiments, *Pten$^{Flox/+}$* mice (*Pten^{tm1Hwu}/J* [102])

were crossed to *Isl1^{Cre}* mice (*Isl1$^{tm1(cre)}$Sev/J* [103]). To examine activation of the GSK-

3β/β-Catenin pathway, *Pten^{Het}* breeders were crossed to a *TCF/Lef:H2B-EGFP* reporter

line (*Tg(TCF/Lef1-HIST1H2BB/EGFP)61Hadj/J* [69]). All animals were group housed

with littermates and maintained on a mixed genetic background. Mice of both sexes

were used for all experiments, and wildtype littermates were used as controls.

Proliferation assays. A single intraperitoneal (IP) injection of EdU in sterile PBS (10μg

EdU/g body weight) was given to pregnant dams on one day between E9.5 and E14.5.

Two hours after injection, embryos were collected and fixed as described below.

Tissue processing and immunolabeling. All immunolabeling was performed on

cryosections following standard protocols [105]. Whole mouse embryos, from litters

ranging from E9.5 to E14.5, were fixed overnight at 4°C in 4% paraformaldehyde with

gentle agitation. For analysis of P21 DRGs, intact spinal columns were first dissected

and fixed overnight at 4°C in 4% paraformaldehyde. The next day, spinal columns were

washed in PBS for 30 minutes at room temperature, and spinal cords and attached

DRGs were dissected from the spinal column for further processing. All tissue was

cryoprotected by incubating in 15% sucrose in PBS for 1 hour, followed by 20% sucrose

in PBS overnight at 4°C with gentle agitation. Samples were mounted in OCT and

rapidly frozen in methylbutane on dry ice. 20 μm consecutive sections were cut on a

Leica CM3050S cryostat, mounted on slides, and allowed to dry for 2 hours at room

temperature.

For immunolabeling, cryosections mounted on slides were washed 3 times for 5

minutes in PBS at room temperature and incubated for 1 hour in blocking solution (5%

v/v Normal Donkey Serum, 5% v/v DMSO, 0.25% v/v Triton X-100 in PBS). Slides were

then incubated in primary antibodies diluted in blocking solution overnight at 4°C in a

humidified staining box. Concentrations for primary antibodies are listed on Table 1.

Slides were then washed three times for 5 minutes in PBS, and incubated for 2 hours at

room temperature in secondary antibodies diluted at 1:500 in blocking solution. EdU

incorporation was detected using Click-iT™ EdU Cell Proliferation Kit for Imaging, Alexa

Fluor™ 647 (Cat#C10340). Slides were incubated in EdU detection solution (100 µM

Tris pH7.5, 4 mM $CuSO_4$, 5 µM sulfo-Cy5 azide, 100 mM sodium ascorbate) for 30 min

at RT, protected from light. Slides were washed 3 times for 10 minutes in PBS, with

1:5000 Hoescht added to the second wash to label nuclei. Slides were then

coverslipped with Flourmount-G and imaged.

Cell counts and statistical analysis. To quantify immunoreactive DRG neurons, level-

matched DRGs from lumbar segments were serially sectioned. For analysis of total

number of neurons, sections of entire DRGs were counted. For population proportions,

cell proliferation, and cell death assays, 10 consecutive sections of level-matched DRGs

were counted. Sections were co-immunostained with antibodies against specific cell

makers and Isl1/2. The proportions of each population were calculated by dividing the

number of marker-positive cells over the number of total neurons in serial sections from

level-matched DRGs of control and mutant mice. Values were averaged for each

experiment, with a minimum of 3 animals of mixed sex for each genotype at each

timepoint and condition across different litters. The average ± SEM is reported in the

figures. Statistical analysis for each experiment was performed by Student's t-test with

post-hoc Tukey test. P-values are reported in figure legends and results section.

Table 1: Antibodies used:

Antibody	Species	Supplier	Cat #	RRID	Concentration
Primary Antibodies					
Calbindin	Rabbit	Swant	CB 38	AB_10000340	1:1000
Tyrosine Hydroxylase (TH)	Rabbit	Millipore	AB152	AB_390204	1:1000
TrkA	Goat	Fisher (R&D Systems)	AF1056	AB_2283049	1:100
TrkB	Goat	Fisher (R&D Systems)	AF1494	AB_2155264	1:100
TrkC	Goat	Fisher (R&D Systems)	AF1404	AB_2155412	1:100
Casp3	Rabbit	Cell Signaling	9661S	AB_2341188	1:500
Parvalbumin	Rabbit	Swant	PVG213	AB_2721207	1:1000
Isl1/2	Mouse Igg2b	DHSB	39.4D5	AB_2314683	1:250
GFP	Chicken	Abcam	ab13970	AB_300798	1:1000
Secondary Antibodies					
Anti-Rabbit IgG 488	Donkey	ThermoFisher	A-21206		1:500

Anti-Rabbit IgG 546	Donkey	ThermoFisher	A10040		1:500
Anti-Rabbit IgG 647	Donkey	ThermoFisher	A-31573		1:500
Anti-Mouse Igg2b 488	Goat	ThermoFisher	A-21141		1:500
Anti-Chicken IgY 488	Donkey	Jackson Immuno Research	NC0215979		1:500
Anti-Goat 647	Donkey	ThermoFisher	A-21447		1:500

References:

1. Marmigere, F. and P. Ernfors, *Specification and connectivity of neuronal subtypes in the sensory lineage.* Nat Rev Neurosci, 2007. **8**(2): p. 114-27.
2. Meltzer, S., C. Santiago, N. Sharma, and D.D. Ginty, *The cellular and molecular basis of somatosensory neuron development.* Neuron, 2021. **109**(23): p. 3736-3757.
3. Faure, L., et al., *Single cell RNA sequencing identifies early diversity of sensory neurons forming via bi-potential intermediates.* Nat Commun, 2020. **11**(1): p. 4175.
4. Sharma, N., et al., *The emergence of transcriptional identity in somatosensory neurons.* Nature, 2020. **577**(7790): p. 392-398.
5. Li, C.L., et al., *Somatosensory neuron types identified by high-coverage single-cell RNA-sequencing and functional heterogeneity.* Cell Res, 2016. **26**(1): p. 83-102.
6. Snider, W.D. and I. Silos-Santiago, *Dorsal root ganglion neurons require functional neurotrophin receptors for survival during development.* Philos Trans R Soc Lond B Biol Sci, 1996. **351**(1338): p. 395-403.
7. Ma, Q., C. Fode, F. Guillemot, and D.J. Anderson, *Neurogenin1 and neurogenin2 control two distinct waves of neurogenesis in developing dorsal root ganglia.* Genes Dev, 1999. **13**(13): p. 1717-28.
8. Meechan, D.W., et al., *Modeling a model: Mouse genetics, 22q11.2 Deletion Syndrome, and disorders of cortical circuit development.* Prog Neurobiol, 2015. **130**: p. 1-28.
9. Liu, J.A., et al., *Fbxo9 functions downstream of Sox10 to determine neuron-glial fate choice in the dorsal root ganglia through Neurog2 destabilization.* Proc Natl Acad Sci U S A, 2020. **117**(8): p. 4199-4210.
10. Mapps, A.A., et al., *Diversity of satellite glia in sympathetic and sensory ganglia.* Cell Rep, 2022. **38**(5): p. 110328.
11. Liu, Y. and Q. Ma, *Generation of somatic sensory neuron diversity and implications on sensory coding.* Curr Opin Neurobiol, 2011. **21**(1): p. 52-60.
12. Usoskin, D., et al., *Unbiased classification of sensory neuron types by large-scale single-cell RNA sequencing.* Nat Neurosci, 2015. **18**(1): p. 145-53.
13. Huang, E.J. and L.F. Reichardt, *Trk receptors: roles in neuronal signal transduction.* Annu Rev Biochem, 2003. **72**: p. 609-42.
14. Mu, X., I. Silos-Santiago, S.L. Carroll, and W.D. Snider, *Neurotrophin receptor genes are expressed in distinct patterns in developing dorsal root ganglia.* J Neurosci, 1993. **13**(9): p. 4029-41.
15. Landy, M.A., M. Goyal, and H.C. Lai, *Nociceptor subtypes are born continuously over DRG development.* Dev Biol, 2021. **479**: p. 91-98.
16. Huang, E.J. and L.F. Reichardt, *Neurotrophins: roles in neuronal development and function.* Annu Rev Neurosci, 2001. **24**: p. 677-736.
17. Gorokhova, S., et al., *Uncoupling of molecular maturation from peripheral target innervation in nociceptors expressing a chimeric TrkA/TrkC receptor.* PLoS Genet, 2014. **10**(2): p. e1004081.
18. Segal, R.A. and M.E. Greenberg, *Intracellular signaling pathways activated by neurotrophic factors.* Annu Rev Neurosci, 1996. **19**: p. 463-89.

19. Reichardt, L.F., *Neurotrophin-regulated signalling pathways.* Philos Trans R Soc Lond B Biol Sci, 2006. **361**(1473): p. 1545-64.

20. Postigo, A., et al., *Distinct requirements for TrkB and TrkC signaling in target innervation by sensory neurons.* Genes Dev, 2002. **16**(5): p. 633-45.

21. York, R.D., et al., *Role of phosphoinositide 3-kinase and endocytosis in nerve growth factor-induced extracellular signal-regulated kinase activation via Ras and Rap1.* Mol Cell Biol, 2000. **20**(21): p. 8069-83.

22. Philpott, K.L., M.J. McCarthy, A. Klippel, and L.L. Rubin, *Activated phosphatidylinositol 3-kinase and Akt kinase promote survival of superior cervical neurons.* J Cell Biol, 1997. **139**(3): p. 809-15.

23. Song, W., et al., *ProNGF induces PTEN via p75NTR to suppress Trk-mediated survival signaling in brain neurons.* J Neurosci, 2010. **30**(46): p. 15608-15.

24. Kreis, P., G. Leondaritis, I. Lieberam, and B.J. Eickholt, *Subcellular targeting and dynamic regulation of PTEN: implications for neuronal cells and neurological disorders.* Front Mol Neurosci, 2014. **7**: p. 23.

25. Stambolic, V., et al., *Negative regulation of PKB/Akt-dependent cell survival by the tumor suppressor PTEN.* Cell, 1998. **95**(1): p. 29-39.

26. Tilot, A.K., T.W. Frazier, 2nd, and C. Eng, *Balancing Proliferation and Connectivity in PTEN-associated Autism Spectrum Disorder.* Neurotherapeutics, 2015. **12**(3): p. 609-19.

27. Milella, M., et al., *PTEN: Multiple Functions in Human Malignant Tumors.* Front Oncol, 2015. **5**: p. 24.

28. Buxbaum, J.D., et al., *Mutation screening of the PTEN gene in patients with autism spectrum disorders and macrocephaly.* Am J Med Genet B Neuropsychiatr Genet, 2007. **144B**(4): p. 484-91.

29. Goffin, A., et al., *PTEN mutation in a family with Cowden syndrome and autism.* Am J Med Genet, 2001. **105**(6): p. 521-4.

30. Butler, M.G., et al., *Subset of individuals with autism spectrum disorders and extreme macrocephaly associated with germline PTEN tumour suppressor gene mutations.* J Med Genet, 2005. **42**(4): p. 318-21.

31. Skelton, P.D., R.V. Stan, and B.W. Luikart, *The Role of PTEN in Neurodevelopment.* Mol Neuropsychiatry, 2020. **5**(Suppl 1): p. 60-71.

32. Sun, H., et al., *PTEN modulates cell cycle progression and cell survival by regulating phosphatidylinositol 3,4,5,-trisphosphate and Akt/protein kinase B signaling pathway.* Proc Natl Acad Sci U S A, 1999. **96**(11): p. 6199-204.

33. Lachyankar, M.B., et al., *A role for nuclear PTEN in neuronal differentiation.* J Neurosci, 2000. **20**(4): p. 1404-13.

34. Chen, Y., et al., *Pten Mutations Alter Brain Growth Trajectory and Allocation of Cell Types through Elevated beta-Catenin Signaling.* J Neurosci, 2015. **35**(28): p. 10252-67.

35. Tilot, A.K., et al., *Germline disruption of Pten localization causes enhanced sex-dependent social motivation and increased glial production.* Hum Mol Genet, 2014. **23**(12): p. 3212-27.

36. Vogt, D., et al., *The parvalbumin/somatostatin ratio is increased in Pten mutant mice and by human PTEN ASD alleles.* Cell Rep, 2015. **11**(6): p. 944-956.

37. Huang, W.C., Y. Chen, and D.T. Page, *Hyperconnectivity of prefrontal cortex to amygdala projections in a mouse model of macrocephaly/autism syndrome.* Nat Commun, 2016. **7**: p. 13421.

38. LaSarge, C.L., V.R. Santos, and S.C. Danzer, *PTEN deletion from adult-generated dentate granule cells disrupts granule cell mossy fiber axon structure.* Neurobiol Dis, 2015. **75**: p. 142-50.

39. Chadborn, N.H., et al., *PTEN couples Sema3A signalling to growth cone collapse.* J Cell Sci, 2006. **119**(Pt 5): p. 951-7.

40. Henle, S.J., L.P. Carlstrom, T.R. Cheever, and J.R. Henley, *Differential role of PTEN phosphatase in chemotactic growth cone guidance.* J Biol Chem, 2013. **288**(29): p. 20837-20842.

41. Backman, S.A., et al., *Deletion of Pten in mouse brain causes seizures, ataxia and defects in soma size resembling Lhermitte-Duclos disease.* Nat Genet, 2001. **29**(4): p. 396-403.

42. Kwon, C.H., et al., *Pten regulates neuronal soma size: a mouse model of Lhermitte-Duclos disease.* Nat Genet, 2001. **29**(4): p. 404-11.

43. Kwon, C.H., et al., *Pten regulates neuronal arborization and social interaction in mice.* Neuron, 2006. **50**(3): p. 377-88.

44. Fraser, M.M., I.T. Bayazitov, S.S. Zakharenko, and S.J. Baker, *Phosphatase and tensin homolog, deleted on chromosome 10 deficiency in brain causes defects in synaptic structure, transmission and plasticity, and myelination abnormalities.* Neuroscience, 2008. **151**(2): p. 476-88.

45. Williams, M.R., et al., *Hyperactivity of newborn Pten knock-out neurons results from increased excitatory synaptic drive.* J Neurosci, 2015. **35**(3): p. 943-59.

46. Xiong, Q., H.V. Oviedo, L.C. Trotman, and A.M. Zador, *PTEN regulation of local and long-range connections in mouse auditory cortex.* J Neurosci, 2012. **32**(5): p. 1643-52.

47. Christie, K.J., et al., *PTEN inhibition to facilitate intrinsic regenerative outgrowth of adult peripheral axons.* J Neurosci, 2010. **30**(27): p. 9306-15.

48. Cotter, L., et al., *Dlg1-PTEN interaction regulates myelin thickness to prevent damaging peripheral nerve overmyelination.* Science, 2010. **328**(5984): p. 1415-8.

49. Lobo, G.P., et al., *Germline and somatic cancer-associated mutations in the ATP-binding motifs of PTEN influence its subcellular localization and tumor suppressive function.* Hum Mol Genet, 2009. **18**(15): p. 2851-62.

50. He, X., et al., *Naturally occurring germline and tumor-associated mutations within the ATP-binding motifs of PTEN lead to oxidative damage of DNA associated with decreased nuclear p53.* Hum Mol Genet, 2011. **20**(1): p. 80-9.

51. Sarn, N., S. Thacker, H. Lee, and C. Eng, *Germline nuclear-predominant Pten murine model exhibits impaired social and perseverative behavior, microglial activation, and increased oxytocinergic activity.* Mol Autism, 2021. **12**(1): p. 41.

52. Orefice, L.L., et al., *Peripheral Mechanosensory Neuron Dysfunction Underlies Tactile and Behavioral Deficits in Mouse Models of ASDs.* Cell, 2016. **166**(2): p. 299-313.

53. Tasnim, A., et al., *The developmental timing of spinal touch processing alterations and its relation to ASD-associated behaviors in mouse models.* bioRxiv, 2023: p. 2023.05.09.539589.

54. Orefice, L.L., et al., *Targeting Peripheral Somatosensory Neurons to Improve Tactile-Related Phenotypes in ASD Models.* Cell, 2019. **178**(4): p. 867-886 e24.

55. Bhattacherjee, A., et al., *Neuronal cytoskeletal gene dysregulation and mechanical hypersensitivity in a rat model of Rett syndrome.* Proc Natl Acad Sci U S A, 2017. **114**(33): p. E6952-E6961.

56.   Segal, R.A., *Selectivity in neurotrophin signaling: theme and variations.* Annu Rev Neurosci, 2003. **26**: p. 299-330.

57.   Thomas, S.D., N.K. Jha, S. Ojha, and B. Sadek, *mTOR Signaling Disruption and Its Association with the Development of Autism Spectrum Disorder.* Molecules, 2023. **28**(4).

58.   Lawson, S.N. and T.J. Biscoe, *Development of mouse dorsal root ganglia: an autoradiographic and quantitative study.* J Neurocytol, 1979. **8**(3): p. 265-74.

59.   Venteo, S., et al., *Neurog2 Deficiency Uncovers a Critical Period of Cell Fate Plasticity and Vulnerability among Neural-Crest-Derived Somatosensory Progenitors.* Cell Rep, 2019. **29**(10): p. 2953-2960 e2.

60.   Serinagaoglu, Y., J. Pare, M. Giovannini, and X. Cao, *Nf2-Yap signaling controls the expansion of DRG progenitors and glia during DRG development.* Dev Biol, 2015. **398**(1): p. 97-109.

61.   Klesse, L.J. and L.F. Parada, *p21 ras and phosphatidylinositol-3 kinase are required for survival of wild-type and NF1 mutant sensory neurons.* J Neurosci, 1998. **18**(24): p. 10420-8.

62.   Datta, S.R., et al., *Akt phosphorylation of BAD couples survival signals to the cell-intrinsic death machinery.* Cell, 1997. **91**(2): p. 231-41.

63.   Georgescu, M.M., *PTEN Tumor Suppressor Network in PI3K-Akt Pathway Control.* Genes Cancer, 2010. **1**(12): p. 1170-7.

64.   Zhang, Q., et al., *Temporal requirements for ISL1 in sympathetic neuron proliferation, differentiation, and diversification.* Cell Death Dis, 2018. **9**(2): p. 247.

65.   McKeown, S.J., et al., *Sox10 overexpression induces neural crest-like cells from all dorsoventral levels of the neural tube but inhibits differentiation.* Dev Dyn, 2005. **233**(2): p. 430-44.

66.   Sonnenberg-Riethmacher, E., et al., *Development and degeneration of dorsal root ganglia in the absence of the HMG-domain transcription factor Sox10.* Mech Dev, 2001. **109**(2): p. 253-65.

67.   Sun, Y., et al., *A central role for Islet1 in sensory neuron development linking sensory and spinal gene regulatory programs.* Nat Neurosci, 2008. **11**(11): p. 1283-93.

68.   Huang, W., et al., *GSK3 beta mediates suppression of cyclin D2 expression by tumor suppressor PTEN.* Oncogene, 2007. **26**(17): p. 2471-82.

69.   Ferrer-Vaquer, A., et al., *A sensitive and bright single-cell resolution live imaging reporter of Wnt/ss-catenin signaling in the mouse.* BMC Dev Biol, 2010. **10**: p. 121.

70.   Ben-Sasson, A., et al., *A meta-analysis of sensory modulation symptoms in individuals with autism spectrum disorders.* J Autism Dev Disord, 2009. **39**(1): p. 1-11.

71.   Han, S.Y., et al., *Functional evaluation of PTEN missense mutations using in vitro phosphoinositide phosphatase assay.* Cancer Res, 2000. **60**(12): p. 3147-51.

72.   Dietrich, S., et al., *Molecular identity of proprioceptor subtypes innervating different muscle groups in mice.* Nat Commun, 2022. **13**(1): p. 6867.

73.   Lallemend, F. and P. Ernfors, *Molecular interactions underlying the specification of sensory neurons.* Trends Neurosci, 2012. **35**(6): p. 373-81.

74.   Bourane, S., et al., *Low-threshold mechanoreceptor subtypes selectively express MafA and are specified by Ret signaling.* Neuron, 2009. **64**(6): p. 857-70.

75.   Lawson, S.N., X. Fang, and L. Djouhri, *Nociceptor subtypes and their incidence in rat lumbar dorsal root ganglia (DRGs): focussing on C-polymodal nociceptors, Abeta-*

nociceptors, moderate pressure receptors and their receptive field depths. Curr Opin Physiol, 2019. **11**: p. 125-146.

76. Fang, X., et al., *trkA is expressed in nociceptive neurons and influences electrophysiological properties via Nav1.8 expression in rapidly conducting nociceptors.* J Neurosci, 2005. **25**(19): p. 4868-78.

77. Marmigere, F., et al., *The Runx1/AML1 transcription factor selectively regulates development and survival of TrkA nociceptive sensory neurons.* Nat Neurosci, 2006. **9**(2): p. 180-7.

78. Luo, W., et al., *A hierarchical NGF signaling cascade controls Ret-dependent and Ret-independent events during development of nonpeptidergic DRG neurons.* Neuron, 2007. **54**(5): p. 739-54.

79. Keeler, A.B., et al., *A developmental atlas of somatosensory diversification and maturation in the dorsal root ganglia by single-cell mass cytometry.* Nat Neurosci, 2022. **25**(11): p. 1543-1558.

80. Sun, C., et al., *PTEN regulation of the proliferation and differentiation of auditory progenitors through the PTEN/PI3K/Akt-signaling pathway in mice.* Neuroreport, 2014. **25**(3): p. 177-83.

81. Lyu, J., et al., *The protein phosphatase activity of PTEN is essential for regulating neural stem cell differentiation.* Mol Brain, 2015. **8**: p. 26.

82. Tachibana, N., et al., *Pten Regulates Retinal Amacrine Cell Number by Modulating Akt, Tgfbeta, and Erk Signaling.* J Neurosci, 2016. **36**(36): p. 9454-71.

83. Malik, R., et al., *Tsc1 represses parvalbumin expression and fast-spiking properties in somatostatin lineage cortical interneurons.* Nat Commun, 2019. **10**(1): p. 4994.

84. Cheung, S.K.K., et al., *Neuropathological signatures revealed by transcriptomic and proteomic analysis in Pten-deficient mouse models.* Sci Rep, 2023. **13**(1): p. 6763.

85. Andreska, T., P. Luningschror, and M. Sendtner, *Regulation of TrkB cell surface expression-a mechanism for modulation of neuronal responsiveness to brain-derived neurotrophic factor.* Cell Tissue Res, 2020. **382**(1): p. 5-14.

86. Mitre, M., et al., *Transactivation of TrkB Receptors by Oxytocin and Its G Protein-Coupled Receptor.* Front Mol Neurosci, 2022. **15**: p. 891537.

87. Danilov, C.A. and O. Steward, *Conditional genetic deletion of PTEN after a spinal cord injury enhances regenerative growth of CST axons and motor function recovery in mice.* Exp Neurol, 2015. **266**: p. 147-60.

88. Willenberg, R., et al., *Variable laterality of corticospinal tract axons that regenerate after spinal cord injury as a result of PTEN deletion or knock-down.* J Comp Neurol, 2016. **524**(13): p. 2654-76.

89. Park, K.K., et al., *Promoting axon regeneration in the adult CNS by modulation of the PTEN/mTOR pathway.* Science, 2008. **322**(5903): p. 963-6.

90. Liu, K., et al., *PTEN deletion enhances the regenerative ability of adult corticospinal neurons.* Nat Neurosci, 2010. **13**(9): p. 1075-81.

91. Chandran, V., et al., *A Systems-Level Analysis of the Peripheral Nerve Intrinsic Axonal Growth Program.* Neuron, 2016. **89**(5): p. 956-70.

92. Hu, L., et al., *The role of PTEN in primary sensory neurons in processing itch and thermal information in mice.* Cell Rep, 2022. **39**(3): p. 110724.

93. Carlin, D., et al., *Deletion of Tsc2 in Nociceptors Reduces Target Innervation, Ion Channel Expression, and Sensitivity to Heat.* eNeuro, 2018. **5**(2).

94. Abe, N., et al., *Mammalian target of rapamycin (mTOR) activation increases axonal growth capacity of injured peripheral nerves.* J Biol Chem, 2010. **285**(36): p. 28034-43.

95. Price, T.J. and O.K. Melemedjian, *Fragile X mental retardation protein (FMRP) and the spinal sensory system.* Results Probl Cell Differ, 2012. **54**: p. 41-59.

96. He, C.X., et al., *Tactile Defensiveness and Impaired Adaptation of Neuronal Activity in the Fmr1 Knock-Out Mouse Model of Autism.* J Neurosci, 2017. **37**(27): p. 6475-6487.

97. Chelini, G., et al., *Aberrant Somatosensory Processing and Connectivity in Mice Lacking Engrailed-2.* J Neurosci, 2019. **39**(8): p. 1525-1538.

98. Deng, P.Y., O. Avraham, V. Cavalli, and V.A. Klyachko, *Hyperexcitability of Sensory Neurons in Fragile X Mouse Model.* Front Mol Neurosci, 2021. **14**: p. 796053.

99. Han, Q., et al., *SHANK3 Deficiency Impairs Heat Hyperalgesia and TRPV1 Signaling in Primary Sensory Neurons.* Neuron, 2016. **92**(6): p. 1279-1293.

100. Dawes, J.M., et al., *Immune or Genetic-Mediated Disruption of CASPR2 Causes Pain Hypersensitivity Due to Enhanced Primary Afferent Excitability.* Neuron, 2018. **97**(4): p. 806-822 e10.

101. Suzuki, A., et al., *High cancer susceptibility and embryonic lethality associated with mutation of the PTEN tumor suppressor gene in mice.* Curr Biol, 1998. **8**(21): p. 1169-78.

102. Lesche, R., et al., *Cre/loxP-mediated inactivation of the murine Pten tumor suppressor gene.* Genesis, 2002. **32**(2): p. 148-9.

103. Yang, L., et al., *Isl1Cre reveals a common Bmp pathway in heart and limb development.* Development, 2006. **133**(8): p. 1575-85.

104. Madisen, L., et al., *A robust and high-throughput Cre reporting and characterization system for the whole mouse brain.* Nat Neurosci, 2010. **13**(1): p. 133-40.

105. Pomaville, M.B. and K.M. Wright, *Follicle-innervating Adelta-low threshold mechanoreceptive neurons form receptive fields through homotypic competition.* Neural Dev, 2023. **18**(1): p. 2.

Figure 1. Population diversity in DRGs is altered in *Pten^Het* mice. A) Schematic representation of spinal cord with attached DRGs. All quantifications were done in L4 DRGs in P21 animals. **B)** Images and quantification of total number of neurons using ISL1 as a marker. Total number of neurons was the same in WT and *Pten^Het* mutants (n=3, p=0.57, Student's t-test). **C)** Schematic of DRG primary sensory neuron diversity subpopulations of proprioceptors, mechanoreceptors and nociceptors are labeled with each of the markers outlined in schematic. **D-I)** Representative images of cryosections of WT and *Pten^Het* adult DRGs with specific markers (left panels) and quantification (right panels). Inserts show magnification of squared area. **D)** TrkA positive peptidergic nociceptors are decreased in *Pten^Het* mutants (n=3, p<0.002, Student's t-test). **E)** Quantification of IB4-positive non-peptidergic nociceptors (right panel) shows no change in *Pten^Het* mutants compared to WT controls (n=3, p=0.20, Student's t-test). **F)** Tyrosine hydroxylase (TH) positive LTMRs are increased in *Pten^Het* mutants compared to WT littermates (n=3, p<0.05, Student's t-test). **G-H)** There were no significant changes in the number of TrkB-lineage mechanoreceptors (TrkB+ Aδ-LTMRs:, n=3, p=0.34, Student's t-test; Calbindin+ Aβ RA-LTMR: n=3, p=0.41, Student's t-test). **I)** Parvalbumin positive (PV+) proprioceptors were increased in *Pten^Het* mutants compared to WT littermates (n=3 p<0.002, Student's t-test).

Figure S1: Control of population diversification by *Pten* is neuron-autonomous.

A-F) Representative images of cryosections of DRGs from P21 *Isl1^{Cre};Pten$^{fl/+}$* and control littermates with specific markers. **G)** Quantification of proportions of each population subtypes in *Isl1^{Cre};Pten$^{fl/+}$* normalized to control. TrkA+ peptidergic nociceptors are reduced in *Isl1^{Cre};Pten$^{fl/+}$* mutants (n=3 p<0.015, Student's t-test). IB4+ non-peptidergic nociceptors were unchanged in mutants when compared to controls (n=3, p=0.69, Student's t-test).TH+ c-LTMRs show an increase in *Isl1^{Cre};Pten$^{fl/+}$* mutants compared to controls (n=3, p<0.001, Student's t-test). There were no significant changes in the number of TrkB+ or Calb1+ mechanoreceptors (TrkB: n=3, p=0.83; Calb1: n=3, p=0.49, Student's t-test). PV+ neurons showed an increase in *Isl1^{Cre};Pten$^{fl/+}$* mutants compared to controls (n=3 p<0.05, Student's t-test).

Figure 2: Neuronal population diversity is altered in *Pten^Het^* mutants throughout development. A) Schematic of DRG primary sensory neuron diversification during development. Neural crest cells (NCCs) differentiate and diversify into subpopulations that give rise to proprioceptors, mechanoreceptors, and nociceptors; schematic shows markers for specific subpopulations. **B-D)** Representative images of cryosections of WT and *Pten^Het^* E11.5 DRGs with markers of each subtype (left panels) and quantification (right panels). **B)** Representative images of DRGs labeled with TrkA (left), quantification (right) shows a decrease in the number of TrkA positive neurons in *Pten^Het^* mutants throughout development (n=3, E10.5: $p<0.004$; E11.5 : $p<0.03$; E12.5: $p<0.02$, E13.5: $p<0.045$, E14.5: $p<0.03$, Student's t-test). **C)** Images showing DRGs labeled with TrkB (left), quantification (right) shows no changes the number of TrkB+ neurons in *Pten^Het^* mutants (n=3, E10.5: $p=0.46$; E11.5: $p=0.32$; E12.5: $p=0.45$; E13.5: $p=0.47$; E14.5: $p=0.28$, Student's t-test). **D)** Images of E11.5 DRGs labeled with TrkC (left), quantification (right) shows an increase in the proportion of TrkC+ neurons in *Pten^Het^* mutants after E10.5 (E10.5: $p=0.14$; E11.5 $p<0.02$; E12.5: $p<0.05$; E13.5: $p<0.05$; E14.5: $p<0.01$, Student's t-test).

Figure 3: Pten levels are elevated in newly generated DRG neurons. A)
Representative images of embryonic DRGs (left to right from E9.5-14.5) of WT animals labeled with PTEN by immunohistochemistry. At E9.5, when nascent DRGs are primarily compromised of mitotic precursors, PTEN levels are relatively low. PTEN levels rise sharply at E10.5 and E11.5, as the majority of DRG neurons differentiate. **B)** Representative images of embryonic DRGs (left to right, from E10.5 to E14.5) labeled by immunofluorescence with antibodies against PTEN, Islet1/2 (Isl1/2) and Sox10. Arrows indicate Sox10+ progenitors, which are not Isl1/2+ or Pten+. **C)** Quantification of the average intensity levels of PTEN within Isl1/2+ neurons and Sox10+ positive precursors. At all ages, PTEN levels were significantly higher in Isl1/2+ neurons compared to Sox10+ progenitors (n=3, E10.5: $p<0.03$; E11.5: $p<0.001$; E12.5: $p<0.02$; E13.5: $p<0.01$; E14.5: $p<0.02$; Student's t-test).

Figure 4: Proliferation and developmental cell death are normal in *Pten^Het*

mutants. A) Representative images of E11.5 DRGs from WT and *Pten^Het* littermates

birth-dated with EdU and labeled with Sox10 immunostaining (left panels).

Quantification (right panel) shows percentage of Sox10+ progenitors undergoing

proliferation at each developmental time point (E9.5-E14.5). At all ages quantified,

there is no difference in the percentage of EdU+;Sox10+ precursors in *Pten^Het*

compared to WT controls (n=3, E9.5: p=0.055; E10.5: p=0.11; E11.5: p=0.45; E12.5:

p=0.06; E13.5: p=0.484; E14.5: p=0.481, at E14.5; Student's t-test). **B)** Representative

images from E11.5 DRGs of WT and *Pten^Het* littermates showing activated Caspase 3

(Casp3; green) and Islet 1/2 immunoreactivity (left panels). Quantification (right panel)

shows percentage of Isl1/2+ neurons co-labeled with activated Caspase 3. At all ages

quantified (E10.5- E14.5), there is no difference in the percentage of activated

Casp3+;Islet1+ neurons in *Pten^Het* DRGs compared to WT controls (n=3, E10.5: p=0.54;

E11.5: p=0.16; at E12.5: p=0.069; at E13.5: p=0.91; at E14.5: p=0.84, Student's t-test).

Figure 5: Progression of neuronal progenitors to neurons is altered in *Pten^Het*mutants. **A)** Representative images of E12.5 DRGs from WT and *Pten^Het* littermates co-stained with markers of neuronal progenitors (Sox10) and neurons (Isl1/2). **B)** Quantification of proportions of neuronal precursors (left) and neurons (right) within DRGs of WT and *Pten^Het* littermates at different developmental ages from E9.5 to E14.5. At E12.5 and E13.5 there was a small significant increase in the proportion of Sox10+ progenitors and a corresponding decrease in the proportion of Isl1/2+ neurons in *Pten^Het* mutants (n=3, E9.5: p=0.13; E10.5: p=0.48; E11.5: p=0.45; E12.5: p<0.04; E13.5: p<0.03; E14.5: p=0.28, Student's t-test). **C)** Quantification of the proportion of cells co-labeled with markers for neuronal precursors (Sox10) and neurons (Isl1/2) from E9.5 to E14.5. The proportion of cells that were both Sox10+ and Isl1/2+ increased dramatically in *Pten^Het* mice compared to WT littermates at each timepoint examined (n=3, E9.5: p<0.02; E10.5: p<0.04; E11.5: p<0.001; E12.5: p<0.001; E13.5: p<0.001; E14.5: p<0.04, Student's t-test).

Figure 6: Altered signaling downstream of PI3K/AKT in TrkA and TrkC lineages in

Pten^{Het} **mutants. A)** Schematic representation of mTOR and GSK-3β signaling

pathways downstream of PI3K/AKT. **B-E)** Representative images (left) and

quantification (right) of E11.5 DRGs of WT and *Pten^{Het}* littermates immunolabeled with

Isl1 (B) TrkA (C) TrkB (D) or TrkC (E) and phospho-S6 (pS6) to measure activity of the

PI3K/AKT/mTOR pathway. **B)** Quantification (right) shows an increase in the proportion

of pS6+/Isl1+ neurons in *Pten^{Het}* compared to WT (n=3, p<0.041, Student's t-test). **C)**

Within the TrkA population, pS6+/TrkA+ neurons were increased in *Pten^{Het}* compared to

WT littermates (n=3, p<0.02, Student's t-test). **D)** Quantification in TrkB+ neurons (right)

shows no significant change in the proportions of pS6+/TrkB+ neurons between *Pten^{Het}*

and WT controls (n=3, p=0.82, Student's t-test). **E)** Quantification shows increased

number of pS6+/TrkC+ neurons in *Pten^{Het}* compared to WT littermates (n=3, p<0.03,

Student's t-test). **F-I)** Representative images (left) and quantification (right) of E11.5

DRGs from *TCF/Lef:H2B-GFP;Pten^{Het}* DRGs and control littermates immunolabeled

with Isl1 (F) TrkA (H) TrkB (H) or TrkC (I) to measure activity of the GSK-3β/β-catenin

pathway. **F)** The proportion of GFP+/Isl1+ neurons was increased in *TCF/Lef:H2B-*

GFP;Pten^{Het} when compared to controls (n=3, p<0.005, Student's t-test). **G)** Within the

TrkA population, there was no significant change in the number of GFP+/TrkA+ neurons

between *TCF/Lef:H2B-GFP;Pten^{Het}* and *TCF/Lef:H2B-GFP;Pten^{WT}* controls (n=3,

p=0.11, Student's t-test). **H)** There was no significant change in the proportion of

GFP+/TrkB+ neurons between *TCF/Lef:H2B-GFP;Pten^{Het}* and *TCF/Lef:H2B-*

GFP;Pten^{WT} WT controls (p=0.96, Student's t-test). **I)** The TrkC lineage shows a 127%

increase in GFP+/TrkC+ neurons in *TCF/Lef:H2B-GFP;Pten^Het* compared to

TCF/Lef:H2B-GFP;Pten^WT controls (n=3, p<0.002, Student's t-test).

Figure 7: DRG neuron diversification is abnormal in *Pten^Y68H/+* mice. A) Representative images of cryosections of WT (top row) and *Pten^Y68H* (bottom row) P21 DRGs labeled with specific markers for select DRG subtypes. Inserts show magnification of squared area. **B)** The total number of Isl1/2+ DRG neurons was not altered in *Pten^Y68H* mutants compared to WT controls (n=3 p=0.18, Student's t-test). **C)** Quantification shows that the number of TrkA+ peptidergic nociceptors is decreased in *Pten^Y68H* mutants (n=3 p<0.04, Student's t-test), whereas TH+ c-LTMRs and PV+ neurons are increased in *Pten^Y68H* mutants (TH: n=3 p<0.005, PV: n=3 p<0.02, Student's t-test). There was no significant difference in the number of IB4+ non-peptidergic nociceptors, TrkB+ Aδ-LTMRs, or Calb1+ Aβ-LTMRs in *Pten^Y68H* mutants compared to WT littermates (IB4: n=3 p=0.09, TrkB: n=3 p=0.47; Calb1: n=3 p=0.38, Student's t-test).

Figure 1

Figure S1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7