Using human genetics to develop strategies to increase erythropoietic output from

genome-edited hematopoietic stem and progenitor cells

Joab Camarena[1,2]*, Sofia E. Luna[1,2]*, Jessica P. Hampton[1,2], Kiran R. Majeti[1,2], Carsten

T. Charlesworth[1,2], Eric Soupene[3], Vivien A. Sheehan[4], M. Kyle Cromer[5,6,7]†, & Matthew

H. Porteus[1,2]†

[1]Department of Pediatrics, Stanford University, Stanford, CA.

[2]Institute for Stem Cell Biology and Regenerative Medicine, Stanford University Stanford,

CA.

[3]Department of Pediatrics, University of California, San Francisco, Oakland, CA.

[4]Aflac Cancer Center and Blood Disorders, Department of Pediatrics, Emory University

School of Medicine, Atlanta, GA.

[5]Department of Surgery, University of California, San Francisco, San Francisco, CA.

[6]Department of Bioengineering & Therapeutic Sciences, University of California, San

Francisco, San Francisco, CA, USA.

[7]Eli & Edythe Broad Center for Regeneration Medicine, University of California, San

Francisco, San Francisco, CA, USA.

*Contributed equally

†Correspondence: kyle.cromer@uscf.edu, mporteus@stanford.edu

Abstract

Human genetic polymorphisms result in a diversity of phenotypes. Some sequences are pathologic and lead to monogenic diseases, while others may confer beneficial traits. Genome editing is a powerful tool to recreate genotypes found in the population, including the ability to correct pathologic mutations. One interesting naturally occurring mutation is a truncation in the erythropoietin receptor (tEPOR) causing congenital erythrocytosis— characterized by non-pathogenic hyper-production of red blood cells (RBCs). Using the precision of CRISPR/Cas9 genome editing, we have recreated tEPOR and studied the effect of variations of the genotype on RBC development. We then combined tEPOR with a correction strategy developed for β-thalassemia and demonstrated that coupling the two genome editing events gave RBCs a significant selective advantage. This demonstrates the potential of combining human genetics with the precision of genome editing to enable safer and more effective genome editing therapies for patients with serious genetic diseases.

Main

Many of the initial applications of clinical genome editing have aimed to correct or compensate for disease-causing mutations of monogenic diseases. However, human genetic variation is more nuanced than monogenic diseases, as there are variants that appear to confer positive health benefits as well. Examples are people that have bi-allelic deletions in *CCR5* causing resistance to HIV infection, a variety of polymorphisms causing upregulation of fetal hemoglobin, and mutations in *PCSK9* causing low cholesterol levels[1–3]. It is important, however, to broadly assess variations that occur in only small numbers of people for both their potential risks as well as for their potential benefits.

Congenital erythrocytosis (CE) is a rare phenotype in which people have higher than normal levels of red blood cells and consequently elevated hemoglobin. The genetic cause of this phenotype was identified by studying the family of a Finnish Olympic gold medal-winning cross-country skier who was found to have levels of hemoglobin >50% higher than normal[4]. It is caused by truncations in the erythropoietin receptor (tEPOR) in which the intracellular inhibitory domain to erythropoietin (EPO) signaling is eliminated[4,5]. This domain contains binding sites for SHP1 which normally leads to downregulation of EPO-dependent JAK2/STAT5 signaling (Extended Data Figure 1a). Further studies have shown that tEPOR does not create a constitutively active EPOR signaling cascade, but rather imparts hypersensitivity to EPO[5]. As a consequence, these kindreds with tEPOR typically present with abnormally low levels of EPO, indicating a new homeostasis is attained to prevent CE from becoming pathogenic. Finally, studies of families with CE show that it is not pre-disposing to cancer and is thus considered a benign trait[6].

Genome editing is a powerful method that enables the precise changing of nucleotides in the DNA of a cell. There are multiple genome-editing strategies including nuclease-based insertion/deletion (indel) formation, base editing, and prime editing, but the most versatile approach to genome-editing is homology-directed repair (HDR). In HDR, a nuclease-induced double-strand break (DSB) is repaired using a donor template. The natural donor template for HDR is the sister chromatid and the natural repair pathway is homologous recombination. By providing a donor template that resembles a sister chromatid with large homology arms flanking the intended cut site, the homologous recombination machinery can use this "substitute" sister chromatid to repair the DSB. HDR editing is the most flexible because it can create single nucleotide changes, precisely insert large gene cassettes, and even swap out large genomic regions for other gene sequences[7–10]. We utilize all of these applications of HDR in this work.

One of the challenges in hematopoietic stem cell gene therapy is to achieve sufficient engraftment of the genetically engineered cells to have a beneficial clinical effect without increasing risk. To make this possible, effort is exerted to maximize editing frequencies in hematopoietic stem and progenitor cells (HSPCs)[11–13]. Even if clinically relevant editing frequencies are achieved, high-morbidity chemo-therapeutic regimens are currently required to create niche space in the bone marrow (BM) for these edited HSPCs, which can create toxicities, including oncogenic risk[14–16]. In this work, we aimed to give edited cells a selective advantage such that low levels of engraftment might still result in a clinical benefit, perhaps enabling less toxic conditioning, through use of genome editing to recreate the CE phenotype by engineering tEPOR into HSPCs in different ways. We find that when tEPOR is engineered into human HSPCs using

genome-editing, there is a significant selective advantage to the derived red blood cells

(RBCs). We then show that this selective advantage can be coupled to a therapeutic gene

edit to give the cells with the therapeutic edit a selective advantage in RBC development,

without affecting the stem and progenitor cells. In this way, we demonstrate the power of

combining human genetics with precision genome editing to potentially enable safer and

more effective genome editing therapies for patients with serious genetic diseases,

particularly those of red blood cells.

Results

Cas9-guided EPOR truncation in HSPCs enhances erythroid proliferation

Truncating mutations in the *EPOR* gene that cause clinically benign erythrocytosis[4,6]

provide a potentially safe avenue to increase erythropoietic output from genome-edited

HSPCs. In this study we designed Cas9 single-guide RNAs (sgRNAs)[11] (termed *EPOR*-

sg1 and *EPOR*-sg2) that overlap the location of the originally identified nonsense

mutation, *EPOR* c.1316G>A (p.W439X) ("Mäntyranta variant").[4] (Figure 1a, Extended

Data Figure 2a). Our hypothesis was that targeting this site in exon 8 with Cas9 would

create a spectrum of indels, a subset of which would result in a frameshift of the reading

frame and yield premature downstream stop codons in the *EPOR* gene.

To test this hypothesis, we pre-complexed each sgRNA with high-fidelity Cas9

protein[17] and delivered these ribonucleoprotein (RNP) complexes to human CD34[+]

HSPCs. At 2-3 days post-editing, we transferred cells into culture media that promotes

erythroid differentiation over the course of two weeks[18] (Figure 1a). To determine whether

edited HSPCs have a proliferative advantage compared to unedited cells, we harvested

genomic DNA at days 0, 4, 7, 11, and 14 of RBC differentiation. We then quantified indel

frequency by PCR amplification followed by Sanger sequencing and decomposition

analysis using TIDE[19]. In the absence of a selective advantage or disadvantage, percent

edited alleles in cells at the beginning and end of RBC differentiation will be roughly

equivalent—which is what we observe for the editing frequency of the *HBB* sgRNA used

for correction of sickle cell disease. However, we observe that the editing frequency of

EPOR-targeting sgRNAs increases significantly over the course of erythroid

differentiation, to a greater extent in *EPOR*-sg1 than *EPOR*-sg2 (Figure 1b, Extended

Data Figure 2b) (P=0.0016 for *EPOR*-sg1 from day 0 to day 14). Additionally, we show

that the increase in indels for both *EPOR*-sg1 and *EPOR*-sg2 are predominantly driven

by indels that yield downstream stop codons (Extended Data Figure 2c-d, Extended Data

Figure 3a-b). This indicates that edited cells, particularly those with premature stop

codons, are outcompeting unedited cells because of the EPO hypersensitivity of tEPOR-

expressing cells in culture.

Because not all indels created by the sgRNAs cause truncations in *EPOR*, we

speculated that we could increase this proliferative effect by using HDR to insert a stop

codon at the exact location of the original variant (c.1316G>A). To accomplish this, we

designed an AAV6 repair template vector that introduces a stop codon into *EPOR* at the

439th amino acid (W439X) followed by a BGH-polyA tail to terminate transcription. We

also included a downstream GFP marker driven by the constitutive human UbC promoter

to ensure that each GFP⁺ allele harbors the intended *EPOR*-truncating mutation. The

entire integration cassette was flanked by 950-base pair homology arms that

corresponded to the genomic DNA immediately upstream and downstream of the

intended Cas9 cut site created by the more effective *EPOR*-sg1 (Figure 1c). To determine

whether this editing strategy was also able to drive enrichment of genome-edited RBCs,

we complexed *EPOR*-sg1 with Cas9 protein and delivered this by electroporation to

human CD34[+] HSPCs followed by transduction with AAV6 DNA repair template. Two to

three days post-editing we either maintained cells in HSPC media or began erythroid

differentiation with 3 U/mL of EPO (+EPO), as has been previously described[8], or with 0

U/mL of EPO (-EPO) in order to determine whether *tEPOR*-expressing cells retain EPO

sensitivity or became EPO independent during their differentiation. At day 14 of erythroid

differentiation, we stained for established RBC markers[8] and analyzed cells using flow

cytometry. We observed no differentiation when cells were kept in HSPC media and

efficient RBC differentiation in all treatments with EPO. In the edited conditions in the

absence of EPO, we observed moderate differentiation which may indicate the

hypersensitivity of tEPOR-expressing cells to trace amounts of EPO in the media, as has

been previously observed[4] (Figure 1d). By analyzing GFP[+] cells over the course of RBC

differentiation, in the *EPOR*[W439X]-edited conditions we observed a significant increase in

the frequency of edited cells in only the +EPO conditions (P=0.0006 comparing day 0 to

day 14) (Figure 1e). At the end of differentiation, we estimate that almost all of the RBCs

are derived from cells with either an INDEL or UBC-GFP knock-in. In addition to the

competitive advantage that *tEPOR* expression gives to edited cells over the course of

RBC differentiation, we also observed increased RBC production in both *EPOR*-sg1 and

EPOR-sg1 + UbC-GFP-BGH conditions compared to Mock control in the +EPO condition

(average 1.45×10^3 total fold increase in mock edited cells versus 4.71×10^3 in *EPOR*-sg1

+ UbC-GFP-BGH over the 14-day RBC differentiation) (Figure 1f). These increased cell

counts were not observed in the -EPO condition or when cells were maintained in HSPC

media, indicating an EPO-driven increase in erythroid proliferation in cells expressing

tEPOR (Extended Data Figure 4a).

In terms of the safety of this editing strategy, we show that introduction of these

EPOR indels yields RBCs with production of both fetal and adult hemoglobin tetramers

following hemoglobin tetramer high-performance liquid chromatography (HPLC)

(Extended Data Figure 4b). We also found similar colony number and lineage distribution

from CD34$^+$ HSPCs plated into wells containing methylcellulose media either with or

without EPO that were scored for colony-formation ability after 14 days (Extended Data

Figure 4c). As expected, there was a marked decrease in the ability to form BFU-E

colonies in the absence of EPO even if the cells contained the tEPOR. These data

reinforce the idea that truncation of EPOR does not alter the lineage bias of HSPCs, but

rather has an effect only after commitment to the erythroid lineage.

Integration of *tEPOR* cDNA at safe-harbor locus recapitulates the proliferative

effect

Given the therapeutic utility of transgene integration at safe harbor sites[20], we

hypothesized that integration of a *tEPOR* cDNA at a safe harbor site may also enable

increased erythroid production from edited HSPCs while leaving the endogenous *EPOR*

locus intact. Given the fact that integration at the *CCR5* locus is an established method

for delivery of therapeutic transgenes in HSPCs[21], we developed a custom AAV6-

packaged DNA repair template that would facilitate integration of an exogenous human

UbC promoter driving expression of *tEPOR* cDNA followed by a T2A-YFP-BGH reporter

(Figure 2a). Given the strong, constitutive expression of the UbC promoter, this method

of insertion is expected to express *tEPOR* ubiquitously in all hematopoietic cell types,

regardless of lineage.

To test this strategy for *tEPOR* expression, we edited HSPCs with Cas9

complexed with an established sgRNA targeting exon 2 of *CCR5*[21] (*CCR5*-sg3),

immediately followed by transduction with our custom DNA repair template. We then

performed RBC differentiation post-editing and analyzed the kinetics of editing frequency,

YFP expression, and erythroid differentiation using droplet digital PCR (ddPCR) and flow

cytometry. As with endogenous *EPOR* truncation strategies, we observed more efficient

erythroid differentiation in all treatments with EPO compared to the minus EPO conditions

(Figure 2b). While flow cytometry confirmed the ubiquitous expression of YFP in edited

cells, regardless of presence of CD71 and GPA erythroid markers during differentiation

(Figure 2c), we did observe significant enrichment of YFP-expressing RBCs in the

presence of EPO ($P<0.0001$ when comparing day 0 to day 14 of RBC differentiation

(Figure 2d)). This enrichment was confirmed at the genomic level by ddPCR which

showed an increase in percent edited alleles when *tEPOR*-expressing cells were

subjected to erythroid differentiation, increasing by an average of 7.3-fold in the presence

of EPO, as well as enrichment to a limited degree enrichment in the -EPO condition

(Extended Data Figure 5a). With this editing strategy, we observed increased RBC

production in the CCR5-sg3 + *tEPOR* condition and no increase in proliferation of CCR5-

sg3 alone compared to mock control cells in the +EPO condition. We saw no increased

proliferation in the cells maintained in HSPC media or in the -EPO condition (Extended

Data Figure 5b).

We once again observed that edited and unedited cells had no noticeable

difference in lineage commitment or colony-forming ability following a colony-forming unit

assay (Extended Data Figure 5c). In addition, we analyzed RBCs post-differentiation

using hemoglobin tetramer HPLC and found that UbC-mediated expression of *tEPOR*

resulted in both HgbF and HgbA expression but a relative increase in HgbF expression

(Extended Data Figure 5d). These results indicate that expression of *tEPOR* from a safe

harbor site is an effective means of driving increased RBC production from genome-

edited HSPCs.

Integration of *tEPOR* cDNA at *HBA1* demonstrates an erythroid-specific

proliferative effect

While integration at a safe harbor locus effectively increased erythropoietic output from

edited HSPCs, there is the concern that constitutive expression in all cell types could

disrupt stem-ness or lead to other unintended effects. As an alternative, we can introduce

promoter-less transgenes into endogenous genes in order for the integration cassette to

be regulated by endogenous expression machinery. For instance, in prior work we

designed a genome editing strategy to fully replace the *HBA1* gene with an *HBB*

transgene in order to correct β-thalassemia[8]. We found that because α-globin is produced

by duplicate genes, *HBA1* may serve as a safe harbor site to deliver custom payloads

with strong erythroid-specific expression.

We therefore hypothesized that integration of the *tEPOR* cDNA at the *HBA1* locus

could further enhance production of edited RBCs while avoiding potential complications

with ubiquitous transgene expression because of the specificity of expression of *HBA1* in

the RBC lineage. To test this, we designed a custom integration cassette (also packaged

in AAV6) to use with an established sgRNA (*HBA1*-sg4) to introduce a promoter-less

tEPOR cDNA followed by a T2A-YFP reporter under expression of endogenous *HBA1*

regulatory machinery (Figure 2e). Following editing, we observed efficient erythroid

differentiation in all treatments with EPO and little to no differentiation in the absence of

EPO (Figure 2f). We found that this integration strategy indeed yielded RBC-specific

expression of YFP, which was only detectable as cells gained CD71 and GPA erythroid

markers (Figure 2g). Over the course of RBC differentiation, we observed dramatic

enrichment of GFP$^+$ cells exclusively in the +EPO condition (Figure 2h). While we

observed a mild degree of enrichment of edited alleles using ddPCR in the absence of

EPO, this effect was more pronounced in the presence of EPO, eliciting an average 4.4-

fold increase in percentage of edited alleles over the course of RBC differentiation

(Extended Data Figure 6a). Once again, we found that cells edited with *tEPOR* at *HBA1*

demonstrated similar colony-forming ability as mock edited cells (Extended Data Figure

6b). Additionally, we found that *tEPOR*-expressing cells produce ratios of human

hemoglobin that are similar to unedited RBCs (Extended Data Figure 6c).

Combined *HBB-tEPOR* bicistronic cassettes simultaneously correct β-thalassemia

and increase production of edited RBCs

Because the above results demonstrate that *tEPOR* expression yields increased RBC

production from edited HSPCs, we then sought to couple this selective advantage with a

therapeutic gene edit. We chose to combine tEPOR with our prior β-thalassemia

correction approach to simultaneously correct the disease and increase production of

clinically meaningful RBCs from these corrected HSPCs. One way this can be

accomplished is by creating a bicistronic cassette that links expression of the therapeutic

full-length *HBB* transgene with a *tEPOR* cDNA. Prior studies have found that the type of

linker domain used can have a great bearing on transgene expression and protein

function—particularly when function is dependent on formation of protein complexes, as

is the case with the globin genes[8]. Therefore, we designed and tested a variety of AAV6

repair template vectors linking the two genes using standard T2A peptides, optimized T2A

peptides with Furin cleavage sites[22] (referred to as "FuT2A"), internal ribosome entry sites

(referred to as "IRES"), as well by driving *tEPOR* expression from a separate exogenous

promoter—human PGK1 (referred to as "PGK"). To evaluate the different vectors, we

edited healthy donor HSPCs as previously described at *HBA1* using the bicistronic AAV6

repair templates and evaluated their ability to differentiate and enrich for edited RBCs

over the course of erythroid differentiation (Figure 3a). While efficient RBC differentiation

was achieved in all editing conditions (Figure 3b), we found that all four bicistronic

cassettes drove >2-fold enrichment of edited alleles (range: 2.1-fold to 3.5-fold) (P=0.003

for PGK-tEPOR, P=0.0055 for T2A-tEPOR, P=0.0259 for F2A-tEPOR, and P=0.0003 for

IRES-tEPOR when comparing day 0 to day 14) (Figure 3c-d). We note that allele targeting

frequencies of 60% translates into >80% of the cells having at least one allele targeted

(cell targeting frequency) and thus would not expect to see much more enrichment than

we observed.

To ensure this strategy was also effective in patient-derived cells, we tested these

bicistronic vectors in HSPCs derived from sickle cell disease (SCD) patients, this time

comparing them to a therapeutic full-length *HBB* transgene[8]. All of the constructs are

knocked-in to the *HBA1* locus without disrupting the endogenous *HBB* locus expressing

HgbS. We found that vectors did not disrupt erythroid differentiation when compared to

mock edited cells in the same donor, although their ability to differentiate was likely

impacted by the variable quality of the frozen patient samples (Extended Data Figure 7a).

Again, we observed >2-fold enrichment of edited alleles for all four bicistronic vectors

(range: 2.0-fold to 3.1-fold), but no change in editing frequency for the original β-

thalassemia correction vector (P=0.0061 for PGK-tEPOR, P=0.011 for T2A-tEPOR,

P=0.0016 for Fu2A-tEPOR, P=0.0153 for IRES-tEPOR when comparing day 0 to day 14)

(Figure 3e-f). We estimate that at the end of the differentiation almost all of the cells have

at least one allele with the HBB-tEPOR knock-in and thus no biologic drive for further

enrichment. When we analyzed differentiated RBCs for hemoglobin tetramers by HPLC,

we found that the 2A vectors showed almost no HgbA expression (consistent with what

has been seen previously in which addition of a 2A can disrupt HBB protein function[23]).

In contrast, we found that PGK-tEPOR and IRES-tEPOR vectors showed an improvement

in HgbA production relative to the *HBB*-only edited cells (Extended Data Figure 7b). There

was no change in the HgbF expression in these samples.

Multiplexed genome editing to introduce *tEPOR* and *HBB* leads to robust increase

in *β-globin* mRNA in edited RBCs

In lieu of coupling the *tEPOR* cDNA and therapeutic edit at the same locus, an alternative

strategy would be to multiplex two editing events at different loci to simultaneously

truncate the endogenous *EPOR* and introduce the original β-thalassemia correction

vector at *HBA1*. This may have the additional advantage that the endogenous *EPOR*

truncation will more reliably recapitulate congenital erythrocytosis. We hypothesized that

we could simultaneously deliver Cas9 separately pre-complexed with *EPOR*-truncating

EPOR-sg1 and *HBA1*-sg4 gRNA and then transduce HSPCs with both the β-thalassemia

correction vector as well as the W439X *EPOR*-targeting vector. Because homology arms

of each vector are specific for each site—*HBB* for the *HBA1* locus and W439X for the

EPOR locus—integration for each vector will occur only at the intended locus. This

strategy may allow simultaneous correction of β-thalassemia and increased erythropoietic

output from corrected HSPCs. For this to be maximally effective, the two editing events

must be present in the same cell. Therefore, during editing we used a DNA-PKcs inhibitor

to increase the frequency of template integration at each locus[24,25]. Following editing, we

determined whether this multiplexed editing strategy increases the frequency of corrected

RBCs over the course of erythroid differentiation.

To model the clinical setting where both edited and unedited HSPCs would occupy

the patient's bone marrow, we introduced unedited cells at various concentrations at the

start of erythroid differentiation (Figure 4a). Importantly, none of the multiplexed

conditions disrupted erythroid differentiation compared to single-edited *HBB* or mock

conditions (Figure 4b). In all of the multiplexed conditions, we observed an increase in

the frequency of corrected alleles over the course of RBC differentiation (P=0.0332 for

HBB+tEPOR 100%, P=0.0086 for *HBB+tEPOR* 30%, P=0.0122 for *HBB+tEPOR* 10%

when comparing day 0 to day 14) (Figure 4c). In fact, in both the 30% and 10%

multiplexed conditions, we achieved a higher frequency of edited alleles by the conclusion

of erythroid differentiation compared to single-edited conditions with unedited cells

introduced at an equivalent concentration (P=0.0113 for *HBB* versus *HBB+tEPOR* 30%

at day 14 and P=0.008 for *HBB* versus *HBB+tEPOR* 10% at day 14) (Figure 4d). We

confirmed truncation at the *EPOR* locus measured by GFP[+] cells at day 14 of RBC

differentiation (Extended Data Figure 8a). We also observed a corresponding increase in

HBB mRNA expression from the *HBA1* locus in the 30% and 10% multiplexed edited

conditions on day 14 of RBC differentiation compared to single-edited conditions (Figure

4e), indicating an improved ability for multiplexed editing to increase therapeutic potential

of this β-thalassemia correction strategy. When we measured the colony-forming ability

of the edited cells and found that differentiation into the various lineages was similar

between mock cells and cells edited with *HBB* alone or *HBB+tEPOR*, although we did

observe a decrease in total colonies produced in both edited conditions as would be

expected from the increased amount of AAV used to target two loci[26,27] (Extended Data

Figure 8b-c).

Discussion

While insights from clinical genetics have typically implicated novel genes and

pathways in disease, here we sought to use human genetics to develop novel strategies

to *treat* disease. We used the precision of genome editing to capitalize on a previously

characterized disorder called congenital erythrocytosis, which leads to EPO

hypersensitivity and hyperproduction of erythrocytes, without causing pathology[4]. As

demonstrated previously with variants identified in human genetics, such as those found

in *CCR5*, *PCSK9*, and the gamma-globin promoter region, we hypothesized that we could

use CRISPR to introduce this natural *EPOR* variant (*tEPOR*) to increase erythropoietic

output from edited HSPCs. Prior work has highlighted the challenge of achieving long-

term correction of disease following delivery of gene therapy or genome editing correction

strategies[2,28,29]. While many efforts are underway to improve editing and engraftment

frequencies, we hypothesized that we could develop a strategy to increase production of

the clinically relevant cell type—the red blood cell—from edited HSPCs. If successful,

then lower editing and engraftment frequencies could yield sufficient production of RBCs

to achieve therapeutic benefit, and thus be curative for patients. Uchida *et al*. have shown

that introducing the *tEPOR* variant using lentiviral delivery enhanced the efficacy of

shRNA knockdown of *BCL11A* in upregulating HgbF, confirming the beneficial function of

this variant[30]. However, our work deploys the precision of genome editing to generate

tEPOR which may have broad utility across a spectrum of blood disorders.

We explored multiple genome-editing strategies to create the *tEPOR* variant, either

through truncation of the endogenous *EPOR* gene or integration of a *tEPOR* cDNA at

safe harbor loci. We found that HSPCs expressing *tEPOR* consistently demonstrated

increased erythropoiesis but otherwise normal, EPO-dependent production of

hemoglobin. In order to increase RBC production of genome-edited HSPCs in the context

of disease correction, we combined the *tEPOR* cassette with a previously described β-

thalassemia strategy in which the *HBB* gene replaces the *HBA1* gene using homology-

directed repair-based genome editing[8]. This allowed us to simultaneously introduce an

HBB transgene to restore normal hemoglobin production and to increase erythropoietic

output from edited HSPCs. To demonstrate the flexibility of the various *tEPOR*-

introduction strategies, we also developed an alternative multiplexed, site-specific

genome editing strategy to pair the original β-thalassemia correction strategy with

introduction of the *EPOR* truncation at the endogenous locus. Both strategies led to

enrichment of genome-edited RBCs over the course of erythroid differentiation compared

to the traditional β-thalassemia correction strategy. Because we found the effects to be

EPO-dependent, it is possible the selective advantage *in vivo* may be even more

pronounced because patients suffering from the hemoglobinopathies display elevated

EPO levels due to their severe anemia[31].

In terms of safety, the strategy to use CRISPR to introduce natural variants has

the benefit of having been already "tested" *in vivo* in humans. However, it must be noted

that several of the genome editing strategies introduce *tEPOR* under non-native

regulation which could alter the normal function of tEPOR. In considering this possibility,

we note while every cell in patients with CE harbor an *EPOR* truncation, therapeutic

deployment of the genome editing strategy will result in the introduction of *tEPOR* in only

a subset of HSPCs resident in the BM. Therefore, any aberrant effects of non-native

tEPOR expression (e.g., bias away from lymphoid or other cell types) are unlikely to lead

to cytopenia given the large number of unedited HSPCs remaining in the bone marrow

post-transplant. Furthermore, by increasing erythropoietic output from edited HSPCs, we

believe this work could enable the reduction or elimination of high-morbidity myeloablation

regimens that are currently required to attain therapeutic levels of edited HSPCs.

Expression of *tEPOR* could therefore be integrated into any treatment for blood disorders

that involve transplantation of HSPCs. For example, even in an allogeneic hematopoietic

stem cell transplant for red blood cell disorders, a truncation in the natural *EPOR* could

be created using indel-based genome editing to give the derived transplanted erythroid

progenitors a selective advantage. This strategy could thereby enable less toxic

myeloablative conditioning to be effectively utilized where mixed chimerism might be the

result.

385   Taken together, these results demonstrate the power of combining knowledge

from human genetics with the precision of CRISPR genome editing technology to

introduce clinically meaningful variants. As human genome sequencing becomes more

commonplace and clinically routine[32,33], it is likely that a greater number of variants of

unknown significance will be discovered and characterized. We therefore believe that the

strategy defined in this work—using CRISPR to introduce natural human variants—may

be deployed to amplify the therapeutic potential of current and future cell therapies.

Methods

AAV6 vector design, production, and purification. Adeno-associated virus, serotype 6 (AAV6) vector plasmids were cloned into the pAAV-MCS plasmid (Agilent Technologies, Santa Clara, CA, USA), comprised of inverted terminal repeats (ITRs) derived from AAV2. Gibson Assembly Mastermix (New England Biolabs, Ipswich, MA, USA) was used for the creation of all DNA repair vectors as per manufacturer's instructions. AAV6 vector was produced and purified with little variation from previously described processes[34]. 293T cells (Life Technologies, Carlsbad, CA, USA) were seeded in five 15 cm^2 dishes with 13-15×10^6 cells per plate 24 hours pre-transfection. Then, each dish was transfected with a standard polyethylenimine (PEI) transfection of 6µg ITR-containing plasmid and 22µg pDGM6 (gift from David Russell, University of Washington, Seattle, WA, USA), which holds the AAV6 cap, AAV2 rep, and Ad5 helper genes. Following a 48-72h incubation, cells were harvested and vectors were purified using the AAVpro purification kit (cat.: 6666; Takara Bio, Kusatsu, Shiga, Japan) as per manufacturer's instructions and stored at -80 °C until further use. AAV6 vectors were titered using ddPCR to measure the number of vector genomes as previously described[35].

***In vitro* culture of CD34$^+$ HSPCs.** Human CD34$^+$ HSPCs were cultured in conditions as previously described[7,36–39]. CD34$^+$ HSPCs were isolated from cord blood (provided by Stanford Binns Program for Cord Blood Research) or sourced from Plerixafor- and/or G-CSF-mobilized peripheral blood (AllCells, Alameda, CA, USA and STEMCELL Technologies, Vancouver, Canada). Frozen Plerixafor- and/or G-CSF-mobilized peripheral blood of patients with sickle cell disease were provided by Dr. Vivien Sheehan,

Emory University. CD34$^+$ HSPCs were cultured at 1×10^5–5×10^5 cells/mL in StemSpan

Serum-Free Expansion Medium II (STEMCELL Technologies, Vancouver, Canada) or

Good Manufacturing Practice Stem Cell Growth Medium (SCGM, CellGenix, Freiburg,

Germany) supplemented with a human cytokine (PeproTech, Rocky Hill, NJ, USA)

cocktail: stem cell factor (100 ng/mL), thrombopoietin (100 ng/mL), Fms-like tyrosine

kinase 3 ligand (100 ng/ml), interleukin 6 (100 ng/mL), streptomycin (20 mg/mL), and

penicillin (20 U/mL), and 35 nM of UM171 (cat.: A89505; APExBIO, Houston, TX, USA).

The cell incubator conditions were 37°C, 5% CO_2, and 5% O_2.

Electroporation-aided transduction of cells. The synthetic chemically modified

sgRNAs used to edit CD34$^+$ HSPCs were purchased from Synthego (Menlo Park, CA,

USA) or TriLink Biotechnologies (San Diego, CA, USA) and were purified by HPLC.

These modifications are comprised of 2′-O-methyl-3′-phosphorothioate at the three

terminal nucleotides of the 5′ and 3′ ends described previously[11]. The target sequence for

the gRNAs were as follows:

EPOR gRNA (*EPOR*-sg1)

5′-AGCTCAGGGCACAGTGTCCA-3′

EPOR gRNA (*EPOR*-sg2)

5′-GCTCCCAGCTCTTGCGTCCA-3′

CCR5 gRNA (*CCR5*-sg3)

5′-GCAGCATAGTGAGCCCAGAA-3′

HBA1 gRNA (*HBA1*-sg4)

5′-GGCAAGAAGCATGGCCACCG-3′

The HiFi Cas9 protein was purchased from Integrated DNA Technologies (IDT)

(Coralville, Iowa, USA) or Aldevron (Fargo, ND, USA). Preceding electroporation, RNPs

were complexed at a Cas9:sgRNA molar ratio of 1:2.5 at 25°C for 10-20 min. Next, CD34$^+$

cells were resuspended in P3 buffer (Lonza, Basel, Switzerland) with complexed RNPs

and subsequently electroporated using the Lonza 4D Nucleofector and 4D-Nucleofector

X Unit (program DZ-100). Electroporated cells were then plated at 1×10^5-5×10^5 cells/mL

in the previously described cytokine-supplemented media. Immediately following

electroporation, AAV6 was dispensed onto cells at 2.5×10^3-1×10^4 vector genomes/cell

based on titers determined by ddPCR. For multiplex editing experiments, in addition to

the steps described above, cells were incubated with 0.5 µM of a DNA-PKcs inhibitor,

AZD7648 (cat.: S8843; Selleck Chemicals, Houston, TX) for 24 hours, as previously

described[24,25].

Allelic modification analysis using ddPCR. Edited HSPCs were harvested within 2-3

days post-electroporation and at each media change throughout erythrocyte

differentiation and then analyzed for modification frequencies of the alleles of interest. To

quantify editing frequencies, we created custom ddPCR primers and probes to quantify

HDR alleles (using in-out PCR and probe corresponding to the expected integration

event) compared to an established genomic DNA reference (REF) at the *CCRL2* locus[8].

QuickExtract DNA extraction solution (Epicentre, Madison, WI, USA) was used to collect

gDNA input, which was then digested using BamHI-HF or HindIII-HF as per the

manufacturer's instructions (New England Biolabs, Ipswich, MA, USA). The percentage

of targeted alleles within a cell population was measured with a Bio-Rad QX200 ddPCR

machine and QuantaSoft software (v.1.7; Bio-Rad, Hercules, CA, USA) using the

following reaction mixture: 1-4 µL of digested gDNA input, 10 µL of ddPCR SuperMix for

Probes (no dUTP) (Bio-Rad), primer/probes (1:3.6 ratio; Integrated DNA Technologies),

and volume up to 20 µL with H_2O. ddPCR droplets were then generated following the

manufacturer's instructions (Bio-Rad): 20 µL of ddPCR reaction, 70 µL of droplet

generation oil, and 40 µL of droplet sample. Thermocycler (Bio-Rad) settings were as

follows: 98°C (10 min), 94°C (30 s), 55.7-60°C (30 s), 72°C (2 min), return to step 2 × 40–

50 cycles, and 98°C (10 min). ddPCR analysis was carried out as described above with

the PCR product generated as the template (20 fg total amplicon). Analysis of droplet

samples was then performed using the QX200 Droplet Digital PCR System (Bio-Rad).

We next divided the copies/µL for HDR (%): HDR (FAM) / REF (HEX). The following

primers and probes were used in the ddPCR reaction:

CCR5 (for tEPOR-YFP construct)

Forward Primer (FP): 5'-GGGAGGATTGGGAAGACA-3'

Reverse Primer (RP): 5'-AGGTGTTCAGGAGAAGGACA-3'

Probe: 5'-6-FAM/AGCAGGCATGCTGGGGATGCGGTGG/3IABkFQ-3'

HBA1 (for tEPOR-YFP construct)

FP: 5'-AGTCCAAGCTGAGCAAAGA-3'

RP: 5'-ATCACAAACGCAGGCAGAG-3'

Probe: 5'-CGAGAAGCGCGATCACATGGTCCTGC-3'

HBA1 (for HBB construct and tEPOR-HBB constructs)

FP: 5'-GTGGCTGGTGTGGCTAATG-3'

RP: 5'-CAGAAAGCCAGCCAGTTCTT-3'

Probe: 5'-6-FAM/CCTGGCCCACAAGTATCACT/3IABkFQ-3'

HBA1 (for HBB-tEPOR constructs)

FP: 5'-TCTGCTGCCAGCTTTGAGTA-3'

RP: 5'-GCTGGAGTGGGACTTCTCTG-3'

Probe: 5'-6-FAM/ACTATCCTGGACCCCAGCTC/3IABkFQ-3'

CCRL2 (reference)

FP: 5'- GCTGTATGAATCCAGGTCC-3',

RP: 5'- CCTCCTGGCTGAGAAAAAG -3'

Probe: 5'- HEX/TGTTTCCTC/ZEN/CAGGATAAGGCAGCTGT/3IABkFQ -3'

Indel analysis using TIDE software. Within 2-4 days post-electroporation, HSPCs were

harvested with QuickExtract DNA extraction solution (Epicentre, Madison, WI, USA) to

collect gDNA. The following primer sequences were used to amplify the respective cut

site at the *EPOR* locus*:*

FP: 5'-CAGCTGTGGCTGTACCAGAA-3'

RP: 5'-CAGCCTGGTGTCCTAAGAGC-3'

Sanger sequencing of respective samples was then used as input for indel frequency

analysis relative to a mock, unedited sample using TIDE as previously described[19].

***In vitro* differentiation of CD34⁺ HSPCs into erythrocytes.** Following editing, HSPCs

derived from healthy or sickle cell disease patients were cultured 2-3 days as described

above. Subsequently, a 14-day *in vitro* differentiation was performed in supplemented

SFEMII medium as previously described[18,40]. SFEMII base medium was supplemented

with 100 U/mL penicillin–streptomycin, 10 ng/mL SCF (PeproTech, Rocky Hill, NJ, USA),

1 ng/mL IL-3 (PeproTech, Rocky Hill, NJ, USA), 3 U/mL EPO (eBiosciences, San Diego,

CA, USA), 200 µg/mL transferrin (Sigma-Aldrich, St. Louis, MO, USA), 3% human serum

(heat-inactivated from Sigma-Aldrich or ThermoFisherScientific), 2% human plasma

(isolated from umbilical cord blood provided by Stanford Binns Cord Blood Program), 10

µg/mL insulin (Sigma-Aldrich, St. Louis, MO, USA), and 3 U/mL heparin (Sigma-Aldrich,

St. Louis, MO, USA). Cells were cultured in the first phase of medium for seven days at

1×10^5 cells/mL. In the second phase of medium, day 7-10, cells were maintained at 1×10^5

cells/mL, and IL-3 was removed from the culture. In the third phase of medium, day 11-

14, cells were cultured at 1×10^6 cells/mL, with a transferrin increase to 1 mg/mL.

Immunophenotyping of differentiated erythrocytes. Differentiated erythrocytes were

analyzed by flow cytometry on day 14 for erythrocyte lineage-specific markers using a

FACS Aria II (BD Biosciences, San Jose, CA, USA). Edited and unedited cells were

analyzed using the following antibodies: hCD45-V450 (HI30; BD Biosciences, San Jose,

CA, USA), CD34-APC (561; BioLegend, San Diego, CA, USA), CD71-PE-Cy7 (OKT9;

Affymetrix, Santa Clara, CA, USA), and CD235a-PE (GPA) (GA-R2; BD Biosciences, San

Jose, CA, USA). In addition to cell-specific markers, cells were also stained with Ghost

Dye Red 780 (Tonbo Biosciences, San Diego, CA, USA) to measure viability.

Hemoglobin tetramer analysis. Frozen pellets of approximately 1×10^6 cells *in vitro-*

differentiated erythrocytes were thawed and lysed in 30 µL of RIPA buffer with 1x Halt™

Protease Inhibitor Cocktail (Thermo Fisher Scientific, Waltham, MA, USA) for 5 minutes

on ice. The mixture was vigorously vortexed and cell debris were removed by

centrifugation at 13,000 RPM for 10 minutes at 4°C. HPLC analysis of hemoglobins in

their native form was performed on a cation-exchange PolyCAT A column (35×4.6 mm^2,

3 µm, 1,500 Å) (PolyLC Inc., Columbia, MD, USA) using a Perkin-Elmer Flexar HPLC

system (Perkin-Elmer, Waltham, MA, USA) at room temperature and detection at 415 nm.

Mobile phase A consisted of 20 mM Bis-tris and 2 mM KCN at pH 6.94, adjusted with HCl.

Mobile phase B consisted of 20 mM Bis-tris, 2 mM KCN, and 200 mM NaCl at pH 6.55.

Hemolysate was diluted in buffer A prior to injection of 20 µL onto the column with 8%

buffer B and eluted at a flow rate of 2 mL/min with a gradient made to 40% B in 6 min,

increased to 100% B in 1.5 min, and return to 8% B in 1 min and equilibrated for 3.5 min.

Quantification of the area under the curve of the peaks was performed with TotalChrom

software (Perkin-Elmer, Waltham, MA, USA) and raw values were exported to GraphPad

Prism 9 for plotting and further analysis.

mRNA analysis. Following differentiation of HSPCs into erythrocytes, cells were

harvested, and RNA was extracted using the RNeasy Plus Mini Kit (Qiagen, Hilden,

Germany). Subsequently, complementary DNA was made from approximately 100 ng of

RNA using the iScript Reverse Transcription Supermix for quantitative PCR with reverse

transcription (Bio-Rad, Hercules, CA, USA). Expression levels of β-globin transgene and

α-globin mRNA were quantified with a Bio-Rad QX200 ddPCR machine and QuantaSoft

software (v.1.7; Bio-Rad, Hercules, CA, USA) using the following primers and 6-FAM/

ZEN/IBFQ-labeled hydrolysis probes, purchased as custom-designed PrimeTime qPCR

Assays from IDT:

HBB and HBB-tEPOR into HBA1:

FP: 5'-GGTCCCCACAGACTCAGAGA-3'

RP: 5'-CAGCATCAGGAGTGGACAGA

Probe: 5'-6-FAM/AACCCACCATGGTGCATCTG/3IABkFQ -3'

To normalize for RNA input, levels of the RBC-specific reference gene GPA were

determined in each sample using the following primers and HEX/ZEN/IBFQ-labeled

hydrolysis probes, purchased as custom-designed PrimeTime qPCR Assays from IDT:

GPA (reference):

FP: 5′-ATATGCAGCCACTCCTAGAGCTC-3′

RP: 5′-CTGGTTCAGAGAAATGATGGGCA-3′

Probe: 5'- HEX/AGGAAACCGGAGAAAGGGTA/3IABkFQ -3'

ddPCR reactions were created using the respective primers and probes, and droplets

were generated as described above. Thermocycler (Bio-Rad; settings were as follows:

98°C (10 min), 94°C (30 s), 54°C (30 s), 72°C (30 s), return to step 2 × 50 cycles, and

98°C (10 min). Analysis of droplet samples was done using the QX200 Droplet Digital

PCR System (Bio-Rad). To determine relative expression levels, the numbers of *HBB*

transgene copies/mL were divided by the numbers of *GPA* copies/mL.

Methylcellulose colony forming unit (CFU) assay. 2-3 days post-electroporation

HSPCs were plated in SmartDish 6-well plates (cat.: 27370; STEMCELL Technologies,

Vancouver, Canada) containing MethoCult H4434 Classic or MethoCult H4434 Classic

without EPO (cat.: 04444; cat: 04544.; STEMCELL Technologies, Vancouver, Canada).

After 14 days, the wells were imaged using the STEMvision Hematopoietic Colony

Counter (STEMCELL Technologies, Vancouver, Canada). Colonies were counted and

scored to determine the number of BFU-E, CFU-E, CFU-GM, and CFU-GEMM colonies.

Statistical analysis. GraphPad Prism 9 software was used for all statistical analysis.

References

1. Hütter G, Nowak D, Mossner M, et al. Long-Term Control of HIV by CCR5
Delta32/Delta32 Stem-Cell Transplantation. *N Engl J Med*. 2009;360(7):692-698.
doi:10.1056/NEJMoa0802905

2. Frangoul H, Altshuler D, Cappellini MD, et al. CRISPR-Cas9 Gene Editing for Sickle
Cell Disease and β-Thalassemia. *N Engl J Med*. 2021;384(3):252-260.
doi:10.1056/NEJMoa2031054

3. Fitzgerald K, White S, Borodovsky A, et al. A Highly Durable RNAi Therapeutic
Inhibitor of PCSK9. *N Engl J Med*. 2017;376(1):41-51. doi:10.1056/NEJMoa1609243

4. de la Chapelle A, Träskelin AL, Juvonen E. Truncated erythropoietin receptor
causes dominantly inherited benign human erythrocytosis. *Proc Natl Acad Sci U S
A*. 1993;90(10):4495-4499.

5. Sokol L, Luhovy M, Guan Y, Prchal J, Semenza G, Prchal J. Primary familial
polycythemia: a frameshift mutation in the erythropoietin receptor gene and
increased sensitivity of erythroid progenitors to erythropoietin. *Blood*. 1995;86(1):15-
22. doi:10.1182/blood.V86.1.15.bloodjournal86115

6. Arcasoy MO, Degar BA, Harris KW, Forget BG. Familial erythrocytosis associated
with a short deletion in the erythropoietin receptor gene. *Blood*. 1997;89(12):4628-
4635.

7. Dever DP, Bak RO, Reinisch A, et al. CRISPR/Cas9 β-globin gene targeting in
human haematopoietic stem cells. *Nature*. 2016;539(7629):384-389.
doi:10.1038/nature20134

8. Cromer MK, Camarena J, Martin RM, et al. Gene replacement of α-globin with β-
globin restores hemoglobin balance in β-thalassemia-derived hematopoietic stem
and progenitor cells. *Nat Med*. 2021;27(4):677-687. doi:10.1038/s41591-021-01284-
y

9. Vaidyanathan S, Salahudeen AA, Sellers ZM, et al. High-Efficiency, Selection-free
Gene Repair in Airway Stem Cells from Cystic Fibrosis Patients Rescues CFTR
Function in Differentiated Epithelia. *Cell Stem Cell*. 2020;26(2):161-171.e4.
doi:10.1016/j.stem.2019.11.002

10. Choi E, Koo T. CRISPR technologies for the treatment of Duchenne muscular
dystrophy. *Mol Ther*. 2021;29(11):3179-3191. doi:10.1016/j.ymthe.2021.04.002

11. Hendel A, Bak RO, Clark JT, et al. Chemically modified guide RNAs enhance
CRISPR-Cas genome editing in human primary cells. *Nat Biotechnol*.
2015;33(9):985-989. doi:10.1038/nbt.3290

12. Shin JJ, Schröder MS, Caiado F, et al. Controlled Cycling and Quiescence Enables
Efficient HDR in Engraftment-Enriched Adult Hematopoietic Stem and Progenitor
Cells. *Cell Rep*. 2020;32(9):108093. doi:10.1016/j.celrep.2020.108093

13. Transient Inhibition of 53BP1 Increases the Frequency of Targeted Integration in
Human Hematopoietic Stem and Progenitor Cells. doi:10.21203/rs.3.rs-2621625/v1

14. Chakraborty NG, Bilgrami S, Maness LJ, et al. Myeloablative chemotherapy with
autologous peripheral blood stem cell transplantation for metastatic breast cancer:
immunologic consequences affecting clinical outcome. *Bone Marrow Transplant*.
1999;24(8):837-843. doi:10.1038/sj.bmt.1701999

15. Bhatia S. Long-term health impacts of hematopoietic stem cell transplantation inform
recommendations for follow-up. *Expert Rev Hematol*. 2011;4(4):437-454.
doi:10.1586/ehm.11.39

16. Inamoto Y, Shah N, Savani B, et al. Secondary solid cancer screening following
hematopoietic cell transplantation. *Bone Marrow Transplant*. 2015;50(8):1013-1023.
doi:10.1038/bmt.2015.63

17. Vakulskas CA, Dever DP, Rettig GR, et al. A high-fidelity Cas9 mutant delivered as
a ribonucleoprotein complex enables efficient gene editing in human hematopoietic
stem and progenitor cells. *Nat Med*. 2018;24(8):1216-1224. doi:10.1038/s41591-
018-0137-0

18. Dulmovits BM, Appiah-Kubi AO, Papoin J, et al. Pomalidomide reverses γ-globin
silencing through the transcriptional reprogramming of adult hematopoietic
progenitors. *Blood*. 2016;127(11):1481-1492. doi:10.1182/blood-2015-09-667923

19. Brinkman EK, Chen T, Amendola M, van Steensel B. Easy quantitative assessment
of genome editing by sequence trace decomposition. *Nucleic Acids Res*.
2014;42(22):e168. doi:10.1093/nar/gku936

20. Papapetrou EP, Schambach A. Gene Insertion Into Genomic Safe Harbors for
Human Gene Therapy. *Mol Ther J Am Soc Gene Ther*. 2016;24(4):678-684.
doi:10.1038/mt.2016.38

21. Gomez-Ospina N, Scharenberg SG, Mostrel N, et al. Human genome-edited
hematopoietic stem cells phenotypically correct Mucopolysaccharidosis type I. *Nat
Commun*. 2019;10(1):4045. doi:10.1038/s41467-019-11962-8

22. Fang J, Yi S, Simmons A, et al. An antibody delivery system for regulated
expression of therapeutic levels of monoclonal antibodies in vivo. *Mol Ther J Am
Soc Gene Ther*. 2007;15(6):1153-1159. doi:10.1038/sj.mt.6300142

23. Reinhardt A, Kagawa H, Woltjen K. N-Terminal Amino Acids Determine KLF4
Protein Stability in 2A Peptide-Linked Polycistronic Reprogramming Constructs.
Stem Cell Rep. 2020;14(3):520-527. doi:10.1016/j.stemcr.2020.01.014

24. Selvaraj S, Feist WN, Viel S, et al. High-efficiency transgene integration by
homology-directed repair in human primary cells using DNA-PKcs inhibition. *Press
Nat Biotechnol*.

25. Wimberger S, Akrap N, Firth M, et al. Simultaneous inhibition of DNA-PK and PolΘ
improves integration efficiency and precision of genome editing. Published online
December 15, 2022:2022.12.15.520396. doi:10.1101/2022.12.15.520396

26. Schiroli G, Conti A, Ferrari S, et al. Precise Gene Editing Preserves Hematopoietic
Stem Cell Function following Transient p53-Mediated DNA Damage Response. *Cell
Stem Cell*. 2019;24(4):551-565.e8. doi:10.1016/j.stem.2019.02.019

27. Ferrari S, Jacob A, Cesana D, et al. Choice of template delivery mitigates the
genotoxic risk and adverse impact of editing in human hematopoietic stem cells. *Cell*
Stem Cell. 2022;29(10):1428-1444.e9. doi:10.1016/j.stem.2022.09.001

28. Kanter J, Walters MC, Krishnamurti L, et al. Biologic and Clinical Efficacy of
LentiGlobin for Sickle Cell Disease. *N Engl J Med*. 2022;386(7):617-628.
doi:10.1056/NEJMoa2117175

29. George LA, Sullivan SK, Giermasz A, et al. Hemophilia B Gene Therapy with a
High-Specific-Activity Factor IX Variant. *N Engl J Med*. 2017;377(23):2215-2227.
doi:10.1056/NEJMoa1708538

30. Uchida N, Ferrara F, Drysdale CM, et al. Sustained fetal hemoglobin induction in
vivo is achieved by BCL11A interference and coexpressed truncated erythropoietin
receptor. *Sci Transl Med*. 2021;13(591):eabb0411.
doi:10.1126/scitranslmed.abb0411

31. Nişli G, Kavakli K, Aydinok Y, Oztop S, Cetingül N. Serum erythropoietin levels in
patients with beta thalassemia major and intermedia. *Pediatr Hematol Oncol*.
1997;14(2):161-167. doi:10.3109/08880019709030902

32. Halldorsson BV, Eggertsson HP, Moore KHS, et al. The sequences of 150,119
genomes in the UK Biobank. *Nature*. 2022;607(7920):732-740. doi:10.1038/s41586-
022-04965-x

33. Gudbjartsson DF, Helgason H, Gudjonsson SA, et al. Large-scale whole-genome
sequencing of the Icelandic population. *Nat Genet*. 2015;47(5):435-444.
doi:10.1038/ng.3247

34. Khan IF, Hirata RK, Russell DW. AAV-mediated gene targeting methods for human
cells. *Nat Protoc*. 2011;6(4):482-501. doi:10.1038/nprot.2011.301

35. Aurnhammer C, Haase M, Muether N, et al. Universal real-time PCR for the
detection and quantification of adeno-associated virus serotype 2-derived inverted
terminal repeat sequences. *Hum Gene Ther Methods*. 2012;23(1):18-28.
doi:10.1089/hgtb.2011.034

36. Bak RO, Porteus MH. CRISPR-Mediated Integration of Large Gene Cassettes Using
AAV Donor Vectors. *Cell Rep*. 2017;20(3):750-756.
doi:10.1016/j.celrep.2017.06.064

37. Bak RO, Dever DP, Porteus MH. CRISPR/Cas9 genome editing in human
hematopoietic stem cells. *Nat Protoc*. 2018;13(2):358-376.
doi:10.1038/nprot.2017.143

38. Charlesworth CT, Camarena J, Cromer MK, et al. Priming Human Repopulating
Hematopoietic Stem and Progenitor Cells for Cas9/sgRNA Gene Targeting. *Mol*
Ther Nucleic Acids. 2018;12:89-104. doi:10.1016/j.omtn.2018.04.017

39. Cromer MK, Vaidyanathan S, Ryan DE, et al. Global Transcriptional Response to
CRISPR/Cas9-AAV6-Based Genome Editing in CD34+ Hematopoietic Stem and
Progenitor Cells. *Mol Ther*. 2018;26(10):2431-2442.
doi:10.1016/j.ymthe.2018.06.002

40. Hu J, Liu J, Xue F, et al. Isolation and functional characterization of human
erythroblasts at distinct stages: implications for understanding of normal and
disordered erythropoiesis in vivo. *Blood*. 2013;121(16):3246-3253.
doi:10.1182/blood-2013-01-476390

Figure 1

Figure 1: Cas9-guided *EPOR* truncation in HSPCs enhances erythroid proliferation.

a. Schematic of HSPC editing and model of tEPOR effect. Representation of *EPOR* gene and location of the candidate sgRNA (*EPOR*-sg1) indicated by a line. Location of c.1316G>A mutation is denoted by the star. **b.** Frequency of indels created by sg1 in primary human CD34$^+$ HSPCs over the course of erythroid differentiation compared to control *HBB* sgRNA. Points represent median ± interquartile range. Values represent biologically independent HSPC donors: N=5 for *EPOR*-sg1, N=1 for control *HBB* sgRNA. *P=0.0016 of day 1 vs. day 14 by unpaired two-tailed *t*-test. **c.** Genome editing strategy when using an AAV6 DNA repair template to introduce the *EPOR* c.1316G>A mutation followed by a BGH-polyA region and UbC-driven GFP reporter. **d.** Percentage of GPA$^+$/CD71$^+$ of live single cells on day 14 of differentiation. Bars represent median ± interquartile range. Values represent biologically independent HSPC donors: N=2-3 for HSPC, N=3-4 for -EPO and +EPO conditions. *P=0.0016 of -EPO vs. HSPC conditions; **P=0.003 and ***P=0.0001 of -EPO vs. +EPO conditions by unpaired two-tailed *t*-test. **e.** Percentage of GFP$^+$ cells of live single cells maintained in RBC media +/- EPO or HSPC media as determined by flow cytometry. Points represent median ± interquartile range. Values represent biologically independent HSPC donors N=2 for HSPC condition and N=3-4 for -EPO and +EPO conditions. *P=0.04, **P=0.0006 of day 0 vs. day 14 by unpaired two-tailed *t*-test. **f.** Fold change in cell count throughout RBC differentiation (e.g., if at day 0 starting cell numbers were 1E5 cells total, therefore a fold count change of 1000 would yield a total cell number of 1E8 at day 14). Points represent mean ± SEM. Values represent biologically independent HSPC donors N=3 for Mock and *EPOR*-sg1 + BGH and N=2 for *EPOR*-sg1.

Figure 2

Figure 2: Integration of *tEPOR* cDNA demonstrates an erythroid-specific proliferative effect.

a. Genome editing strategy to introduce *tEPOR*-T2A-YFP-BGH-polyA cDNA at the *CCR5* locus with expression driven by a ubiquitous UbC promoter. **b.** Percentage of GPA$^+$/CD71$^+$ of live single cells on day 14 of differentiation following introduction of *tEPOR* at *CCR5* locus. Bars represent median ± interquartile range. Values represent biologically independent HSPC donors: N=2-3 for HSPC condition, N=2-4 for -EPO and +EPO conditions. *P=0.0018 of -EPO vs. HSPC conditions; **P<0.0001 of -EPO vs. +EPO conditions by unpaired two-tailed *t*-test. **c.** Representative flow cytometry plots of *CCR5*-edited HSPCs following differentiation into RBCs. **d.** Percentage of GFP$^+$ cells of live single cells as determined by flow cytometry. Points represent mean ± SEM. Values represent biologically independent HSPC donors: N=2 for HSPC condition, N=3 for -EPO condition, N=3-4 for +EPO condition. *P = 0.0003 of day 0 vs. day 14 by unpaired two-tailed *t*-test. **e.** Genome editing strategy to introduce *tEPOR*-T2A-YFP cDNA at the *HBA1* locus by whole gene replacement to place integration cassette under regulation of the endogenous *HBA1* promoter. **f.** Percentage of GPA$^+$/CD71$^+$ of live single cells on day 14 of differentiation following introduction of *tEPOR* cDNA at the *HBA1* locus. Bars represent median ± 95% confidence interval. Values represent biologically independent HSPC donors: N=2 for HSPC condition, N=2-3 for -EPO and +EPO condition. *P=0.0002 of -EPO to +EPO condition by unpaired two-tailed *t*-test. **g.** Representative flow cytometry plot of *HBA1*-edited HSPCs following differentiation into RBCs. **h.** Percentage of GFP$^+$ cells of live single cells as determined by flow cytometry. Points represent mean ± SEM.

Values represent biologically independent HSPC donors: N=2 for HSPC condition, N=3 for -EPO and +EPO condition.

Figure 3

Figure 3: Therapeutic editing frequencies are achieved using bicistronic *HBB-tEPOR* cassette.

a. Design of *HBB* (control) and *HBB-tEPOR* (bicistronic) AAV6 donor cassettes targeted to the *HBA1* locus by whole gene replacement. **b.** Percentage of GPA$^+$/CD71$^+$of CD34$^-$/CD45$^-$ cells on day 14 as determined by flow cytometry. Points shown as median ± 95% confidence interval. Values represent N=3 biologically independent HSPC donors. **c.** Percentage of edited alleles for each bicistronic *tEPOR* vector in cord-blood derived CD34$^+$ cells over the course of RBC differentiation. Points shown as median ± 95% confidence interval. Values represent N=3 biologically independent HSPC donors. ***P=0.003 for PGK-*tEPOR*, P=0.0055 for T2A-*tEPOR*, P=0.0259 for Fu2A-*tEPOR*, and P=0.0003 for IRES-*tEPOR* (day 0 vs. day 14) by unpaired two-tailed *t*-test. **d.** Fold change in edited alleles from beginning (day 0) to end (day 14) of RBC differentiation. Bars represent median ± 95% confidence interval. **e.** Percentage of edited alleles for control (*HBB*) and bicistronic *HBB-tEPOR* vectors in sickle cell disease patient cells over the course of RBC differentiation. Points shown as median ± 95% confidence interval. Values represent biologically independent HSPC donors: N=3 for PGK-*tEPOR*, N=4 for all other vectors. ***P=0.0061 for PGK-*tEPOR*, P=0.011 for T2A-*tEPOR*, P=0.0016 for Fu2A-*tEPOR*, P=0.0153 for IRES-*tEPOR* (day 0 vs. day 14) by unpaired two-tailed *t*-test. **f.** Fold change in edited alleles from beginning (day 0) to end (day 14) of RBC differentiation. Bars represent median ± standard deviation.

Figure 4

Figure 4: Multiplexed editing of *EPOR* and *HBA1* leads to robust increase in *HBB* mRNA within editing HSPCs.

a. Schematic of multiplexed editing strategy with spike-in of unedited cells at start of erythroid differentiation to model HSPC transplantation. **b.** Percentage of GPA$^+$/CD71$^+$ of CD34$^-$/CD45$^-$ cells on day 14 of RBC differentiation as determined by flow cytometry. Points shown as median ± interquartile range. N=3 biologically independent HSPC donors. **c.** Percentage of edited alleles at *HBA1* in all multiplexed editing/spike-in conditions throughout RBC differentiation. Points shown as median ± 95% confidence interval. N=3 biologically independent HSPC donors. *P=0.0332 for *HBB+tEPOR* 100%, P=0.0086 for *HBB+tEPOR* 30%, P=0.0122 for *HBB+tEPOR* 10% (day 0 vs. day 14) by unpaired two-tailed *t*-test. **P=0.0113 for *HBB* versus *HBB+tEPOR* 30% at day 14 and P=0.008 for HBB versus *HBB+tEPOR* 10% at day 14 by unpaired two-tailed *t*-test. **d.** Fold increase in edited alleles on day 14 of differentiation of multiplexed conditions versus *HBB* only. Bars represent median ± 95% confidence interval. *P=0.0315 and **P=0.0123 by unpaired two-tailed *t*-test. **e.** mRNA expression of integrated *HBB* at *HBA1* locus normalized to *HBB* expression from mock. *GPA* mRNA expression was used as a reference. N=3 biologically independent HSPC donors. Bars represent median ± 95% confidence interval.

Extended Data Figure 1

a

Extended Data Figure 1. Schematic of wild-type (WT) and tEPOR signaling cascade.

a. Visual representation of EPOR signaling cascade. SHP1 binds to intracellular inhibitory domain in WT cells, downregulating EPOR signaling. This domain is truncated in tEPOR cells making them hypersensitive to EPO.

Extended Data Figure 2

Extended Data Figure 2. Indel analysis of HSPCs edited with second candidate _EPOR_ sgRNA (sg2) over course of erythroid differentiation.

a. Schematic of _EPOR_ gene and location of the second candidate sgRNA (_EPOR_-sg2) indicated by a line. Location of c.1316G>A mutation is denoted by the star **b.** Frequency of indels created by _EPOR_-sg2 in primary human CD34$^+$ HSPCs over the course of erythroid differentiation. Points represent median ± interquartile range. Values represent N=4 biologically independent HSPC donors. **c.** Frequency of five most common indels found in one HSPC donor targeted with _EPOR_-sg2 over the course of RBC differentiation. **d.** Fold enrichment of five most common indels over course of RBC differentiation in one HSPC donor targeted with _EPOR_-sg2.

Extended Data Figure 3

Extended Data Figure 3. Indel analysis of HSPCs edited with most effective *EPOR* sgRNA (sg1) over course of erythroid differentiation.

a. Frequency of five most common indels found in one HSPC donor targeted with *EPOR*-sg1over the course of RBC differentiation. **b.** Fold enrichment of five most common indels over course of RBC differentiation in one HSPC donor targeted with *EPOR*-sg1.

Extended Data Figure 4

Extended Data Figure 4. HSPCs edited at *EPOR* locus show increased erythroid proliferation and maintain lineage formation.

a. Cell count fold change in mock and edited cells maintained in RBC media +/- EPO or HSPC media. Points represent mean ± SEM. Values represent biologically independent HSPC donors N=3 for Mock and *EPOR*-sg1+ BGH and N=2 for sg1. **b.** Representative HPLC plots of cells targeted with *EPOR*-sg1at day 14 of RBC differentiation. HgbF=fetal hemoglobin, HgbA=adult hemoglobin. **c.** Colony formatting unit (CFU) assay of mock edited cells versus cells edited with *EPOR*-sg1 + BGH. Bars represent total number of colonies of each type: CFU-GEMM (multi-potential granulocyte, erythroid, macrophage, megakaryocyte progenitor cells), CFU-GM (colony forming unit-granulocytes and monocytes), BFU-E (erythroid burst forming units), CFU-E (colony forming unit-erythroid) colonies. N=1.

Extended Data Figure 5

a

CCR5-sg3 + tEPOR

- HSPC
- -EPO
- +EPO

Fold increase in edited alleles vs Day of RBC differentiation

b

Cell count fold change vs Day of RBC differentiation

Mock | *CCR5*-sg3 | *CCR5*-sg3 + tEPOR

- HSPC
- -EPO
- +EPO

c

Number of colonies

- CFU-GEMM
- CFU-GM
- BFU-E
- CFU-E

Cas9	-	sg3	-	sg3
AAV6	-	+	-	+
EPO	-	-	+	+

d

Mock | *CCR5*-sg3 | *CCR5*-sg3 + tEPOR

HgbF, HgbA vs Time (min)

Extended Data Figure 5. Safe harbor integration of *tEPOR* leads to increased erythroid proliferation and editing frequencies without affecting HSPC lineage or hemoglobin formation.

a. Fold change in edited alleles at *CCR5* over course of RBC differentiation +/- EPO or maintained in HSPC media measured by ddPCR. Points shown as median ± interquartile range. N=3-4 biologically independent HSPC donors. **b.** Cell count fold change in mock and edited cells maintained in RBC media +/- EPO or HSPC media. Points represent mean ± SEM. Values represent biologically independent HSPC donors N=3 for Mock and *CCR5*-sg3 + *tEPOR* and N=2 for *CCR5*-sg3. **c.** CFU assay of mock edited cells versus cells edited with *CCR5*-sg3 + *tEPOR.* Bars represent total number of colonies of each type: CFU-GEMM (multi-potential granulocyte, erythroid, macrophage, megakaryocyte progenitor cells), CFU-GM (colony forming unit-granulocytes and monocytes), BFU-E (erythroid burst forming units), CFU-E (colony forming unit-erythroid) colonies. N=1. Mock condition from Extended Data 3c shown here for comparison. **d.** Representative HPLC plots of cells targeted with *CCR5*-sg3 or *CCR5*-sg3 + *tEPOR* at day 14 of RBC differentiation. HgbF=fetal hemoglobin, HgbA=adult hemoglobin.

Extended Data Figure 6

a *HBA1*-sg4 + tEPOR

-EPO
+EPO

Fold increase in edited alleles vs. Day of RBC differentiation

b

Number of colonies

Cas9	-	sg4	-	sg4
AAV6	-	+	-	+
EPO	-	-	+	+

CFU-GEMM
CFU-GM
BFU-E
CFU-E

c

Mock — HgbF, HgbA, mV vs. Time (min)

HBA1-sg4 + tEPOR — HgbF, HgbA, Time (min)

Extended Data Figure 6. Erythroid specific expression of *tEPOR* leads to increased editing frequencies without affecting HSPC lineage or hemoglobin formation.

a. Fold change in edited alleles at *HBA1* over course of RBC differentiation +/- EPO measured by ddPCR. Points shown as median ± 95% CI. N=3 biologically independent HSPC donors. **b.** CFU assay of mock edited cells versus cells edited with *HBA1*-sg4 + *tEPOR.* Bars represent total number of colonies of each type: CFU-GEMM (multi-potential granulocyte, erythroid, macrophage, megakaryocyte progenitor cells), CFU-GM (colony forming unit-granulocytes and monocytes), BFU-E (erythroid burst forming units), CFU-E (colony forming unit-erythroid) colonies. N=1. Mock condition from Extended Data 3c shown here for comparison. **c.** Representative HPLC plots of cells targeted with *HBA1*-sg4 + *tEPOR* at day 14 of RBC differentiation. HgbF=fetal hemoglobin, HgbA=adult hemoglobin.

Extended Data Figure 7

a

SCD CD34⁺ donors RBC differentiation

b

Extended Data Figure 7. Hemoglobin tetramer analysis of *tEPOR* edited SCD patient HSPCs following RBC differentiation.

a. Percentage of GPA$^+$/CD71$^+$ of CD34$^-$/CD45$^-$ cells on day 14 of RBC differentiation as determined by flow cytometry. Points shown as median ± 95% confidence interval. N=4 biologically independent HSPC donors. **b.** Ratio of adult hemoglobin to sickle hemoglobin from HPLC analysis of hemoglobin tetramers from differentiated SCD patient HSPCs. Bars shown as mean ± SD. N=4 biologically independent HSPC donors.

Extended Data Figure 8

a

AAV6: -, HBB, HBB, HBB, HBB + tEPOR, HBB + tEPOR, HBB + tEPOR

% Edited: -, 100%, 30%, 10%, 100%, 30%, 10%

Extended Data Figure 8. Multiplexed edited strategy shows maintenance of *EPOR* truncation at day 14 of RBC differentiation and does not disrupt HSPC lineage formation.

a. Percentage of GFP+ cells of live single cells on day 14 of RBC differentiation as determined by flow cytometry. Points represent median ± interquartile range. Values represent N=3 biologically independent HSPC donors. **b.** CFU assay of mock, single edited, and multiplex edited HSPCs. Bars represent percent of total colonies: CFU-GEMM (multi-potential granulocyte, erythroid, macrophage, megakaryocyte progenitor cells), CFU-GM (colony forming unit-granulocytes and monocytes), and BFU-E (erythroid burst forming units). N=1. **c.** Number of total colonies in CFU assay produced per 500 plated cells in each condition.