Functional potential and evolutionary response to long-term heat selection of bacterial
associates of coral photosymbionts

Running title (54 characters): Functions of Symbiodiniaceae-associated bacteria

Justin Maire[1,#], Gayle K. Philip[2], Jadzia Livingston[1], Louise M. Judd[3], Linda L. Blackall[1],
Madeleine J. H. van Oppen[1,4]

[1] School of BioSciences, The University of Melbourne, Parkville, VIC, 3010, Australia.

[2] Melbourne Bioinformatics, The University of Melbourne, Parkville, VIC, 3010, Australia.

[3] Doherty Applied Microbial Genomics, Department of Microbiology and Immunology, The
University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Parkville,
VIC, 3010, Australia.

[4] Australian Institute of Marine Science, PMB No 3, Townsville, 4810, QLD, Australia.

[#] Corresponding author: justin.maire@unimelb.edu.au

Abstract

Corals rely on a wide range of microorganisms for their functioning, including intracellular
dinoflagellates (Symbiodiniaceae) and bacteria. Marine heatwaves trigger the loss of
Symbiodiniaceae from coral tissues - coral bleaching - often leading to death. While coral-
bacteria interactions are widely studied, Symbiodiniaceae-bacteria interactions have remained
largely uninvestigated. Here, we provide a genomic analysis of 49 bacteria, spanning 16
genera, that closely associate with six cultured Symbiodiniaceae species. We analyzed
bacterial functional potential by focusing on potentially beneficial functions for the
Symbiodiniaceae host, including B vitamin synthesis and antioxidant abilities, which may be
crucial for Symbiodiniaceae heat tolerance and in turn coral resistance to thermal bleaching.
These analyses suggest a wide potential for B vitamin synthesis and the scavenging of
reactive oxygen species (through the production of carotenoids or antioxidant enzymes), and
possibly the transfer of organic carbon to host cells. Single nucleotide polymorphism analysis
between bacteria isolated from wild-type and heat-evolved Symbiodiniaceae cultures
revealed that exposure to long-term elevated temperature has resulted in mutations in genes
known to be involved in host-symbiont interactions, such as secretion systems. Climate
change may therefore modify how Symbiodiniaceae and bacteria interact. This study
provides an overview of the possible roles of Symbiodiniaceae-associated bacteria in
Symbiodiniaceae functioning and heat tolerance, reinforcing the need for further studies of
such interactions to fully understand coral biology and climate resilience.

Importance

Symbiotic microorganisms are crucial for the survival of corals and their resistance to coral bleaching in the face of climate change. However, the impact of microbe-microbe interactions on coral functioning is mostly unknown, but could be essential factors for coral adaption to future climates. Here, we investigated interactions between cultured dinoflagellates of the Symbiodiniaceae family, essential photosymbionts of corals, and associated bacteria. By assessing the genomic potential of 49 bacteria, we found that they are likely beneficial for Symbiodiniaceae, through the production of B vitamins and antioxidants. Additionally, bacterial genes involved in host-symbiont interactions, such as secretion systems, accumulated mutations following long-term exposure to heat, suggesting symbiotic interactions may change under climate change. This highlights the importance of microbe-microbe interactions in coral functioning.

Word count: 4556/5000

Introduction

Coral reefs are valuable ecosystems, as they provide a home to about a third of all eukaryotic marine species as well as benefits to human populations, such as food security and coastal protection (1, 2). Reef-building corals rely on a diverse suite of microorganisms for their survival (3–5). Endosymbiotic dinoflagellates of the Symbiodiniaceae family (Suessiales, Dinophyta) are probably the most crucial and most studied member of this microbial consortium (6). By providing photosynthate to their host, Symbiodiniaceae meet most of the energy needs of corals (7), and are essential for the health of coral reefs. Unfortunately, this symbiotic association is extremely fragile and is particularly sensitive to temperature fluctuations. Increases in sea surface temperatures can result in the loss of Symbiodiniaceae from coral tissues, a process known as coral bleaching, which can result in coral death (8). The main mechanism thought to trigger coral bleaching is an increase in the production of reactive oxygen species (ROS) by Symbiodiniaceae, which leak into coral cells, overwhelm the coral's antioxidant response, and trigger a cellular cascade that results in Symbiodiniaceae loss (9). Mass coral bleaching events are increasing in frequency and intensity because of anthropogenic climate change (10), thus highlighting the need to increase knowledge on coral and Symbiodiniaceae biology and to develop novel coral conservation and restoration methods, while greenhouse gas emissions are curbed.

Corals also house a wide diversity of bacteria, which colonize all microhabitats within their host (4, 11–13). Putative bacterial functions include nutrient cycling and protection against pathogens (4, 12). Recently, Symbiodiniaceae were also found to associate with bacteria, both extra- and intracellularly (14–16). Symbiodiniaceae-associated bacterial communities are highly diverse and change after both short-term thermal stress (17–19) and long-term heat selection (19), pointing at a potential bacterial involvement in Symbiodiniaceae thermal tolerance. Two studies have shown that inoculation of Symbiodiniaceae cultures with a single bacterial strain (belonging to the *Muricauda* and *Roseovarius* genera, respectively) can enhance Symbiodiniaceae (*Durusdinium* sp. and *Breviolum minutum*, respectively) performance under temperature and/or light stress (18, 20). In the case of *Durusdinium/Muricauda*, this was attributed to the production of a carotenoid by *Muricauda*, a pigment able to scavenge ROS (20). Nevertheless, the functions and functional potential of most Symbiodiniaceae-associated bacteria remain mostly unexplored (13, 21).

Here, we provide the first comparative genomic analysis of Symbiodiniaceae-associated bacteria. From a culture collection of more than 200 bacterial isolates, cultured from six Symbiodiniaceae species, we sequenced the genomes of 49 representative isolates. We present an in-depth analysis of bacterial functional potential that may be beneficial for Symbiodiniaceae hosts, such as B vitamin synthesis and ROS scavenging. We also analyze the effect of Symbiodiniaceae thermal selection on bacterial genomic evolution. This study provides an important step to understanding the contribution of bacteria to Symbiodiniaceae functioning, and in turn their roles in coral health and climate resilience.

Results and discussion

Isolation of bacteria from Symbiodiniaceae cultures

In a previous study, 141 pure bacterial isolates, spanning 20 genera, were obtained from six Symbiodiniaceae species (15). We recently showed that long-term heat selection of the species *Cladocopium proliferum* affected the composition of Symbiodiniaceae-associated bacteria (19), both at the extracellular and intracellular level (22). Thus, we expanded our culture collection by pure culturing bacteria from one wild-type (WT10) and two heat-evolved (SS03 and SS08) *C. proliferum* strains. For each *C. proliferum* strain, samples were washed, and either bead-beat, in order to release potentially intracellular bacteria, or not. A total of 65 pure bacterial isolates were obtained and identified through 16S rRNA gene sequence analysis (Table S1). These newly obtained isolates spanned ten genera, five of which were previously isolated from Symbiodiniaceae cultures (15). Four genera (*Marinobacter, Muricauda, Mamelliela,* and *Roseitalea*) were cultured from all three *C. proliferum* strains.

Six genera (*Janibacter, Aestuariicoccus, Muricauda, Roseibium*, *Roseitalea*, and *Oceaniradius*) were isolated exclusively from *C. proliferum* samples that were bead beaten prior to spreading on agar plates, suggesting they may be intracellular associates. Nonetheless, *Muricauda, Roseitalea*, and *Roseibium* (previously referred to as *Labrenzia*) were previously isolated from *C. proliferum* WT10 samples that were not bead beaten (15), suggesting they can be extracellular. Therefore, only *Janibacter, Aestuariicoccus,* and *Oceaniradius* are strong candidates for being intracellular symbionts of *C. proliferum*, while all others are likely extracellular, though still closely associated.

Genome sequencing of 49 Symbiodiniaceae-associated bacteria

To explore the genomic abilities of bacteria that closely associate with Symbiodiniaceae, the whole genome of 49 isolates were sequenced, assembled, and annotated. The 49 isolates were chosen to span both the bacterial and Symbiodiniaceae diversity of our culture collection. All 49 assembled genomes exhibited a completeness above 98% and contamination below 2.5%, and ranged from 2.98-6.88 Mb in size (Table S2). The taxonomy of these genomes was confirmed with the GTDB-Tk tool, the construction of a phylogenetic tree, and the calculation of average nucleotide and amino acid identity (ANI and AAI, respectively) (Figures 1 and S1, Table S2). This corroborated the taxonomy previously obtained through 16S rRNA gene sequence analysis (see "Taxonomy notes" in Table S2). It is worth noting that, within one genus, isolates from distinct Symbiodiniaceae hosts often show >99.9% ANI and 100% AAI with each other (*e.g.*, *Muricauda, Mameliella, Roseitalea,* and *Marinobacter*; Figure S1), suggesting a lack of host-symbiont specificity.

Figure 1: Maximum-likelihood phylogenetic tree of the 49 pure-cultured Symbiodiniaceae-associated bacteria based on 120 marker genes (GTDB-Tk). The 49 isolates were assigned to 16 bacterial genera, as stipulated on the right-hand side. Bootstrap support values based on 1000 replications are provided.

The metabolic abilities of the 49 isolates were reconstructed using a KEGG annotation (Table S3), and pathways of interest are summarized in Figure 2. Pathways that were more than 75% complete were considered functional. Again, no host specificity or co-diversification was observed: isolates of a given genus all had almost identical metabolic potentials, regardless of the Symbiodiniaceae host they were cultured from. This is intriguing as the range of Symbiodiniaceae strains used in the study were originally isolated from four coral species, one sea anemone species, and one giant clam species. This is consistent with the fact that community composition of intracellular bacteria was highly similar across these Symbiodiniaceae cultures (15). Therefore, these bacteria may be opportunistic associates (as opposed to long-term symbionts), perhaps selected during the initial isolation of Symbiodiniaceae or accidentally introduced in Symbiodiniaceae cultures later on, which have

been maintained over the years, presumably because they were not detrimental, and
potentially beneficial, to their Symbiodiniaceae host.

Almost all isolates possessed all genes for glycolysis, the tricarboxylic acid (TCA) cycle, the
pentose phosphate pathway, the biosynthesis of all nucleotides, and the biosynthesis of
several amino acids (threonine, valine, isoleucine, leucine, lysine histidine). Only two
Roseibium isolates had the genes to perform denitrification, while all four *Roseibium* isolates
had the genetic potential for dissimilatory nitrate reduction. Denitrifying potential is
commonly observed in other *Roseibium* strains (sometimes referred to as *Labrenzia*) (23, 24).
Several isolates also possessed complete secretion systems, which may assist in host-bacteria
and bacteria-bacteria interactions, including Symbiodiniaceae infection: a type II secretion
system (T2SS) was found in three isolates (belonging to *Marinobacter*); a type IV secretion
system (T4SS) was found in 11 isolates (belonging to *Roseovarius, Roseivivax, Mameliella,*
Aquicoccus); a type VI secretion system (T6SS) was found in seven isolates (belonging to
Halomonas, Marinobacter, Roseibium). We also interrogated the presence of eukaryotic-like
proteins, which may facilitate host-bacteria interactions by mediating bacterial protein-
eukaryotic host protein interactions (25), but did not detect any in any of the 49 genomes
reported here.

Figure 2: Metabolic potential of 49 Symbiodiniaceae-associated bacteria. The
completeness of metabolic pathways (KEGG module database) of interest were annotated
using METABOLIC-G and pathway completeness (in %) was estimated in EnrichM. See
Table S3 for the full list of pathways.

Potential for B vitamin synthesis

B vitamins are required by most microalgae (26–28), including Symbiodiniaceae (21, 29),
and promote algal growth. For example, biotin (vitamin B_7) is essential for fatty acid
metabolism, while thiamine (vitamin B_1) is important for carbohydrate and amino acid
metabolism (27). Commonly used Symbiodiniaceae culture media, such as Daigo's IMK
medium used in this study, contain biotin, thiamine and cobalamin (vitamin B_{12}), although
some B vitamins have been shown to be provided by bacteria in some cases. For example,

Halomonas sp. and *Sinorhizobium meliloti* provide cobalamin to the unicellular algae
Porphyridium purpureum and *Chlamydomonas reinhardtii*, respectively (26, 30). Therefore,
we interrogated the completeness of B vitamin biosynthesis pathways in Symbiodiniaceae-
associated bacteria (Figure 2, Tables S3 and S4).

All but one isolate were found to be able to synthesize riboflavin (vitamin B_2), while 43
possessed the genes for tetrahydrofolate (vitamin B_9) synthesis (only missing in one
Oceaniradius, two *Roseibium* and three *Roseovarius*), 28 possessed the genes for
pantothenate (vitamin B_5) synthesis (belonging to *Bacillus, Roseivirga, Muricauda,*
Halomonas, Marinobacter, Oceanicaulis, Mameliella) and 23 isolates showed potential to
synthesize cobalamin (belonging to *Halomonas, Oceaniradius, Roseitalea, Stappia,*
Roseovarius, Mameliella, Aquicoccus, Roseibium, Nioella). The potential for the synthesis of
other vitamins was scarcer: 12 isolates for thiamine synthesis (belonging to *Bacillus,*
Halomonas, Qipengyuania, Roseibium, Oceanicaulis, Aquicoccus, Roseovarius); four isolates
for pyridoxal (vitamin B_6) synthesis (belonging to *Halomonas* and *Marinobacter*); five for
pimeloyl-ACP and biotin synthesis, including synthesis of the biotin precursor pimeloyl-ACP
(belonging to *Bacillus, Halomonas, Marinobacter*). Only MMSF_3323 (*Halomonas* sp.) had
the potential to synthesize the seven B vitamins that were investigated, while MMSF_3353
(*Bacillus* sp.) and all three *Marinobacter* isolates could synthesize five (Table S4).
Provisioning of cobalamin by a separate *Halomonas* strain was demonstrated in the red alga
P. purpureum (26). Therefore, there is high potential for B vitamin provisioning from
Symbiodiniaceae-associated bacteria. How these nutrients are exchanged between the
partners remains unknown, although some isolates did possess ABC transporters for thiamine
that may facilitate its export. Intracellular bacteria may be able to directly export the vitamins
in the Symbiodiniaceae cell's cytoplasm, making it readily available.

The potential for bacterial B vitamin provisioning is important for several reasons. First, as
shown in several other microalgae, B vitamin supplementation, specifically cobalamin, can
increase algal growth (26, 30, 31). Second, cobalamin supplementation by bacteria was
linked to increased thermal tolerance in *C. reinhardtii* (30), which is relevant to
Symbiodiniaceae in the context of thermal stress and coral bleaching. Third, coral hosts of
Symbiodiniaceae cannot synthesize B vitamins either. Some coral-associated bacteria, such
as *Endozoicomonas*, have been hypothesized to synthesize B vitamins for their coral hosts
(32–34). Symbiodiniaceae-associated bacteria may also provide B vitamins to the wider coral
holobiont, rather than just Symbiodiniaceae.

Potential for reactive oxygen species scavenging during thermal stress

ROS have been implicated in the thermal stress response of Symbiodiniaceae and in turn in
coral bleaching (8, 9, 35). Because of this, the production of ROS is a key trait in the design
of probiotics for improving thermal tolerance in Symbiodiniaceae and corals (18, 20, 36–39).
While Symbiodiniaceae continuously produce ROS (such as singlet oxygen ($^1O_2$), superoxide
(O_2^-), hydrogen peroxide (H_2O_2) and hydroxyl radicals (OH^-)), thermal stress results in an
increased production of ROS that may ultimately leak into coral cells and trigger coral

bleaching (9). Thus, we looked at pathways and proteins potentially involved in ROS
scavenging and/or the production of antioxidants. Pathways for the synthesis of glutathione
and ubiquinone, two antioxidants, were complete in 30 and 4 isolates, respectively (Figure 2,
Table S3). All Proteobacteria were able to produce glutathione, while only the four
Gammaproteobacteria isolates (*Halomonas* and *Marinobacter*) were able to produce
ubiquinone. Additionally, specific protein families (Pfam) or their synthesis pathways related
to ROS and RNS scavenging were manually searched in the 49 genomes (Table 1): synthesis
of zeaxhanthin, a carotenoid that quenches singlet oxygen and can scavenge other ROS (40);
dimethylsulfonioproprionate (DMSP) and dimethylsulfate (DMS), OH⁻ scavengers (41);
mannitol, another OH⁻ scavenger (42); ROS-scavenging enzymes, including superoxide
dismutases (O_2^- scavengers), peroxiredoxins and peroxidases (H_2O_2 scavengers),
glutaredoxins and thioredoxins (general ROS scavengers). These are summarized in Figure 3.

Table 1: Genes and respective protein families (Pfam) involved for which Symbiodiniaceae-
associated bacterial genomes were screened. DMSP: dimethylsulfonioproprionate; DMS:
dimethylsulfate; DMSO: dimethylsulfoxide; ROS: reactive oxygen species; SWEET: sugars
will eventually be exported transporter.

Pfam ID	Protein	Process
PF00494	Phytoene synthase	Zeaxanthin and β-carotene synthesis
PF18916	Lycopene β-cyclase	
PF04116	β-carotene hydroxylase	
PF00891	Methyltranferase (DsyB)	DMSP and DMS synthesis
PF16864	Methyltranferase (DsyB)	
PF00384	DMSO reductase	
PF16867	DMSP lyase	
PF08125	Mannitol 2-dehydrogenase	Mannitol synthesis
PF00081	Superoxide dismutase [Mn]	ROS scavenging
PF02777	Superoxide dismutase [Mn]	
PF00080	Superoxide dismutase [Cu-Zn]	
PF00578	Peroxiredoxin	
PF00462	Glutaredoxin	
PF13192	Thioredoxin	
PF00141	Peroxidase	

PF04193	SemiSWEET protein	Carbon transport

Figure 3: Putatively beneficial functions in Symbiodiniaceae-associated bacterial genomes. Protein families (Pfam) IDs were retrieved in each genome based on an InterProScan annotation. See Table 1 for full details on each Pfam ID. DMSP: dimethylsulfonioproprionate; DMS: dimethylsulfate; ROS: reactive oxygen species.

Only 14 isolates, all belonging to *Muricauda*, were found to be able to synthesize zeaxanthin. This is consistent with a previous study showing *Durusdinium*-associated *Muricauda* can produce zeaxanthin and contributes to Symbiodiniaceae tolerance to thermal and light stress (20). Twenty-two isolates possessed the *dsyB* gene, involved in the production of DMSP (belonging to *Roseovarius, Mameliella, Roseovivax, Roseibium, Stappia, Roseitalea, Oceaniradius,* and *Marinobacter*), while all isolates except the two *Roseivirga* isolates were able to synthesize DMS, through DMSO reduction and/or DMSP cleavage. Interestingly, a *Roseovarius* isolate, MMSF_3448, which enhanced thermal tolerance in *Breviolum minutum* cultures (18) had the genes to produce both DMSP and DMS, which may be part of the mechanism underlying the benefit found in that study. Additionally, bacterial DMSP and DMS metabolism was shown to increase in the proximity of Symbiodiniaceae cells (43), suggesting sulfur-based interactions between Symbiodiniaceae and their associated bacteria. A gene encoding for a mannitol-1-phosphate dehydrogenase, necessary for mannitol production, was found in 22 isolates (belonging to *Bacillus, Halomonas, Qipengyuania, Oceaniradius, Aquicoccus, Mameliella, Roseivivax, Nioella, Roseibium, Stappia,* and *Roseitalea*). Mannitol is has been shown to mitigate thermal bleaching in corals and sea anemones (44, 45). Finally, while all isolates possessed ROS-scavenging enzymes, some isolates had particularly high numbers, including isolates belonging to *Halomonas, Muricauda, Roseibium*, and *Roseivirga*. Isolates of the *Muricauda* and *Roseitalea* genera were the only bacteria possessing genes encoding thioredoxins, while only *Roseitalea* isolates

possessed all six queried enzymes. Hence, there is a high potential for Symbiodiniaceae-
associated bacteria to assist with ROS removal during thermal stress via various mechanisms.
This is particularly important for intracellular bacteria, as they may be able to export ROS-
scavenging compounds directly into the Symbiodiniaceae cytoplasm, strengthen the
antioxidant response, and minimize ROS leakage out of the Symbiodiniaceae cells. Most of
the bacterial genera studied here are commonly detected in and cultured from cnidarian hosts
(11, 46, 47) and may therefore participate in the *in hospite* Symbiodiniaceae response to
thermal stress, and thereby influence the outcome of coral thermal stress. Nonetheless,
whether these bacteria truly associate with Symbiodiniaceae *in hospite* remains unknown.

269 Potential for carbon exchange

Finally, we investigated the presence of carbon exporters in the bacterial genomes. The
"sugars will eventually be exported transporters" (SWEET) are bidirectional sugar
transporters found in eukaryotes, and are hypothesized to regulate carbon exchange between
Symbiodiniaceae and corals (48). SemiSWEET proteins are the bacterial homologues of
SWEET proteins, although sugar export has not yet been confirmed in bacteria (49). Five
isolates were found to have SemiSWEET genes (Figure 3): two *Roseivirga* isolates, one
Aquicoccus isolate, and two *Roseovarius* isolates. In a healthy state, it is unlikely that bacteria
would provide glucose to Symbiodiniaceae, which can make its own through photosynthesis,
and the bacterial transporters may instead be used to import photosynthate. However, sugars
may be exported via the same transporter, which may help compensate for the decrease in
photosynthesis, and thereby glucose production, by Symbiodiniaceae during thermal stress.
This is the first report of genes encoding for SemiSWEET proteins in Symbiodiniaceae-
associated bacteria, and their functional analysis is needed to shed light on their role towards
Symbiodiniaceae functioning and thermal tolerance.

285 Genomic evolution in response to Symbiodiniaceae experimental evolution

Among the bacterial genomes sequenced from *C. proliferum* cultures, we obtained a
representative from each of the three *C. proliferum* strains (WT10, SS03, and SS08) for four
bacterial species, *Marinobacter* sp. (MMSF_3506, MMSF_3519, MMSF_3550), *Muricauda*
sp. (MMSF_3501, MMSF_3509, MMSF_3543), *Mameliella* sp. (MMSF_3510,
MMSF_3537, MMSF_3552), and *Roseitalea* sp. (MMSF_3504, MMSF_3516,
MMSF_3546). Within a given bacterial species, all three genomes were >99.9% identical
(based on ANI and AAI calculations, Figure S1), suggesting they belong to the same strain.
Therefore, we assumed that the different bacterial isolates were present in the original *C.
proliferum* culture, before experimental evolution started, and independently evolved in the
three *C. proliferum* strains, under elevated temperature (SS strains) versus ambient
temperature (WT10 strain). We performed a single nucleotide polymorphism (SNP) analysis
to uncover which bacterial genes, if any, were impacted by host heat selection.

Across the four bacterial strains, we detected 32 SNPs between bacteria isolated from WT10
C. proliferum and the corresponding bacteria isolated from SS *C. proliferum* (Table S5). Six
SNPs were synonymous mutations and therefore not expected to bear phenotypic effects. For
each given bacterial strain, we next chose to focus on the non-synonymous mutations that
were present in both bacteria isolated from SS *C. proliferum* (Table S5, boldened rows), as
there is a higher chance these would be true SNPs and may point to convergent evolution
under heat selection. No non-synonymous SNPs were detected in *Roseitalea* sp.

In *Mameliella* sp., a missense mutation affected the *chvI* gene, a transcriptional regulator
known to be essential for the virulence of the plan pathogen *Agrobacterium tumefaciens* (50),
as well as successful establishment of symbiosis in the nodule-forming and nitrogen-fixing
plant associates *Rhizobium leguminosarum* and *Sinorhizobium meliloti* (51, 52).

In *Muricauda* sp., missense mutations affected genes encoding a heme A synthase, a histidine
kinase, and a DNA gyrase. Histidine kinases have a broad range of actions in signal
transduction, while DNA gyrases are involved in DNA replication. The impact of mutations
on those two genes is therefore hard to assess. Heme A synthases are involved in heme
synthesis, a cofactor involved in many cellular processes, including antioxidant functions
through its incorporation into hemoproteins, as well as infectivity and virulence. Intracellular
Brucella abortus are dependent on heme production for their survival (53), while heme
deficiency negatively impacts mouse tissue colonization by *Staphylococcus aureus* (54).
Additionally, excess free heme catalyzes the formation of ROS, resulting in oxidative stress
(55). If this mutation resulted in decreased heme synthesis, it may contribute to lower ROS
production during heat stress and thereby participate in the heat tolerance of SS *C.*
proliferum.

In *Marinobacter* sp., a missense mutation affected *gspE*, which encodes an ATPase involved
in the T2SS. As is the case with most secretion systems, the T2SS is required for the
virulence of a variety of pathogens (56), as well as the colonization of leech gut by the
mutualistic symbiont *Aeromonas veronii* (57). Additionally, a frameshift mutation affected
pilO, while a mutation in *pilM* resulted in the loss of a stop codon; both genes are involved in
the type IV pilus assembly. Type IV pili are involved in the formation of biofilms, but also
support bacterial motility and may play a role in host colonization (58).

In all three cases, the detected mutations in response to heat selection affected genes
potentially involved in host-symbiont interactions, such as host infection or symbiosis
establishment. Whether they enhance or are detrimental to these interactions remains
unknown, and functional studies will be needed to uncover the phenotypic effects of these
mutations.

Conclusion

We provide the first extensive genomic analysis of Symbiodiniaceae-associated bacteria, by
presenting the genomes of 49 bacteria isolated from six Symbiodiniaceae genera. Most
bacteria have the potential to be beneficial for their hosts, through the production of B

vitamins or antioxidants. *Halomonas* sp. MMSF_3323 is particularly interesting as it was
predicted to synthesize seven essential B vitamins. Isolates of the *Muricauda* genus may also
be crucial as they were the only bacteria in our data set that are able to produce antioxidant
carotenoids. *Marinobacter* sp. are also likely important for Symbiodiniaceae health as they
can produce four B vitamins, as well as many ROS-scavenging compounds, such as
ubiquinone and glutathione. It is noteworthy that the functional potential of these bacteria
was independent of their Symbiodiniaceae host. Given that these Symbiodiniaceae were
initially obtained from corals, clams, and anemones, it is unclear whether the bacteria
analyzed here would interact directly with an animal host *in hospite*, or rather only with
Symbiodiniaceae. Genomic analyses of Symbiodiniaceae-associated bacteria *in hospite* are
therefore needed to assess potential lifestyle differences.

Finally, we analyzed the effect of heat selection on genome evolution in four bacterial
species. In *Muricauda* sp., *Marinobacter* sp., and *Mameliella* sp., mutations observed in
bacteria isolated from heat-evolved Symbiodiniaceae were observed in genes potentially
involved in host-symbiont interactions. While the effect of the mutations was not
investigated, it is possible that future climate conditions will reinforce or damage interactions
between bacteria and Symbiodiniaceae. Whether this may affect coral climate resilience
remains to be investigated. More extensive genomic comparisons, using long-read
sequencing for increased accuracy, are needed to fully understand the response of
Symbiodiniaceae-associated bacteria to host heat selection. Overall, our study reveals the
beneficial potential of Symbiodiniaceae-associated bacteria.

Material and methods

Cultures of bacteria from three *Cladocopium proliferum* strains

Three *Cladocopium proliferum* cultures were selected to isolate pure bacteria from: SCF055-
01.10 (WT10), SCF055-01.03 (SS03), and SCF055-01.08 (SS08). These three *C. proliferum*
strains originated from a monoclonal culture isolated from *Acropora tenuis* (59). SS03 and
SS08 have been evolving in the laboratory after ratcheting to 31°C since 2011 while WT10
was kept at 27°C. At the time of sampling for this study (August - October 2022), cultures
were experimentally evolved for an estimated 300-330 generations (based on generation time
of the WT cultures). Symbiodiniaceae cultures were maintained in 15 mL Daigo's IMK
medium (1X), prepared with filtered Red Sea Salt water (fRSSW, 34 ppt) in sterile 50 mL
polypropylene culture flasks where media was changed fortnightly. These flasks were kept in
a 12 h light:12 h dark incubator with lighting at 50–60 μmol photons $m^{-2} s^{-1}$ of
photosynthetically active radiation (Taiwan Hipoint Corporation, model 740FHC LED, light
chambers) and temperature at either 27°C (WT) or 31°C (SS).

Symbiodiniaceae cultures cell counts were quantified (Life Technologies Countess II FL) and
an aliquot of 10^6 cells was deposited in a 5-μM mesh size strainer (pluriSelect, Germany) to
separate the Symbiodiniaceae (ranging from 6-9 μm in diameter) and their closely associated
(physically attached/intracellular) bacterial cells from loosely associated (planktonic)
bacterial cells. Strainers were sealed with parafilm and centrifuged at $10,000 \times g$ for 3 min. A

volume of 500 µL of fRSS was added to the strainers, which were sealed and centrifuged at
10,000 × g for 3 min. Symbiodiniaceae cells and their associated bacteria were resuspended
from the filter in 1 mL of fRSS. For each culture, one aliquot was not processed any further
and one other aliquot was bead beaten with 100 mg of sterile glass beads (400-600 nm) for 30
s at 30 Hz in a Tissue-lyser II (Qiagen, Germany), in order to open up Symbiodiniaceae cells
and release intracellular bacteria to facilitate their growth on agar plates. Both aliquots were
serially diluted spread onto BD Difco™ Marine Agar 2216 (MA) and three Oxoid R2A Agar
(R2A) prepared with fRSS, culture plates. Plates were incubated at 27°C (for the WT10
samples) or at 31°C (SS03 and SS08 samples) for 7 d to facilitate bacterial colony growth.
Representatives of all morphologies were subcultured to attain culture purity. Pure cultures
were stored in sterile 40% glycerol at -80°C.

16S rRNA gene sequencing from cultured bacteria

Individual pure freshly grown bacterial colonies were suspended in 20 µL sterile Milli-Q®
water, incubated for 10 min at 95°C then used as templates in PCRs. PCR amplification of
the bacterial 16S rRNA gene was performed with primers 27F and 1492R (60) in reactions
containing 2 µL template DNA, 1 µL of each primer at a 10 µM concentration (0.25 µM final
concentration), 20 µL of 2X Mango Mix (Bioline 25034) and 16 µL of PCR grade water. The
amplification cycle was: 5 min at 94°C; 30 cycles of 60 s at 94°C, 45 s at 50°C, and 90 s at
72°C; 10 min at 72°C; with a final holding temperature of 4°C. The PCR products were
purified Sanger sequenced with primer 1492R at Macrogen (South Korea). Raw sequences
were trimmed and proofread in Geneious Prime v2019.1.3. Sequences were analyzed in
BLAST. For each sequence, the closest hit and percent identity was recorded in Table S1.

DNA extraction and whole-genome sequencing

For each isolate selected for whole-genome sequencing, a single colony was picked with an
inoculation loop and snap-frozen. DNA extraction was performed on the QIAsymphony
using the DSP Virus/Pathogen Mini Kit (Qiagen). Library preparation performed using
Nextera XT (Illumina Inc.) according to manufacturer's instructions. Whole-genome
sequencing was performed on NextSeq 500/550 with a 150bp PE kit.

Genome assembly and annotation

Raw reads were trimmed and quality-filtered using Trimmomatic v0.36 (61)
(HEADCROP:10 LEADING:5 TRAILING:5 SLIDINGWINDOW:4:28 MINLEN:30). The
quality of raw reads before and after trimming was checked with FASTQC v0.11.9 (62).
Trimmed and quality-filtered reads were de novo assembled into contigs using SPAdes
v3.15.5 (63) with 21, 33, 55 and 77 k-mers and the option "--careful" was applied to
minimise the number of mismatches and short indels. All contigs with a length of <1000 bp
were removed using BBMap v38.96 (64). Subsequently, the levels of completeness and

contamination of assembled genomes were assessed using CheckM v1.2.2 using the

"lineage_wf" workflow (65). Taxonomic assignment of all assembled genomes was carried

out using GTDB-Tk v2.1.0 (66) using the "classify_wf" workflow. GTDB-Tk assigned the

taxonomy of the genomes based on 120 bacterial marker genes. A phylogenetic tree was built

in IQ-Tree v2.0.3 (67) using the best model LG+F+R4, selected by ModelFinder wrapped in

IQ-tree (68), and 1000 ultrafast bootstrap replicates (69). The tree was visualized in iTOL v6

(70). Average Nucleotide Identities (ANI) and Average Amino acid Identities (AAI) of all

assembled genomes were calculated using a genome-based matrix calculator (71) and plotted

in R using the ggplot2 package (72).

Gene prediction was performed in Bakta v1.7.0 (73) and InterProScan v5.55 using with the

parameter "-appl pfam" and evalue <1e-5 (74). Metabolic pathways, transport systems and

secretion systems were annotated via METABOLIC-G v4.0 (75) applying "-m-cutoff 0.75"

to include pathways which are ≥75% complete. The completeness of metabolic pathways

(KEGG module database, https://www.genome.jp/kegg/module.html), transporters and

secretion systems was estimated in EnrichM v0.6.4 (76). KEGG modules of interest were

then plotted in GraphPad Prism. Pfam IDs of interest (Table 1) were queried in the

InterProScan annotation, and the number of hits for each Pfam IDs per genome was plotted in

GraphPad Prism.

Single nucleotide polymorphism analysis

Sequencing reads were assembled using the genome assembler pipeline Shovill v1.1.0 (77).

Briefly, the Shovill pipeline included read trimming using Trimmomatic v0.39 (61), de novo

assembly with SPAdes v3.15.5 (63) and genome polishing with Pilon v1.24 (78). After the

pipeline, additional polishing was performed by mapping the reads back to the contigs with

BWA v0.7.17 (79) and sorting the resulting SAM/BAM files using SAMtools v1.15.1 (80).

Pilon v1.24 was then used to correct bases, fix mis-assemblies and fill gaps. The reformat.sh

script from the Bbmap package v38.76 (-minlength=1000) was used to filter out contigs less

than 1000bp (sourceforge.net/projects/bbmap/). The quality and completeness of the

assemblies was assessed with CheckM v1.2.1 (65), BUSCO v5.3.2 , and CAT/BAT v5.2.3.

The draft genome assemblies were then annotated with Bakta v1.7.0 (73).

Single nucleotide polymorphism (SNP) detection between WT and SS samples were then

performed using snippy v4.6.0 (81), where both WT and SS samples were inputted as the

reference genome in turn. Variants were then selected if they were found in both the

reciprocal snippy analyses.

Data availability

454   Genbank accession numbers for 16S rRNA sequences of cultured bacteria are OQ945078-

455   OQ945142 (see Table S1). Genome assemblies and raw NextSeq reads are available under

456   NCBI BioProject ID PRJNA971760 (see Table S2 for individual accession numbers). A

457   subset of the snippy output files is available on Zenodo at 10.5281/zenodo.8210291.

Acknowledgements

This research was supported by the Australian Research Council (FL180100036 to MJHvO) and the Native Australian Animals Trust (to JM). JL was supported by a Future Fulbright scholarship. We thank Glen Carter for assistance with genome sequencing; Kshitij Tandon and Torsten Seemann for assistance with genome analyses; the Melbourne Research Cloud, the University of Melbourne's Research Computing Services and the Petascale Campus Initiative for providing the high-performance computing instances needed for this work.

Author contributions

Conceptualization: JM, LLB, MJHvO; Formal analysis: JM, GKP; Funding acquisition: JM, MJHvO; Investigation: JM, JL, LMJ; Methodology: JM, JL; Visualization: JM, GKP; Writing - Original Draft: JM, MJHvO. All authors read and edited the manuscript.

Competing interests

The authors declare that they have no competing interests.

References

1. Fisher R, O'Leary RA, Low-Choy S, Mengersen K, Knowlton N, Brainard RE, Caley MJ. 2015. Species richness on coral reefs and the pursuit of convergent global estimates. Current Biology 25:500–505.

2. Sing Wong A, Vrontos S, Taylor ML. 2022. An assessment of people living by coral reefs over space and time. Global Change Biology https://doi.org/10.1111/gcb.16391.

3. Blackall LL, Wilson B, van Oppen MJH. 2015. Coral-the world's most diverse symbiotic ecosystem. Molecular Ecology 24:5330–5347.

4. Bourne DG, Morrow KM, Webster NS. 2016. Insights into the coral microbiome: Underpinning the health and resilience of reef ecosystems. Annual Review of Microbiology 70:317–340.

5.   Maire J, Buerger P, Chan WY, Deore P, Dungan AM, Nitschke MR, van Oppen MJH. 2022.

Effects of Ocean Warming on the Underexplored Members of the Coral Microbiome.

Integrative and Comparative Biology icac005.

6.   LaJeunesse TC, Parkinson JE, Gabrielson PW, Jeong HJ, Reimer JD, Voolstra CR, Santos SR. 2018.

Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral

endosymbionts. Current Biology 28:2570-2580.e6.

7.   Muscatine L, R. McCloskey L, E. Marian R. 1981. Estimating the daily contribution of carbon

from zooxanthellae to coral animal respiration1. Limnology and Oceanography 26:601–611.

8.   Suggett DJ, Smith DJ. 2020. Coral bleaching patterns are the outcome of complex biological and

environmental networking. Global Change Biology 26:68–79.

9.   Szabó M, Larkum AWD, Vass I. 2020. A Review: The role of reactive oxygen species in mass

coral bleaching, p. 459–488. *In* Advances in Photosynthesis and Respiration. Springer.

10.   Setter RO, Franklin EC, Mora C. 2022. Co-occurring anthropogenic stressors reduce the

timeframe of environmental viability for the world's coral reefs. PLOS Biology 20:e3001821.

11.   Huggett MJ, Apprill A. 2019. Coral microbiome database: Integration of sequences reveals high

diversity and relatedness of coral-associated microbes. Environmental Microbiology Reports

11:372–385.

12.   van Oppen MJH, Blackall LL. 2019. Coral microbiome dynamics, functions and design in a

changing world. Nature Reviews Microbiology 17:557.

13.   Maire J, Blackall LL, van Oppen MJH. 2021. Intracellular Bacterial Symbionts in Corals:

Challenges and Future Directions. Microorganisms 9:2209.

14. Lawson CA, Raina J-B, Kahlke T, Seymour JR, Suggett DJ. 2018. Defining the core microbiome of

the symbiotic dinoflagellate, *Symbiodinium*. Environmental Microbiology Reports 10:7–11.

15. Maire J, Girvan SK, Barkla SE, Perez-Gonzalez A, Suggett DJ, Blackall LL, van Oppen MJH. 2021.

Intracellular bacteria are common and taxonomically diverse in cultured and in hospite algal

endosymbionts of coral reefs. The ISME Journal 15:2028–2042.

16. Frommlet JC, Sousa ML, Alves A, Vieira SI, Suggett DJ, Serôdio J. 2015. Coral symbiotic algae

calcify ex hospite in partnership with bacteria. Proceedings of the National Academy of

Sciences of the United States of America 112:6158–6163.

17. Camp EF, Kahlke T, Nitschke MR, Varkey D, Fisher NL, Fujise L, Goyen S, Hughes DJ, Lawson CA,

Ros M, Woodcock S, Xiao K, Leggatt W, Suggett DJ. 2020. Revealing changes in the microbiome

of Symbiodiniaceae under thermal stress. Environmental Microbiology 22:1294–1309.

18. Heric K, Maire J, Deore P, Perez-Gonzalez A, Van Oppen MJH. 2023. Inoculation with

Roseovarius increases thermal tolerance of the coral photosymbiont, Breviolum minutum.

Frontiers in Ecology and Evolution 11:1079271.

19. Buerger P, Vanstone RT, Maire J, van Oppen MJH. 2022. Long-Term Heat Selection of the Coral

Endosymbiont Cladocopium C1acro (Symbiodiniaceae) Stabilizes Associated Bacterial

Communities. International Journal of Molecular Sciences 23:4913.

20. Motone K, Takagi T, Aburaya S, Miura N, Aoki W, Ueda M. 2020. A Zeaxanthin-producing

bacterium isolated from the algal phycosphere protects coral endosymbionts from

environmental stress. mBio 11:e01019-19.

21. Matthews JL, Raina J, Kahlke T, Seymour JR, van Oppen MJH, Suggett DJ. 2020.

Symbiodiniaceae-bacteria interactions: rethinking metabolite exchange in reef-building corals

as multi-partner metabolic networks. Environmental Microbiology 22:1675.

22.   Maire J, Deore P, Jameson VJ, Sakkas M, Perez-Gonzalez A, Blackall LL, Van Oppen MJH.

Assessing the contribution of bacteria to the heat tolerance of experimentally evolved coral

photosymbionts.

23.   Wyman M, Hodgson S, Bird C. 2013. Denitrifying alphaproteobacteria from the Arabian Sea

that express nosZ, the gene encoding nitrous oxide reductase, in oxic and suboxic waters. Appl

Environ Microbiol 79:2670–2681.

24.   Zhong H, Sun H, Liu R, Zhan Y, Huang X, Ju F, Zhang X-H. 2021. Comparative Genomic Analysis

of Labrenzia aggregata (Alphaproteobacteria) Strains Isolated From the Mariana Trench:

Insights Into the Metabolic Potentials and Biogeochemical Functions. Front Microbiol 12.

25.   Martyn JE, Gomez-Valero L, Buchrieser C. 2022. The evolution and role of eukaryotic-like

domains in environmental intracellular bacteria: the battle with a eukaryotic cell. FEMS

Microbiology Reviews 46:fuac012.

26.   Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG. 2005. Algae acquire vitamin B12

through a symbiotic relationship with bacteria. Nature 438:90–93.

27.   Croft MT, Warren MJ, Smith AG. 2006. Algae Need Their Vitamins. Eukaryotic Cell 5:1175.

28.   Helliwell KE. 2017. The roles of B vitamins in phytoplankton nutrition: new perspectives and

prospects. New Phytologist 216:62–68.

29.   Agostini S, Suzuki Y, Casareto BE, Nakano Y, Hidaka M, Badrun N. 2009. Coral symbiotic

complex: Hypothesis through vitamin B12 for a new evaluation. Galaxea, Journal of Coral Reef

Studies 11:1–11.

30.   Xie B, Bishop S, Stessman D, Wright D, Spalding MH, Halverson LJ. 2013. Chlamydomonas

550   reinhardtii thermal tolerance enhancement mediated by a mutualistic interaction with vitamin

551   B12-producing bacteria. ISME Journal 7:1544–1555.

552   31.   Grant MAA, Kazamia E, Cicuta P, Smith AG. 2014. Direct exchange of vitamin B12 is

553   demonstrated by modelling the growth dynamics of algal-bacterial cocultures. ISME Journal

554   8:1418–1427.

555   32.   Maire J, Tandon K, Collingro A, van de Meene A, Damjanovic K, Gotze CR, Stephenson S, Philip

556   GK, Horn M, Cantin NE, Blackall LL, Van Oppen MJH. 2023. Colocalization and potential

557   interactions of Endozoicomonas and chlamydiae in microbial aggregates of the coral

558   Pocillopora acuta. Science Advances 9:eadg0773.

559   33.   Pogoreutz C, Oakley CA, Rädecker N, Cárdenas A, Perna G, Xiang N, Peng L, Davy SK, Ngugi DK,

560   Voolstra CR. 2022. Coral holobiont cues prime Endozoicomonas for a symbiotic lifestyle. The

561   ISME Journal 2022 1–13.

562   34.   Wada N, Hsu M-T, Tandon K, Hsiao SS-Y, Chen H-J, Chen Y-H, Chiang P-W, Yu S-P, Lu C-Y, Chiou

563   Y-J, Tu Y-C, Tian X, Chen B-C, Lee D-C, Yamashiro H, Bourne DG, Tang S-L. 2022. High-resolution

564   spatial and genomic characterization of coral-associated microbial aggregates in the coral

565   Stylophora pistillata. Science Advances 8:eabo2431.

566   35.   Doering T, Maire J, Chan WY, Perez-Gonzalez A, Meyers L, Sakamoto R, Buthgamuwa I, Blackall

567   LL, van Oppen MJH. 2023. Comparing the Role of ROS and RNS in the Thermal Stress Response

568   of Two Cnidarian Models, Exaiptasia diaphana and Galaxea fascicularis. 5. Antioxidants

569   12:1057.

570   36.   Peixoto RS, Rosado PM, Leite DC de A, Rosado AS, Bourne DG. 2017. Beneficial Microorganisms

571   for Corals (BMC): Proposed mechanisms for coral health and resilience. Frontiers in

572   Microbiology 8:341.

573   37.   Dungan AM, Bulach D, Lin H, van Oppen MJH, Blackall LL. 2021. Development of a free radical

574   scavenging bacterial consortium to mitigate oxidative stress in cnidarians. Microbial

575   biotechnology 14:2025–2040.

576   38.   Maire J, van Oppen MJH. 2022. A role for bacterial experimental evolution in coral bleaching

577   mitigation? Trends in Microbiology 30:217–228.

578   39.   Doering T, Maire J, Oppen MJH van, Blackall LL. 2023. Advancing coral microbiome

579   manipulation to build long-term climate resilience. Microbiol Aust 44:36–40.

580   40.   Fiedor J, Burda K. 2014. Potential Role of Carotenoids as Antioxidants in Human Health and

581   Disease. Nutrients 6:466–488.

582   41.   Sunda W, Kieber DJ, Kiene RP, Huntsman S. 2002. An antioxidant function for DMSP and DMS

583   in marine algae. Nature 418:317–320.

584   42.   Shen B, Jensen RG, Bohnert HJ. 1997. Mannitol Protects against Oxidation by Hydroxyl

585   Radicals. Plant Physiol 115:527–532.

586   43.   Gao C, Fernandez VI, Lee KS, Fenizia S, Pohnert G, Seymour JR, Raina J-B, Stocker R. 2020.

587   Single-cell bacterial transcription measurements reveal the importance of

588   dimethylsulfoniopropionate (DMSP) hotspots in ocean sulfur cycling. Nature Communications

589   11:1942.

590   44.   Lesser MP. 1997. Oxidative stress causes coral bleaching during exposure to elevated

591   temperatures. Coral Reefs 16:187–192.

592   45.   Dungan AM, Maire J, Perez-Gonzalez A, Blackall LL, van Oppen MJH. 2022. Lack of evidence for

593   the oxidative stress theory of bleaching in the sea anemone, Exaiptasia diaphana, under

594   elevated temperature. Coral Reefs 1–12.

595   46.   Maire J, Blackall LL, van Oppen MJH. 2021. Microbiome characterization of defensive tissues in

596   the model anemone Exaiptasia diaphana. BMC Microbiology 21:152.

597   47.   Sweet M, Villela H, Keller-Costa T, Costa R, Romano S, Bourne DG, Cárdenas A, Huggett MJ,

598   Kerwin AH, Kuek F, Medina M, Meyer JL, Müller M, Pollock FJ, Rappé MS, Sere M, Sharp KH,

599   Voolstra CR, Zaccardi N, Ziegler M, Peixoto R. 2021. Insights into the Cultured Bacterial Fraction

600   of Corals. mSystems 6:e0124920.

601   48.   Maor-Landaw K, Eisenhut M, Tortorelli G, van de Meene A, Kurz S, Segal G, van Oppen MJH,

602   Weber APM, McFadden GI. 2023. A candidate transporter allowing symbiotic dinoflagellates to

603   feed their coral hosts. 1. ISME COMMUN 3:1–7.

604   49.   Jia B, Hao L, Xuan YH, Jeon CO. 2018. New Insight Into the Diversity of SemiSWEET Sugar

605   Transporters and the Homologs in Prokaryotes. Frontiers in Genetics 9.

606   50.   Mantis NJ, Winans SC. 1993. The chromosomal response regulatory gene chvI of

607   Agrobacterium tumefaciens complements an Escherichia coli phoB mutation and is required

608   for virulence. J Bacteriol 175:6626–6636.

609   51.   Bélanger L, Dimmick KA, Fleming JS, Charles TC. 2009. Null mutations in Sinorhizobium meliloti

610   exoS and chvI demonstrate the importance of this two-component regulatory system for

611   symbiosis. Mol Microbiol 74:1223–1237.

612   52.   Vanderlinde EM, Yost CK. 2012. Mutation of the sensor kinase chvG in Rhizobium

613   leguminosarum negatively impacts cellular metabolism, outer membrane stability, and

614   symbiosis. J Bacteriol 194:768–777.

53.   Almirón M, Martínez M, Sanjuan N, Ugalde RA. 2001. Ferrochelatase is present in Brucella

abortus and is critical for its intracellular survival and virulence. Infect Immun 69:6225–6230.

54.   Hammer ND, Reniere ML, Cassat JE, Zhang Y, Hirsch AO, Indriati Hood M, Skaar EP. 2013. Two

heme-dependent terminal oxidases power Staphylococcus aureus organ-specific colonization

of the vertebrate host. mBio 4:e00241-13.

55.   Chiabrando D, Vinchi F, Fiorito V, Mercurio S, Tolosano E. 2014. Heme in pathophysiology: a

matter of scavenging, metabolism and trafficking across cell membranes. Frontiers in

Pharmacology 5.

56.   Cianciotto NP, White RC. 2017. Expanding Role of Type II Secretion in Bacterial Pathogenesis

and Beyond. Infect Immun 85:e00014-17.

57.   Maltz M, Graf J. 2011. The Type II Secretion System Is Essential for Erythrocyte Lysis and Gut

Colonization by the Leech Digestive Tract Symbiont Aeromonas veronii. Appl Environ Microbiol

77:597–603.

58.   Craig L, Forest KT, Maier B. 2019. Type IV pili: dynamics, biophysics and functional

consequences. 7. Nat Rev Microbiol 17:429–440.

59.   Chakravarti LJ, Beltran VH, van Oppen MJH. 2017. Rapid thermal adaptation in photosymbionts

of reef-building corals. Global Change Biology 23:4675–4688.

60.   Lane D. 1991. 16S/23S rRNA sequencing, p. 115–47. *In* Stackebrandt, E, Goodfellow, M (eds.),

Nucleic Acid techniques in bacterial systematics. John Wiley & Sons, New York.

61.   Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics (Oxford, England) 30:2114–2120.

62.   Andrews S. 2010. FASTQC. A quality control tool for high throughput sequence data |

637   BibSonomy. https://www.bibsonomy.org/bibtex/f230a919c34360709aa298734d63dca3.

638   Retrieved 17 October 2022.

639   63.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI,

640   Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA.

641   2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing.

642   J Comput Biol 19:455–477.

643   64.   Bushnell B, Rood J, Singer E. 2017. BBMerge – Accurate paired shotgun read merging via

644   overlap. PLOS ONE 12:e0185056.

645   65.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the

646   quality of microbial genomes recovered from isolates, single cells, and metagenomes. Genome

647   Research 25:1043–1055.

648   66.   Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. 2020. GTDB-Tk: a toolkit to classify genomes

649   with the Genome Taxonomy Database. Bioinformatics 36:1925–1927.

650   67.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A, Lanfear R,

651   Teeling E. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in

652   the Genomic Era. Molecular Biology and Evolution 37:1530–1534.

653   68.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast

654   model selection for accurate phylogenetic estimates. 6. Nat Methods 14:587–589.

655   69.   Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the

656   Ultrafast Bootstrap Approximation. Molecular Biology and Evolution 35:518–522.

70. Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. Nucleic Acids Research 49:W293–W296.

71. Rodriguez-R LM, Konstantinidis KT. 2016. The enveomics collection: a toolbox for specialized
analyses of microbial genomes and metagenomes
https://doi.org/10.7287/PEERJ.PREPRINTS.1900V1.

72. Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

73. Schwengers O, Jelonek L, Dieckmann MA, Beyvers S, Blom J, Goesmann A. 2021. Bakta: rapid
and standardized annotation of bacterial genomes via alignment-free sequence identification.
Microb Genom 7:000685.

74. Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A,
Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong SY, Lopez R, Hunter S.
2014. InterProScan 5: genome-scale protein function classification. Bioinformatics 30:1236–
1240.

75. Zhou Z, Tran PQ, Breister AM, Liu Y, Kieft K, Cowley ES, Karaoz U, Anantharaman K. 2022.
METABOLIC: high-throughput profiling of microbial genomes for functional traits, metabolism,
biogeochemistry, and community-scale functional networks. Microbiome 10:33.

76. Boyd J, Woodcroft B, Tyson GW. 2019. Comparative genomics using EnrichM.
https://github.com/geronimp/enrichM. Retrieved 4 May 2023.

77. Seemann T. 2016. Shovill. GitHub Available at: github.com/tseemann/shovill.

78. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman
J, Young SK, Earl AM. 2014. Pilon: An Integrated Tool for Comprehensive Microbial Variant
Detection and Genome Assembly Improvement. PLOS ONE 9:e112963.

79.   Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.

680   Bioinformatics 25:1754–1760.

681   80.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000

682   Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and

683   SAMtools. Bioinformatics 25:2078–2079.

684   81.   Seemann T. 2015. Snippy: rapid haploid variant calling and core SNP phylogeny. GitHub

685   Available at: github.com/tseemann/snippy.