Title: Somatosensory cortex microstimulation modulates primary motor and ventral premotor cortex neurons with extensive spatial convergence and divergence

Abbreviated Title: S1-ICMS effects in M1 and PMv

Authors: Brandon Ruszala[1], Kevin A. Mazurek[2,3], and Marc H. Schieber[1,2,3*]

Affiliations:
[1]Department of Biomedical Engineering, University of Rochester, Rochester, NY, 14627
[2]Department of Neurology, University of Rochester, Rochester, NY, 14642
[3]Department of Neuroscience, University of Rochester, Rochester, NY 14642

*Corresponding Author:
       Marc H. Schieber
       E-mail: mschiebe@ur.rochester.edu

Number of:
    Pages -   **19**
    Figures -   **9**
    Tables -   **2**
    Multimedia -   **0**
    3D models -   **0**
    Words in:
        Abstract -   **170**
        Significance Statement   **100**
        Introduction -   **638**
        Discussion -   **1,497**

Conflict of interest:

The authors declare no conflicts of interest.

Acknowledgements:

The authors thank Gil Rivlis for custom task-control and stimulation control software, and Marsha Hayles for editorial comments. This work was supported by grants F31NS129099 (BR), F32NS093709 (KAM), and R01NS107271 (MHS) from the National Institute of Neurological Disorders and Stroke. The funders had no role in study design, data collection or interpretation, nor in the decision to submit the work for publication. KAM is currently a Data Scientist in the Neurology Artificial Intelligence Program, Department of Neurology, Mayo Clinic, Rochester, MN, 55905

Author contributions

BR, KAM and MHS designed the study. KAM collected and processed the data and performed some preliminary analyses. BR performed the analyses and wrote the manuscript. MHS supervised, provided resources, and edited the manuscript.

48 **Abstract**
49
Intracortical microstimulation (ICMS) is known to affect distant neurons transynaptically, yet the extent
to which ICMS pulses delivered in one cortical area modulate neurons in other cortical areas remains
largely unknown. Here we assessed how the individual pulses of multi-channel ICMS trains delivered in
the upper extremity representation of the macaque primary somatosensory area (S1) modulate neuron
firing in the primary motor cortex (M1) and in the ventral premotor cortex (PMv). S1-ICMS pulses
modulated the majority of units recorded both in the M1 upper extremity representation and in PMv,
producing more inhibition than excitation. Effects converged on individual neurons in both M1 and PMv
from extensive S1 territories. Conversely, effects of ICMS delivered in a small region of S1 diverged to
wide territories in both M1 and PMv. The effects of this direct modulation of M1 and PMv neurons
produced by multi-electrode S1-ICMS like that used here may need to be taken into account by
bidirectional brain-computer interfaces that decode intended movements from neural activity in these
cortical motor areas.

Significance Statement

Although ICMS is known to produce effects transynaptically, relatively little is known about how ICMS
in one cortical area affects neurons in other cortical areas. We show that the effects of multi-channel
ICMS in a small patch of S1 diverge to affect neurons distributed widely in both M1 and PMv, and
conversely, individual neurons in each of these areas can be affected by ICMS converging from much of
the S1 upper extremity representation. Such direct effects of ICMS may complicate the decoding of
motor intent from M1 or PMv when artificial sensation is delivered via S1-ICMS in bidirectional brain-
computer interfaces.

Keywords

Brain-computer interface, Brain stimulation, Instruction, Intracortical microstimulation, Excitation,
Inhibition

Introduction

Intracortical microstimulation (ICMS) injects brief microampere current pulses into cortical gray matter, eliciting action potentials in neural elements in the vicinity of the electrode tip. Early work found that single 40 µA pulses delivered in cat motorsensory cortex elicited action potentials in neuron somata up to 400 µm away (Stoney et al., 1968). Subsequent studies have demonstrated that ICMS pulses delivered in mouse and cat visual cortex also elicit action potentials in local axons of passage, producing antidromic action potentials in sparse populations of cortical neurons up to millimeters from the electrode tip (Histed et al., 2009). A recent computational model of macaque somatosensory cortex further suggests that all ICMS-elicited action potentials originate in axons, and that the radius of activated axons increases with increasing current through the typical range from 15 to 100 µA. Antidromic propagation of those elicited action potentials leads to an increasing density of antidromically activated somata over a distance up to ~2 mm, and while less frequent, orthodromic propagation of those action potentials can elicit somatic action potentials transsynaptically, which may affect neurons at even further distances (Kumaravelu et al., 2022). Indeed, individual ICMS pulses delivered in macaque primary motor cortex (M1) or ventral premotor cortex (PMv) elicit action potentials in local corticospinal neurons transsynaptically (Maier et al., 2013).

However they originate, orthodromically conducting action potentials elicited by ICMS can produce transsynaptic effects in additional neurons at considerable distances. While individual ICMS pulses generally may be subthreshold for eliciting action potentials transsynaptically, such effects become prominent when ICMS pulses are delivered in high-frequency trains ($\geq$ 200 Hz), enabling temporal summation of successive excitatory postsynaptic potentials (Jankowska et al., 1975; Hussin et al., 2015). Trains of ICMS pulses delivered in motor cortex excite distant spinal motoneurons and produce muscle contractions (Asanuma and Ward, 1971; Andersen et al., 1975). Delivered at lower frequencies in the primary somatosensory cortex (S1), ICMS pulse trains elicit somatosensory percepts (Romo et al., 1998; Flesher et al., 2016; Callier et al., 2020).

Though examined in relatively few studies, transsynaptic effects from individual ICMS pulses also can modulate the firing of neurons well beyond the cortical locale of the stimulating electrode. Single pulses delivered in macaque ventral premotor cortex (PMv) modulate the firing of many M1 neurons, and vice versa, single pulses delivered in M1 modulate the firing of many PMv neurons (Kraskov et al., 2011). ICMS pulses delivered in rodent somatosensory can produce activity dependent plasticity in neurons of the rostral forelimb motor area (Averna et al., 2019). Understanding such cortico-cortical effects of ICMS pulses, in addition to advancing our knowledge of both how ICMS is perceived and how ICMS influences other unperceived brain activity (Mazurek and Schieber, 2019; Mazurek and Schieber, 2021), may have practical applications. In bidirectional brain-computer interfaces (O'Doherty et al., 2011; Flesher et al., 2021; Pandarinath and Bensmaia, 2022), for example, ICMS delivered in the primary somatosensory cortex (S1) to provide artificial tactile feedback might modulate the discharge of M1 neurons being used to decode movement intent. Such ICMS-induced modulation could reduce the performance of a decoding algorithm that uses the activity of those M1 neurons as inputs (Shelchkova et al., 2022).

We therefore tested the hypothesis that ICMS pulses delivered in macaque S1 directly modulate the firing of neurons in two cortical motor areas known to receive somatosensory input. In M1, neurons respond to both cutaneous and deep peripheral stimulation (Rosen and Asanuma, 1972; Lemon and Porter, 1976; Wong et al., 1978; Fetz et al., 1980). In PMv, neurons also respond to both cutaneous and deep peripheral

125   stimulation, but have larger and more complex somatosensory receptive fields (Rizzolatti et al., 1981;
126   Gentilucci et al., 1988; Rizzolatti et al., 1988; Graziano and Gandhi, 2000). We went on to examine the
127   spatial distribution of the modulation that S1-ICMS pulses produced in each of these cortical motor areas.
128
129

Materials and Methods

We performed new analyses on data collected during a previously reported study (Mazurek and Schieber,
2017a). Methods for behavioral training, neural recording, and delivering ICMS, all have been described
there in detail and are summarized here only briefly, whereas the new experimental design and statistical
analyses for the present work are described here in detail.

Non-human Primates

Two male rhesus monkeys (Monkeys L and X, weight 9 and 11 kg, age 11 and 13 years old, respectively)
were subjects in the present study. All procedures for the care and use of these non-human primates
followed the Guide for the Care and Use of Laboratory Animals and were approved by the University
Committee on Animal Resources at the University of Rochester, Rochester, New York.

Initial Training with Visual Instructions

Each monkey was initially trained to perform a reach-grasp-manipulate (RGM) task as described in detail
previously (Mollazadeh et al., 2011; Mazurek and Schieber, 2017a). The monkey initiated each trial by
reaching to a central home object, grasping it, and pulling it for a variable initial hold period (700-1500
ms). Then the monkey was instructed to reach to, grasp, and manipulate 1 of 4 peripheral objects by
illumination of a ring of blue LEDs around the base of that object. The 4 objects were arranged at 45º
intervals 13 cm radially from the home object. The monkey pushed a button, turned a sphere, pulled a
coaxial cylinder (coax), or pulled a perpendicular cylinder (perp). The monkey then held the object in the
manipulated position for a final hold period of 700-1000 ms. Three hundred milliseconds after the
successful completion of a trial, a water reward was delivered, and the trial ended. A 1000 ms inter-trial
interval occurred before another trial could be initiated. Trials for each target object were presented
pseudo-randomly in blocks. Each block included one trial for each target object, and the order of the 4
instructed objects within a block was re-randomized between blocks. Errors occurred if the monkey failed
to hold the home object for the required duration, failed to release the home object within 1000 ms of the
instruction onset, contacted the wrong target object, failed to contact a peripheral object within 1000 ms
of releasing the center object, or failed to hold the target object for the required duration. When an error
occurred, trials were aborted immediately, no reward was given, and the same object was instructed again
on the next trial to prevent the monkey from avoiding trials involving a particular peripheral object. Trials
following an error were excluded from analysis because the monkey could have known which object
would be instructed. The RGM task was controlled by a custom software written in TEMPO (Reflective
Computing, Olympia, WA).
166

Neural Recordings

Once trained to perform the RGM task instructed with visual cues, each monkey was implanted with
multiple 16-electrode (impedance ~0.5 MΩ, 70% Pt, 30% Ir) floating microelectrode arrays (FMAs,
Microprobes for Life Sciences, Gaithersburg, MD) in the primary somatosensory cortex (S1), primary
motor cortex (M1), and premotor cortex (PM) of
the left hemisphere, as described previously
(Mollazadeh et al., 2011). The present analyses
are based on ICMS pulses delivered via 4 S1
arrays in each monkey while spikes were recorded
from 4 M1 arrays in each monkey and from 4 PM
arrays in monkey L, but from only 2 PM arrays in
monkey X. In monkey L, all 4 PM FMAs were
located in the ventral premotor cortex (PMv),
while in monkey X no arcuate spur was present
and the more medial array (D) may have been in
the dorsal premotor cortex. Nevertheless, for
present purposes we will refer to all of these PM
arrays as having been in PMv. The intracortical
locations of electrode tips were confirmed by the
presence of single- or multi-unit activity. Neural
data were recorded with a Multi Acquistion
Processor and SortClient software (Plexon, Inc.,
Dallas, TX). After 1,000 – 32,000x amplification,
waveforms that crossed a threshold set online by
the user were sampled at 40 kHz and saved for
off-line sorting along with behavioral event
markers generated by the TEMPO system. After
off-line sorting (Plexon Offline Sorter), the signal-
to-noise ratio (SNR) and the estimated percentage
of inter-spike interval (ISI) violations (< 1ms)
were used to classify each sorted unit as a definite
single unit (SNR ≥ 3 and no ISIs < 1ms), a

Figure 1. Location of microelectrode arrays implanted in Monkey L (A) and Monkey X (B). Arrays (~2x2 mm) were implanted in primary motor cortex (M1, blue), ventral premotor cortex (PMv, green), and primary somatosensory cortex (S1, yellow). In Monkey X, data from ICMS delivered via array M (pale yellow) were not used in the present study because the high frequency (225 Hz) resulted in inter-pulse intervals (4.4 ms) too short for reliable assessment of modulation. Letters next to each array are used to identify that array. AS – arcuate sulcus, CS – central sulcus, and IPS – intraparietal sulcus. Orientation arrows indicate medial (M) and caudal (C) directions. Modified with permission from Mazurek & Schieber, 2017.

probable single unit (SNR ≥ 2.5 and <10% ISIs < 1ms), a multi-unit (SNR ≥ 1.5 and <90% ISIs < 1ms),
or noise (SNR < 1.5 or ≥ 90% ISIs < 1ms) (Rouse and Schieber, 2016).

Learning to Use ICMS Instructions

Several months after the microelectrode arrays had been implanted, each monkey was trained to perform
the RGM task using ICMS pulse trains delivered in S1 as instructions, instead of the visual LED cues
(Mazurek and Schieber, 2017a). ICMS trains consisted of symmetric, biphasic, cathodic-leading, 200 µs
per phase, 1-64 µA pulses delivered at a constant frequency. For monkey L, ICMS pulse frequency was
100 Hz for all target objects, whereas for monkey X, ICMS pulse trains had a different frequency for each
object: button – 75 Hz, sphere – 100 Hz, perp – 150 Hz, coax – 225 Hz. ICMS pulses were produced by a
constant-current IZ2 Neural Stimulator controlled by an RZ5 BioAmp Processor hosted by a PC running
the OpenEx Software Suite (Tucker-Davis Technologies, Gainesville, FL).

Each of the 4 objects was instructed with a train of ICMS pulses delivered through electrodes on a
different S1 array. ICMS pulses for each instruction were delivered simultaneously through a set of 3-7
electrodes all on the same S1 array. At first, these ICMS pulse trains were delivered concurrently with the
LED cues, i.e. stimulation on the S1 array assigned to the instructed target was delivered simultaneously
with illumination of the LED ring surrounding that target. Then, over multiple subsequent daily sessions,
the LEDs were gradually dimmed until they could remain completely off while the monkey continued to
perform well. At this point, ICMS trains served as the only instruction for the correct target object on each
ICMS-instructed trial, with no visual cue. ICMS trains were delivered throughout the reaction time and
movement time on each trial (~400 — 600 ms), from instruction onset (the first ICMS pulse) until the
monkey had manipulated an object. Ultimately, trials with only visual instructions or only ICMS
instructions could be interleaved randomly and the monkeys each continued to perform well. The data
used in the present analyses were recorded in four such sessions from monkey L and three from monkey
X.

227
228 *Experimental Design and Statistical Analysis*

229
230 All analyses presented here were performed off-line using MATLAB (MathWorks, Natick, MA).

231
232   Peristimulus Time Histograms

233
To determine whether M1 and PMv neurons were directly modulated by pulses of S1-ICMS, we
constructed four peristimulus time histograms (PSTHs) for each recorded unit – one PSTH triggered on
pulse artifacts from each of the four different ICMS instructions. The time of each ICMS pulse artifact
was identified offline by discriminating the waveforms of stimulation artifacts on recording channels
from M1 or PMv in which any spikes were much smaller than the recorded artifacts. Each rhythmic train
of artifacts could be unambiguously assigned to ICMS pulses from a given S1 array based on the shapes
of the four different artifact waveforms and knowledge of which object was instructed on each trial. The
time window of each PSTH was chosen to span from 20% of the inter-pulse interval before the timestamp
of the triggering ICMS pulse artifact until the
timestamp of the next ICMS pulse artifact. For
example, a 100 Hz pulse train would have 10 ms
inter-pulse intervals, and therefore the PSTH
window spanned from -2 ms before to +10 ms
after the trigger time. A PSTH bin-width of 0.1 ms
was chosen to provide high temporal resolution.
Raw spike counts in each bin were divided by the
total number of ICMS trigger pulses, such that the
PSTH ordinate represented the probability of the
neuron firing a spike in each 0.1 ms bin
(Guggenmos et al., 2013). Figure 2 illustrates one
of the four PSTHs (for instruction 1, the button)
for a definite single unit in M1.

256
257 In forming these PSTHs, we blanked any neural
258 data from -0.2 ms before to +0.6 ms after the
259 timestamp assigned to each stimulus artifact by

Figure 2. Example peristimulus time histogram. Across multiple RGM trials, this definite single unit from M1 fired a total of 1011 spikes (nSpikes) during ICMS Instruction 1 (button). Spike counts in each 0.1ms bin were converted to probabilities by dividing the count in each bin by the number of ICMS pulse artifacts used as triggers to form the PSTH (nTriggers). Pink rectangles show the artifact window during which spikes could not be reliably discriminated. Horizontal black lines represent the mean (solid line) ±2 standard deviations (dashed lines) of the baseline bins preceding the first artifact window. The average waveform of the unit is shown in the upper right corner of the PSTH, with overlapped individual spikes in gray and a 1ms timescale bar below in black. Ch78U2, L, 20150729.

Offline Sorter, as we considered any spike waveform on an M1 or PMv recording channel that occurred
during this interval to have been potentially distorted by collision with the ICMS pulse artifact. This
produced a minimum blanked window in each PSTH from -0.2 ms to +0.6 ms around the trigger time at
0.0 ms. ICMS pulse artifacts varied across different recording channels, however, often having
waveforms different from the artifacts in the channels used to trigger each PSTH. In some recording
channels, the artifacts could begin more than -0.2 ms before the timestamp or last longer than +0.6 ms
after the timestamp. We therefore expanded the duration of the artifact window separately for each PSTH,
using the 15th percentile of all the bins in each PSTH as a threshold. We defined the first bin that
exceeded the 15th percentile proceeding backward from time -0.2 ms to be the start and the first bin that
exceeded the 15th percentile proceeding forward from time +0.6 ms to be the end of the artifact window
on that channel. Any spikes within this "final artifact window" (shaded pink in all PSTH figures) were
excluded from subsequent analyses.

In one situation, however, we terminated the final
artifact window earlier than the 15th percentile
threshold crossing proceeding forward from time
+0.6 ms. As illustrated by the example PSTH of
Fig. 3A, the artifact trigger time at 0.0 ms
sometimes was followed by a prolonged series of
bins with zero spike counts, suggesting that the
ICMS pulses produced early inhibition of the unit.
We therefore set a maximum termination time for
artifact windows to distinguish early inhibition
following ICMS pulses from spikes lost due to
collisions with the ICMS artifacts. To determine
this maximum termination time, we identified the
time of the earliest spike discriminated more than
the +0.6 ms minimum after the artifact trigger for
each of the 596 constructed PSTHs. We then
compiled a histogram of those earliest spike times
(Fig. 3B), which appeared bimodal. We
interpreted the first mode to represent PSTHs in
which spikes occurred promptly following the
ICMS pulse artifact (i.e. without early inhibition)
and the second mode to represent PSTHs in which
spikes resumed after a period of early inhibition.
We fit a generalized Gaussian mixture model to
the bimodal distribution and used the nadir
between the two components at 2.3 ms after the
trigger time at 0.0 ms as the maximum possible
termination of an ICMS pulse artifact window.
Any artifact window (determined using the 15th
percentile criterion) that lasted longer than 2.3 ms

Figure 3. Distinguishing early inhibition from collisions with ICMS pulse artifacts. *A) Example PSTH in which a period of early inhibition followed the ICMS pulse artifact. The final artifact window began -0.5 ms before and ended 2.3 ms after the trigger time at 0.0 ms. (Ch114U1, L, 20150729). B) For every PSTH, the time of the earliest spike discriminated more than the +0.6 ms minimum after the artifact trigger at 0.0 ms (indicated by the dark grey rectangle) was collected and binned in 0.1 ms steps. The resulting distribution appeared bimodal, and a generalized Gaussian mixture model with two components was fit to the data. The first component (dashed red curve) was interpreted as spikes resuming as soon as the artifact ended with no immediately following inhibition, whereas the second component (solid red curve) was interpreted as spikes resuming after an early inhibitory period immediately following the artifact. The nadir between the two components at 2.3 ms therefore was set as the maximum time following the trigger at which the final artifact window could terminate.*

303 therefore was terminated at 2.3 ms. Whereas bins during the final artifact window were excluded from
304 subsequent analyses, a series of zero-count bins beyond 2.3 ms was considered to indicate a period of
305 early inhibition rather than collisions with artifacts, and therefore was included in subsequent analyses.

The number of triggers used to construct PSTHs varied for a number of reasons. First, while ICMS was delivered at constant frequencies through the reaction and movement epochs, the monkeys' reaction and movement times varied from trial to trial. When reaction and movement times were longer, more ICMS pulses were available as triggers. Second, for monkey X the instruction for each object had a different frequency (button – 75 Hz, sphere – 100 Hz, or perp – 150 Hz) and higher frequency pulse trains therefore provided more triggers. And third, ICMS artifacts occasionally were distorted by collision with other waveforms and consequently were unrecoverable in the Offline Sorter. For each object, Table 1 summarizes the medians and interquartile ranges across all trials in all sessions for each monkey's reaction times (Rx), movement times (Mv), and response times (Rsp, reaction + movement times), as well as the number of triggers (nTrig) used to construct PSTHs.

Behavioral Metrics		Button			Sphere			Perp			Coax		
		Med	25th	75th	Med	25th	75th	Med	25th	75th	Med	25th	75th
L	Rx	390	343	480	293	262	322	302	274	332	356	318	409
	Mv	275	249	303	249	226	280	223	201	243	223	209	241
	Rsp	668	622	774	542	510	574	522	493	551	581	541	639
	nTrig	**12519**	**11243**	**13804**	**7906**	**5576**	**10264**	**7958**	**6742**	**9074**	**10463**	**8355**	**12773**
X	Rx	597	429	974	581	520	685	466	417	531			
	Mv	225	205	251	135	127	147	131	123	143			
	Rsp	841	653	1207	724	656	850	604	547	671			
	nTrig	**11683**	**9586**	**13620**	**8507**	**7869**	**8727**	**7726**	**6900**	**10453**			

Table 1. Behavioral metrics during ICMS-instructed trials determining the number of triggers available for PSTH construction. For each object and each monkey, the median, the 25th percentile, and the 75th percentile across all trials in all sessions is given for: Rx – reaction time, Mv – movement time, Rsp – total response time, nTrig – number of triggers.

Modulation in PSTHs

In the PSTH of each neuron-array pair, we identified excitatory peaks and/or inhibitory troughs when ≥ 3 consecutive bins went beyond ± 2 standard deviations of the baseline mean calculated from the bins preceding the final artifact window. In cases for which the mean minus 2 standard deviations were less than zero, we required 7 consecutive bins with counts of zero to identify a trough. For 59 pairs we did not attempt to identify troughs because the total spike count was less than the number of bins in the PSTH (i.e. <100 spikes for 100 Hz pulse trains leading to 100 bins, <133 spikes for 75 Hz pulse trains leading to 133 bins, etc.), producing many bins that contained zero spikes by chance alone.

We then measured the latency, full width at half maximum (FWHM), and amplitude of each peak or trough, as illustrated in Figure 4. Latencies were calculated as the time from the beginning of the final artifact window to the first bin in which a peak or trough crossed above or below 2 standard deviations from the mean of the baseline spike probability. FWHM was calculated as the difference in time between the bins in which a peak or trough crossed 50% of its maximum or minimum value with respect to the baseline mean. We quantified amplitude as the maximum of a peak or the minimum of a trough minus the baseline mean, normalized as a percentage of the baseline mean. For peaks, we also calculated the cumulative probability as the sum of the probability in each bin from onset to offset of the peak.

Code Accessibility

Analysis-specific code and data are available from the Lead Contact upon request.

Figure 4. Peak and trough characteristics. *Peaks and troughs were identified when the PSTH crossed 2 standard deviations (horizontal dashed lines) beyond the mean (horizontal solid line) of the baseline period before the final artifact window (pink rectangles). The onset latency, offset time, FWHM, and amplitude of each peak (purple) or trough (orange) were calculated as described in the text. In this example, the onset and offset times of the trough happened to coincide with those of the half minimum.*

Results

During each session, neuron spiking data were collected from all the 16-channel microelectrode arrays implanted in M1 and PMv of each monkey. After off-line sorting, each sorted waveform was classified as a definite single unit (DSU), probable single unit (PSU), or multi-unit (MU) according to criteria described in the Methods. Units were tracked across consecutive days based on their mean waveforms, mean firing rates, autocorrelograms, and cross-correlograms with other units (Fraser and Schwartz, 2012). Units found to be present in multiple sessions were analyzed only once, from the first session in which they appeared. In Monkey L, 81 units were analyzed – 38 from M1 and 42 from PMv. The 38 M1 units were comprised of 9 DSUs, 8 PSUs, and 21 MUs. The 42 PMv units were comprised of 5 DSUs, 12 PSUs, and 25 MUs. In Monkey X, 69 units were analyzed – 36 from M1 and 33 from PMv. The 36 M1 units were comprised of 6 DSUs, 14 PSUs, and 16 MUs. The 33 PMv units were comprised of 2 DSUs, 6 PSUs, and 25 MUs. To obtain the best available estimate of the prevalence of excitatory and inhibitory effects, we chose to include not only DSUs and PSUs, but also MUs in the analyzed population. The presence of a peak or trough in the PSTH of an MU demonstrated that at least one neuron recorded at that location was modulated by the S1-ICMS pulses.

Direct modulation of neurons in M1 and PMv produced by S1-ICMS

Figure 5 displays four PSTHs constructed with spikes from the same definite single unit recorded in M1 of Monkey L. Each PSTH was triggered on the artifacts of ICMS pulses from one of the four ICMS instructions, each delivered via electrodes on a different S1 array (colored red). Whereas the ICMS pulses of instructions 1 and 4 did not modulate this neuron's discharge (Fig. 5A, 5D), the ICMS pulses of instructions 2 and 3 did,

Figure 5. Peristimulus time histograms triggered on ICMS pulse artifacts from the instruction for each target object. *All four PSTHs show spikes from the same definite single unit in M1, triggered on ICMS pulse artifacts from the A) button, B) sphere, C) perpendicular cylinder, or D) coaxial cylinder instruction. Spike counts in each 0.1 ms bin were converted to probabilities by dividing the count in each bin by the number of ICMS pulse artifacts used as triggers to form the PSTH (nTriggers). Pink rectangles show the artifact window during which spikes could not be reliably discriminated. Insets within the left pink rectangle illustrate the arrays implanted in S1 posterior to the central sulcus (CS), with the stimulated array colored red and the others yellow. Horizontal black lines represent the mean (solid line) ± 2 standard deviations (dashed lines) of the baseline bins preceding the first artifact window. The average waveform of the unit is shown in the upper right corner of the PSTH in A, with overlapped individual spikes in gray and a 1 ms timescale bar below in black. Note that the baseline firing probability varied among the four PSTHs because the neuron fired differently during trials involving the four different objects. Ch109U1, L, 20150729*

384 producing different patterns of peaks and/or troughs in their respective PSTHs (Fig. 5B, 5C). We refer to
385 such evidence of increased and/or decreased spike probability as "direct modulation."
386
ICMS instructions delivered through electrodes on the four different S1 arrays in each monkey provided
152 M1 neuron-array pairs and 168 PMv pairs in monkey L; 144 M1 pairs and 132 PMv pairs in monkey
X. The pairs involving one S1 array in monkey X (colored pale yellow in Figure 1) were excluded from
analysis, however, because the high stimulation frequency used for that array (225 Hz) resulted in inter-
pulse intervals (4.4 ms) too short for reliable assessment of modulation. The remaining 108 M1 and 99
PMv neuron-array pairs from monkey X were included in the subsequent analyses.
393
For M1 neurons, we identified at least one peak or trough that exceeded ± 2 standard deviations of the
pre-stimulus baseline (see Methods) in the PSTHs of 85/152 (55.9%) neuron-array pairs from monkey L
and 44/108 (40.7%) pairs from monkey X; for PMv neurons, 45/168 (26.8%) pairs from monkey L and
35/99 (35.4%) pairs from monkey X. We also assessed the percentage of neurons that were directly
modulated by ICMS delivered through electrodes on at least one S1 array. For M1 neurons, 33/38 (87%)
in monkey L and 28/36 (78%) in monkey X were directly modulated by ICMS delivered on at least one
S1 array; for PMv neurons, 26/42 (62%) in monkey L and 29/33 (88%) in monkey X. The majority of
neurons recorded in both M1 and PMv were directly modulated by ICMS pulses delivered on at least one
S1 array.
403
404 Combining the data from the two monkeys, we identified 46 peaks in M1 and 44 in PMv; 160 troughs in
405 M1 and 77 in PMv. The frequencies of peaks vs. troughs differed in M1 vs PMv (X^2-test, p = 0.0052).
406 Overall, inhibition was more common than excitation, more so in M1 than in PMv.
407
Although single peaks or troughs were most
frequent, many PSTHs showed multiple effects, as
exemplified by the two troughs in Fig. 5B or the
peak and trough in Fig. 5C. Table 2 gives the
numbers of PSTHs with single peaks, single
troughs, or multiple effects in M1, in PMv, and in
the two areas combined, each broken down by
whether the unit was a DSU, PSU, or MU. We
considered the possibility that MUs might be
more likely to show multiple effects, but pooling
the effects from M1 and PMv showed no
dependence between unit quality and single peak
vs. single trough vs. multiple effects (X^2-test,
p=0.088). Comparing the DSU+PSU+MU totals
for M1 vs. PMv, however, did confirm that the
frequencies of single peak vs. single trough vs.

Region	Unit Quality	1 peak	1 trough	Multiple	Total
M1	DSU	1	6	6	**13**
	PSU	2	23	9	**34**
	MU	2	43	37	**82**
	Total	**5**	**72**	**52**	**129**
PMv	DSU	3	0	1	**4**
	PSU	6	9	4	**19**
	MU	11	25	21	**57**
	Total	**20**	**34**	**26**	**80**
M1 + PMv	DSU	4	5	8	**17**
	PSU	8	32	13	**53**
	MU	13	68	58	**139**
	Total	**25**	**105**	**79**	**209**

Table 2. Types of direct modulation. *M1 – primary motor cortex, PMv – premotor cortex, DSU – definite single unit, PSU – probable single unit, MU – multi-unit*

424 multiple effects differed in the two cortical areas (X^2 test, p=2.82e-05), with troughs and multiple effects
425 being more frequent in M1 and peaks more frequent in PMv.
426
427 *Latency, full-width half-maximum, and amplitude of direct modulation effects in M1 and PMv neurons*
428
429 We then quantified the latency, full-width at half maximum, and amplitude of each peak or trough. Fig.
430 6A shows the distributions of peak (left) and trough (right) latencies in M1 (top row) and PMv (bottom

row). Although the median latency of troughs was
not different in M1 versus PMv (1.8 ms in M1 vs.
1.6 ms in PMv, p=0.49, Mann-Whitney U-Test),
the median latency of peaks was longer in M1 than
in PMv (4.3 ms in M1 vs. 3.4 ms in PMv, p=0.04,
Mann-Whitney U-Test). Inspection of the
distributions shows that this difference was not
due to fewer short-latency peaks produced in M1,
but rather due to more peaks at longer latencies (>
4ms) – an unexpected finding given that M1 is
physically closer than PMv to S1.

We quantified the duration of each peak or trough
by measuring its full-width at half maximum
(FWHM). The distributions of FWHMs (Fig. 6B)
show that virtually all peaks and many troughs in
both M1 and PMv were relatively short-lived, with
FWHMs < 1 ms. In contrast to peaks, though,
many troughs had FWHMs > 1 ms, making the
median duration of troughs longer than that of
peaks in both areas (1.0 ms vs. 0.4 ms in M1,
p=1.0e-06, Mann-Whitney U-Test; 0.7 ms vs 0.4
ms in PMv, p=2.8e-05, Mann-Whitney U-Test).
Between M1 and PMv, no differences were found
in FWHM for either peaks (p=0.34, Mann-
Whitney U-test) or troughs (p=0.13, Mann-
Whitney U-test).

We quantified the amplitudes of peaks/troughs as
the maximal/minimal value expressed as a percent
change from the baseline. Distributions are shown
in Fig. 6C. Peak amplitudes were as high as 869%
in one M1 neuron and 2650% in one PMv neuron
(corresponding to the probability of a spike in the
peak bin of 2.5e-03 and 7.5e-03, respectively).
The median percent increase for peaks was
significantly higher in PMv than in M1 (320% in
PMv vs 160% in M1, p=0.009, Mann-Whitney U-
Test).

Our ability to assess trough amplitude was
constrained by the spike probability floor of zero,
which was reached by most troughs, resulting in a
median percent decrease of 100% for both areas.
We recalculated the median percent decrease after

***Figure 6. Properties of excitatory and inhibitory effects
observed in M1 (blue) and PMv (green) neurons following S1-
ICMS pulses.*** *Separate histograms are shown for peaks (left
column) and troughs (right column). Dashed vertical lines mark
the median of each distribution. A) Latencies. The range of
latencies was similar for excitation and inhibition in both M1
and PMv, Short-latency troughs were more common than short-
latency peaks in both areas. Histogram bin size is 0.3 ms. B)
Full-Width Half-Maxima (FWHM). Most peaks and troughs in
both M1 and PMv were of relatively short duration, though
some troughs lasted longer than any peaks. Histogram bin size
is 0.1 ms. C) Amplitude (percent change from baseline). Black
dashed lines mark medians after excluding all values of 100%,
i.e. the floor for inhibition, 0 spikes. Histogram bin size is 60%
for peaks and 3% for troughs.*

476 excluding values of 100%. The median percent
477 decrease of the remaining troughs was still
478 substantial in both areas: 90% for M1 and 80% for
479 PMv.
480
The relatively short latency of some peaks raised
the possibility that S1-ICMS might have excited
some neurons in M1 or PMv antidromically.
Antidromic responses would be expected to
produce very narrow peaks with cumulative
probabilities approaching 1.0, reflecting one-to-
one following. However, antidromic responses
evoked by stimulation of thin, unmyelinated axon
collaterals may not follow stimulation at
frequencies ~100 Hz, with as few as 10% of
stimulus pulses evoking a somatic action potential,
i.e. a spike probability of 0.1 (Chomiak and Hu,
2007). In Fig. 7A we have plotted the FWHM of
each peak we evoked with S1-ICMS pulses
against its cumulative probability. Most peaks had
cumulative probabilities < 0.05. Of the 6 peaks
with cumulative probabilities > 0.05, an M1
neuron had the narrowest FHWM (arrow). The
PSTH with this peak is shown above in Fig. 5C.
Fig. 7B illustrates the jitter of the first action
potentials of this M1 unit (black) following 10
stimulus artifacts (red), indicating that excitation
of this M1 neuron was orthodromic and
transsynaptic. We thus infer that most, if not all, of
the peaks we found represented orthodromic,
transsynaptic excitation.
507
508 *Spatial convergence and divergence of S1-ICMS*
509 *effects on neurons across M1 and PMv*
510
511 Next, we examined the spatial convergence of
512 effects from multiple S1 arrays onto individual
513 units in M1 and PMv, as well as the spatial

Figure 7. Cumulative probability vs. FWHM and Jitter of S1-ICMS effects. A) Cumulative probability versus full-width half-maximum of each peak from M1 (blue) and PMv (green). Cumulative probabilities were calculated by summing each bin in the PSTH from the onset of the peak to the offset (described in the Methods). The black arrow indicates a point that represents a peak with both a relatively high cumulative probability and a narrow FWHM, as might be expected of an antidromic response. B) Overlay of 10 ICMS pulse artifacts (red traces) and the 10 following spikes (black traces) from the neuron-array pair denoted by the black arrow in A. Both waveforms were aligned on the timestamps of the ICMS artifacts from the designated artifact channel as described in Methods (0 ms, vertical dashed line). The jitter in the latencies of the spike waveforms relative to the artifact waveforms indicates orthodromic, transsynaptic excitation of the neuron. The PSTH of this neuron-array pair is shown in Figure 5C.

514 divergence of effects from stimulation of each S1 array to units across each cortical area. We present our
515 findings on convergence and divergence in both tabular (Fig. 8) and graphic (Fig. 9) form.
516
In both monkeys, ICMS effects from multiple different S1 arrays converged on individual neurons in both
M1 and PMv. Figure 8 (Convergence, left) gives the percentage (rounded to the nearest whole number) of
neurons recorded on each array in M1 or PMv that had at least one significant peak or trough following
ICMS delivered through different numbers of arrays in S1. Focusing on the right column (4) in monkey
L, for example, shows that ICMS effects converged from all four S1 arrays on some individual units
recorded from each of the four M1 arrays and from one of the four PMv arrays. In monkey X, ICMS

M1
PMv

Convergence

Monkey L — # S1 Arrays Producing Peak/Trough

# Units	Array	0	1	2	3	4
11	G	18%	36%	18%	0%	27%
5	I	0%	80%	0%	0%	20%
7	J	0%	14%	29%	29%	29%
15	K	20%	13%	7%	13%	47%
10	A	60%	10%	30%	0%	0%
7	B	57%	29%	14%	0%	0%
12	C	17%	42%	17%	25%	0%
13	D	31%	38%	15%	8%	8%

M1 vs. PMv: p=0.002, χ^2-test

Monkey X — # S1 Arrays Producing Peak/Trough

# Units	Array	0	1	2	3
5	F	0%	60%	20%	20%
11	G	18%	55%	27%	0%
2	H	0%	0%	0%	100%
18	I	33%	33%	33%	0%
12	C	8%	83%	0%	8%
18	D	17%	61%	22%	0%

M1 vs. PMv: p=0.14, χ^2-test

Divergence

Monkey L — S1-ICMS Arrays

# Units	Array	M	N	O	P
11	G	73%	27%	55%	27%
5	I	20%	80%	40%	20%
8	J	71%	71%	43%	86%
15	K	67%	60%	73%	53%
9	A	33%	11%	0%	11%
7	B	29%	29%	0%	0%
12	C	50%	67%	17%	17%
13	D	31%	38%	23%	31%

M1 vs. PMv: p=1.0e-7, χ^2-test

Monkey X — S1-ICMS Arrays

# Units	Array	K	L	O
5	F	60%	40%	60%
11	G	73%	18%	18%
2	H	100%	100%	100%
18	I	50%	50%	0%
12	C	92%	8%	8%
18	D	78%	17%	11%

M1 vs. PMv: p=0.25 χ^2-test

Figure 8. S1-ICMS effects by array. For convergence tables (left), each entry gives the percent of units on each M1 (blue) or PMv (green) array that had a significant peak and/or trough in the PSTHs triggered on artifacts of ICMS pulses delivered through different numbers of S1 arrays, from 0 to 4. For divergence tables (right), each entry gives the percent of units on each M1 or PMv array with a peak and/or trough in the PSTH triggered on artifacts from ICMS pulses delivered through the indicated S1 array. All values have been rounded to the nearest whole number and entries are color coded on a scale from white (0%) to dark blue/green (100%, M1/PMv). Results for monkey L are displayed above and results for monkey X below.

effects from converged from all three S1 arrays on individual units from two of the four M1 arrays and
from one of the two PMv arrays. In total, 13/38 (34%) M1 units and 1/42 (2%) PMv units received effects
converging from all four S1 arrays in monkey L, and 3/36 (8%) M1 and 1/30 (3%) PMv units received
effects converging from all three S1 arrays in monkey X. In each monkey, other units received effects
converging from fewer S1 arrays. ICMS effects from wide territories in S1 thus converged on many units
in both M1 and PMv.
529
Conversely, ICMS effects from each S1 array diverged to neurons recorded on multiple different arrays in
both M1 and PMv. Figure 8 (Divergence, right) gives the percentage of neurons recorded on each M1 or
PMv array that had at least one peak or trough following ICMS pulses delivered through each S1 array.
Looking down the columns shows that in monkey L, ICMS effects diverged to neurons on all 4 M1 arrays
from each of the 4 S1 arrays and diverged to all 4 PMv arrays from 2 of the 4 S1 arrays (M and N). In
monkey X, ICMS effects diverged to neurons on all 4 M1 arrays from 2 of the 3 S1 arrays (K and L) and
diverged to neurons from both PMv arrays from all three S1 arrays.
537
538

We then compared the
convergence and
divergence of S1-ICMS
effects between M1 and
PMv. By comparing counts
of units receiving direct
modulation from 0, 1, 2, 3,
or 4 S1 arrays, we found
that in both monkeys
individual M1 units were
directly modulated by
ICMS pulses delivered on
more S1 arrays than were
individual PMv units. This
trend was significant in
monkey L (p=0.002, χ^2
test), though not in monkey
X (p=0.14, χ^2 test). We also
found that in both monkeys

Figure 9. Graphic topography of direct modulation effects from each S1 array. Each square represents an FMA implanted in M1 (blue), PMv (green), or S1 (yellow). The thickness of each solid black line is proportional to the percentage of units recorded from each M1 or PMv array that showed a peak and/or trough following ICMS pulses delivered through the connected S1 array. The letters next to the arrays correspond to those in Figures 1 and 8. Brown dotted lines indicate sulci: Arcuate Sulcus (AS) and Central Sulcus (CS).

558 S1-ICMS delivered on individual arrays diverged to modulate units across M1 more commonly than
559 PMv, though again this trend was significant in monkey L (p=1.0e-7, χ^2 test), but not in monkey X
560 (p=0.25, χ^2 test). As illustrated by the thickness of the black lines in Figure 9, both convergence and
561 divergence of ICMS effects were more extensive from S1 to M1 than from S1 to PMv.
562
563 **Discussion**
564
565 We found that ICMS pulses delivered in S1 modulated the spiking activity of a majority of units recorded
566 in M1 and PMv. In both areas, we found inhibitory effects to be more common than excitatory effects. In
567 addition, the duration of inhibitory effects was often longer than that of excitatory effects. S1 outputs thus
568 may produce more inhibition than excitation in both M1 and PMv.
569
We also found extensive divergence of effects from S1 to both M1 and PMv. ICMS pulses delivered
simultaneously via a few electrodes on a single ~2x2 mm array in S1 produced effects in neurons
recorded on multiple other ~2x2 mm microelectrode arrays spread throughout the upper extremity
representation in M1 and over a considerable territory in PMv. Conversely, many single- or multi-units
anywhere in these regions of M1 and PMv received effects converging from all arrays spread across the
S1 upper extremity representation. Although macaque S1 has a detailed somatotopic organization (Pons et
al., 1985; Pons et al., 1987), its outputs to M1 and PMv appear to have less somatotopic segregation.
577
To our knowledge, few if any previous studies have examined the responses produced in multiple cortical
areas by ICMS in another cortical area. Our simultaneous recordings in M1 and PMv during S1-ICMS
permit certain comparisons of the effects evoked in these two regions. In particular, although the
underlying reasons are not apparent, M1 neurons showed a larger preponderance of inhibitory troughs
than PMv neurons, whereas PMv neurons showed on average larger excitatory peaks than M1 neurons
(Figure 6).
584
585 *Physiological and anatomical bases of distant effects produced by S1-ICMS*

586
Consistent with the natural sensory inputs known to reach M1 and PMv (see Introduction), we found that
more than 60% of single- and multi-units in both areas were directly modulated by ICMS pulses delivered
through at least one S1 microelectrode array. The pathways via which natural somatosensory input
reaches M1 and PMv are not entirely understood, though a substantial fraction may be processed through
S1. We were not able to obtain histological verification of the locations of our microelectrode tips, but
based on the distance of our S1 arrays from the central sulcus (< 3 mm) and their lengths (1.5 – 6.0 mm),
the tips of our S1 electrodes were probably located in areas 3b and 1, with some possibly in area 3a. Area
3a projects directly to M1 (area 4), but areas 3a, 3b and 1 all project to area 2, which in turn projects to
area 4 (Jones et al., 1978; Pons and Kaas, 1986; Dariansmith et al., 1993; Tokuno and Tanji, 1993;
Padberg et al., 2019). Area 2 also projects to area 5, which projects more heavily to area 4 (Strick and
Kim, 1978; Zarzecki et al., 1978). All of these parietal areas also project to the secondary somatosensory
area (S2) in the insular cortex, which in turn projects to area 4 (Friedman et al., 1986; Dariansmith et al.,
1993).
600
601 Whereas multiple transsynaptic routes are available from S1 to M1, routes from S1 to PMv are more
602 limited. Known somatosensory projections to PMv arise only from area 3a and from S2 (Godschalk et al.,
603 1984; Matelli et al., 1986; Kurata, 1991). The greater variety of transsynaptic routes from S1 to M1 may
604 account in part for the distribution of latencies for peaks in M1 being slightly wider, with a slightly longer
605 median latency compared to PMv (Fig. 6A).
606
The latencies and durations we observed were consistent with those observed in other studies following
natural and electrical stimulation of peripheral and cortical neurons. The latencies of M1 responses to
natural stimulation of the upper extremity or electrical stimulation of the median nerve range from 8 to 15
ms (Lemon, 1981). As would be expected, we found the latencies of effects produced in M1 by ICMS
pulses delivered in S1 (0.9 – 9.9 ms for peaks, 0.6 – 7.2 ms for troughs) to be shorter than latencies from
stimulation in the periphery. Although our S1-ICMS pulses were delivered at frequencies of ~100 Hz,
providing some potential to miss long latency effects, the latencies we observed in M1 neurons with S1-
ICMS were in the same range as those found in M1 neurons in a study using PMv-ICMS pulses delivered
at 10 Hz (1.8 –3.0 ms for peaks, 2–5 ms for troughs) (Kraskov et al., 2011). In humans, S1-ICMS at 100
Hz produced effects in M1 neurons at similar latencies (2–6 ms) (Shelchkova et al., 2022). Moreover, the
durations of the S1-ICMS effects we found in M1 (FWHMs 0.2 – 1.7 ms for peaks, 0.3 – 4.3 ms for
troughs), were similar to those obtained with PMv-ICMS (durations ~1ms for peaks, 5–7 ms for troughs)
(Kraskov et al., 2011).
620
A surprising observation was that the range of latencies we found for excitatory peaks in PMv neurons
(0.6 – 5.8 ms) was somewhat shorter than for those in M1 neurons (0.9 – 9.9 ms), even though the
physical distance from S1 to PMv is longer than from M1 to S1. Though the present latencies in PMv
neurons might seem unexpectedly short, they are similar to the range reported from area 5 to M1 (0.9 –
4.4 ms) (Zarzecki et al., 1978), which covers a similar physical distance. These observations may indicate
that conduction velocities are faster from S1 to PMv than to M1. In addition, the longer latencies of some
effects in M1 neurons from S1-ICMS may reflect the presence of more diverse transsynaptic routes from
S1 to M1 than to PMv.
629
We consider it unlikely that the peaks we identified represented antidromic conduction. To our
knowledge, there is no evidence that PMv projects directly to S1. For M1, antidromic responses would be
unlikely because axons from area 4, though projecting to areas 3a and 2, do not project to areas 3b or 1

(Godschalk et al., 1984; Matelli et al., 1986; Dariansmith et al., 1993), where most of our electrode tips
probably were situated. Furthermore, for the neuron-array pair having a peak with both a high cumulative
spike probability and a narrow FWHM (Figure 7), the first spike following the stimulus artifact showed
temporal jitter consistent with transsynaptic excitation. Therefore, the effects that S1-ICMS pulses evoked
in M1 and PMv neurons in this study were most likely orthodromic.
638

639   *Implications of Distant S1-ICMS Effects*
640

S1-ICMS produced more inhibition than excitation in both M1 and PMv. Somatosensory inputs typically
have been assessed by determining the receptive fields from which neurons in S1, M1, or PMv can be
excited. We therefore had not anticipated that S1-ICMS would produce both more frequent and longer
duration inhibitory effects than excitatory effects in both M1 and PMv neurons. In contrast to the present
results with S1-ICMS, PMv-ICMS has been found to produce more excitatory than inhibitory modulation
of M1 neurons (Kraskov et al., 2011). We suggest that, while PMv generally facilitates activity in M1, S1
may provide inhibition of undesired activity in both M1 and PMv.
648

We also observed extensive convergence and divergence between our arrays that spanned the upper
extremity representation in S1 and our arrays in both M1 and PMv. This might have been expected in
PMv given the large somatosensory receptive fields of PMv neurons (Rizzolatti et al., 1981; Graziano and
Gandhi, 2000). Less expected perhaps, was the similarly extensive convergence and divergence between
our S1 arrays and those in M1, which spanned the M1 upper extremity representation. A recent study in
humans likewise found that ICMS on many S1 electrodes evoked responses in M1 neurons, mostly
consistent with orthodromic, transsynaptic responses (Shelchkova et al., 2022).   Interestingly, focal
injection of tracers in area 3a, 3b, 1, or 2 results in terminal labeling spread mediolaterally over several
millimeters in the same cortical area (Jones et al., 1978). Likewise neurons at any focal location in the M1
upper extremity representation send horizontal collaterals throughout that representation (Huntley and
Jones, 1991). Horizontal collaterals in both S1 and M1 thus may provide an anatomical substrate for the
extensive convergence and divergence we observed physiologically between S1 and M1.
661

662   *Limitations of the Present Study*
663

The data analyzed here were collected in a study that deterimined whether ICMS delivered through
microelectrodes in different S1 locations produced experiences monkeys could distinguish. To maximize
the likelihood that the monkeys would distinguish S1-ICMS at different locations, we delivered current
pulses of 1-64 µA simultaneously on a set of 3-7 electrodes all on the same 16-electrode array. In
monkey X, we additionally varied the frequency of the pulse trains on different arrays from 75 to 225 Hz,
though in monkey Q all pulse trains were delivered at 100 Hz. The data analyzed here was collected after
we found S1-ICMS patterns that produced experiences each monkey could distinguish and report by
performing different movements. Subsequent testing showed that the monkeys could reliably detect S1-
ICMS on some single electrodes with currents as low as 10-15 µA, at frequencies as low as 30-50 Hz, or
in trains as short as 400 ms. Hence the present results likely approach the upper bounds of the strength
and spatial distribution of direct modulation in M1 and PM neurons produced by multichannel S1-ICMS.
Stimulating simultaneously on fewer electrodes and/or at lower currents could be expected to produce
fewer and weaker effects, though not necessarily with narrower spatial distribution.
677

Furthermore, we focused here on S1-ICMS because ICMS in S1 currently is being developed to provide
somatosensory feedback for bidirectional brain-computer interfaces. We plan subsequent studies of the
effects evoked by ICMS in other cortical areas as well. In addition, future studies will be needed to
address questions such as i) whether the amplitude of direct modulation produced by individual ICMS
pulses varies depending on task-related variation in neuron firing rates, ii) whether the use of ICMS
versus visual cues as instructions for different movements affects the average firing rates of M1 and PMv
neurons during task performance, and iii) how the instructional information provided by ICMS
propagates across cortical areas.
686
Conclusions

Whereas previous studies typically have delivered ICMS through a single microelectrode, the present
results are based on ICMS pulses delivered simultaneously through multiple microelectrodes on the same
~2x2 mm array. We chose to deliver ICMS in this manner to make it more likely that our monkeys would
perceive some component of the multi-electrode S1-ICMS and thereby be able to use it as an instruction
(Mazurek and Schieber, 2017a). Interestingly, recent studies of macaque and human S1 have suggested
that stimulating simultaneously through multiple microelectrodes may be a valuable approach for evoking
artificial sensations. Multi-channel S1-ICMS evokes tactile sensations more focal (Greenspon et al.,
2023b), at shorter latencies (Sombeck and Miller, 2019; Bjanes et al., 2022), and with more
distinguishable levels of perceived force (Greenspon et al., 2023a) than stimulating through a single
microelectrode. Multi-electrode ICMS therefore may be used increasingly in bidirectional brain-computer
interfaces (O'Doherty et al., 2011; Flesher et al., 2021). Our results show that such multi-electrode S1-
ICMS can directly modulate a majority M1 and PMv neurons, with extensive spatial divergence and
convergence across large cortical territories. In bidirectional BCIs that use S1-ICMS to deliver artificial
feedback, these effects will need to be taken into account when decoding the intended movement from
neural activity in M1 and PMv.
703

References

Andersen P, Hagan PJ, Phillips CG, Powell TP (1975) Mapping by microstimulation of overlapping projections from area 4 to motor units of the baboon's hand. Proc R Soc Lond B Biol Sci 188:31-36.

Asanuma H, Ward JE (1971) Patterns of contraction of distal forelimb muscles produced by intracortical stimulation in cats. Brain Res 27:97-109.

Averna A, Pasquale V, Murphy MD, Rogantin MP, Van Acker GM, Nudo RJ, Chiappalone M, Guggenmos DJ (2019) Differential Effects of Open- and Closed-Loop Intracortical Microstimulation on Firing Patterns of Neurons in Distant Cortical Areas. Cereb Cortex.

Bjanes DA, Bashford L, Pejsa K, Lee B, Liu CY, Andersen RA (2022) Multi-channel intra-cortical micro-stimulation yields quick reaction times and evokes natural somatosensations in a human participant. medRxiv.

Callier T, Brantly NW, Caravelli A, Bensmaia SJ (2020) The frequency of cortical microstimulation shapes artificial touch. Proc Natl Acad Sci U S A 117:1191-1200.

Chomiak T, Hu B (2007) Axonal and somatic filtering of antidromically evoked cortical excitation by simulated deep brain stimulation in rat brain. J Physiol 579:403-412.

Dariansmith C, Dariansmith I, Burman K, Ratcliffe N (1993) Ipsilateral Cortical Projections to Area-3a, Area-3b, and Area-4 in the Macaque Monkey. Journal of Comparative Neurology 335:200-213.

Fetz EE, Finocchio DV, Baker MA, Soso MJ (1980) Sensory and motor responses of precentral cortex cells during comparable passive and active joint movements. Journal of neurophysiology 43:1070-1089.

Flesher SN, Downey JE, Weiss JM, Hughes CL, Herrera AJ, Tyler-Kabara EC, Boninger ML, Collinger JL, Gaunt RA (2021) A brain-computer interface that evokes tactile sensations improves robotic arm control. Science (American Association for the Advancement of Science) 372:831-836.

Flesher SN, Collinger JL, Foldes ST, Weiss JM, Downey JE, Tyler-Kabara EC, Bensmaia SJ, Schwartz AB, Boninger ML, Gaunt RA (2016) Intracortical microstimulation of human somatosensory cortex. Sci Transl Med 8:361ra141.

Fraser GW, Schwartz AB (2012) Recording from the same neurons chronically in motor cortex. Journal of neurophysiology 107:1970-1978.

Friedman DP, Murray EA, Oneill JB, Mishkin M (1986) Cortical Connections of the Somatosensory Fields of the Lateral Sulcus of Macaques - Evidence for a Corticolimbic Pathway for Touch. Journal of Comparative Neurology 252:323-347.

Gentilucci M, Fogassi L, Luppino G, Matelli M, Camarda R, Rizzolatti G (1988) Functional organization of inferior area 6 in the macaque monkey. I. Somatotopy and the control of proximal movements. ExpBrain Res 71:475-490.

Godschalk M, Lemon RN, Kuypers HG, Ronday HK (1984) Cortical afferents and efferents of monkey postarcuate area: an anatomical and electrophysiological study. Exp Brain Res 56:410-424.

Graziano MS, Gandhi S (2000) Location of the polysensory zone in the precentral gyrus of anesthetized monkeys. ExpBrain Res 135:259-266.

Greenspon CM, Valle G, Hobbs TG, Verbaarschot C, Callier T, Okorokova EV, Shelchkova ND, Sobinov AR, Jordan PM, Gonzalez-Martinez J, Warnke PC, Miller LE, Boninger ML, Collinger JL, Gaunt RA, Downey JE, Hatsopoulos NG, Bensmaia SJ (2023a) Biomimetic multi-channel microstimulation of somatosensory cortex conveys high resolution force feedback for bionic hands. Neuroscience.

Greenspon CM et al. (2023b) Tesslation of artificial touch via microstimulation of human somatosensory cortex. bioRxiv.

Guggenmos DJ, Azin M, Barbay S, Mahnken JD, Dunham C, Mohseni P, Nudo RJ (2013) Restoration of function after brain damage using a neural prosthesis. Proceedings of the National Academy of Sciences - PNAS 110:21177-21182.

Histed MH, Bonin V, Reid RC (2009) Direct Activation of Sparse, Distributed Populations of Cortical Neurons by Electrical Microstimulation. Neuron (Cambridge, Mass) 63:508-522.

Huntley GW, Jones EG (1991) Relationship of intrinsic connections to forelimb movement representations in monkey motor cortex: a correlative anatomic and physiological study. Journal of Neurophysiology 66:390-413.

Hussin AT, Boychuk JA, Brown AR, Pittman QJ, Teskey GC (2015) Intracortical Microstimulation (ICMS) Activates Motor Cortex Layer 5 Pyramidal Neurons Mainly Transsynaptically. Brain stimulation 8:742-750.

Jankowska E, Padel Y, Tanaka R (1975) The mode of activation of pyramidal tract cells by intracortical stimuli. J Physiol 249:617-636.

Jones EG, Coulter JD, Hendry SH (1978) Intracortical connectivity of architectonic fields in the somatic sensory, motor and parietal cortex of monkeys. J Comp Neurol 181:291-347.

Kraskov A, Prabhu G, Quallo MM, Lemon RN, Brochier T (2011) Ventral premotor-motor cortex interactions in the macaque monkey during grasp: response of single neurons to intracortical microstimulation. J Neurosci 31:8812-8821.

Kumaravelu K, Sombeck J, Miller LE, Bensmaia SJ, Grill WM (2022) Stoney vs. Histed: Quantifying the spatial effects of intracortical microstimulation. Brain Stimul 15:141-151.

Kurata K (1991) Corticocortical inputs to the dorsal and ventral aspects of the premotor cortex of macaque monkeys. Neuroscience Research 12:263-280.

Lemon RN (1981) Functional properties of monkey motor cortex neurones receiving afferent input from the hand and fingers. Journal of Physiology - London 311:497-519.

Lemon RN, Porter R (1976) Afferent input to movement-related precentral neurones in conscious monkeys. ProcRSocLond 194:313-339.

Maier MA, Kirkwood PA, Brochier T, Lemon RN (2013) Responses of single corticospinal neurons to intracortical stimulation of primary motor and premotor cortex in the anesthetized macaque monkey. J Neurophysiol 109:2982-2998.

Matelli M, Camarda R, Glickstein M, Rizzolatti G (1986) Afferent and efferent projections of the inferior area 6 in the macaque monkey. JComp Neurol 251:281-298.

Mazurek KA, Schieber MH (2017a) Injecting Instructions into Premotor Cortex. Neuron 96:1282-1289 e1284.

Mazurek KA, Schieber MH (2017b) Injecting Instructions into Premotor Cortex. Neuron (Cambridge, Mass) 96:1282-1289.e1284.

Mazurek KA, Schieber MH (2019) How is electrical stimulation of the brain experienced, and how can we tell? Selected considerations on sensorimotor function and speech. Cogn Neuropsychol 36:103-116.

Mazurek KA, Schieber MH (2021) Injecting Information into the Mammalian Cortex: Progress, Challenges, and Promise. The Neuroscientist 27:129-142.

Mollazadeh M, Aggarwal V, Davidson AG, Law AJ, Thakor NV, Schieber MH (2011) Spatiotemporal variation of multiple neurophysiological signals in the primary motor cortex during dexterous reach-to-grasp movements. J Neurosci 31:15531-15543.

O'Doherty JE, Lebedev MA, Ifft PJ, Zhuang KZ, Shokur S, Bleuler H, Nicolelis MAL (2011) Active tactile exploration using a brain-machine-brain interface. Nature (London) 479:228-231.

Padberg J, Cooke DF, Cerkevich CM, Kaas JH, Krubitzer L (2019) Cortical connections of area 2 and posterior parietal area 5 in macaque monkeys. J Comp Neurol 527:718-737.

Pandarinath C, Bensmaia SJ (2022) The science and engineering behind sensitized brain-controlled bionic hands. Physiol Rev 102:551-604.

Pons TP, Kaas JH (1986) Corticocortical connections of area 2 of somatosensory cortex in macaque monkeys: a correlative anatomical and electrophysiological study. J Comp Neurol 248:313-335.

Pons TP, Garraghty PE, Cusick CG, Kaas JH (1985) The somatotopic organization of area 2 in macaque monkeys. Journal of Comparative Neurology 241:445-466.

Pons TP, Wall JT, Garraghty PE, Cusick CG, Kaas JH (1987) Consistent features of the representation of
the hand in area 3b of macaque monkeys. Somatosensory Research 4:309-331.
Rizzolatti G, Scandolara C, Matelli M, Gentilucci M (1981) Afferent properties of periarcuate neurons in
macaque monkeys. I. Somatosensory responses. Behavioural Brain Research 2:125-146.
Rizzolatti G, Camarda R, Fogassi L, Gentilucci M, Luppino G, Matelli M (1988) Functional organization
of inferior area 6 in the macaque monkey. II. Area F5 and the control of distal movements.
ExpBrain Res 71:491-507.
Romo R, Hernandez A, Zainos A, Salinas E (1998) Somatosensory discrimination based on cortical
microstimulation. Nature 392:387-390.
Rosen I, Asanuma H (1972) Peripheral afferent inputs to the forelimb area of the monkey motor cortex:
input-output relations. Experimental Brain Research 14:257-273.
Rouse AG, Schieber MH (2016) Spatiotemporal distribution of location and object effects in primary
motor cortex neurons during reach-to-grasp. The Journal of neuroscience 36:10640-10653.
Shelchkova ND et al. (2022) Microstimulation of human somatosensory cortex evokes taskdependent,
spatially patterned responses in motor cortex. bioRxiv.
Sombeck JT, Miller LE (2019) Short reaction times in response to multi-electrode intracortical
microstimulation may provide a basis for rapid movement-related feedback. J Neural Eng
17:016013.
Stoney SD, Jr., Thompson WD, Asanuma H (1968) Excitation of pyramidal tract cells by intracortical
microstimulation: effective extent of stimulating current. J Neurophysiol 31:659-669.
Strick PL, Kim CC (1978) Input to primate motor cortex from posterior parietal cortex (area 5). I.
Demonstration by retrograde transport. Brain Res 157:325-330.
Tokuno H, Tanji J (1993) Input organization of distal and proximal forelimb areas in the monkey primary
motor cortex: a retrograde double labeling study. Journal of Comparative Neurology 333:199-
209.
Wong YC, Kwan HC, MacKay WA, Murphy JT (1978) Spatial organization of precentral cortex in awake
primates. I. Somatosensory inputs. Journal of neurophysiology 41:1107-1119.
Zarzecki P, Strick PL, Asanuma H (1978) Input to primate motor cortex from posterior parietal cortex
(area 5). II. Identification by antidromic activation. Brain Res 157:331-335.