1   **Postzygotic barriers persist despite ongoing introgression in hybridizing *Mimulus* species**
2

3   Samuel J. Mantel[1*] and Andrea L. Sweigart[1]
4
5   [1]Department of Genetics, University of Georgia, Athens, GA, 30602, USA
6
7   [*]Corresponding author:
8   Samuel Mantel
9   120 E. Green St.
10   Athens, GA 30602
11   smantel1@uga.edu

Abstract

The evolution of postzygotic isolation is thought to be a key step in maintaining species

boundaries upon secondary contact. Yet the dynamics and persistence of hybrid

incompatibilities in naturally hybridizing species is not well understood. Here, we explore these

issues with genetic mapping in three independent populations of recombinant inbred lines

between naturally hybridizing monkeyflowers *Mimulus guttatus* and *M. nasutus* from the

sympatric Catherine Creek population. We discovered that the three *M. guttatus* founders differ

dramatically in admixture history, with nearly a quarter of one founder's genome introgressed

from *M. nasutus*. Comparative genetic mapping in the three RIL populations revealed three new

putative inversions, each one segregating among the *M. guttatus* founders, two due to

admixture. We also found strong, genome-wide transmission ratio distortion in all RILs, but

patterns were highly variable among the three populations. At least some of this distortion

appears to be explained by epistatic selection favoring parental genotypes, but tests of inter-

chromosomal linkage disequilibrium also revealed multiple candidate Dobzhansky-Muller

incompatibilities. We also mapped several genetic loci for hybrid pollen viability, including two

interacting pairs that coincide with peaks of distortion. Remarkably, even with this limited

sample of three *M. guttatus* lines, we discovered abundant segregating variation for hybrid

incompatibilities with *M. nasutus,* suggesting this population harbors diverse contributors to

postzygotic isolation. Moreover, even with substantial admixture, hybrid incompatibilities

between Mimulus species persist, suggesting postzygotic isolation might be a potent force in

maintaining species barriers in this system.

Introduction

Despite often being conceptualized as a single event, it has long been understood that speciation is a complex and dynamic process (Darwin, 1859; Dobzhansky, 1937). As new species form (generally in allopatry), the cumulative impacts of adaptive divergence, genetic drift, and the sorting of ancestral variation can lead to the evolution of reproductive isolation, which limits gene flow between these species when they come into secondary contact (Dobzhansky, 1937; Mayr, 1942; Coyne & Orr, 2004). Nevertheless, gene flow between recognized species – at rates high enough to have substantial genomic consequences – is not uncommon, even when strong reproductive isolating barriers exist (Nolte *et al.*, 2009; Teeter *et al.*, 2010; Ellegren *et al.*, 2012; Renaut *et al.*, 2013; Larson *et al.*, 2014; Delmore *et al.*, 2015; Caeiro-Dias *et al.*, 2023). At the extremes, hybridization can break down established barriers causing previously isolated lineages to fuse (Taylor *et al.*, 2006) or to strengthen reproductive isolation through reinforcement (Servedio & Kirkpatrick, 1997; Jaenike et al., 2006; Hopkins & Rausher, 2012). On the other hand, theory suggests that hybridization might sometimes reflect partial reproductive isolation as a stable optimum(Servedio & Hermisson, 2020). Studies of naturally hybridizing species can provide a window into the causes of these alternative evolutionary outcomes, allowing us to identify and measure the impact of the genetic loci involved in reproductive isolation.

One powerful approach to this problem is experimental hybridization of incompletely isolated species that occur together in secondary sympatry. In particular, the use of late-generation hybrids such as recombinant inbred lines (RILs) can enable replication of wild, admixed genotypes and facilitate genetic mapping of barrier loci under controlled

environmental conditions. This approach should be especially fruitful for identifying loci

involved in postzygotic isolation because strongly incompatible allele combinations are likely to

be purged during RIL formation. Indeed, selection against hybrid incompatibilities can be a

major source of transmission ratio distortion (TRD) in RIL populations (Colomé-Tatché &

Johannes, 2016). Although there are often additional contributors to TRD in experimental

hybrid populations (reviewed in Fishman & McIntosh, 2019), hybrid incompatibilities are

expected to leave a particular signature of distortion, generating both local peaks and linkage

disequilibrium between unlinked loci caused by underrepresented or absent multi-locus

genotypes. This signature might be particularly apparent in RILs, which have the advantage of

multiple generations of recombination, increasing power to detect and localize incompatibilities

by amplifying their effects (and revealing recessive phenotypes) through inbreeding (Moyle &

Nakazato, 2008).

Despite the clear value of experimental hybridization and RILs for illuminating the

genetic causes of reproductive isolation, inferences to natural populations can be challenging

because such studies are often based on initial crosses between only two individuals or inbred

lines. This limitation might be serious for investigations of postzygotic isolation between species

in the early stages of divergence, as hybrid incompatibilities are often polymorphic (Sweigart *et*

al., 2007; Good *et al.*, 2008; Cutter, 2012), particularly in populations with ongoing hybridization

(Zuellig & Sweigart, 2018). Variation in chromosomal structure is also a common feature of

natural populations (Hoffmann & Rieseberg, 2008), and polymorphic inversions are often

associated with adaptive differences (Feder *et al.*, 2003; Lowry & Willis, 2010; Jones *et al.*, 2012)

and reproductive isolation (Noor *et al.*, 2001; Sotola *et al.*, 2023). One way to capture a more

representative set of natural variation is to generate a multi-parent mapping population, for

instance by performing multiple biparental crosses to a common founder (Scott *et al.*, 2020).

Although this strategy is now routinely used in plant breeding programs (Scott *et al.*, 2020) and

to dissect the genetics of complex traits (Aylor *et al.*, 2011; Long *et al.*, 2014) it has rarely been

applied to studies of speciation.

In this study we investigate patterns and mechanisms of reproductive isolation in three

independent RIL populations constructed from crosses between *Mimulus guttatus* and *M.*

nasutus (also referred to as *Erythranthe guttata* and *E. nasuta*: see Lowry *et al.*, 2019 and

Nesom *et al.*, 2019). These two species are closely related (~200 KYA diverged: Brandvain *et al.*,

2014), and broadly allopatric across western North America, but they also occur in secondary

sympatry at many sites in their shared range. One such sympatric site occurs at Catherine Creek

(CAC) in Washington state, a system of ephemeral seeps and streams that flow into the

Columbia River. The two *Mimulus* species have diverged in mating system (*M. guttatus* is large

flowered and predominantly outcrossing, whereas *M. nasutus* is a small flowered and primarily

selfing), which acts as a strong premating barrier where the two species co-occur (Martin &

Willis, 2007). At Catherine Creek, divergent flowering phenology (Kenney & Sweigart, 2016) and

microhabitat adaptation (Mantel & Sweigart, 2019) are also important contributors to

premating isolation. Despite these substantial barriers to interspecific mating, studies of

population genomic variation at Catherine Creek have revealed clear signatures of historical and

contemporary hybridization (Kenney & Sweigart, 2016), resulting in largely unidirectional

introgression from *M. nasutus* into *M. guttatus* (Sweigart & Willis, 2003; Brandvain *et al.*, 2014;

Kenney & Sweigart, 2016). Even in the absence of interspecific gene flow, *M. guttatus*

populations are exceptionally genetically diverse (Flagel *et al.*, 2014; Puzey *et al.*, 2017) and

maintain substantial variation for complex phenotypes (Troth et al., 2018).

Here, we attempt to capture and leverage this diversity at Catherine Creek to

understand the patterns and mechanisms of hybrid breakdown in naturally hybridizing species.

We generate three independent RIL populations by crossing distinct CAC *M. guttatus*

progenitors to a common *M. nasutus* parent. Using this powerful crossing design, we discover

that differences in admixture have cascading effects on structural variation and reproductive

barriers. Moreover, even in this sample of three *M. guttatus* lines, we find segregating variation

for hybrid incompatibilities with *M. nasutus*, suggesting this population harbors diverse

contributors to postzygotic isolation. Finally, we discover that even with substantial admixture,

hybrid incompatibilities between *Mimulus* species persist, suggesting postzygotic isolation might

be a potent force in maintaining species barriers in the face of ongoing hybridization.

Materials and Methods

Plant Material and Crossing Design

The plants used in this study were derived from a collection of inbred lines (>10

generations inbred, single seed decent) produced from wild-collected seed from Catherine

Creek (CAC). The CAC site consists of a system of gently sloping streams and seeps carrying

snowmelt and spring rain out of the Cascade Mountains southward through a rocky meadow

into the Columbia River Gorge. While *M. nasutus* and *M. guttatus* grow sympatrically at CAC,

often within centimeters of each other, they occupy somewhat distinct microhabitats and

flower asynchronously (Kenney & Sweigart, 2016). Still, reproductive isolation is incomplete

between the species and ongoing introgression is detectable, mostly from *M. nasutus* into *M.*

guttatus (Brandvain *et al.*, 2014; Kenney & Sweigart, 2016).

We produced three parallel RIL populations from crosses between CAC *M. guttatus* and

M. nasutus. We selected three previously studied CAC *M. guttatus* inbred lines (CAC110,

CAC162, and CAC415; TableS1; Mantel & Sweigart, 2019) with stereotypic *M. guttatus*

morphological and phenological characteristics and which produced highly fertile F1s with *M.*

nasutus. One *M. nasutus* inbred line (CAC9) served as a common maternal parent, which we

emasculated before pollen development, and crossed by hand-pollination with pollen from

each of the *M guttatus* inbred lines, producing three F1 hybrids. For each of these F1 hybrids,

we self-fertilized a single plant to generate three corresponding F2 populations (*N* = 500 each).

We planted seeds onto moist Fafard 3-B potting mix (Sun Gro Horticulture, Agawam, MA) in 2.5"

pots, and cold stratified them for seven days at 4°C before moving them to a Conviron growth

chamber (16-hr days; 23°C days/16°C nights) for germination. We transplanted seedlings into

individual cells of 96-cell plug trays filled with potting mix and moved them to the University of

Georgia greenhouses under 16-hr supplemental light, 23°C days/16°C nights. Under these

germination and growing conditions, we performed four additional rounds of self-fertilization

using single-seed-descent to produce three populations of F6 recombinant inbred lines (RILs;

Figure S1). We generated a total of 368 RILs (151 CAC110 RILs, 128 CAC162 RILs, and 89 CAC415

RILs), much lower than the initial 1500 due to line extinction during the inbreeding process.

Library Preparation, Sequencing, and Processing

We grew one individual of each of the 368 RILs (F6 generation or later) for tissue

collection and DNA extraction in the greenhouse at the University of Georgia under the

conditions described above. We collected bud and meristem tissue from each individual into 96-

well plates on dry ice, stored them at -80°C, and extracted genomic DNA using a modified CTAB

method in a 96-well format (dx.doi.org/10.17504/protocols.io.bgv6jw9e). We then constructed

ddRADseq libraries using a modified version of RAD Capture (Ali *et al.*, 2016), "BestRAD"

(dx.doi.org/10.17504/protocols.io.6awhafe). Briefly, we digested 50-100 ng of genomic DNA

with BfaI and PstI, and used the 'NEBNext Ultra II' library preparation kits for Illumina (New

England BioLabs, Ipswich, MA) to ligate 48 unique barcoded i5 adaptors to each sample, isolated

tagged fragments, and pooled samples into 8 pools of 48 or fewer samples each. We attached a

unique 'NEBNext Adaptor for Illumina' i7 adaptor to each pool of samples, and enriched the

samples with PCR. We size-selected the libraries to 300-900bp using BluePippin 2% agarose

cassettes (Sage Science, Beverly, MA) and pooled the libraries in equimolar concentrations

following quantification via qPCR. We prepared libraries for, and sequenced the same RILs

twice, each on two partial lanes of Illumina HiSeq4000 for paired end 150-bp reads at the

Genome Sequencing Facility at Duke University. We also extracted genomic DNA from bud and

leaf tissue of the three parental CAC *M. guttatus* lines using a modified CTAB method as above.

Whole genome sequence (WGS) libraries were prepared and sequenced on a partial lane of

Illumina HiSeq4000 for paired end 150-bp reads at the Genome Sequencing Facility at Duke

University.

After trimming Illumina adaptors from reads using Trimmomatic (v0.36; Bolger *et al.*,

2014), we aligned WGS reads to the *Mimulus guttatus* IM62v3 refence genome (phytozome.org)

using BWA-MEM (v0.7.17; Li & Durbin, 2009; Li, 2013). We then removed reads with alignment

qualities <Q29 with SAMtools view (v1.6; Li *et al.*, 2009) and added read groups, checked mate-

pair information, and removed PCR duplicates with Picard (v2.16.0;

broadinstitute.github.io/picard/). We then removed paired reads which had been mapped to

different contigs using SAMtools view (v0.7.17; Li *et al.*, 2009). Finally, using GATK

HaplotypeCaller (v3.8.1; Van der Auwera & O'Connor, 2020) in GVCF mode we called SNPs, and

filtered SNPs to remove heterozygous calls and retain sites with a minimum depth of 3x, and a

maximum depth of 2 standard deviations above the mean (calculated with Qualimap2 (v2.2.1;

Okonechnikov *et al.*, 2016)) for each line. To allow for genotyping of *M. guttatus* and *M. nasutus*

SNPs across the genome, we identified a full list of diagnostic SNPs (those homozygous for

different alleles in each of the parental pairs, CAC9 & CAC110, CAC9 & CAC162, and CAC9 &

CAC415). We joint genotyped each pair of parental lines using GATK GenotypeGVCFs (v3.8.1;

Van der Auwera & O'Connor, 2020). We quality filtered the called SNPs (QD < 2.0, FS > 60.0, MQ

< 40.0, MQRankSum < -12.5, ReadPosRankSum < -8.0) and retained only biallelic SNPs using

VariantFiltration and SelectVariants. From these three VCFs we extracted a list of high quality

diagnostic SNPs with VCFtools (v0.1.16; Danecek *et al.*, 2011).

We demultiplexed the ddRADseq data by i7 index following sequencing and removed

PCR duplicates from each library, using the unique molecular barcode of the i5 adaptor with a

custom Python script (dx.doi.org/10.17504/protocols.io.bjnbkman). We demultiplexed each

library to the individual sample level using Flip2BeRAD (github.com/tylerhether/Flip2BeRAD)

and Stacks process_radtags (v2.55; Catchen *et al.*, 2011, 2013). We removed Illumina adaptors

and aligned reads to the *Mimulus guttatus* IM62 v3 reference assembly (IM: Iron Mountain, an

allopatric *M. guttatus* population in Oregon, ~200km south of CAC). 6 samples were removed

from further analysis (4 from the CAC162 RILs, and 2 from the CAC415 RILs) due to low

similarity between duplicate BAMs produced from the two independent ddRAD library

preparations. We then merged the remaining duplicate BAMs with SAMtools merge (v0.7.17; Li

et al., 2009), leaving a total of 362 RIL individuals (151 CAC110 RILs, 124 CAC162 RILs, and 87

CAC415 RILs). We then processed reads with Picard and SAMtools as above (with the exception

of MarkDuplicates, as PCR duplicates were already removed). We called SNPs, joint genotyped

each RIL population with the parental lines from which it was derived, and filtered called SNPs

as described above. We then removed sites with only parental coverage with VariantFiltration

and SelectVariants in GATK (v3.8.1; Van der Auwera & O'Connor, 2020). Finally, using the list of

diagnostic SNPs between each pair of parental lines, we removed non-diagnostic sites with

VCFtools (v0.1.16; Danecek *et al.*, 2011).

To assess levels of residual heterozygosity in the RIL populations and ensure its removal

would not skew our RIL genotypes, we filtered the VCFs of diagnostic SNPs to a minimum depth

of 10x (GATK VariantFiltration and SelectVariants as above). We then estimated genome wide

heterozygosity at the SNP level. Apart from a few very small regions of intermediate

heterozygote frequency which may warrant future investigation, heterozygosity was low and

uniform across the genome: CAC110 RILs: 1.95%, N=13,975; CAC162 RILs: 2.34%, N=33,340;

CAC415 RILs: 2.79%, N=23,178 and consistent with the Mendelian expectation (~1.5-3%, in

F6/F7 individuals).

Assessment of Introgression in Parental Lines

As previous studies have shown that admixture between *M. guttatus* and *M. nasutus* is

common at CAC (Brandvain *et al.*, 2014; Kenney & Sweigart, 2016) we first checked the selected

CAC *M. guttatus* parental lines for evidence of introgression from *M. nasutus*. To coarsely

identify putative regions of introgression between CAC *M. nasutus* and the CAC *M. guttatus*

parental lines, we used a custom perl script to calculate Dxy between CAC9 and the three

parental CAC *M. guttatus* lines in non-overlapping 50kb windows which contained 5000 or more

called sites.

To more concretely asses evidence of *M. nasutus* introgression in the parental CAC *M.*

guttatus lines we used the newly generated WGS data, plus WGS data from 12 other inbred

Mimulus accessions downloaded from the SRA (https://www.ncbi.nlm.nih.gov/sra): the parental

M. nasutus line CAC9, 3 non-CAC *M. nasutus* lines, 5 additional non-parental *M. guttatus* lines

from CAC, 2 non-CAC *M. guttatus* lines*,* and 1 *M. dentilobus* line as an outgroup) (Table S1). We

processed reads from additional (non-parental) lines as above and joint genotyped all lines

together using GATK GenotypeGVCFs (v3.8.1; Van der Auwera & O'Connor, 2020). We used

GATK to filter called SNPs based on quality as above, and retained only biallelic SNPs with

VariantFiltration and SelectVariants, producing a high-quality set of called SNPs.

We identified putative regions of introgression using a Hidden Markov Model (HMM,

modified from (Durbin *et al.*, 1998) developed in (Brandvain *et al.*, 2014)). As an input for the

model, we identified fourfold degenerate sites in the IM62 v3 reference assembly using the

associated gene annotations extracted with a Perl script developed by Tim Sackton

(https://github.com/tsackton/linked-selection/tree/master/misc_scripts), and identified non-

degenerate sites in the same way using modified versions of the script. With these lists and the

VCF of high-quality filtered SNPs in parental and non-parental *Mimulus* lines described above,

we used custom Perl scripts to count the total number of sites, as well as the number of non-

degenerate and fourfold degenerate sites, with genotype calls in each 1kb window across the

genome of each sequenced line. We then counted the number of those sites which differed in

their SNP call between each pairwise comparison of lines. With these counts and the HMM, we

calculated posterior probabilities of *M. nasutus* and *M. guttatus* ancestry across the genomes of

the parental CAC *M. guttatus* lines. We considered 1kb windows with a calculated posterior

probability of *M. nasutus* ancestry >0.95 to be windows showing evidence of introgression.

RIL Genotyping and Linkage Mapping

To genotype the RILs we first used custom python scripts to count the number of reads

supporting reference and alternate SNP calls at each site, in each RIL individual. We then

assigned parental ancestry to these calls, with a custom R script leveraging the parental SNP

information to convert ref/alt calls to *M. guttatus* or *M. nasutus* allele calls. We then used a

custom python script to average SNP calls across non-overlapping 50kb windows, producing a

consensus call leveraging the read depth at each called SNP. We called windows with >75% *M.*

guttatus calls as homozygous *M. guttatus*, and vice versa, and called windows with between 25

and 75% of either genotype heterozygous. This produced windowed genotype calls for many

50kb windows across the genome, but left gaps where windows containing no sequencing reads

could not be genotyped. To fill these gaps we used Python scripts from the processing steps of

the program GOOGA (Flagel *et al.*, 2019) to calculate genotype probabilities for each window in

each RIL individual. We first used the windowed genotypes to estimate genotype error rates and

recombination rates for each RIL sample, using the advanced generation RIL expectation of 1:1

homozygous genotype frequencies. During this process we dropped 39 samples from further

analysis due to a low number of called genotypes (from low original coverage) leading to high

($e1$ or $e2 > 0.2$) estimated error rates (8 from the CAC110 RILs, 26 from the CAC162 RILs, and 5

from the CAC415 RILs). We used an HMM (Flagel *et al.*, 2019) leveraging this error and

recombination information to estimate genotype probabilities at each 50kb window across the

genome in each RIL individual. We converted genotype probabilities to hard genotype calls by

excluding calls with an estimated genotype probability of <0.75. Of the original 368 RILs, we

generated genotype data for 323 of them (143 of the original 151 from the CAC110 RILs, 98/128

from the CAC162 RILs, and 82/89 from the CAC415 RILs).

We used LepMAP3 (Rastas, 2017) to construct a high density linkage map for each of

the three RIL families using the genotype matrices produced above. We used the affx2post.awk

script to calculate posterior genotype probabilities, and then ran ParentCall2 with default

parameters. We ran SeparateChromosomes2 to group markers from the same chromosome,

followed by OrderMarkers2 (grandparentPhase=1, useKosambi=1, sexAveraged=1, 10

iterations/per chromosome) to create a linkage map for each chromosome of the reference

genome (details in Table S2). Of the 10 iterations of OrderMarkers2, (or using the physical order

of markers in the reference genome if marker order was poorly resolved), we selected the order

with the highest likelihood for each chromosome (Table S2). We then manually removed

standalone markers whose physical position differed greatly with their calculated genetic

position on the chromosome. This procedure produced three linkage maps containing 3,828,

4,389 and 4,379 markers respectively. Of these markers, we retained 794, 606, and 690 which

mapped to recombination-informative sites.

Assessment of Transmission Ratio Distortion and Linkage Disequilibrium

276   Following linkage mapping, we calculated genotype frequencies at each marker in each

map and identified regions of transmission ratio distortion (TRD) by testing for deviations from

Mendelian 1:1 expectation with χ^2 tests. We used both an uncorrected $\alpha = 0.01$ and a more

stringent Bonferonni-corrected α (unique α for each chromosome in each RIL population

depending on the number of markers tested). Because the linkage maps for the three RIL

populations varied both in size and marker order, we investigated TRD colocalization among the

three papulations on a marker-by-marker basis. To investigate whether peaks of TRD were

associated with major two-locus hybrid incompatibilities we used the package pegas (Paradis,

2010) to characterize pairwise linkage disequilibrium (LD) between all markers with unique map

positions in the three RIL populations. To minimize false positives driven by the large number of

pairwise comparisons, we first identified putative interacting loci associated with inter-

chromosomal LD based on r values greater than the 95% percentile (CAC110 RILs: r > 0.37,

CAC162 RILs: r > 0.41, CAC415 RILs: r > 0.39). At these interacting loci we then performed

likelihood ratio tests comparing observed genotype ratios to expected ratios (calculated by cross

tabulation of observed single locus allele frequencies). We describe only the most significant

epistatic interactions where p<0.0001.

Assessment of Hybrid Male and Female Sterility

294   To measure male fertility, we collected pollen from three flowers (second flower pair or

later) of each RIL individual. We suspended the pollen of each flower separately in lactophenol-

aniline blue stain (Kearns & Inouye, 1993) and inspected the pollen grains using a compound

microscope. For each flower, we estimated pollen fertility by calculating the proportion of darkly

stained (viable) pollen grains from a haphazardly selected group of >100 pollen grains within a

field of view in of the microscope. When fertility estimates from flowers of the same plant

differed by more than 0.2, pollen from up to 3 more flowers was collected and counted. We

then calculated the average pollen viability from the collected flowers for each RIL individual.

Because the distribution of pollen viability in each RIL population was left-skewed, we logit

transformed the data. We also collected pollen from and estimated viability in each parental

line, and the three F1s between them.

305   To estimate hybrid female sterility, we intercrossed random pairs of RILs in at least

duplicate, and collected all recombinant inbred intercrossed (RIX) seeds produced. We counted

the number of seeds produced per fruit for each cross made and ran a linear mixed model

(LMM, using the lmer command in the 'lme4' package). In addition to maternal parent genotype

(fixed effect), the model included average viable pollen proportion of the paternal parent (as

calculated above, fixed effect), a unique RIX identifier nested within RIX cross type (CAC110 RIL x

CAC162 RIL, CAC162 RIL x CAC415 RIL, etc.; random effect) and the year that the cross was

made (crosses were made during two separate grow-outs in separate years; random effect). We

then calculated least square means (LSMs) of female viability (using seeds per fruit as a proxy)

for each RIL genotype using the emmeans function in the 'emmeans' package. For each of the

three RIL families this produced distributions with a long right tail, so we used a square root

transformation to more closely match a normal distribution.

QTL Mapping

Given the small sample sizes of each RIL population, we took a multistep approach to

genetically mapping hybrid male and female sterility. For each trait in each RIL population, we

first used r/QTL (Broman & Sen, 2009; Arends *et al.*, 2010) to perform 10 iterations of composite

interval mapping (CIM; Zeng, 1993, 1994; Jansen & Stam, 1994). Because CIM requires no

missing data, missing genotypes are imputed before the scan is run, leading to potential

differences between iterations. We set lenient LOD thresholds (α = 0.15) for QTL significance, by

performing 5,000 permutations, and calculated 1.5-LOD intervals ('lodint' command) for each

significant peak position in any of the 10 iterations. We then performed single-QTL scans using

each QTL identified with CIM as an interactive covariate ('intcovar' option) in separate runs. We

again we set LOD thresholds (α = 0.15, 10,000 permutations) for interacting QTL significance,

and calculated 1.5-LOD intervals for each significant peak position. With these identified peaks,

we then fit a multiple-QTL model ('fitqtl' command; HK method) and directly estimated additive

QTL effects (at each peak position) and identified epistatic effects. We then removed any

nonsignificant (p > 0.05) effects, and re-fit the model.

Results

Admixture in M. guttatus parental lines

All three *M. guttatus* parents show clear evidence of admixture from *M. nasutus*, but the

genomic location and proportion of introgression vary considerably across lines (Figure 1, Figure

S2). Of the three parents, CAC110 is the most admixed with ~24% of its genome showing very

low genetic divergence (*Dxy*) and a high posterior probability of recent coalescence with *M.*

nasutus via probabilistic inference (HMM, see Methods; Brandvain *et al.*, 2014). At least some

of this introgression in CAC110 appears to be recent, with large portions of entire chromosome

arms showing evidence of introgression (Figure S2). Although the largest inferred contiguous

blocks in CAC110 are only ~1.1Mb (Figure S3B), the HMM may sometimes break up true

contiguous blocks of introgression (or incorrectly identify short regions with low *Dxy* as

admixture blocks – see Brandvain et al. 2014). Given the improbability of inheriting a large

number of small introgressions immediately adjacent to one another, we think it likely that

several regions in CAC110 (on Chromosomes 1, 4, 8, 11, and 13 – Figure 1 and Figure S2)

represent true contiguous blocks of admixture. In both CAC162 and CAC415, the overall

proportion of admixture is lower (~9% in each) and blocks are generally smaller and more

evenly dispersed throughout the genome (Figure S2; Figure S3B), consistent with introgression

events that occurred in the more distant past. Even in CAC110, however, there are many very

small regions of inferred introgression (Figure S3), suggesting *M. guttatus* genomes at Catherine

Creek (CAC) are shaped by a long history of introgression from *M. nasutus*.

Results

Admixture in M. guttatus parental lines

All three *M. guttatus* parents show clear evidence of admixture from *M. nasutus*, but the

genomic location and proportion of introgression vary considerably across lines (Figure 1, Figure

S2). Of the three parents, CAC110 is the most admixed with ~24% of its genome showing very

low genetic divergence (*Dxy*) and a high posterior probability of recent coalescence with *M.*

nasutus via probabilistic inference (HMM, see Methods; Brandvain *et al.*, 2014). At least some

of this introgression in CAC110 appears to be recent, with large portions of entire chromosome

arms showing evidence of introgression (Figure S2). Although the largest inferred contiguous

blocks in CAC110 are only ~1.1Mb (Figure S3B), the HMM may sometimes break up true

contiguous blocks of introgression (or incorrectly identify short regions with low *Dxy* as

admixture blocks – see Brandvain et al. 2014). Given the improbability of inheriting a large

number of small introgressions immediately adjacent to one another, we think it likely that

several regions in CAC110 (on Chromosomes 1, 4, 8, 11, and 13 – Figure 1 and Figure S2)

represent true contiguous blocks of admixture. In both CAC162 and CAC415, the overall

proportion of admixture is lower (~9% in each) and blocks are generally smaller and more

evenly dispersed throughout the genome (Figure S2; Figure S3B), consistent with introgression

events that occurred in the more distant past. Even in CAC110, however, there are many very

small regions of inferred introgression (Figure S3), suggesting *M. guttatus* genomes at Catherine

Creek (CAC) are shaped by a long history of introgression from *M. nasutus*.

Comparative linkage mapping

Across most of the genome, the three RIL linkage maps are colinear with similar lengths

(Table 1), and marker order generally follows the physical order of the *M. guttatus* IM62 v3

reference assembly (Figure 2, Figure S4). However, we identified three clear exceptions to this

pattern, where the maps differ due to parental variation in putative inversions, characterized by

a large number of physically dispersed markers mapping to the same genetic position. Two of

these putative inversions are on LG1 (*inv1.1* and *inv1.2*) and are potentially unique to Catherine

Creek (or locally distributed), having not been reported in other crosses with these *Mimulus*

species (Fishman *et al.*, 2014; Flagel *et al.*, 2019). In both cases, the CAC9 *M. nasutus* parent

carries the ancestral orientation and the *M. guttatus* parents segregate for the derived variant.

At *inv1.1*, recombination is suppressed in the CAC110 and CAC415 RILs, but not in the CAC162

RIL, indicating collinearity between the CAC9 *M. nasutus* and CAC162 *M. guttatus* parents. At

inv1.2, on the other hand, the genetic maps suggest CAC9 is collinear only with CAC110.

Remarkably, this *M. guttatus* parent also carries a large block of introgression spanning this

entire region (Figure 1; Figure S2), suggesting that gene flow from *M. nasutus* has re-introduced

the ancestral variant into CAC110. We discovered a similar situation on LG13, where

introgression from *M. nasutus* accounts for collinearity between CAC9 and CAC110 at *inv13*

(recombination rates in the CAC110 RIL are much higher in this region than in the CAC162 and

CAC415 RILs; Figure S4). Additionally, *inv13* might have arisen in *M. nasutus*, as it is also appears

to be present in the reference genome assembly of *M. nasutus* SF5 (an accession derived from

Shear's Falls, an allopatric *M. nasutus* population ~70 km from CAC), but not in other recently

assembled *M. guttatus* genomes (phytozome.org). Taken together, these mapping results point

to introgression from *M. nasutus* as a major source of large-scale structural variation at

Catherine Creek, responsible both for restoring ancestral variants and for introducing new

derived inversions into *M. guttatus*.

In addition to the effects of structural variation, we observed some instances of map

length variation that appear to be associated only with parental differences in introgression. In

the CAC110 RILs, introgressed regions on LGs 1, 11, 12, and 13 have elevated genetic distances

relative the same (non-introgressed) regions in the other two RIL populations (Table 1, Figure 2,

Figure S4). Rather than reflecting a biological difference in recombination frequency between

introgressed and non-introgressed regions, we attribute this effect to technical differences

arising from the much lower marker densities in introgressed regions, which are highly similar

to the CAC9 parent (Table 1, Figure 2, Figure S4).

Genome-wide patterns of transmission ratio distortion (TRD)

With the exception of LG9, we observed strong TRD in at least one RIL population on all

linkage groups (Figure 3, Figure S6, Table S3). Globally, patterns of TRD are highly variable across

RIL populations. In the CAC110 and CAC415 RILs, the majority of distorted markers show an

excess of *M. guttatus* genotypes (59% in CAC110 and 80% in CAC162), whereas in the CAC162

RILs, there is a strong deficit of *M. guttatus* across the genome (80% of distorted markers show

an excess of *M. nasutus*). In a few cases, the same region is even distorted in opposite directions

(e.g., on the end of LG4: toward *M. guttatus* in CAC110 RILs and *M. nasutus* in CAC162 RILs;

Figure 3, Figure S6). Despite this general variability, there are a few instances of shared

distortion in multiple RIL populations, potentially implying the same causal mechanisms. Most

notably, the proximal end of LG14 shows an excess of *M. guttatus* alleles in all three

populations. There is also an excess of *M. guttatus* alleles on LGs 1 and 4 and an excess of *M.*

nasutus alleles on LGs 8, 12, and 13 in two of the three RIL populations (CAC110 and CAC162).

These regions of shared distortion are largely confined to segments of the genome that are non-

introgressed in parental lines. In addition to local peaks of TRD among homozygous marker

genotypes, we also investigated levels of residual heterozygosity in the three RIL populations. At

the SNP level we observed low levels of heterozygosity consistent with Mendelian expectation.

Residual heterozygosity was broadly uniform across the genome, with no clear regions enriched

for heterozygous SNP calls.

Hybrid incompatibilities as a source of TRD

In each RIL population, we identified several instances of two-locus epistasis (10 in the

CAC110 RILs, 4 in the CAC162 RILs, 9 in the CAC415 RILs: Table S4, Table 2). All but one of these

23 cases are due to a deficit of heterospecific allele combinations, consistent with the action of

hybrid incompatibilities reducing viability or fertility during RIL formation. In 16 of these cases,

the underlying mechanism seems to be epistatic selection favoring parental genotypes – that is,

we observed an excess of *M. guttatus* genotypes or *M. nasutus* genotypes at *both* loci. In six

cases, however, there are clear deficits of only one of the two heterospecific genotype

combinations *and* the expected distortion stemming from selection against the incompatible

genotype at each locus (Table 2). This asymmetry is a hallmark of Dobzhansky-Muller

incompatibilities. In four of the six cases, loci involved in these incompatibilities occur

exclusively in non-introgressed regions. However, in two of the pairs, one or both loci reside in

introgressed blocks, suggesting there may also be segregating incompatibilities within CAC *M.*

nasutus. As with overall patterns of TRD, regions involved in two-locus epistasis are largely

population specific (although in some cases, we likely have limited power to detect shared

epistasis due to small sample sizes and genetic background effects).

Genetics of hybrid fertility traits

Despite six generations of selfing, all three RIL populations exhibited substantial

variation in male fertility. Some of this variation might be due to inbreeding depression in *M.*

guttatus parental lines, which showed lower pollen viability than the *M. nasutus* parent (Figure

S13). However, heterospecific interactions are likely also involved, as all three RIL populations

included a considerable fraction of individuals (~10-22%) with pollen viability <50% (well below

the *M. guttatus* average of 72%). In the CAC110 RIL population, we mapped five pollen viability

QTLs accounting for ~36% of variance, including two separate epistatic pairs (Table 3, Figure 4).

One of these epistatic pairs leads to reduced male fertility in RILs with *M. nasutus* alleles on LG4

and *M. guttatus* alleles on LG14 (Figure S14A). The pattern of TRD at these same loci (excess of

M. guttatus on LG4 and *M. nasutus* alleles on LG14, Figure 3) might suggest the action of a

gametic incompatibility that reduces pollen viability, but other genetic mechanisms are also

possible (e.g., a sporophytic incompatibility that reduces pollen viability to the point of

preventing effective self-fertilization/seed production). The second epistatic pair in the CAC110

RIL population – between QTLs on LG12 and LG14 – also arises due to a negative heterospecific

interaction (RILs homozygous for *M. nasutus* alleles on LG12 and *M. guttatus* on LG14 have the

lowest fertility: Figure S14A), although *M. nasutus* genotypes have higher pollen fertility at each

QTL individually (Table 3). At the remaining QTL for male fertility on LG11, *M. nasutus* alleles

result in lower pollen viabilities (opposite to parental values), suggesting there could be

additional unmapped loci that interact with this QTL to decrease male fertility. In the CAC162

RIL population, we mapped two QTLs for pollen viability (accounting for ~28% of the variance),

but because we detected no epistasis between them and *M. nasutus* alleles increased fertility at

both, these effects might be due to inbreeding depression or to incompatibilities with additional

loci below our detection thresholds. We mapped no QTLs for male fertility in the CAC415 RIL

population.

For female fertility, all three *M. guttatus* parents had much lower fruit production than

the *M. nasutus* parent, suggesting substantial inbreeding depression affecting this trait (Figure

S13). Accordingly, all QTLs we identified – three in the CAC110 RIL and one in the CAC415 RIL –

were additive with *M. nasutus* alleles increasing fertility (Table 3).

Discussion

Closely related species living in secondary sympatry, especially those displaying

incomplete reproductive isolation and still experiencing gene flow, are an excellent opportunity

to investigate the evolution and maintenance of reproductive barriers in nature. Here we

construct three RIL populations from crosses in such a system, sister species *Mimulus guttatus*

and *M. nasutus,* from the sympatric site Catherine Creek. This multi-parent design allows for a

uniquely powerful assessment of the hybrid incompatibilities maintaining species boundaries in

a naturally admixed and highly genetically diverse population, assessing the effects of genetic

variation in *M. guttatus* on the outcomes of hybridization. Overall, patterns of transmission ratio

distortion and long-range linkage disequilibrium, paired with mapping of fertility in the

experimental hybrid populations reveal the hallmarks of substantial segregating hybrid

incompatibilities, as well as structural variation between this pair of naturally hybridizing taxa.

Additionally, these RILs are now a permanent resource to be used in research on adaptation and

speciation at CAC, allowing for continued investigations into the genetic basis of hybrid

incompatibilities, and hybrid fitness in replicate, genetically identical individuals across

environmental conditions.

Introgression in sympatry: M. nasutus ancestry in CAC M. guttatus

Suggestive of a long history of admixture, including recent hybridization events at

Catherine Creek, and consistent with previous analyses (Brandvain *et al.*, 2014; Kenney &

Sweigart, 2016) we identified substantial introgression from *M. nasutus* across the genomes of

three CAC *M. guttatus* inbred lines. Both the amount and location of introgression varies greatly

among the *M. guttatus* lines. Although variable, introgression within the genome does appear

non-random, with some consistent loci showing evidence of introgression in all three lines

(Figure S2). This is consistent with previous findings, which identified frequent introgression in a

large sample of wild individuals, with some loci more commonly introgressed than expected by

chance (Kenney & Sweigart, 2016). These overrepresented areas of introgression may indicate

adaptive introgression (Suarez-Gonzalez *et al.*, 2018) and therefore warrant further

investigation of additional sequenced *M. guttatus* lines from CAC, especially wild collected

individuals (as the process of creating inbred lines will likely influence observed levels of

admixture in lab created lines). While our data suggest adaptive introgression is likely not

ubiquitous, there may be small regions of *M. nasutus* variation (most notably, the distal regions

of Chr03 Chr04; Figure S2) that are being preferentially retained in *M. guttatus* genomes.

M. nasutus as a source of structural variation in sympatric M. guttatus

The ongoing and prevalent hybridization occurring at CAC also appears to have strong

impacts on segregating variation in genome architecture in CAC *M. guttatus*. Genome wide, the

three linkage maps produced are largely colinear, and have similar lengths (Table 1), following

the physical order of the reference genome. However, there are notable local differences

associated with segregating structural variation in CAC *M. guttatus* (Figure 2, Figure S4). We

identified three previously unidentified structural variants segregating within *M. guttatus* at CAC

with high confidence. Incredibly, two of these variants appear to have been introduced into *M.*

guttatus through admixture with *M. nasutus* (Figure S5). The most admixed *M. guttatus* line,

CAC110, carries the *M. nasutus* orientation of inv1.2 and inv13 in regions which show a high

probability of recent introgression (Figure S4, Figure S5, Figure S2). As inversions can help to

prevent the breakup of coevolved alleles, such as in the known inversion distinguishing

perennial and annual ecotypes of *M. guttatus* (Lowry & Willis, 2010; Twyford & Friedman,

2015), these inverted regions, will be worth comparing to past and future QTL mapping studies,

as introgression of these loci may be expected to have profound effects on species boundaries

and fitness.

Variation in transmission ratio distortion across experimental hybrid populations

TRD is frequently variable across the RIL populations, with the *M. guttatus* parent greatly

influencing patterns of distortion. In many instances, one *M. guttatus* allele is favored over the

M. nasutus allele in one population, while others are highly disfavored, or inherited in a

Mendelian fashion. Similar patterns, in which distinct crosses show vastly different patterns of

TRD, is consistent with previous crosses between species/ecotypes in *Mimulus* (Hall & Willis,

2005), however, such large scale differences have rarely been investigated within populations.

Of particular interest, however, are the small number of consistently distorted regions across

the RIL populations, which may point to a conserved mechanism of TRD in these instances. For

example, across all three maps, a ~2Mb region on the proximal arm of chromosome 14 shows

an excess of *M. guttatus* alleles. Interestingly, this same region shows evidence of distortion in

one previous analyses of TRD between *M. guttatus* ecotypes, but *not* between allopatric *M.*

nasutus and *M. guttatus* populations in that study (Hall & Willis, 2005). It therefore, may be an

especially good candidate for further investigations of loci contributing to distortion specifically

at CAC.

Further analysis will be needed to begin to determine likely agents causing the pervasive

TRD observed in this experiment. Distortion may arise for a number of reasons, each of which

have potentially important implications for species maintenance. Among others, meiotic drive

(Fishman & Saunders, 2008), gamete killers (Koide *et al.*, 2008), pollen competition (Fishman *et*

al., 2008), and Dobzhansky-Muller incompatibilities which effect the viability of gametes or

zygotes (Leppälä *et al.*, 2013; Kerwin & Sweigart, 2017) can all effect genomic patterns of

distortion. Additionally, cytonuclear incompatibilities, which we are unable to identify given our

unidirectional crossing design, may impact patterns of TRD if the three *M. guttatus* genomes

interact differently with CAC9 cytoplasmic factors. Reciprocal crosses in these same CAC

individuals would allow for the investigation of this possibility. Here we focus on instances

where transmission patterns at individual loci suggest selection against epistatic incompatible

alleles, indicative of DMI loci, but these other causes certainly influence variation in distortion

across the genome.

Evidence for multiple epistatic hybrid incompatibility loci

Overall, patterns of observed TRD and linkage disequilibrium in the three RIL

populations, paired with QTL mapping of fertility, reveal highly complex epistatic relationships

between alleles across the genome. It is likely that more complicated multi-locus

incompatibilities, that we don't have the sample size or power to full detect, are present,

contributing to the observed discordance between two-locus epistasis and patterns of single

locus TRD. Further explaining these patterns, many regions of high interchromosomal LD display

patterns of TRD consistent with epistatic selection for one parental genotype. Such patterns can

be explained by diverse mechanisms including selection against trait mismatches (Thompson *et*

al., 2022), in the case of selection for *M. nasutus* alleles, avoidance of inbreeding depression, or

environmental selection during greenhouse growth. These factors are all likely to influence

patterns of distortion, further complicating the interpretation of trends. The patterns we

identified of LD and distortion, caused by putative DMIs, could be the result of negative

epistasis occurring at either the gametic or zygotic stage. Although we are not able to

definitively distinguish between these different potential mechanisms leading to incompatible

epistatic interactions, because incompatibilities acting at the gametic stage would require both

a strong effect on both pollen genotype *and* pollen number to effectively purge incompatibility

loci, this option seems more in line with our observation of significant distortion without

complete purging.

Of the *M. guttatus* lines we crossed, CAC110 had the greatest proportion of *M. nasutus*

ancestry. We may have expected that this pervasive introgression might limit the potential

effects of hybrid incompatibilities, as *M. nasutus* alleles, entirely compatible with each other,

increase in frequency. However, we do not observe this trend. Even with nearly ¼ of its genome

introgressed from *M. nasutus,* multiple postzygotic barriers are still present in the CAC110 RIL

population. As such, long term hybrid persistence at CAC does not seem likely, as the

maintenance of a high proportion of *M. nasutus* ancestry following hybridization (through self-

fertilization, or mating with other admixed individuals) is likely to retain a substantial load of

hybrid incompatibilities. Rather, backcrossing to *M. guttatus* and purging *M. nasutus* ancestry

would seem to be most stable. This finding, along with the relatively small effect sizes of the

identified fertility QTLs, supports previous evidence for broad genome wide selection against *M.*

nasutus ancestry in an *M. guttatus* background (Brandvain *et al.*, 2014), supporting a complex

and dynamic genetic basis of hybrid incompatibility at CAC, with many small effect loci across

the genome decreasing hybrid fitness in concert.

Despite six generations of selfing, which is expected to purge deleterious heterospecific

combinations, the CAC110 RILs retained at least two hybrid incompatibilities with effects on

pollen viability, both of which lie in non-introgressed regions as expected. Despite their

maintenance in the CAC110 RILs, likely due to supplemental pollination and our concerted effort

to retain lines, these genotypes are likely to be under strong selection in nature, where pollen

limitation is more likely. Interestingly, these interacting loci did not appear in our tests for two-

locus epistasis. Although these loci are associated with pollen viability, and may be expected to

cause two-locus distortion, in these cases even the incompatible genotype class retained

intermediate pollen viability (~55-70% viable), and therefore was maintained in the RIL

population during the selfing process. If this strong, but incomplete sterility is the norm, it may

also explain why seemingly incompatible genotypes have not been entirely purged from the

population during RIL construction. Additionally, incompatibilities acting in gametes rather than

zygotes wouldn't necessarily lead to complete purging of incompatible genotypes even if

selection is strong.

Although we found no significant epistatic loci interacting with the remainder of the

QTLs we identified, they did frequently show some evidence of epistasis among them, although

it did not reach significance. This seems to be evidence of other subtle epistatic interactions

influencing hybrid fertility that warrant further investigation with larger sample sizes.

Additionally, the fact that we still see DMI-like epistasis in areas of introgression suggests that

there may also be segregating incompatibilities within CAC *M. nasutus*. Variation in

incompatibility loci has been identified previously between populations of *M. nasutus* (Sweigart

et al., 2007), but this may be unlikely within this highly inbred population. Alternatively, there

may be small blocks of true *M. guttatus* variation within these diagnosed blocks of introgression

which are harboring incompatible alleles.

Together, these results provide a wealth of evidence for the maintenance of substantial

postzygotic reproductive isolation between *M. nasutus* and *M. guttatus* in sympatry. Despite a

long history of admixture, these intrinsic barriers remain pervasive, limiting the homogenizing

effects of geneflow between species, and are likely integral to the maintenance of these species.

Author Contributions

Research conceived and designed by SJM and ALS. Data collected and analyzed by SJM.

Manuscript written by SJM and ALS.

Data Accessibility

All whole genome sequence data and RADseq data will be deposited at the NCBI Sequence Read

Archive upon acceptance. Other data and scripts will be made available from the Dryad Digital

Repository.

Acknowledgements

We thank Amanda Kenney for seed collection at Catherine Creek. Jill Anderson, Lisa Donovan,

Bob Schmitz, Mike White provided helpful discussions and guidance. This work was supported

by a National Science Foundation grant [IOS-1827645 to ALS]; [Rules of Life]; funds from the

University of Georgia Research Foundation [to ALS]; the National Institute of General Medical

Sciences of the National Institute of Health [T32GM007103 to SJM]; and the UGA Genetics Jan

and Kirby Alton Graduate Fellowship [to SJM].

Conflict of Interest

The authors declare no conflicts of interest.

ORCID

Samuel J. Mantel https://orcid.org/0000-0001-6392-4022

References

Ali OA, O'Rourke SM, Amish SJ, Meek MH, Luikart G, Jeffres C, Miller MR. **2016**. RAD Capture (Rapture): Flexible and Efficient Sequence-Based Genotyping. *Genetics* **202**: 389–400.

Arends D, Prins P, Jansen RC, Broman KW. **2010**. R/qtl: high-throughput multiple QTL mapping. *Bioinformatics* **26**: 2990–2992.

Aylor DL, Valdar W, Foulds-Mathes W, Buus RJ, Verdugo RA, Baric RS, Ferris MT, Frelinger JA, Heise M, Frieman MB, *et al.* 2011. Genetic analysis of complex traits in the emerging Collaborative Cross. *Genome Research* **21**: 1213–1222.

Barker WR, Nesom GL, Beardsley PM, Fraga NS. **2012**. A taxonomic conspectus of Phrymaceae: A narrowed circumscription for Mimulus, new and resurrected genera, and new names and combinations. *Phytoneuron* **2012–39**: 1–60.

Bolger AM, Lohse M, Usadel B. **2014**. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.

Brandvain Y, Kenney AM, Flagel L, Coop G, Sweigart AL. **2014**. Speciation and Introgression between Mimulus nasutus and Mimulus guttatus. *PLOS Genetics* **10**: e1004410.

Broman KW, Sen S. **2009**. *A Guide to QTL Mapping with R/qtl*. New York, NY: Springer New York.

Caeiro-Dias G, Brelsford A, Meneses-Ribeiro M, Crochet P-A, Pinho C. **2023**. Hybridization in late stages of speciation: Strong but incomplete genome-wide reproductive isolation and 'large Z-effect' in a moving hybrid zone. *Molecular Ecology* **32**: 4362–4380.

Catchen JM, Amores A, Hohenlohe P, Cresko W, Postlethwait JH. **2011**. Stacks: Building and Genotyping Loci De Novo From Short-Read Sequences. *G3 Genes|Genomes|Genetics* **1**: 171–182.

Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA. **2013**. Stacks: an analysis tool set for population genomics. *Molecular Ecology* **22**: 3124–3140.

Colomé-Tatché M, Johannes F. **2016**. Signatures of Dobzhansky–Muller Incompatibilities in the Genomes of Recombinant Inbred Lines. *Genetics* **202**: 825–841.

Coyne JA, Orr HA. **2004**. *Speciation*. Oxford, New York: Oxford University Press.

Cutter AD. **2012**. The polymorphic prelude to Bateson–Dobzhansky–Muller incompatibilities. *Trends in Ecology & Evolution* **27**: 209–218.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST, *et al.* 2011. The variant call format and VCFtools. *Bioinformatics* **27**: 2156–2158.

Darwin C. **1859**. *On the Origin of Species*.

Delmore KE, Hübner S, Kane NC, Schuster R, Andrew RL, Câmara F, Guigó R, Irwin DE. **2015**.
Genomic analysis of a migratory divide reveals candidate genes for migration and implicates
selective sweeps in generating islands of differentiation. *Molecular Ecology* **24**: 1873–1888.

Dobzhansky T. **1937**. *Genetics and the Origin of Species*. New York, NY: Columbia University
Press.

Durbin R, Eddy SR, Krogh A, Mitchison G. **1998**. *Biological Sequence Analysis: Probabilistic*
Models of Proteins and Nucleic Acids. Cambridge: Cambridge University Press.

Ellegren H, Smeds L, Burri R, Olason PI, Backström N, Kawakami T, Künstner A, Mäkinen H,
Nadachowska-Brzyska K, Qvarnström A, *et al.* 2012. The genomic landscape of species
divergence in Ficedula flycatchers. *Nature* **491**: 756–760.

Feder JL, Roethele JB, Filchak K, Niedbalski J, Romero-Severson J. **2003**. Evidence for Inversion
Polymorphism Related to Sympatric Host Race Formation in the Apple Maggot Fly, Rhagoletis
pomonella. *Genetics* **163**: 939–953.

Fishman L, Aagaard J, Tuthill JC. **2008**. TOWARD THE EVOLUTIONARY GENOMICS OF
GAMETOPHYTIC DIVERGENCE: PATTERNS OF TRANSMISSION RATIO DISTORTION IN
MONKEYFLOWER (MIMULUS) HYBRIDS REVEAL A COMPLEX GENETIC BASIS FOR CONSPECIFIC
POLLEN PRECEDENCE. *Evolution* **62**: 2958–2970.

Fishman L, McIntosh M. **2019**. Standard Deviations: The Biological Bases of Transmission Ratio
Distortion. *Annual Review of Genetics* **53**: 347–372.

Fishman L, Saunders A. **2008**. Centromere-Associated Female Meiotic Drive Entails Male Fitness
Costs in Monkeyflowers. *Science* **322**: 1559–1562.

Fishman L, Willis JH, Wu CA, Lee Y-W. **2014**. Comparative linkage maps suggest that fission, not
polyploidy, underlies near-doubling of chromosome number within monkeyflowers (Mimulus;
Phrymaceae). *Heredity* **112**: 562–568.

Flagel LE, Blackman BK, Fishman L, Monnahan PJ, Sweigart A, Kelly JK. **2019**. GOOGA: A
platform to synthesize mapping experiments and identify genomic structural diversity. *PLOS*
Computational Biology **15**: e1006949.

Flagel LE, Willis JH, Vision TJ. **2014**. The Standing Pool of Genomic Structural Variation in a
Natural Population of Mimulus guttatus. *Genome Biology and Evolution* **6**: 53–64.

Good JM, Handel MA, Nachman MW. **2008**. ASYMMETRY AND POLYMORPHISM OF HYBRID
MALE STERILITY DURING THE EARLY STAGES OF SPECIATION IN HOUSE MICE. *Evolution* **62**: 50–
65.

Hall MC, Willis JH. **2005**. Transmission Ratio Distortion in Intraspecific Hybrids of *Mimulus*
guttatus: Implications for Genomic Divergence. *Genetics* **170**: 375–386.

Hoffmann AA, Rieseberg LH. **2008**. Revisiting the Impact of Inversions in Evolution: From
Population Genetic Markers to Drivers of Adaptive Shifts and Speciation? *Annual Review of*
Ecology, Evolution, and Systematics **39**: 21–42.

Jansen RC, Stam P. **1994**. High resolution of quantitative traits into multiple loci via interval
mapping. *Genetics* **136**: 1447–1455.

Jones FC, Grabherr MG, Chan YF, Russell P, Mauceli E, Johnson J, Swofford R, Pirun M, Zody
MC, White S, *et al.* 2012. The genomic basis of adaptive evolution in threespine sticklebacks.
Nature **484**: 55–61.

Kearns C, Inouye D. **1993**. *Techniques for Pollination Biologists*. Niwot, CO: University of
Colorado Press.

Kenney AM, Sweigart AL. **2016**. Reproductive isolation and introgression between sympatric
Mimulus species. *Molecular Ecology* **25**: 2499–2517.

Kerwin RE, Sweigart AL. **2017**. Mechanisms of Transmission Ratio Distortion at Hybrid Sterility
Loci Within and Between Mimulus Species. *G3 Genes|Genomes|Genetics* **7**: 3719–3730.

Koide Y, Ikenaga M, Sawamura N, Nishimoto D, Matsubara K, Onishi K, Kanazawa A, Sano Y.
2008. The Evolution of Sex-Independent Transmission Ratio Distortion Involving Multiple Allelic
Interactions at a Single Locus in Rice. *Genetics* **180**: 409–420.

Larson EL, White TA, Ross CL, Harrison RG. **2014**. Gene flow and the maintenance of species
boundaries. *Molecular Ecology* **23**: 1668–1678.

Leppälä J, Bokma F, Savolainen O. **2013**. Investigating Incipient Speciation in Arabidopsis lyrata
from Patterns of Transmission Ratio Distortion. *Genetics* **194**: 697–708.

Li H. **2013**. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv:1303.3997 [q-bio].

Li H, Durbin R. **2009**. Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics **25**: 1754–1760.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
1000 Genome Project Data Processing Subgroup. **2009**. The Sequence Alignment/Map format
and SAMtools. *Bioinformatics* **25**: 2078–2079.

Long AD, Macdonald SJ, King EG. **2014**. Dissecting complex traits using the Drosophila Synthetic
Population Resource. *Trends in Genetics* **30**: 488–495.

Lowry DB, Sobel JM, Angert AL, Ashman T-L, Baker RL, Blackman BK, Brandvain Y, Byers KJRP,
Cooley AM, Coughlan JM, *et al.* 2019. The case for the continued use of the genus name
Mimulus for all monkeyflowers. *TAXON* **68**: 617–623.

Lowry DB, Willis JH. **2010**. A Widespread Chromosomal Inversion Polymorphism Contributes to
a Major Life-History Transition, Local Adaptation, and Reproductive Isolation. *PLOS Biology* **8**:
e1000500.

Mantel SJ, Sweigart AL. **2019**. Divergence in drought-response traits between sympatric species
of Mimulus. *Ecology and Evolution* **9**: 10291–10304.

Martin NH, Willis JH. **2007**. Ecological Divergence Associated with Mating System Causes Nearly
Complete Reproductive Isolation between Sympatric Mimulus Species. *Evolution* **61**: 68–82.

Mayr E. **1942**. *Systematics And The Origin Of Species*.

Mojica JP, Lee YW, Willis JH, Kelly JK. **2012**. Spatially and temporally varying selection on
intrapopulation quantitative trait loci for a life history trade-off in Mimulus guttatus. *Molecular*
Ecology **21**: 3718–3728.

Monnahan PJ, Kelly JK. **2015**. Naturally segregating loci exhibit epistasis for fitness. *Biology*
Letters **11**: 20150498.

Monnahan PJ, Kelly JK. **2017**. The Genomic Architecture of Flowering Time Varies Across Space
and Time in Mimulus guttatus. *Genetics* **206**: 1621–1635.

Moyle LC, Nakazato T. **2008**. Comparative Genetics of Hybrid Incompatibility: Sterility in Two
Solanum Species Crosses. *Genetics* **179**: 1437–1453.

Nesom GL, Fraga NS, Barker WR, Beardsley PM, Tank DC, Baldwin BG, Olmstead RG. **2019**.
Response to "The case for the continued use of the genus name Mimulus for all
monkeyflowers". *TAXON* **68**: 624–627.

Nolte AW, Gompert Z, Buerkle CA. **2009**. Variable patterns of introgression in two sculpin hybrid
zones suggest that genomic isolation differs among populations. *Molecular Ecology* **18**: 2615–
2627.

Noor MAF, Grams KL, Bertucci LA, Reiland J. **2001**. Chromosomal inversions and the
reproductive isolation of species. *Proceedings of the National Academy of Sciences* **98**: 12084–
12088.

Okonechnikov K, Conesa A, García-Alcalde F. **2016**. Qualimap 2: advanced multi-sample quality
control for high-throughput sequencing data. *Bioinformatics* **32**: 292–294.

Paradis E. **2010**. pegas: an R package for population genetics with an integrated–modular
approach. *Bioinformatics* **26**: 419–420.

Puzey JR, Willis JH, Kelly JK. **2017**. Population structure and local selection yield high genomic
variation in Mimulus guttatus. *Molecular Ecology* **26**: 519–535.

Rastas P. **2017**. Lep-MAP3: robust linkage mapping even for low-coverage whole genome
sequencing data. *Bioinformatics* **33**: 3726–3732.

Renaut S, Grassa CJ, Yeaman S, Moyers BT, Lai Z, Kane NC, Bowers JE, Burke JM, Rieseberg LH.
2013. Genomic islands of divergence are not affected by geography of speciation in sunflowers.
Nature Communications **4**: 1827.

Scott MF, Ladejobi O, Amer S, Bentley AR, Biernaskie J, Boden SA, Clark M, Dell'Acqua M,
Dixon LE, Filippi CV, *et al.* 2020. Multi-parent populations in crops: a toolbox integrating
genomics and genetic mapping with breeding. *Heredity* **125**: 396–416.

Servedio MR, Hermisson J. **2020**. The evolution of partial reproductive isolation as an adaptive
optimum. *Evolution* **74**: 4–14.

Servedio MR, Kirkpatrick M. **1997**. The Effects of Gene Flow on Reinforcement. *Evolution* **51**:
1764–1772.

Sotola VA, Berg CS, Samuli M, Chen H, Mantel SJ, Beardsley PA, Yuan Y-W, Sweigart AL,
Fishman L. **2023**. Genomic mechanisms and consequences of diverse postzygotic barriers
between monkeyflower species. : 2023.04.21.537885.

Suarez-Gonzalez A, Lexer C, Cronk QCB. **2018**. Adaptive introgression: a plant perspective.
Biology Letters **14**: 20170688.

Sweigart AL, Mason AR, Willis JH. **2007**. NATURAL VARIATION FOR A HYBRID INCOMPATIBILITY
BETWEEN TWO SPECIES OF MIMULUS. *Evolution* **61**: 141–151.

Sweigart AL, Willis JH. **2003**. Patterns of Nucleotide Diversity in Two Species of Mimulus Are
Affected by Mating System and Asymmetric Introgression. *Evolution* **57**: 2490–2506.

Taylor EB, Boughman JW, Groenenboom M, Sniatynski M, Schluter D, Gow JL. **2006**. Speciation
in reverse: morphological and genetic evidence of the collapse of a three-spined stickleback
(Gasterosteus aculeatus) species pair. *Molecular Ecology* **15**: 343–355.

Teeter KC, Thibodeau LM, Gompert Z, Buerkle CA, Nachman MW, Tucker PK. **2010**. The
Variable Genomic Architecture of Isolation Between Hybridizing Species of House Mice.
Evolution **64**: 472–485.

Thompson KA, Peichel CL, Rennison DJ, McGee MD, Albert AYK, Vines TH, Greenwood AK,
Wark AR, Brandvain Y, Schumer M, *et al.* 2022. Analysis of ancestry heterozygosity suggests
that hybrid incompatibilities in threespine stickleback are environment dependent. *PLOS Biology*
20: e3001469.

Twyford AD, Friedman J. **2015**. Adaptive divergence in the monkey flower Mimulus guttatus is
maintained by a chromosomal inversion. *Evolution; International Journal of Organic Evolution*
69: 1476–1486.

Van der Auwera G, O'Connor B. **2020**. *Genomics in the Cloud: Using Docker, GATK, and WDL in*
Terra (1st Edition). O'Reilly Media.

Zeng ZB. **1993**. Theoretical basis for separation of multiple linked gene effects in mapping
quantitative trait loci. *Proceedings of the National Academy of Sciences* **90**: 10972–10976.

Zeng ZB. **1994**. Precision mapping of quantitative trait loci. *Genetics* **136**: 1457–1468.

Zuellig MP, Sweigart AL. **2018**. A two-locus hybrid incompatibility is widespread, polymorphic,
and active in natural populations of Mimulus*. *Evolution* **72**: 2394–2405.

Figure 1. Introgression from *M. nasutus* into sympatric *M. guttatus*. A) *Dxy* calculated in 50kb windows across the first two chromosomes of the IM62 v3 *Mimulus guttatus* reference genome, between the parental lines of the three RIL populations (CAC9 v. CAC110, CAC9 v. CAC162, and CAC9 v. CAC415). Blue shaded regions indicate putatively admixed regions in the three focal CAC *M. guttatus* lines tested, diagnosed by a greater than a 95% posterior probability of *M. nasutus* ancestry (in 1kb windows) as inferred via the HMM (i.e. more recent coalescence with included *M. nasutus* samples than with included *M. guttatus* samples). Grey shaded regions indicate missing data (fewer than 5000 genotyped sites within any 50kb window). Red heatmaps above each chromosome indicate the density of centromeric repeats in the same 50kb windows. All chromosomes are shown in Figure S2. **B)** Heatmap showing the proportion of *M. nasutus* ancestry in each chromosome, and across the genome, of each of the three CAC *M. guttatus* lines used as parents of the RIL populations, as estimated as the proportion of 1kb windows with a greater than a 95% posterior probability of *M. nasutus* ancestry as inferred via the HMM. Darker blue shading indicates a higher proportion of *M. nasutus* ancestry.

Figure 2. Map position (cM) versus physical marker positions (Mb) for the first two chromosomes of the IM62 v3 *Mimulus guttatus* reference genome in the three RIL populations. Red heatmaps below each curve indicate the density of markers, measured as the distance in Mb to the nearest neighboring marker. Low SNP density, and the subsequent inability to genotype markers in areas of the *M. guttatus* parental genomes showing evidence of introgression from *M. nasutus* manifests as regions of low marker density on some chromosomes.

816
Figure 3. Transmission Ratio Distortion, cM, A) LG1 – LG7, B) LG8 – LG14. Genotype frequencies across the 14 linkage groups of the three RIL populations. Red dots indicate homozygous *M. guttatus* genotypes, yellow dots, homozygous *M. nasutus* genotypes. Bars at the top of each plot indicate regions showing significant transmission ratio distortion from the Mendelian 1:1 expectation by χ^2 tests at α = 0.01 (thin bars), and a more stringent Bonferonni-corrected α (thick bars, unique α for each chromosome in each RIL population depending on the number of markers tested, see Table S3 for values). Bar color indicates an excess of *M. guttatus* (red), or *M. nasutus* (yellow) genotypes. Markers are ordered by genetic position within each map (cM).

Figure 4. Male and female fertility QTL locations in each mapping population. Physical position (Mb) of the 1.5-LOD interval of each significant QTL is shown. Centromeres are indicated in black. Phenotype and RIL population indicated in key. Arrows indicate the position of peak marker(s). Arrows pointing left indicate the *M. guttatus* allele has the lower value, arrows pointing right indicate the *M. nasutus* allele has the lower value. QTLs with significant epistatic effects are indicated with shared symbols. The 1.5-LOD interval of the QTL underlying female fertility on Chr10 in the CAC415 RILs spans a breakpoint of the known derived inversion in the reference *M. guttatus* line IM62, leading to a disjunct interval.

830   **Table 1. Summary of linkage map length and intrascaffold recombination rate estimates.** Total
831   map length and intrascaffold recombination rate estimates for each chromosome, plus total
832   map length and average recombination rate in each of the three RIL populations.

		CAC110 RILs		CAC162 RILs		CAC415 RILs	
Chromosome	Reference Length (Mb)	Map Length (cM)	cM/Mb	Map Length (cM)	cM/Mb	Map Length (cM)	cM/Mb
1	13.88	134.368	9.68	66.451	4.79	74.470	5.37
2	20.54	86.970	4.23	79.725	3.88	85.500	4.16
3	23.80	82.263	3.46	64.360	2.70	67.161	2.82
4	22.81	82.362	3.61	84.446	3.70	92.822	4.07
5	21.77	90.440	4.15	82.863	3.81	92.963	4.27
6	20.89	99.115	4.74	95.776	4.58	104.195	4.99
7	21.88	72.643	3.32	60.311	2.76	89.304	4.08
8	27.69	109.184	3.94	117.963	4.26	117.700	4.25
9	25.15	63.801	2.54	92.286	3.67	98.934	3.93
10	22.12	89.656	4.05	106.661	4.82	104.910	4.74
11	33.60	137.499	4.09	88.907	2.65	91.633	2.73
12	25.84	101.548	3.93	82.763	3.20	81.868	3.17
13	30.77	117.730	3.83	78.866	2.56	105.060	3.41
14	28.29	134.029	4.74	109.580	3.87	128.236	4.53
Global	339.04	1401.608	4.13	1210.958	3.57	1334.756	3.94

Table 2. Positions of two-locus epistasis. Physical (Mb) positions of pairs of loci on each linkage group (LG) in the three RIL
populations that show LD (r) in the 95th percentile, as well as Chi-Square statistics from likelihood ratio tests with p<0.0001. Counts of
each genotypic class in the population are shown. Loci marked with an asterisk have peak r and χ^2 values at adjacent markers. Single
locus proportions of *M. guttatus* alleles (G Prop.) were used to determine if the pattern of TRD at each epistatic locus could be
explained by a two locus hybrid incompatibility (NG- or GN-), or if single locus distortion in the same direction better explains the
pattern (GG++, GG+, NN++, or NN+). The best candidates for two locus hybrid incompatibilities are shaded grey. See Table S5 for full
list of epistatic loci in LD 95th percentile.

Population	Locus 1 LG	Mb	Locus 2 LG	Mb	r	GG	GN	NG	NN	N	χ2	p	Locus 1 G Prop.	Locus 2 G Prop.	Pattern
CAC110	03	20.40 - 22.65	13	3.05 - 5.20	0.56	64	3	17	19	103	32.552	<0.0001	0.65	0.79	GG++
	05	3.10 - 20.75	11	2.55 - 4.90	0.54	39	21	1	22	83	29.035	<0.0001	0.72	0.48	NG-
	01	11.15 - 12.85	04	1.20 - 19.65	0.46	73	13	12	19	117	22.858	<0.0001	0.74	0.73	GG++
	02	2.85 - 18.95	13	21.75 - 23.65	0.45	54	1	26	14	95	21.079	<0.0001	0.58	0.84	GG++
	09	0.80 - 0.90	11	26.30 - 26.60	0.38	29	37	6	54	126	19.359	<0.0001	0.52	0.28	NG-
	05	2.40 - 20.90	11	5.05 - 10.20*	0.50	43	14	5	17	79	18.697	<0.0001	0.72	0.61	GG++
	03	0.15 - 18.95	11	24.35 - 25.30	0.40	29	24	12	56	121	18.567	<0.0001	0.44	0.34	NN++
	11	8.80 - 24.40	12	1.00 - 19.2	0.44	18	12	12	59	101	17.988	<0.0001	0.30	0.30	NN++
	01	3.30 - 5.50	11	4.00 - 5.75	0.47	29	9	10	24	72	16.515	<0.0001	0.53	0.54	GG+
	05	2.25 - 3.50	13	24.00 - 25.00	0.44	54	13	7	14	88	15.846	<0.0001	0.76	0.69	GG++
CAC162	04	19.90 - 22.80	14	2.35 - 18.15	0.50	29	10	14	44	97	24.708	<0.0001	0.40	0.44	NN+
	01	1.40 - 5.55	10	19.65 - 19.95	0.54	27	9	8	31	75	23.571	<0.0001	0.48	0.47	NN+
	06	6.30 - 15.10	12	6.20 - 7.75	0.47	9	50	23	15	97	21.637	<0.0001	0.61	0.33	HV
	01	2.35 - 5.45	03	0.00 - 20.65	0.45	28	21	7	42	98	20.628	<0.0001	0.50	0.36	NG-
CAC415	04	1.90 - 22.30	14	2.15 - 25.60	0.61	34	9	7	31	81	31.85	<0.0001	0.53	0.51	GG+
	06	17.00 - 19.25	10	19.50 - 20.25	0.43	38	0	22	13	73	22.217	<0.0001	0.52	0.82	GG+
	05	3.50 - 20.40	14	16.50 - 23.00	0.49	26	5	17	32	80	19.799	<0.0001	0.39	0.54	GN-
	07	13.20 - 21.80	08	22.30 - 27.70	0.48	23	7	14	38	82	19.719	<0.0001	0.37	0.45	NN+
	13	22.05 - 25.55	14	0.00 - 3.00	0.47	41	6	15	20	82	18.738	<0.0001	0.57	0.68	GG+
	06	9.65 - 16.00	08	23.60 - 25.55	0.44	22	10	12	37	81	15.892	<0.0001	0.40	0.42	NN+
	01	5.85 - 7.40	04	4.10 - 16.55	0.43	32	6	18	24	80	15.338	<0.0001	0.48	0.63	GN-
	01	7.70 - 11.70	13	24.20 - 25.15	0.42	24	4	22	30	80	15.279	<0.0001	0.35	0.58	GN-
	05	20.85 - 21.75	06	7.10 - 16.05	0.43	20	11	11	40	82	15.242	<0.0001	0.38	0.38	NN++

Table 3. Male and female fertility QTL locations and effects in each mapping population. Linkage group (LG), map (cM) and physical
(Mb) position of each significant QTL is shown. Additive effects (a) are negative when the *M. guttatus* allele has the lower value,
indicated in the summary column. QTLs with significant epistatic effects are shaded together. Percentage of RIL population variance
(PVE) is shown for each additive and epistatic effect. The 1.5-LOD interval of the QTLs underlying female fertility on LG10 in the
CAC415 RILs, marked with an asterisk, spans a breakpoint of the known derived inversion in the reference *M. guttatus* line IM62,
leading to a disjunct interval.

Population	Trait	LG	cM Position (1.5 LOD)	Mb Position (1.5 LOD)	Additive effect				Epistatic effect			
					a	PVE (%)	p	Summary	a x a	PVE (%)	p	Summary
CAC110 RILs	Male Fertility	4	26.975 (21.030 - 36.595)	16.70 - 16.75 (4.50 - 19.65)	-0.145	7.746	0.0008	G+	0.347	6.786	0.0004	NN+ (NG-)
		14	42.717 (37.467 - 48.317)	3.40 - 3.85 (3.05 - 4.30)	0.474	13.876	<0.0001	N+				
		11	132.951 (131.376 - 137.499)	26.65 - 26.70 (26.60 - 32.20)	-0.278	4.054	0.0059	G+				
		12	87.711 (80.538 - 101.548)	24.45 - 24.55 (23.25 - 25.80)	0.080	5.620	0.0055	N+	0.302	5.037	0.0022	NN+ (NG-)
		14	114.441 (112.343 - 116.889)	25.20 - 25.40 (25.00 - 25.60)	0.230	10.317	<0.0001	N+				
	Female Fertility	2	71.214 (62.399 - 86.970)	18.90 - 18.95 (18.40 - 20.55)	0.991	9.736	0.0001	N+				
		3	15.741 (11.890 - 20.758)	18.95 - 19.10 (18.40 - 19.95)	0.747	6.820	0.0012	N+				
		4	21.030 (16.833 - 26.101)	4.50 - 4.75 (3.85 - 16.70)	0.968	10.245	<0.0001	N+				
CAC162 RILs	Male Fertility	6	95.776 (90.673 - 95.776)	20.70 - 20.90 (19.25 - 20.90)	0.490	10.010	0.0011	N+				
		14	57.509 (54.192 - 61.594)	15.90 - 16.20 (7.05 - 16.80)	0.680	19.320	<0.0001	N+				
	Female Fertility	NA										
CAC415 RILs	Male Fertility	NA										
	Female Fertility	10	51.596 (48.660 - 53.554)	5.70 - 5.75 (4.50 - 5.75, 1.40 - 1.80)*	1.473	12.439	0.0022	N+				

849
850 **Figure S1. Recombinant inbred line (RIL) population crossing designs.** We constructed three
851 parallel RIL populations by crossing maternal *M. nasutus* line CAC9 to three paternal *M. guttatus*
852 lines: CAC110, CAC162, and CAC415. We self-pollinated one F1 from each cross, producing a
853 population of 500 F2s. We self-pollinated each F2 for at least 4 more generations with single
854 seed decent to produce three RIL populations of F6 (or later generation) inbred individuals.

856
Figure S2. Introgression from *M. nasutus* into sympatric *M. guttatus*. A) Chr01-Chr08, B) Chr09-Chr14. Dxy calculated in 50kb
windows across each chromosome of the IM62 v3 *Mimulus guttatus* reference genome, between the parental lines of the three RIL
populations (CAC9 v. CAC110, CAC9 v. CAC162, and CAC9 v. CAC415). Blue shaded regions indicate putatively admixed regions of the
three focal CAC *M. guttatus* lines tested, diagnosed by a greater than a 95% posterior probability of *M. nasutus* ancestry (in 1kb
windows) as inferred via the HMM (i.e. more recent coalescence with included *M. nasutus* samples than with included *M. guttatus*
samples). Grey shaded regions indicate missing data (fewer than 5000 genotyped sites within any 50kb window). Red heatmaps
above each chromosome indicate the density of centromeric repeats in the same 50kb windows.

864
Figure S3. Block size and characteristics of introgression from *M. nasutus* into sympatric *M. guttatus*. A) A fine scale look at
diagnosed blocks of introgression on the first chromosome in sympatric *M. guttatus* line CAC415. Blue shaded regions indicate
putatively admixed regions diagnosed by a greater than a 95% posterior probability of *M. nasutus* ancestry (in 1kb windows) as
inferred via the HMM (i.e. more recent coalescence with included *M. nasutus* samples than with included *M. guttatus* samples). Grey
shaded regions indicate missing data (fewer than 5000 genotyped sites within any 50kb window). Red heatmaps above the
chromosome indicate the density of centromeric repeats in the same 50kb windows. The shaded region between ~3 and 7.5 Mb is
shown in finer detail, with Dxy calculated in 5kb windows across this region. **B)** Histograms showing the length of each inferred block
of *M. nasutus* introgression in each of the sympatric *M. guttatus* lines used as parents for the three RIL populations. CAC110 average
block length: ~125kb, n=645 blocks, CAC162: ~66kb, n=476, CAC415: ~48bk, n=603. **C)** Dxy in each 50kb window in the three
sympatric *M. guttatus* lines used as parents for the three RIL populations binned by if that window has no inferred *M. nasutus*
introgression within it (No), part of the window contains inferred introgression (Partial), or the entire window is covered by an
inferred block of introgression (Full). Wilcox test, **** p<0.0001

Figure S4. Map position (cM) versus physical marker positions (Mb) of the IM62 v3 *Mimulus guttatus* **reference genome in the**

three RIL populations. A) LG1-7, B) LG8-14. Red heatmaps below each curve indicate the density of markers, measured as the

distance in Mb to the nearest neighboring marker. Low SNP density, and the subsequent inability to genotype markers in areas of the

M. guttatus parental genomes showing evidence of introgression from *M. nasutus* manifests as regions of low marker density on

some chromosomes. Recombination rate is substantially lower across the center of each chromosome, with a gap in genotyped

markers that colocalize with a metacentric (dicentric in the case of Chr11) chromosomal structure, identified via a high density of

centromeric repeats. Negative slope suggests inverted marker order relative to the IM62 v3 *M. guttatus* reference genome in both

the *M. nasutus* and *M. guttatus* parents. Blocks of markers spanning large physical distances but that map to nearly the same genetic

position suggest inversions between the *M. guttatus* and *M. nasutus* parents used to produce that RIL population.

Figure S5. Locations of newly identified and known inversions in *M. gutattus*. Physical positions (Mb) of each of the identified putative chromosomal inversions among the CAC *M. nasutus* and *M. guttatus* lines crossed. Centromeres are indicated in black. Arrows to the right of each chromosome indicate the observed orientation of each inversion in CAC9, CAC110, CAC162, and CAC415 from left to right. Arrows in yellow show evidence of *M. nasutus* ancestry, arrows in red show evidence of *M. guttatus* ancestry. The locations of previously identified inversions in *M. guttatus* are shown to the right of each chromosome. Brackets marked with question marks indicate further areas of interest, due to unique marker order, but which could not be confidently identified as inversions.

Figure S6. Transmission Ratio Distortion, Mb, A) Chr01-Chr07, B) Chr07-Chr14. Genotype frequencies across the first seven chromosomes of the three RIL populations. Red dots indicate homozygous *M. guttatus* genotypes, yellow dots, homozygous *M. nasutus* genotypes. Bars at the top of each plot indicate regions showing significant transmission ratio distortion from the Mendelian 1:1 expectation by χ^2 tests at α = 0.01 (thin bars), and a more stringent Bonferonni-corrected α (thick bars, unique α for each chromosome in each RIL population depending on the number of markers tested, see Table S3 for values). Bar color indicates and excess of *M. guttatus* (red), or *M. nasutus* (yellow) genotypes. Markers are ordered by physical position (Mb) within the IM62 v3 *Mimulus guttatus* reference genome.

905
906 **Figure S7. CAC110 RILs linkage disequilibrium heatmap.** LD heatmap for each pairwise marker
907 comparison across all 14 chromosomes, showing r values greater than the 95% quantile (>0.37)
908 in the first RIL population. Values range from 0 (white) to 1 (red).

919
Figure S8. CAC110 RILs linkage disequilibrium heatmap. LD heatmaps for each pairwise marker
comparison, showing r values greater than the 95% quantile (>0.37) in chromosome
comparisons with at least 1 marker above the cutoff in the first RIL population. Values range
from 0 (white) to 1 (red). Arrows indicates significant marker comparisons, when distorted
regions are small. **A)** Chr01 and Chr04 **B)** Chr01 and Chr11 **C)** Chr01 and Chr13 **D)** Chr02 and
Chr13 **E)** Chr03 and Chr11 **F)** Chr03 and Chr13 **G)** Chr04 and Chr13 **H)** Chr05 and Chr11 **I)** Chr05
and Chr13 **J)** Chr09 and Chr11 **K)** Chr11 and Chr12.

927
928 **Figure S9. CAC162 RILs linkage disequilibrium heatmap.** LD heatmap for each pairwise marker
929 comparison across all 14 chromosomes, showing r values greater than the 95% quantile (>0.41)
930 in the second RIL population. Values range from 0 (white) to 1 (red).

934
Figure S10. CAC162 RILs linkage disequilibrium heatmap. LD heatmaps for each pairwise
marker comparison, showing r values greater than the 95% quantile (>0.41) in chromosome
comparisons with at least 1 marker above the cutoff in the second RIL population. Values range
from 0 (white) to 1 (red). Arrows indicates significant marker comparisons, when distorted
regions are small. **A)** Chr01 and Chr03 **B)** Chr01 and Chr10 **C)** Chr04 and Chr14 **D)** Chr06 and
Chr12.

941
942 **Figure S11. CAC415 RILs linkage disequilibrium heatmap.** LD heatmap for each pairwise marker
943 comparison across all 14 chromosomes, showing r values greater than the 95% quantile (>0.39)
944 in the third RIL population. Values range from 0 (white) to 1 (red).

Figure S12. CAC415 RILs linkage disequilibrium heatmap.
LD heatmaps for each pairwise marker comparison, showing r values greater than the 95%
quantile (>0.39) in chromosome comparisons with at least 1 marker above the cutoff in the
third RIL population. Values range from 0 (white) to 1 (red). Arrows indicates significant marker
comparisons, when distorted regions are small. A) Chr01 and Chr04 B) Chr01 and Chr13 C)
Chr01 and Chr14 D) Chr03 and Chr06 E) Chr04 and Chr05 F) Chr04 and Chr08 G) Chr04 and
Chr14 H) Chr05 and Chr06 I) Chr05 and Chr07 J) Chr05 and Chr12 K) Chr05 and Chr14 L) Chr06
and Chr08 M) Chr06 and Chr10 N) Chr06 and Chr14 O) Chr07 and Chr08 P) Chr13 and Chr14.

Figure S13. Male and female viability distributions in the three RIL populations. Yellow lines indicate *M. nasutus* average values, red lines indicate *M. guttatus* average values.

974
975 **Figure S14. CAC110 RILs male fertility effect plots.** Average proportion viable pollen grains of
976 each genotypic class for each significant additive and epistatic QTL.

979
980 **Figure S15. CAC110 RILs female fertility effect plots.** Average seeds per fruit of each genotypic
981 class for each significant additive QTL.

984 **Figure S16. CAC162 RILs male fertility effect plots.** Average proportion viable pollen grains of
985 each genotypic class for each significant additive QTL.

986
987 **Figure S17. CAC415 RILs female fertility effect plots.** Average seeds per fruit of each genotypic
988 class for each significant additive QTL.

989 **Table S1. Geographic locations of *Mimulus* populations used.**

990 Information about the biology, geography, and sequencing of each sample used in this manuscript.

Line abbreviation	Species/Patry	Population	Latitude (N)	Longitude (W)	NCBI SRA Accession No.
CAC110	sympatric *M. guttatus*	Catherine Creek, Washington side of the Columbia River Gorge off of Hwy. 14	45°42′42′′	121°21′55′′	
CAC112					
CAC134					
CAC141					
CAC162					
CAC262					
CAC415					
CAC6					SRX525044
CAC9	sympatric *M. nasutus*	Catherine Creek, Washington side of the Columbia River Gorge off of Hwy. 14			SRX525048
IM62	allopatric *M. guttatus*	Iron Mountain, Hwy. 20, Linn County, Oregon	44° 24′ 03"	122° 08′ 16"	SRX115898
DPRG102	sympatric *M. guttatus*	Don Pedro Reservoir, Stanislaus National Forest, Junction of Hwy. 120 and Jacksonville Road, Tuolumne County, California	37° 49′ 44"	120° 20′ 38"	
SF5	allopatric *M. nasutus*	Sherar's Falls, Tygh Valley, Wasco County, Oregon	45°15′52′′	121°01′21′′	SRX116529
KOOT	allopatric *M. nasutus*	West of Kootenai National Forest, off MT 200, north of Cabinet Gorge Reservoir	48° 06′ 14"	115° 58′ 59"	SRX525049
NHN26	sympatric *M. nasutus*	Notch Hill, north of Nanoose Harbour, Vancouver Island, BC	49° 16′ 23′	124° 09′ 36"	SRX525051
DENT13007	*M. dentilobus*	Arizona	unknown	unknown	SRX030541

Table S2. Linkage map construction. Information regarding the number of markers, and
program options in LepMAP3 used to construct each of the three linkage maps featured in this
manuscript. Each chromosome was either run through SeparateChromosomes2 on its own
(independently) or as a group with other chromosomes in the same family (together). Physical
orders with better higher likelihoods when run through OrderMarkers2 are shown in bold. The
three linkage maps contain as total of 4,967 unique markers among them, 3,141 (~63%) of
which are present in all three maps. 513 markers (~10%) are unique to a single map, leaving
1,330 markers (~27%) shared between two of the three maps.

	LG	Number of markers	SeparateChromosomes2			OrderMarkers2		Retained positions	Unique positions
			Run conditions	LodLimit	Theta	Highest Likelihood 10 iterations	Physical Order Likelihood		
CAC110 RILs	1	156	independently	0.1	0.15	-7,008.42	-	156	60
	2	314	independently	0.1	0.15	-6,946.23	-	314	49
	3	292	together	20	0.3	-6,326.90	-	271	40
	4	279	independently	0.1	0.15	-6,730.64	-	279	47
	5	305	together	20	0.3	-10,574.70	-	277	42
	6	335	together	20	0.3	-7,543.75	-	332	57
	7	279	together	20	0.3	-4,606.16	-	252	46
	8	352	independently	0.1	0.15	-15,013.73	-15,514.09	349	88
	9	255	together	20	0.3	-4,355.72	-	255	41
	10	322	together	20	0.3	-8,102.44	-16,185.25	319	55
	11	115	independently	0.1	0.15	-9,735.51	**-8,846.31**	115	67
	12	317	independently	0.1	0.15	-11,752.58	-	304	68
	13	166	independently	0.1	0.15	-6,879.29	-7,732.36	138	50
	14	467	together	20	0.3	-14,236.16	-	467	84
	Global	3954	-	-	-	-	-	3828	794
CAC162 RILs	1	218	together	15	0.25	-3,311.21	-	213	36
	2	340	together	15	0.25	-5,011.07	-	340	47
	3	309	together	15	0.25	-8,771.58	-	291	54
	4	359	independently	0.1	0.15	-6,319.86	-	359	36
	5	331	together	15	0.25	-7,611.04	-	307	36
	6	315	independently	0.1	0.15	-7,782.91	-	315	48
	7	243	together	15	0.25	-6,165.75	-	222	36
	8	392	independently	0.1	0.15	-11,342.19	**-11,224.64**	392	54
	9	265	together	15	0.25	-6,308.35	-	265	45
	10	327	independently	0.1	0.15	-7,708.27	-	303	25
	11	260	independently	0.1	0.15	-8,465.66	-9,211.12	260	35
	12	338	together	15	0.25	-9,874.55	-	316	47
	13	340	together	15	0.25	-4,384.34	-	340	38
	14	466	together	15	0.25	-12,223.42	-	466	69
	Global	4503	-	-	-	-	-	4389	606
CAC415 RILs	1	186	together	15	0.25	-5,020.23	-	174	41
	2	314	together	15	0.25	-6,706.58	-	314	44
	3	302	independently	0.1	0.15	-7,280.74	-	289	34
	4	364	independently	0.1	0.15	-8,592.37	-	363	54
	5	327	independently	0.1	0.15	-7,690.35	-	325	50
	6	323	independently	0.1	0.15	-7,628.57	-	322	51
	7	245	together	15	0.25	-7,388.27	-	237	46
	8	438	together	15	0.25	-12,658.23	-	436	56
	9	268	independently	0.1	0.15	6,441.53	-	266	49
	10	258	together	15	0.25	-5,819.33	-	252	33
	11	289	together	15	0.25	-10,483.63	-11,605.05	255	50
	12	333	independently	0.1	0.15	-7,276.95	-	322	46
	13	343	independently	0.1	0.15	-8,136.78	-	343	58
	14	489	independently	0.1	0.15	-15,535.39	-	481	78
	Global	4479	-	-	-	-	-	4379	690

1000

1001 **Table S3. Positions of significant transmission ratio distortion.** Genetic (cM) and physical (Mb)
1002 position on each chromosome in the three RIL populations that show significant distortion
1003 (Direction) via Chi-Square test at α = 0.01 and Bonferonni-corrected α. Shading indicates
1004 putative contiguous regions of TRD.

Chromosome	cM	Mb	Direction	RIL Population	α Threshold
01	0.000 - 15.128	0.05 - 3.90	*M. guttatus*	CAC110	0.01
01	10.930 - 12.679	3.30 - 3.45	*M. guttatus*	CAC110	0.000167
01	32.397	4.10 - 4.25	*M. guttatus*	CAC110	0.01
01	122.469 - 134.368	11.05 - 12.85	*M. guttatus*	CAC110	0.01
01	124.218 - 134.368	11.10 - 12.85	*M. guttatus*	CAC110	0.000167
01	NA	NA	NA	CAC162	0.01
01	NA	NA	NA	CAC162	0.000278
01	1.220 - 28.693	0.50 - 3.60	*M. guttatus*	CAC415	0.01
01	12.204	3.40 - 3.45	*M. guttatus*	CAC415	0.000244
01	72.640 - 74.470	11.20 - 11.70	*M. nasutus*	CAC415	0.01
02	64.202 - 64.551	18.40 - 18.80	*M. nasutus*	CAC110	0.01
02	74.023 - 86.970	18.95 - 20.55	*M. nasutus*	CAC110	0.01
02	81.370 - 83.468	19.90 - 20.10	*M. nasutus*	CAC110	0.000204
02	34.741 - 36.783	3.40 - 5.25	*M. nasutus*	CAC162	0.01
02	38.823 - 39.844	5.50 - 12.05	*M. nasutus*	CAC162	0.01
02	40.864 - 43.926	12.15 - 15.70	*M. nasutus*	CAC162	0.01
02	NA	NA	NA	CAC162	0.000213
02	56.154	18.50 - 18.75	*M. guttatus*	CAC415	0.01
02	NA	NA	NA	CAC415	0.000227
03	NA	NA	NA	CAC110	0.01
03	NA	NA	NA	CAC110	0.000250
03	22.452 - 31.637	19.65 - 20.85	*M. nasutus*	CAC162	0.01
03	44.449 - 64.360	21.15 - 23.80	*M. nasutus*	CAC162	0.01
03	54.660 - 56.702	22.75 - 23.15	*M. nasutus*	CAC162	0.000185
03	0.000 - 17.078	0.25 - 19.95	*M. guttatus*	CAC415	0.01
03	0.000 - 10.978	1.20 - 19.95	*M. guttatus*	CAC415	0.000294
04	0.000 - 35.719	0.00 - 19.65	*M. guttatus*	CAC110	0.01
04	0.000 - 35.719	0.00 - 19.65	*M. guttatus*	CAC110	0.000213
04	39.221 - 44.119	19.80 - 20.15	*M. guttatus*	CAC110	0.01
04	0.000 - 16.873	0.05 - 3.70	*M. nasutus*	CAC162	0.01
04	0.000	0.05 - 0.45	*M. nasutus*	CAC162	0.000278
04	3.062 - 13.298	1.20 - 3.20	*M. nasutus*	CAC162	0.000278
04	16.873	3.45 - 3.70	*M. nasutus*	CAC162	0.000278
04	42.725 - 54.929	6.45 - 18.95	*M. guttatus*	CAC415	0.01
04	NA	NA	NA	CAC415	0.000185

1005

05	0.000 - 82.002	0.00 - 20.90	M. guttatus	CAC110	0.01
05	0.000 - 82.002	0.00 - 20.90	M. guttatus	CAC110	0.000238
05	NA	NA	NA	CAC162	0.01
05	NA	NA	NA	CAC162	0.000278
05	2.441 - 9.765	0.50 - 1.75	M. guttatus	CAC415	0.01
05	78.303 - 89.303	19.45 - 21.25	M. nasutus	CAC415	0.01
05	NA	NA	NA	CAC415	0.000200
06	NA	NA	NA	CAC110	0.01
06	NA	NA	NA	CAC110	0.000175
06	0.000 - 27.785	0.00 - 4.05	M. guttatus	CAC162	0.01
06	8.689 - 27.785	1.80 - 4.05	M. guttatus	CAC162	0.000208
06	67.161	8.50 - 9.25	M. guttatus	CAC162	0.01
06	69.203	13.90 - 14.30	M. guttatus	CAC162	0.01
06	69.713	14.35 - 14.40	M. guttatus	CAC162	0.01
06	69.713	14.60 - 14.70	M. guttatus	CAC162	0.01
06	57.782	6.9 - 7.05	M. nasutus	CAC415	0.01
06	60.221	7.60 - 9.20	M. nasutus	CAC415	0.01
06	60.221	9.5 - 9.65	M. nasutus	CAC415	0.01
06	60.221	10.25 - 15.40	M. nasutus	CAC415	0.01
06	NA	NA	NA	CAC415	0.000196
07	45.655 - 72.643	3.05 - 21.80	M. guttatus	CAC110	0.01
07	53.409 - 64.250	3.10 - 8.75	M. guttatus	CAC110	0.000217
07	65.299 - 72.643	15.00 - 21.80	M. guttatus	CAC110	0.000217
07	NA	NA	NA	CAC162	0.01
07	NA	NA	NA	CAC162	0.000278
07	56.933	6.90 - 6.95	M. nasutus	CAC415	0.01
07	55.713 - 73.440	8.30 - 14.40	M. nasutus	CAC415	0.01
07	64.900	12.85 - 13.05	M. nasutus	CAC415	0.000217
07	68.561 - 72.22	15.20 - 16.25	M. nasutus	CAC415	0.01
08	56.691 - 67.882	6.60 - 9.00	M. nasutus	CAC110	0.01
08	56.691 - 62.287	6.60 - 7.20	M. nasutus	CAC110	0.000114
08	63.686	7.90 - 7.95	M. nasutus	CAC110	0.000114
08	63.686	8.95 - 9.00	M. nasutus	CAC110	0.000114
08	53.543 - 53.893	14.90 - 15.15	M. nasutus	CAC110	0.01
08	50.745 - 66.133	15.85 - 17.05	M. nasutus	CAC110	0.01
08	65.084 - 66.133	15.85 - 16.05	M. nasutus	CAC110	0.000114
08	72.088 - 109.184	18.35 - 27.70	M. nasutus	CAC110	0.01
08	82.934 - 87.492	21.25 - 21.65	M. nasutus	CAC110	0.000114
08	93.096 - 100.091	21.95 - 23.80	M. nasutus	CAC110	0.000114
08	102.889	24.10 - 24.25	M. nasutus	CAC110	0.000114
08	104.288	25.35 - 26.40	M. nasutus	CAC110	0.000114
08	39.841 - 55.513	4.25 - 7.55	M. nasutus	CAC162	0.01
08	71.385 - 117.963	21.50 - 27.35	M. nasutus	CAC162	0.01
08	80.576 - 83.129	22.10 - 22.20	M. nasutus	CAC162	0.000185
08	95.938 - 117.963	22.70 - 27.35	M. nasutus	CAC162	0.000185
08	9.765	1.70 - 1.80	M. nasutus	CAC415	0.01
08	NA	NA	NA	CAC415	0.000179

1006

09	NA	NA	NA	CAC110	0.01
09	NA	NA	NA	CAC110	0.000244
09	NA	NA	NA	CAC162	0.01
09	NA	NA	NA	CAC162	0.000222
09	NA	NA	NA	CAC415	0.01
09	NA	NA	NA	CAC415	0.000204
10	19.249 - 20.648	5.10 - 5.30	M. nasutus	CAC110	0.01
10	88.257 - 89.656	14.90 - 15.80	M. guttatus	CAC110	0.01
10	NA	NA	NA	CAC110	0.000182
10	NA	NA	NA	CAC162	0.01
10	NA	NA	NA	CAC162	0.000400
10	0.000 - 1.220	0.00 - 0.77	M. nasutus	CAC415	0.01
10	53.554 - 104.91	1.80 - 22.00	M. guttatus	CAC415	0.01
10	53.554 - 95.746	1.80 - 20.25	M. guttatus	CAC415	0.000303
10	98.797	20.30 - 20.40	M. guttatus	CAC415	0.000303
11	54.170 - 56.618	5.05 - 5.75	M. guttatus	CAC110	0.01
11	82.175 - 85.323	9.60 - 9.90	M. nasutus	CAC110	0.01
11	94.429 - 132.951	10.35 - 26.70	M. nasutus	CAC110	0.01
11	102.823 - 103.523	23.50 - 23.65	M. nasutus	CAC110	0.000149
11	119.282 - 124.889	25.30 - 26.60	M. nasutus	CAC110	0.000149
11	0.000 - 2.551	0.05 - 1.10	M. nasutus	CAC162	0.01
11	1.021 - 2.551	0.75 - 1.10	M. nasutus	CAC162	0.000286
11	NA	NA	NA	CAC415	0.01
11	NA	NA	NA	CAC415	0.000200
12	15.399 - 87.186	1.05 - 24.50	M. nasutus	CAC110	0.01
12	21.705 - 25.202	1.70 - 5.20	M. nasutus	CAC110	0.000147
12	22.404 - 29.398	5.30 - 17.60	M. nasutus	CAC110	0.000147
12	38.49 - 60.193	18.60 - 21.70	M. nasutus	CAC110	0.000147
12	65.090	22.55 - 22.70	M. nasutus	CAC110	0.000147
12	68.239 - 80.538	22.75 - 23.60	M. nasutus	CAC110	0.000147
12	8.168 - 9.188	1.30 - 1.90	M. nasutus	CAC162	0.01
12	15.823 - 16.334	2.00 - 2.55	M. nasutus	CAC162	0.01
12	18.375 - 18.885	2.70 - 3.10	M. nasutus	CAC162	0.01
12	17.864 - 30.110	5.05 - 7.95	M. nasutus	CAC162	0.01
12	30.110	6.20 - 6.25	M. nasutus	CAC162	0.000213
12	30.621	18.55 - 18.60	M. nasutus	CAC162	0.01
12	30.621	18.55 - 18.60	M. nasutus	CAC162	0.000213
12	32.152 - 67.944	18.65 - 24.20	M. nasutus	CAC162	0.01
12	32.152 - 32.152	18.65 - 19.20	M. nasutus	CAC162	0.000213
12	37.259 - 45.425	19.80 - 23.40	M. nasutus	CAC162	0.000213
12	69.986 - 75.611	24.30 - 24.90	M. nasutus	CAC162	0.01
12	NA	NA	NA	CAC415	0.01
12	NA	NA	NA	CAC415	0.000217

1007

13	0.000 - 53.932	2.10 - 25.00	*M. guttatus*	CAC110	0.01
13	0.000 - 45.185	2.10 - 24.00	*M. guttatus*	CAC110	0.000200
13	88.949 - 117.730	26.50 - 30.80	*M. nasutus*	CAC110	0.01
13	88.949 - 117.730	26.50 - 30.80	*M. nasutus*	CAC110	0.000200
13	38.986 - 78.866	25.55 - 30.80	*M. nasutus*	CAC162	0.01
13	54.367 - 78.866	26.30 - 30.80	*M. nasutus*	CAC162	0.000263
13	61.649	25.50 - 25.55	*M. guttatus*	CAC415	0.01
13	NA	NA	NA	CAC415	0.000172
14	0.000 - 22.761	0.00 - 2.35	*M. guttatus*	CAC110	0.01
14	0.000 - 9.099	0.00 - 1.90	*M. guttatus*	CAC110	0.000119
14	0.000 - 3.062	0.00 - 0.60	*M. guttatus*	CAC162	0.01
14	NA	NA	NA	CAC162	0.000145
14	0.000 - 13.422	0.00 - 2.65	*M. guttatus*	CAC415	0.01
14	NA	NA	NA	CAC415	0.000128

1008

Table S4. Average Interchromosomal linkage disequilibrium (r) across all pairwise marker comparisons for each chromosome pair in each RIL population. N is the total number of pairwise marker comparisons. Average ILD of those marker comparisons which are greater than the 95% quantile (r95). N95 is the number of those pairwise comparisons that are greater than the 95% quantile. NA indicates there were no pairwise marker comparisons greater than the 95% quantile.

LG Comparison	r	N	r95	N95	Proportion	RIL Population
01,02	0.1003	2940	NA	0	0.0000	CAC110
01,02	0.0871	1692	NA	0	0.0000	CAC162
01,02	0.0822	1804	NA	0	0.0000	CAC415
01,03	0.0948	2400	NA	0	0.0000	CAC110
01,03	**0.2244**	**1944**	**0.4251**	**25**	**0.0129**	**CAC162**
01,03	0.0618	1394	NA	0	0.0000	CAC415
01,04	**0.0991**	**2820**	**0.4027**	**69**	**0.0245**	**CAC110**
01,04	0.1335	1296	NA	0	0.0000	CAC162
01,04	**0.1431**	**2214**	**0.4041**	**19**	**0.0086**	**CAC415**
01,05	0.1085	2520	NA	0	0.0000	CAC110
01,05	0.1097	1296	NA	0	0.0000	CAC162
01,05	0.1040	2050	NA	0	0.0000	CAC415
01,06	0.0697	3420	NA	0	0.0000	CAC110
01,06	0.0808	1728	NA	0	0.0000	CAC162
01,06	0.0677	2091	NA	0	0.0000	CAC415
01,07	0.1052	2760	NA	0	0.0000	CAC110
01,07	0.0818	1296	NA	0	0.0000	CAC162
01,07	0.0673	1886	NA	0	0.0000	CAC415
01,08	0.0745	5280	NA	0	0.0000	CAC110
01,08	0.1135	1944	NA	0	0.0000	CAC162
01,08	0.1408	2296	NA	0	0.0000	CAC415
01,09	0.0883	2460	NA	0	0.0000	CAC110
01,09	0.1120	1620	NA	0	0.0000	CAC162
01,09	0.0620	2009	NA	0	0.0000	CAC415
01,10	0.0722	3300	NA	0	0.0000	CAC110
01,10	**0.1510**	**900**	**0.4638**	**43**	**0.0478**	**CAC162**
01,10	0.0770	1353	NA	0	0.0000	CAC415
01,11	**0.1260**	**4020**	**0.3961**	**64**	**0.0159**	**CAC110**
01,11	0.0539	1260	NA	0	0.0000	CAC162
01,11	0.0785	2050	NA	0	0.0000	CAC415
01,12	0.0832	4080	NA	0	0.0000	CAC110

01,12	0.1656	1692	NA	0	0.0000	CAC162
01,12	0.0490	1886	NA	0	0.0000	CAC415
01,13	**0.1026**	**3000**	**0.3831**	**24**	**0.0080**	**CAC110**
01,13	0.0816	1368	NA	0	0.0000	CAC162
01,13	**0.1492**	**2378**	**0.4022**	**9**	**0.0038**	**CAC415**
01,14	0.0864	5040	NA	0	0.0000	CAC110
01,14	0.1142	2484	NA	0	0.0000	CAC162
01,14	**0.1319**	**3198**	**0.3934**	**5**	**0.0016**	**CAC415**
02,03	0.0867	1960	NA	0	0.0000	CAC110
02,03	0.0784	2538	NA	0	0.0000	CAC162
02,03	0.0911	1496	NA	0	0.0000	CAC415
02,04	0.0808	2303	NA	0	0.0000	CAC110
02,04	0.0946	1692	NA	0	0.0000	CAC162
02,04	0.0634	2376	NA	0	0.0000	CAC415
02,05	0.0616	2058	NA	0	0.0000	CAC110
02,05	0.1019	1692	NA	0	0.0000	CAC162
02,05	0.0842	2200	NA	0	0.0000	CAC415
02,06	0.0682	2793	NA	0	0.0000	CAC110
02,06	0.0975	2256	NA	0	0.0000	CAC162
02,06	0.0884	2244	NA	0	0.0000	CAC415
02,07	0.0848	2254	NA	0	0.0000	CAC110
02,07	0.0742	1692	NA	0	0.0000	CAC162
02,07	0.1123	2024	NA	0	0.0000	CAC415
02,08	0.0958	4312	NA	0	0.0000	CAC110
02,08	0.1219	2538	NA	0	0.0000	CAC162
02,08	0.0753	2464	NA	0	0.0000	CAC415
02,09	0.0952	2009	NA	0	0.0000	CAC110
02,09	0.0915	2115	NA	0	0.0000	CAC162
02,09	0.0974	2156	NA	0	0.0000	CAC415
02,10	0.0520	2695	NA	0	0.0000	CAC110
02,10	0.0531	1175	NA	0	0.0000	CAC162
02,10	0.1337	1452	NA	0	0.0000	CAC415
02,11	0.0792	3283	NA	0	0.0000	CAC110
02,11	0.1718	1645	NA	0	0.0000	CAC162
02,11	0.0674	2200	NA	0	0.0000	CAC415
02,12	0.0401	3332	NA	0	0.0000	CAC110
02,12	0.0882	2209	NA	0	0.0000	CAC162
02,12	0.0780	2024	NA	0	0.0000	CAC415
02,13	**0.1408**	**2450**	**0.4083**	**90**	**0.0367**	**CAC110**
02,13	0.1008	1786	NA	0	0.0000	CAC162

02,13	0.0684	2552	NA	0	0.0000	CAC415
02,14	0.0945	4116	NA	0	0.0000	CAC110
02,14	0.1041	3243	NA	0	0.0000	CAC162
02,14	0.0707	3432	NA	0	0.0000	CAC415
03,04	0.0496	1880	NA	0	0.0000	CAC110
03,04	0.0856	1944	NA	0	0.0000	CAC162
03,04	0.1287	1836	NA	0	0.0000	CAC415
03,05	0.1039	1680	NA	0	0.0000	CAC110
03,05	0.1435	1944	NA	0	0.0000	CAC162
03,05	0.0768	1700	NA	0	0.0000	CAC415
03,06	0.0940	2280	NA	0	0.0000	CAC110
03,06	0.0809	2592	NA	0	0.0000	CAC162
03,06	**0.1284**	**1734**	**0.4024**	**5**	**0.0029**	**CAC415**
03,07	0.0752	1840	NA	0	0.0000	CAC110
03,07	0.0582	1944	NA	0	0.0000	CAC162
03,07	0.1069	1564	NA	0	0.0000	CAC415
03,08	0.0851	3520	NA	0	0.0000	CAC110
03,08	0.1265	2916	NA	0	0.0000	CAC162
03,08	0.0799	1904	NA	0	0.0000	CAC415
03,09	0.0865	1640	NA	0	0.0000	CAC110
03,09	0.1260	2430	NA	0	0.0000	CAC162
03,09	0.1058	1666	NA	0	0.0000	CAC415
03,10	0.0897	2200	NA	0	0.0000	CAC110
03,10	0.1326	1350	NA	0	0.0000	CAC162
03,10	0.0958	1122	NA	0	0.0000	CAC415
03,11	**0.1867**	**2680**	**0.3860**	**69**	**0.0257**	**CAC110**
03,11	0.1129	1890	NA	0	0.0000	CAC162
03,11	0.0647	1700	NA	0	0.0000	CAC415
03,12	0.0964	2720	NA	0	0.0000	CAC110
03,12	0.0840	2538	NA	0	0.0000	CAC162
03,12	0.0761	1564	NA	0	0.0000	CAC415
03,13	**0.1453**	**2000**	**0.4824**	**64**	**0.0320**	**CAC110**
03,13	0.0684	2052	NA	0	0.0000	CAC162
03,13	0.1268	1972	NA	0	0.0000	CAC415
03,14	0.0543	3360	NA	0	0.0000	CAC110
03,14	0.1271	3726	NA	0	0.0000	CAC162
03,14	0.0718	2652	NA	0	0.0000	CAC415
04,05	0.0623	1974	NA	0	0.0000	CAC110
04,05	0.0706	1296	NA	0	0.0000	CAC162
04,05	**0.1399**	**2700**	**0.3988**	**3**	**0.0011**	**CAC415**

04,06	0.0690	2679	NA	0	0.0000	CAC110
04,06	0.0548	1728	NA	0	0.0000	CAC162
04,06	0.0895	2754	NA	0	0.0000	CAC415
04,07	0.0548	2162	NA	0	0.0000	CAC110
04,07	0.1111	1296	NA	0	0.0000	CAC162
04,07	0.0962	2484	NA	0	0.0000	CAC415
04,08	0.0768	4136	NA	0	0.0000	CAC110
04,08	0.1041	1944	NA	0	0.0000	CAC162
04,08	**0.1439**	**3024**	**0.3979**	**1**	**0.0003**	**CAC415**
04,09	0.0735	1927	NA	0	0.0000	CAC110
04,09	0.1238	1620	NA	0	0.0000	CAC162
04,09	0.1040	2646	NA	0	0.0000	CAC415
04,10	0.0626	2585	NA	0	0.0000	CAC110
04,10	0.1461	900	NA	0	0.0000	CAC162
04,10	0.0672	1782	NA	0	0.0000	CAC415
04,11	0.0755	3149	NA	0	0.0000	CAC110
04,11	0.0728	1260	NA	0	0.0000	CAC162
04,11	0.1005	2700	NA	0	0.0000	CAC415
04,12	0.0470	3196	NA	0	0.0000	CAC110
04,12	0.1031	1692	NA	0	0.0000	CAC162
04,12	0.1035	2484	NA	0	0.0000	CAC415
04,13	**0.1303**	**2350**	**0.3716**	**2**	**0.0009**	**CAC110**
04,13	0.1510	1368	NA	0	0.0000	CAC162
04,13	0.1289	3132	NA	0	0.0000	CAC415
04,14	0.0531	3948	NA	0	0.0000	CAC110
04,14	**0.1708**	**2484**	**0.4401**	**17**	**0.0068**	**CAC162**
04,14	**0.2961**	**4212**	**0.4512**	**832**	**0.1975**	**CAC415**
05,06	0.0675	2394	NA	0	0.0000	CAC110
05,06	0.1297	1728	NA	0	0.0000	CAC162
05,06	**0.0983**	**2550**	**0.4037**	**15**	**0.0059**	**CAC415**
05,07	0.0628	1932	NA	0	0.0000	CAC110
05,07	0.1130	1296	NA	0	0.0000	CAC162
05,07	**0.1240**	**2300**	**0.4012**	**4**	**0.0017**	**CAC415**
05,08	0.0508	3696	NA	0	0.0000	CAC110
05,08	0.0812	1944	NA	0	0.0000	CAC162
05,08	**0.1105**	**2800**	**0.3940**	**1**	**0.0004**	**CAC415**
05,09	0.0724	1722	NA	0	0.0000	CAC110
05,09	0.1111	1620	NA	0	0.0000	CAC162
05,09	0.1016	2450	NA	0	0.0000	CAC415
05,10	0.0599	2310	NA	0	0.0000	CAC110

05,10	0.0734	900	NA	0	0.0000	CAC162
05,10	0.1091	1650	NA	0	0.0000	CAC415
05,11	**0.2175**	**2814**	**0.4226**	**474**	**0.1684**	**CAC110**
05,11	0.1003	1260	NA	0	0.0000	CAC162
05,11	0.1079	2500	NA	0	0.0000	CAC415
05,12	0.0704	2856	NA	0	0.0000	CAC110
05,12	0.1033	1692	NA	0	0.0000	CAC162
05,12	**0.1176**	**2300**	**0.3957**	**4**	**0.0017**	**CAC415**
05,13	**0.1133**	**2100**	**0.3957**	**13**	**0.0062**	**CAC110**
05,13	0.0733	1368	NA	0	0.0000	CAC162
05,13	0.1121	2900	NA	0	0.0000	CAC415
05,14	0.0616	3528	NA	0	0.0000	CAC110
05,14	0.0864	2484	NA	0	0.0000	CAC162
05,14	**0.1589**	**3900**	**0.4167**	**94**	**0.0241**	**CAC415**
06,07	0.0763	2622	NA	0	0.0000	CAC110
06,07	0.0821	1728	NA	0	0.0000	CAC162
06,07	0.1154	2346	NA	0	0.0000	CAC415
06,08	0.0677	5016	NA	0	0.0000	CAC110
06,08	0.1192	2592	NA	0	0.0000	CAC162
06,08	**0.1273**	**2856**	**0.4106**	**23**	**0.0081**	**CAC415**
06,09	0.0496	2337	NA	0	0.0000	CAC110
06,09	0.1248	2160	NA	0	0.0000	CAC162
06,09	0.0741	2499	NA	0	0.0000	CAC415
06,10	0.0717	3135	NA	0	0.0000	CAC110
06,10	0.0845	1200	NA	0	0.0000	CAC162
06,10	**0.1326**	**1683**	**0.4304**	**10**	**0.0059**	**CAC415**
06,11	0.1348	3819	NA	0	0.0000	CAC110
06,11	0.0890	1680	NA	0	0.0000	CAC162
06,11	0.0867	2550	NA	0	0.0000	CAC415
06,12	0.0634	3876	NA	0	0.0000	CAC110
06,12	**0.2184**	**2256**	**0.4341**	**83**	**0.0368**	**CAC162**
06,12	0.1248	2346	NA	0	0.0000	CAC415
06,13	0.0843	2850	NA	0	0.0000	CAC110
06,13	0.0812	1824	NA	0	0.0000	CAC162
06,13	0.0995	2958	NA	0	0.0000	CAC415
06,14	0.0485	4788	NA	0	0.0000	CAC110
06,14	0.0892	3312	NA	0	0.0000	CAC162
06,14	**0.1300**	**3978**	**0.4029**	**1**	**0.0003**	**CAC415**
07,08	0.0530	4048	NA	0	0.0000	CAC110
07,08	0.1047	1944	NA	0	0.0000	CAC162

07,08	0.1350	2576	0.4180	32	0.0124	CAC415
07,09	0.0974	1886	NA	0	0.0000	CAC110
07,09	0.1520	1620	NA	0	0.0000	CAC162
07,09	0.0977	2254	NA	0	0.0000	CAC415
07,10	0.0691	2530	NA	0	0.0000	CAC110
07,10	0.0948	900	NA	0	0.0000	CAC162
07,10	0.0828	1518	NA	0	0.0000	CAC415
07,11	0.0750	3082	NA	0	0.0000	CAC110
07,11	0.0862	1260	NA	0	0.0000	CAC162
07,11	0.1393	2300	NA	0	0.0000	CAC415
07,12	0.0747	3128	NA	0	0.0000	CAC110
07,12	0.0811	1692	NA	0	0.0000	CAC162
07,12	0.0939	2116	NA	0	0.0000	CAC415
07,13	0.0689	2300	NA	0	0.0000	CAC110
07,13	0.1137	1368	NA	0	0.0000	CAC162
07,13	0.0472	2668	NA	0	0.0000	CAC415
07,14	0.0760	3864	NA	0	0.0000	CAC110
07,14	0.1120	2484	NA	0	0.0000	CAC162
07,14	0.0740	3588	NA	0	0.0000	CAC415
08,09	0.0781	3608	NA	0	0.0000	CAC110
08,09	0.1185	2430	NA	0	0.0000	CAC162
08,09	0.1120	2744	NA	0	0.0000	CAC415
08,10	0.0597	4840	NA	0	0.0000	CAC110
08,10	0.1139	1350	NA	0	0.0000	CAC162
08,10	0.0935	1848	NA	0	0.0000	CAC415
08,11	0.0939	5896	NA	0	0.0000	CAC110
08,11	0.0979	1890	NA	0	0.0000	CAC162
08,11	0.0908	2800	NA	0	0.0000	CAC415
08,12	0.0849	5984	NA	0	0.0000	CAC110
08,12	0.0863	2538	NA	0	0.0000	CAC162
08,12	0.0740	2576	NA	0	0.0000	CAC415
08,13	0.1100	4400	NA	0	0.0000	CAC110
08,13	0.0866	2052	NA	0	0.0000	CAC162
08,13	0.1011	3248	NA	0	0.0000	CAC415
08,14	0.0686	7392	NA	0	0.0000	CAC110
08,14	0.1274	3726	NA	0	0.0000	CAC162
08,14	0.0995	4368	NA	0	0.0000	CAC415
09,10	0.0904	2255	NA	0	0.0000	CAC110
09,10	0.1074	1125	NA	0	0.0000	CAC162
09,10	0.0758	1617	NA	0	0.0000	CAC415

09,11	**0.1469**	**2747**	**0.3752**	**2**	**0.0007**	**CAC110**
09,11	0.0711	1575	NA	0	0.0000	CAC162
09,11	0.0896	2450	NA	0	0.0000	CAC415
09,12	0.0639	2788	NA	0	0.0000	CAC110
09,12	0.1226	2115	NA	0	0.0000	CAC162
09,12	0.1003	2254	NA	0	0.0000	CAC415
09,13	0.0699	2050	NA	0	0.0000	CAC110
09,13	0.0717	1710	NA	0	0.0000	CAC162
09,13	**0.1417**	**2842**	**0.4009**	**1**	**0.0004**	**CAC415**
09,14	0.0612	3444	NA	0	0.0000	CAC110
09,14	0.1531	3105	NA	0	0.0000	CAC162
09,14	0.0679	3822	NA	0	0.0000	CAC415
10,11	0.0930	3685	NA	0	0.0000	CAC110
10,11	0.1011	875	NA	0	0.0000	CAC162
10,11	0.0721	1650	NA	0	0.0000	CAC415
10,12	0.0570	3740	NA	0	0.0000	CAC110
10,12	0.0980	1175	NA	0	0.0000	CAC162
10,12	0.1084	1518	NA	0	0.0000	CAC415
10,13	0.0757	2750	NA	0	0.0000	CAC110
10,13	0.1372	950	NA	0	0.0000	CAC162
10,13	0.0864	1914	NA	0	0.0000	CAC415
10,14	0.0664	4620	NA	0	0.0000	CAC110
10,14	0.1520	1725	NA	0	0.0000	CAC162
10,14	0.1194	2574	NA	0	0.0000	CAC415
11,12	**0.1682**	**4556**	**0.3902**	**135**	**0.0296**	**CAC110**
11,12	0.0680	1645	NA	0	0.0000	CAC162
11,12	0.0685	2300	NA	0	0.0000	CAC415
11,13	0.0919	3350	NA	0	0.0000	CAC110
11,13	0.0740	1330	NA	0	0.0000	CAC162
11,13	0.0794	2900	NA	0	0.0000	CAC415
11,14	0.0930	5628	NA	0	0.0000	CAC110
11,14	0.1356	2415	NA	0	0.0000	CAC162
11,14	**0.1458**	**3900**	**0.3904**	**1**	**0.0003**	**CAC415**
12,13	0.0909	3400	NA	0	0.0000	CAC110
12,13	0.0714	1786	NA	0	0.0000	CAC162
12,13	0.0941	2668	NA	0	0.0000	CAC415
12,14	0.1328	5712	NA	0	0.0000	CAC110
12,14	0.1018	3243	NA	0	0.0000	CAC162
12,14	0.1522	3588	NA	0	0.0000	CAC415
13,14	**0.1047**	**4200**	**0.3966**	**10**	**0.0024**	**CAC110**

13,14	0.1072	2622	NA	0	0.0000	CAC162
13,14	**0.1665**	**4524**	**0.4149**	**50**	**0.0111**	**CAC415**

1015

Population	Locus 1 LG	Locus 1 Mb	Locus 2 LG	Locus 2 Mb	r	GG	GN	NG	NN	N	χ2	p	Locus 1 G Prop.	Locus 2 G Prop.	Pattern
	03	20.40 - 22.65	13	3.05 - 5.20	0.56	64	3	17	19	103	32.552	<0.0001	0.65	0.79	GG++
	05	3.10 - 20.75	11	2.55 - 4.90	0.54	39	21	1	22	83	29.035	<0.0001	0.72	0.48	NG-
	01	11.15 - 12.85	04	1.20 - 19.65	0.46	73	13	12	19	117	22.858	<0.0001	0.74	0.73	GG++
	02	2.85 - 18.95	13	21.75 - 23.65	0.45	54	1	26	14	95	21.079	<0.0001	0.58	0.84	GG++
	09	0.80 - 0.90	11	26.30 - 26.60	0.38	29	37	6	54	126	19.359	<0.0001	0.52	0.28	NG-
	05	2.40 - 20.90	11	5.05 - 10.20*	0.50	43	14	5	17	79	18.697	<0.0001	0.72	0.61	GG++
	03	0.15 - 18.95	11	24.35 - 25.30	0.40	29	24	12	56	121	18.567	<0.0001	0.44	0.34	NN++
CAC110	11	8.80 - 24.40	12	1.00 - 19.2	0.44	18	12	12	59	101	17.988	<0.0001	0.30	0.30	NN++
	01	3.30 - 5.50	11	4.00 - 5.75	0.47	29	9	10	24	72	16.515	<0.0001	0.53	0.54	GG+
	05	2.25 - 3.50	13	24.00 - 25.00	0.44	54	13	7	14	88	15.846	<0.0001	0.76	0.69	GG++
	13	25.00 - 25.70	14	1.25 - 2.15	0.41	41	7	15	17	80	13.622	0.0002	0.60	0.70	GG+
	03	18.40 - 21.55	11	3.35 - 6.25	0.42	34	7	12	17	70	13.194	0.0003	0.59	0.66	GG+
	01	7.80 - 9.60	13	21.75 - 23.65	0.42	40	2	25	12	79	11.101	0.0009	0.53	0.82	GG+
	04	3.80 - 3.90	13	21.40 - 21.55	0.37	60	1	16	5	82	9.671	0.0019	0.74	0.93	GG++
	01	11.10 - 11.15	11	4.10 - 4.55	0.39	26	13	5	15	59	9.497	0.0021	0.66	0.53	GG+
	04	19.90 - 22.80	14	2.35 - 18.15	0.50	29	10	14	44	97	24.708	<0.0001	0.40	0.44	NN+
CAC162	01	1.40 - 5.55	10	19.65 - 19.95	0.54	27	9	8	31	75	23.571	<0.0001	0.48	0.47	NN+
	06	6.30 - 15.10	12	6.20 - 7.75	0.47	9	50	23	15	97	21.637	<0.0001	0.61	0.33	HV
	01	2.35 - 5.45	03	0.00 - 20.65	0.45	28	21	7	42	98	20.628	<0.0001	0.50	0.36	NG-
	04	1.90 - 22.30	14	2.15 - 25.60	0.61	34	9	7	31	81	31.85	<0.0001	0.53	0.51	GG+
	06	17.00 - 19.25	10	19.50 - 20.25	0.43	38	0	22	13	73	22.217	<0.0001	0.52	0.82	GG+
	05	3.50 - 20.40	14	16.50 - 23.00	0.49	26	5	17	32	80	19.799	<0.0001	0.39	0.54	GN-
	07	13.20 - 21.80	08	22.30 - 27.70	0.48	23	7	14	38	82	19.719	<0.0001	0.37	0.45	NN+
	13	22.05 - 25.55	14	0.00 - 3.00	0.47	41	6	15	20	82	18.738	<0.0001	0.57	0.68	GG+
	06	9.65 - 16.00	08	23.60 - 25.55	0.44	22	10	12	37	81	15.892	<0.0001	0.40	0.42	NN+
	01	5.85 - 7.40	04	4.10 - 16.55	0.43	32	6	18	24	80	15.338	<0.0001	0.48	0.63	GN-
	01	7.70 - 11.70	13	24.20 - 25.15	0.42	24	4	22	30	80	15.279	<0.0001	0.35	0.58	GN-
	05	20.85 - 21.75	06	7.10 - 16.05	0.43	20	11	11	40	82	15.242	<0.0001	0.38	0.38	NN++
	05	20.05 - 20.40	07	20.05 - 21.80	0.42	17	7	15	42	81	14.019	0.0002	0.30	0.40	NN+
CAC415	06	6.30 - 6.45	14	8.85 - 9.80	0.49	24	7	18	32	81	13.719	0.0002	0.38	0.52	GN-
	05	8.50 - 17.80	12	0.30 - 0.70	0.40	23	7	18	33	81	13.458	0.0002	0.37	0.51	GN-
	03	15.95 - 19.85	06	17.65 - 20.05	0.41	44	18	5	15	82	13.338	0.0003	0.76	0.60	GG+
	04	19.75 - 19.90	08	2.20 - 2.50	0.40	13	36	22	11	82	13.207	0.0003	0.60	0.43	HV
	05	21.40 - 21.75	08	22.10 - 22.15*	0.39	22	9	16	35	82	12.436	0.0004	0.38	0.46	NN+
	01	3.35 - 3.40	14	20.65 - 22.70	0.40	35	18	5	17	75	12.133	0.0005	0.71	0.53	GG+
	11	24.70 - 25.50	14	25.10 - 25.15*	0.39	32	15	10	24	81	12.119	0.0005	0.58	0.52	GG+
	07	1.00 - 1.45	08	18.50 - 18.55*	0.40	13	30	26	12	81	12.078	0.0005	0.53	0.48	HV
	04	20.05 - 20.10	05	2.40 - 3.50	0.40	33	11	10	19	73	12.024	0.0005	0.60	0.59	GG+
	05	0.50 - 1.20	06	1.45 - 1.50	0.40	23	21	2	18	64	11.726	0.0006	0.69	0.39	NG-
	09	23.70 - 23.85	13	26.20 - 27.70*	0.40	14	25	30	11	80	11.494	0.0007	0.49	0.55	HV

1025