TITLE:

A Novel Simulation Framework for Validation of Ecological Network Inference

RUNNING TITLE:

Network Inference Validation

AUTHORS:

Erik Kusch[1,2,3,*], Anna C. Vinton[4,*]

AFFILIATIONS:

* … Authors have contributed equally to this research

[1]Machine Readable Nature Research Group, Natural History Museum, University of Oslo,
Oslo, Norway, Norway

[2]Center for Biodiversity Dynamics in a Changing World (BIOCHANGE), Department of
Biology, Arhus University, Aarhus, Denmark

[3]Section for Ecoinformatics & Biodiversity, Department of Biology, Arhus University,
Aarhus, Denmark

[4]Department of Biology, University of Oxford, Oxford, UK

CORRESPONDING AUTHOR:

Erik Kusch; erik.kusch@nhm.uio.no

KEYWORDS

Ecological Networks, Network Inference, Demographic Stochasticity, Biological Interaction,

Interaction Simulation, Model Validation

ABSTRACT

1. Understanding how the differential magnitude and sign of ecological interactions vary

across space is vital to assessing ecosystem resilience to biodiversity loss and predict

community assemblies. This necessity for ecological network knowledge and their

labour-intensive sampling requirements has spurred the creation of ecological

network inference methodology. Recent research has identified inconsistencies in

networks inferred using different approaches thus necessitating quantification of

inference performance to facilitate choice of network inference approach.

2. Here we develop a data simulation method to generate data products fit for network

inference and subsequently quantify the validity of two well-established ecological

interaction network inference methods – HMSC and COOCCUR. The simulation

framework we present here can be parameterised using real-world information (e.g.,

biological interactions observed in-situ and bioclimatic niche preferences) thus

representing network inference capabilities in real-world applications. Using this

framework, it is thus possible to evaluate the performance of any ecological network

inference approach.

3. We find that network inference performance scores highlight concerningly low

accuracy of inferred networks as compared to true association networks and suggest

analysis procedures with which to explore network inference reliability with respect

to bioclimatic niche preferences and association strength of association partner-

species.

4. With this study, we provide the groundwork with which to validate and compare

ecological network inference methods, and ultimately vastly increase our ability to

predict species biodiversity across space.

INTRODUCTION

Despite their complexity (e.g., in directionality, sign, and strength), biological interactions

(e.g., competition, mutualism, predation) can be described readily – even among and across

large assemblages of interacting species – via ecological networks. Ecological networks

quantify the direction and sign of interactions (i.e., links) between species (i.e., nodes) as

either positive, negative, or absent while also describing their strength/magnitude (Morales-

Castilla et al., 2015). Real-world biological interactions and their respective networks are

undergoing changes at local and macroecological scales simultaneously, but traditional in-

situ quantifications become intractable across large scales due to sampling effort

requirements (Jordano, 2016). Nevertheless, it is at these macroecological scales that climate

change impacts ought to be evaluated (Lavorel & Garnier, 2002; Rapacciuolo & Blois, 2019;

Wang et al., 2016) and so macroecological networks have to be inferred from proxies. This

has led to the development of interaction inference approaches, most of which utilize spatial

patterns of species co-occurrence (e.g.; Griffith, Veech, & Marsh, 2016; Blonder & Morueta-

Holme, 2017; Damgaard, Ehlers, Ransijn, Schmidt, & Svenning, 2018; Lamonica, Pagel, &

Schurr, 2021; Swain et al., 2021; Tikhonov et al., 2021; Weiss-Lehman et al., 2021; Bimler,

Mayfield, Martyn, & Stouffer, 2022) to identify ecological networks. Due to their

demonstrated usefulness in studying and predicting (1) community assembly patterns

(Freilich et al., 2018; Ovaskainen et al., 2017), (2) ecosystem processes (Dongli et al., 2022)

and services (Keyes et al., 2021) as well as (3) ecosystem stability (Korkmazhan & Dunn,

2022; Thébault & Fontaine, 2010), it is pivotal that networks inferred via these approaches

are accurate representations of realised, biological interactions.

A recent comparison of network inference outcomes has demonstrated that – even when

analysing the same ecosystem – inferred networks are inconsistent across inference

approaches or geographic scales (Kusch et al., 2023). This means that practitioners of

ecological network inference need to carefully consider which network inference approach to

use as inferred networks differ considerably across scales and approaches. To streamline

choice of inference methodology for specific study purposes, criteria of inference approach

applicability have been proposed (Kusch et al., 2023). However, objective measurements of

inference performance to judge quality of ecological network inference remain absent. To

account for the ongoing development of network inference approaches, a generalizable

inference validation framework that can take into account ongoing developments is vital.

Such a framework would allow researchers to judge the quality and accuracy of inferred

ecological networks of any inference approach.

To study performance of ecological network inference approaches, we require a framework

which produces testable networks and data which may be treated as network hypotheses to

compare to inferred networks. While one previous attempt has been made at creating such a

comparison framework (Lavender et al., 2019), this comparison arguably suffers from a lack

of biological realism which has also been identified as a set of pitfalls in a recent overview of

ecological network inference practices (Blanchet et al., 2020). First, limiting data simulations

and network inference to undirected links (i.e., biological associations) fails to represent a

large part of the ecological interaction type spectrum (Harris, 2016). To study the effect of

link directionality on ecological network inference, a network inference performance

assessment framework must therefore be capable of generating synthetic networks containing

directed (i.e., interactions) or undirected (i.e., associations) links. Second, neglecting to

simulate and infer data according to bioclimatic niche preferences may lead to

overestimations of the impact of ecological interactions (Turnbull et al., 2013). Thus, a

simulation framework for network inference validation must explicitly consider bioclimatic

preferences of ecological network-contained species to represent more realistic biodiversity

products for network inference. Third, functional trait expressions may inform ecological

interaction outcomes (Zakharova et al., 2019) but are rarely considered in ecological network

inference practices. To assess their impacts, a data simulation framework should incorporate

their adjustments to interaction/association outcomes. Fourth, spatial biodiversity products

beyond co-occurrence records (i.e., abundance or performance matrices) have been

demonstrated to lead to vastly different inference outcomes when compared to co-abundance-

or co-performance-based approaches (Kusch et al., 2023). Conclusively, to quantify the

accuracy-changes in network inference owing to differences in spatial input types, the

simulation framework must be capable of generating biodiversity products beyond co-

occurrence. Finally, to mirror real-world data as closely as possible, such a simulation

framework ought to be capable of incorporating considerations of population dynamics such

as density dependence and dispersal as these factors have been demonstrated to shape spatial

patterns of biodiversity alongside biological interactions (Saupe et al., 2015).

Here we present a novel simulation-based toolbox for ecological network inference

performance assessments overcoming the above-mentioned limitations. This framework

centres on a spatially explicit, individual-based simulation for the generation of data products

ready for network inference. With this framework, we address the impact of bioclimatic

niches and biological interactions as well as density dependence and dispersal dynamics in

shaping species distributions through time and space. This data generation framework can be

parameterised either with real-world network and species data (e.g., species-specific

bioclimatic niches and dispersal potential), or randomly generated network and species data

(for which we provide supporting functionality). While the former will allow for precise

quantification of network inference performance given specific study systems, the latter

enables a more general assessment of network inference reliability across a range of systems

and species. Because of the flexibility of the data input and robustness of the mathematical

simulation approach, this framework is generalisable across study ecosystems and species

pools. Moreover, this simulation framework provides a flexible toolbox with which

additional considerations beyond the ones presented here may be addressed (e.g., different

types of species interactions, environmental change impacts). Using this simulation

framework, we subsequently carry out ecological network inference using two well-

established approaches of ecological network inference – HMSC (Ovaskainen et al., 2017)

and COOCCUR (Veech, 2013). While the former allows us to assess the performance of

network inference while accounting for bioclimatic preferences and spatial products beyond

simple co-occurrence records, the latter inference approach is blind to climatological

gradients over space and restricted in input data to co-occurrence matrices. We expect that

network inference will more accurately identify true pairwise associations when bioclimatic

preferences are accounted for thus making HMSC inference more accurate. Consequently,

the framework for data generation and process of network inference performance assessments

we present here represent an advancement in our understanding of performance and

applicability of ecological interaction inference methodology.

MATERIAL & METHODS

ECOLOGICAL NETWORK MATRICES

When encoding ecological networks, it is common practice to represent nodes (i.e., species)

and their links (i.e., interactions or associations) as matrices (Ávila-Thieme et al., 2023;

Opedal & Hegland, 2020). Within these matrices, columns and rows represent species-

identities while cells contain the link weight (interaction/association strength) between

corresponding species pairs. Network inference approaches can detect network matrices as

either representing directed (Bimler et al., 2022) or undirected networks (Morueta-Holme et

al., 2016; Tikhonov et al., 2020; Veech, 2013).

Directedness of ecological network links is identified through arrangement of values within

the corresponding network matrix. Undirected, association matrix entries are contained either

in the upper or lower triangle of the species-by-species matrix (i.e., every species in an

association pair affects the partner equally). Directed, interaction matrices, on the other hand,

contain the interaction strength and sign of an actor species (stored in columns) onto a subject

species (stored in rows) and vice versa. While undirected network matrices identify

biological associations (e.g., probability of co-occurrence of two species), directed network

matrices allow for the identification of more complex biological processes like predator-prey

interactions. Conclusively, for our framework of network inference validation, we have

implemented support for interaction and association matrices.

While our simulation framework can be parameterised with ecological network matrices as

observed in nature, we provide functionality for creation of random network interaction or

association matrices. Optional arguments of this helper function control maximum value of

association/interaction strengths and sparsity of the matrix (i.e., how many of all possible

values should be exactly 0).

SIMULATION FRAMEWORK

In order to generate data to assess network inference performance, we develop a spatially

explicit birth-death model. Here, we iteratively simulate birth and death events of individuals

belonging to interacting species (as defined via a network matrix) across a two-dimensional

(x and y), continuous landscape.

THE MODEL

Individuals in this model have constant birth rates (b_0), while their death rate over time

($d(t)$) is determined by their (1) population density and carrying capacity (P), (2)

environmental maladaptation (E), and (3) interaction effects of neighbouring individuals (Ω).

$$d_i(t) = d_0 + P_i \times E_i - \Omega_i \qquad (1)$$

167   Where $d_i(t)$ is the dynamic death rate for individual i at time t and d_0 denotes the

168   background death rate.

Population Dynamics

170   To account for logistic growth of individual populations, our simulation evaluates the impact

171   of density and carrying capacity (P_i) within the dynamic death rate calculation (equation 1).

172   Here, P_i is determined through species-specific (S) population sizes and carrying capacities

173   thus ensuring $P_i = P_S$ for all individuals i belonging to the same species S:

$$P_S(t) = \frac{N_S(t) \times (b_0 - d_0)}{K_S} \qquad (2)$$

174   Where d_0 and b_0 represents the background rate of death and birth respectively. K_S and $N_S(t)$

175   represent the carrying capacity and population size at time t for an individual of species S

176   respectively. The higher the value of P_S, the higher the dynamic death rate of individual of

177   species S.

Impact of Environment and Trait

179   The maladaptation of an individual with trait μ at location (x, y), $E_\mu(x, y)$ is related to the

180   difference between the individual's trait value μ and the optimal trait value, or the

181   environment at its location ($u_0(x, y)$):

$$E_\mu(x, y) = M_{\sigma_E}(\mu_i - u_0(x, y)) \qquad (3)$$

182   The larger $E_\mu(x, y)$, the worse the individual is adapted to its local abiotic conditions and the

183   higher its dynamic death rate will be. $M_{\sigma_E}(z)$ is a Gaussian function that determines the

184   effect of being maladapted in the current environment. This maladaptedness is weighted by

185   the model parameter σ_E which controls the severity by which the difference between an

186   individual's trait and their environment affects the dynamic death rate:

$$M_{\sigma_E}(z) = e^{\left(\frac{z}{\sigma_E}\right)^2} \qquad (4)$$

187   The optimal phenotype, or $u_0(x, y)$, is determined via a simulation-user defined function. For

188   the study presented here, we assigned a linear function of the one-dimensional environmental

189   gradient (see Figure S1 for a visual representation) but note that this framework can be

190   adapted to more complex landscapes:

$$u_0(x, y) = x \qquad (5)$$

In evaluating environmental maladaptedness as an outcome only of the x dimension our

simulations enable the formation of alternative, competing community assemblies where

species of similar environmental preference, but with competitive association between them

may both persevere.

<u>Species Interactions/Associations</u>

The strength of the impact of species interactions/associations an individual experiences is

given by the number and species identity of individuals within a defined area around each

individual. These individual interactions vary between species according to network matrices.

Our framework quantifies the effect of interactions Ω_i for a user-defined distance around

each individual i as follows. The interaction coefficient, $l_{i,j}$ denotes the impact of species j

on species i drawn from the $i \, X \, j$ network matrix. To account for effects of neighbouring

individuals by their abundance, the interaction coefficients are multiplied by the respective

abundances of the j-th species in the user-defined area surrounding the i-th individual (a_j).

The resulting sum of abundance-weighted interaction coefficients is subsequently

standardised by dividing its value by the number of individuals in the i-neighbourhood (a_j):

$$\Omega_i = \frac{\sum_1^J l_{i,j} * a_j}{\sum_1^J a_j} \tag{6}$$

Positive values of Ω_i reflect beneficial interaction effects and thus decrease the dynamic

death rate while negative Ω_i values increase the dynamic death rate.

STOCHASTIC SIMULATION ALGORITHM

Our simulation framework tracks birth and death events of individuals belonging to

interacting species through time and space as directed by Gillespie, 1977. At each time step t,

the simulation framework computes the dynamic death rate $d_i(t)$ for each individual as

described above. A birth or death event is selected to occur at random (demographic

stochasticity), weighted by the individual's static birth and dynamic death rates.

If a death event is selected, the chosen individual is removed from the population. When a

birth event occurs, the offspring is a clone of the parent individual in their trait value μ_i.

Thus, reproduction in our framework is asexual. Natal dispersal is the only form of

movement in our framework with offspring location coordinates (x, y) given as the parental

x and y location, with the addition of a dispersal magnitude drawn from a normal distribution

with mean equal to 0 and a user-defined standard deviation. The boundary conditions of the

continuous landscape in either dimension are defined by the simulation user. Any dispersal

attempt beyond these limits will result in dispersal to the limit.

Next, the time until the next event is determined by randomly sampling from an exponential

distribution with a mean of the inverse of the sum of all possible events. This is done to

ensure that simulation time represents realistic passage of time in biological systems (i.e.,

more individual birth and/or death events happen in shorter time in larger populations).

Finally, the simulation starts over with the updated pool of individuals. This framework is

similarly developed in Vinton & Vasseur, 2020.

Lastly, simulations are initialised with a user-defined or randomly generated (1) network

matrix, (2) table of initial individuals containing spatial location and trait values for each

individual, (3) a vector of carrying capacities for each species, and (4) a function which

defines how coordinates relate to environmental values/optimal phenotype.

IMPLEMENTATION IN THE R ENVIRONMENT

We coded and executed our simulation framework in R version 4.0.2. (R Core Team, 2021).

The simulation inputs (see below) parameterize the simulation framework described above.

To ensure reproducibility, we set a seed for random computation processes. For faster

computation of several simulation instances at once we enable parallel processing of several

simulation runs using the doParallel (Microsoft & Weston, 2020a) and foreach (Microsoft &

Weston, 2020b) libraries in R.

Network Matrices

Using the randcorr R package (Schmidt & Makalic, 2018), we created supporting

functionality with which we establish a set of 500 random association matrices containing 20

species with a maximum strength of association of 1 and a matrix-sparsity of 50%.

Initial Individuals & Carrying Capacities

Using each random network we have created, additional functionality of our simulation

framework creates a user-defined number of individuals (either in total or for each species)

and allocates trait values to each individual which are drawn from a random distribution

around a randomly selected trait mean for each species given a user-defined standard

deviation for these random trait distributions. Additionally, individuals are placed randomly

within a user-defined spatial extent in x and y dimensions. Lastly, carrying capacities for

each species are chosen from a uniform distribution with user-defined boundaries.

For the analyses presented here, we chose random trait mean values between 0 and 10 for

each species and set carrying capacities of 200 for all species. We subsequently initialised

each simulation with 500 individuals per species in random locations ranging between 0 and

10 in both x and y dimensions. Individual trait values were drawn from random distributions

around individual-species-specific trait mean values and a standard deviation of one.

<u>Simulation Parameters</u>

To use the initialising data (i.e., network matrices, data tables of individuals and carrying

capacities) to generate spatially explicit products ready for network inference, we executed

our simulation framework with the following parameters. Background rates of birth (b_0) and

death (d_0) were set to 0.4 and 0.6 respectively. To calculate local environmental effects on

the dynamic death rate, we defined σ_E as 1 and supplied a linear function which translates x-

values directly into environmental values/optimal phenotypes (see equation 5). When

calculating association effects on the dynamic death rate, we defined the size of bounding

box around each individual by which association effects and respective species abundances

are calculated as 0.5 resulting in a quadratic extent where each side measures 1 unit in x and

y dimensions. Dispersal distance of individuals originating from a birth event were randomly

drawn for each dimension (x and y), for each birth event, from a normal distribution centred

on 0 with a standard deviation of 0.2. Finally, we ran each simulation until simulation time 20

was reached, recording individual locations and trait values in time intervals of 0.1 (see

Figure S2 for a visual representation of how these individual-time-step information can be

used for studying time-series of abundance which a number of network inference approaches

may use; e.g.; Damgaard & Weiner, 2021).

<u>INFERENCE APPROACHES</u>

We carried out ecological network inference using two approaches – HMSC and COOCCUR.

We chose these two approaches for their unique placements along a co-occurrence to

performance spectrum which characterises network inference input data types and which has

been demonstrated to affect network inference outcomes (Kusch et al., 2023). While

COOCCUR incorporates co-occurrence data, HMSC can consider species performance

implicitly through abundance products. Additionally, while COOCCUR is blind to

environmental gradients and preferences, HMSC aims to identify these as part of its inference

approach. Lastly, COOCCUR has also been part of a previous simulation-based assessment

(Lavender et al., 2019).

<u>COOCCUR</u>

COOCCUR is a probabilistic framework first introduced by Veech, 2013 which we

implemented in our analyses via the cooccur R package (Griffith et al., 2016). COOCCUR

identifies species associations in sign and strength through comparison of analytical estimates

of species co-occurrence probabilities to purely random species distributions and resulting

co-occurrence patterns. Probabilities of observed co-occurrence rates being higher or lower

than expected at random are then used as p-values to denote statistical significance of

identified associations.

<u>HMSC</u>

We executed the HMSC method (Ovaskainen et al., 2017) through the hmsc R package

(Tikhonov et al., 2021). HMSC is designed as a hierarchical species distribution modelling

framework which aims to explain inputs of site-by-species matrices through environmental

niches of species and site-specific environmental characteristics, functional traits of species,

and phylogenetic relatedness. Residual covariance of species is then treated as association

inference (Ovaskainen et al., 2017). The statistical significance of these inferred associations

is available via the credible intervals of their posterior distributions (Opedal & Hegland,

2020). For all subsequent analyses, we only retained those associations per HMSC model

which were identified with 95% credible interval not overlapping 0.

DATA PREPARATION FOR NETWORK INFERENCE

To prepare the simulation outputs for network inference, for each final spatial output, we

extracted trait means for each species which correspond to species- but not individual-

specific optimal location along the x dimension. We then identified species abundances

across the continuous two-dimensional landscape in 100 evenly spaced grids (i.e., 10

segments along each axis resulting in grid cells with a length of 1 on either side). This

information was subsequently stored as a species-by-site abundance matrix (see Figure S3 for

a visual representation). Lastly, we noted the coordinates of each grid cell and assigned a

unique ID to each grid cell. This product was used as a study design identifier and to

establish random effect levels with respect to cell ID in HMSC models. For COOCCUR

analyses, the species-by-site abundance matrix was transformed into a matrix of

presence/absence.

INTERACTION/ASSOCIATION REALISATION

As a product of the random generation of environmental preferences and association motifs

per species, some of the simulated associations may never be realised due to association

partners not being able to survive in each other's presence due to their environmental

preferences. As such, the simulated ecological networks represent "potential" association

patterns which network inference approaches cannot reasonably be expected to identify.

Therefore, we distinguish between true, potential ecological networks (i.e., the randomly

generated input networks for our population dynamics simulations) and true, realised

ecological networks. The latter are established by assigning as absent (i.e., magnitude of 0)

all associations whose association partners exhibit environmental niche preferences (i.e.,

observed trait values at the final simulation step) further apart than the radius of the

association effect bounding box (i.e., 0.5) plus the niche breadth parameter (σ_E; i.e., 2.5). See

Figure 1 for a visual representation of how this process works.

PERFORMANCE ASSESSMENT

To facilitate quantification of network inference accuracy, we used the concept of network

beta diversity which partitions ecological network dissimilarity according to differences in

links between nodes shared by two networks and links which differ because of a difference in

node set (i.e., species) (Poisot et al., 2012). Since species may go extinct throughout our

simulation runs either due to maladaptation to the environment or competitive exclusion in

their preferred environment, we focus on the dissimilarity in links between nodes shared by

the true, realised and the inferred networks. To simplify this comparison of true, realised (i.e.,

initialising) and inferred networks, we transformed all network links into a binary

representation of -1 for negative associations and +1 for positive associations thus omitting

link strength from our analysis of network dissimilarities. Subsequently, we use the betalink

R package (Poisot et al., 2012) to quantify network dissimilarity components for each set of

known network as well as informed and naïve inferred network. Ultimately, we use these

dissimilarity scores to assess network inference performance at a whole-network level.

Additionally, we calculate rates of (1) correct identification of positive, negative, and absent

associations, (2) incorrect identification of positive, negative, and absent associations, and (3)

failure of identification of positive, negative and absent associations.

Finally, we model accuracy of network inference approaches with respect to identity

(present/absent) and sign (positive or negative) of associations with respect to true

association strength and environmental niche preferences of associations partners. We do so

to create an understanding of how erroneous results of network inference arise and what kind

of inferred associations should be treated more carefully than others. To do this, we establish

logistic regression models using the brms R package (Bürkner, 2021) which model correct

and incorrect identification of a specific association type (negative, positive, absent) as an

outcome of the difference in trait expression per pair of association partners as well as the

true strength of associations.

RESULTS

Our simulations yielded 500 distinct random association networks containing 20 species each

with random mean trait/phenotype expressions per species per simulation product. These are

accompanied by their corresponding 500 lists of data tables containing individual IDs,

phenotypes/traits and locational information which we use for network inference. Due to the

complexity of these outputs, we focus on a reduced conceptual simulation run to demonstrate

the functionality of the data generation simulation framework before we explore validation

efforts of network inference.

SIMULATION OUTPUTS

To demonstrate our simulation framework, we established a conceptual random association

matrix containing five species (with a maximum association strength of 10 and a network

sparsity of 0.1). The simulation was then executed with $\sigma_E = 2.5$. See Figure 1 for a visual

representation of the conceptual true potential association network matrix as well as the

process by which we establish an understanding of the true realised version of this network.

The random association network matrix is the used to calculate the interaction effect on

dynamic death rates at each simulation step and shapes the final spatial product of species

abundances across the artificial continuous landscape as can be seen in Figure 2.

FIGURE 1 - CONCEPTUAL RANDOM NETWORK MATRIX AND REALISATION OF ASSOCIATIONS. A) A true potential association matrix for five distinct species. Note that maximum association

strength was set to 10 and network sparsity to 0.1 resulting in 1 out of the possible ten associations having a strength of exactly 0 while all other associations may range between -10 and +10. B) The pairwise difference

in environmental difference as identified through mean and standard deviation of species-specific trait values at the final simulation step (see vertical lines in right-hand panel of Figure 2). Black squares identify

associations which can be realised due to shared environmental preferences. C) The true, realised association matrix corresponding to A). All associations which cannot be realised have been set to a magnitude of 0

(i.e., absent associations).

375   If desired, our simulation framework also returns products containing all extant individuals as

376   well as their traits and locational information at pre-defined simulation-time intervals. This

377   enables the analyses of time-series of species abundances as shown in Figure S2.

378
FIGURE 2 - CONCEPTUAL SPATIAL PRODUCTS. Simulations are initialised (left panel) with individuals belonging to interacting

species which are placed randomly across the continuous landscape. This placement is irrespective of trait expressions specific to each

species (coloured lines showing the mean trait expression per species in either panel). The simulation terminates at a user-defined

simulation-time and returns the final output of individuals across the landscape (right panel) which have shifted with respect to (1)

population dynamics, (2) individual environmental maladaptation, and (3) interaction effects.

<u>NETWORK INFERENCE</u>

The network inference approaches we use here analyse species-by-site matrices of spatial

biodiversity products (i.e., abundance or presence/absence) in evenly spaced sites of one unit

length in   and   dimension respectively. A conceptual overview of how these products are

generated is provided in Figure S3.

INFERRED NETWORKS

Using COOCCUR and HMSC, we infer ecological networks and retain statistically

significant associations as realised links whose strength we set to -1 and +1 with respect to

the sign of the inferred association. Potential associations which were not identified

statistically significantly, we set to 0 to denote absence of an association. The inferred

networks are notably different to the true, original networks which we used to drive our

simulations (Figure 3).

396
397 **FIGURE 3 - CONCEPTUAL INFERRED NETWORKS.** Inferred networks of statistically significant pairwise associations (whose

398 association strengths have been set to -1 and +1 respective to association inference sign) show great dissimilarity to original network (Figure

399 1, C) which affected the simulation framework. Black squares identify correctly identified positive, negative, and absent associations.

PERFORMANCE OF NETWORK INFERENCE METHODS

Using the beta diversity concept of assessing Network dissimilarities, we find that, within our

analyses both network inference approaches perform equally when contrasting inferred and

true realised networks, but COOCCUR outperforms HMSC when assessing true, potential

networks (Figure 4). We focus on dissimilarities in links between shared species to mitigate

any biasing effects of species extinction throughout the simulation run which effectively

leads to smaller node sets for inferred networks than original networks. However, overall

network dissimilarity remains high for both network inference methods we assess here.

Despite their similar performance at correctly identifying true realised associations (Figure 4,

right panel), individual networks rendered by HMSC and COCCUR per simulation run are

markedly dissimilar (Figure S4) highlighting that the network inference approaches identify

different ecological networks which are equally dissimilar to the actual associations affecting

the individuals across simulations.

FIGURE 4 - NETWORK INFERENCE ACCURACY AS ASSESSED THROUGH BETA-DIVERSITY NETWORK

DISSIMILARITY. When analysing dissimilarity scores of links of shared species between inferred networks and the original, true

association networks for each inference approach across all simulation runs, we find that COOCCUR produces more accurate inference than

HMSC at true potential network comparisons due to its lower OS (link dissimilarity) score which denotes dissimilarity in link identity and

sign between the vertices present in the known and the inferred networks.

Further deconstruction of network dissimilarities into rates of true positive, negative, and

absent as well as false positive, negative, and absent associations identified in each inferred

network (see Figure 5 for comparison to true realised network; Figure S5 for comparison to

true potential network) reveals that (1) positive associations inferred by COOCCUR are more

likely to be true realised positive associations, but HMSC tends to miss less positive

associations in its inference, (2) neither HMSC nor COOCCUR perform well at correctly

identifying negative associations, but HMSC again missing less negative associations in its

inference, and (3) COOCCUR, on average, performing better at correctly identifying absence

of   pairwise   associations.

FIGURE 5 – RATES OF TRUE REALISED ASSOCIATION DETECTION. Decomposition of overall network dissimilarities expressed as density distributions reveal similar and low performance of the two

network inference approaches for our simulations. Two-letter codes denote type of error rate: TP…True positive, FP…False positive, MP…Missed positive (a positive association that was failed to be identified),

TN…True negative, FN… False Negative, MN… Missed negative, TA…True absent, FA…False absent, and MA…Missed absent.

To further understand network inference approach behaviour, we executed logistic regression

models using the difference in trait expression per pair of association partners as well as the

strength of associations to predict the probability of assigning correctly either positive,

negative, or absent associations to species-pairs for COOCCUR and HMSC. See Figure 6 for

a visualisation of these model outcomes for performance of inferring true realised networks.

Model outcomes for performance of true potential network inference is presented in Figure

S6.

We find that HMSC and COOCCUR are similarly competent at correctly identifying realised

positive associations so long as the association partners closely share environmental

preferences and the true association is strong (Figure 6A). When assessing correct

identification of realised negative associations, however, COOCCUR failed to render

correctly negative, realised associations to such a degree (see also Figure 5, second row) that

logistic models were not able to reliably establish under which conditions COOCCUR

inference of negative associations may be reliable. For HMSC, on the other hand, correct

identification of negative associations becomes more likely as true realised negative

associations increase in magnitude and species-niche preferences become more dissimilar.

Lastly, absence of associations are correctly identified by either network inference approach

predominantly when species-niche preferences of potential association partners are

dissimilar.

FIGURE 6 - PROBABILITY OF CORRECT IDENTIFICATION OF TRUE REALISED ASSOCIATIONS. Logistic regression

coefficients (left column) show clear effects of association magnitude, environmental difference in species niches as well as the interaction

of these two which translate to distinct probabilities of correctly identifying a pairwise associations (right column). Effects correct

identification likelihood of association magnitude and environmental niche difference between association partners depends on sign of the

underlying association. Note that COOCCUR identified so few negative associations correctly that logistic models did not converge well.

DISCUSSION

We find that overall accuracy of network inference using either approach is low when compared to known and inferred ecological networks. This highlights that current ecological network inference approaches fail at identifying pairwise ecological associations. This concern is exacerbated when considering that our performance analyses was carried out using networks which have been transformed into association networks with link strength -1, 0, and +1 which is an oversimplification of the potential link strength spectrum which has been demonstrated to shape ecosystem stability and community diversity (Kusch & Ordonez, 2023; Ushio, 2022). However, we do not claim our network inference performance scores to be necessarily representative of real-world applications, but merely that they remain accurate for our specific simulation parameters. For example, increasing σ_E (i.e., making the environmental constraint more forgiving) or increasing association magnitudes would likely increase the performance of either network inference approach as the effects of associations will become greater in comparison to environmental preference and location-specific environmental mismatches. Additionally, we suspect that increasing the distance which defines the i-neighbourhood may make identification of associations easier for HMSC as interaction effects will have greater reach, thus affecting each individual site in the site-by-species matrix used for network inference more than before. Alternatively, restricting interactive effects not to the i-neighbourhood but strictly to the grid cells which define sites should improve performance of either network inference approach and be more representative of study setups with distinct sites (López-Segoviano et al., 2021) rather than continuous megaplot designs (Lutz et al., 2012). We expect that using our simulation framework in conjunction with real-world association networks and bioclimatic niche preferences and environmental suitability constraints will yield a more realistic overview of

network inference performances. However, such a study will require extensive data collection

to allow more precise parameterisation of the simulation framework we are presenting here.

Irrespective of the overall low accuracy of network inference, our analyses reveal network

inference performances of COOCCUR and HMSC which seemingly disprove our hypothesis

of HMSC outperforming COOCCUR due to its incorporation of environmental information

and an abundance input data type rather than one of presence/absence. Our analyses of

association-type (i.e., positive, negative, or absent) error and detection rates highlight that

COOCCUR and HMSC differ predominantly in their respective capability of correctly

identifying associations of different signs suggesting the two may be used in an ensemble-

like inference approach for more accurate network inference than could be rendered by using

either in isolation.

While both network inference approaches successfully identify positive pairwise associations

so long as the associations partner-species share environmental preferences (Figure),

COOCCUR fails to correctly identify negative associations (Figure 4, TN panel) with HMSC

likewise struggling to correctly identify negative associations. It stands to reason that

COOCCUR's performance is driven by environmental differences rather than association

strength itself thus assigning negative associations to species pairs who do not share an

environment due to differences in environmental preferences rather than competitive

interactions (i.e., negative associations). This will lead to assignment of negative associations

to species-pairs whose true associations (negative or not) may never be realised due to their

dissimilar environmental preferences. The analysis of detection probability of true potential

negative associations (Figure S6B) supports this hypothesis. Consequently, this highlights the

fallacy of neglecting environmental gradients in network inference with positive and negative

associations assigned incorrectly because of differences in environmental preferences rather

than actual interactions (Blanchet et al., 2020).

507 Conclusively, we find that network inference accuracy is largely driven by differences in

508 environmental preferences of association partner-species. While the overall low network

509 inference performance scores we have identified via network dissimilarity scores (Figure 4)

510 may indicate that ecological network inference is performing below expectations or necessary

511 thresholds of accuracy, we argue that these scores highlight an important distinction that has

512 yet to be fully appreciated and explored in the study of network inferences:

513 association/interaction sign. Instead of selecting one "best-performing" network inference

514 approach, we believe that it may be more prudent to quantify capabilities of network

515 inference approaches to identify association types with respect to association sign,

516 magnitude, and differences in environmental preferences of association partner-species as we

517 have done with our logistic models in Figure . Doing so will create ranges of environmental

518 and biological constellations within which individual network inference approaches can be

519 treated as reliable. Ultimately, we believe that extensive real-world parameterisation of our

520 simulation framework will be required to arrive at conclusive assessments of network

521 inference performance given such considerations of realised and inferred

522 associations/interactions.

523 While our simulation framework represents a major advancement of our capability of

524 identifying network inference performance and validate inferred networks, we also realise

525 that the simulation framework we present here may be regarded as an oversimplification of

526 natural processes. For examples, non-natal movement, which is absent from our framework,

527 has been identified as a confounding effect to interactions studies of animals (Blanchet et al.,

528 2020; Matthiopoulos et al., 2015). Subsequently, we expect the current version of our

529 simulation framework to be most useful to studies of plant-plant interactions/associations or

530 those of other sessile organisms. Furthermore, our simulation framework only includes one

531 functional trait per individual despite organisms inhabiting a multi-dimensional space of

functional trait expressions (Blonder, 2018). Our framework neglects the impacts of

functional trait expression on association/interaction outcomes which have been

demonstrated particularly among tree species (Kunstler et al., 2012). Thus, we believe that

our simulation may be further enhanced by the inclusion of additional functional trait metrics

which determine the dynamical death rate through the interaction/association coefficient (Ω).

Additionally, many species reproduce sexually rather than clonally and there may thus exist

considerable variation in such functional traits between parents and offspring. Despite these

avenues for future development, our simulation framework remains the most realistic for

network inference data generation purposes particularly when compared to previous attempts

at quantifying pairwise association inference performance (Lavender et al., 2019).

CONCLUSION

Network inference approaches have been developed to address the labour intensity of

ecological network research which requires impractical sampling efforts particularly at large

geographical scales or for large species pools. Despite the large number and diversity in

approaches of network inference frameworks, no general framework for network inference

validation or performance quantification has been established. Here, we have presented an

easy-to-use, mathematically robust simulation framework which can be used for data

generation with which to carry out network inference validation. While we find overall low

accuracy of network inference, we argue that this may be further remedied by parametrising

our simulation with more realistic real-world data and exploring performance of network

inference through a lens potential and realised interactions/associations. Thus, our simulation

framework for network inference validation represents an important addition to the toolbox

of network inference method development which fills a crucial knowledge gap of network

inference performance assessments.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGMENTS

This research was supported by the Aarhus University Research Foundation Start-up Grant

(grant no. AUFF-2018-7-8) to Alejandro Ordonez.

AUTHORS' CONTRIBUTIONS

E.K. conceptualised the study. A.V. developed the simulation framework used in this study.

E.K developed the network inference and network validation framework used in this study.

E.K. created all R scripts necessary for analyses. E.K. led the analysis and manuscript

writing. All authors contributed critically to the drafts and gave final approval for publication.

DATA AVAILABILITY

All code required for data generation, network inference, and inference performance

quantification is available at https://github.com/ErikKusch/Scaleable-Inference-of-

Ecological-Networks (please note that this is currently a private repository until submission

to a relevant journal so that the simulation framework specification remains protected). We

are ready to submit code and data to dryad upon acceptance of the manuscript.

REFERENCES

Ávila⬜Thieme, M. I., Kusch, E., Corcoran, D., Castillo, S. P., Valdovinos, F. S., Navarrete,
S. A., & Marquet, P. A. (2023). NetworkExtinction: An R package to simulate
extinction propagation and rewiring potential in ecological networks. *Methods in*
Ecology and Evolution, 2020.10.17.305391. https://doi.org/10.1111/2041-210X.14126
Bimler, M. D., Mayfield, M. M., Martyn, T. E., & Stouffer, D. B. (2022). *Estimating*
interaction   matrices   from   performance   data   for   diverse   systems.
https://doi.org/doi.org/10.1101/2022.03.28.486154
Blanchet, F. G., Cazelles, K., & Gravel, D. (2020). Co⬜occurrence is not evidence of
ecological interactions. *Ecology Letters*, ele.13525. https://doi.org/10.1111/ele.13525
Blonder, B. (2018). Hypervolume concepts in niche- and trait-based ecology. *Ecography*,
41(9), 1441–1455. https://doi.org/10.1111/ecog.03187
Blonder, B., & Morueta-Holme, N. (2017). *netassoc: Inference of Species Associations from*
Co-Occurrence Data (p. 1).
Bürkner, P. C. (2021). Bayesian Item Response Modeling in R with brms and Stan. *Journal*
of Statistical Software, *100*(5). https://doi.org/10.18637/JSS.V100.I05
Damgaard, C., Ehlers, B. K., Ransijn, J. C. G., Schmidt, I. K., & Svenning, J. C. (2018).
Insights on plant interaction between dominating species from patterns of plant
association: expected covariance of pin-point cover measurements of two species.
Environmental   and   Ecological   Statistics,   *25*(2),   221–235.
https://doi.org/10.1007/S10651-018-0400-Z/FIGURES/6
Damgaard, C., & Weiner, J. (2021). The need for alternative plant species interaction models.
Journal of Plant Ecology, *14*(5), 771–780. https://doi.org/10.1093/jpe/rtab030
Dongli, D., Chengxing, W., Yuchen, Z., Changchun, L., & Ning, W. (2022). Coexistence
mechanism of alien species and local ecosystem based on network dimensionality
reduction   method.   *Chaos,   Solitons   &   Fractals*,   *159*,   112077.
https://doi.org/10.1016/j.chaos.2022.112077
Freilich, M. A., Wieters, E., Broitman, B. R., Marquet, P. A., & Navarrete, S. A. (2018).
Species co-occurrence networks: Can they reveal trophic and non-trophic interactions in
ecological communities? *Ecology*, *99*(3), 690–699. https://doi.org/10.1002/ecy.2142
Gillespie, D. T. (1977). Exact stochastic simulation of coupled chemical reactions. *The*
Journal of Physical Chemistry, *81*(25), 2340–2361. https://doi.org/10.1021/j100540a008
Griffith, D. M., Veech, J. A., & Marsh, C. J. (2016). cooccur: Probabilistic species co-
occurrence   analysis   in   R.   *Journal   of   Statistical   Software*,   *69*(1),   1–17.
https://doi.org/10.18637/jss.v069.c02
Harris, D. J. (2016). Inferring species interactions from co-occurrence data with Markov
networks. *Ecology*, *97*(12), 3308–3314. https://doi.org/10.1002/ecy.1605
Jordano, P. (2016). Sampling networks of ecological interactions. *Functional Ecology*,
30(12), 1883–1893. https://doi.org/10.1111/1365-2435.12763
Keyes, A. A., McLaughlin, J. P., Barner, A. K., & Dee, L. E. (2021). An ecological network
approach to predict ecosystem service vulnerability to species losses. *Nature*
Communications, *12*(1), 1586. https://doi.org/10.1038/s41467-021-21824-x
Korkmazhan, E., & Dunn, A. R. (2022). High-order correlations in species interactions lead
to complex diversity-stability relationships for ecosystems. *Physical Review E*, *105*(1),
1–10. https://doi.org/10.1103/physreve.105.014406
Kunstler, G., Lavergne, S., Courbaud, B., Thuiller, W., Vieilledent, G., Zimmermann, N. E.,
Kattge, J., & Coomes, D. A. (2012). Competitive interactions between forest trees are
driven by species' trait hierarchy, not phylogenetic or functional similarity: Implications
for   forest   community   assembly.   *Ecology   Letters*,   *15*(8),   831–840.

https://doi.org/10.1111/j.1461-0248.2012.01803.x

Kusch, E., Bimler, M. D., Lutz, J. A., & Ordonez, A. (2023). *Ecological network inference is not consistent across scales or approaches approaches*. https://doi.org/10.1101/2023.07.13.548816

Kusch, E., & Ordonez, A. (2023). *Ecological Network Resilience & Extinction Proxies - Updating Projections of Ecological Networks*. https://doi.org/10.1101/2023.08.02.551629

Lamonica, D., Pagel, J., & Schurr, F. M. (2021). Predicting the dynamics of establishing tree populations: a framework for statistical inference and lessons for data collection. *Methods in Ecology and Evolution*, 2041-210X.13656. https://doi.org/10.1111/2041-210x.13656

Lavender, T. M., Schamp, B. S., Arnott, S. E., & Rusak, J. A. (2019). A comparative evaluation of five common pairwise tests of species association. *Ecology*, *100*(4). https://doi.org/10.1002/ecy.2640

Lavorel, S., & Garnier, E. (2002). Predicting changes in community composition and ecosystem functioning from plant traits: revisting the Holy Grail. *Functional Ecology*, *16*(Essay Review), 545–556. https://doi.org/Doi 10.1046/J.1365-2435.2002.00664.X

López-Segoviano, G., Arenas-Navarro, M., Villa-Galaviz, E., Díaz-Infante, S., & Arizmendi, M. del C. (2021). Hummingbird-plant interactions along an altitudinal gradient in northwestern Mexico. *Acta Oecologica*, *112*, 103762. https://doi.org/10.1016/j.actao.2021.103762

Lutz, J. A., Larson, A. J., Swanson, M. E., & Freund, J. A. (2012). Ecological importance of large-diameter trees in a temperate mixed-conifer forest. *PLoS ONE*, *7*(5). https://doi.org/10.1371/journal.pone.0036131

Matthiopoulos, J., Fieberg, J., Aarts, G., Beyer, H. L., Morales, J. M., & Haydon, D. T. (2015). Establishing the link between habitat selection and animal population dynamics. *Ecological Monographs*, *85*(3), 413–436. https://doi.org/10.1890/14-2244.1

Microsoft, & Weston, S. (2020a). *doParallel: Foreach Parallel*.

Microsoft, & Weston, S. (2020b). *foreach: Provides Foreach Looping Construct*.

Morales-Castilla, I., Matias, M. G., Gravel, D., & Araújo, M. B. (2015). Inferring biotic interactions from proxies. *Trends in Ecology and Evolution*, *30*(6), 347–356. https://doi.org/10.1016/j.tree.2015.03.014

Morueta-Holme, N., Blonder, B., Sandel, B., McGill, B. J., Peet, R. K., Ott, J. E., Violle, C., Enquist, B. J., Jørgensen, P. M., & Svenning, J. C. (2016). A network approach for inferring species associations from co-occurrence data. *Ecography*, *39*(12), 1139–1150. https://doi.org/10.1111/ecog.01892

Opedal, Ø. H., & Hegland, S. J. (2020). Using hierarchical joint models to study reproductive interactions in plant communities. *Journal of Ecology*, *108*(2), 485–495. https://doi.org/10.1111/1365-2745.13301

Ovaskainen, O., Tikhonov, G., Norberg, A., Guillaume Blanchet, F., Duan, L., Dunson, D., Roslin, T., & Abrego, N. (2017). How to make more out of community data? A conceptual framework and its implementation as models and software. *Ecology Letters*, *20*(5), 561–576. https://doi.org/10.1111/ele.12757

Poisot, T., Canard, E., Mouillot, D., Mouquet, N., & Gravel, D. (2012). The dissimilarity of species interaction networks. In *Ecology Letters* (Vol. 15, Issue 12, pp. 1353–1361). https://doi.org/10.1111/ele.12002

R Core Team. (2021). *R: A Language and Environment for Statistical Computing* (4.0.5). R Foundation for Statistical Computing.

Rapacciuolo, G., & Blois, J. L. (2019). Understanding ecological change across large spatial, temporal and taxonomic scales: integrating data and methods in light of theory.

Ecography, *42*(7), 1247–1266. https://doi.org/10.1111/ecog.04616

Saupe, E. E., Portell, R. W., Qiao, H., Soberón, J., Hendricks, J. R., Lieberman, B. S., &
Hunter, S. J. (2015). Niche breadth and geographic range size as determinants of species
survival on geological time scales. *Global Ecology and Biogeography*, *24*(10), 1159–
1169. https://doi.org/10.1111/geb.12333

Schmidt, D. F., & Makalic, E. (2018). *randcorr: Generate a Random p x p Correlation*
Matrix (1.0).

Swain, A., Byrum, T., Zhuang, Z., Perry, L., Lin, M., & Fagan, W. (2021). *Inferring,*
comparing and exploring ecological networks from time-series data through R packages
constructnet, disgraph and dynet. 1–15.

Thébault, E., & Fontaine, C. (2010). Stability of ecological communities and the architecture
of mutualistic and trophic networks. *Science*, *329*(5993), 853–856.
https://doi.org/10.1126/science.1188321

Tikhonov, G., Opedal, Ø. H., Abrego, N., Lehikoinen, A., Jonge, M. M. J., Oksanen, J., &
Ovaskainen, O. (2020). Joint species distribution modelling with the R-package Hmsc.
Methods in Ecology and Evolution, *11*(3), 442–447. https://doi.org/10.1111/2041-
210X.13345

Tikhonov, G., Ovaskainen, O., Oksanen, J., de Jonge, M., Opedal, O., & Dallas, T. (2021).
hmsc: Hierarchical Model of Species Communities. https://doi.org/10.1111/2041-
210X.13345

Turnbull, L. A., Levine, J. M., Loreau, M., & Hector, A. (2013). Coexistence, niches and
biodiversity effects on ecosystem functioning. *Ecology Letters*, *16*(SUPPL.1), 116–127.
https://doi.org/10.1111/ele.12056

Ushio, M. (2022). Interaction capacity as a potential driver of community diversity.
Proceedings of the Royal Society B: Biological Sciences, *289*(1969).
https://doi.org/https://doi.org/10.1098/rspb.2021.2690

Veech, J. A. (2013). A probabilistic model for analysing species co-occurrence. *Global*
Ecology and Biogeography, *22*(2), 252–260. https://doi.org/10.1111/j.1466-
8238.2012.00789.x

Vinton, A. C., & Vasseur, D. A. (2020). Evolutionary tracking is determined by differential
selection on demographic rates and density dependence. *Ecology and Evolution*, *10*(12),
5725–5736. https://doi.org/10.1002/ece3.6311

Wang, D., Heckathorn, S. A., Mainali, K., & Tripathee, R. (2016). Timing effects of heat-
stress on plant ecophysiological characteristics and growth. *Frontiers in Plant Science*,
7(NOVEMBER2016), 1–11. https://doi.org/10.3389/fpls.2016.01629

Weiss-Lehman, C. P., Werner, C. M., Bowler, C. H., Hallett, L. M., Mayfield, M. M., Godoy,
O., Aoyama, L., Barabás, G., Chu, C., Ladouceur, E., Larios, L., & Shoemaker, L. G.
(2021). Disentangling key species interactions in diverse and heterogeneous
communities: A Bayesian sparse modeling approach. In *bioRxiv*. Cold Spring Harbor
Laboratory. https://doi.org/10.1101/2021.07.23.453227

Zakharova, L., Meyer, K. M., & Seifan, M. (2019). Trait-based modelling in ecology: A
review of two decades of research. *Ecological Modelling*, *407*(April), 108703.
https://doi.org/10.1016/j.ecolmodel.2019.05.008

SUPPLEMENTARY MATERIAL

SIMULATION FRAMEWORK OUTPUTS

3
FIGURE S1 – SPATIAL GRADIENT USED IN SIMULATIONS. Our analyses were executed using a linear environmental gradient

along the x-axis which established the optimal phenotype/trait value for each location of (x, y). Note that this visualisation is exaggerated to

highlight how the optimal phenotype is affected only by x. The spatial gradient used in the simulation framework is continuous rather than

categorical as shown above.

FIGURE S2 – ABUNDANCE OF CONCEPTUAL SPECIES THROUGHOUT THE SIMULATION RUN. Our simulation framework can export information on extant individuals throughout the simulation at

user-defined intervals with which additional studies can be carried out such as network inference via analyses of time-series of abundances.

FIGURE S3 – CONCEPTUAL SPATIAL PRODUCT AND MATRICES FOR NETWORK INFERENCE. Network inference approaches leverage species-by-site matrices of presence/absence or abundance. To

establish these, we divide our continuous landscape (A) into evenly spaced grid cells for which the abundance (B) of each species is then recorded and stored in a species-by-site matrix.

14   N<small>ETWORK</small> I<small>NFERENCE</small>

16   **FIGURE S4 – DISSIMILARITY OF INFERRED NETWORKS.** Networks inferred by HMSC and COOCCUR per simulation are

17   markedly dissimilar.

FIGURE S5 – RATES OF TRUE POTENTIAL ASSOCIATION DETECTION. Decomposition of overall network dissimilarities reveal that COOCCUR identifies a considerably higher rate of negative and positive associations than does HMSC while HMSC identifies far more absent associations. Two-letter codes denote type of error rate: TP…True positive, FP…False positive, MP…Missed positive (a positive association that was failed to be identified), TN…True negative, FN… False Negative, MN… Missed negative, TA…True absent, FA…False absent, and MA…Missed absent.

DETECTION PROBABILITIES OF TRUE POTENTIAL ASSOCIATIONS

We find that COOCCUR and HMSC perform similarly well at correctly identifying a

positive association as long as such associations occur between species who overlap

considerably in their environmental preferences (i.e., low environmental differences in

Error! Reference source not found.). However, HMSC struggles to identify weak

associations more so than COOCCUR whose performance is less affected by

strength/magnitude of positive associations. Meanwhile, HMSC is capable of correctly

identifying positive associations between species (provided they are strong positive

associations) for species-pairs which are quite dissimilar in their environmental preferences

which COOCCUR is incapable of.

When assessing mechanisms of accurate identification of negative associations, we find that

HMSC struggles to identify any true negative associations while COOCCUR does so

particularly when association partners have dissimilar phenotypes/environmental preferences.

Lastly, when aiming to understand how accurate inferences of absent associations come

about, we find that HMSC manages to identify absent interactions more readily. While both

network inference approaches struggle to identify true absent associations for species who

share environmental preferences, HMSC is capable of identifying true absent associations

more readily than COOCCUR particularly when environmental preferences of association

partner-species are disparate.

FIGURE S6 – PROBABILITY OF CORRECT IDENTIFICATION OF TRUE POTENTIAL ASSOCIATIONS. Logistic regression

coefficients (left column) show clear effects of association magnitude, environmental difference in species niches as well as the interaction

of these two which translate to distinct probabilities of correctly identifying a pairwise associations (right column). Effects correct

identification likelihood of association magnitude and environmental niche difference between association partners depends on sign of the

underlying association.