Full title: Mucilage Produced by Sorghum (*Sorghum bicolor*) Aerial Roots Hosts

Diazotrophs that Provide Significant Amounts of Nitrogen to the Plant

Short title: Biological Nitrogen Fixation on the Aerial Roots of Sorghum

Rafael E. Venado[1], Jennifer Wilker[1*], Vania Pankievicz[1#], Valentina Infante[1], April MacIntyre[1&],

Emily S. A. Wolf[2**], Saddie Vela[2], Fletcher Robbins[1], Paulo Ivan Fernandes-Júnior[1,3], Wilfred

Vermerris[4], and Jean-Michel Ané[1,5]

[1]Department of Bacteriology, University of Wisconsin-Madison, Madison, Wisconsin, USA

[2]Plant Molecular and Cellular Biology Graduate Program, University of Florida, Gainesville,

Florida, USA

[3]Embrapa Semiárido, Petrolina, Brazil

[4]Department of Microbiology & Cell Science and UF Genetics Institute, University of Florida,

Gainesville, Florida, USA

[5]Department of Plant and Agroecosystem Sciences, University of Wisconsin-Madison, Madison,

Wisconsin, USA

[*]Now at the Alliance of Bioversity International and CIAT, Palmira, Colombia

[#]Now at GoGenetics, Curitiba, Paraná, Brazil

[&]Now at Valent BioSciences, Libertyville, Illinois, USA

[**]Now at USDA-ARS, Beltsville, MD, USA

Corresponding Author:

Jean-Michel Ané

Tel: +1 608-262-6457

Email: jeanmichel.ane@wisc.edu

Abstract

Sorghum (*Sorghum bicolor*) is a significant crop globally, serving as an important source of food, feed, and fodder, and is increasingly recognized as an energy crop due to its high potential for biomass production. Certain sorghum accessions exhibit prolific aerial root development and produce abundant carbohydrate-rich mucilage after precipitation. This aerial root mucilage bears resemblance to that found in landraces of maize (*Zea mays*) from southern Mexico, which have previously been found to harbor diazotrophs. In this study, we examined the aerial root development of specific sorghum accessions and investigated the influence of humidity on this trait. Our microbiome analysis of the aerial root mucilage of maize and sorghum revealed the presence of numerous diazotrophs in sorghum mucilage, with *Pseudomonadota*, *Bacillota*, and *Bacteriodota* being the predominant phyla observed. However, the community composition varied significantly depending on the host plant and location. Through acetylene reduction, $^{15}N_2$ gas feeding, and ^{15}N isotope dilution assays, we determined that these sorghum accessions can acquire approximately 40% of their nitrogen from the atmosphere through these symbiotic associations on aerial roots. The nitrogen fixation occurring in sorghum aerial root mucilage presents a promising opportunity to reduce reliance on synthetic fertilizers and advance sustainable agricultural practices for food, feed, fodder, and bioenergy production.

Introduction

Nitrogen (N) is an essential nutrient for plant growth and development. However, despite its abundance in the atmosphere as dinitrogen (N_2), plants cannot access it directly [1]. Biological nitrogen fixation is the conversion of dinitrogen into ammonium performed by bacteria and archaea known as diazotrophs that express the nitrogenase enzyme [2,3]. This process is essential in agriculture, enabling certain plants, such as legumes, that associate with diazotrophs called rhizobia, to obtain large amounts of nitrogen directly from the atmosphere. Legumes host rhizobia in specialized organs called root nodules, inside which rhizobia fix nitrogen and transfer it to the plant in exchange for carbon from photosynthesis [4]. This symbiosis is critical for the productivity and sustainability of agricultural systems worldwide by reducing the need for synthetic fertilizers and enhancing soil fertility [5–7]. In contrast, cereal crops like maize (*Zea mays* L.) and sorghum (*Sorghum bicolor* (L.) Moench) generally require substantial nitrogen inputs. While synthetic nitrogen fertilizers, derived from natural gas using the Haber-Bosch process, can address this limitation, their extensive use has led to significant economic and environmental concerns, including nitrogen leaching and the generation of greenhouse gases. For decades, approaches have been proposed to enhance biological nitrogen fixation in cereals, including engineering root nodules in cereals or the expression of the nitrogenase directly in plant cells [8,9]. Another strategy is to manipulate the diazotrophs naturally associated with cereals. Such associative diazotrophs have been documented, for instance, in the rhizosphere, the xylem, or the root mucilage of cereals [10,11]. A synthetic community comprising *Klebsiella variicola* MNAZ1050 and *Citrobacter sp* MNAZ1397 was reported recently to contribute 11% of nitrogen maize through biological nitrogen fixation [10]. Some approaches also rely on altering the genetic circuits controlling nitrogenase expression in associative diazotrophs to induce the release of ammonium on the root surface [12]. While exciting, all these engineering approaches require long development times and are often

based on complex genetic manipulations that limit their development and adoption. To enhance

the benefits of biological nitrogen fixation for cereal crops in the shorter term, we explored plant

natural diversity to identify genotypes of cereals that could be better hosts for diazotrophs than

currently available cultivars.

In 2018, we and colleagues published the characterization of maize landraces cultivated and

selected by Indigenous communities in the Sierra Mixe, Oaxaca, Mexico, that secrete viscous

mucilage from their aerial roots. Unlike brace roots, these aerial roots are adventitious nodal roots

that do not reach the ground. We demonstrated that this mucilage supports communities of

diazotrophs that can provide 29-82% of the plant's nitrogen needs [13–16]. These landraces could

serve as the foundation for nitrogen fixation in maize through conventional breeding with limited

impact on yield [17]. However, the production of aerial roots is not unique to maize. Such roots

have been reported throughout the Andropogoneae and Paniceae tribes that include maize,

sorghum, sugarcane (*Saccharum* spp.), foxtail millet (*Setaria italica* L.*)*, and pearl millet

(*Pennisetum glaucum* (L) R. Br.). These roots serve anchorage functions, especially under flooded

conditions, and are important for water and nutrient acquisition [18,19]. Sorghum is a versatile and

resilient crop used for food, feed, and fodder production. It has recently gained significant attention

as a promising bioenergy crop due to the high biomass productivity of some genotypes [20,21]. Its

ability to thrive in diverse climatic conditions, including drought-prone regions, makes it an

attractive option for sustainable food and bioenergy production worldwide [22,23].

Despite the similarity between sorghum aerial roots and those in maize known to host diazotrophs,

evidence of biological nitrogen fixation in sorghum has yet to be well characterized. Images of

aerial roots of sorghum covered with mucilage, as well as a micrograph depicting the bacterium

Azospirillum brasilense, were featured in the abstract book of a conference on nitrogen fixation in cereals hosted by the International Crops Research Institute for the Semi-Arid Tropics, India (ICRISAT) in 1984. Despite this decades-old report, no data have been published on biological nitrogen fixation in sorghum aerial roots [24]. The similarity between these sorghum aerial roots and those from the Sierra Mixe maize prompted us to explore nitrogen fixation in sorghum. We examined the diversity of aerial roots in sorghum accessions, mucilage production, and their microbiome. We demonstrated that, like the maize landraces from the Sierra Mixe, some sorghum accessions can acquire significant amounts of nitrogen from their aerial roots via the diazotrophs established in their mucilage.

Results

Specific sorghum accessions developed aerial roots.

Sorghum accessions selected from the sorghum minicore [25] and augmented with accessions provided by ICRISAT exhibited extensive development of thick aerial roots, similar to those in Sierra Mixe maize landraces [13,25]. Fourteen accessions were grown in a greenhouse at the University of Wisconsin-Madison and assessed for the number of nodes with aerial roots to characterize this trait better (**Supplementary Table 1**). The number of nodes with aerial roots varied from 0 to 8 between accessions. Accessions IS17348 and IS24453 exhibited minimal and no development of aerial roots, respectively. (**Figures 1A**). Upon exposure to water, a number of accessions demonstrated substantial mucilage production from their aerial roots. Notably, accessions IS10757, IS11026, IS15170, IS15466, IS23992, IS25089, and IS28025 consistently produced an average of 200 µl of mucilage per aerial root with comparable lengths. However, accessions IS17348, IS24453, IS28023, IS28033, IS28262, and IS29078 exhibited lower mucilage

production, averaging less than 100 µl per root with similar lengths. (**Figure 1B and C**). Sorghum

aerial root diameters were measured to determine if the sorghum accessions exhibited thick aerial

roots, as shown for the Sierra Mixe maize landraces. The diameter of aerial roots ranged from 0

(when absent, only the accession IS24453) to 8 mm (**Figures 1D and 1E**). Through this screening,

we identified IS10757, IS11026, IS15170, and IS25089 as the most promising accessions,

distinguished by both thick aerial roots and high mucilage production. These results indicate

significant variation in the number and diameter of aerial roots and mucilage production among

sorghum accessions. A positive correlation was observed between the number of nodes forming

aerial roots, root diameter, and mucilage production volume (**Supplementary Figure 1A**). The

correlation between the aerial root diameter and mucilage volume was modest, $r^2 = 0.4$ (p-value =

0.0024), but low for the number of nodes producing aerial roots and volume, $r^2 = 0.2$ (p-value =

0.1517, non-significant) (**Supplementary Figure 1B**). This result indicates that, as we reported

for maize, thicker roots are associated with higher mucilage production in these sorghum

accessions, too [15].

Fig 1. Sorghum screening of selected accessions under greenhouse conditions.

Different sorghum accessions display diverse traits associated with aerial roots and mucilage

production. A) Number of nodes in different sorghum accessions (n = 3-6). Most of the accessions

evaluated showed more than 5 nodes with aerial roots. B) Representative image of selected

sorghum accessions. Scale bar = 28 cm. C) Volume of mucilage per root (µl) produced by different

sorghum accessions (n = 4-9). The production of mucilage varied among all the accessions. D)

Diameter (mm) of aerial roots in different sorghum accessions (n = 6-15). Thick roots with

diameters bigger than 4 mm were noted in nearly all the accessions. E) Aerial roots of the sorghum

accessions in D. Scale bar = 28 cm. An ANOVA (F-test) to calculate significant differences

between means was conducted with the package multcompView (0.1-10).

To test these findings under field conditions, the same set of 14 sorghum accessions were grown

at the University of Florida North Florida Research and Education Center-Suwannee Valley in

2021 (Live Oak, Florida, USA) and the University of Wisconsin-Madison West Madison

Agricultural Research Station in 2022 (Madison, Wisconsin, USA). Florida has a humid

subtropical climate, whereas Wisconsin has a continental climate that is affected by the humidity

from Lake Superior and Lake Michigan. In these field experiments, the number of nodes with

aerial roots, the root diameter, and the number of roots at the top node were recorded

(**Supplementary Figure 2**). Environmental conditions, likely a combination of soil properties and

humidity, played a significant role, as not all accessions developed aerial roots in both locations.

The sorghum accessions generally produced more aerial roots in Florida than in Wisconsin. In

Florida, approximately half of the 14 accessions produced nodes with aerial roots; in Wisconsin,

fewer than a quarter of the accessions did (**Supplementary Figures 2A and B**). Furthermore,

larger aerial root diameters were recorded in Florida compared to Wisconsin (**Supplementary**

Figures 2C and D). In Florida, the count of aerial roots at the top node exceeded that in Wisconsin,

averaging six roots compared to three roots at the top node, respectively (**Supplementary Figures**

2E and F). Overall, accessions IS11026, IS15170, and IS23992 demonstrated the potential to

develop traits related to aerial root formation and mucilage production, highlighting the significant

influence of environmental factors on these characteristics.

Humidity influences the number of nodes in sorghum.

Environmental factors can affect plant growth and development, including moisture, light

intensity, and temperature [26]. We investigated the impact of humidity on our previous findings

under greenhouse and field conditions,1 we evaluated two sorghum accessions: IS23992, which

produces many aerial roots, and IS24453, which lacks aerial roots under field conditions (**Figure**

1 and Supplementary Figure 2). We grew these accessions in a greenhouse under either high

humidity (75%) or low humidity (30%) conditions. For both sorghum accessions, we observed a

significant increase in the number of nodes with aerial roots when grown under high humidity,

with a more pronounced effect observed in IS23992 (**Figure 2A**). Interestingly, IS24453 exhibited

aerial root development in this experiment, contrasting with the results observed in the field

experiments (**Supplementary Figure 2**). Moreover, the number of aerial roots at the top node was

only significantly affected by humidity in IS24453 (**Figure 2B**). To determine whether the increase

in the number of nodes with aerial roots in humid conditions was due to differences in root

diameter, we measured the diameter of aerial roots at the top node under both humidity conditions.

The aerial root diameter did not significantly differ between these accessions (**Figure 2C**). These

findings highlight the critical role of humidity as a key environmental factor in aerial root

development in sorghum.

Fig 2. Effect of humidity in sorghum accession IS23992 and IS24453.

Boxplots of different phenotypes measured in the sorghum accessions under high (75%) and low

(30%) relative humidity. A) Number of nodes with aerial roots (n = 8). The number of nodes with

aerial roots significantly increased in both accessions. B) Number of roots on the top node (n = 8).

Only IS24453 increased the number of roots at the top node. C) Average root diameter on the top

node (n = 8). The root diameter was not affected by the humidity in both accessions. A Wilcoxon

test was performed in R (Ver 4.2.1) with the package ggpubr (Ver 0.6.0). Significance levels * p-

value ≤ 0.05, ** p-value ≤ 0.01, and n.s. not significant.

Sorghum mucilage is composed of carbohydrates.

The viscous mucilage produced by the Sierra Mixe maize landraces consists of a complex

polysaccharide with a galactose backbone and side chains of fucose, arabinose, xylose, mannose,

and glucuronic acid [13,14]. Visually the mucilage from different sorghum accessions, such as

IS2392 (**Figure 3A**), looked similar. To identify any chemical differences between maize and

sorghum mucilage, we conducted a total carbohydrate composition analysis using gas

chromatography/mass spectrometry (GC/MS). We collected mucilage from accession IS11026 in

the greenhouse, as this accession produced a substantial amount of mucilage under greenhouse

conditions (**Figure 1C**). For a clear distinction between sorghum and maize mucilage comparison,

we presented the proportions (mole %) of the quantified monosaccharides. (**Figure 3B and**

Supplementary Table 2). Arabinose, galactose, glucuronic acid, and mannose had higher ratios

in sorghum than in maize mucilage. In particular, galactose accounts for nearly 50% of the total

sugars in sorghum mucilage, and the proportion of fucose in sorghum mucilage was approximately

half of that in maize mucilage. Using this approach, we did not detect the presence of other

carbohydrates, such as glucose, rhamnose, ribose, and galacturonic acid. These results indicate

that the aerial root mucilage of the two species consists of the same monosaccharides, but that their

proportions differ.

Fig 3. Sorghum aerial root mucilage composition.

Sorghum aerial roots release mucilage . A) Mucilage production is triggered by water and under

high-humidity conditions. The mucilage from different accessions appears similar. Accession

IS2392 serves as a representative example of sorghum mucilage. Scale bar (5 cm) B)

Monosaccharide composition of sorghum accession IS11026 and Sierra Mixe maize mucilage.

Both mucilages consist of the same monosaccharides but in different proportions (mole %). C)

Concentration of reducing sugars in sorghum mucilage from the accession IS23992 determined

with benedict's reaction. A Wilcoxon test was performed to compare means *** p-value ≤ 0.001

(n = 8). D) Principal component analysis using Sparse Partial Least Squares. Two clusters are

depicted based on the species studied: sorghum accessions IS29091 and IS2245 and maize

landraces CB017456 and GRIN 19897.

In maize landraces from the Sierra Mixe, water triggered the expression of plant genes associated

with carbohydrate synthesis and degradation [15]. To investigate this degradation process further,

we assessed the levels of free sugars in sorghum mucilage at 1 and 5 hours after adding water to

the aerial roots of the accession IS23992. Using Benedict's reaction to quantify free sugars, we

observed that the free sugar concentration increased over time, rising from an average of 86 mM

at 1 hour to 109 mM after 5 hours (**Figure 3C**)[27]. This finding suggests that sorghum aerial roots

produce enzymes that break down the mucilage polysaccharide into simple sugars. We cannot

exclude the contribution of microbes in this degradation process as the aerial roots from plants

grown in the greenhouse were not germ-free. However, compared to field conditions, very few

microbes could be isolated from sorghum mucilage when grown in this greenhouse.

To explore the differences in the metabolites between maize and sorghum aerial root mucilage, we

performed a metabolomics analysis using gas chromatography coupled with time-of-flight mass

spectrometry (GC/TOF-MS) (**Supplementary Table 3**). We utilized the sorghum accessions

IS29091 and IS2245 from the Wolf *et al.* (2023) [28] study, which investigated the diversity of

aerial roots in sorghum, along with maize landraces CB017456 and GRIN19897, which display

aerial root. In total, 252 metabolites were detected in maize and sorghum mucilage (**Dataset S1)**.

A principal component analysis (PCA) revealed two clusters where samples were grouped based

on plant species (**Figure 3D**). We used a Significance Analysis of Microarray (SAM) to identify

metabolites that differed among the mucilage samples of the two species. The top 10 most

differentially abundant metabolites between sorghum and maize included 6 unknown metabolites

but also octadecanoic acid, montanic acid (octacosanoic acid), pinitol (1D-*chiro*-inositol), and *N*-

acetyl aspartate diethyl ester (**Supplementary Table 4**). These metabolomic differences may

contribute to functional differences between sorghum and maize mucilage.

Aerial root mucilage from sorghum contains abundant and large border cells.

The root cap of underground roots produces small quantities of mucilage [29–31]. In particular,

border cells (BCs) that detach from the root cap are responsible for mucilage production [15,32].

To investigate sorghum mucilage production, we examined mucilage derived from both

underground and aerial roots in sorghum accession IS23992 (**Supplementary Figures 3A and C**).

BCs derived from underground roots exhibited a distinct and elongated morphology similar to

those observed in BCs from pea (*Pisum sativum* L.), maize, and alfalfa (*Medicago sativa* L.)[33–

35]. In contrast, BCs from sorghum aerial roots appeared much larger and elongated, with irregular

shapes and sizes (**Supplementary Figure 3C**). Interestingly, these BCs from sorghum aerial roots

resembled those recently reported in mucilage from maize aerial roots [15]. These larger BCs may

contribute to the abundant mucilage production by maize and sorghum aerial roots.

Border cells detached from the underground root cap can remain active for several weeks [32,36].

To determine the viability of sorghum border cells, we used a live-dead staining protocol on both

types of sorghum mucilage collected after 72 hours. We used a combination of SYTO^{TM} 13 and

propidium iodine, which are permeating and non-permeating cell dyes, respectively [37,38]. Most

cells were viable, as indicated by only SYTO^{TM} 13 in the nuclei. Additionally, BCs from

underground roots displayed regular and well-defined nuclei, while BCs from aerial roots

exhibited irregular nuclei (**Supplementary Figures 3B and D**). To quantify the morphological

differences between the two types of BCs, we measured their length between the two cellular poles

and their cell area. Significant differences were observed in both measurements. BCs from aerial

roots are generally twice the size of BCs from underground roots. Although both types of BCs

maintain the function of mucilage production, the observed morphological differences may

account for the variation in the amount of mucilage secreted by these cells.

Sorghum mucilage hosts a wide range of diazotrophs.

To explore the microbial diversity within sorghum mucilage, we isolated and identified bacteria

using 16S amplicon sequencing. The sorghum association panel [39] and sorghum minicore [25]

were used to explore the genetic control of aerial root development in sorghum [28], obtain large

amounts of mucilage, and characterize aerial root microbiomes [28,39]. We used mucilage

samples from accessions IS2245, IS23992, IS29092 and IS2902 from the fields at the University

of Wisconsin West Madison Agricultural Research Station and the UF North Florida Research &

Education Center - Suwannee Valley. We also included mucilage samples from Sierra Mixe

landraces at a neighboring West Madison Agricultural Research Station field as reference

(**Supplementary Table 5**). To analyze the bacterial composition, we amplified and sequenced the

16S V3-V4 region and grouped the reads into Operational Taxonomic Units (OTUs) to provide

information on the relative abundance of different taxa [40]. We performed principal coordinate

analysis to examine the similarity between samples. As expected, the plant origin of the mucilage

was the main factor influencing beta diversity, although location also plays a role in this diversity

(**Figure 4A**). However, Analysis of Molecular Variance (AMOVA) uncovered instances where no

statistically significant differences were detected between the microbiomes of maize and sorghum

(**Supplementary Table 6**). We also investigated the taxonomy of bacteria in sorghum and maize

mucilage. The phylum Pseudomonadota (former Proteobacteria) was dominant in the mucilage of

both maize and sorghum, followed by Bacillota and Bacteroidota (**Figure 4B**). The profiles of

samples from Wisconsin were relatively similar regardless of the plant of origin, while samples

from Florida exhibited enrichment in Cyanobacteria (**Figure 4B**). In both mucilage samples, 34

distinct bacterial genera were identified, showing, as expected, differences in their abundance

across the samples (**Supplementary Figure 4**).

Fig 4. Microbiome profile of sorghum mucilage.

The community profile shows the relative abundance of bacteria present in sorghum (S) and maize

(M) mucilages. Samples S1 to S4 (squares) and M1 to M6 (triangles) were collected from the field

at the University of Wisconsin-Madison (UW-Madison), and samples S5 to S7 (circles) were

collected at the University of Florida (UF). A) Principal coordinates analysis plot of sorghum and

maize samples based on 16S amplicon sequences. B) Relative abundance of sorghum and maize

samples based on 16S amplicons and OTUs. C) Unrooted maximum-likelihood phylogenetic tree

of 103 bacteria isolated from sorghum aerial roots mucilage in BMGM nitrogen-free semisolid

medium. The evolutionary distances were computed using the Jukes-Cantor method and are in the

units of the number of base substitutions per site. This analysis involved 103 nucleotide sequences.

All positions containing gaps and missing data were eliminated (complete deletion). There were

908 positions in the final dataset. Numbers at nodes are bootstrap values (1000 replications, values

< 50% are not shown). Colors: Red: Gammaproteobacteria, Purple: Betaproteobacteria, Blue:

Alphaproteobacteria, Brown: Bacillota, Green: Actinomycetota, Yellow: Bacteroidota. D) Nitrogenase activity of 34 isolated from sorghum aerial roots mucilage in BMGM nitrogen-free semisolid medium (n=3). The red arrows show the reference strains for comparison. Bacteria: *Klebsiella variicola* A3 Δ*nifH*. Species codes (alphabetic order): *Agrobacterium divergens*, *Agrobacterium fabaceae*, *Agrobacterium larrymoorei*, *Agrobacterium pusense*, *Azospirillum brasilense*, *Azospirillum humicireducens*, *Azospirillum palustre*, *Bacillus amyloliquefaciens*, *Enterobacter cancerogenus*, *Epilithonimonas hungarica*, *Herbaspirillum seropedicae*, *Klebsiella michiganensis*, *Klebsiella oxytoca*, *Klebsiella variicola*, *Microbacterium aerolatum*, *Microbacterium binotii*, *Microbacterium hominis*, *Microbacterium neimengense*, *Microbacterium oleivorans*, *Microbacterium testaceum*, *Novosphingobium kaempferiae*, *Phytobacter diazotrophicus*, *Pseudoacidovorax intermedius*, *Pseudomonas bharatica*, *Pseudomonas campi*, *Pseudomonas lutea*, *Pseudomonas sediminis*, *Pseudomonas turukhanskensis*, *Pseudoxanthomonas winnipegensis*, *Rhizobium cellulosilyticum*, *Siphonobacter intestinalis*, *Stenotrophomonas lactitubi*, *Stenotrophomonas nematodicola*, *Stenotrophomonas rhizophila*, and *Stenotrophomonas terrae*.

In a second experiment, we isolated bacteria from the same sorghum association panel but expanded the number of accessions (**Supplementary Table 7**) [28,39]. The isolation of diazotrophs was performed using the nitrogen-free semi-solid BMGM medium, which revealed a high diversity of potential diazotrophs. We isolated 103 non-clonal strains from 34 different bacterial species with the prevalence of Pseudomonadota (Gammaproteobacteria - 31, Alphaproteobacteria - 33, Betaproteobacteria - 5), Actinomycetota (28), Bacteroidota (5), and Bacillota (1) (**Figure 4C and Supplementary Table 7**). Subsequently, we assessed the nitrogenase activity via the acetylene reduction assay (ARA) in one representative strain selected

from our prior isolates. Additionally, we introduced the reference strains *Azospirillum brasilense*

FP2[41], a bacterium known for promoting plant growth and commonly associated with crops, and

Klebsiella variicola A3, isolated from sorghum mucilage and established as a standard for nitrogen

fixation in our laboratory. As a negative control, we included the *Klebsiella variicola* A3 Δ*nifH*

mutant, unable to fix nitrogen. The strains *Agrobacterium pusense* 20.4, *Bacillus*

amyloliquefaciens 7.6, *Pseudomonas turukhanskensis* 14.7, *Microbacterium aerolatum* 14.4*,*

Agrobacterium fabacearum 6.8*, Microbacterium neimengense* 10.2*, Enterobacter cancerogenus*

14.3*, Epilithonimonas hungarica* 19.2*, Shinella oryzae* 21.1*, and Pseudacidovorax intermedius*

2.1 showed higher nitrogenase activity than the reference diazotrophs *K. variicola* A3 and *A.*

brasilense FP2 (**Figure 4D**). These findings validate that sorghum mucilage hosts a variety of

diazotrophs capable of potentially aiding in nitrogen fixation[42].

Mucilage enables efficient biological nitrogen fixation in sorghum.

In Sierra Mixe maize landraces, the mucilage fosters a nitrogen-acquiring habitat for diazotrophs

with low oxygen levels and an abundance of carbohydrates [13]. Using ARA, we tested whether

sorghum mucilage could serve as a proper medium for nitrogen fixation. A panel of nine

diazotrophs isolated from sorghum and maize mucilage along with other isolates from cereals (e.g.,

maize, rice (*Oryza sativa* L.), wheat (*Triticum aestivum* L.), and sugarcane) were used for this

experiment (**Supplementary Table 8**). As a negative control, we included *A. brasiliense* FP10, a

strain lacking nitrogenase activity yet exhibiting a growth rate equivalent to the wild-type FP2

strain [41]. To limit the abundance of environmental diazotrophic bacteria in the sorghum

mucilage, we collected mucilage from sorghum plants grown in the greenhouse. We selected

sorghum accession IS23992 because it produced numerous nodes with aerial roots in greenhouse

and field conditions (**Figure 1 and Supplementary Figure 2**). ARA indicated significant

nitrogenase activity in mucilage inoculated with the strains *Azotobacter vinelandii* DJ, a reference

diazotroph for decades, and *Klebsiella variicola* A3 that we isolated sorghum mucilage (**Figure**

5A). However, all seven other strains tested exhibited limited nitrogenase activity, with only a

Klebsiella michiganensis strain isolated from maize mucilage showing modest ARA activity. This

experiment indicated that the sorghum aerial root mucilage provides a proper environment for

some diazotrophs to fix nitrogen, particularly those isolated from mucilage.

Fig 5. Biological nitrogen fixation on aerial roots from sorghum.

A) Acetylene reduction assay in mucilage generated in the greenhouse. Mucilage was used as

media to grow 9 nitrogen-fixing strains (*Azospirillum brasilense* FP2, *Azospirillum baldaniorum*,

Azorhizobium caulinodans, *Azotobacter vinelandii* DJ, *Herbaspirillum seropedicae*, *Klebsiella*

michiganensis, *Klebsiella variicola* A3, *Paraburkholderia silvatlantica*, and *Pseudomonas*

stutzeri) and *Azospirillum brasilense* FP10 as a non-fixing control. Tukey's honestly significant

difference (HSD) test was performed in R, and significant group means are displayed as lower

case letters (n = 6). B) Analysis of ^{15}N gas feeding experiment on aerial roots of sorghum accession

IS23992. Aerial roots were inoculated with *K. variicola* Δ*nifH* (negative control) or *K. variicola*

A3 (positive control). *** indicates p-value ≤ 0.001 based on Wilcoxon signed-rank test (n = 8).

C) ^{15}N isotope dilution experiment on aerial roots of sorghum accessions IS23992 and IS24453.

Accessions IS23992 and IS24453 were inoculated with the nine strains from the ARA experiment

(diazotrophs), and three non-fixing mutant strains *Azospirillum brasilense* FP10, *Azotobacter*

vinelandii DJ100 Δ*nifH*, *Azorhizobium caulinodans* ORS571 Δ*nifA*. Whole plants at the flowering

stage were ground, and the average of three replicates was plotted. Percent Ndfa was calculated

for both accessions using the plants inoculated with diazotrophs and using as reference the plants

inoculated with mutant strains. An ANOVA was performed to compare means between treatments

and genotypes using the package multcompView. D) Scatterplot of the percentage of nitrogen

derived from the atmosphere (%Ndfa) in different sorghum samples from the accessions IS23992

and IS24453, indicated with circles and triangles, respectively. Solid and dotted lines are

regression lines for IS23992 and IS24453, respectively. Samples were collected every two weeks,

starting at stage 3 (Samples S3 to S5) and after flowering (S6). Error bars display standard

deviations. Pair-wise comparisons were made with Wilcoxon's test;**** means p-value ≤ 0.0001

and n.s. Indicates non-significant.

To test the plant's ability to assimilate nitrogen fixed by diazotrophs, we conducted a $^{15}N_2$ gas

feeding experiment. Aerial roots producing mucilage were collected from the greenhouse-grown

sorghum plants of the accession IS23992. Based on the ARA results, we inoculated the roots with

wild-type fixing *Klebsiella variicola* A3 or its non-fixing $\Delta nifH$ mutant strain as a negative

control[42]. Isotope-ratio mass spectrometry (IRMS) was used to quantify the $\delta^{15}N$ in chlorophyll

from sorghum aerial roots. Roots inoculated with wild-type fixing *K. variicola* exhibited

significant incorporation of ^{15}N in plant chlorophyll compared to aerial roots inoculated with the

non-fixing strain (**Figure 5B**). To provide additional evidence of fixed nitrogen uptake, we

conducted a ^{15}N isotope dilution assay. Two sorghum accessions, IS23992 and IS24453, exhibiting

high and low numbers of aerial roots, respectively, were grown in pots in the greenhouse. These

accessions were inoculated either with a panel of diazotrophs from the ARA experiment or three

non-fixing mutants (**Supplementary Table 8**). Atom%^{15}Nexcess was evaluated by IRMS in

leaves collected at four vegetative stages (S3, S4, S5, and S6) and whole plant samples at S6.

Significantly lower Atom%^{15}Nexcess values were observed in the IS23992 accession, which

produces many aerial roots, when inoculated with nitrogen-fixing bacteria compared to plants

inoculated with the non-fixing controls. Atom%^{15}Nexcess values of the IS24453 accession, which

makes few aerial roots, inoculated with fixers or non-fixers, were similar to those of the IS23992

genotype inoculated with non-fixers (**Figure 5C**). Based on these ^{15}N dilution data, we evaluated

the percentage of nitrogen derived from the atmosphere (%Ndfa) at these four stages of sorghum

development (**Supplementary Figure 5**). %Ndfa increased steadily over time, reaching 43% on

whole plants at S6 (**Figure 5D**). Even though all the techniques to quantify nitrogen fixation (ARA,

^{15}N$_2$ gas feeding, and ^{15}N isotope dilution) were used in the greenhouse and have their own biases,

our results collectively indicate that sorghum accessions with a large number of aerial roots can

support an active community of diazotrophs and obtain a significant fraction of their nitrogen from

the biological nitrogen fixation.

Discussion

Some sorghum accessions produce abundant and thick aerial roots under humid conditions.

Previous studies have reported variations in the number of nodes adventitious nodal roots among

different sorghum genotypes. For example, the Chinese landrace Sansui produces 6-8 nodes with

aerial roots, while the elite cultivar Jiliand only has a single node with roots [43]. Our study based

on select sorghum accessions from different sources such as the sorghum mincore panel, the

sorghum association panel and ICRISAT confirmed significant variation in the number of nodes

with aerial roots (**Figure 1A**). We observed differences in diameter and mucilage production in

these accessions, and, as in maize, a positive correlation among these traits was confirmed [13].

The presence of aerial roots in other members of the Andropogoneae and Paniceae tribes, such as

sugarcane and foxtail millet, suggests a common evolutionary origin of this trait [18]. It will be

interesting to explore if similar genes control this trait across these species.

Water availability, hypoxia, and humidity impact plant growth and development by producing

phytohormones, including abscisic acid and ethylene [44]. In the case of rice, high humidity

stimulates the development of lateral roots, which was attributed to increased levels of ethylene

[45–47]. Abscisic acid also controls plant responses to water availability [48]. Under high

humidity, we observed more nodes with aerial roots across sorghum genotypes, possibly by

regulating phytohormones like ethylene and abscisic acid. The accession IS24453 failed to develop

aerial roots in the initial experiment, where humidity remained at base levels. However, it

eventually developed aerial roots upon increasing humidity, likely due to increased phytohormone

levels (**Figure 2**). Subsequent experiments should prioritize quantifying these components to

mitigate unidentified experimental variables that may influence aerial root development.

Some sorghum accessions produce abundant mucilage from aerial roots.

Plants secrete mucilage from various tissues to adapt to abiotic and biotic stresses. In underground

roots, mucilage production is carried out by border cells. In the case of the maize landraces, the

mucilage from their aerial roots resembles the mucilage commonly found at the tips of

underground roots. Pankievicz *et al*. [15] showed the abundance of border cells in the mucilage

from the aerial roots of Sierra Mixe maize. They noted differences in border cell sizes compared

to border cells from underground roots. This result explained the difference in mucilage production

between the two types of roots [15]. Recently, a study involving 146 sorghum accessions described

the generation of mucilage from the aerial roots[49]. Our data corroborated these sorghum

findings, showing differences between border cell morphologies between underground and aerial

roots and mucilage production (**Supplementary Figure 3 and Figure 3A**). Based on these

findings, we hypothesize that the border cells in sorghum and maize exhibit greater similarity than

those in underground roots. The difference in quantity and size explains the volume generated by

each type of root. Additionally, we hypothesize these specialized cells are more abundant and

functionally active in younger roots, as observed in maize aerial roots.

Carbohydrates are the primary components of both underground and aerial root mucilage

[13,14,50,51]. In this study, we investigated the carbohydrate composition of sorghum mucilage

and showed the presence of similar monosaccharides in maize and sorghum aerial root mucilage,

albeit in different proportions (**Figure 3**). These findings are consistent with a prior study on maize

genotypes, which revealed changes in the carbohydrate profile based on their geographic locations

and agroecological systems [51]. Wolf *et al.* also analyzed the monosaccharide composition in

mucilage from sorghum aerial roots, and we found that both studies reported a similar range of

values [28]. To fully understand variations in monosaccharide composition in sorghum mucilage,

a broader range of environments and genotypes needs to be considered. Interestingly, our analysis

did not identify the presence of fructose in the sorghum mucilage, as reported in other studies

(**Figure 3B**) [49]. This mucilage composition may suggest that environmental factors significantly

influence maize and sorghum mucilage. However, the genetic aspect cannot be disregarded

entirely, as evidenced by our principal component analysis, which displayed a clear separation

based on species (**Figure 3**). Overall, inter-species differences are more pronounced than inter-

accession differences (**Figure 3B and 3D)**, supporting reports that highlight variations in the

microbiomes of cereals [52]. Additionally, there must be differences within genotypes, as observed

in maize underground mucilage[51].

Sorghum aerial roots mucilage hosts specific microbial communities containing diazotrophs.

Several studies have documented variations in plant microbiome composition based on host and

environmental factors [53–56]. Our microbiome survey revealed that location and host influence

microbial composition in the aerial root mucilage (**Figure 4**). Certain plant enzymes that break

down mucilage into simple sugars are expressed in the Sierra Mixe landraces and may play a role

in selecting specific microbes. The microbiome may contribute to mucilage degradation through

specialized enzymes [15]. Pseudomonadota, Acidobacteriota, Cyanobacteriota, and Bacillota were

the most abundant phyla in sorghum, aligning with findings from other sorghum microbiome

studies [57]. Our study sheds light on the microbial profiles of sorghum and maize for nitrogen

fixation and offers valuable strains for engineering diazotrophs and enhancing nitrogen fixation.

These resources will serve as a foundation for investigating microbial community interactions in

the mucilage. Such insights could pave the way for inoculation strategies with diazotrophic

consortia and/or engineered diazotrophs.

Sorghum growth promotion enabled by inoculation with diazotrophs has been reported several

years ago [58–60]. There was evidence that sorghum could acquire variable amounts of nitrogen

from the air after inoculation with diazotrophs or in association with native bacterial communities

[61,62]. However, these reports provided no information about the ability of these genotypes to

produce aerial roots or mucilage. Our study provides evidence that the production of aerial roots

and mucilage is a mechanism in sorghum that facilitates association with diazotrophs, enabling the

plant to obtain nitrogen from the atmosphere through biological nitrogen fixation. Microbial

communities can thrive in plant mucilages due to the abundance of carbon, and diazotrophs often

benefit from the microaerobic conditions. The mucilage viscosity is strikingly similar to nitrogen-

free semi-solid media, which is classically used to isolate diazotrophs from non-legumes [63]. The

presence of diazotrophs in sorghum aerial root mucilage was confirmed in our study using bacterial

isolation in nitrogen-free semisolid media and nitrogenase activity assays.

Efficient nitrogen fixation occurs in sorghum aerial root mucilage.

Biological nitrogen fixation in cereals can reduce agriculture's dependence on synthetic nitrogen-

based fertilizers. Our research has shown that mucilage is crucial in creating a favorable

environment for diazotrophs to thrive, enabling plants to acquire nitrogen (**Figure 5A, B, and C**).

Two independent studies have revealed that specific diazotrophs form associations with the

underground roots of sorghum plants, delivering nitrogen to the plant. We observed this unique

specificity as only the diazotrophs isolated from mucilage exhibited higher nitrogenase activity

than those isolated from other grasses, considered the benchmark for studying nitrogen fixation in

cereals (**Figure 5A**). The biological nitrogen fixation in sorghum was previously studied through

ARA, ^{15}N isotope dilution, and ^{15}N abundance assays [62,64,65]. This study used three

complementary approaches (ARA, ^{15}N$_2$ gas feeding, and ^{15}N isotope dilution) to provide evidence

of efficient nitrogen fixation in sorghum aerial roots (**Figure 5**). While these methods can suffer

from inherent biases, they collectively point towards efficient nitrogen fixation in the sorghum

mucilage and the acquisition of this nitrogen by the plant. The mucilage offers a favorable

environment for diazotrophs; therefore, obtaining mucilage or mucilage-producing plants devoid

of diazotrophs can be challenging. Thus, for these experiments, rather than using non-inoculated

mucilage or plants as negative controls, we favored inoculation with non-fixing mutants to exclude

as much as possible diazotrophs from the environment. ARA revealed that diazotrophs can actively

fix nitrogen in this environment (**Figure 5A**). However, ARA cannot prove that the plant can

acquire this nitrogen. This was demonstrated in ^{15}N$_2$ gas feeding and ^{15}N isotope dilution (**Figures**

5B and 5C). Previous field studies using ^{15}N isotope dilution in sorghum suggested that

diazotrophs can provide between 36 and 78 % of nitrogen depending on the edaphoclimatic

conditions [62]. In this study using ^{15}N isotope dilution in greenhouse conditions, we observed that

sorghum with aerial roots can acquire up to 43% of the atmospheric nitrogen (**Figure 5C**). ^{15}N

dilution can be biased due to the segregation of ^{15}N within the soil, and different plants may acquire

nitrogen from different soil layers. In our case, the sorghum root system filled the pots, acquiring

nitrogen from all the soil layers and alleviating this concern. However, we should be careful to

extrapolate results from greenhouse experiments to field conditions where the environment will

undoubtedly positively or negatively affect this plant-microbe relationship.

Sorghum,maize and their relatives with abundant and thick aerial roots as promising avenues to

reduce the use of synthetic nitrogen fertilizers

Farmers in Oaxaca, Mexico, have preserved and cultivated the Sierra Mixe maize landraces

for centuries. Field experiments indicated that soils with more than ten years of maize

cultivation showed higher %Ndfa values than those with zero to one year of maize history.

This consistent cultivation has promoted biological nitrogen fixation in these maize

landraces[13]. Consequently, this selection process has significantly enhanced biological

nitrogen fixation. To our knowledge, the sorghum accessions we identified have undergone

minimal selection for biological nitrogen fixation, which may explain the differences in

%Ndfa between sorghum and maize. By adopting modern cultivation practices, we may

achieve increased levels of %Ndfa in sorghum.

Sorghum is renowned for its resilience to drought, which might restrict the areas suitable

for exploiting biological nitrogen fixation from aerial roots with mucilage. In maize,

drought-adapted cultivars can release mucilage that retains water effectively [66].

Therefore, sorghum accessions that produce aerial roots and that originate from drought-

prone regions may also produce mucilage under drought conditions. However,

(sub)tropical regions or areas with high relative humidity might be ideal for these sorghum

accessions.

This study examined the diversity within the minicore and ICRISAT sorghum collections

and identified accessions that developed aerial roots. Sorghum, maize, and sugarcane are

all members of the Andropogoneae tribe, and genera from the Paniceae tribe (Setaria and

Pennisetum) are known to develop brace roots [18]. In sugarcane, thin roots emerging from

nodes have been observed [67]. During a screening of Setaria, we discovered *Setaria italica*

and *Setaria magna* accessions with thin aerial roots lacking mucilage. Additionally, we

found that Johnsongrass (*Sorghum halepense*) and sudangrass (*Sorghum bicolor* ×

Sorghum arundinaceum) developed aerial roots with mucilage (**Supplementary Figure**

6). Notably, these grasses exhibited thicker roots and mucilage. Our research indicates

thicker roots are necessary for mucilage production, as seen in maize and sorghum. Two

loci associated with aerial root diameter in sorghum were reported by Wolf et al. [39].

Based on the observations of aerial root formation in other grasses, additional screenings

of germplasm collections can provide more information on the genetic basis of the aerial

root formation and aerial root diameter.

To achieve efficient nitrogen fixation in cereals, synthetic biology offers exciting approaches

through transferring nitrogenase genes to plants, engineering root nodule symbiosis, or

remodeling associative diazotrophs [68,69]. While exploring these approaches is undeniably

valuable, it is important to explore the untapped potential present in the natural diversity of plants

to identify and breed superior plant hosts for diazotrophs using non-transgenic methods. Certain

maize landraces in Sierra Mixe can obtain significant amounts of nitrogen from the atmosphere

through their aerial roots. Our research suggests that a similar trait is present in sorghum, and

investigating the natural diversity in cereal crops for nitrogen fixation provides a near-term, non-

transgenic pathway to enhance biological nitrogen fixation in these crops. Leveraging our genetic

understanding of this trait, we can engage in breeding efforts in sorghum and maize to promote

aerial development and mucilage secretion [28]. These natural traits are quite promising to reduce

the dependence on synthetic nitrogen fertilizers for cereal crops and improve the sustainability of

our agriculture for food, feed, fodder and biofuel production.

Materials and methods

Plant material and phenotyping

Sorghum accessions were selected from the sorghum minicore collection [25] and augmented with

accessions kindly provided by Vetriventhan Mani and Vania C. R. Azevedo at the International

Crops Research Institute for the Semi-Arid Tropics (ICRISAT, Hyderabad, Telangana, India)

(**Supplementary Table 1**). The plants were cultivated in C1200 nursery pots containing Metro-

Mix® 360 SUN-COIR™ (Burton, Ohio) medium in the Walnut Street Greenhouses at the

University of Wisconsin-Madison with temperatures at 25 ˚C during the day and 17 ˚C at night.

The plants were started on a long daylight regimen (14/10), and once they were 1-1.5 m tall, they

were switched to short days (12/12) due to their photoperiod sensitivity. Relative humidity was set

at 20%. Sorghum plants were watered with municipal water or NPK fertilizer (20-10-20). At the

flowering stage, when aerial roots were visible on many nodes, the root diameter of three roots at

the top node was measured with a caliper, as well as the number of nodes forming aerial roots and

the number of aerial roots formed on the top node. Overhead water was applied, and 30 minutes

after the water application, the mucilage volume was quantified from three young and three old

aerial roots with similar lengths, of approximately 10 cm. Mucilage was collected into 2 ml tubes

by simply removing it from the base to the tip of the root. The same accessions were grown at the

University of Wisconsin-Madison West Madison Agricultural Research Station (43.0610329663,

-89.5325872441) and the University of Florida North Florida Research and Education Center -

Suwannee Valley near Live Oak, FL (30.313277, -82.902158) in 1.5 m rows, with 10 cm between

plants, and 76 cm between rows. Only the number of nodes and root diameter were phenotyped in

the field experiments.

Greenhouse experiments on the influence of humidity levels

Sorghum accessions IS23992 and IS24453 were grown in individual 6-gallon (23-L) pots

(Custom-Tainer 2800C-14") with Pro-Mix LP15 medium (Quakertown, Pennsylvania) at the

Walnut Street Greenhouses at the University of Wisconsin-Madison. Greenhouse temperature was

set at 28 ˚C during the day and 25 ˚C at night. Plants were grown under a light regime of 12-hour

photoperiod from 6 am to 6 pm. Greenhouses were kept at high (75%) or low relative humidity

30%, controlled with a humidifier system (Smart Fog, Reno, Nevada). The phenotypes evaluated

included number of nodes with roots, the number of roots at the top node, and the root diameter.

The measurements were obtained with the previously described procedure.

Bacterial growth conditions

Twelve strains were used for different experiments (**Supplementary Table 8**). *Azospirillum*

brasilense FP2 and FP10 (*nifA*⁻) were kindly provided by Fábio Pedrosa and Emanuel M. de Souza

from UFPR-Brazil[41]. The *Klebsiella variicola* Δ*nifH* mutant was developed by Maya

Venkataraman (UW-Madison). *Azospirillum baldaniorum* Sp 245^T, *Klebsiella variicola* A3,

Klebsiella michiganensis A2, *Azorhizobium caulinodans* ORS571^T, *Stutzerimonas stutzeri* A1501,

Paraburkholderia silvatlantica SRMrh-20^T, *Azotobacter vinelandii* DJ, *Herbaspirillum*

seropedicae SmR1, *Azorhizobium caulinodans* ORS571 Δ*nifA* were grown in LB medium, except

for *A. vinelandii* DJ and *A. vinelandii* DJ100 Δ*nifH*, which were grown in Burk's sucrose

medium[70,71]. The cells were grown overnight at 30 ˚C while shaking at 130 rpm.

Mucilage sugar composition

Root mucilage from accessions IS11026, was collected from cut aerial roots of different sorghum

genotypes from the greenhouse experiment. The roots underwent a 24-hour water exposure in a

50 ml tube, and the mucilage from two separate experiments was sent to the Complex

Carbohydrate Center Research Center (CCRC) at the University of Georgia, Athens, USA.

Glycosyl composition analysis was performed by combined gas chromatography/mass

spectrometry (GC/MS) of the per-*O*-trimethylsilyl (TMS) derivatives from the monosaccharide

methyl glycosides produced by acidic methanolysis as described previously [13]. The analysis

included rabinose, rhamnose, ribose, fucose, xylose, glucuronic acid, galacturonic acid, mannose,

galactose, glucose, *N*-acetyl mannosamine, *N*-acetyl galactosamine, *N*-acetyl glucosamine were

measured but only monosaccharides that were detected are reported.

Total free sugar quantification

Mucilage was collected at two time points from the sorghum accession IS23992 grown under

greenhouse conditions. Benedict's reaction was used to quantify free sugars, as described in

Hernandez-Lopez *et al*.[27]. Benedict's reagent (700 µl) was incubated with sorghum mucilage

(30 µl) for 10 min at 95 ºC. Subsequently, the sample was centrifuged at 10,000 g for 5 min. The

supernatant was collected, and the absorbance at 740 nm was determined using a BioRad

SmartSpect[TM] Plus spectrophotometer. Sugars were quantified using a calibration based on

solutions with different concentrations of D-glucose.

Metabolomic analyses

Metabolome analysis of mucilage samples from sorghum accessions IS29091 and IS2241, as well

as maize landraces CB017456 and GRIN 19897, was conducted at the West Coast Metabolomics

Center at UC Davis following their standard protocols (https://metabolomics.ucdavis.edu/). A

minumum of 75 mg of mucilage was sent per sample (Supplementary Table 3). A primary

metabole analysis was performed to detect sugar phosphates, amino acids, hydroxyl acids, free

fatty acids, purines, pyrimidines, aromatics compounds, and unidentified compounds. Metabolites

were quantified by peak heights, normalized, and processed with MetaboAnalyst[72]

(https://www.metaboanalyst.ca/). This algorithm offers several different methods. Sparse partial

least squares - discriminant analysis and significance analysis of microarray were used to analyze

the data.

Microscopy

Border cells from underground roots were collected from the root tip of six-day-old seedlings from

sorghum accession IS23992 that grew in 1.2% agar at 24 ºC in a growth chamber. Border cells

from aerial roots were collected from mucilage produced from the humidity study. The mucilage

was triggered by spraying the aerial roots directly with municipal water 5 times every 5 minutes.

The mucilage was collected after 1 hour after the final spraying. Approximately 5 µl of mucilage

were spread on a slide and combined with 5 µl of distilled water, 10 µl of SYTO green (5 µM),

and 10 μl of propidium iodine (10 μg/mL). The cell suspension was observed under a Leica DMi8

model S inverted fluorescence microscope. GFP and mCherry filters were used to examine SYTO

green and propidium iodide fluorescent signals, respectively. Light microscopy was used to obtain

images of border cells from both underground and aerial roots. ImageJ - Fiji (version 2.9.0) [73]

was used to measure the area and length of the border cells.

Microbiome study and 16S rRNA amplicon sequencing

DNA was isolated from the mucilage samples from sorghum and maize using the FastDNA™

SPIN Kit for Soil (MP Biomedicals, Irvine, California) following the manufacturer's instructions.

DNA was quantified using a Nanodrop One (Thermofisher Scientific, Waltham, Massachusetts).

The V3-V4 region of the *16S rRNA* gene was amplified with the standard bacterial primers 335F

and 769R [74] with Go*Taq* Mastermix (Promega, Madison, Wisconsin) according to the

manufacturer's instructions. Amplification reaction volumes were 50 μl containing 10 pmol

of each primer and 4 ng of template DNA. Thermocycler conditions were as follows: 95°C

for 1 minute, then 35 cycles of 95°C for 10 s, 55°C for 30 s, and 72°C for 30 s, with a final

step at 72°C for 5 min. PCR products were separated on an agarose gel and extracted using

the GeneJET Gel Extraction Kit (Thermofisher Scientific). Novogene (Santa Clara,

California, USA) conducted library preparation and sequencing. Amplicons were sequenced

with an Illumina paired-end platform generating 250 bp paired-end raw reads. OTUs

abundance was normalized using a standard sequence number corresponding to the sample

with the fewest sequences. Sequences exhibiting a similarity of ≥97% were grouped into

identical OTUs. A representative sequence was chosen for each OTU to undergo subsequent

annotation. Principal component and beta diversity analyses were performed based on these

normalized data. These data were processed with QIIME2 (Version 1.7.0) [75] and displayed

with R software (Version 2.15.3) package ggplot2 (Version 3.4.3.) [76]. Sequencing data are

available at the Short Read Archive of NCBI under BioProject: PRJNA1009066.

Isolation of bacteria from sorghum samples

Mucilage samples were collected from a West Madison Agricultural Research Station field trial in

2022 from mucilage after rain (**Supplementary Tables 5 and 7**). The sampling was conducted

after a continuous rainy weekend with an accumulated rainfall of around 60 mm. In the field, we

looked for homogeneous mucilage production in the sorghum association panel [28,39]. A sample

was composed of a mix of two replicates from the panel. Mucilage (2-3 mL per plant) was collected

with a 1-mL pipette in a 15 mL tube. The samples were kept cold (0 ºC) until analysis. The

mucilage samples were added to BMGM semisolid medium and incubated at room temperature

for one week [77]. After that, the microaerophilic pellicles that formed were transferred to a fresh

medium and incubated for the same period seven more times. After the final incubation period,

solid BMGM medium supplemented with 0.5 g/l of yeast extract was inoculated with the bacteria.

A total of 164 isolates were screened by clonal strains using the Box-PCR approach with the A1

primer (5′-CTACGGCAAGGCGACGCTGACG-3′) and subjected to agarose gel electrophoresis

(1.2%) [78]. Identical profiles were excluded, and 103 non-clonal strains were kept. The bacterial

strains were identified by sequencing 16S rRNA genes using the primers 27F (5'-

AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-GGTTACCTTGTTACGACTT-3')[79]. The

PCR was initiated with a denaturation cycle at 95°C for 10 minutes, followed by 35 cycles

consisting of denaturation at 95°C for 30 seconds, annealing at 55°C for 30 seconds, and extension

at 72°C for 1 minute and 30 seconds. The process concluded with a final extension at 72°C for 10

minutes. PCR product was sequenced in an Applied Biosystems 3500 genetic analyzer at the

Functional Biosciences facilities. Sequence Scanner v.2.0 (Informer Technologies, Inc.) was used

for quality assessment, and the contigs of almost complete 16S rRNA sequences were assembled

in BioEdit v.7.7[80]. The sequences were compared against those deposited within the 16S rRNA

reference database of GeneBank using the BLASTn tool [81]. The maximum likelihood

phylogenetic tree was constructed using MEGA 11 software [82]. The aligning used 103 different

16S rRNA gene sequences from the bacteria isolated from the sorghum mucilage.

Acetylene reduction assays (ARA) to quantify nitrogenase activity

Nitrogenase activity was evaluated in ten bacterial strains (**Supplementary Table 8**) grown in

sorghum mucilage collected from the greenhouse. The bacteria were grown in 3 ml of LB medium

overnight at 30 ºC and shaken at 180 rpm. The culture was centrifuged at 5000 g for 5 minutes,

and the pellet was resuspended in 1 ml of sterile water. Then 30 µl (OD_{600nm} = 0.1) were inoculated

in sterile vials containing 3 ml of mucilage. Vials were capped and incubated for 24 hours at 30

ºC. The cap was replaced with a crimp cap, and 1 ml of air was replaced with acetylene, followed

by a 3-day incubation period at 30 ºC. Gas Chromatography - Flame Ionization Detection was

performed in a GC-2010 (Shimadzu) with an HS20 autosampler by injecting 1 ml of air sample.

The injector and the detector were set at 200 ºC with a flow rate of 86.6 ml/min nitrogen carrier

gas. The temperature of the column was 100 ºC. Controls included a vial with acetylene to check

for traces of ethylene and *A. brasilense* FP2 without fixing activity. The indirect measure of the

nitrogenase activity was expressed in nmol of ethylene produced per hour per mL of mucilage.

ARA in the 34 isolated diazotrophs was performed similarly but using a semi-solid BMGM

medium.

^{15}N$_2$ gas feeding experiments

Aerial roots from sorghum accession IS23992 were collected in 150 ml sterile flasks. These roots were obtained from greenhouse-grown plants with copious amounts of mucilage. *K. variicola A3* and the *K. variicola ΔnifH* mutant were grown overnight at 30 ºC and shaken at 180 rpm in LB medium. The culture underwent centrifugation at 5000 *g* for 5 minutes, and the resulting pellet was then resuspended in 1 ml of sterile water. Bacteria were directly inoculated onto the surface of mucilage-coated roots (100 µl, OD$_{600nm}$ = 0.1). Flasks were injected with ^{15}N$_2$ gas (98%) to reach a concentration of 15% (v/v), and flasks were incubated for three days at room temperature (20ºC). Non-fixing bacteria were included as negative controls to confirm the absence of other sources of ^{15}N in the system. The roots with mucilage were excised from the plant for practicality and proper incubation in ^{15}N$_2$ gas. To determine the ^{15}N enrichment specifically in plant tissue, chlorophyll was extracted by peeling the green cell layers from the aerial roots after removal of mucilage. The extraction followed the protocol described by Kahn *et al.* [83]. The samples were analyzed by isotope ratio mass spectrometry (IRMS) at the Department of Soil Science at the University of Wisconsin-Madison to evaluate the δ^{15}N of the samples.

^{15}N isotope dilution experiments

A total of 40 sorghum plants from each of the two accessions, IS23992 and IS24453, were grown in 6-gallon (23-L) pots at the Walnut Street Greenhouses in two adjacent greenhouse rooms to prevent contamination between the two types of bacterial inoculants (fixing and non-fixing). An equal number of plants were placed in the two treatments, and standard humidity conditions were set (relative humidity 50%) with a humidifier system (Smart Fog, Reno, Nevada). After four weeks

post-germination, pots were fertilized weekly with 500 ml of Hoagland's No.2 Basal Salt mixture

(1.34 g per litter) and 80% ammonium-^{15}N sulfate (^{15}NH$_4$)$_2$SO$_4$ as the only source of nitrogen.

Five weeks after germination, aerial roots of plants in one room were inoculated with a mixture of

nine fixing strains (OD$_{600}$ = 0.1), and the aerial roots on the plants in the other room was inoculated

with three non-fixing strains (OD$_{600}$ = 0.1) (**Supplementary Table 8**). Plants were sprayed with

250 ml of the inoculum resuspended in a phosphate-buffered saline (PBS) buffer using a backpack

sprayer. The third leaf counting from the top was collected before inoculation (stage 3) and then

three times every two weeks (stages 4 and 5). At flowering (stage 6), whole plants were collected,

ground, and homogenized to a fine powder . Quantification of the samples' ^{15}N content above the

atmospheric concentration of N$_2$ (Atom% ^{15}N excess) was measured by isotope ratio mass

spectrometry (IRMS) at the Department of Soil Science at the University of Wisconsin-Madison.

The values of whole-plant samples were averaged. The percentage of nitrogen derived from the

atmosphere (%Ndfa) was calculated according to the equation described in van Deynze *et al*.[13].

The fixers were the accessions inoculated with diazotrophs, whereas the references were those

inoculated with the mutant strains.

***Setaria* growth conditions**

Setaria accessions PI43384 and PI689086 were cultivated at the Walnut Street Greenhouses under

high relative humidity (80%) using the same system as described for the humidity experiment with

sorghum. The plants were grown in 1-gallon (3.8-L) Pro Cal Premium Nursery Pots with Pro-Mix

LP15 medium from Quakertown, Pennsylvania. Plants were maintained at 28 ˚C during the day

and 25 ˚C at night, with a 12-hour photoperiod from 6 am to 6 pm. The phenotype evaluated

included the number of nodes with roots.

Raw data

All raw data are available at DOI: 10.6084/m9.figshare.25193696

Statistical analyses

All statistical analyses were conducted in R version 4.2.1 and 2.15.3[84]. ANOVA and Tukey's HSD were conducted with the package ggpubr (Version 0.6.0) and multcompView (Version 0.1-9). Wilcoxon's test was also calculated with package ggpubr (Version 0.6.0). Microbiome studies were performed with QUIIME2 (Version 1.7.0) [85].

Acknowledgments

We thank Sanhita Chakraborty and Kimberly Gibson for their valuable input and engaging discussions, Alyssa Davis and Julio Quiñones for their technical support, and Ben Broughton and Mike Boyette at the UF North Florida Research & Education Center-Suwannee Valley for managing the field plots. We thank Janet Hedtcke, the field manager at UW-Madison Agricultural Research Station. We thank Maya Venkataraman and Brian F. Pfleger for providing the *K. variicola ΔnifH* strain, David Brenner for providing us with seeds from different *Setaria* accessions, and Daniel Carmeli for providing pictures from Johnsongrass and sudangrass. Funding for this project was provided by the United States Department of Energy (DOE), Office of Science (BER), grant no. DE-SC0021052 (JMA, WV).

Fig 1. Sorghum screening of selected accessions under greenhouse conditions.

Different sorghum accessions display diverse traits associated with aerial roots and mucilage production. A) Number of nodes in different sorghum accessions (n = 3-6). Most of the accessions evaluated showed more than 5 nodes with aerial roots. B) Representative image of selected sorghum accessions. Scale bar = 28 cm. C) Volume of mucilage per root (µl) produced by different sorghum accessions (n = 4-9). The production of mucilage varied among all the accessions. D)

807   Diameter (mm) of aerial roots in different sorghum accessions (n = 6-15). Thick roots with

808   diameters bigger than 4 mm were noted in nearly all the accessions. E) Aerial roots of the sorghum

809   accessions in D. Scale bar = 28 cm. An ANOVA (F-test) to calculate significant differences

810   between means was conducted with the package multcompView (0.1-10).

Fig 2. Effect of humidity in sorghum accession IS23992 and IS24453.

Boxplots of different phenotypes measured in the sorghum accessions under high (75%) and low (30%) relative humidity. A) Number of nodes with aerial roots (n = 8). The number of nodes with aerial roots significantly increased in both accessions. B) Number of roots on the top node (n = 8). Only IS24453 increased the number of roots at the top node. C) Average root diameter on the top node (n = 8). The root diameter was not affected by the humidity in both accessions. A Wilcoxon test was performed in R (Ver 4.2.1) with the package ggpubr (Ver 0.6.0). Significance levels * p-value ≤ 0.05, ** p-value ≤ 0.01, and n.s. not significant.

A

B

Monosaccharide	Amount *S. bicolor* (%)[1]	Amount *Z. mays* (%)[2]
Arabinose (Ara)	20.22	13.7
Fucose (Fuc)	15.23	41.8
Xylose (Xyl)	2.66	3.3
Glucuronic acid (GlcA)	7.4	2.7
Mannose (Man)	6.4	2.7
Galactose (Gal)	47.16	35.8
Total	**100**	**100**

[1] Data obtained from the average of two biological replicates
[2] Data from Van Deynze *et al.* 2018

C

D

Fig 3. Sorghum aerial root mucilage composition.

Sorghum aerial roots release mucilage . A) Mucilage production is triggered by water and under

high-humidity conditions. The mucilage from different accessions appears similar. Accession

IS2392 serves as a representative example of sorghum mucilage. Scale bar (5 cm) B)

Monosaccharide composition of sorghum accession IS11026 and Sierra Mixe maize mucilage.

Both mucilages consist of the same monosaccharides but in different proportions (mole %). C)

829   Concentration of reducing sugars in sorghum mucilage from the accession IS23992 determined

830   with benedict's reaction. A Wilcoxon test was performed to compare means *** p-value ≤ 0.001

831   (n = 8). D) Principal component analysis using Sparse Partial Least Squares. Two clusters are

832   depicted based on the species studied: sorghum accessions IS29091 and IS2245 and maize

833   landraces CB017456 and GRIN 19897.

Fig 4. Microbiome profile of sorghum mucilage.

The community profile shows the relative abundance of bacteria present in sorghum (S) and maize

(M) mucilages. Samples S1 to S4 (squares) and M1 to M6 (triangles) were collected from the field

at the University of Wisconsin-Madison (UW-Madison), and samples S5 to S7 (circles) were

collected at the University of Florida (UF). A) Principal coordinates analysis plot of sorghum and

maize samples based on 16S amplicon sequences. B) Relative abundance of sorghum and maize

samples based on 16S amplicons and OTUs. C) Unrooted maximum-likelihood phylogenetic tree

of 103 bacteria isolated from sorghum aerial roots mucilage in BMGM nitrogen-free semisolid

medium. The evolutionary distances were computed using the Jukes-Cantor method and are in the

units of the number of base substitutions per site. This analysis involved 103 nucleotide sequences.

All positions containing gaps and missing data were eliminated (complete deletion). There were

908 positions in the final dataset. Numbers at nodes are bootstrap values (1000 replications, values

< 50% are not shown). Colors: Red: Gammaproteobacteria, Purple: Betaproteobacteria, Blue:

Alphaproteobacteria, Brown: Bacillota, Green: Actinomycetota, Yellow: Bacteroidota. D)

Nitrogenase activity of 34 isolated from sorghum aerial roots mucilage in BMGM nitrogen-free

semisolid medium (n=3). The red arrows show the reference strains for comparison. Bacteria:

Klebsiella variicola A3 Δ*nifH*. Species codes (alphabetic order): *Agrobacterium divergens*,

Agrobacterium fabaceae, *Agrobacterium larrymoorei*, *Agrobacterium pusense*, *Azospirillum*

brasilense, *Azospirillum humicireducens*, *Azospirillum palustre*, *Bacillus amyloliquefaciens*,

Enterobacter cancerogenus, *Epilithonimonas hungarica*, *Herbaspirillum seropedicae*, *Klebsiella*

michiganensis, *Klebsiella oxytoca*, *Klebsiella variicola*, *Microbacterium aerolatum*,

Microbacterium binotii, *Microbacterium hominis*, *Microbacterium neimengense*, *Microbacterium*

oleivorans, *Microbacterium testaceum*, *Novosphingobium kaempferiae*, *Phytobacter*

diazotrophicus, *Pseudoacidovorax intermedius*, *Pseudomonas bharatica*, *Pseudomonas campi*,

Pseudomonas lutea, *Pseudomonas sediminis*, *Pseudomonas turukhanskensis*,

Pseudoxanthomonas winnipegensis, *Rhizobium cellulosilyticum*, *Siphonobacter intestinalis*,

Stenotrophomonas lactitubi, *Stenotrophomonas nematodicola*, *Stenotrophomonas rhizophila*, and

Stenotrophomonas terrae.

Fig 5. Biological nitrogen fixation on aerial roots from sorghum.

A) Acetylene reduction assay in mucilage generated in the greenhouse. Mucilage was used as media to grow 9 nitrogen-fixing strains (*Azospirillum brasilense* FP2, *Azospirillum baldaniorum*, *Azorhizobium caulinodans*, *Azotobacter vinelandii* DJ, *Herbaspirillum seropedicae*, *Klebsiella michiganensis*, *Klebsiella variicola* A3, *Paraburkholderia silvatlantica*, and *Pseudomonas stutzeri*) and *Azospirillum brasilense* FP10 as a non-fixing control. Tukey's honestly significant difference (HSD) test was performed in R, and significant group means are displayed as lower

case letters (n = 6). B) Analysis of ^{15}N gas feeding experiment on aerial roots of sorghum accession IS23992. Aerial roots were inoculated with *K. variicola* Δ*nifH* (negative control) or *K. variicola* A3 (positive control). *** indicates p-value ≤ 0.001 based on Wilcoxon signed-rank test (n = 8). C) ^{15}N isotope dilution experiment on aerial roots of sorghum accessions IS23992 and IS24453. Accessions IS23992 and IS24453 were inoculated with the nine strains from the ARA experiment (diazotrophs), and three non-fixing mutant strains *Azospirillum brasilense* FP10, *Azotobacter vinelandii* DJ100 Δ*nifH*, *Azorhizobium caulinodans* ORS571 Δ*nifA*. Whole plants at the flowering stage were ground, and the average of three replicates was plotted. Percent Ndfa was calculated for both accessions using the plants inoculated with diazotrophs and using as reference the plants inoculated with mutant strains. An ANOVA was performed to compare means between treatments and genotypes using the package multcompView. D) Scatterplot of the percentage of nitrogen derived from the atmosphere (%Ndfa) in different sorghum samples from the accessions IS23992 and IS24453, indicated with circles and triangles, respectively. Solid and dotted lines are regression lines for IS23992 and IS24453, respectively. Samples were collected every two weeks, starting at stage 3 (Samples S3 to S5) and after flowering (S6). Error bars display standard deviations. Pair-wise comparisons were made with Wilcoxon's test;**** means p-value ≤ 0.0001 and n.s. Indicates non-significant.

Suppl. Fig 1. Trait correlations.

A) Scatter plot shows the correlation between the number of nodes, mucilage volume, and

diameter. A 3D regression line is dot plotted (purple) to observe the correlation among the traits.

B) The correlation matrix for the number of nodes, mucilage volume, and diameter reveals a

modest positive correlation across all traits. The correlation coefficient (r) values indicate the

897 correlation strength between each trait. The correlation plot and values were calculated with the

898 packages "scatterplot3d" [86] and "psych" [87], respectively.

Suppl. Fig 2. Sorghum screening of selected genotypes in two locations.

A) Number of nodes in different sorghum accessions grown at the University of Florida during

2021 (n = 5). Generally, two groups can be observed, one with nodes with aerial roots and another

without. B) Number of nodes in different sorghum accessions grown at the University of

Wisconsin during 2022 (n = 3). Only specific accessions developed fewer aerial roots compared

with the conditions in Florida. C) Diameter (mm) of aerial roots of different sorghum accessions

grown at the University of Florida (n = 5). The accessions with aerial roots have thick diameters.

D) Diameter (mm) of aerial roots in different sorghum accessions grown at the University of

Wisconsin during 2022 (n = 3). Thickness was still observed in the accession with aerial roots. E)

Number of aerial roots at the top node in different sorghum accessions grown at the University of

Florida (n = 5). F) Number of aerial roots at the top node in different sorghum accessions grown

at the University of Wisconsin during 2022 (n = 3). * Data from these accessions were obtained

under a low nitrogen regimen. A subset of data for the Florida plots were obtained from the Wolf

et al. study [28].

Underground root border cells - *S.bicolor* Aerial root border cells - *S.bicolor*

A

C

B

D

Bright field

SYTO 13 combined with propidium iodine

E

F

Suppl. Fig 3. Border cells in sorghum mucilage.

A) Border cells from sorghum underground roots (S=scale bar 100 μm). Border cells have a bean

shape. B) Border cells from sorghum aerial roots (scale bar 100 μm). Irregular cells are observed

in a more elongated form. C) Live-dead viability staining in border cells from sorghum

underground roots (scale bar 100 μm). The presence of live cells was recorded using the green

fluorescence of their nuclei. D) Live-dead viability staining in border cells from sorghum aerial

roots (scale bar 100 μ m). Live cells were identified using the green fluorescence of their nuclei.

E) Length comparison between border cells in μm. Border cells from mucilage were more

elongated than border cells from underground roots. F) Area comparison between border cells in

μm^2. ** and *** denote p< 0.01 and < 0.001, respectively, based on a Wilcoxon test (n = 20).

Suppl. Fig 4: Bacterial genera from different sorghum and maize mucilage.

The hierarchical clustering heatmap represents the microbiome community composition at the genus level for 13 samples obtained from the mucilage of sorghum (S) and maize (M). The samples have been grouped through clustering based on their community composition. The microbiome of each sample was obtained from different genotypes and locations, thus showing unique profiles.

Suppl. Fig 5. ^{15}N isotope dilution experiment on aerial roots of sorghum accessions from leaf samples.

Accessions IS23992 and IS24453 were inoculated with the nine strains from the ARA experiment (diazotrophs), and three mutant strains *Azospirillum brasilense* FP10, *Azotobacter vinelandii* DJ100 Δ*nifH*, *Azorhizobium caulinodans* ORS571 Δ*nifA*. The third topmost leave was analyzed. The samples were collected before inoculation (sample one - stage three) and every two weeks (sample 3 to 4 - stage 4 to 6). An ANOVA was conducted using the package multcompViewto determine the factors contributing to differences in means.

Suppl. Fig 6. Aerial roots in other grass species

A preliminary screening in *Setaria* (*Setaria italica* and *Setaria magna*) and other species of the

Sorghum clade revealed the presence of aerial roots. A) *Setaria magna* accession PI689086

displayed few aerial roots. Scale bar: 20 cm. B) Close-up of an aerial root at the base of *Setaria*

italica accession PI689086. C) The *Setaria magna* accession PI433384 exhibited more aerial roots.

Scale bar: 20 cm. D) Close-up of an aerial root at the base of *Setaria magna* accession PI433384.

E) Quantification of aerial roots in Setaria accessions PI433384 and PI689086. F) Johnsongrass

(*Sorghum halapense*) grown in Tallahassee, Florida, displayed aerial roots with mucilage. G)

Sudangrass (*Sorghum bicolor × Sorghum arundinaceum*) grown in western North Carolina

displayed aerial roots with mucilage. Asterisks mark the nodes with aerial roots; white arrows

indicate aerial roots with mucilage.

Supplementary Table 1. Sorghum accessions to evaluate aerial root formation in greenhouse and

field experiments

Accession	Collection	Origin
IS10757	Mini core	Chad
IS11026	Mini core	Ethiopia
IS15170	Mini core	Cameron
IS15466	Mini core	Cameron
IS15478	Mini core	Cameron
IS17348	ICRISAT	Ethiopia
IS23992	Mini core	Yemen
IS24453	Mini core	South Africa
IS25089	Mini core	Ghana
IS28023	ICRISAT	Yemen
IS28025	ICRISAT	Yemen
IS28033	ICRISAT	Yemen
IS28262	ICRISAT	Yemen
IS29078	ICRISAT	Yemen

963   **Supplementary Table 2.** Mucilage sugar composition of accession IS11026 in two independent

964   experiments. Monosaccharide composition was determined by GC-MS. * data obtained from 1.64

965   mg sample; ** data obtained from a 2.16 mg sample; nd = not detected.

	Replicate 1*		Replicate 2**	
Monosaccharide	**Amount (µg)**	**% by mole**	**Amount (µg)**	**% by mole**
Arabinose	163.9	19.77	254.4	20.66
Rhamnose	nd	-	nd	-
Ribose	nd	-	nd	-
Fucose	124.9	13.77	224.6	16.68
Xylose	21.1	2.54	34.1	2.77
Glucuronic Acid	75.9	7.08	121.8	7.65
Galacturonic acid	nd	-	nd	-
Mannose	67.8	6.81	87.1	5.89
Galactose	484.9	48.73	674.0	45.59
Glucose	12.9	1.29	11.4	0.77
N-Acetyl Mannosamine	nd	-	nd	-
N-Acetyl Galactosamine	nd	-	nd	-
N-Acetyl Glucosamine	nd	-	nd	-
Total	951.4	100	1,407.4	100

968   **Supplementary Table 3.** Mucilage samples from maize and sorghum for primary metabolism

969   analysis

Genotype	Plant	Volume (mg)
IS 29091 biological replicate 1	Sorghum	1500
IS 29091 biological replicate 2	Sorghum	200
IS 29091 biological replicate 3	Sorghum	1000
IS 2245 biological replicate 1	Sorghum	150
IS 2245 biological replicate 2	Sorghum	75
IS 2245 biological replicate 3	Sorghum	250
CIMMYT-BANK-017456 biological replicate 1	Maize	750
CIMMYT-BANK-017456 biological replicate 2	Maize	750
CIMMYT BANK 017456 biological replicate 3	Maize	500
CIMMYT BANK 017456 biological replicate 4	Maize	500
GRIN AMES 19897 biological replicate 1	Maize	400
GRIN AMES 19897 biological replicate 2	Maize	500
GRIN AMES 19897 biological replicate 3	Maize	250
GRIN AMES 19897 biological replicate 4	Maize	100

972   **Supplementary Table 4.** Top metabolites in aerial root mucilage. Peak (mz/rt) is the mass by

973   charge (mz) divided by the retention time (rt). d.value denotes the comparison between sorghum

974   and maize, where a positive value signifies that the metabolite is more abundant in sorghum than

975   maize, and negative values indicate higher abundance in maize. Other acronyms include standard

976   deviation (stdev), raw p-value (rawp), and q-value (q.value) for statistical analysis.

	Peak (mz/rt)	d.value	stdev	rawp	q.value
1	4850	8.6391	22.282	0	0
2	montanic acid	5.9485	1276.4	0.00011905	0.0032409
3	1064	5.825	2328.1	0.00011905	0.0032409
4	4901	-5.7521	469.11	0.00015873	0.0032409
5	168	5.3462	3634.3	0.00015873	0.0032409
6	Pinitol	-5.1539	375.29	0.0002381	0.0040512
7	Octadecanol	4.7778	248.11	0.00043651	0.0049514
8	N-acetylaspartate dyethilester	4.7568	18986	0.00043651	0.0049514
9	84161	4.7356	631.58	0.00043651	0.0049514
10	505155	4.5206	19490	0.00059524	0.0057391
11	328824	-4.4684	166.07	0.00063492	0.0057391
12	505146	4.4321	19227	0.0006746	0.0057391
13	100956	-4.2263	150.2	0.00087302	0.0068558
14	levoglucosan	4.0558	195.21	0.0011111	0.0081023
15	dodecanol	3.7188	293.75	0.001746	0.011883
16	1725	3.676	14665	0.0019841	0.01263
17	14724	3.6622	143.57	0.0021032	0.01263
18	azelaic acid	3.5732	108.45	0.0025397	0.014404
19	103182	3.5634	85.184	0.0026984	0.014499
20	21885	3.4903	1123.9	0.0032937	0.015836
21	thymine	-3.4522	216.9	0.0034127	0.015836
22	lignoceric acid	3.4511	465.46	0.0034127	0.015836
23	127661	3.4037	393.86	0.0037698	0.016733
24	2551	3.3414	130.98	0.0043254	0.018399
25	putrescine	-3.2171	1115.5	0.0057937	0.023659
26	threonine	-3.1399	2661.8	0.0071429	0.028046
27	169611	-3.1161	750.57	0.0074603	0.028208
28	42357	3.0631	321.56	0.0080159	0.028218
29	87877	3.0619	710.68	0.0080159	0.028218
30	4-hydroxymandelic acid	3.0201	290.67	0.0084921	0.028898
31	1-octacosanol	2.8541	963.42	0.012143	0.035571
32	isopalmitic acid	2.8421	218.03	0.01246	0.035571
33	4-hydroxybenzoic acid	2.8215	5841.1	0.012937	0.035571
34	adipic acid	2.8159	1130.8	0.013175	0.035571
35	47197	2.7985	335	0.013611	0.035571
36	131202	2.7821	258.41	0.014048	0.035571

981   **Supplementary Table 5.** Maize and sorghum accession for microbiome study. Number of

982   replicates is indicated in the sample ID.

Sample ID	Genotype	Plant	Location
S1 (4 samples)	IS 2245	sorghum	West Madison
S2 (3 samples)	IS 2245*	sorghum	West Madison
S3 (4 samples)	IS 23992	sorghum	West Madison
S4 (4 samples)	IS 29092	sorghum	West Madison
S5 (4 samples)	IS 2902	sorghum	UF
S6 (4 samples)	IS 2902*	sorghum	UF
S7 (4 samples)	IS 2245	sorghum	UF
M1 (4 samples)	CIMMYT-BANK-017456	maize	West Madison
M2 (4 samples)	CIMMYT-BANK-017456	maize	West Madison
M3 (4 samples)	CIMMYTA-BANK-014019	maize	West Madison
M4 (4 samples)	GRIN AMES 19897	maize	West Madison
M5 (3 samples)	GRIN AMES 19897	maize	West Madison
M6 (4 samples)	GRIN AMES 19897	maize	West Madison

*Samples were obtained from a low nitrogen pilot study

985 **Supplementary Table 6.** AMOVA for each sample comparison. S = sorghum

986 sample, M = maize sample, and df = degrees of freedom. Statistically

987 significant differences are indicated with asterisks: * p ≤ 0.05; ** p ≤

988 0.01, and *** p ≤ 0.001

Group	Sum of Square	df	Mean Square	Fixation Index	p-value
S1-M3	0.144504	1	0.144504	5.1731	0.005**
M4-M5	0.0855831	1	0.0855831	2.22751	0.095
S2-M4	0.0864243	1	0.0864243	3.0695	0.038*
M2-M5	0.115087	1	0.115087	3.71158	0.029*
S6-S7	0.0614475	1	0.0614475	0.486006	0.61
S3-M5	0.180132	1	0.180132	5.09055	0.026*
S1-S6	0.184116	1	0.184116	1.95325	0.263
S5-M6	0.450519	1	0.450519	5.24901	0.01**
S3-M1	0.054311	1	0.054311	1.92444	0.114
S2-M3	0.186426	1	0.186426	7.09958	0.03*
S2-S7	0.266473	1	0.266473	4.39073	0.068
S4-S6	0.22921	1	0.22921	2.53538	0.198
M4-M6	0.138883	1	0.138883	3.15016	0.05*
S2-S6	0.137483	1	0.137483	1.29883	0.413
S5-M2	0.223533	1	0.223533	3.187	0.107
M2-M4	0.0199211	1	0.0199211	0.70151	0.589

M3-M5	0.0460803	1	0.0460803	1.26165	0.341
M1-M5	0.1314	1	0.1314	3.8839	0.013*
S7-M6	0.740479	1	0.740479	10.4001	0.02*
S6-M2	0.225729	1	0.225729	2.42344	0.184
S4-M6	0.183315	1	0.183315	5.21219	0.011*
S4-M1	0.0626517	1	0.0626517	2.86947	0.023*
S3-S4	0.0656918	1	0.0656918	2.8403	0.072
S2-S4	0.0574829	1	0.0574829	3.29319	0.066
S4-S7	0.436925	1	0.436925	8.27984	0.029
S5-M3	0.407831	1	0.407831	5.45696	0.017
S5-M1	0.270468	1	0.270468	3.73097	0.03*
M1-M2	0.0309953	1	0.0309953	1.26133	0.189
S4-S5	0.231663	1	0.231663	3.43716	0.019**
S3-M3	0.16078	1	0.16078	5.27755	0.005**
S5-S7	0.0721351	1	0.0721351	0.697439	0.438
S4-M2	0.0571673	1	0.0571673	2.93463	0.028*
M1-M3	0.115701	1	0.115701	3.96639	0.003**
S1-S5	0.163211	1	0.163211	2.29049	0.184
S4-M5	0.0948554	1	0.0948554	3.42189	0.034*
S7-M5	0.549234	1	0.549234	7.74059	0.04*
S2-S5	0.111242	1	0.111242	1.4217	0.269
S7-M4	0.436884	1	0.436884	7.08226	0.001***
S1-M4	0.0736667	1	0.0736667	2.49594	0.089

S7-M3	0.655788	1	0.655788	10.9105	0.007**
S3-M2	0.0690644	1	0.0690644	2.66988	0.034*
M3-M6	0.0679816	1	0.0679816	1.59931	0.201
S3-S7	0.418749	1	0.418749	7.07855	0.011*
M5-M6	0.10333	1	0.10333	2.07341	0.044*
S2-M1	0.0962802	1	0.0962802	4.08539	0.032*
M2-M6	0.201935	1	0.201935	5.3267	0.014*
S5-M4	0.245334	1	0.245334	3.21468	0.023*
S5-M5	0.348851	1	0.348851	3.94134	0.027*
S4-M4	0.0506137	1	0.0506137	1.97263	0.121
S6-M6	0.437969	1	0.437969	4.02419	<0.001*
S4-M3	0.108391	1	0.108391	4.50182	0.012*
S2-M2	0.0652358	1	0.0652358	3.14503	0.025*
S6-M1	0.272552	1	0.272552	2.85402	0.048*
S1-S3	0.0566845	1	0.0566845	2.10059	0.052*
M1-M6	0.221378	1	0.221378	5.49822	0.015*
S1-S4	0.0218245	1	0.0218245	1.05958	0.349
M2-M3	0.0991976	1	0.0991976	3.69911	0.033*
S2-M5	0.16442	1	0.16442	5.24333	0.105
S1-M5	0.123894	1	0.123894	3.83004	0.016*
S6-M3	0.381372	1	0.381372	3.90187	0.007**
S1-M6	0.217747	1	0.217747	5.57941	0.009**
S2-M6	0.264406	1	0.264406	6.68187	0.015*

S3-S6	0.219448	1	0.219448	2.26721	0.309
S1-M2	0.0740045	1	0.0740045	3.17115	0.014*
S6-M4	0.23653	1	0.23653	2.38145	0.211
S7-M1	0.485294	1	0.485294	8.38696	0.007**
S1-S2	0.0290423	1	0.0290423	1.31515	0.245
M1-M4	0.0441517	1	0.0441517	1.43577	0.138
M3-M4	0.0719655	1	0.0719655	2.18114	0.125
S1-S7	0.390742	1	0.390742	6.90035	0.006**
S5-S6	0.0159133	1	0.0159133	0.11281	0.878
S6-M5	0.325414	1	0.325414	2.80247	0.157
S3-S5	0.194684	1	0.194684	2.63844	0.07
S2-S3	0.0518463	1	0.0518463	2.06391	0.13
S3-M6	0.241142	1	0.241142	5.80253	0.023*
S1-M1	0.0850429	1	0.0850429	3.31029	0.016*
S7-M2	0.406276	1	0.406276	7.31907	0.032*
S3-M4	0.0837043	1	0.0837043	2.61203	0.001***
All samples	2.31435	12	0.192863	3.6472	<0.001***
S1-S2-S3-S4-S5-S6-S7-M1-M2-M3-M4-M5-M6					

991 **Supplementary Table 7.** Diazotrophs isolated from the sorghum mucilage.

Strain	Sorghum Genotype	Phylum/Class	Closest type strain	Length (bp)	Similarity (%)	Accession Number
1.1	IS_29091	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas lactitubi* M15^{T}	1495	99.78	NR_179509.1
1.2	IS_29091	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas lactitubi* M15^{T}	1320	100	NR_179509.1
1.3	IS_29091	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas lactitubi* M15^{T}	1333	100	NR_179509.1
1.6	IS_29091	*Pseudomonadota/Gammaproteobacteria*	*Klebisiella michiganensis* W14^{T}	1319	99.7	MT572941.1
1.7	IS_29091	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas lactitubi* M15^{T}	1377	99.49	NR_179509.1
1.9	IS_29091	*Pseudomonadota/Gammaproteobacteria*	*Klebisiella oxitoca* JCM 1665^{T}	1286	99.61	LC133345.1
2.1	IS_31706	*Pseudomonadota/Betaproteobacteria*	*Pseudacidovorax intermedius* CC-21	1298	100	NR_044241.1
2.2	IS_31706	*Actinomycetota/Actinomycetes*	*Microbacterium testaceum* DSM 20166^{T}	1310	100	NR_026163.1
2.5	IS_31706	*Actinomycetota/Actinomycetes*	*Microbacterium testaceum* DSM 20166^{T}	1355	99.78	NR_026163.1
2.6	IS_31706	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas rhizophila* e-p10^{T}	1356	99.93	NR_028930.1
3.1	IS_15744	*Pseudomonadota/Alphaproteobacteria*	*Novosphingobium resinovorum* NCIMB 8767^{T}	1236	100	NR_044045.1
3.2	IS_15744	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas lactitubi* M15^{T}	1353	99.78	NR_179509.1

3.3	IS_15744	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas lactitubi* M15[T]	1250	100	NR_179509.1
3.5	IS_15744	*Pseudomonadota/Alphaproteobacteria*	*Novosphingobium resinovorum* NCIMB 8767[T]	1220	100	NR_044045.1
3.6	IS_15744	*Actinomycetota/Actinomycetes*	*Microbacterium neimengense* 7087[T]	1330	99.55	NR_118272.1
4.2	IS_31706	*Actinomycetota/Actinomycetes*	*Microbacterium testaceum* DSM 20166[T]	1388	99.77	NR_026163.1
4.4	IS_31706	*Pseudomonadota/Gammaproteobacteria*	*Phytobacter diazotrophicus* DSM 17806[T]	1441	99.74	KY288669.1
4.6	IS_31706	*Pseudomonadota/Alphaproteobacteria*	*Sphingobium limneticum* 301[T]	1369	99.7	NR_109484.1
5.1	IS_31706	*Pseudomonadota/Alphaproteobacteria*	*Agrobacterium fabacearum* CNPSo 675[T]	1295	100	MN741112.1
5.2	IS_31706	*Pseudomonadota/Gammaproteobacteria*	*Pseudoxanthomonas winnipegensis* NML 130738[T]	1352	99.93	MH795530.1
5.3	IS_31706	*Bacteroidota/Flavobacteriia*	*Epilithonimonas hungarica* CHB-20p[T]	1265	99.84	NR_044354.1
5.4	IS_31706	*Pseudomonadota/Alphaproteobacteria*	*Sphingobium yanoikuyae* HAMBI 1842[T]	1317	96.82	LT899948.1
6.1	IS_15744	*Pseudomonadota/Alphaproteobacteria*	*Agrobacterium fabacearum* CNPSo 675[T]	1290	100	MN741112.1
6.11	IS_15744	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas nematodicola* W5[T]	1365	99.93	NR_181111.1

6.2	IS_15744	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas cyclobalanopsidis* TPQG1-4^T	1356	99.85	MN036524.2
6.3	IS_15744	*Pseudomonadota/Alphaproteobacteria*	*Novosphingobium kaempferiae* Sx8-5^T	1298	99.54	CP089301.1
6.4	IS_15744	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas cyclobalanopsidis* TPQG1-4^T	1356	99.7	MN036524.2
6.7	IS_15744	*Pseudomonadota/Alphaproteobacteria*	*Agrobacterium divergens* LMG 31531^T	1350	100	AM403584.1
6.8	IS_15744	*Pseudomonadota/Alphaproteobacteria*	*Agrobacterium fabacearum* CNPSo 675^T	1292	100	MN741112.1
6.9	IS_15744	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas nematodicola* W5^T	1377	99.85	NR_181111.1
7.2	IS_31706	*Actinomycetota/Actinomycetes*	*Microbacterium oleivorans* BAS69^T	1379	98.7	NR_042262.1
7.3	IS_31706	*Pseudomonadota/Gammaproteobacteria*	*Pseudomonas campi* S1-A32-2	1284	98.68	MT415401.1
7.6	IS_31706	*Bacillota/Bacilli*	*Bacillus amyloliquefaciens* DSM 7^T	1389	100	FN597644.1
7.8	IS_31706	*Actinomycetota/Actinomycetes*	*Microbacterium oleivorans* BAS69^T	1329	98.87	NR_042262.1
8.1	IS_29314	*Pseudomonadota/Alphaproteobacteria*	*Azospirillum humicireducens* SgZ-5^T	1181	96.21	CP015285.1
8.4	IS_29314	*Pseudomonadota/Gammaproteobacteria*	*Pseudomonas bharatica* CSV86^T	1331	99.77	MN866057.2
10.2	IS_29091	*Actinomycetota/Actinomycetes*	*Microbacterium neimengense* 7087^T	1291	99.84	NR_118272.1
10.3	IS_29091	*Bacteroidota/Chitinophagia*	*Filimonas endophytica* SR 2-06^T	1292	98.14	NR_145918.1
10.4	IS_29091	*Bacteroidota/Chitinophagia*	*Filimonas zeae* 772^T	1329	97.9	NR_149799.1
10.5	IS_29091	*Actinomycetota/Actinomycetes*	*Microbacterium testaceum* DSM 20166^T	1349	100	NR_026163.1

10.6	IS_29091	*Pseudomonadota/Alphaproteobacteria*	*Agrobacterium divergens* LMG 31531[T]	1365	100	AM403584.1
10.7	IS_29091	*Pseudomonadota/Alphaproteobacteria*	*Agrobacterium fabacearum* CNPSo 675[T]	1231	100	MN741112.1
11.1	IS_15744	*Actinomycetota/Actinomycetes*	*Microbacterium binotii* CIP 101303[T]	1330	100	NR_044290.1
11.1	IS_15744	*Actinomycetota/Actinomycetes*	*Microbacterium testaceum* DSM 20166[T]	1338	99.18	NR_026163.1
11.2	IS_15744	*Pseudomonadota/Alphaproteobacteria*	*Sphingobium yanoikuyae* HAMBI 1842[T]	1297	99.92	LT899948.1
11.4	IS_15744	*Pseudomonadota/Gammaproteobacteria*	*Pseudomonas sediminis* PI11[T]	1145	94.53	KP319033.1
11.7	IS_15744	*Pseudomonadota/Alphaproteobacteria*	*Azospirillum palustre* B2[T]	1232	99.35	NR_178321.1
11.8	IS_15744	*Pseudomonadota/Alphaproteobacteria*	*Sphingobium yanoikuyae* HAMBI 1842[T]	1311	100	LT899948.1
12.2	PI_534063	*Pseudomonadota/Alphaproteobacteria*	*Rhizobium cellulosilyticum* ALA10B2	1183	99.32	NR_043985.1
12.4	PI_534063	*Pseudomonadota/Betaproteobacteria*	*Herbaspirillum seropedicae* Z67[T]	1388	99.71	CP011930.1
12.6	PI_534063	*Actinomycetota/Actinomycetes*	*Microbacterium neimengense* 7087[T]	1378	99.78	NR_118272.1
12.7	PI_534063	*Pseudomonadota/Betaproteobacteria*	*Herbaspirillum seropedicae* Z67[T]	1341	99.85	CP011930.1
12.9	PI_534063	*Pseudomonadota/Betaproteobacteria*	*Herbaspirillum seropedicae* Z67[T]	1281	99.84	CP011930.1
13.1	IS_9745	*Actinomycetota/Actinomycetes*	*Microbacterium testaceum* DSM 20166[T]	1379	99.78	NR_026163.1
13.3	IS_9745	*Actinomycetota/Actinomycetes*	*Microbacterium hominis* DSM 12509[T]	1325	98.19	NR_042480.1

13.4	IS_9745	*Actinomycetota/Actinomycetes*	*Microbacterium hominis* DSM 12509^T	1360	99.76	NR_042480.1
13.5	IS_9745	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas terrae* R-32768	1361	99.34	NR_042569.1
13.7	IS_9745	*Actinomycetota/Actinomycetes*	*Microbacterium testaceum* DSM 20166^T	1305	99.16	NR_026163.1
13.8	IS_9745	*Actinomycetota/Actinomycetes*	*Microbacterium testaceum* DSM 20166^T	1322	99.17	NR_026163.1
13.9	IS_9745	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas lactitubi* M15^T	1339	99.78	NR_179509.1
14.1	IS_29091	*Actinomycetota/Actinomycetes*	*Microbacterium testaceum* DSM 20166^T	1270	98.98	NR_026163.1
14.2	IS_29091	*Pseudomonadota/Gammaproteobacteria*	*Pseudomonas turukhanskensis* IB1.1^T	1331	98.5	NR_152710.1
14.3	IS_29091	*Pseudomonadota/Gammaproteobacteria*	*Enterobacter cancerogenus* FDAARGOS 1428^T	1287	99.92	CP077290.1
14.4	IS_29091	*Actinomycetota/Actinomycetes*	*Microbacterium aerolatum* CCM 4955^T	1289	99.07	MT760116.1
14.5	IS_29091	*Actinomycetota/Actinomycetes*	*Microbacterium testaceum* DSM 20166^T	1373	99.05	NR_026163.1
14.6	IS_29091	*Actinomycetota/Actinomycetes*	*Microbacterium testaceum* DSM 20166^T	1355	99.12	NR_026163.1
14.7	IS_29091	*Pseudomonadota/Gammaproteobacteria*	*Pseudomonas turukhanskensis* IB1.1^T	1299	100	NR_152710.1

14.9	IS_29091	*Actinomycetota/Actinomycetes*	*Microbacterium testaceum* DSM 20166[T]	1315	100	NR_026163.1
15.1	IS_13971	*Actinomycetota/Actinomycetes*	*Microbacterium laevaniformans* DSM 20140[T]	1270	99.21	MN543870.1
15.2	IS_13971	*Pseudomonadota/Alphaproteobacteria*	*Novosphingobium resinovorum* NCIMB 8767[T]	1257	100	EF029110.2
15.5	IS_13971	*Pseudomonadota/Alphaproteobacteria*	*Devosia riboflavina* NBRC 13584[T]	1191	100	NR_113618.1
15.7	IS_13971	*Pseudomonadota/Alphaproteobacteria*	*Ketogulonicigenium vulgare* DSM 4025T	1259	100	DQ915606.1
15.8	IS_13971	*Pseudomonadota/Alphaproteobacteria*	*Devosia riboflavina* NBRC 13584[T]	1347	99.85	NR_113618.1
15.9	IS_13971	*Pseudomonadota/Betaproteobacteria*	*Pseudacidovorax intermedius* CC-21	1213	96.83	NR_044241.1
16.1	IS_15744	*Actinomycetota/Actinomycetes*	*Microbacterium oleivorans* BAS69[T]	1269	99.13	NR_042262.1
16.2	IS_15744	*Actinomycetota/Actinomycetes*	*Microbacterium oleivorans* BAS69[T]	1276	98.82	NR_042262.1
16.3	IS_15744	*Pseudomonadota/Alphaproteobacteria*	*Sphingobium yanoikuyae* HAMBI 1842[T]	1355	99.92	LT899948.1
16.4	IS_15744	*Actinomycetota/Actinomycetes*	*Microbacterium binotii* CIP 101303[T]	1319	100	NR_044290.1
16.5	IS_15744	*Actinomycetota/Actinomycetes*	*Microbacterium oleivorans* BAS69[T]	1377	98.77	NR_042262.1
16.7	IS_15744	*Actinomycetota/Actinomycetes*	*Microbacterium neimengense* 7087[T]	1370	99.89	NR_118272.1
17.2	IS_29091	*Pseudomonadota/Alphaproteobacteria*	*Agrobacterium divergens* LMG 31531[T]	1231	100	AM403584.1
17.3	IS_29091	*Pseudomonadota/Alphaproteobacteria*	*Novosphingobium kaempferiae* Sx8-5[T]	1193	99.92	CP089301.1

17.4	IS_29091	*Pseudomonadota/Gammaproteobacteria*	*Phytobacter diazotrophicus* DSM 17806^T	1288	99.74	KY288669.1
17.5	IS_29091	*Pseudomonadota/Alphaproteobacteria*	*Agrobacterium divergens* LMG 31531^T	1291	98.99	AM403584.1
17.6	IS_29091	*Pseudomonadota/Gammaproteobacteria*	*Pseudomonas turukhanskensis* IB1.1^T	1331	98.81	NR_152710.1
18.1	IS_31706	*Pseudomonadota/Gammaproteobacteria*	*Enterobacter cancerogenus* FDAARGOS 1428^T	1247	99.68	CP077290.1
18.2	IS_31706	*Pseudomonadota/Alphaproteobacteria*	*Agrobacterium larrymoorei* ATCC 51759^T	1230	100	MT780298.1
18.3	IS_31706	*Pseudomonadota/Alphaproteobacteria*	*Sphingobium yanoikuyae* HAMBI 1842^T	1320	100	LT899948.1
18.4	IS_31706	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas maltophilia* IAM 12423	1295	99.85	MN240936.1
18.6	IS_31706	*Pseudomonadota/Alphaproteobacteria*	*Novosphingobium kaempferiae* Sx8-5^T	1311	99.39	CP089301.1
18.7	IS_31706	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas lactitubi* M15^T	1404	99.52	NR_179509.1
18.8	IS_31706	*Pseudomonadota/Alphaproteobacteria*	*Azospirillum palustre* B2^T	1267	98.97	NR_178321.1
18.9	IS_31706	*Pseudomonadota/Alphaproteobacteria*	*Azospirillum humicireducens* SgZ-5^T	1298	99.03	CP015285.1
19.2	IS_31706	*Bacteroidota/Flavobacteriia*	*Epilithonimonas hungarica* CHB-20p^T	1339	99.71	NR_044354.1
19.4	IS_31706	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas cyclobalanopsidis* TPQG1-4^T	1403	99.86	MN036524.2
19.6	IS_31706	*Bacteroidota/Chitinophagia*	*Siphonobacter intestinalis* 63MJ-2	1335	99.1	NR_165683.1

19.7	IS_31706	*Pseudomonadota/Gammaproteobacteria*	*Klebsiella variicola* DSM 15968	1410	99.65	CP010523.2
20.1	IS_15744	*Pseudomonadota/Alphaproteobacteria*	*Sphingobium yanoikuyae* HAMBI 1842[T]	1311	100	LT899948.1
20.2	IS_15744	*Pseudomonadota/Alphaproteobacteria*	*Sphingobium yanoikuyae* HAMBI 1842[T]	1299	100	LT899948.1
20.3	IS_15744	*Actinomycetota/Actinomycetes*	*Microbacterium testaceum* DSM 20166[T]	1373	99.16	NR_026163.1
20.4	IS_15744	*Pseudomonadota/Alphaproteobacteria*	*Agrobacterium pusense* NRCPB10[T]	1291	99.92	NR_116874.1
21.1	IS_29091	*Pseudomonadota/Alphaproteobacteria*	*Shinella oryzae* Z-25[T]	1341	98.44	NR_029103.1
21.3	IS_29091	*Pseudomonadota/Gammaproteobacteria*	*Stenotrophomonas lactitubi* M15[T]	1293	99.61	NR_179509.1

994 **Supplementary Table 8.** Diazotrophs that have been used for biological nitrogen fixation

995 experiments. References to the origin of these strains and the source of isolation are provided.

Diazotroph	Phenotype	Reference	Source*
Azospirillum brasilense FP2 [#]	Fixer strain	(Pedrosa and Yates 1984)	Wheat, maize and forage grass (*Poa pratense*)
Azospirillum baldaniorum Sp 245[T#]	Fixer strain	(Baldani et al. 1983)	Wheat
Klebisiela variicola A3[#]	Fixer strain	This study	Sorghum mucilage
Klebisiela michiganensis A2 [#]	Fixer strain	This study	Maize mucilage
Azorhizobium caulinodans ORS571[T#]	Fixer strain	(Dreyfus and Dommergues 1981)	Nodules in *Sesbania rostrata*
Stutzerimonas stutzeri A1501 [#]	Fixer strain	(Qiu et al. 1981)	Rice
Paraburkholderia silvatlantica SRMrh-20[T#]	Fixer strain	(Perin et al. 2006)	Maize
Azotobacter vinelandii DJ[#]	Fixer strain	(Setubal et al. 2009)	Ubiquitous
Herbaspirillum seropedicae SmR1[#]	Fixer strain	(Pedrosa et al. 2011)	Maize, sorghum, rice, wheat and sugarcane
Azospirillum brasilense FP10	Non-fixer strain	(Pedrosa and Yates 1984)	Wheat, maize and forage grass (*Poa pratense*)
Azotobacter vinelandii DJ100 Δ*nifD*	Non-fixer strain	(Robinson et al. 1986)	Ubiquitous
Azorhizobium caulinodans ORS571 Δ*nifA*	Non-fixer strain	(Ryu et al. 2020)	Nodules in *Sesbania rostrata*

* Plant(s) where it has been isolated.

[#] For ARA experiment

References

1. Burt TP. Nitrogen Cycle☆. In: Fath B, editor. Encyclopedia of Ecology (Second Edition). Oxford: Elsevier; 2013. pp. 135–142.

2. Wagner SC. Biological Nitrogen Fixation. Nature Education Knowledge. 3, 15. 2011.

3. Zehr JP, Jenkins BD, Short SM, Steward GF. Nitrogenase gene diversity and microbial community structure: a cross-system comparison. Environ Microbiol. 2003;5: 539–554.

4. Oldroyd GED. Speak, friend, and enter: signalling systems that promote beneficial symbiotic associations in plants. Nat Rev Microbiol. 2013;11: 252–263.

5. Saikia SP, Jain V. Biological nitrogen fixation with non-legumes: An achievable target or a dogma? Curr Sci. 2007;92: 317–322.

6. Werner D, Newton WE. Nitrogen Fixation in Agriculture, Forestry, Ecology, and the Environment. Springer Netherlands; 2010.

7. Kumar A, Verma JP. The Role of Microbes to Improve Crop Productivity and Soil Health. In: Achal V, Mukherjee A, editors. Ecological Wisdom Inspired Restoration Engineering. Singapore: Springer Singapore; 2019. pp. 249–265.

8. Pankievicz VCS, Irving TB, Maia LGS, Ané J-M. Are we there yet? The long walk towards the development of efficient symbiotic associations between nitrogen-fixing bacteria and non-leguminous crops. BMC Biol. 2019;17: 99.

9. Oldroyd GED, Dixon R. Biotechnological solutions to the nitrogen problem. Curr Opin Biotechnol. 2014;26: 19–24.

10. Zhang L, Zhang M, Huang S, Li L, Gao Q, Wang Y, *et al*. A highly conserved core bacterial microbiota with nitrogen-fixation capacity inhabits the xylem sap in maize plants. Nat Commun. 2022;13: 3361.

11. Mechan-Llontop ME, Mullet J, Shade A. Phyllosphere Exudates Select for Distinct Microbiome Members in Sorghum Epicuticular Wax and Aerial Root Mucilage. Phytobiomes Journal. 2023;7: 184–197.

12. Chakraborty S, Venkataraman M, Infante V, Pfleger BF, Ané J-M. Scripting a new dialogue between diazotrophs and crops. Trends Microbiol. 2023. doi:10.1016/j.tim.2023.08.007

13. Van Deynze A, Zamora P, Delaux P-M, Heitmann C, Jayaraman D, Rajasekar S, *et al*. Nitrogen fixation in a landrace of maize is supported by a mucilage-associated diazotrophic microbiota. PLoS Biol. 2018;16: e2006352.

14. Amicucci MJ, Galermo AG, Guerrero A, Treves G, Nandita E, Kailemia MJ, *et al*. Strategy for Structural Elucidation of Polysaccharides: Elucidation of a Maize Mucilage that Harbors

Diazotrophic Bacteria. Anal Chem. 2019;91: 7254–7265.

15. Pankievicz VCS, Delaux P-M, Infante V, Hirsch HH, Rajasekar S, Zamora P, *et al*.
Nitrogen fixation and mucilage production on maize aerial roots is controlled by aerial root
development and border cell functions. Front Plant Sci. 2022;13: 977056.

16. Bennett AB, Pankievicz VCS, Ané J-M. A Model for Nitrogen Fixation in Cereal Crops.
Trends Plant Sci. 2020;25: 226–235.

17. Van Gelder K, Oliveira-Filho ER, Messina CD, Venado RE, Wilker J, Rajasekar S, *et al*.
Running the numbers on plant synthetic biology solutions to global problems. Plant Sci.
2023;335: 111815.

18. Hostetler AN, Khangura RS, Dilkes BP, Sparks EE. Bracing for sustainable agriculture: the
development and function of brace roots in members of Poaceae. Curr Opin Plant Biol.
2021;59: 101985.

19. Sparks EE. Maize plants and the brace roots that support them. New Phytol. 2023;237: 48–
52.

20. Mullet J, Morishige D, McCormick R, Truong S, Hilley J, McKinley B, *et a*l. Energy
Sorghum—a genetic model for the design of C4 grass bioenergy crops. J Exp Bot. 2014;65:
3479–3489.

21. Silva TN, Vermerris W. High-Biomass Sorghums as a Feedstock for Renewable Fuels and
Chemicals. In: Tonapi VA, Talwar HS, Are AK, Bhat BV, Reddy CR, Dalton TJ, editors.
Sorghum in the 21st Century: Food – Fodder – Feed – Fuel for a Rapidly Changing World.
Singapore: Springer Singapore; 2020. pp. 723–754.

22. Regassa TH, Wortmann CS. Sweet sorghum as a bioenergy crop: Literature review.
Biomass Bioenergy. 2014;64: 348–355.

23. Rao PS, Vinutha KS, Kumar GSA, Chiranjeevi T, Uma A, Lal P, *et al*. Sorghum: A
multipurpose bioenergy crop. Sorghum. Madison, WI, USA: Soil Science Society of
America; 2019. pp. 399–424.

24. Wani SP. Cereal nitrogen fixation: proceedings of the Working Group Meeting held at
ICRISAT Center, India, 9-12 October 1984. International Crops Research Institute for the
Semi-Arid Tropics; 1986.

25. Upadhyaya HD, Pundir RPS, Dwivedi SL, Gowda CLL, Reddy VG, Singh S. Developing a
mini core collection of sorghum for diversified utilization of germplasm. Crop Sci. 2009;49:
1769–1780.

26. Ford MA, Thorne GN. Effects of Atmospheric Humidity on Plant Growth. Ann Bot.
1974;38: 441–452.

27. Hernández-López A, Sánchez Félix DA, Zuñiga Sierra Z, García Bravo I, Dinkova TD,

Avila-Alejandre AX. Quantification of Reducing Sugars Based on the Qualitative Technique of Benedict. ACS Omega. 2020;5: 32403–32410.

28. Wolf ESA, Vela S, Wilker J, Davis A, Robert M, Infante V, *et al*. Identification of genetic and environmental factors influencing aerial root traits that support biological nitrogen fixation in sorghum. G3 . 2023. doi:10.1093/g3journal/jkad285

29. Hawes MC, Lin HJ. Correlation of Pectolytic Enzyme Activity with the Programmed Release of Cells from Root Caps of Pea (Pisum sativum). Plant Physiol. 1990;94: 1855–1859.

30. Brigham LA, Woo HH, Nicoll SM, Hawes MC. Differential Expression of Proteins and mRNAs from Border Cells and Root Tips of Pea. Plant Physiol. 1995;109: 457–463.

31. Knee EM, Gong FC, Gao M, Teplitski M, Jones AR, Foxworthy A, *et al.* Root mucilage from pea and its utilization by rhizosphere bacteria as a sole carbon source. Mol Plant Microbe Interact. 2001;14: 775–784.

32. Hawes MC, Pueppke SG. Sloughed peripheral root cap cells: Yield from different species and callus formation from single cells. Am J Bot. 1986;73: 1466–1473.

33. Hawes MC, Pueppke SG. Correlation between binding of *Agrobacterium tumefaciens* by root cap cells and susceptibility of plants to crown gall. Plant Cell Rep. 1987;6: 287–290.

34. Woo H-H, Hirsch AM, Hawes MC. Altered susceptibility to infection by *Sinorhizobium meliloti* and *Nectria haematococca* in alfalfa roots with altered cell cycle. Plant Cell Rep. 2004;22: 967–973.

35. Zhang Y, Chen B, Gao S, Rong T. Biological characters of root border cells development in maize (*Zea mays*). Biotechnology. 2014;13: 89–98.

36. Vicré M, Santaella C, Blanchet S, Gateau A, Driouich A. Root border-like cells of Arabidopsis. Microscopical characterization and role in the interaction with rhizobacteria. Plant Physiol. 2005;138: 998–1008.

37. Boulos L, Prévost M, Barbeau B, Coallier J, Desjardins R. LIVE/DEAD® BacLight^{TM}: application of a new rapid staining method for direct enumeration of viable and total bacteria in drinking water. J Microbiol Methods. 1999;37: 77–86.

38. Ullal AJ, Pisetsky DS, Reich CF 3rd. Use of SYTO 13, a fluorescent dye binding nucleic acids, for the detection of microparticles in in vitro systems. Cytometry A. 2010;77: 294–301.

39. Casa AM, Pressoir G, Brown PJ, Mitchell SE, Rooney WL, Tuinstra MR, *et al*. Community resources and strategies for association mapping in sorghum. Crop Sci. 2008;48: 30–40.

40. He Y, Caporaso JG, Jiang X-T, Sheng H-F, Huse SM, Rideout JR, *et al*. Stability of operational taxonomic units: an important but neglected property for analyzing microbial

diversity. Microbiome. 2015;3: 20.

41. Pedrosa FO, Yates MG. Regulation of nitrogen fixation (*nif*) genes of *Azospirillum*
brasilense by *nifA* and *ntr* (*gln*) type gene products. FEMS Microbiol Lett. 1984;23: 95–
101.

42. Venkataraman M, Yñigez-Gutierrez A, Infante V, MacIntyre A, Fernandes-Júnior PI, Ané
J-M, *et al*. Synthetic Biology Toolbox for Nitrogen-Fixing Soil Microbes. ACS Synth Biol.
2023;12: 3623–3634.

43. Li R, Han Y, Lv P, Du R, Liu G. Molecular mapping of the brace root traits in sorghum
(Sorghum bicolor L. Moench). Breed Sci. 2014;64: 193–198.

44. Arve LE, Torre S. Ethylene is involved in high air humidity promoted stomatal opening of
tomato (*Lycopersicon esculentum*) leaves. Funct Plant Biol. 2015;42: 376–386.

45. McDonald MP, Visser EJW. A Study of the Interaction between Auxin and Ethylene in
Wild Type and Transgenic Ethylene-Insensitive Tobacco during Adventitious Root
Formation Induced by Stagnant Root Zone Conditions. Plant Biol . 2003;5: 550–556.

46. Visser EJW, Voesenek LACJ. Acclimation to Soil Flooding–sensing and signal-
transduction. Plant Soil. 2005;274: 197–214.

47. Chhun T, Uno Y, Taketa S, Azuma T, Ichii M, Okamoto T, *et al*. Saturated humidity
accelerates lateral root development in rice (Oryza sativa L.) seedlings by increasing
phloem-based auxin transport. J Exp Bot. 2007;58: 1695–1704.

48. Kavi Kishor PB, Tiozon RN Jr, Fernie AR, Sreenivasulu N. Abscisic acid and its role in the
modulation of plant growth, development, and yield stability. Trends Plant Sci. 2022;27:
1283–1295.

49. Xu S, Li X-Q, Guo H, Wu X-Y, Wang N, Liu Z-Q, *et al*. Mucilage secretion by aerial roots
in sorghum (Sorghum bicolor): sugar profile, genetic diversity, GWAS and transcriptomic
analysis. Plant Mol Biol. 2023. doi:10.1007/s11103-023-01365-1

50. Osborn HMI, Lochey F, Mosley L, Read D. Analysis of polysaccharides and
monosaccharides in the root mucilage of maize (*Zea mays* L.) by gas chromatography. J
Chromatogr A. 1999;831: 267–276.

51. Nazari M, Riebeling S, Banfield CC, Akale A, Crosta M, Mason-Jones K, *et al*. Mucilage
Polysaccharide Composition and Exudation in Maize From Contrasting Climatic Regions.
Front Plant Sci. 2020;11: 587610.

52. Bouffaud M-L, Poirier M-A, Muller D, Moënne-Loccoz Y. Root microbiome relates to
plant host evolution in maize and other Poaceae: Poaceae evolution and root bacteria.
Environ Microbiol. 2014;16: 2804–2814.

53. Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, *et al*. Defining

1136   the core Arabidopsis thaliana root microbiome. Nature. 2012;488: 86–90.

1137 54. Edwards J, Johnson C, Santos-Medellín C, Lurie E, Podishetty NK, Bhatnagar S, *et al*.
1138   Structure, variation, and assembly of the root-associated microbiomes of rice. Proc Natl
1139   Acad Sci U S A. 2015;112: E911–20.

1140 55. Hartman K, van der Heijden MG, Roussely-Provent V, Walser J-C, Schlaeppi K.
1141   Deciphering composition and function of the root microbiome of a legume plant.
1142   Microbiome. 2017;5: 2.

1143 56. Crosbie DB, Mahmoudi M, Radl V, Brachmann A, Schloter M, Kemen E, *et al*.
1144   Microbiome profiling reveals that Pseudomonas antagonises parasitic nodule colonisation
1145   of cheater rhizobia in Lotus. New Phytol. 2022;234: 242–255.

1146 57. Xu L, Naylor D, Dong Z, Simmons T, Pierroz G, Hixson KK, *et al*. Drought delays
1147   development of the sorghum root microbiome and enriches for monoderm bacteria. Proc
1148   Natl Acad Sci U S A. 2018;115: E4284–E4293.

1149 58. Wong PP, Stenberg NE. Characterization of Azospirillum isolated from nitrogen-fixing
1150   roots of harvested sorghum plants. Appl Environ Microbiol. 1979;38: 1189–1191.

1151 59. Pal UR, Malik HS. Contribution of *Azospirillum brasilense* to the nitrogen needs of grain
1152   sorghum (*Sorghum bicolor* (L) Moench) in humid sub-tropics. Plant Soil. 1981;63: 501–
1153   504.

1154 60. Morgenstern E, Okon Y. Promotion of plant growth and NO_3^- and Rb^+ uptake in *Sorghum*
1155   *bicolor* × *Sorghum sudanense* inoculated with *Azospirillum brasilense*-Cd. Arid Soil
1156   Research and Rehabilitation. 1987;1: 211–217.

1157 61. dos Santos CLR, Alves GC, de Matos Macedo AV, Giori FG, Pereira W, Urquiaga S, *et al*.
1158   Contribution of a mixed inoculant containing strains of *Burkholderia spp*. and
1159   *Herbaspirillum ssp*. to the growth of three sorghum genotypes under increased nitrogen
1160   fertilization levels. Appl Soil Ecol. 2017;113: 96–106.

1161 62. Barros BG de F, Freitas ADS de, Tabosa JN, Lyra M do CCP de, Mergulhão AC do ES,
1162   Silva AF da, *et al*. Biological nitrogen fixation in field-grown sorghum under different
1163   edaphoclimatic conditions is confirmed by N isotopic signatures. Nutr Cycling Agroecosyst.
1164   2020;117: 93–101.

1165 63. Baldani JI, Reis VM, Videira SS, Boddey LH, Baldani VLD. The art of isolating nitrogen-
1166   fixing bacteria from non-leguminous plants using N-free semi-solid media: a practical guide
1167   for microbiologists. Plant Soil. 2014;384: 413–431.

1168 64. Hara S, Morikawa T, Wasai S, Kasahara Y, Koshiba T, Yamazaki K, *et al*. Identification of
1169   Nitrogen-Fixing Bradyrhizobium Associated With Roots of Field-Grown Sorghum by
1170   Metagenome and Proteome Analyses. Front Microbiol. 2019;10: 407.

1171 65. Gao J, Feng P, Zhang J, Dong C, Wang Z, Chen M, *et al*. Enhancing Maize's Nitrogen-

Fixing Potential through Zm*SBT3*, a Gene Suppressing Mucilage Secretion. J Integr Plant
Biol. 2023. doi:10.1111/jipb.13581

66. Nazari M, Bickel S, Kuzyakov Y, Bilyera N, Zarebanadkouki M, Wassermann B, *et al*.
Root mucilage nitrogen for rhizosphere microorganisms under drought. Biol Fertil Soils.
2024. doi:10.1007/s00374-024-01827-8

67. Ryker TC. Studies on sugar cane roots. 1931 [cited 15 May 2024]. Available:
https://repository.lsu.edu/agexp/104/?utm_source=repository.lsu.edu%2Fagexp%2F104&ut
m_medium=PDF&utm_campaign=PDFCoverPages

68. Allen RS, Tilbrook K, Warden AC, Campbell PC, Rolland V, Singh SP, *et al*. Expression of
16 Nitrogenase Proteins within the Plant Mitochondrial Matrix. Front Plant Sci. 2017;8:
287.

69. Xiang N, Guo C, Liu J, Xu H, Dixon R, Yang J, *et al*. Using synthetic biology to overcome
barriers to stable expression of nitrogenase in eukaryotic organelles. Proc Natl Acad Sci U S
A. 2020;117: 16537–16545.

70. Vervliet G, Holsters M, Teuchy H, Van Montagu M, Schell J. Characterization of different
plaque-forming and defective temperate phages in Agrobacterium. J Gen Virol. 1975;26:
33–48.

71. Toukdarian A, Kennedy C. Regulation of nitrogen metabolism in *Azotobacter vinelandii*:
isolation of ntr and glnA genes and construction of ntr mutants. EMBO J. 1986;5: 399–407.

72. Xia J, Psychogios N, Young N, Wishart DS. MetaboAnalyst: a web server for metabolomic
data analysis and interpretation. Nucleic Acids Res. 2009;37: W652–60.

73. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, *et al*. Fiji: an
open-source platform for biological-image analysis. Nat Methods. 2012;9: 676–682.

74. Dorn-In S, Bassitta R, Schwaiger K, Bauer J, Hölzel CS. Specific amplification of bacterial
DNA by optimized so-called universal bacterial primers in samples rich of plant DNA. J
Microbiol Methods. 2015;113: 50–56.

75. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, *et al*.
Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2.
Nat Biotechnol. 2019;37: 852–857.

76. Wickham H. Getting Started with ggplot2. In: Wickham H, editor. ggplot2: Elegant
Graphics for Data Analysis. Cham: Springer International Publishing; 2016. pp. 11–31.

77. Estrada-De Los Santos P, Bustillos-Cristales R, Caballero-Mellado J. *Burkholderia*, a genus
rich in plant-associated nitrogen fixers with wide environmental and geographic
distribution. Appl Environ Microbiol. 2001;67: 2790–2798.

78. Versalovic J, Schneider M, De Bruijn FJ, Lupski JR. Genomic fingerprinting of bacteria

using repetitive sequence-based polymerase chain reaction. Methods Mol Cell Biol. 1994;5: 25–40.

79. Heuer H, Krsek M, Baker P, Smalla K, Wellington EM. Analysis of actinomycete communities by specific amplification of genes encoding 16S rRNA and gel-electrophoretic separation in denaturing gradients. Appl Environ Microbiol. 1997;63: 3233–3241.

80. Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic acids symposium series. Oxford; 1999. pp. 95–98.

81. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol. 1990;215: 403–410.

82. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol. 2018;35: 1547–1549.

83. Kahn ML, Parra-Colmenares A, Ford CL, Kaser F, McCaskill D, Ketchum RE. A mass spectrometry method for measuring ^{15}N incorporation into pheophytin. Anal Biochem. 2002;307: 219–225.

84. Foundation for Statistical Computing RR. R: A language and environment for statistical computing. RA Lang Environ Stat Comput.

85. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, *et al*. QIIME allows analysis of high-throughput community sequencing data. Nat Methods. 2010;7: 335–336.

86. Ligges U, Maechler M. Scatterplot3d - an R package for visualizing multivariate data. J Stat Softw. 2003;008: 1–20.

87. Revelle W, Others. psych: Procedures for psychological, psychometric, and personality research. R package version. 2018;1: 195.