Costs of being a diet generalist in the super-generalist protist predator *Dictyostelium*

discoideum

P. M. Shreenidhi *, Debra A. Brock, Rachel I. McCabe, Joan E. Strassmann and David C. Queller

Department of Biology, Washington University in St. Louis, St. Louis, MO 63130

Corresponding author: *P. M. Shreenidhi

Email: shreenidhi@wustl.edu

Author Contributions: P.M.S., J.E.S., and D.C.Q. designed research. P.M.S., D.A.B., and R.I.M.

performed the experiments. P.M.S., J.E.S., and D.C.Q. analyzed data. P.M.S., J.E.S., and D.C.Q.

wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Classification: Major - Biological Sciences Minor- Ecology

Keywords: Diet breadth, Generalist, Predation, Protists, Switching costs

This PDF file includes:

Main Text
Figures 1 to 6
Tables 1
Supplementary tables
18

Abstract

Organisms can be classified as diet specialists or generalists based on the number of prey types

they eat. Since different prey species have different constitutions and defenses, eating a wide

variety of prey presents challenges. We explore three potential costs of generalism in a super-

generalist amoeba predator, *Dictyostelium discoideum*, feeding on naturally co-occuring

bacterial prey. First, we found that amoebas experience sub-optimal growth on many of their

prey bacteria. Second, amoebas also exhibit a reduction in growth rate when they switch from

one species of prey bacteria to another, something we call switching costs. These switching

costs disappear within a day, indicating adjustment to new prey bacteria. Third, amoebas usually

divide more slowly on mixtures of bacteria compared to their average on single bacteria,

something we call concurrent costs. Our results support that idea that, although amoebas can

consume a huge variety of prey, they must use partially different methods and thus must pay

costs to handle multiple prey, either sequentially or simultaneously.

Significance Statement

Perhaps the most fundamental conflict in nature involves consumption of one organism by

another. Diet generalists benefit from the advantage of eating many prey species but deal with

many prey defences. We explore costs associated with a broad diet in a protist microbial

predator, *Dictyostelium discoideum*. These predators of bacteria show a delay in growth when

switched from one prey to another, supporting the hypothesis that they must deploy different

strategies. They also experience costs when grown on many bacteria at once, suggesting that

the alternative strategies for different prey are partly incompatible with each other. Our findings

shed light on the nature of diet generalism and highlight the complexity of predation in the

microbial world, where most organisms reside.

Main Text

Introduction

Organisms vary widely in their diet breadth. Some are diet specialists that eat one or few prey

types, such as koalas that feed only on eucalyptus leaves (1) or snail kites that exclusively hunt

apple snails (2). Some organisms are diet generalists that eat many different prey, such as

coyotes that feed on many small mammals (3) or spiders that feed on many species of

arthropods (4, 5). Diet generalists benefit from the ability to exploit diverse prey species,

especially when prey species are scarce and fluctuate in their availability (6). But they must deal

with prey that often differ in nutritional quality and defensive mechanisms. Efficiently handling

these diverse prey is potentially hindered by trade-offs, which could select for diet specialization

(7, 8). Have generalist predators overcome costs associated with consuming multiple prey

species or do they experience costly trade-offs?

Many macroscopic predators consume a formidable number of prey species, but few

can match the diversity of prey consumed by some microbial protist predators (9, 10). Predation

by protists is a major factor accounting for bacterial mortality in the environment and plays an

important role in nutrient cycling (9, 11, 12). This can determine the composition and properties

of bacterial communities (13). Predation by protists is also an important selective pressure for

bacterial defenses such as biofilm formation, antibiotic production, and secretion systems (14,

15).

Dictyostelium discoideum is one such super-generalist protist predator. *D. discoideum* is

a social amoeba that lives in forest soils. It is a unicellular amoeba when bacterial prey are

abundant and transitions to a non-feeding multicellular dispersal stage upon starvation. *D.*

discoideum has been the subject of extensive research for this fascinating multicellular stage

(16, 17). Its feeding behavior has been less extensively studied, but it appears to qualify as a

super-generalist predator. It can eat the majority of bacteria it is presented with (18–20). For

example one study tested 159 bacterial strains found in close association with fruiting bodies of

D. discoideum from forest soil habitats and found that the amoebas were able to consume 77%

of them (20). Given the sheer diversity of bacteria in the soil – estimated to be as high as a

million species in 30 g of forest soil (21) – the number of edible species must be very large.

Specializing on one or few species of bacteria is probably not a viable option for *D.*

discoideum. This is because soil bacterial communities are spatially and temporally variable,

such that no one bacterium may be consistently sufficiently abundant (22). Indeed, even as

generalists starvation must occur often enough for it to evolve as the trigger for their

multicellular dispersal stage. Additionally, amoebas can travel considerable distances during

their social cycle through slug migration and especially through the dispersal of spores by animal

vectors (23). This can introduce them to new soil environments with different bacterial

communities. Thus, environmental heterogeneity likely plays a large role in the maintenance of

a generalist strategy in *D. discoideum*.

But handling a highly diverse array of prey bacteria is likely to be challenging for *D.*

discoideum. These prey bacteria are not identical and can range across four highly divergent

bacterial phyla: Actinobacteria, Bacteroidetes, Firmicutes and Proteobacteria (19). These shared

a common ancestor ~3000 million years ago, far older than the divergence time of diverse insect

prey (~400 mya) that generalist vertebrates feed on (24). Bacteria also possess highly varied

defensive mechanisms against microbial predators (14, 15). They can produce many kinds of

secondary metabolite toxins to repel, disable, or kill their enemies. They can use secretion

systems and effectors that can kill or allow for intracellular survival within protists. Some

bacteria can swim away at high speeds to escape predators. Some can group together to form

biofilms to prevent ingestion by predators.

We investigated three potential costs of *D. discoideum*'s broad diet. Each could reduce

the obvious benefit of being able to eat many species of prey bacteria. First, it is likely that not

all bacteria are equally profitable. The bacteria can differ in their nutritional quality and/or in

their susceptibility to the amoeba's hunting and processing techniques. Therefore, the first step

to understanding generalism of *D. discoideum* is to assess its performance on diverse prey.

Second, although generalist amoebas benefit from the ability to switch to a new prey

species when prey abundance changes, prey switching could be costly. If amoebas use different

methods to hunt and consume different prey, then changing between these methods can result

in temporary costs. These methods may be related to prey detection, foraging, detoxification,

and processing (25, 26) and may involve changes from the transcriptional level on up. Amoebas

do express partially different transcriptomes on different bacteria (27, 28). We refer to this

reduction in foraging efficiency as a switching cost.

Finally, in spatially fine-grained soil habitats, amoebas that consume multiple prey at

once could experience costs if specific techniques work best for different prey. It may be

impossible to effectively deploy multiple optimal strategies at the same time. We refer to the

reduction in foraging efficiency in the presence of multiple prey species as concurrent costs.

Alternatively, amoebas could benefit in this scenario due to improved nutritional balance. Some

studies have found an increase in protist growth rates in the presence of multiple prey species,

where mixed prey communities represent a balanced diet and can better fulfill the nutritional

requirements of the predator (29, 30).

Results

Growth rate of *D. discoideum* amoebas varies on different species of bacteria.

We measured the growth rate of *D. discoideum* on 22 species of soil bacteria and the commonly

used food bacterium *K. pneumoniae*. *D. discoideum* amoebas showed wide variation in their

doubling times on different soil bacteria (Test bacterium: $F_{22,45}$ = 23.88, p-value < 2.2×10^{-16}) (Fig

1). They doubled within four hours on some of the most edible soil bacteria and took longer

than nine hours on some of the least edible soil bacteria.

D. discoideum **amoebas experience switching costs on all tested bacteria pairs.**

We conducted switching cost experiments to investigate how *D. discoideum* amoebas

proliferate on a given bacterium when previously conditioned to a different bacterium (Fig 2). As

expected given previously observed variation in growth rate of *D. discoideum* on different

species of bacteria (Fig 1), the total number of amoebas at 3 hours for both switches and

controls depended on the test bacterium (Test bacterium: $F_{5,106}$ = 28.39, p-value < 2×10^{-16}). But

independent of this effect, there were also switching costs; we found that the total number of

amoebas at 3 hours from transfer significantly differed between the control and switches

(Treatment: $F_{1,106}$ = 27.621, p-value = 7.69×10^{-7}). The switched amoebas performed poorly on a

given bacterium compared to controls that did not experience a prey switch (Fig 3). There was

no evidence that bacteria differed in switching costs as we found no significant interaction

between bacterial species and Treatment (Treatment x Test bacterium: $F_{1,106}$= 0.838, p-value =

0.526). There was no evidence that conditioning time of amoebas (2 days or 5 days) affected

switching costs (Treatment x Conditioning length: $F_{1,106}$ = 0.543, p-value = 0.463). This is

inconsistent with switching costs being due to evolutionary changes during conditioning since

evolutionary changes should be greater over 5 days than 2.

***D. discoideum* do not experience switching costs if they undergo spore formation before the**

prey switch

If switching costs are because of an evolved response in amoebas, then these genetic changes

should be passed on through the spores. But if the costs are physiological, then undergoing the

social cycle will likely erase the effects of food conditioning and therefore eliminate switching

costs. We therefore tested if *D. discoideum* still show switching costs when they undergo the

social cycle and spore formation before changing between prey bacteria. The switched and

control spores resulted in a similar number of amoebas (Fig 4, Treatment: $F_{1,48}$ = 0.263, p-value =

0.61). Thus, we find no support for the hypothesis that switching costs are due to an evolved

response. But as expected, the total number of amoebas depends on the test bacterium the

spores were grown on (Test bacterium: $F_{5,48}$ = 186.4, p-value < $2x10^{-16}$).

Switching costs are temporary

If switching costs are not due to an evolved response, as supported by our previous results, then

the switched amoebas should eventually recover their growth rates to the level of the controls

that did not switch to new food bacteria. We conducted a time course experiment to test this

for one pair of bacteria (*A. tumefaciens*, *P. vranovensis*) and compared early (zero to six hours)

and late (24 to 27 hours) growth rates. The switched treatments had significantly lower early

growth rates compared to controls (Fig 5A, Treatment: $F_{1,17}$ = 9.458, p – value =0.007), as

expected from our earlier experiments. But there was no difference in late growth rate of

switches and controls at the later time points (Fig 5B, Treatment: $F_{1,17}$ = 0.111, p-value = 0.73).

Thus, though switched amoebas proliferate slower than controls during early growth, they

eventually catch up and perform similarly to controls, consistent with physiological adjustment

to the new prey.

***D. discoideum* amoebas experience concurrent costs in some multi-prey communities**

We tested if *D. discoideum* experiences costs when grown in multi-species prey communities

compared to expectations from their growth in single-species communities. We refer to these

costs as concurrent costs. Consistent with concurrent costs, we observed significantly fewer

amoebas than expected in multi-species communities (Fig 6, Treatment: $F_{1,59}$=11.83, p-value =

0.001). We also found that total number of amoebas depended on which bacterial community it

was plated with (Bacterial community: $F_{5,59}$ = 8.74, p-value = 3.01x 10^{-6}). The interaction

between treatment and bacterial community was also significant, indicating that the identity of

the bacterial community influenced the degree of concurrent costs (Treatment x Bacterial

Community: $F_{5,59}$ = 2.557 p-value = 0.036). The general trend is in the expected direction in five

out of the six bacterial communities with one community trending in the opposite direction.

Discussion

Predators and prey engage in the most ancient of battles. Though largely unseen and understudied, the microbial stage for this conflict may be the largest of all. Here we have examined a key feature of this struggle: costs and benefits of specializing on few prey as compared to embracing a catholic diet. The eukaryote *D. discoideum* lives in heterogenous soil environments where it encounters many species of prey bacteria. In such a variable environment with shifting prey abundance, the benefit of being a diet generalist is obvious. The cost to generalism is the struggle to defeat a wide variety of defense mechanism as prey work to defend themselves. We found that amoebas experience varying growth rates when presented with the same quantity of different prey bacteria (Fig 2). We opted to equalize bacterial quantity across different prey by using optical density. Bacteria can differ in how they influence optical density, especially through their varying cell sizes (31). But optical density is a much more robust measure of dry weight (31) which is probably a more relevant measure of nutritional potential of bacterial prey. Previous studies of *E. coli* show that different strains vary substantially in nutritional quality to *D. discoideum*, and this result holds whether the bacteria are measured by OD (32) or by cell number (33). Thus, we have evidence that bacterial prey differ in nutritional quality for the amoebas. This is perhaps not surprising, especially given that there are also some bacteria that *D. discoideum* cannot grow on at all (20).

Amoebas experience switching costs when moved from one prey bacterium to another, even after controlling for variation in bacterial edibility (Fig 3). We did three experiments to test if these costs were physiological in nature or due to an evolved response. First, we found no evidence that the conditioning time (duration spent by amoebas on a given bacterium before the prey switch) affected the magnitude switching costs. If the costs were due to evolution

during the conditioning period, they should become stronger with longer conditioning time.

Second, we found no evidence that amoebas experience switching costs if they undergo spore

formation before prey switch. If this was due to an evolved response, then these effects would

be genetically passed on through the spores. Third, we found that switching costs occur during

early growth but disappear within a handful of generations, faster than they would change by

selection. Taken together these experiments show that these costs are physiological in nature,

likely to be associated with the need to alter gene expression on different prey bacteria, and not

due to evolution.

We also found that *D. discoideum* amoebas experience concurrent costs, where they

proliferate less than expected in communities with multiple species of prey bacteria. It is

interesting that we observe these costs despite two factors that may obscure them. First, some

mixed bacterial prey communities may provide a more balanced diet that could increase growth

rate of *D. discoideum*. Some studies on other protists that have looked at growth-rate in multi-

prey communities have found an increase in growth rates (29, 30). Second, the amoebas may

also choose to eat the most profitable bacteria first. All but late measurements would reflect the

growth rate of the best bacterium and any concurrent costs might therefore be obscured.

Amoebas do show preferential attraction towards Gram-negative bacteria compared to Gram-

positive bacteria (34). Thus, it would be interesting to test if *D. discoideum* amoebas can avoid

some concurrent costs by preferentially eating the most profitable prey bacteria first, but if they

do, our data show that it is apparently not enough to fully overcome such costs.

Both switching and concurrent costs are consistent with the idea that *D. discoideum*

amoebas use partially different methods to hunt and process different prey. Previous studies

have shown that *D. discoideum* amoebas transcribe partially distinct sets of genes on Gram-

positive (*B. subtilis*, *S. aureus*) and Gram-negative bacteria (*K. pneumoniae*, *P. aeruginosa*) (27).

The transcriptome of *D. discoideum* was also found to be highly species-specific when tested on

3 species of bacteria: *K. pneumoniae*, *B. subtlis*, *M. luteus* (28). These variable transcriptional

responses suggest that there are different mechanisms for hunting or processing these distantly

related bacteria. Another study found that mutations that affect the ability of *D. discoideum* to

grow on different bacteria were highly prey-specific (35).

Predation by amoebas is a complex process that can be divided into 4 broad steps:

search, encounter, attack, and digestion (14). Both switching and concurrent costs could arise in

any of these steps. Eukaryotic phagocytes use G-protein coupled receptors (GPCRs) to detect

and chase bacteria and use pattern recognition receptors to recognize and eliminate bacteria

(36, 37). *D. discoideum* is equipped with 61 GPCRs (38). Some GPCRs are involved in the

amoebas' social cycle, but at least one GPCR, the folate receptor fAR1, has been implicated in

both chasing and engulfing *Klebsiella* (39). Other GPCRs may play a role chasing and recognizing

different species of bacteria. *D. discoideum* is also equipped with 22 genes that may encode 4

different types of lysozymes (40). Lysozymes play an important role in attacking and digesting

bacteria. Knocking out some lysozyme genes generally reduces the ability of amoebas to feed on

all tested bacteria, while deletion of others reduces growth only on specific gram-negative

bacteria (41). Future research could involve identifying at what stage of predation these costs

are incurred.

Why can't amoebas evolve one predation technique that works optimally on all prey

bacteria to eliminate the costs of diet generalism? After all, amoebas are a few hundred times

larger than the average bacterium by mass (42). They should be able to overwhelm most

bacteria with their size advantage. But bacteria possess many mechanisms to resist their

eukaryotic predators (14, 15, 43, 44). Morphological adaptations such as formation of

microcolonies, biofilms, and filamentation can prevent attack or ingestion by predators. Bacteria

can modify their membrane properties by changing the lipopolysaccharides on the outer

membrane, secreting an S-layer and sporulating among others that can help them avoid

recognition, ingestion and digestion by predators. They can also produce a huge array of

secondary metabolites that can deter predation by protists (45, 46). Thus, each species of

bacterium may possess a unique combination of defenses that is unlikely to yield to a common

predation strategy. It would be interesting to test how many different predation strategies *D.*

discoideum can employ. *D. discoideum* may treat some groups of bacteria similarly and some as

different.

In this paper, we have demonstrated that *D. discoideum* experiences switching costs,

concurrent costs, and costs of feeding on sub-optimal prey. Many interesting questions remain

for future research. It remains to be seen how often amoebas experience these costs in nature.

This depends on the patchiness of bacteria on a microscale. One might expect that amoebas

have plenty of choices because soil habitats are extremely high in both bacterial diversity and

abundance. But on a microscale, soil is made up of small aggregated particles that are

connected via a network of air- and water-filled pores (47). The three-dimensional nature of

these particles increases soil surface area such that as little of 10^{-6} percent of soil surfaces may

contain bacteria (48). These particles generally contain bacterial patches of a limited number of

clonal cells and these patches can be separated by distances that are large on a microbial scale

(49). However, upper layers of soil can contain large bacterial patches comprised of different

morphotypes. Thus, it is possible that amoebas are likely to feed on sub-optimal prey bacteria

sometimes and may experience switching costs as they move between bacterial patches.

Most of what is known about predation comes from large organisms, but most

organisms are microbes. The study of interactions between *D. discoideum* and its prey bacteria

presents an exciting new frontier to investigate the mechanisms that underlie the costs and

benefits associated with a generalist strategy. Microbial systems are particularly amenable to

experimentation of all kinds in the lab. Questions on the strategies of predator and prey can be

answered. Choice among bacterial prey, genetic bases of predation and resistance to it, and

ecological foraging decisions can all be addressed. Our findings here contribute to this rich and

exciting area of research.

Materials and Methods

Growth rate of *D. discoideum* on different species of bacteria isolated from forest soil

environments

To investigate costs of eating sub-optimal prey, we first measured the growth rate of wild clones

of *D. discoideum* on *Klebsiella pneumoniae,* the lab food bacterium, and on 22 species of

bacteria found in close-association with *D. discoideum* isolated by Brock et al. 2018 from forest

soil environments (Table 1). The bacterial species identification is based on closest partial 16S

BLAST hit by Brock et al. 2018. In this paper, we refer to the bacteria using these closest 16s

BLAST hits. We performed this experiment on 3 *D. discoideum* strains: QS1, QS6, and QS9. We

plated 100,000 amoebas with 200μl of 10 OD_{600} bacterial suspension on starving agar plates. We

estimated amoeba numbers at 20 hours and calculated doubling times as:

285
$$Doubling\ time\ (T_D)\ =\ ln(2)\ \times\ \frac{20}{ln(N_{t=20})-\ ln(N_{t=0})}$$

This assay also helped us identify reasonably edible bacteria for other experiments.

Switching cost assay in *D. discoideum* amoebas

To test if switching costs occur when *D. discoideum* amoebas are switched from Bacterium A to

Bacterium B, we first conditioned separate populations of the amoebas to each species of

bacterium (Fig 2, details below for three separate experiments). The switched treatment for

these experiments used amoebas conditioned to Bacterium A and replated with Bacterium B.

We plated 100,000 amoebas conditioned to Bacterium A with 200μl 20 OD_{600} suspension of

Bacterium B on SM/20 plates and then measured amoeba numbers after 3 hours. The controls

for this experiment were identical except that amoebas conditioned to Bacterium B were

replated with Bacterium B. If we observe fewer amoebas in switched plates compared to control

plates, then *D. discoideum* amoebas experience switching costs. We measured the costs of prey

switching between 3 pairs of bacteria with 2 reciprocal switches for each pair. We replicated this

experiment with 5 *D. discoideum* strains: QS6, QS9, QS14, QS17, QS160.

We did three sets of experiments to rule out the possibility that switching costs are

instead due to an evolved response in the amoebas during the conditioning period to either

bacterium. Switched amoebas could experience poor growth on a Bacterium B because of trade-

offs associated with new evolutionary adaptation to the Bacterium A during the conditioning

period. Control amoebas may also experience better growth rates if they have evolved and

adapted to Bacterium B during the conditioning period. We ruled out this complication in three

ways.

Effect of conditioning length of amoebas on switching costs

First, we tested if the length of the conditioning period of amoebas on the bacterium affected

switching costs. If switching costs are due to either adaptation in switched amoebas that results

in evolutionary trade-offs, or to direct improvement in growth rate because of adaptation in

control amoebas, we expect stronger costs with longer conditioning (i.e. evolving) time. Thus,

we conditioned amoebas on a given bacterium for a) 2 days and b) 5 days before measuring

switching costs. In the 2 day treatment, we conditioned the amoebas for 40 hours by plating

200,000 spores of *D. discoideum* with 200ul of 1.5 OD_{600} bacterial suspension on SM/5 plates.

After ~40 hours, we washed off the plates with 10ml of ice cold KK2 to collect the amoebae-

bacteria suspension. Next, we centrifuged the suspension for 3 minutes at 300g at 10°C to spin

down the amoebas and discarded the supernatant containing the bacteria. We resuspended the

amoeba pellet in ice-cold KK2 and washed off two more times until all the bacteria were

discarded. Finally, we suspended the amoeba pellet in 500μl – 1000μl of ice-cold KK2. We

conditioned the amoebas for the 5-day experiment similarly, but collected amoebas from the

resulting vegetative front after 5 days with a sterile loop and resuspending in 600ul ice-cold KK2.

We performed washing centrifuging steps as described above to thoroughly wash the amoebas

off the conditioning bacterium. A conditioning time of 2 days translates to 10-14 amoeba

divisions and 5 days to 30-40 amoeba divisions on good prey. If switching costs are because of

an evolutionary response, then we expect costs to be stronger for the 5-day conditioning period

than the 2-day period.

Switching costs assay with *D. discoideum* spores

To further distinguish between a temporary switching cost and an evolved response, we tested

if allowing amoebas to go through the social cycle and subsequent sporulation before prey

bacteria switches erases switching costs (Fig 2). If switching costs are because of an evolved

response in amoebas, then these changes should still be evident after the social stage. But if the

costs are because of conditioning of the amoebas that results in temporary mismatch in

transcriptional tools on the new bacterium, then undergoing the social cycle should erase most

switching costs. This is because *D. discoideum* experiences a huge turn-over in gene expression

when transitioning from a unicellular lifestyle into the multicellular cycle (Parikh et al. 2010,

Rosengarten et al. 2015). The abundance of almost every transcribed mRNA changes at least 2-

fold throughout development starting from vegetative amoebas to multicellular fruiting bodies

(50). Therefore, any transcriptional conditioning of amoebas towards a bacterium should be

largely erased by development. The switched and control spores are at the same transcriptional

start line of dormancy.

We used the same design as the switching experiment for amoebas described above but

with sporulation at the end of the conditioning phase (~7 days). *D. discoideum* amoebas feed on

bacteria for the first 2-4 days of the conditioning phase, and then transition to the social cycle

once the bacteria are depleted. We then plated 100,000 spores from the resulting fruiting

bodies with bacterial suspensions on SM/20 (Fig 2). Since the spores require some time to

germinate into amoebas and start feeding on the bacteria, we counted the total number of

amoebas on these plates after 36 hours.

Time course data of switching costs in *D. discoideum* amoebas

As a third check on whether what we see is an evolved or a conditioned response, we conducted

a time series study on one pair of bacteria (*A. tumefaciens* and *P. vranovensis*) to check if the

growth rates of the switched treatments quickly catch up with control growth rates, which

would indicate a physiological lag rather than an evolutionary change. We tracked the number

of amoebas in switched and control plates after 6, 24 and 27 hours from plating. We calculated

early growth rate between 0h and 6h time points and calculated late growth rate between 24h

and 27h time points. We assumed exponential growth and used this formula to calculate growth

rate between time points t1 and t2, where N_t stands for number of amoebas at time t:

360
$$Growth\ rate\ (r)\ =\ \frac{ln(N_{t2})\ -\ ln\ (N_{t1})}{(t2\ -\ t1)}$$

If switching costs are not an evolved response, then switches and controls should not differ in

their late growth rates. We performed this on SM/5 plates with a starting number of 100,000

amoebas with 200μl of 1.5 OD_{600} suspension of the test bacterium on the *D. discoideum* clone

QS9. We replicated this experiment five times.

Concurrent costs in *D. discoideum* amoebas

Next, we determined whether *D. discoideum* amoebas experience concurrent costs of preying

on multiple bacteria. For this experiment, we chose 5 prey bacteria that were generally highly

edible from our growth rate assay of amoebas on 23 species of bacteria. We generated 5 single-

species communities, 6 three-species communities and 1 five-species community from the 5

prey bacteria. We made OD10 suspensions of each bacterial species and mixed these

suspensions in equal proportions to make a given multi-species suspension. We used AX4

amoebas that were grown axenically in bacteria-free cultures to preclude any effect of prior

conditioning of amoebas to a given species of bacterium on the assay. We plated 100,000

amoebas on starving agar plates with 200μl of bacterial suspensions and measured amoeba

numbers in all the different treatments after 20 hours. We replicated this experiment 6 times.

To test if amoebas experience concurrent costs, we first calculated the expected number of

amoebas in multi-prey treatments with data from our single-prey treatments. For example, the

expected number of amoebas in multi-prey treatment containing *A. tumefaciens*, *S. maltophilia*

and *C. testosteroni* is the average of observed numbers of amoebas in single-prey treatment of

those bacteria. We then compared these expected numbers to our observed number of

amoebas in these multi-species treatments to infer concurrent costs.

Statistics:

We performed all statistical analysis in R (version 4.2.1) (52). To test if *D. discoideum* amoebas

experience varying doubling times on different species of bacteria, we used a one-way analysis

of variance with amoeba doubling times as the response variable and the Test bacterium (23

species of bacteria listed in Table 1) as the fixed factor (Table S1). We define Test bacterium as

the bacterium on which we measured amoeba growth rates. To test if *D. discoideum* amoebas

experience a switching cost on a given bacterium when previously grown on a different

bacterium we used a three-way analysis of variance (ANOVA) with log transformed amoeba

numbers after 3 hours of growth as the response variable and Treatment (Control/Switch), Test

bacterium (At, Pv, Fg, Sm, Sz, Ct), and Conditioning length (2days/5days) as fixed factors. We

also included interaction effects for Treatment x Test bacterium, Treatment x Conditioning

length (Table S2). For the similar experiment that includes a spore stage after conditioning, we

used a two-way analysis of variance (ANOVA) because there was no variation in conditioning

length. Log transformed amoeba numbers after 36 hours was the response variable and

Treatment and Test bacterium were the fixed factors. We included interaction effects for

Treatment x Test bacterium (Table S3).

We performed the following statistical tests on time-course data collected on (At, Pv)

switches. To test if there are switching costs during early proliferation we used a two-way

analysis of variance (ANOVA) with early growth rate calculated between 0 and 6 hours as the

response variable and Treatment (Control/Switch) and Test bacterium (At, Pv) as fixed factor

(Table S4). We performed a similar two-way ANOVA with late growth rate calculated between

24 and 27 hours as the response variable to test if switching costs persist during late

proliferation (Table S5).

To test if *D. discoideum* amoebas experience concurrent costs, we tested if amoebas

performed worse than expected in multi-species prey communities. We used a three—way

analysis of variance (ANOVA) with log transformed amoeba numbers at 20 hours as the

response variable, Treatment (Expected numbers/Observed numbers) and Bacterial Community

(6 Communities) and Experimenter (2 Experimenters) as fixed factors. We included an

interaction effect between Treatment x Bacterial Community (Table S6).

Acknowledgments

This material is based upon work funded by the National Science Foundation under grants

IOS 16-56756, DEB 17-53743, and DEB 2237266. We thank Margaret Steele and Calum

Stephenson for comments on the manuscript. We thank Trey Scott for feedback on statistical

analysis. We would also like to thank the Queller-Strassmann lab for their helpful insights on

experimental design.

419
420
References

1.   B. D. Moore, W. J. Foley, A review of feeding and diet selection in koalas (Phascolarctos
cinereus). *Aust. J. Zool.* **48**, 317 (2000).

2.   S. R. Beissinger, Hunting behavior, prey selection, and energetics of snail kites in Guyana:
consumer choice by a specialist. *The Auk* **100**, 84–92 (1983).

3.   J. M. McVey, *et al.*, Diets of sympatric red wolves and coyotes in northeastern North
Carolina. *Journal of Mammalogy* **94**, 1141–1148 (2013).

4.   J. M. Schmidt, P. Sebastian, S. M. Wilder, A. L. Rypstra, The nutritional content of prey
affects the foraging of a generalist arthropod predator. *PLOS ONE* **7**, e49223 (2012).

5.   B. Eitzinger, *et al.*, Assessing changes in arthropod predator–prey interactions through
DNA-based gut content analysis—variable environment, stable diet. *Molecular Ecology* **28**,
266–280 (2019).

6.   E. A. Bernays, O. P. J. M. Minkenberg, Insect herbivores: different reasons for being a
generalist. *Ecology* **78**, 1157–1169 (1997).

7.   D. J. Futuyma, G. Moreno, The evolution of ecological specialization. *Annual Review of
Ecology and Systematics*, 29 (1998).

8.   M. L. Forister, L. A. Dyer, M. S. Singer, J. O. Stireman III, J. T. Lill, Revisiting the evolution of
ecological specialization, with emphasis on insect–plant interactions. *Ecology* **93**, 981–991
(2012).

9.   E. B. Sherr, B. F. Sherr, Significance of predation by protists in aquatic microbial food webs.
Antonie van Leeuwenhoek **81**, 16 (2002).

10.   J. Johnke, *et al.*, A generalist protist predator enables coexistence in multitrophic predator-
prey systems containing a phage and the bacterial predator Bdellovibrio. *Front. Ecol. Evol.*
5 (2017).

11.   S. Geisen, *et al.*, Soil protists: a fertile frontier in soil biology research. *FEMS Microbiology
Reviews* **42**, 293–323 (2018).

12.   T. Fenchel, *Ecology of Protozoa* (Springer Berlin Heidelberg, 1987)
448   https:/doi.org/10.1007/978-3-662-06817-5 (August 4, 2023).

449   13.   G. Corno, K. Jürgens, Structural and functional patterns of bacterial communities in
450   response to protist predation along an experimental productivity gradient: Top-down and
451   bottom-up control of bacteria. *Environmental Microbiology* **10**, 2857–2871 (2008).

452   14.   A. Jousset, Ecological and evolutive implications of bacterial defences against predators:
453   antipredator defences and bacterial ecology. *Environmental Microbiology* **14**, 1830–1843
454   (2012).

455   15.   W. P. J. Smith, B. R. Wucher, C. D. Nadell, K. R. Foster, Bacterial defences: mechanisms,
456   evolution and antimicrobial resistance. *Nat Rev Microbiol* (2023)
457   https:/doi.org/10.1038/s41579-023-00877-3 (May 11, 2023).

458   16.   R. H. Kessin, *Dictyostelium: Evolution, Cell Biology, and the Development of Multicellularity*
459   (Cambridge University Press, 2001).

460   17.   J. E. Strassmann, D. C. Queller, Evolution of cooperation and control of cheating in a social
461   microbe. *Proceedings of the National Academy of Sciences* **108**, 10855–10862 (2011).

462   18.   K. B. Raper, N. R. Smith, The growth of Dictyostelium discoideum upon pathogenic
463   bacteria. 15 (1939).

464   19.   V. E. Paquet, S. J. Charette, Amoeba-resisting bacteria found in multilamellar bodies
465   secreted by *Dictyostelium discoideum:* social amoebae can also package bacteria. *FEMS*
466   *Microbiology Ecology* **92**, fiw025 (2016).

467   20.   D. A. Brock, *et al.*, Diversity of free-living environmental bacteria and their interactions
468   with a bactivorous amoeba. *Front Cell Infect Microbiol* **8** (2018).

469   21.   D. Dykhuizen, Species numbers in bacteria. *Proc Calif Acad Sci* **56**, 62–71 (2005).

470   22.   C. L. Lauber, K. S. Ramirez, Z. Aanderud, J. Lennon, N. Fierer, Temporal variability in soil
471   microbial communities across land-use types. *ISME J* **7**, 1641–1650 (2013).

472   23.   jeff smith, D. C. Queller, J. E. Strassmann, Fruiting bodies of the social amoeba
473   Dictyostelium discoideum increase spore transport by Drosophila. *BMC Evolutionary*
474   *Biology* **14**, 105 (2014).

475   24.   S. Kumar, G. Stecher, M. Suleski, S. B. Hedges, Timetree: a resource for timelines,
476   timetrees, and divergence times. *Molecular Biology and Evolution* **34**, 1812–1819 (2017).

477   25.   H. Cornell, Search strategies and the adaptive significance of switching in some general
478   predators. *The American Naturalist* **110**, 317–320 (1976).

479   26.   R. N. Hughes, Optimal diets under the energy maximization premise: the effects of
480   recognition time and learning. *The American Naturalist* **113**, 209–221 (1979).

481   27.   W. Nasser, *et al.*, Bacterial discrimination by Dictyostelid amoebae reveals the complexity
482   of ancient interspecies interactions. *Current Biology* **23**, 862–872 (2013).

483   28.   O. Lamrabet, *et al.*, Transcriptional responses of Dictyostelium discoideum exposed to
484   different classes of bacteria. *Front. Microbiol.* **11**, 410 (2020).

485   29.   L. Gamfeldt, H. Hillebrand, P. R. Jonsson, Species richness changes across two trophic
486   levels simultaneously affect prey and consumer biomass. *Ecology Letters* **8**, 696–703
487   (2005).

488   30.   M. Saleem, I. Fetzer, H. Harms, A. Chatzinotas, Diversity of protists and bacteria
489   determines predation performance and stability. *ISME J* **7**, 1912–1921 (2013).

490   31.   J. A. Myers, B. S. Curtis, W. R. Curtis, Improving accuracy of cell and chromophore
491   concentration measurements using optical density. *BMC Biophys* **6**, 4 (2013).

492   32.   S. DiSalvo, D. A. Brock, jeff smith, D. C. Queller, J. E. Strassmann, In the social amoeba
493   Dictyostelium discoideum, density, not farming status, determines predatory success on
494   unpalatable Escherichia coli. *BMC Microbiology* **14**, 328 (2014).

495   33.   S. Adiba, C. Nizak, M. van Baalen, E. Denamur, F. Depaulis, From grazing resistance to
496   pathogenesis: The coincidental evolution of virulence factors. *PLOS ONE* **5**, e11882 (2010).

497   34.   G. Rashidi, E. A. Ostrowski, Phagocyte chase behaviours: discrimination between Gram-
498   negative and Gram-positive bacteria by amoebae. *Biol. Lett.* **15**, 20180607 (2019).

499   35.   B. Stewart, *et al.*, The genetic architecture underlying prey-dependent performance in a
500   microbial predator. *Nat Commun* **13**, 319 (2022).

501   36.   V. Jaumouillé, S. Grinstein, Receptor mobility, the cytoskeleton, and particle binding during
502   phagocytosis. *Current Opinion in Cell Biology* **23**, 22–29 (2011).

503   37.   P. N. Devreotes, *et al.*, Excitable signal transduction networks in directed cell migration.
504   *Annual Review of Cell and Developmental Biology* **33**, 103–125 (2017).

505   38.   A. J. Heidel, *et al.*, Phylogeny-wide analysis of social amoeba genomes highlights ancient
506   origins for complex intercellular communication. *Genome Research* **21**, 1882–1891 (2011).

507   39.   M. Pan, *et al.*, A G-protein-coupled chemoattractant receptor recognizes
508   lipopolysaccharide for bacterial phagocytosis. *PLOS Biology* **16**, e2005754 (2018).

509   40.   O. Lamrabet, T. Jauslin, W. C. Lima, M. Leippe, P. Cosson, The multifarious lysozyme
510   arsenal of Dictyostelium discoideum. *Developmental & Comparative Immunology* **107**,
511   103645 (2020).

512   41.   T. Jauslin, *et al.*, How phagocytic cells kill different bacteria: a quantitative analysis using
513   Dictyostelium discoideum. *mBio* **12**, e03169-20, /mbio/12/1/mBio.03169-20.atom (2021).

42. R. Kumar, R. Maurya, S. Saran, Investigating the role of translationally control tumor
protein in growth, development and differentiation of Dictyostelium discoideum. *Frontiers*
in Cell and Developmental Biology **8** (2020).

43. C. Matz, S. Kjelleberg, Off the hook – how bacteria survive protozoan grazing. *Trends in*
Microbiology **13**, 302–307 (2005).

44. D. W. Kikuchi, *et al.*, The evolution and ecology of multiple antipredator defences. *J of*
Evolutionary Biology **36**, 975–991 (2023).

45. O. Tyc, C. Song, J. S. Dickschat, M. Vos, P. Garbeva, The ecological role of volatile and
soluble secondary metabolites produced by soil bacteria. *Trends in Microbiology* **25**, 280–
292 (2017).

46. J. M. Raaijmakers, M. Mazzola, Diversity and natural functions of antibiotics produced by
beneficial and plant pathogenic bacteria. *Annu. Rev. Phytopathol.* **50**, 403–424 (2012).

47. M. Vos, A. B. Wolf, S. J. Jennings, G. A. Kowalchuk, Micro-scale determinants of bacterial
diversity in soil. *FEMS Microbiol Rev* **37**, 936–954 (2013).

48. I. M. Young, J. W. Crawford, Interactions and self-organization in the soil-microbe complex.
Science **304**, 1634–1637 (2004).

49. N. Nunan, K. Wu, I. M. Young, J. W. Crawford, K. Ritz, Spatial distribution of bacterial
communities and their relationships with the micro-architecture of soil. *FEMS*
Microbiology Ecology **44**, 203–215 (2003).

50. A. Parikh, *et al.*, Conserved developmental transcriptomes in evolutionarily divergent
species. *Genome Biology* **11**, R35 (2010).

51. R. D. Rosengarten, *et al.*, Leaps and lulls in the developmental transcriptome of
Dictyostelium discoideum. *BMC Genomics* **16**, 294 (2015).

52. A. R Core Team, R. C. Team, *R: A language and environment for statistical computing. R*
Foundation for Statistical Computing, Vienna, Austria. 2012 (2022).

539
540

Figures and Tables

Figure 1. Doubling time of *D. discoideum* on *K. pneumoniae* and 22 species of soil bacteria closely

associated with *D. discoideum*. Error bars are mean amoeba doubling time ± S.D. The points

represent the 3 *D. discoideum* strains. The circles represent the 8 species of bacteria used in

other parts of this study. Bacteria species identities by Brock et al. 2018 based on closest partial

16S BLAST hit are listed in Table 1.

Figure 2. Schematic on the left shows switching costs experiment in amoebas. Amoebas in

switched treatment are conditioned on bacterium A and are switched to bacterium B for the

experiment. The controls for this experiment are amoebas conditioned to bacterium B plated

with a fresh culture of the same. We also varied conditioning time of amoebas on a given

bacterium (2 days, 5 days). Schematic on the right outlines switching costs experiment on

spores. We used the same basic switching design as described for the amoebas, except we let

the amoebas undergo multicellular development on the conditioning bacterium. We then

collected spores from these fruiting bodies and performed the experiment.

Figure 3: *D. discoideum* amoebas experience prey switching costs: Mean amoeba numbers ± S.D

of controls and switches after 3 hours of growth. The shapes represent the different

conditioning times of amoebas (• - 2 days, ▲ - 5 days). The points represent the five *D.*

discoideum strains. Total amoeba numbers are significantly different between controls and

switches pooled across all treatments ($F_{1,106}$ = 27.621, p-value = 7.69 x 10^{-7}). There was no

significant interaction between conditioning time and switching costs (Treatment x Conditioning

length: $F_{1,106}$ = 0.543, p-value = 0.463). Interpreting x-axis labels: For example - Sm-Sm refers to

amoebas grown on *S. maltophilia* moved to *S. maltophilia*, while Fg-Sm refers to amoebas

grown on *F. ginsengisoli* moved to *S. maltophilia*.

Figure 4: Switching costs disappear when spore formation precedes prey switch: Mean amoeba

numbers ± S.D of controls and switches at 36 hours from plating. The points represent the five

D. discoideum strains. We found no significant difference in amoeba numbers between controls

and switches ($F_{1,48}$ = 0.263, p-value = 0.61).

Figure 5: Switching costs occur during early growth period rather than later: A) Early growth rate

estimates of controls and switches between 0 and 6 hours from set-up. Switched treatments

have significantly lower early growth rates compared to controls $F_{1,17}$ = 9.458, $p-value$ =0.007)

B) Late growth rate estimates of controls and switches between 24 to 27 hours from set-up for

(*Agrobacterium tumefaciens*, *Pseudomonas vranovensis*). We found no difference in growth

rates switches and controls at the later time points ($F_{1,17}$ = 0.111, p-value = 0.73) The points

represent 5 technical replicates of *D. discoideum* clone QS9.

Figure 6: *D. discoideum* amoebas experience concurrent costs: Ratios of observed vs. expected

amoeba numbers in different bacterial communities. A ratio lower than one indicates

concurrent costs. Blue dot marks the mean. There are significantly fewer amoebas than

expected in multi-species communities ($F_{1,59}$=11.83, p-value = 0.001). At- *Agrobacterium*

tumefaciens, Bw- *Buttiauxella warmboldiae*, Ct - *Comamonas testosteroni*, Sm -

Stenotrophomonas maltophilia, Pk - *Pseudomonas kuykendallii*.

Table 1: List of 23 species of bacteria used in *D. discoideum* growth rate experiments. *
Species identification of bacterial isolates by Brock et al. 2018 based on closest partial 16S
BLAST hit.

Strain number	Closest 16S sequence	Phylum/Class	Abbreviation
Lab clone	*Klebsiella pneumoniae*	Gammaproteobacteria	Kp
14P 8.1.1	*Pseudomonas kuykendallii*	Gammaproteobacteria	Pk
18P 8.2.2	*Buttiauxella warmboldiae*	Gammaproteobacteria	Bw
18P 2.2.1	*Agrobacterium tumefaciens*	Alphaproteobacteria	At
20P 9.1.2	*Agrobacterium rubi*	Alphaproteobacteria	Ar
14P 4.3.1	*Serratia liquefaciens*	Gammaproteobacteria	Sl
14P 6.2.3	*Stenotrophomonas maltophilia*	Gammaproteobacteria	Sm
18P 6.2.3	*Comamonas testosteroni*	Betaproteobacteria	Ct
20P 10.2.4	*Variovorax boronicumulans*	Betaproteobacteria	Vb
14P 8.1.4	*Comamonas kerstersii*	Betaproteobacteria	Ck
20P 3.2.2	*Flavobacterium ginsengisoli*	Bacteroidetes	Fg
20P 9.1.1	*Shinella zoogloeoides*	Alphaproteobacteria	Sz
20P 6.1.2	*Brucella papionis*	Alphaproteobacteria	Bp
5S 2.1.1	*Staphlycoccus saprophyticus*	Firmicutes	Ss
5P 5.1.1	*Pseudomonas vranovensis*	Gammaproteobacteria	Pv
18P 8.1.1	*Achromobacter aegrifaciens*	Betaproteobacteria	Aa
14P 5.3.2	*Pseudomonas migulae*	Gammaproteobacteria	Pm
20P 10.2.1	*Sphingobacterium ginsensidimutans*	Bacteroidetes	Sg
20P 2.1.2	*Pandoraea oxalativorans*	Betaproteobacteria	Po
14P 4.3.2	*Chryseobacterium rhizosphaerae*	Bacteroidetes	Cr
14P 6.2.1	*Microbacterium maritypicum*	Actinobacteria	Mm
20P 7.2.1	*Bacillus aryabhattai*	Firmicutes	Ba
20P 2.1.1	*Paenibacillus pabuli*	Firmicutes	Pp

596   **Supplemental information:**

597   **Table S1: ANOVA of amoeba growth rate assay**

598   **aov(Doubling_time ~ Test_bacterium)**

	Df	Sum sq	Mean Sq	F_value	Pr(>F)
Test_bacterium	22	299.346	13.3794	24.949	2.20E-16
Residuals	46	24.668	0.5363		

600   **Table S2: ANOVA of amoeba switching assay**

601   **aov(log(amoeba_number) ~ Treatment x Test_bacterium + Treatment x Conditioning_length)**

	Df	Sum sq	Mean Sq	F_value	Pr(>F)
Treatment	1	0.965	0.9648	27.621	7.69E-07
Test_bacterium	5	4.958	0.9917	28.392	2.00E-16
Conditioning_length	1	0.057	0.0567	1.623	0.205
Treatment x Test_bacterium	5	0.146	0.0293	0.838	0.526
Treatment x Conditioning_length	1	0.019	0.019	0.543	0.463
Residuals	106	3.702	0.0349		

603   **Table S3: ANOVA of spore switching assay**

604   **aov(log(amoeba_number) ~ Treatment x Test_bacterium)**

	Df	Sum sq	Mean Sq	F_value	Pr(>F)
Treatment	1	0.03	0.03	0.264	0.61
Test_bacterium	5	106.78	21.355	186.402	<2e-16
Treatment x Test_bacterium	5	0.68	0.136	1.191	0.328
Residuals	48	5.5	0.115		

606   **Table S4: ANOVA of early growth rate data**

607 **aov (early.growth.rate ~ Treatment + Test_bacterium)**

	Df	Sum_sq	Mean sq	F value	Pr(>F)
Treatment	1	0.012	0.012	9.458	0.007
Test_Bacterium	1	0.005	0.005	3.931	0.064
Residuals	17	0.022	0.001		

609 **Table S5: ANOVA of late growth rate data**

610 **aov(late.growth.rate ~ Treatment + Test_bacterium)**

	Df	Sum_sq	Mean sq	F value	Pr(>F)
Treatment	1	0.000	0.000	0.111	0.743
Bacteria	1	0.095	0.095	42.836	5.00E-06
Residuals	17	0.038	0.002		

612 **Table S6: ANOVA of concurrent costs data**

613 **aov (log(Total.amoeba.number) ~ Treatment x Community + Experimenter)**

	Df	SumSq	MeanSq	F-value	Pr(>F)
Treatment	1	0.4564	0.4564	11.8394	0.001071
Community	5	1.6847	0.3369	8.74	3.01E-06
Experimenter	1	24.3675	24.3675	632.0609	2.20E-16
Treatment x Community	5	0.4929	0.0986	2.5571	0.036767
Residuals	59	2.2746	0.0386		