Resource-combining costs of being a diet generalist in the super-generalist protist

predator *Dictyostelium discoideum*

P. M. Shreenidhi *, Debra A. Brock, Rachel I. McCabe, Joan E. Strassmann and David C. Queller

Department of Biology, Washington University in St. Louis, St. Louis, MO 63130

Corresponding author: *P. M. Shreenidhi

Email: shreenidhi@wustl.edu

Author Contributions: P.M.S., J.E.S., and D.C.Q. designed research. P.M.S., D.A.B., and R.I.M.

performed the experiments. P.M.S., J.E.S., and D.C.Q. analyzed data. P.M.S., J.E.S., and D.C.Q.

wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Classification: Major - Biological Sciences, Minor- Ecology

Keywords: Diet breadth, Generalism, Predation, Protists, Resource-switching costs

This PDF file includes:

Main Text
Figures 1 to 6
Tables 1
Supplementary figures and tables
18

Abstract

Consumers lie on a continuum between diet specialization on few resources to being generalist

feeders on many resources. Generalism has the clear advantage of having more resources to

exploit, but the costs that limit generalism are less clear. We explore two understudied costs of

generalism in a super-generalist amoeba predator, *Dictyostelium discoideum*, feeding on

naturally co-occurring bacterial prey. Both involve costs of combining different prey. First,

amoebas exhibit a reduction in growth rate when they switch from one species of prey bacteria

to another, something we call resource-switching costs. These switching costs typically

disappear within a day, indicating adjustment to new prey bacteria. Second, amoebas usually

divide more slowly on mixtures of bacteria compared to on single bacteria, something we call

resource-mixing costs. Both results support that idea that, although amoebas can consume a

huge variety of prey, they must use partially different methods and thus must pay costs to

handle multiple prey, either sequentially or simultaneously.

Significance Statement

Perhaps the most fundamental conflict in nature occurs when one organism consumes another.

Diet generalists benefit from the advantage of eating many prey but then must deal with many

prey defences. We explore costs associated with a broad diet in a protist microbial predator,

Dictyostelium discoideum. These predators of bacteria show a delay in growth when switched

from one bacteria to another, supporting the hypothesis that they must deploy different

strategies. They also experience costs when grown on many bacteria at once, suggesting that

the alternative strategies for consuming different prey are partly incompatible with each other.

Our findings shed light on the nature of diet generalism and highlight the complexity of

predation in the microbial world.

Main Text

Introduction

Consumers vary widely in their diet breadth. Some are diet specialists that eat one or few

resources, such as koalas that feed only on eucalyptus leaves (1) or snail kites that exclusively

hunt apple snails (2). Some are diet generalists that consume many different resources, such as

coyotes that feed on many small mammals (3) or spiders that feed on many species of

arthropods (4, 5). This variation in diet breadth has important ecological and evolutionary

consequences that impact the structure and stability of food-webs (6), the stability of

communities to perturbations (7), within- and between-species competition (8), the strength of

coevolutionary dynamics (9) and speciation (10).

The obvious benefit to being a diet generalist is the ability to exploit diverse resources,

especially when resources are scarce and fluctuate in their availability (11). But there must be

associated costs with diet generalism or else all consumers would be generalists. One classic

explanation for diet specialization is that "the jack of all-trades is the master of none." We

expect generalists to suffer from trade-offs in resource utilization because of antagonistic

pleiotropy, where mutations that are good for exploiting one resource maybe bad on another

(12–14). Such trade-offs can also occur via mutation accumulation, where mutations that are

neutral on one resource may be detrimental on new resources (15). Generally, there has been

surprisingly little support for trade-offs in performance on different resources (12, 16, 17).

Other, less studied, costs include that generalists may also be slower to adapt to a given

resource because they spend less time on a given resource compared to specialists (18), and

that generalists can suffer from information costs from having to track more information about

their resource environment (19).

Another kind of cost that seems to be rarely considered is the cost of combining

resources. We distinguish two types of combining costs. First, generalists occur in environments

with shifting resource abundances and may need to switch among different resources. If

different resources require different methods of exploitation, then changing among these

methods can result in costs. These methods may be related to resource recognition, handling

time, processing or detoxification (20, 21). We refer to these as resource-switching costs. Such

costs have been observed in arctic charr, where the fish experience lower metabolic rates when

fed a diet different from the one they were raised on compared to fish that did not have a

dietary change (22).

A second kind of combining cost is when generalists try to feed on multiple resources at

once and the specific techniques that work best for different resources are partly incompatible.

It may be impossible to effectively deploy multiple optimal strategies at the same time. For

example, synergistic interactions between defensive chemicals of different resources could

occur such that the combined toxins are more detrimental (23). We refer to the reduction in

foraging efficiency in the presence of multiple resources as resource-mixing costs. However,

generalists could alternatively benefit from resource-mixing because of improved nutritional

balance or through reduction in the amount of toxins ingested from any one resource or prey

(24). These two kinds of combining costs aren't the same as standard evolutionary trade-offs

because a generalist may do just as well on every individual resource as a specialist but do more

poorly when it tries to mix or switch among resources.

The costs and benefits associated with diet breadth evolution have been extensively

studied in herbivorous insects. They are one of the most abundant and diverse eukaryotic life-

forms and often possess highly specialized diets (25, 26). But the study of consumers with highly

generalized diets provides an equally important perspective. Predators are often among the

most general of consumers. By definition, predators are consumers that kill more than one

victim in their lifetime (27), usually many more. They also tend to be much larger than their

victims. Thus, to avoid starvation and minimize variance in energy intake, predators may need to

consume many sub-optimal prey types when the most profitable prey is not abundant enough

(28). Many macroscopic predators consume a formidable number of prey species, but few can

match the diversity of prey consumed by some microbial protist predators. We call species with

extreme diet breadth "super-generalists". Predation by protists is a major factor accounting for

bacterial mortality in the environment and as a consequence plays an important role in nutrient

cycling (29–31). Predation by protists can determine the composition and properties of bacterial

communities (32) and can be an important selective pressure for bacterial defenses such as

biofilm formation, antibiotic production, and secretion systems (33, 34).

Dictyostelium discoideum is one such super-generalist protist predator. It is a social

amoeba that lives in forest soils. It is a unicellular amoeba when bacterial prey are abundant and

transitions to a non-feeding multicellular dispersal stage upon starvation. *D. discoideum* has

been the subject of extensive research because of this fascinating multicellular stage (35, 36). Its

feeding behavior has been less extensively studied, but it appears to qualify as a super-

generalist predator. It can eat the majority of bacteria it is presented with (37–39). For example,

one study tested 159 bacterial strains found in close association with fruiting bodies of *D.*

discoideum from forest soil habitats and found that the amoebas were able to consume 77% of

them (39).

The prey bacteria of *D. discoideum* are very diverse, ranging across at least four highly

divergent bacterial phyla: Actinobacteria, Bacteroidetes, Firmicutes, and Proteobacteria (38).

These shared a common ancestor about three billion years ago, far older than the divergence

time of diverse insect prey (~400 mya) that generalist invertebrates feed on (40). Bacteria also

possess highly varied defensive mechanisms against microbial predators (33, 34). They can

produce many kinds of secondary metabolite toxins to repel, disable, or kill their enemies. They

can use secretion systems and effectors that can kill or allow for intracellular survival within

protists. Some bacteria can swim away at high speeds to escape predators. Some can group

together to form biofilms to prevent ingestion by predators.

Here we investigate the two kinds of costs of combining resources in *D. discoideum*.

Amoebas occur in spatially and temporally variable communities of soil bacteria (41). Amoebas

can encounter patchy bacterial distributions such that they switch from hunting one species of

bacteria to another. We therefore tested for resource-switching costs by seeing if amoebas

perform worse when switched to a new species of bacteria compared to controls that continued

to grow on the same bacterium. Amoebas will often encounter mixed communities of prey, so

we also looked at resource-mixing costs, testing if amoebas performed worse than expected in

multi-species bacterial communities compared to their growth in single-species communities.

Results

Growth rate of *D. discoideum* amoebas varies on different species of bacteria.

As a preliminary step we confirmed that *D. discoideum* is a generalist. We measured the growth

rate of three *D. discoideum* clones on the commonly used lab food bacterium, *Klebsiella*

pneumoniae and on 22 species of bacteria that had been collected as transient associates on *D.*

discoideum fruiting bodies. These fruiting bodies emerged from soil and deer feces collected in

the field, so the 22 bacterial species represent biologically relevant prey species for the

amoebas. *D. discoideum* amoebas showed wide variation in their doubling times on different

soil bacteria (Test bacterium: $F_{22,45}$ = 23.88, p-value < 2.2×10^{-16}). On each of the 23 bacteria, all

three *D. discoideum* clones grew similarly (Figure 1), ruling out the possibility that the

generalism of the species might be due to a mixture of individually specialized clones (42). The

results also confirm that the amoebas are generalist feeders on prey bacteria that are likely to

be encountered in nature.

***D. discoideum* amoebas experience resource-switching costs**

We conducted resource-switching cost experiments to investigate how *D. discoideum* amoebas

proliferate on a given bacterium when previously conditioned to a different bacterium (Figure

2). We found that there were fewer amoebas at 3 hours in switched treatments compared to

controls that did not experience a prey switch (Figure 3, Treatment $F_{1,106}$ = 27.621, p -value =

7.69×10^{-7}; Effect size = - 0.96, 95% CI = [-1.34, -0.575], df = 106; all effect sizes in this paper are

the standardized measure Cohen's d). Although only some bacterial species showed individually

significant switching costs (Figure S1A), there was no interaction of treatment with prey species

(Treatment x Test bacterium: $F_{1,106}$= 0.838, p-value = 0.526) and the overall effect size was

strong. Possible percentage change in amoeba numbers ranges from -24.04% to -8.03% (95%

CI). We also found that conditioning time of amoebas (2 days or 5 days) had a similar effect on

switching costs (Figure S1B, Conditioning time of 2 days: Treatment p-value < 0.001; Effect size =

-1.094, 95% CI = [-1.63, -0.561], df =106, Conditioning time of 5 days: Treatment p-value =

0.0018; Effect size = -0.825, 95% CI = [-1.35, -0.301], df =106). This is inconsistent with switching

costs being due to evolutionary changes during conditioning since evolutionary changes should

be greater over 5 days than 2. The trend in the magnitude of effect sizes is also inconsistent with

the evolutionary hypothesis because we would expect larger effect with longer conditioning

time.

No evidence for resource-switching costs if amoebas undergo spore formation before the prey

switch

If resource-switching costs are due to an evolved response in amoebas, then these genetic

changes should be passed on through the spores. But if the costs are physiological, then

undergoing the social cycle would be likely to erase the effects of food conditioning and

therefore eliminate switching costs, because spores have very different gene expression profiles

compared to amoebas (43, 44). We therefore tested if *D. discoideum* still show switching costs

when they undergo the social cycle and spore formation before changing between prey

bacteria. The switched and control spores resulted in a similar number of amoebas (Figure 4,

Treatment: $F_{1,48}$ = 0.263, p-value = 0.61; Effect size = -0.133, 95% CI = [-0.653,0.387], df = 48).

Possible percentage change in amoeba numbers ranges from -25.43% to 22.58%. Thus, we find

no support for the hypothesis that switching costs are due to an evolved response, though the

lower end of confidence interval of effect size does not rule out some cost.

No evidence that resource-switching cost persists over the long term

If switching costs are not due to an evolved response, as supported by our previous results, then

the switched amoebas should eventually recover their growth rates to the level of the controls

that did not switch to new food bacteria. We conducted a time-course experiment to test this

for all three pairs of bacteria and compared early (zero to six hours) and late (24 to 27 hours)

growth rates of the switched and control treatments. As before (though tested here at 6 hours

rather than 3 hours), the switched treatments have significantly lower early growth rates

compared to controls (Figure 5A, Treatment $F_{1,53}$ = 12.98, p-value = 6.9 x 10^{-4}; Effect size =-0.931,

95% CI = (-1.48, -0.382), df= 53). Possible percentage change in growth rates ranges from -

27.47% to -3.75%. But there was no difference in late growth rate of switches and controls

(Figure 5B, Treatment $F_{1,53}$ = 0.014, p-value = 0.90; Effect size = -0.0311 , 95% CI = (-0.549,

0.487), df= 53). Possible percentage change in growth rates ranges from -23.72% to 28.31%.

Thus, though switched amoebas proliferate slower than controls during early growth, they

appear to eventually catch up and perform similarly to controls, consistent with physiological

adjustment to the new prey. Again, the confidence interval of effect size cannot rule out some

cost at the late stage.

***D. discoideum* amoebas experience resource-mixing costs in some multi-prey communities**

We tested if *D. discoideum* experiences costs when grown in multi-species prey communities

compared to expectations from their growth in single-species communities. Consistent with

resource-mixing costs, we observed significantly fewer amoebas than expected after growing in

multi-species communities (Figure 6, Treatment $F_{1,59}$ = 11.84, p-value = 0.001; Effect size = -

0.811, 95% CI = [-1.31, -0.316], df = 59). Possible percentage change in observed and expected

amoeba numbers ranges from -25.19% to -2.79%. But there seems to be variation in these

mixing costs between the different bacterial communities (Treatment x Bacterial Community:

$F_{5,59}$ = 2.557 p-value = 0.036). Three of the prey communities cause significant costs as judged by

non-overlap of effect size confidence intervals with zero, and none show a significant mixing

benefit (Figure S2).

Discussion

Consumers lie on a spectrum between specialization on a few resources or generalism on a wide

variety of resources. The balance between costs and benefits associated with these two

strategies determines where the consumer falls in that spectrum and has important ecological

and evolutionary consequences. But the costs of generalism are not well understood. We

examined an understudied type of cost of diet generalism in the social amoeba, *D. discoideum*.

As a preliminary step, we showed that the different strains of *D. discoideum* amoebas are true

generalists that grow similarly on any given species of bacterium (Figure 1). Some have

hypothesized that a generalist species could be a collection of individual specialists (23, 42), but

our results show that this is not the case in *D. discoideum*.

Amoebas experience early resource-switching costs when moved from one prey

bacterium to another, even after controlling for variation in bacterial edibility (Figures 3, 5A,

S1A). The smallest costs consistent with the 95% confidence intervals were 8% and 4% and even

these smallest effects seem biologically significant. We also did three experiments to test if

these costs were physiological in nature or due to an evolved response. First, we found no

evidence that the conditioning time (duration spent by amoebas on a given bacterium before

the resource switch) affected the magnitude of switching costs. If the costs were due to

evolution during the conditioning period, they should become stronger with longer conditioning

time. Second, we found no evidence that amoebas experience switching costs if they undergo

spore formation before prey switch (Figure 4). If this was due to an evolved response, then

these effects would be genetically passed on through the spores. Third, we found that switching

costs occur during early growth but are no longer apparent after a handful of additional

generations (Figure 5), faster than they would change by selection. Our conclusions from these

three experiments must be drawn from their failure to find significant costs, and the confidence

intervals of each cannot exclude some costs. However, we have three different kinds of

experiment, each one failing to support the hypothesis of evolutionary change and we have two

separate experiments showing significant costs during the early stages where physiological

effects were expected (Figures 3, 5A). Taken together these five experiments make a strong case

that these costs are physiological in nature and not due to evolution.

Resource-switching costs have not been extensively studied and rarely enter into

broader discussions of generalism, but there are other examples. Arctic charrs show reduced

metabolic rate when switched among amphipods, bloodworms, and *Daphnia* (22). These costs

might be related to irreversible changes in their digestive tracts during development which

increase efficiency on the specific prey they were reared, but seem to be maladaptive on new

prey. Some songbirds, such as American Robins and European Starlings, exhibit costs when

switched between very different categories of foods in their natural diets, such as insects and

fruit (45, 46). These switching costs can be reflected in utilization efficiencies, gut retention

time, and body mass. Cabbage butterflies, *Pieris rapae*, take longer to extract nectar from a

flower species if they are moved to it from a different flower (47). Switching between different

flower species has a learning cost and may be related formation of a search-image for a specific

flower. *Escherichia coli*, the model bacterium shows a distinct lag phase when shifting from one

carbon resource to another (48). The lag phase may be a result of time required to switch on

relevant metabolic genes or related to the ability of the cells to accurately detect the depletion

of the primary carbon resource and presence of the secondary resource (49).

248   We also found that *D. discoideum* amoebas experience resource-mixing costs, where

they proliferate less than expected in some communities with multiple species of prey bacteria.

Our findings are in the opposite direction from the trend in other studies that have looked at the

effect of mixing resources in generalists. Some studies on other protists that have looked at

growth rate in multi-prey communities have found an increase in growth rates (50, 51).

Similarly, a study on marine amphipods found that fitness on mixed diets of algal species and

animal matter was either improved or at least comparable to fitness on the best monospecific

diet (52). A meta-analysis on the effect of diet mixing that included diverse consumers found

that consumers grew better on mixed diets compared to the averages from single-species diet

(53). However, if they considered only defended prey, then mixed diet was not different from

the average of single-species diets. This might help explain our results. There could be negative

synergistic interactions between defensive chemicals of different prey bacteria such that

combined toxins are more detrimental to the amoebas. Another possible explanation of lower

success on mixtures is that the different bacterial species compete and lower their overall

numbers. However, we minimized this effect by not providing nutrients for bacterial growth. It is

not clear why *D. discoideum* shows mixing costs when other taxa do not, but it is consistent with

their switching costs.

It is interesting that we observe resource-mixing costs despite another factor that may

obscure them. That factor is that amoebas may choose to eat the most profitable bacteria first.

All but late measurements would then reflect the growth rate on the best bacterium and any

mixing costs might therefore be obscured. Amoebas do show preferential attraction towards

Gram-negative bacteria compared to Gram-positive bacteria (54). Thus, it would be interesting

to test if *D. discoideum* amoebas can avoid some mixing costs by preferentially eating the most

profitable prey bacteria first, but if they do, our data show that it is apparently not enough to

fully overcome such costs.

These costs of combining resources are consistent with the idea that *D. discoideum*

amoebas use partially different methods to hunt and process different prey. The costs probably

involve changing gene expression. *D. discoideum* amoebas transcribe partially distinct sets of

genes on Gram-positive (*Bacillus subtilis*, *Staphylococcus aureus*) and Gram-negative bacteria (*K.*

pneumoniae, *Pseudomonas aeruginosa*) (55). The transcriptome of *D. discoideum* was also

found to be highly species-specific when tested on 3 species of bacteria: *K. pneumoniae*, *B.*

subtlis, and *Mycobacterium marinum* (56). Another study found that mutations that affect the

ability of *D. discoideum* to grow on different bacteria were highly prey-specific (57). This

suggests that there are different mechanisms for hunting or processing these distantly related

bacteria.

Predation by amoebas is a complex process that can be divided into 4 broad steps:

search, encounter, attack, and digestion (33). The costs of combining resources could arise in

any of these steps. Eukaryotic phagocytes use G-protein coupled receptors (GPCRs) to detect

and chase bacteria and use pattern recognition receptors to recognize and eliminate bacteria

(58, 59). *D. discoideum* is equipped with 61 GPCRs (60). Some GPCRs are involved in the

amoebas' social cycle, but at least one GPCR, the folate receptor fAR1, has been implicated in

both chasing and engulfing *Klebsiella* bacteria (61). Other GPCRs may play a role in other aspects

of prey capture including chase and recognition. *D. discoideum* is also equipped with 22 genes

that encode as many as 4 different types of lysozymes (62). Lysozyme genes play an important

role in digesting bacteria. Knocking out some lysozyme genes generally reduces the ability of

amoebas to feed on all tested bacteria, while deletion of other lysozyme genes reduces growth

only on specific gram-negative bacteria (63). Future research on the mechanisms underlying

these costs would be valuable.

Why can't amoebas evolve one predation technique that works optimally on all prey

bacteria to eliminate the costs of diet generalism? After all, amoebas are a few hundred times

larger than the average bacterium by mass (64). They should be able to overwhelm most

bacteria with their size advantage. But bacteria possess many mechanisms to resist their

eukaryotic predators (33, 34, 65, 66). Morphological adaptations such as formation of

microcolonies, biofilms, and filamentation can prevent attack or ingestion by predators. Bacteria

can modify their membrane properties by changing the lipopolysaccharides on the outer

membrane, secreting an S-layer and sporulating among others that can help them avoid

recognition, ingestion, and digestion by predators. They can also produce a huge array of

secondary metabolites that can deter predation by protists (67, 68). Thus, each bacterial species

may possess a unique combination of defenses that is unlikely to yield to a common predation

strategy. It would be interesting to test how many different predation strategies *D. discoideum*

can employ. *D. discoideum* may treat some groups of bacteria similarly and some as different.

We suspect that environmental heterogeneity plays a large role in the maintenance of a

generalist strategy in *D. discoideum*. Environmental heterogeneity, especially temporal variation

favors generalists (69). The scale of variation can also influence the nature of generalism (70).

Coarse-grained environments may select for early developmental plasticity, but for a fixed

phenotype later in life. Fine-grained environments select for versatile generalists that are

capable of reversing their phenotypic response.

How much environmental heterogeneity amoebas experience in nature is not fully

known. Soil bacterial communities are certainly spatially and temporally variable, such that no

one bacterium may be consistently sufficiently abundant (41). But the scale is important. On a

microscale, soil is made up of small aggregated particles that are connected via a network of air-

and water-filled pores (71). The three-dimensional nature of these particles increases soil

surface area such that as little of 10^{-6} percent of soil surfaces may contain bacteria (72). These

particles generally contain bacterial patches of a limited number of clonal cells and these

patches can be separated by distances that are large on a microbial scale (73). Thus, it is possible

that amoebas are likely to experience switching costs as they move between bacterial patches.

Amoebas can also travel considerable distances during their social cycle through slug migration

and especially through the dispersal of spores by animal vectors (74). This can introduce them to

new soil environments with different bacterial communities. Thus, amoebas are in a selective

environment that probably favors generalists.

Resource-combining costs may be less relevant to consumers that can find one big

resource unit to spend their whole life on, such as phytophagous insects or parasitoid wasps.

However, the generalist parasitoid wasp, *Venturia canescens* shows a cost of resource switching

across generations, perhaps due to epigenetic or maternal effects (75). Consumers that

encounter diverse and defended prey species in their lifetimes are more likely to be plagued by

these costs. One example is wolf spiders that experience costs when fed a mixed diet of toxic

prey (76).

Resource-combining costs are different from standard trade-offs because here

generalists can handle each resource well, but they experience costs when combining different

resources. These costs, like trade-offs, could eventually lead to resource specialization. We have

shown that resource-combining costs are prominent in a resource generalist but it would be

interesting to also study resource specialists. Are they sometimes constrained in their diets less

by being unable to feed on a more general diet, but by the inability to readily switch to another

suitable resource? Are they kept from having a broader diet by resource-mixing costs? We

suggest that resource-combining costs deserve greater consideration than they have received in

the debate on the evolution of generalism and specialism.

Materials and Methods

Growth rate of *D. discoideum* on different species of bacteria isolated from forest soil

environments

We first measured the growth rate of wild clones of *D. discoideum* on *Klebsiella pneumoniae,*

the lab food bacterium, and on 22 species of bacteria found in close-association with *D.*

discoideum isolated by Brock et al. 2018 from forest soil environments (Table 1). These bacteria

were isolated from fruiting bodies of *D. discoideum* that developed from field-collected samples

of soil and deer feces from Mountain Lake Biological Station, Virginia in 2014. All *D. discoideum*

strains used in this paper were also isolated from Mountain Lake Biological Station, Virginia. The

bacterial species identification is based on closest partial 16S BLAST hit by Brock et al. 2018. In

this paper, we refer to the bacteria using names from these closest 16s BLAST hits. We

performed this experiment on 3 *D. discoideum* strains: QS1, QS6, and QS9. We plated 100,000

amoebas with 200μl of 10 OD_{600} bacterial suspension on starving agar plates. We estimated

amoeba numbers at 20 hours and calculated doubling times as:

$$Doubling\ time\ (T_D)\ = ln(2)\ \times\ \frac{20}{ln(N_{t=20}) -\ ln(N_{t=0})}$$

This assay also helped us identify reasonably edible bacteria for other experiments.

Resource-switching cost assay in *D. discoideum* amoebas

To test if resource-switching costs occur when *D. discoideum* amoebas are switched from

Bacterium A to Bacterium B, we first conditioned separate populations of the amoebas to each

species of bacterium (Figure 2, details below for three separate experiments). The switched

treatment for these experiments used amoebas conditioned to Bacterium A and replated with

Bacterium B. We plated 100,000 amoebas conditioned to Bacterium A with 200μl 20 OD_{600}

suspension of Bacterium B on SM/20 plates and then measured amoeba numbers after 3 hours.

The controls for this experiment were identical except that amoebas conditioned to Bacterium B

were replated with Bacterium B. If we observe fewer amoebas in switched plates compared to

control plates, then *D. discoideum* amoebas experience resource-switching costs. We measured

the costs of switching between 3 pairs of bacteria with 2 reciprocal switches for each pair. We

replicated this experiment with 5 *D. discoideum* strains: QS6, QS9, QS14, QS17, QS160.

We did three sets of experiments to rule out the possibility that these costs are instead

due to an evolved response in the amoebas during the conditioning period to either bacterium.

Switched amoebas could experience poor growth on a Bacterium B because of trade-offs

associated with new evolutionary adaptation to the Bacterium A during the conditioning period.

Control amoebas may also experience better growth rates if they have evolved and adapted to

Bacterium B during the conditioning period. We ruled out this complication in three ways.

Effect of conditioning length of amoebas on resource-switching costs

First, we tested if the length of the conditioning period of amoebas on the bacterium affected

resource-switching costs. If costs are due to either adaptation in switched amoebas that results

in evolutionary trade-offs, or to direct improvement in growth rate because of adaptation in

control amoebas, we expect stronger costs with longer conditioning (i.e. evolving) time. Thus,

we conditioned amoebas on a given bacterium for a) 2 days and b) 5 days before measuring

switching costs. In the 2-day treatment, we conditioned the amoebas for 40 hours by plating

200,000 spores of *D. discoideum* with 200ul of 1.5 OD_{600} bacterial suspension on SM/5 plates.

After ~40 hours, we washed off the plates with 10ml of ice cold KK2 to collect the amoebae-

bacteria suspension. Next, we centrifuged the suspension for 3 minutes at 300g at 10°C to spin

down the amoebas and discarded the supernatant containing the bacteria. We resuspended the

amoeba pellet in ice-cold KK2 and washed it off two more times until all the bacteria were

discarded. Finally, we suspended the amoeba pellet in 500μl – 1000μl of ice-cold KK2. We

conditioned the amoebas for the 5-day experiment similarly, but collected amoebas from the

resulting vegetative front after 5 days with a sterile loop and resuspending in 600ul ice-cold KK2.

We performed washing centrifuging steps as described above to thoroughly wash the amoebas

off the conditioning bacterium. A conditioning time of 2 days translates to 10-14 amoeba

divisions and 5 days to 25-35 amoeba divisions on good prey. If resource-switching costs are

because of an evolutionary response, then we expect costs to be stronger for the 5-day

conditioning period than the 2-day period.

Resource-switching costs assay with *D. discoideum* spores

To further distinguish between a temporary switching cost and an evolved response, we tested

whether allowing amoebas to go through the social cycle and subsequent sporulation before

prey bacteria switches erases switching costs (Figure 2). If costs are because of an evolved

response in amoebas, then these changes should still be evident after the social stage. But if the

costs are because of conditioning of the amoebas that results in temporary mismatch in

transcriptional tools on the new bacterium, then undergoing the social cycle should erase most

switching costs. This is because *D. discoideum* experiences a huge turn-over in gene expression

when transitioning from a unicellular lifestyle into the multicellular cycle (Parikh et al. 2010,

Rosengarten et al. 2015). The abundance of almost every transcribed mRNA changes at least 2-

fold throughout development starting from vegetative amoebas to multicellular fruiting bodies

(43). Therefore, any transcriptional conditioning of amoebas towards a bacterium should be

largely erased by development. The switched and control spores are at the same transcriptional

start line of dormancy.

We used the same design as the switching experiment for amoebas described above but

with sporulation at the end of the conditioning phase (~7 days). *D. discoideum* amoebas feed on

bacteria for the first 2-4 days of the conditioning phase, and then transition to the social cycle

once the bacteria are depleted. We then plated 100,000 spores from the resulting fruiting

bodies with bacterial suspensions on SM/20 (Figure 2). Since the spores require some time to

germinate into amoebas and start feeding on the bacteria, we counted the total number of

amoebas on these plates after 36 hours.

Time course data of resource-switching costs in *D. discoideum* amoebas

As a third check on whether what we see is an evolved or a conditioned response, we conducted

a time series study of the amoebas on all three pairs of bacteria to check if the growth rates of

the switched treatments quickly catch up with control growth rates, which would indicate a

physiological lag rather than an evolutionary change. We tracked the number of amoebas in

switched and control plates after 6, 24 and 27 hours from plating. We calculated early growth

rate between 0h and 6h time points and calculated late growth rate between 24h and 27h time

points. We assumed exponential growth and used this formula to calculate growth rate between

time points t1 and t2, where N_t stands for number of amoebas at time t:

432
$$Growth\ rate\ (r)\ =\ \frac{ln(N_{t2})\ -\ ln\ (N_{t1})}{(t2\ -\ t1)}$$

If switching costs are not an evolved response, then switches and controls should not differ in

their late growth rates. We performed this on SM/5 plates with a starting number of 100,000

amoebas with 200μl of 1.5 OD_{600} suspension of the test bacterium on the *D. discoideum* clone

QS9. We replicated this experiment five times.

Resource-mixing costs in *D. discoideum* amoebas

Next, we determined whether *D. discoideum* amoebas experience resource-mixing costs when

feeding on multiple species of bacteria. For this experiment, we chose 5 prey bacteria that were

generally highly edible from our growth rate assay of amoebas on 23 species of bacteria. We

generated 5 single-species communities, 6 three-species communities and 1 five-species

community from the 5 prey bacteria. We made 10 OD_{600} suspensions of each bacterial species

and mixed these suspensions in equal proportions to make a given multi-species suspension. We

used AX4 amoebas that were grown axenically in bacteria-free cultures to preclude any effect of

prior conditioning of amoebas to a given species of bacterium on the assay. We plated 100,000

amoebas on starving agar plates with 200µl of bacterial suspensions and measured amoeba

numbers in all the different treatments after 20 hours. We replicated this experiment 6 times.

We expect bacterial growth and consequently competition to be minimal because we

performed this experiment on starving agar. To test if amoebas experience resource-mixing

costs, we first calculated the expected number of amoebas in multi-prey treatments with data

from our single-prey treatments. For example, the expected number of amoebas in multi-prey

treatment containing *A. tumefaciens*, *S. maltophilia* and *C. testosteroni* is the average of

observed numbers of amoebas in single-prey treatment of those bacteria. We then compared

these expected numbers to our observed number of amoebas in these multi-species treatments

to infer costs.

Statistics:

We performed all statistical analysis in R (version 4.2.1) (77). We used general linear models

(with log link functions for count data) to analyze our data after testing for normality of residuals

using the Shapiro-Wilk's test and examining Q-Q plots. We used the "*emmeans*" package to

calculate estimated effect sizes (Cohen's d, a standardized measure) and 95% confidence

intervals for relevant contrasts (78). We calculated the largest and the smallest percentage

difference between treatments from the 95% C.I range of the estimated marginal means. To test

if *D. discoideum* amoebas experience varying doubling times on different species of bacteria, we

built a linear model with amoeba doubling times as the response variable and the Test

bacterium (23 species of bacteria listed in Table 1) as the fixed factor. We define test bacterium

as the bacterium on which we measured amoeba growth rates. To test if *D. discoideum*

amoebas experience a resource-switching cost on a given bacterium when previously grown on

a different bacterium, we built a linear model with log-transformed amoeba numbers after 3

hours of growth as the response variable and Treatment (Control, Switch), Test bacterium (At,

Pv, Fg, Sm, Sz, Ct), and Conditioning length (2days, 5days) as fixed factors. We also included

interaction effects for Treatment x Test bacterium, Treatment x Conditioning length. For the

similar experiment that includes a spore stage after conditioning, we used a similar linear model

with log transformed amoeba numbers after 36 hours as the response variable and Treatment

and Test bacterium as the fixed factors. We included interaction effects for Treatment x Test

bacterium in this model.

We performed the following statistical tests on time-course data collected on switches.

To test if there are switching costs during early proliferation we used a linear model with early

growth rate calculated between 0 and 6 hours as the response variable and Treatment (Control,

Switch) and Test bacterium (At, Pv, Fg, Sm, Sz, Ct) as fixed factor. To test if resource-switching

costs persist during late proliferation, we used a similar linear model with late growth rate

calculated between 24 and 27 hours as the response variable.

To test if *D. discoideum* amoebas experience resource-mixing costs, we tested if

amoebas performed worse than expected in multi-species prey communities. We built a linear

model with log transformed amoeba numbers at 20 hours as the response variable, Treatment

(categorical variable for whether an observation is expected or observed) and Bacterial

Community (6 Communities) and Experimenter (2 Experimenters) as fixed factors. We included

an interaction effect between Treatment x Bacterial Community. ANOVA tables for all models

are included in the supplement.

Acknowledgments

This material is based upon work supported by the National Science Foundation under grants

IOS 16-56756, DEB 17-53743, and DEB 2237266. We thank Trey Scott for feedback on statistical

analysis. We thank Margaret Steele and Calum Stephenson for comments on the manuscript.

We also thank the Queller-Strassmann lab for their helpful insights on experimental design.

496
497
498   **References**
499

1.   B. D. Moore, W. J. Foley, A review of feeding and diet selection in koalas (Phascolarctos
cinereus). *Aust. J. Zool.* **48**, 317 (2000).

2.   S. R. Beissinger, Hunting behavior, prey selection, and energetics of snail kites in Guyana:
consumer choice by a specialist. *The Auk* **100**, 84–92 (1983).

3.   J. M. McVey, *et al.*, Diets of sympatric red wolves and coyotes in northeastern North
Carolina. *Journal of Mammalogy* **94**, 1141–1148 (2013).

4.   J. M. Schmidt, P. Sebastian, S. M. Wilder, A. L. Rypstra, The nutritional content of prey
affects the foraging of a generalist arthropod predator. *PLOS ONE* **7**, e49223 (2012).

5.   B. Eitzinger, *et al.*, Assessing changes in arthropod predator–prey interactions through
DNA-based gut content analysis—variable environment, stable diet. *Molecular Ecology* **28**,
266–280 (2019).

6.   J. Bascompte, P. Jordano, Plant-animal mutualistic networks: the architecture of
biodiversity. *Annu. Rev. Ecol. Evol. Syst.* **38**, 567–593 (2007).

7.   A. Mougi, M. Kondoh, Diversity of interaction types and ecological community stability.
Science **337**, 349–351 (2012).

8.   R. Svanbäck, D. I. Bolnick, Intraspecific competition affects the strength of individual
specialization: an optimal diet theory method. *Evolutionary Ecology Research* **7**, 993–1012
(2005).

9.   P. R. Guimarães, M. M. Pires, P. Jordano, J. Bascompte, J. N. Thompson, Indirect effects
drive coevolution in mutualistic networks. *Nature* **550**, 511–514 (2017).

10.   N. B. Hardy, S. P. Otto, Specialization and generalization in the diversification of
phytophagous insects: tests of the musical chairs and oscillation hypotheses. *Proceedings*
of the Royal Society B: Biological Sciences **281**, 20132960 (2014).

11.   E. A. Bernays, O. P. J. M. Minkenberg, Insect herbivores: different reasons for being a
generalist. *Ecology* **78**, 1157–1169 (1997).

12.   D. J. Futuyma, G. Moreno, The evolution of ecological specialization. *Annual Review of*
Ecology and Systematics **19**, 207–33 (1998).

13.   J. P. Sexton, J. Montiel, J. E. Shay, M. R. Stephens, R. A. Slatyer, Evolution of ecological
niche breadth. *Annu. Rev. Ecol. Evol. Syst.* **48**, 183–206 (2017).

14.   N. B. Hardy, C. Kaczvinsky, G. Bird, B. B. Normark, What we don't know about diet-breadth
evolution in herbivorous insects. *Annual Review of Ecology, Evolution, and Systematics* **51**,
103–122 (2020).

15.   T. J. Kawecki, Accumulation of deleterious mutations and the evolutionary cost of being a
generalist. *The American Naturalist* **144**, 833–838 (1994).

16.   J. D. Fry, The evolution of host specialization: are trade-offs overrated? *The American*
Naturalist **148**, S84–S107 (1996).

17.   S. J. Agosta, J. A. Klemens, Resource specialization in a phytophagous insect: no evidence
for genetically based performance trade-offs across hosts in the field or laboratory. *Journal*
of Evolutionary Biology **22**, 907–912 (2009).

18.   M. C. Whitlock, The red queen beats the jack-of-all-trades: the limitations on the evolution
of phenotypic plasticity and niche breadth. *The American Naturalist* **148**, S65–S77 (1996).

19.   S. R. X. Dall, I. C. Cuthill, The information costs of generalism. *Oikos* **80**, 197–202 (1997).

20.   H. Cornell, Search strategies and the adaptive significance of switching in some general
predators. *The American Naturalist* **110**, 317–320 (1976).

21.   R. N. Hughes, Optimal diets under the energy maximization premise: the effects of
recognition time and learning. *The American Naturalist* **113**, 209–221 (1979).

22. O. E. Hooker, T. E. Van Leeuwen, C. E. Adams, The physiological costs of prey switching
reinforce foraging specialization. *J Anim Ecol* **86**, 605–614 (2017).

23. D. I. Bolnick, *et al.*, The ecology of individuals: incidence and implications of individual
specialization. *The American Naturalist* **161**, 1–28 (2003).

24. W. J. Freeland, D. H. Janzen, Strategies in herbivory by mammals: the role of plant
secondary compounds. *The American Naturalist* **108**, 269–289 (1974).

25. M. L. Forister, *et al.*, The global distribution of diet breadth in insect herbivores. *PNAS* **112**,
442–447 (2015).

26. J. Jaenike, Host specialization in phytophagous insects. *Annual Review of Ecology and
Systematics* **21**, 243–273 (1990).

27. K. D. Lafferty, A. M. Kuris, Trophic strategies, animal diversity and body size. *Trends in
Ecology & Evolution* **17**, 507–513 (2002).

28. L. Real, T. Caraco, Risk and foraging strategies in stochastic environments. *Ann. Rev. Em!.
Syst.*, 371–90 (1986).

29. E. B. Sherr, B. F. Sherr, Significance of predation by protists in aquatic microbial food webs.
Antonie van Leeuwenhoek **81**, 16 (2002).

30. S. Geisen, *et al.*, Soil protists: a fertile frontier in soil biology research. *FEMS Microbiology
Reviews* **42**, 293–323 (2018).

31. T. Fenchel, *Ecology of Protozoa* (Springer Berlin Heidelberg, 1987).

32. G. Corno, K. Jürgens, Structural and functional patterns of bacterial communities in
response to protist predation along an experimental productivity gradient: top-down and
bottom-up control of bacteria. *Environmental Microbiology* **10**, 2857–2871 (2008).

33. A. Jousset, Ecological and evolutive implications of bacterial defences against predators:
antipredator defences and bacterial ecology. *Environmental Microbiology* **14**, 1830–1843
(2012).

34. W. P. J. Smith, B. R. Wucher, C. D. Nadell, K. R. Foster, Bacterial defences: mechanisms,
evolution and antimicrobial resistance. *Nat Rev Microbiol* **21**, 519–534 (2023).

35. R. H. Kessin, *Dictyostelium: Evolution, Cell Biology, and the Development of Multicellularity*
(Cambridge University Press, 2001).

36. J. E. Strassmann, D. C. Queller, Evolution of cooperation and control of cheating in a social
microbe. *Proceedings of the National Academy of Sciences* **108**, 10855–10862 (2011).

37. K. B. Raper, N. R. Smith, The growth of Dictyostelium discoideum upon pathogenic
bacteria. *Journal of bacteriology* **38**, 431–445 (1939).

579 38. V. E. Paquet, S. J. Charette, Amoeba-resisting bacteria found in multilamellar bodies
580   secreted by *Dictyostelium discoideum:* social amoebae can also package bacteria. *FEMS*
581   *Microbiology Ecology* **92**, fiw025 (2016).

582 39. D. A. Brock, *et al.*, Diversity of free-living environmental bacteria and their interactions
583   with a bactivorous amoeba. *Front Cell Infect Microbiol* **8** (2018).

584 40. S. Kumar, G. Stecher, M. Suleski, S. B. Hedges, Timetree: a resource for timelines,
585   timetrees, and divergence times. *Molecular Biology and Evolution* **34**, 1812–1819 (2017).

586 41. C. L. Lauber, K. S. Ramirez, Z. Aanderud, J. Lennon, N. Fierer, Temporal variability in soil
587   microbial communities across land-use types. *ISME J* **7**, 1641–1650 (2013).

588 42. H. D. Loxdale, G. Lushai, J. A. Harvey, The evolutionary improbability of 'generalism' in
589   nature, with special reference to insects. *Biological Journal of the Linnean Society* **103**, 1–
590   18 (2011).

591 43. A. Parikh, *et al.*, Conserved developmental transcriptomes in evolutionarily divergent
592   species. *Genome Biology* **11**, R35 (2010).

593 44. R. D. Rosengarten, *et al.*, Leaps and lulls in the developmental transcriptome of
594   Dictyostelium discoideum. *BMC Genomics* **16**, 294 (2015).

595 45. D. J. Levey, W. H. Karasov, Digestive responses of temperate birds switched to fruit or
596   insect diets. *The Auk* **106**, 675–686 (1989).

597 46. D. J. Levey, W. H. Karasov, Digestive modulation in a seasonal frugivore, the American
598   robin (Turdus migratorius). *American Journal of Physiology-Gastrointestinal and Liver*
599   *Physiology* **262**, G711–G718 (1992).

600 47. A. C. Lewis, Memory constraints and flower choice in Pieris rapae. *Science* **232**, 863–865
601   (1986).

602 48. J. Monod, The growth of bacterial cultures. *Annual Review in Microbiolgy* **3**, 371–349
603   (1949).

604 49. D. Chu, D. J. Barnes, The lag-phase during diauxic growth is a trade-off between fast
605   adaptation and high growth rate. *Sci Rep* **6**, 25191 (2016).

606 50. L. Gamfeldt, H. Hillebrand, P. R. Jonsson, Species richness changes across two trophic
607   levels simultaneously affect prey and consumer biomass. *Ecology Letters* **8**, 696–703
608   (2005).

609 51. M. Saleem, I. Fetzer, H. Harms, A. Chatzinotas, Diversity of protists and bacteria
610   determines predation performance and stability. *ISME J* **7**, 1912–1921 (2013).

52.   E. Cruz-Rivera, M. E. Hay, The effects of diet mixing on consumer fitness: macroalgae,
epiphytes, and animal matter as food for marine amphipods. *Oecologia* **123**, 252–264
(2000).

53.   J. S. Lefcheck, M. A. Whalen, T. M. Davenport, J. P. Stone, J. E. Duffy, Physiological effects
of diet mixing on consumer fitness: a meta-analysis. *Ecology* **94**, 565–572 (2013).

54.   G. Rashidi, E. A. Ostrowski, Phagocyte chase behaviours: discrimination between Gram-
negative and Gram-positive bacteria by amoebae. *Biol. Lett.* **15**, 20180607 (2019).

55.   W. Nasser, *et al.*, Bacterial discrimination by Dictyostelid amoebae reveals the complexity
of ancient interspecies interactions. *Current Biology* **23**, 862–872 (2013).

56.   O. Lamrabet, *et al.*, Transcriptional responses of Dictyostelium discoideum exposed to
different classes of bacteria. *Front. Microbiol.* **11**, 410 (2020).

57.   B. Stewart, *et al.*, The genetic architecture underlying prey-dependent performance in a
microbial predator. *Nat Commun* **13**, 319 (2022).

58.   V. Jaumouillé, S. Grinstein, Receptor mobility, the cytoskeleton, and particle binding during
phagocytosis. *Current Opinion in Cell Biology* **23**, 22–29 (2011).

59.   P. N. Devreotes, *et al.*, Excitable signal transduction networks in directed cell migration.
Annual Review of Cell and Developmental Biology **33**, 103–125 (2017).

60.   A. J. Heidel, *et al.*, Phylogeny-wide analysis of social amoeba genomes highlights ancient
origins for complex intercellular communication. *Genome Research* **21**, 1882–1891 (2011).

61.   M. Pan, *et al.*, A G-protein-coupled chemoattractant receptor recognizes
lipopolysaccharide for bacterial phagocytosis. *PLOS Biology* **16**, e2005754 (2018).

62.   O. Lamrabet, T. Jauslin, W. C. Lima, M. Leippe, P. Cosson, The multifarious lysozyme
arsenal of Dictyostelium discoideum. *Developmental & Comparative Immunology* **107**,
103645 (2020).

63.   T. Jauslin, *et al.*, How phagocytic cells kill different bacteria: a quantitative analysis using
Dictyostelium discoideum. *mBio* **12**, e03169-20 (2021).

64.   R. Kumar, R. Maurya, S. Saran, Investigating the role of translationally control tumor
protein in growth, development and differentiation of Dictyostelium discoideum. *Frontiers
in Cell and Developmental Biology* **8**, 742 (2020).

65.   C. Matz, S. Kjelleberg, Off the hook – how bacteria survive protozoan grazing. *Trends in
Microbiology* **13**, 302–307 (2005).

66.   D. W. Kikuchi, *et al.*, The evolution and ecology of multiple antipredator defences. *J of
Evolutionary Biology* **36**, 975–991 (2023).

67. O. Tyc, C. Song, J. S. Dickschat, M. Vos, P. Garbeva, The ecological role of volatile and
soluble secondary metabolites produced by soil bacteria. *Trends in Microbiology* **25**, 280–
292 (2017).

68. J. M. Raaijmakers, M. Mazzola, Diversity and natural functions of antibiotics produced by
beneficial and plant pathogenic bacteria. *Annu. Rev. Phytopathol.* **50**, 403–424 (2012).

69. R. Kassen, The experimental evolution of specialists, generalists, and the maintenance of
diversity: Experimental evolution in variable environments. *Journal of Evolutionary Biology*
15, 173–190 (2002).

70. R. Levins, *Evolution in changing environments: some theoretical explorations* (Princeton
University Press, 1968).

71. M. Vos, A. B. Wolf, S. J. Jennings, G. A. Kowalchuk, Micro-scale determinants of bacterial
diversity in soil. *FEMS Microbiol Rev* **37**, 936–954 (2013).

72. I. M. Young, J. W. Crawford, Interactions and self-organization in the soil-microbe complex.
Science **304**, 1634–1637 (2004).

73. N. Nunan, K. Wu, I. M. Young, J. W. Crawford, K. Ritz, Spatial distribution of bacterial
communities and their relationships with the micro-architecture of soil. *FEMS*
Microbiology Ecology **44**, 203–215 (2003).

74. jeff smith, D. C. Queller, J. E. Strassmann, Fruiting bodies of the social amoeba
Dictyostelium discoideum increase spore transport by Drosophila. *BMC Evolutionary*
Biology **14**, 105 (2014).

75. T. S. Jones, A. R. Bilton, L. Mak, S. M. Sait, Host switching in a generalist parasitoid:
contrasting transient and transgenerational costs associated with novel and original host
species. *Ecology and Evolution* **5**, 459–465 (2015).

76. S. Toft, D. H. Wise, Growth, development, and survival of a generalist predator fed single-
and mixed-species diets of different quality. *Oecologia* **119**, 191–197 (1999).

77. A. R Core Team, R. C. Team, *R: A language and environment for statistical computing. R*
Foundation for Statistical Computing, Vienna, Austria. 2012 (2022).

78. R. V. Lenth, *et al.*, *Emmeans: Estimated Marginal Means, aka Least-Squares Means,*
Version 1.8. 1-1 (2022).

Figures and Tables

Figure 1. Doubling time of *D. discoideum* on *K. pneumoniae* and 22 species of soil bacteria closely

associated with *D. discoideum*. Error bars are 95% C.I. The points represent the 3 *D. discoideum*

strains. The circles represent the 8 species of bacteria used in other parts of this study. Bacteria

species identities by Brock et al. 2018 based on closest partial 16S BLAST hit are listed in Table 1.

Figure 2. Schematic on the left shows resource-switching costs experiment in amoebas. Amoebas

in switched treatment are conditioned on bacterium A and are switched to bacterium B for the

experiment. The controls for this experiment are amoebas conditioned to bacterium B plated

with a fresh culture of the same. We also varied conditioning time of amoebas on a given

bacterium (2 days, 5 days). Schematic on the right outlines switching costs experiment on

spores. We used the same basic switching design as described for the amoebas, except we let

the amoebas undergo multicellular development on the conditioning bacterium. We then

collected spores from these fruiting bodies and performed the experiment.

Figure 3: *D. discoideum* amoebas experience resource-switching costs: Mean amoeba numbers

± 95% C. I. of controls (blue) and switches (orange) after 3 hours of growth. The shapes

represent the different conditioning times of amoebas (• - 2 days, ▲ - 5 days). The points

represent the five *D. discoideum* strains. There are fewer amoebas after three hours in switched

treatments compared to controls. The two conditioning times are similar in the magnitude of

switching costs. Interpreting x-axis labels: For example - Sm-Sm refers to amoebas grown on *S.*

maltophilia moved to *S. maltophilia*, while Fg-Sm refers to amoebas grown on *F. ginsengisoli*

moved to *S. maltophilia*.

Figure 4: Switching costs disappear when spore formation precedes prey switch: the switched

and control treatment have similar number of amoebas. Graph shows mean amoeba numbers ±

95% C.I. of controls and switches at 36 hours from plating. The points represent the five *D.*

discoideum strains.

Figure 5: Switching costs occur during early growth period rather than later: **A)** Early growth rate

estimates of controls and switches between 0 and 6 hours from set-up. Switched treatments

have lower early growth rates compared to controls. **B)** Late growth rate estimates of controls

and switches between 24 to 27 hours from set-up for. Graph shows mean growth rates ± 95%

714 C.I. We found no difference in growth rates switches and controls at the later time points. The

715 points represent 5 technical replicates of *D. discoideum* clone QS9.

Figure 6: *D. discoideum* amoebas experience resource-mixing costs in some multi-species

bacterial communites: Ratios of observed vs. expected amoeba numbers in different bacterial

communities. A ratio lower than one indicates costs of resource mixing. The blue dot marks the

mean. At- *Agrobacterium tumefaciens*, Bw- *Buttiauxella warmboldiae*, Ct - *Comamonas*

testosteroni, Sm - *Stenotrophomonas maltophilia*, Pk - *Pseudomonas kuykendallii*.

Table 1: List of 23 species of bacteria used in *D. discoideum* growth rate experiments. *
Species identification of bacterial isolates by Brock et al. 2018 based on closest partial 16S
BLAST hit.

Strain number	Closest 16S sequence	Phylum/Class	Abbreviation
Lab clone	*Klebsiella pneumoniae*	Gammaproteobacteria	Kp
14P 8.1.1	*Pseudomonas kuykendallii*	Gammaproteobacteria	Pk
18P 8.2.2	*Buttiauxella warmboldiae*	Gammaproteobacteria	Bw
18P 2.2.1	*Agrobacterium tumefaciens*	Alphaproteobacteria	At
20P 9.1.2	*Agrobacterium rubi*	Alphaproteobacteria	Ar
14P 4.3.1	*Serratia liquefaciens*	Gammaproteobacteria	Sl
14P 6.2.3	*Stenotrophomonas maltophilia*	Gammaproteobacteria	Sm
18P 6.2.3	*Comamonas testosteroni*	Betaproteobacteria	Ct
20P 10.2.4	*Variovorax boronicumulans*	Betaproteobacteria	Vb
14P 8.1.4	*Comamonas kerstersii*	Betaproteobacteria	Ck
20P 3.2.2	*Flavobacterium ginsengisoli*	Bacteroidetes	Fg
20P 9.1.1	*Shinella zoogloeoides*	Alphaproteobacteria	Sz
20P 6.1.2	*Brucella papionis*	Alphaproteobacteria	Bp
5S 2.1.1	*Staphlycoccus saprophyticus*	Firmicutes	Ss
5P 5.1.1	*Pseudomonas vranovensis*	Gammaproteobacteria	Pv
18P 8.1.1	*Achromobacter aegrifaciens*	Betaproteobacteria	Aa
14P 5.3.2	*Pseudomonas migulae*	Gammaproteobacteria	Pm
20P 10.2.1	*Sphingobacterium ginsensidimutans*	Bacteroidetes	Sg
20P 2.1.2	*Pandoraea oxalativorans*	Betaproteobacteria	Po
14P 4.3.2	*Chryseobacterium rhizosphaerae*	Bacteroidetes	Cr
14P 6.2.1	*Microbacterium maritypicum*	Actinobacteria	Mm
20P 7.2.1	*Bacillus aryabhattai*	Firmicutes	Ba
20P 2.1.1	*Paenibacillus pabuli*	Firmicutes	Pp

Supplemental information:

Figure S1: Estimated effect size (Cohen's d) ± 95% C.I. of amoeba numbers of the switched

treatment minus the control treatment. **A)** shows that switched treatment in all test bacterium

generally had a roughly similar negative effects on amoeba numbers. Plot **B)** shows that

switched treatments of the different conditioning time had a similar negative effect on amoeba

number compared to controls.

Figure S2: Estimated effect size (Cohen's d) ± 95% C.I. of observed – expected amoeba numbers

in resource-mixing experiment averaged across observers. We see support for mixing costs

overall and expected trend in four out of the six bacterial communities.

741 **Table S1: ANOVA of amoeba growth rate assay**

742 Formula: anova(lm (log(amoeba_number) ~ Test_bacterium))

	Df	Sum sq	Mean Sq	F_value	Pr(>F)
Test_bacterium	22	299.346	13.3794	24.949	2.20E-16
Residuals	46	24.668	0.5363		

744 **Table S2: ANOVA of amoeba switching assay**

745 Formula: anova(lm(log(amoeba_number) ~ Treatment x Test_bacterium + Treatment x
746 Conditioning_length)

	Df	Sum sq	Mean Sq	F_value	Pr(>F)
Treatment	1	0.965	0.9648	27.621	7.69E-07
Test_bacterium	5	4.958	0.9917	28.392	2.00E-16
Conditioning_length	1	0.057	0.0567	1.623	0.205
Treatment x Test_bacterium	5	0.146	0.0293	0.838	0.526
Treatment x Conditioning_length	1	0.019	0.019	0.543	0.463
Residuals	106	3.702	0.0349		

748 **Table S3: ANOVA of spore switching assay**

749 Formula: anova(lm(log(amoeba_number) ~ Treatment x Test_bacterium + Treatment x
750 Conditioning_length))

	Df	Sum sq	Mean Sq	F_value	Pr(>F)
Treatment	1	0.03	0.03	0.264	0.61
Test_bacterium	5	106.78	21.355	186.402	<2e-16
Treatment x Test_bacterium	5	0.68	0.136	1.191	0.328
Residuals	48	5.5	0.115		

753 **Table S4: ANOVA of early growth rate data**

754 Formula: anova(lm(log(amoeba_number) ~ Treatment + Test_bacterium))

	Df	Sum sq	Mean Sq	F_value	Pr(>F)
Treatment	1	0.012	0.012	12.988	0.0006923
Test_bacterium	5	0.043	0.009	9.561	1.52E-06
Residuals	53	0.048	0.001		

757 **Table S5: ANOVA of late growth rate data**

758 Formula: anova(lm(log(amoeba_number) ~ Treatment + Test_bacterium))

	Df	Sum sq	Mean Sq	F_value	Pr(>F)
Treatment	1	0.000	0.000	0.015	0.9045156
Test_bacterium	5	0.123	0.025	5.469	0.0003944
Residuals	53	0.239	0.005		

761 **Table S6: ANOVA of resource-mixing data**

762 Formula: anova(lm(log(amoeba_number)~Treatment *Community + Experimenter))

	Df	SumSq	MeanSq	F-value	Pr(>F)
Treatment	1	0.4564	0.4564	11.8394	0.001071
Community	5	1.6847	0.3369	8.74	3.01E-06
Experimenter	1	24.3675	24.3675	632.0609	2.20E-16
Treatment x Community	5	0.4929	0.0986	2.5571	0.036767
Residuals	59	2.2746	0.0386		