Title: Deficiency of Orexin Receptor Type 1 in Dopaminergic Neurons Increases Novelty-

Induced Locomotion and Exploration

Xing Xiao[1], Gagik Yeghiazaryan[2,3], Anna L. Cremer[4], Heiko Backes[4], Peter Kloppenburg[2,3],

and A. Christine Hausen[1*]

1   Max Planck Institute for Metabolism Research, Department of Neuronal Control of Metabolism,

Gleueler Strasse 50, 50931 Cologne, Germany

2   Excellence Cluster on Cellular Stress Responses in Aging Associated Diseases (CECAD) and

Center for Molecular Medicine Cologne (CMMC), University of Cologne, Joseph-Stelzmann-

Strasse 26, 50931 Cologne, Germany

3   Department of Biology, Institute for Zoology, University of Cologne, Zülpicher Strasse 47b,

50674 Cologne, Germany

4   Max Planck Institute for Metabolism Research, Multimodal Imaging of Brain Metabolism Group,

Gleueler Strasse 50, 50931 Cologne, Germany

* Correspondence: Christine.hausen@sf.mpg.de

Abstract

Orexin signaling in the ventral tegmental area and substantia nigra promotes locomotion and reward processing, but it is not clear whether dopaminergic cells directly mediate these effects. We show that dopaminergic neurons in these areas mainly express orexin receptor subtype 1 (Ox1R). In contrast, only a minor population in the medial ventral tegmental area express orexin receptor subtype 2 (Ox2R). To analyze the functional role of Ox1R signaling in dopaminergic neurons, we deleted Ox1R specifically in dopamine transporter-expressing neurons of mice and investigated the functional consequences. The deletion of Ox1R increased locomotor activity and exploration during exposure to novel environments or when intracerebroventricularely injected with orexin. Spontaneous activity in home cages, anxiety, reward processing, and energy metabolism did not change. Positron emission tomography imaging revealed that Ox1R signaling in dopaminergic neurons affected distinct neural circuits depending on the stimulation mode. In line with an increase of neural activity in lateral paragigantocellular nucleus (LPGi) of Ox1R$^{\Delta DAT}$ mice, we found that dopaminergic projections innervate the LPGi in regions where the inhibitory dopamine receptor subtype D2 but not the excitatory D1 subtype resides. These data suggest a crucial regulatory role of Ox1R signaling in dopaminergic neurons in novelty-induced locomotion and exploration.

Key words: orexin, Ox1R, Ox2R, dopamine, locomotion, exploration, arousal

Introduction

Orexin neurons are exclusively located in the lateral hypothalamus (LH) and adjacent regions in the brain, including perifornical area, and dorsomedial and posterior hypothalamus (Soya & Sakurai, 2020). Abundant orexin fibers innervate the ventral tegmental area (VTA) and substantia nigra (SN), which together contain the majority of dopaminergic neurons in humans and rodents (Fadel & Deutch, 2002; Hrabovszky et al., 2013; Peyron et al., 1998). There are also some dopaminergic neurons in other brain regions, although not all of these neurons express the dopamine transporter (DAT) (Koblinger et al., 2018; Sharples et al., 2014; Turiault et al., 2007). Both the orexin and dopaminergic systems control food intake, locomotor activity, reward processing as well as energy metabolism (Howe & Dombeck, 2016; Inutsuka & Yamanaka, 2013; Narayanan et al., 2010; Palmiter, 2007; Tsujino & Sakurai, 2013). The nigrostriatal dopaminergic system, containing the dopaminergic projections from SN to the dorsal striatum, predominantly regulates action selection and exploratory behaviors, while the mesolimbic and mesocortical dopaminergic systems, which consist of the dopaminergic pathways from the VTA to ventral striatum and prefrontal cortex, predominantly regulate motivation, cognition, reward, and aversion signaling (Roeper, 2013). Orexin A injection into the VTA increases cocaine- and morphine-related reward processing (Mahler et al., 2012), and intra-SN injection of orexin A induces hyperlocomotion and stereotypic behaviors (Liu et al., 2018; Nakamura et al., 2000). On the other hand, systemic injection of dopamine receptor antagonists reduces orexin A-induced increase of hyperlocomotion, stereotypic and grooming behaviors (Nakamura et al., 2000). However, whether the above effects of orexin signaling in the VTA and SN is mediated by dopaminergic or non-dopaminergic cells is not clear yet.

The orexin system consists of the two peptides orexin A and B, which are derived from the precursor, prepro-orexin (de Lecea et al., 1998; Sakurai et al., 1998). Both are endogenous ligands for two G-protein-coupled receptors, orexin receptor subtype 1 (Ox1R), and 2 (Ox2R) (Sakurai et al., 1998). Orexin A activates both receptors, while orexin B predominantly activates Ox2R (Sakurai et al., 1998). The mRNA detection by *in situ* hybridization revealed the expression of both receptors in VTA and SN (Marcus et al., 2001). However, the cell-type specific expression of Ox1R and Ox2R in the VTA and SN is not well understood. Electrophysiological studies showed that orexin receptors are present in

dopaminergic VTA neurons (Baimel et al., 2017; Korotkova et al., 2003; Tung et al., 2016), whereas

data for the SN seem controversial (Korotkova et al., 2002; Liu et al., 2018).

Here, we aim to map the expression patterns of Ox1R and Ox2R in dopaminergic VTA and SN

neurons and investigate the functional role of Ox1R using conditional knockout mice.

<div align="center">**Results**</div>

Ox1R is the predominant orexin receptor in dopaminergic neurons

To analyze the expression patterns of orexin receptors in dopaminergic VTA and SN neurons, we

performed fluorescent *in situ* hybridization (RNAscope) of tyrosine hydroxylase (*Th*), Ox1R (*Hcrtr1*)

Figure 1 with 2 supplements

Figure 1. Ox1R predominantly mediates orexin-induced activation of dopaminergic neurons in the substantia nigra (SN).

Representative images of RNAscope *in situ* hybridization in SN and paranigral (PN), parainterfascicular (PIF) and parabrachial (PBP) subnuclei of the ventral tegmental area (VTA), of **(A)** a control and **(B)** an Ox1R$^{\Delta DAT}$ mouse. Amplifications of the **(C)** left and **(D)** right marked areas in white squares in (A). Amplifications of the **(E)** left and **(F)** right marked area in white squares in (B). White, tyrosine hydroxylase (Th); magenta, Ox1R; green, Ox2R. Scale bar: 100 μm (A, B) or 50 μm (C-F). Representative images of an overview in SN and VTA were shown in figure 1—figure supplement 1. Percentages of **(G)** Ox1R and **(H)** Ox2R positive neurons in Th positive neurons. IF, interfascicular subnuclei of VTA. Control, n = 3; Ox1R$^{\Delta DAT}$, n = 4. Data are represented as means ± SEM. **p < 0.01; ****p < 0.0001; as determined by two-way ANOVA followed by Sidak's post hoc test. **(I)** Schematic illustration of Ox1R and Ox2R expression in dopaminergic neurons in the VTA and SN. 3 control mice were analyzed and N indicates the mean dopaminergic neuron numbers in the respective subregions. Magenta, dopaminergic neurons only expressing Ox1R; green, dopaminergic neurons only expressing Ox2R; yellow, dopaminergic neurons expressing both Ox1R and Ox2R; grey, dopaminergic neurons expressing neither Ox1R or Ox2R. **(J-L)** Effect of orexin A on dopaminergic SN neurons analyzed by Ca^{2+} imaging with GCaMP6. Recordings were performed in brain slices from control and Ox1R$^{\Delta DAT}$ male mice with GCaMP6 expressed in dopaminergic SN neurons for Ca^{2+} imaging. **(J)** Schematic illustration of the SN. The red frame indicates the region where calcium imaging was performed. **(K, L)** Orexin A effect on $[Ca^{2+}]_i$ of dopaminergic SN neurons. **(K)** Heat maps of 5 individual dopaminergic SN neurons from control (top) and Ox1R$^{\Delta DAT}$ mice. The recordings show the responses to 100 nM orexin A. The dashed lines indicate the range where the responses were quantified. **(L)** Mean Ca^{2+} responses upon 100 nM orexin A application. Data are shown as the percentage of the maximal response to high K^+ saline. Bar graphs represent means ± SEM. The stacked bar shows the percentage of individual dopaminergic neurons in control mice in which the increase in $[Ca^{2+}]_i$ was larger than 3 x SD of the control, thus defining them as orexin A responsive (see Materials and Methods). The significance of mean changes in $[Ca^{2+}]_i$ during orexin A application were compared using one-sample t-tests. ****p < 0.0001. n values are given in brackets.

and Ox2R (*Hcrtr2*) in control (Ox1R flox/flox, DAT wt/wt) mice and Ox1R$^{\Delta DAT}$ (Ox1R fl/fl, DAT-Cre tg/wt) mice, from the same breeding crossing mice carrying a loxP-flanked Ox1R-allele (Ox1R fl/fl) with DAT-Cre mice (Ekstrand et al., 2007).

In control mice, Ox1R was expressed in 44 - 58% of dopaminergic neurons in all VTA and SN sub-regions, including the interfascicular (IF), paranigral (PN), parainterfascicular (PIF) and parabrachial (PBP) subnuclei of the VTA and SN. In contrast, Ox2R-positive dopaminergic neurons were mainly located in the medial VTA, including IF, PN, and PIF, with 25 to 42% of dopaminergic neurons expressing the Ox2R (Figure 1A, C, D, G, H; Figure 1—figure supplement 1A). While the above data were calculated as the mean percentages in the investigated mice, we also calculated the mean dopaminergic neuron numbers, separately for those expressing both or exclusively one of the

orexin receptor types. Thus, we could analyze the distribution of Ox1R-, Ox2R-, Ox1R and Ox2R-, and none (of orexin receptors)-expressing dopaminergic neurons in each VTA subregions and SN, which was shown in the schematic illustration with pie charts (Figure 1I). It showed similar results to the above: 47 - 59% of dopaminergic neurons expressed Ox1R in all VTA and SN sub-regions, and Ox2R was mainly expressed by 26 - 41% of dopaminergic neurons in the medial VTA (Figure 1I). 19 - 25% of dopaminergic neurons in the medial VTA expressed both Ox1R and Ox2R (Figure 1I). Of note, abundant Ox1R and Ox2R expression was present in Th-negative neurons in the VTA and SN and neighboring regions (Figure 1A, C, D, Figure 1—figure supplement 1A).

Since the vast majority of dopaminergic neurons express Ox1R, we focused on the functional analysis of Ox1R signaling in dopaminergic neurons. Ox1R was specifically inactivated in dopaminergic (DAT-expressing) neurons of mice (Ox1R$^{\Delta DAT}$). In Ox1R$^{\Delta DAT}$ mice, Ox1R expression in dopaminergic neurons was significantly reduced and remained unchanged in Th-negative cells. In contrast, the Ox2R expression patterns remained similar to that in the control mice, indicating the successful and specific knockout of Ox1R in dopaminergic neurons (Figure 1B, E-H; Figure 1—figure supplement 1B). To test if orexin activates dopaminergic SN neurons and to functionally validate the deletion of Ox1R signaling, we performed Ca^{2+} imaging on dopaminergic SN neurons from control and Ox1R$^{\Delta DAT}$ mice using the genetically encoded Ca^{2+} indicator GCaMP6 expressed in dopaminergic (DAT expressing) neurons (Figure 1J-L, Figure1—figure supplement 2). During the experiments, GABAergic and glutamatergic synaptic input was blocked. The orexin A-induced Ca^{2+} responses were expressed as a percentage of the maximum fluorescence induced by KCl (Figure 1K, L, Figure1—figure supplement 2). Bath-applied 100 nM orexin A significantly increased [Ca^{2+}]$_i$ in dopaminergic SN control neurons ($p < 0.0001$, n = 71, one-sample t-test). An analysis on the level of individual cells (3 sigma criterion) showed that ~60% of dopaminergic SN neurons responded to 100 nM orexin A. In dopaminergic SN neurons from Ox1R$^{\Delta DAT}$ mice, 100 nM orexin A had no effect ($p = 0.5$, n = 86, one-sample t-test) (Figure 1K, L).

In conclusion, Ox1R is the main orexin receptor expressed in dopaminergic VTA and SN neurons and mediates their activation by orexin. Therefore, we wanted to determine further the functional role of Ox1R signaling in dopaminergic neurons. By comparing data from control and

141 Ox1R$^{\Delta DAT}$ mice, we first identified behaviors modulated by orexin via dopaminergic neurons. In a

142 second step, we used PET imaging to define and map neuronal pathways modulated by orexin..

Ox1R in dopaminergic neurons regulates novelty-induced locomotion and exploration

144 The behavioral test included an analysis of locomotor behavior, novelty exploration, reward processing,

145 anxiety, and energy homeostasis. Since sex differences exist in orexin- and dopamine-mediated

146 behaviors (Durairaja & Fendt, 2020; Freeman et al., 2021; Lippert et al., 2020), we analyzed behavioral

147 **Figure 2** with 5 supplements

Figure 2. Deficiency of Ox1R in dopaminergic neurons increases locomotion and exploration behaviors.
(A) Total traveling distance, (B) ambulating time, (C) vertical activity and (D) stereotypic activity of male mice in the open field test. Control, n = 14; Ox1R$^{\Delta DAT}$, n = 15. (E) Total traveling distance, (F) ambulating time, (G) vertical activity and (H) stereotypic activity of female mice in the open field test. Control, n = 14; Ox1R$^{\Delta DAT}$, n = 12. Locomotor activity upon intracerebroventricular (ICV) injection of saline (NS) or orexin A in (I) male and (K) female control and Ox1R$^{\Delta DAT}$ mice. (J, L) Quantification of the area under curve (AUC) after orexin injection to (J) male and (L) female mice. Male, n = 8; female-control, n = 9; female-Ox1R$^{\Delta DAT}$, n = 10. Data are represented as means ± SEM. *p < 0.05; **p < 0.01; ****p < 0.0001; as determined by unpaired two-tailed Student's t-test (A-C, E-H), or two-way ANOVA followed by Sidak's post hoc test (J, L).

phenotypes in both males and females. Compared to controls, Ox1R$^{\Delta DAT}$ mice did not show significant

changes in spontaneous locomotor activity in home cages (Figure 2—figure supplement 1). When

exposed to a novel open field, both male and female Ox1R$^{\Delta DAT}$ mice exhibited an increase in locomotion

and exploration, with increased traveling distance, ambulating time, and vertical activity in the open

field test (Figure 2A-C, E-G). Female Ox1R$^{\Delta DAT}$ mice also showed increased stereotypic activity (Figure

2D, H). Furthermore, intracerebroventricular (ICV) injection of orexin A (1 nmol) induced a more

pronounced increase of locomotor activity in both female and male Ox1R$^{\Delta DAT}$ mice, compared to control

mice (Figure 2I-L).

In a hole-board test, female Ox1R$^{\Delta DAT}$ mice showed increased nose pokes into the holes in early

(1st and 2nd) sessions compared to control mice (Figure 2—figure supplement 2B). This effect

disappeared later when mice became familiar with the environment (3rd – 8th session) (Figure 2—figure

supplement 2B). In the novel object recognition test, female Ox1R$^{\Delta DAT}$ mice spent more time with each

of the two identical objects in an open field compared to control mice (Figure 2—figure supplement 2E).

One hour later, when female mice were exposed to the same environment except that one object was

replaced by a novel object, there was no significant difference between control and Ox1R$^{\Delta DAT}$ mice in

exploring the novel object (Figure 2—figure supplement 2F). In contrast, male Ox1R$^{\Delta DAT}$ mice did not

behave differently in the hole-board and novel object recognition tests from controls (Figure 2—figure

supplement 2A, C, D). This indicates that Ox1R signaling in dopaminergic neurons is more important

for females in novelty-induced exploration, which was independent of novel object discrimination.

The behavioral changes seemed not related to alterations in anxiety, since there were no

detectable changes in anxiety-related behaviors in an open field test, dark/light box test or 0-maze test

(Figure 2—figure supplement 3).

Ox1R deletion in dopaminergic neurons did not change energy metabolism and reward processing

The orexin and dopaminergic systems are well known to regulate energy metabolism and reward

processing (Howe & Dombeck, 2016; Inutsuka & Yamanaka, 2013; Narayanan et al., 2010; Palmiter,

2007; Tsujino & Sakurai, 2013). Therefore, we measured the impact of Ox1R deletion in dopaminergic

neurons on metabolism and reward-related behaviors. We found that, compared to control mice, neither

male or female Ox1R$^{\Delta DAT}$ mice showed significant changes in the two-bottle preference test (comparing

sweet solution and water drinking), conditional place preference test (comparing cocaine and saline

injection) or energy metabolism, including body weight, fat mass, respiratory exchanging ratio, energy

expenditure, feeding behaviors, glucose tolerance and insulin sensitivity (Figure 2—figure supplement

4 and 5).

Ox1R in dopaminergic neurons regulates specific neural pathways under different conditions

194 To explore and map the underlying neuronal pathways of the orexin-mediated behaviors unbiasedly,

195 we performed positron emission tomography (PET) imaging of control and Ox1R$^{\Delta DAT}$ mice upon ICV

196 injection of orexin A or saline (control).

197 **Figure 3** with 2 supplements

Figure 3. PET imaging studies comparing Ox1R$^{\Delta DAT}$ and control mice.
3D maps of p-values in PET imaging studies comparing Ox1R$^{\Delta DAT}$ and control mice, after
intracerebroventricular (ICV) injection of **(A)** saline (NS) and **(B)** orexin A. Brain areas with significant
changes are indicated. Control-NS, n = 8; control-orexin, n = 6; Ox1R$^{\Delta DAT}$, n = 8. M2, secondary motor
cortex; MPA, medial preoptic area; Pir, piriform cortex; IEn, intermediate endopiriform claustrum; DEn,
dorsal endopiriform claustrum; VEn, ventral endopiriform claustrum; LSS, lateral stripe of the striatum;

BNST, the dorsal bed nucleus of the stria terminalis; HDB, nucleus of the horizontal limb of the diagonal band; MCPO, magnocellular preoptic nucleus; S1Sh, primary somatosensory cortex, shoulder region; S1HL, primary somatosensory cortex, hindlimb region; S1BF, primary somatosensory cortex, barrel field; S1Tr, primary somatosensory cortex, trunk region; V1, primary visual cortex; V2L, secondary visual cortex, lateral area; SubCV, subcoeruleus nucleus, ventral part; Gi, gigantocellular reticular nucleus; IRt, intermediate reticular nucleus; LPGi, lateral paragigantocellular nucleus; Sp5O, spinal trigeminal nucleus, oral part; Sp5I, spinal trigeminal nucleus, interpolar part; sp5, spinal trigeminal tract.

In Ox1R$^{\Delta DAT}$ mice under control conditions (saline injection), we observed higher neuronal activity around the medial preoptic area (MPA), piriform cortex (Pir), endopiriform claustrum, lateral stripe of the striatum (LSS), ventral part of subcoeruleus nucleus (SubCV), spinal trigeminal nucleus, spinal trigeminal tract (sp5), the intermediate reticular zone (IRt) and the gigantocellular reticular nucleus (Gi) than control mice (Figure 3A, Figure 3—figure supplement 1). Compared to control mice, central application of orexin (1 nmol, ICV) to Ox1R$^{\Delta DAT}$ mice induced higher neuronal activity around the lateral paragigantocellular nucleus (LPGi), nucleus of the horizontal limb of the diagonal band (HDB) and magnocellular preoptic nucleus (MCPO), and lower activity around secondary motor cortex (M2), the dorsal bed nucleus of the stria terminalis (BNST), the hindlimb, shoulder, and trunk regions and barrel field of primary somatosensory cortex (S1HL, S1Sh, S1Tr, and S1BF), the lateral area of secondary visual cortex (V2L), primary visual cortex (V1), Pir and dorsal endopiriform claustrum (DEn) (Figure 3B, Figure 3—figure supplement 1). PET imaging did not reveal significant changes in the VTA and SN (Figure 3).

Postmortem analysis of c-Fos staining revealed low c-Fos expression in dopaminergic neurons in the VTA and SN. In contrast, clear orexin-induced c-Fos activity was observed in non-dopaminergic cells (Figure 3—figure supplement 2). Further studies would be needed to investigate how orexin directly and indirectly affect the VTA and SN neurons in the brain.

Dopamine receptors in the dorsal BNST and LPGi

Based on the PET imaging findings, we analyzed the dopaminergic pathways into the dorsal BNST and LPGi that could have been affected by the Ox1R. To this end, we injected Cre-dependent AAV-EYFP bilaterally into the VTA and SN of DAT-Cre (DAT^{EYFP}) and Ox1R$^{\Delta DAT}$ (DAT$^{\Delta Ox1R;\ EYFP}$) mice, and discovered dopaminergic projections in the dorsal BNST and LPGi (Figure 4—figure supplement 1), suggesting that VTA and SN dopaminergic neurons could directly modulate neuronal activity in the

dorsal BNST and LPGi. To further explore this idea, we assessed expression of Fos and dopamine

receptors in the dorsal BNST and LPGi by immunostaining and RNAscope experiments.

Figure 4 with 1 supplement

Figure 4. c-Fos and dopamine receptors in the lateral paragigantocellular nucleus (LPGi) and the

dorsal bed nucleus of the stria terminalis (BNST).

(A) Representative images of c-Fos and tyrosine hydroxylase (Th) staining in LPGi of control and
Ox1R$^{\Delta DAT}$ mice injected (ICV) with saline (NS) and orexin A. Quantification of **(B)** Th fluorescence
and **(C)** c-Fos positive neurons in LPGi. **(D)** Representative images of D1 and D2 subtypes of dopamine
receptor (DRD1 and DRD2) in LPGi of control and Ox1R$^{\Delta DAT}$ mice. **(E)** Quantification of DRD2
fluorescence in LPGi. **(F)** Representative images of a negative control staining of DRD1 and DRD2 in
LPGi of control mice, and **(G)** a positive control staining around the lateral ventricle (LV). **(H)**
Representative images of c-Fos and Th staining in the dorsal BNST of control and Ox1R$^{\Delta DAT}$ mice
injected (ICV) with saline or orexin A. Quantification of **(I)** Th fluorescence and **(J)** c-Fos positive
neurons in dorsal BNST. **(K)** Representative images of DRD1 and DRD2 in the dorsal BNST of control
and Ox1R$^{\Delta DAT}$ mice. Quantification of **(L)** DRD1 and **(M)** DRD2 fluorescence in dorsal BNST. Scale
bar: 200 µm (D, F, G, K), 100 µm (A, H) or 50 µm (insertions in H). Control, n = 3; Ox1R$^{\Delta DAT}$, n = 4.
Cyan, Th; red, c-Fos (A, H). Magenta, DRD1; cyan, DRD2; blue, dapi (D, F, G, K). Data are represented
as means ± SEM. *$p < 0.05$; ***$p < 0.001$; as determined by unpaired two-tailed Student's t-test (L, M),
or two-way ANOVA followed by Sidak's post hoc test (C, J).

257   Orexin injection (1 nmol, ICV) induced higher c-Fos expression levels in LPGi of male

Ox1R$^{\Delta DAT}$ mice compared to control mice (Figure 4A, C). Only the inhibitory D2 but not excitatory D1

subtypes of dopamine receptors (DRD2, DRD1) could be detected in LPGi (Figure 4D, E). Both

receptors were detected around the lateral ventricle (as positive control), whereas no signal was

detectable in LPGi incubated with negative control probe for mice (Figure 4F, G). Th-positive fibers

were detected in LPGi but not Gi (Figure 4A, B), consistent with previous reports (Kitahama et al.,

2000). These data support the hypothesis that Ox1R deletion in dopaminergic neurons results in the

disinhibition of dopaminergic signaling in LPGi.

265   Orexin injection increased c-Fos expression in the dorsal BNST in control but not in Ox1R$^{\Delta DAT}$

266 mice (Figure 4H, J). The expression of DRD1 was significantly reduced, and DRD2 expression levels

267 showed a tendency towards a decrease in dorsal BNST of Ox1R$^{\Delta DAT}$ mice (Figure 4K-M). The Th

268 expression levels were not changed between genotypes (Figure 4H, I). This indicates that dorsal BNST

269 could be less activated, due to the decrease of DRD1 and the lack of orexin-mediated activation of

270 dopaminergic neurons.

Discussion

The main findings of this study are based on RNAscope and optophysiological experiments. These data

show that Ox1R is the most abundant orexin receptor in the midbrain and the only orexin receptor in

the SN. Ca^{2+} imaging experiments on SN dopaminergic neurons from control and $Ox1R^{\Delta DAT}$ mice

functionally confirm these results. They show that orexin-induced activation of SN dopaminergic

neurons is exclusively mediated by Ox1R. In addition, the behavioral experiments, PET imaging and

Fos analyses provide important evidence of these data's systemic relevance. They should serve as a basis

for developing testable working hypotheses on the systemic physiological role of orexin signaling.

Ox1R deficiency in DAT-expressing neurons causes an increase in novelty-induced locomotion and

exploration without altering energy metabolism and anxiety- or reward-processing-related behaviors.

PET imaging, and Fos and DRD expression analysis provide some indications about the underlying

mechanisms.

Earlier studies showed that orexin-induced activation of VTA dopaminergic neurons (Baimel

et al., 2017; Korotkova et al., 2003; Tung et al., 2016). At the end of whole-cell patch clamp recordings,

mRNA measurement of the individual recorded VTA neurons revealed fewer Ox2R- than Ox1R-

positive dopaminergic neurons (Korotkova et al., 2003), which is in line with our findings. While our

data clearly show an orexin A-induced activation of dopaminergic SN neurons when GABAergic and

glutamatergic synaptic input is blocked, previous *in vitro* (Korotkova et al., 2002) or *in vivo*

electrophysiological recordings (Liu et al., 2018) studies were controversial about whether orexin A

could activate dopaminergic neurons in the SN. Since Ox1R and Ox2R are expressed by non-

dopaminergic neurons around the SN, recordings in the absence of synaptic blockers, could detect a

mixed direct and indirect orexin-induced response in dopaminergic neurons (Nair-Roberts et al., 2008).

This may partially explain their controversial findings.

We found that Ox1R deficiency in dopaminergic neurons caused an increase in novelty-induced

locomotion and exploration, which could indicate elevated arousal levels and hyperactivity.

Interestingly, children with attention deficit and hyperactivity disorder (ADHD) have lower orexin A

but unaltered orexin B levels in plasma compared to the healthy controls (Baykal et al., 2019). ADHD

patients exhibit impaired dopaminergic functions, and medicine improving dopaminergic signaling is

used to treat ADHD in clinics (Feldman & Reiff, 2014; Volkow & Swanson, 2013). Ox1R signaling in

dopaminergic neurons might be important to limit novelty-induced arousal and hyperactivity, which are

related to task performance and learning capability.

We did not observe significant impact of Ox1R signaling in dopaminergic neurons on reward

processing. Previous studies reported that Ox1R signaling in VTA promoted reward processing with

pharmacological methods (Tsujino & Sakurai, 2013). The difference could have resulted from the

involvement of non-dopaminergic neurons in VTA or different behavioral models. It would be

interesting to identify the effect under other conditions, such as obesity and operant reward learning, in

the future.

The PET imaging results suggest that Ox1R signaling in dopaminergic neurons affects separate

and distinct neuronal circuits, depending on the mode of stimulation. Orexin and dopaminergic systems

are known to increase arousal in mammals' sleep-wake transitions, locomotor activity, and motivation

(Berke, 2018; Berridge et al., 2010; Furlong et al., 2009; Mahler et al., 2014). The orexin system

responds to various environmental stimuli, such as stress and novelty (Berridge et al., 2010; Gonzalez

et al., 2016). Projection-defined dopaminergic populations in the VTA exhibited different responses to

orexin A (Baimel et al., 2017). Our PET imaging experiments, without orexin application, indicate that

Ox1R signaling in dopaminergic neurons modulates brain regions (MPA, Pir, endopiriform claustrum,

LSS, SubCV, spinal trigeminal nucleus, sp5, IRt and Gi) important for olfaction, eating, arousal and

sleeping (Supplementary table 1).When central orexin levels were increased by orexin injection, Ox1R

signaling in dopaminergic neurons regulates HDB, MCPO, Pir, DEn, S1, V2L, V1, BNST, LPGi and

M2, which are important for sensation, emotion, exploration, and action selection (Supplementary table

1). In future studies, it would be interesting to investigate the direct and indirect downstream pathways

of Ox1R-expressing dopaminergic neurons. Interestingly, in the fruitfly *Drosophila melanogaster*, it

has been shown that the dopaminergic system oppositely regulates sleep-wake arousal and

environmentally stimulated arousal, and the authors discussed the possibility to identify similar patterns

in mammals (Lebestky et al., 2009). It was further suggested that the dopaminergic system differentially

regulates arousal levels under different stimulating conditions (Lebestky et al., 2009; Lima &

Miesenbock, 2005).

We found a potential direct impact of dopaminergic neurons on LPGi and BNST. DRD1 and

DRD2 in the BNST regulate anxiety, fear, and addiction (De Bundel et al., 2016; Eiler et al., 2003;

Giardino et al., 2018; Krawczyk et al., 2013). LGPi regulates exploratory behaviors and locomotor speed

(Takatoh et al., 2013; Arber & Costa, 2022; Capelli et al., 2017). There are few publications on the

functional relevance of dopaminergic projections in the LPGi. Future studies would be very interesting

to functionally analyze DRD2 signaling in the LPGi and to understand how orexin- interacts with DRD2

signaling.

 Materials and Methods

Animal care

All animal procedures were conducted in compliance with protocols approved by the local government authorities (Bezirksregierung Cologne, Germany) and were in accordance with National Institutes of Health guidelines. Mice were housed in groups of 3 - 5 at 22°C - 24°C using a 12-hr light/12-hr dark cycle. Animals had *ad libitum* access to water and food at all times. Animals were fed a NCD (R/M-H, ssniff[R]), which contains 57% calories from carbohydrates, 34% calories from protein, and 9% calories from fat.

Generation of Ox1R$^{\Delta DAT}$ mice

Mice carrying a loxP-flanked Ox1R-allele (Ox1R fl/fl) were generated in our facility (Xiao et al., 2021). Ox1R fl/fl mice were crossed with Dat-Cre mice (Ekstrand et al., 2007) to obtain Ox1R fl/fl, Dat-Cre tg/wt mice, i.e. dopaminergic-specific Ox1R knockout mice on a C57/Bl6 N background (Ox1R$^{\Delta DAT}$). Mice from the same breeding without the Cre transgene were used as controls (Ox1R fl/fl, Dat-Cre wt/wt).

Glucose and insulin tolerance tests

Glucose tolerance tests were performed on 12-weeks-old animals that had been fasted for 6 hours from 9 am. Insulin tolerance tests were performed on 11-weeks-old random-fed mice around 9 am. Animals received an intraperitoneal injection of 20% glucose (10 ml/kg body weight; KabiPac) or insulin (0.75 U/kg body weight; Sanofi-Aventis) into the peritoneal cavity, respectively. Blood glucose values were determined in whole venous blood using an automatic glucose monitor (Contour[R], Bayer). Glucose levels were determined in blood collected from the tail tip immediately before and 15, 30, and 60 minutes after the injection, with an additional value determined after 120 minutes for the glucose tolerance tests.

Analysis of body composition

Nuclear magnetic resonance (NMR) was employed to determine whole body composition of live

animals on NCD at the age of 20 weeks using the NMR Analyzer minispec mq7.5 (Bruker Optik,

Ettlingen, Germany).

Histology

Brains from male mice were removed, post-fixed in 4% paraformaldehyde (PFA) for the indicated time,

and dehydrated in 20% sucrose in 0.1M phosphate-buffered saline (PBS) overnight. Brains were stored

at -80°C until cutting.

Immunostaining

Brains were post-fixed for 6 h at 4 °C and cut (30 μm) with Leica CM3050 S Research Cryostat. Sections

were incubated in 0.3% glycine for 10 min after washing in 0.1 M PBS for 2 × 10 min. After washing

in PBS for another 10 min, sections were incubated in 0.03% SDS (in PBS) for 10 min before they got

blocked with 3% donkey serum (in PBS, 0.25% Triton X-100) for 1 h at RT. Afterwards, they were

incubated with primary antibodies, including goat anti-c-Fos (sc-526, Santa Cruz Biotechnology),

chicken anti-GFP (ab13970, Abcam) and/or rabbit anti-Th (ab112, Abcam), for overnight (GFP) or 48

h (c-Fos and Th) at 4°C, washed in PBS for 3 × 10 min, and incubated with secondary antibodies,

including Alexa Fluro 594 donkey anti-goat, FITC donkey anti-chicken, Alexa Fluro 488 donkey anti-

rabbit (a11058, sa 1-7200 and a21206, Invitrogen), for 1 h at RT. After washing in PBS for 3 × 10 min,

slices were mounted and covered with VECTASHIELD Antifade Mounting Medium with DAPI (Vector

Laboratories). Slices were stained together at one time for each experiment to have identical conditions

for comparable signals. Slides were stored at 4 °C until imaging. Images were obtained with confocal

laser scanning microscope Leica TCS SP8 and manually analyzed with Image J/FIJI (version 1.50d).

RNAscope fluorescent in situ hybridization

Brains from 12-wk-old male mice were post-fixed for 20 - 22 h at RT and cut (20 μm). Slides were

stored at -80°C until staining. All reagents were purchased from Advanced Cell Diagnostics, and the

staining was performed with the RNAscope Multiplex Fluorescent v2 kit (ACD, Advanced Cell

Diagnostics) according to the user manual. Probes for Ox1R (*Hcrtr1*, 471561-C3, ACD) and Ox2R

(*Hcrtr2*, 471551, ACD), DRD1 (*Drd1a*, 406491-C2, ACD), DRD2 (*Drd2*, 406501), and Th (*Th*,

317621-C4) were purchased from ACD. In brief, slides were briefly washed in diethyl-pyrocarbonate (DEPC)-treated Millipore water, air dried, and then dried at 60°C overnight. On the second day, slides were treated with hydrogen peroxide (H_2O_2) for 10 min at RT, washed in water, and boiled in Target Retrieval solution (around 99.4 °C) for 8 -10 min. After a brief washing in water and dehydration in absolute ethanol, slides were incubated with protease IV for 30 min at RT. Slides were washed again in water and hybridized with the mixture of probes in different channels for 2 h in a humidified chamber at 40 °C. Afterwards, the hybridization was amplified with AMP 1 for 30 min, AMP 2 for 30 min and AMP 3 for 15 min. The signal was then developed for each channel. For example, for channel 1, slides were incubated with HRP-C1 for 15 min, the fluorophore for 30 min, and HRP blocker for 15 min. All amplification and development were performed at 40 °C, and 2×2 min of washing in ACD wash buffer was performed after each step. Finally, the slides were counterstained with DAPI for 1 min, mounted with Prolong^{TM} Gold Antifade reagent with DAPI (Invitrogen), and covered with coverslips.

Slides were dried and stored at 4 °C. Imaging was performed with a confocal laser scanning microscope Leica TCS SP8. The images were manually analyzed with Image J/FIJI (version 1.50d).

Ca^{2+} imaging

Calcium dynamics were measured in dopaminergic substantia nigra neurons using the genetically encoded calcium indicator GCaMP6. Experiments were performed on brain slices from 18-19 week-old male control and Ox1R$^{\Delta DAT}$ mice. A minimum of 4 weeks separated the GCaMP6 virus injection and the imaging experiments. Animals were kept under standard laboratory conditions, with tap water and chow available ad libitum, on a 12h light/dark cycle. The animals were lightly anesthetized with isoflurane (B506; AbbVie Deutschland GmbH and Co KG, Ludwigshafen, Germany) and decapitated. Coronal slices (280 μm) containing SN were cut with a vibration microtome (VT1200 S; Leica, Germany) under cold (4°C), carbogenated (95% O_2 and 5% CO_2), glycerol-based modified artificial cerebrospinal fluid (GaCSF). GaCSF contained (in mM): 244 Glycerol, 2.5 KCl, 2 MgCl$_2$, 2 CaCl$_2$, 1.2 NaH$_2$PO$_4$, 10 HEPES, 21 NaHCO$_3$, and 5 Glucose adjusted to pH 7.2 with NaOH. The brain slices were continuously superfused with carbogenated aCSF at a flow rate of ~2.5 ml·min^{-1}. aCSF contained (in mM): 125 NaCl, 2.5 KCl, 2 MgCl$_2$, 2 CaCl$_2$, 1.2 NaH$_2$PO$_4$, 21 NaHCO$_3$, 10 HEPES, and 5 Glucose

adjusted to pH 7.2 with NaOH. To reduce GABAergic and glutamatergic synaptic input the aCSF

contained 10^{-4} M PTX (picrotoxin, P1675; Sigma-Aldrich), 5 x 10^{-6} M CGP (CGP-54626 hydrochloride,

BN0597, Biotrend), 5 x 10^{-5} M DL-AP5 (DL-2-amino-5-phosphonopentanoic acid, BN0086, Biotrend),

and 10^{-5} M CNQX (6-cyano-7-nitroquinoxaline-2,3-dione, C127; Sigma-Aldrich).

The imaging setup consisted of a Zeiss AxioCam/MRm CCD camera with a 1388x1040 chip

and a Polychromator V (Till Photonics, Gräfelfing, Germany) that was coupled via an optical fiber into

the Zeiss AxioExaminer upright microscope (Objective W "Plan-Apochromat" 20x/1.0 DIC D=0.17

M27 75mm). The camera and polychromator were controlled by the software Zen pro, including the

module 'Physiology' (2012 blue edition, Zeiss). The SN neurons were identified according to their

anatomical location and expression of the GCaMP6. Calcium signals in GCaMP6 expressing cells were

monitored by images acquired at 470 nm excitation wavelengths with 20 ms exposure time at ~0.2 Hz.

The emitted fluorescence was detected through a 500-550 nm bandpass filter (BP525/50), and data were

acquired using 4x4 on-chip binning. Images were recorded in arbitrary units (AU) and analyzed as 16-

bit grayscale images.

Orexin A (ab120212, Abcam) in the concentration of 100 nM was applied for 8 min. To analyze

the orexin effect, we compared the fluorescence measured during 4 min intervals that were recorded

immediately before and at the end of the peptide application. This protocol was followed by applying

high K^+ concentration saline (40 mM KCl; osmolarity was adjusted by reducing the NaCl concentration

accordingly) to elicit the maximal calcium response.

The image analysis was performed offline using Image J (version 1.53a), Igor Pro 6

(WaveMetrics, USA), and Prism 8 (GraphPad, California, USA). After the experiments, regions of

interest (ROI) were defined based on the high K^+ saline responses. The mean AU values of the ROIs

were calculated in Image J. Time series analysis was performed with Igor Pro 6. To correct for bleaching

artifacts, the baseline fluorescence (without orexin application) was fit. The extended fit was subtracted

from the raw data. The orexin-induced Ca^{2+} signals were normalized to the high (40 mM) K^+ response.

On the single-cell level, a neuron was considered orexin-responsive if the change in fluorescence

induced by orexin was 3 times larger than the standard deviation (SD) (3 sigma criterion) of the baseline

fluorescence.

Food intake and spontaneous activity

Mice (20 weeks old) were analyzed for food intake, spontaneous activity, respiratory exchanging ratio and energy expenditure in a PhenoMaster System (TSE Systems) as previously described (Jordan et al., 2011).

Behavioral studies

Mice were moved into the behavioral test room, weighed, and handled 1 week before the measurement. On the day of the tests, mice (9-15 weeks old) were moved near to the equipment and handled 1 h before the tests. There was a 1-week break between different behavioral tests for anxiety, exploration or locomotion when using the same mice.

Open field test (OFT) ，dark/light (D/L) box and Hole-board tests

OFT and D/L box tests were performed with a Seamless Open Field Starter Package for Mouse plus the Light/Dark Insert or the hole-board floor (Med Associates, Inc.) between 10 am to 4 pm.

To start OFT, mice were put in the middle of a clear Plexiglas® chamber (27.3 cm × 27.3 cm × 20.3 cm) sitting in a sound-attenuating cubicle, and mouse movement was automatically traced for 30 min with 48 infrared beams divided into the X, Y and Z plane. For the D/L box test, a black Plexiglas® box (half the size of the chamber) was inserted at the left side of the chamber, with an opening at the middle side. Mice were put in the middle of the light side and faced the openings. Mouse movement was traced for 5 min. For the hole-board test, a metal hole-board floor with 16 equally spaced holes for exploration was inserted in the chamber. Mice were put in the middle of the chambers and the nose pokes were monitored automatically for 10 min per session. There was 1 h break between each session. 4 sessions per day were analyzed for 2 consecutive days.

Elevated 0-maze test

An elevated 0-maze test was performed with TSE elevated 0-Maze in combination with the video tracking system TSE VideoMot 3D version 7.01. There are 2 closed and 2 open runways without a center position. The outer diameter is 48 cm, the arm width is 5 cm, the wall height is 10 cm and the base height is 55 cm. Mice were put in the middle of an open arm and faced to the closed arm to start the experiment. Mouse movement was monitored for 5 min.

Novel object recognition test

Novel object recognition test was modified from the publication before (Ruud et al., 2019). The mouse was acclimated and measured for 10 min in the 1st session and for 5 min in the 2nd session on the same day, in a polycarbonate box with non-transparent walls (50 cm × 50 cm × 30 cm). Two objects were present in the box. In the 1st session, two identical objects (glass beaker filled with blue marbles) were present and fixed with odor-free adhesive to the ground flooring at two sides of the box delineated diagonally. In each diagonal corner, the object is 11.5 cm and 14 cm from each wall respectively, and the distance between the two objects is 17.5 cm. Mice always started at the same corner which was between the two fixed objects and were allowed to explore and learn the experimental area and objects. In the 2nd session which started after a 1-h break in the home cage, one of the two identical objects was replaced with a novel object (Lego $^{®}$ Primo brick), and mice were allowed again to explore the environment and objects. The positions of the mouse nose, tail and center were monitored with a video tracking system TSE VideoMot 2. The interaction with a fixed object was identified when the nose was facing the object within a distance of 2 cm.

Conditioned place preference test for cocaine

The test was modified from previous publication (Ruud et al., 2019), and was performed in a polycarbonate box with non-transparent walls. The box was separated into 3 compartments with 2 independent partitions. It contains a left compartment with zebra stripes (white and black) painted on the wall and a rough ground (16.5 x 13.2 x 21.2 cm), a small middle area (7.0 x 13.2 x 21.2 cm), and a right compartment with a white wall and smooth ground. Mouse movement was monitored with TSE VideoMot 2 for 30 min per session, 1 session per day. Mice started from the middle area and had access to both left and right compartments through an opening (4.0 cm x 4.2 cm) in the partition during the habituation and pre-test days (d1-3) and on the test day (d10), and were put into and kept in either the left or right compartment without an opening in the partition during the conditioning phase (d4-9). The preference of the two compartments by nature was analyzed during the pre-test days. At the conditioning phase, on alternative days, mice received i.p. injection of cocaine (5 mg/kg BW) or saline immediately before entering the respective compartments. Cocaine and saline were always paired to the non-preferred and preferred compartments, which were determined by pre-tests, respectively. There was no

injection on the habituation, pre-test or test days. The percentage of time spent in the reward-paired compartment was calculated: 100 x time spent in the compartment / (total time - time spent in the middle area). The CPP score was then analyzed using the calculated percentage of time: 100 x (time on the test day – time on pre-test days)/ time on pre-test days.

Sucrose/sucralose preference test

Two-bottle tests were performed with 2 15-ml falcon tubes connected to sippers. Mice were single caged 1 week before the beginning of the experiment. 2 falcon tubes filled with water (side-by-side) replaced the normal water bottle. Water intake was monitored for a week to make sure mice adjusted to drinking from the sippers. When the test started, 2 new sipper tubes of water were used for 2 days of baseline measurement and were then replaced with one containing water and one containing sucrose/sucralose solution for 2 days of preference measurement. Animals received a 0.5%, 1% and 2% (w/v) sucrose solution and 1 mM sucralose solution in sequential. Between the preference tests of different solutions, 2 days of baseline measurement were performed with 2 new sipper tubes of water as a washout. To avoid side bias, the positions of 2 sipper tubes were switched every day, with the sucrose/sucralose tube always starting on the non-preferred position, which was determined by water intake on the day before the preference measurement. Sipper tubes were weighed every day.

ICV cannula implantation and injection

12-wk-old mice received 1 mg/ml of tramadol in drinking water 2 d before the surgeries. On the day of surgery, mice were anesthetized with isoflurane, and a bolus of buprenorphine (0.1 mg/kg BW) was given (i. p.) to reduce pain. Brain regions were identified according to the atlas (Franklin & Paxinos, 2007). After alignment of the brain in the stereotaxic surgery platform, one small hole was drilled in the skull and the ICV cannula (26 gauge, Plastics One) was immediately implanted (AP: -0.2 mm, ML: +1 mm, DV: -2.4 mm) and fixed to the skull with dental acrylic (Super-Bond C&B). The dummy cannula caps were inserted to close and protect the cannula. Mice received a bolus of meloxicam (5 mg/kg BW, s.c.), and tramadol in drinking water for 3 d after surgeries to relieve pain. BW was checked twice a day. Experiments started at least 1 week later. For injection, the catheter was attached to an injector (Plastics One), which was inserted through the cannula. orexin A (Phoenix Pharmaceuticals) was

injected slowly at the dose of 1 nmol dissolved in 2 ul of saline. Starting from 5 days before the injection,

mice and the dummy cannula caps were handled for acclimation daily. The locomotion activity was

measured in a PhenoMaster System (TSE Systems), and mice were acclimated to the system at least 5

days before the test.

PET imaging

PET imaging was performed using an Inveon preclinical PET/CT system (Siemens). Male mice were

anesthetized with 2% isoflurane in 65%/35% nitrous oxide/oxygen gas, injected with saline or orexin A

and positioned on a dedicated mouse carrier (MEDRES, Germany) carrying two mice. Body

temperature was maintained at $37.0 \pm 0.5°C$ by a thermostatically controlled water heating system. For

injection of the radiotracer, a catheter consisting of a 30G cannula connected to a polythene tubing (ID

= 0.28 mm) was inserted into the tail vein and fixated by a drop of glue. After starting the PET scan, 7-

8 MBq of [18F]FDG in 50-100 µl saline were injected per mouse. Emission data were acquired for 45

minutes. Thereafter, animals were automatically moved into the CT gantry, and a CT scan was

performed (180 projections/360°, 200 ms, 80 kV, 500 µA). The CT data were used for attenuation

correction of the PET data, and the CT image of the scull was used for image co-registration. Plasma

glucose levels were determined from a tail vein blood sample using a standard glucometer (Bayer) after

removing the tail vein catheters. PET data were histogrammed in time frames of 12x30s, 3x60s, 3x120s,

7x240s, Fourier rebinned, and images were reconstructed using the MAP-SP algorithm provided by the

manufacturer. For co-registration, the imaging analysis software Vinci was used (Cizek et al., 2004).

Images were co-registered to a 3D mouse brain atlas constructed from the 2D mouse brain atlas

published by Paxinos (Paxinos et al., 2013).

Kinetic modeling

An image-derived input function was extracted from the PET data of the aorta, which could be identified

in the image of the first time frame of each animal. Input function data were corrected for partial volume

effect by assuming a standardized volume fraction of 0.6 (Green et al., 1998). Parametric images of the

[18F]FDG kinetic constants K_1, k_2, k_3, and k_4 were determined by a voxel-by-voxel (voxel size = 0.4

mm x 0.4 mm x 0.8 mm) fitting of data to a two- tissue-compartment kinetic model. K_1 is the constant

for transport from blood to tissue, k_2 for transport from tissue to blood, k_3 the constant for

phosphorylation of [18F]FDG to [18F]FDG-6-phosphate, and k_4 the constant for dephosphorylation.

The ratio of tissue and plasma glucose concentrations (C_E/C_P) is a measure for glucose transport and is

given by $C_E/C_P = K_1/(k_2 + k_3/0.26)$ (Backes et al., 2011; Jais et al., 2016). Since neuronal activation is

accompanied by increased glucose transport and this parameter is less sensitive to changes in plasma

glucose level, we use alterations of glucose transport (C_E/C_P) as surrogate for alterations in neuronal

activation.

Statistics

Statistical testing was performed by application of a voxel-wise t-test between groups. 3D maps of p-

values allow for the identification of regions with significant differences in the parameters. From these

regions, we defined volumes of interest (VOIs) and performed additional statistical testing for these

VOIs. For presentation only, 3D maps of p-values were re-calculated on a 0.1 mm x 0.1 mm x 0.1 mm

grid from the original dataset using trilinear interpolation.

<u>Virus injection</u>

Male mice received 1 mg/ml of tramadol in drinking water 2 d before the surgeries. On the day of

surgery, mice were anesthetized with isoflurane and a bolus of buprenorphine (0.1 mg/kg BW) was

given (i. p.) to reduce pain. Brain regions were identified according to the atlas (Franklin & Paxinos,

2007). After alignment of the brain in the stereotaxic surgery platform, 1 or 4 small holes were drilled

in the skull at specific coordinates. The virus, pAAV-Ef1a-DIO EYFP (~1×10^{13} GC/ml, 300 nl,

Addgene) was bilaterally injected into VTA and SN of 10 wk old mice, with micropipettes pulled in

house with a heating system. The coordinates from Bregma were: anterior-posterior, AP: -3.4 mm;

medial-lateral, ML: ±0.5/1.25 mm; dorsal-ventral, DV: -4.2 mm. The GCaMP6 virus

(AAV1.Syn.Flex.GCaMP6s.WPRE.SV40, ~4×10^{12} GC/ml, 300 nl, Penn Vector Core) was injected to

left SN (AP: -3.4 mm, ML: 1.1 mm, DV: -4.2 mm) of 12 wk old mice, to specifically express GCaMP6

in dopaminergic neurons. After virus injection, mice received a bolus of meloxicam (5 mg/kg BW, s.c.),

and tramadol in drinking water for 3 d after surgeries to relieve pain. BW was checked twice a day.

Experiments started at least 4 weeks later than virus injection to allow virus expression.

<u>Statistical methods</u>

All values are expressed as the mean $\pm$ SEM. Statistical analyses were conducted using

GraphPad PRISM (version 8) unless stated otherwise. Datasets with only two independent groups were

analyzed for statistical significance using an unpaired two-tailed Student's t-test. Datasets subjected to

two independent factors were analyzed using two-way ANOVA followed by Sidak's post hoc test. All

p-values < 0.05 were considered significant (*$p < 0.05$, **$p < 0.01$, and ***$p < 0.001$, ****$p < 0.0001$).

Acknowledgements

We acknowledge Hella Brönneke for outstanding support, Jens Alber, Pia Scholl, Christiane Schäfer, Nadine Spenrath, Ina Stünkel, Kerstin Marohl, Antonia Lorenz and Helmut Wratil for outstanding technical assistance. We received funding by the DFG within the framework of the TRR 134 (A.C.H., P.K.), CRC 1451 (P.K.) and within the Excellence Initiative by German Federal and State Governments (CECAD). This work was funded (in part) by the Helmholtz Alliance ICEMED (Imaging and Curing Environmental Metabolic Diseases) through the Initiative and Networking Fund of the Helmholtz Association. X.X. received funding from CECAD Family Support. G.Y. gratefully acknowledges financial doctoral support from the DFG-233886668/GRK1960.

Author Contributions

X.X. and A.C.H. conceived the project, designed the experiments, and wrote the manuscript with input from the other authors. A.C.H. generated the Ox1R flox mouse line. X.X. performed all the experiments apart from calcium imaging experiments (G.Y. and P.K.) and PET imaging experiments (A.L.C. and H.B.). All authors discussed the data, commented on the manuscript before submission, and agreed with the final submitted manuscript.

Competing Interests

The authors declare no competing interests.

References

Arber, S., & Costa, R. M. (2022). Networking brainstem and basal ganglia circuits for movement. *Nat Rev Neurosci*, *23*(6), 342-360. https://doi.org/10.1038/s41583-022-00581-w

Backes, H., Walberer, M., Endepols, H., Neumaier, B., Graf, R., Wienhard, K., & Mies, G. (2011). Whiskers area as extracerebral reference tissue for quantification of rat brain metabolism using (18)F-FDG PET: application to focal cerebral ischemia. *Journal of Nuclear Medicine*, *52*(8), 1252-1260. https://doi.org/10.2967/jnumed.110.085266

Baimel, C., Lau, B. K., Qiao, M., & Borgland, S. L. (2017). Projection-target-defined effects of orexin and dynorphin on VTA dopamine neurons. *Cell Rep*, *18*(6), 1346-1355. https://doi.org/10.1016/j.celrep.2017.01.030

Baykal, S., Albayrak, Y., Durankus, F., Guzel, S., Abbak, O., Potas, N., Beyazyuz, M., Karabekiroglu, K., & Donma, M. M. (2019). Decreased serum orexin A levels in drug-naive children with attention deficit and hyperactivity disorder. *Neurol Sci*, *40*(3), 593-602. https://doi.org/10.1007/s10072-018-3692-8

Berke, J. D. (2018). What does dopamine mean? *Nat Neurosci*, *21*(6), 787-793. https://doi.org/10.1038/s41593-018-0152-y

Berridge, C. W., Espana, R. A., & Vittoz, N. M. (2010). Hypocretin/orexin in arousal and stress. *Brain Res*, *1314*, 91-102. https://doi.org/10.1016/j.brainres.2009.09.019

Capelli, P., Pivetta, C., Soledad Esposito, M., & Arber, S. (2017). Locomotor speed control circuits in the caudal brainstem. *Nature*, *551*(7680), 373-377. https://doi.org/10.1038/nature24064

Cizek, J., Herholz, K., Vollmar, S., Schrader, R., Klein, J., & Heiss, W. D. (2004). Fast and robust registration of PET and MR images of human brain. *Neuroimage*, *22*(1), 434-442. https://doi.org/10.1016/j.neuroimage.2004.01.016

De Bundel, D., Zussy, C., Espallergues, J., Gerfen, C. R., Girault, J. A., & Valjent, E. (2016). Dopamine D2 receptors gate generalization of conditioned threat responses through mTORC1 signaling in the extended amygdala. *Molecular Psychiatry*, *21*(11), 1545-1553. https://doi.org/10.1038/mp.2015.210

de Lecea, L., Kilduff, T. S., Peyron, C., Gao, X., Foye, P. E., Danielson, P. E., Fukuhara, C., Battenberg, E. L., Gautvik, V. T., Bartlett, F. S., 2nd, Frankel, W. N., van den Pol, A. N., Bloom, F. E., Gautvik, K. M., & Sutcliffe, J. G. (1998). The hypocretins: hypothalamus-specific peptides with neuroexcitatory activity. *Proc Natl Acad Sci U S A*, *95*(1), 322-327. https://doi.org/10.1073/pnas.95.1.322

Durairaja, A., & Fendt, M. (2020). Orexin deficiency modulates cognitive flexibility in a sex-dependent manner. *Genes Brain Behav*, e12707. https://doi.org/10.1111/gbb.12707

Eiler, W. J., 2nd, Seyoum, R., Foster, K. L., Mailey, C., & June, H. L. (2003). D1 dopamine receptor regulates alcohol-motivated behaviors in the bed nucleus of the stria terminalis in alcohol-preferring (P) rats. *Synapse*, *48*(1), 45-56. https://doi.org/10.1002/syn.10181

Ekstrand, M. I., Terzioglu, M., Galter, D., Zhu, S., Hofstetter, C., Lindqvist, E., Thams, S., Bergstrand, A., Hansson, F. S., Trifunovic, A., Hoffer, B., Cullheim, S., Mohammed, A. H., Olson, L., & Larsson, N. G. (2007). Progressive parkinsonism in mice with respiratory-chain-deficient dopamine neurons. *Proc Natl Acad Sci U S A*, *104*(4), 1325-1330. https://doi.org/10.1073/pnas.0605208103

Fadel, J., & Deutch, A. Y. (2002). Anatomical substrates of orexin–dopamine interactions: lateral hypothalamic projections to the ventral tegmental area. *Neuroscience*, *111*(2), 379-387. https://doi.org/https://doi.org/10.1016/S0306-4522(02)00017-9

Feldman, H. M., & Reiff, M. I. (2014). Clinical practice. Attention deficit-hyperactivity disorder in children and adolescents. *N Engl J Med*, *370*(9), 838-846. https://doi.org/10.1056/NEJMcp1307215

Franklin, K. B. J., & Paxinos, G. (2007). *The mouse brain in stereotaxic coordinates* (3rd ed.). Elsevier/Academic Press.

Freeman, L. R., Bentzley, B. S., James, M. H., & Aston-Jones, G. (2021). Sex differences in demand for highly palatable foods: role of the orexin system. *Int J Neuropsychopharmacol*, *24*(1), 54-63. https://doi.org/10.1093/ijnp/pyaa040

Furlong, T. M., Vianna, D. M., Liu, L., & Carrive, P. (2009). Hypocretin/orexin contributes to the expression of some but not all forms of stress and arousal. *Eur J Neurosci*, *30*(8), 1603-1614. https://doi.org/10.1111/j.1460-9568.2009.06952.x

Giardino, W. J., Eban-Rothschild, A., Christoffel, D. J., Li, S. B., Malenka, R. C., & de Lecea, L. (2018). Parallel circuits from the bed nuclei of stria terminalis to the lateral hypothalamus drive opposing emotional states. *Nat Neurosci*, *21*(8), 1084-1095. https://doi.org/10.1038/s41593-018-0198-x

Gonzalez, J. A., Iordanidou, P., Strom, M., Adamantidis, A., & Burdakov, D. (2016). Awake dynamics and brain-wide direct inputs of hypothalamic MCH and orexin networks. *Nat Commun*, *7*, 11395. https://doi.org/10.1038/ncomms11395

Green, L. A., Gambhir, S. S., Srinivasan, A., Banerjee, P. K., Hoh, C. K., Cherry, S. R., Sharfstein, S., Barrio, J. R., Herschman, H. R., & Phelps, M. E. (1998). Noninvasive methods for quantitating blood time-activity curves from mouse PET images obtained with fluorine-18-fluorodeoxyglucose. *Journal of Nuclear Medicine*, *39*(4), 729-734. <Go to ISI>://WOS:000072956600042

Howe, M. W., & Dombeck, D. A. (2016). Rapid signalling in distinct dopaminergic axons during locomotion and reward. *Nature*, *535*(7613), 505-510. https://doi.org/10.1038/nature18942

Hrabovszky, E., Molnar, C. S., Borsay, B. A., Gergely, P., Herczeg, L., & Liposits, Z. (2013). Orexinergic input to dopaminergic neurons of the human ventral tegmental area. *PLoS One*, *8*(12), e83029. https://doi.org/10.1371/journal.pone.0083029

Inutsuka, A., & Yamanaka, A. (2013). The physiological role of orexin/hypocretin neurons in the regulation of sleep/wakefulness and neuroendocrine functions. *Front Endocrinol (Lausanne)*, *4*, 18. https://doi.org/10.3389/fendo.2013.00018

Jais, A., Solas, M., Backes, H., Chaurasia, B., Kleinridders, A., Theurich, S., Mauer, J., Steculorum, S. M., Hampel, B., Goldau, J., Alber, J., Förster, C. Y., Eming, S. A., Schwaninger, M., Ferrara, N., Karsenty, G., & Brüning, J. C. (2016). Myeloid-cell-derived VEGF maintains brain glucose uptake and limits cognitive impairment in obesity. *Cell*, *165*(4), 882-895. https://doi.org/10.1016/j.cell.2016.03.033

Jordan, S. D., Kruger, M., Willmes, D. M., Redemann, N., Wunderlich, F. T., Bronneke, H. S., Merkwirth, C., Kashkar, H., Olkkonen, V. M., Bottger, T., Braun, T., Seibler, J., & Bruning, J. C. (2011). Obesity-induced overexpression of miRNA-143 inhibits insulin-stimulated AKT activation and impairs glucose metabolism [Research Support, Non-U.S. Gov't]. *Nat Cell Biol*, *13*(4), 434-446. https://doi.org/10.1038/ncb2211

Kitahama, K., Nagatsu, I., Geffard, M., & Maeda, T. (2000). Distribution of dopamine-immunoreactive fibers in the rat brainstem. *Journal of Chemical Neuroanatomy*, *18*(1-2), 1-9. https://doi.org/Doi 10.1016/S0891-0618(99)00047-2

Koblinger, K., Jean-Xavier, C., Sharma, S., Fuzesi, T., Young, L., Eaton, S. E. A., Kwok, C. H. T., Bains, J. S., & Whelan, P. J. (2018). Optogenetic activation of A11 region increases motor activity. *Front Neural Circuits*, *12*, 86. https://doi.org/10.3389/fncir.2018.00086

Korotkova, T. M., Eriksson, K. S., Haas, H. L., & Brown, R. E. (2002). Selective excitation of GABAergic neurons in the substantia nigra of the rat by orexin/hypocretin in vitro. *Regul Pept*, *104*(1-3), 83-89. https://doi.org/10.1016/s0167-0115(01)00323-8

Korotkova, T. M., Sergeeva, O. A., Eriksson, K. S., Haas, H. L., & Brown, R. E. (2003). Excitation of ventral tegmental area dopaminergic and nondopaminergic neurons by orexins/hypocretins. *J Neurosci*, *23*(1), 7-11. https://www.ncbi.nlm.nih.gov/pubmed/12514194

Krawczyk, M., Mason, X., DeBacker, J., Sharma, R., Normandeau, C. P., Hawken, E. R., Di Prospero, C., Chiang, C., Martinez, A., Jones, A. A., Doudnikoff, E., Caille, S., Bezard, E., Georges, F., & Dumont, E. C. (2013). D1 dopamine receptor-mediated LTP at GABA synapses encodes motivation to self-administer cocaine in rats. *Journal of Neuroscience*, *33*(29), 11960-+. https://doi.org/10.1523/Jneurosci.1784-13.2013

Lebestky, T., Chang, J. S., Dankert, H., Zelnik, L., Kim, Y. C., Han, K. A., Wolf, F. W., Perona, P., & Anderson, D. J. (2009). Two different forms of arousal in Drosophila are oppositely regulated by the dopamine D1 receptor ortholog DopR via distinct neural circuits. *Neuron*, *64*(4), 522-536. https://doi.org/10.1016/j.neuron.2009.09.031

Lima, S. Q., & Miesenbock, G. (2005). Remote control of behavior through genetically targeted photostimulation of neurons. *Cell*, *121*(1), 141-152. https://doi.org/10.1016/j.cell.2005.02.004

Lippert, R. N., Hess, S., Klemm, P., Burgeno, L. M., Jahans-Price, T., Walton, M. E., Kloppenburg, P., & Bruning, J. C. (2020). Maternal high-fat diet during lactation reprograms the dopaminergic circuitry in mice. *J Clin Invest*, *130*(7), 3761-3776. https://doi.org/10.1172/JCI134412

Liu, C., Xue, Y., Liu, M. F., Wang, Y., Liu, Z. R., Diao, H. L., & Chen, L. (2018). Orexins increase the firing activity of nigral dopaminergic neurons and participate in motor control in rats. *J Neurochem*, *147*(3), 380-394. https://doi.org/10.1111/jnc.14568

Mahler, S. V., Moorman, D. E., Smith, R. J., James, M. H., & Aston-Jones, G. (2014). Motivational activation: a unifying hypothesis of orexin/hypocretin function. *Nat Neurosci*, *17*(10), 1298-1303. https://doi.org/10.1038/nn.3810

Mahler, S. V., Smith, R. J., Moorman, D. E., Sartor, G. C., & Aston-Jones, G. (2012). Multiple roles for orexin/hypocretin in addiction. *Prog Brain Res*, *198*, 79-121. https://doi.org/10.1016/B978-0-444-59489-1.00007-0

Marcus, J. N., Aschkenasi, C. J., Lee, C. E., Chemelli, R. M., Saper, C. B., Yanagisawa, M., & Elmquist, J. K. (2001). Differential expression of orexin receptors 1 and 2 in the rat brain. *J Comp Neurol*, *435*(1), 6-25. https://doi.org/10.1002/cne.1190

Nair-Roberts, R. G., Chatelain-Badie, S. D., Benson, E., White-Cooper, H., Bolam, J. P., & Ungless, M. A. (2008). Stereological estimates of dopaminergic, GABAergic and glutamatergic neurons in the ventral tegmental area, substantia nigra and retrorubral field in the rat. *Neuroscience*, *152*(4), 1024-1031. https://doi.org/10.1016/j.neuroscience.2008.01.046

Nakamura, T., Uramura, K., Nambu, T., Yada, T., Goto, K., Yanagisawa, M., & Sakurai, T. (2000). Orexin-induced hyperlocomotion and stereotypy are mediated by the dopaminergic system. *Brain Res*, *873*(1), 181-187. https://doi.org/10.1016/s0006-8993(00)02555-5

Narayanan, N. S., Guarnieri, D. J., & DiLeone, R. J. (2010). Metabolic hormones, dopamine circuits, and feeding. *Front Neuroendocrinol*, *31*(1), 104-112. https://doi.org/10.1016/j.yfrne.2009.10.004

Palmiter, R. D. (2007). Is dopamine a physiologically relevant mediator of feeding behavior? *Trends in Neurosciences*, *30*(8), 375-381. https://doi.org/10.1016/j.tins.2007.06.004

Paxinos, G., Franklin, K. B. J., & Franklin, K. B. J. (2013). *Paxinos and Franklin's the mouse brain in stereotaxic coordinates* (4th ed.). Elsevier/Academic Press.

Peyron, C., Tighe, D. K., van den Pol, A. N., de Lecea, L., Heller, H. C., Sutcliffe, J. G., & Kilduff, T. S. (1998). Neurons containing hypocretin (orexin) project to multiple neuronal systems. *J Neurosci*, *18*(23), 9996-10015. https://doi.org/10.1523/JNEUROSCI.18-23-09996.1998

Roeper, J. (2013). Dissecting the diversity of midbrain dopamine neurons. *Trends in Neurosciences*, *36*(6), 336-342. https://doi.org/https://doi.org/10.1016/j.tins.2013.03.003

Ruud, J., Alber, J., Tokarska, A., Engstrom Ruud, L., Nolte, H., Biglari, N., Lippert, R., Lautenschlager, A., Cieslak, P. E., Szumiec, L., Hess, M. E., Bronneke, H. S., Kruger, M., Nissbrandt, H., Korotkova, T., Silberberg, G., Rodriguez Parkitna, J., & Bruning, J. C. (2019). The fat mass and obesity-associated protein (FTO) regulates locomotor responses to novelty via D2R medium spiny neurons. *Cell Rep*, *27*(11), 3182-3198 e3189. https://doi.org/10.1016/j.celrep.2019.05.037

Sakurai, T., Amemiya, A., Ishii, M., Matsuzaki, I., Chemelli, R. M., Tanaka, H., Williams, S. C., Richardson, J. A., Kozlowski, G. P., Wilson, S., Arch, J. R. S., Buckingham, R. E., Haynes, A. C., Carr, S. A., Annan, R. S., McNulty, D. E., Liu, W.-S., Terrett, J. A., Elshourbagy, N. A., . . . Yanagisawa, M. (1998). Orexins and orexin receptors: a family of hypothalamic neuropeptides and G protein-coupled receptors that regulate feeding behavior. *Cell*, *92*(4), 573-585. https://doi.org/https://doi.org/10.1016/S0092-8674(00)80949-6

Sharples, S. A., Koblinger, K., Humphreys, J. M., & Whelan, P. J. (2014). Dopamine: a parallel pathway for the modulation of spinal locomotor networks. *Front Neural Circuits*, *8*, 55. https://doi.org/10.3389/fncir.2014.00055

Soya, S., & Sakurai, T. (2020). Evolution of orexin neuropeptide system: structure and function. *Front Neurosci*, *14*, 691. https://doi.org/10.3389/fnins.2020.00691

Takatoh, J., Nelson, A., Zhou, X., Bolton, M. M., Ehlers, M. D., Arenkiel, B. R., Mooney, R., & Wang, F. (2013). New modules are added to vibrissal premotor circuitry with the emergence of exploratory whisking. *Neuron*, *77*(2), 346-360. https://doi.org/10.1016/j.neuron.2012.11.010

Tsujino, N., & Sakurai, T. (2013). Role of orexin in modulating arousal, feeding, and motivation. *Front Behav Neurosci*, *7*, 28. https://doi.org/10.3389/fnbeh.2013.00028

Tung, L. W., Lu, G. L., Lee, Y. H., Yu, L., Lee, H. J., Leishman, E., Bradshaw, H., Hwang, L. L., Hung, M. S., Mackie, K., Zimmer, A., & Chiou, L. C. (2016). Orexins contribute to restraint stress-induced cocaine relapse by endocannabinoid-mediated disinhibition of dopaminergic neurons. *Nat Commun*, *7*, 12199. https://doi.org/10.1038/ncomms12199

Turiault, M., Parnaudeau, S., Milet, A., Parlato, R., Rouzeau, J. D., Lazar, M., & Tronche, F. (2007). Analysis of dopamine transporter gene expression pattern -- generation of DAT-iCre transgenic mice. *FEBS J*, *274*(14), 3568-3577. https://doi.org/10.1111/j.1742-4658.2007.05886.x

Volkow, N. D., & Swanson, J. M. (2013). Clinical practice: Adult attention deficit-hyperactivity disorder. *N Engl J Med*, *369*(20), 1935-1944. https://doi.org/10.1056/NEJMcp1212625

Xiao, X., Yeghiazaryan, G., Hess, S., Klemm, P., Sieben, A., Kleinridders, A., Morgan, D. A., Wunderlich, F. T., Rahmouni, K., Kong, D., Scammell, T. E., Lowell, B. B., Kloppenburg, P., Brüning,

J. C., & Hausen, A. C. (2021). Orexin receptors 1 and 2 in serotonergic neurons differentially regulate peripheral glucose metabolism in obesity. *Nature Communications*, *12*(1), 5249. https://doi.org/10.1038/s41467-021-25380-2