A parametrized two-domain thermodynamic model explains diverse mutational effects on protein allostery

Zhuang Liu[a], Thomas Gillis[b], Srivatsan Raman[b,c,d], and Qiang Cui[a,e,1]

[a]Department of Physics, Boston University, Boston, MA-02215, United States; [b]Department of Biochemistry, University of Wisconsin, Madison, WI-53706, United States; [c]Department of Chemistry, University of Wisconsin, Madison, WI-53706, United States; [d]Department of Bacteriology, University of Wisconsin, Madison, WI-53706, United States; [e]Department of Chemistry, Boston University, Boston, MA-02215, United States

Our understanding of protein allostery continues to be challenged by new experimental observations. Specifically, recent deep mutational scanning study showed that allosteric hotspots in the tetracycline repressor (TetR) and its homologous transcriptional factors (TFs) are broadly distributed rather than spanning well-defined structural pathways as often assumed. Moreover, the loss of allostery due to mutations of the hotspots was rescued by distributed additional mutations in a degenerate fashion. To gain a physical understanding of these intriguing observations, we develop a two-domain thermodynamic model for TetR, in which the protein is generically divided into ligand- and DNA-binding domains. The model incorporates three biophysical parameters to capture the intra- and inter-domain properties of TetR, and readily rationalizes the distributed network of allosterically coupled residues. Furthermore, we establish a quantitative connection between the model and the in vivo expression level of the TetR-regulated gene, which allows the extraction of the biophysical parameters in diverse mutants based on experimental induction curves using statistical inference. The quantitative mapping of the mutants to the model parameter space reveals the intrinsic connection between intra- and inter-domain allosteric parameters and also enables the evaluation of epistasis between the characterized mutations. The insights gained from this study into the nature of two-domain allostery are expected to have broader implications for other multidomain allosteric proteins.

allostery | thermodynamic model | tetracycline repressor | epistasis

Allostery, a fundamental regulatory mechanism of biomolecular functions, is prevalent in life processes (1–8). The long-range signaling of allostery makes it a fascinating phenomenon, in which binding of an effector molecule (ligand) at the allosteric site alters the function of a distal active site (8–11). Current descriptions of allostery can be largely cast into two categories: one adopts a mechanical view, focusing on the propagation of conformational distortions from the allosteric site to the active site (12–17); the other emphasizes the thermodynamic aspect of the problem, highlighting the effect of ligand binding on shifting the protein population among preexisting conformational states characterized by different ligand binding affinities and active site properties (e.g., the classic MWC model) (1–4, 18–20). Models based on both perspectives have provided insights into the function of prototypical allosteric systems thanks to decades of combined efforts of experiment, computation and theory (4, 5, 8, 20–27). Of note, these two perspectives of allostery are complementary rather than contradictory to each other (28). The conformational coupling between spatially distant functional sites (allosteric site and active site) plays a vital role in regulating allosteric

function, allowing for the transmission of signals from one site to the other (29). While the mechanical view primarily seeks to identify the structural basis for signal transduction, it is implicitly assumed within the population shift perspective, which offers a comprehensive and quantitative description of allostery (30–33).

In recent years, a thermodynamic model referred to as the ensemble allosteric model (EAM) has been applied to conceptualize protein allostery in terms of intra- and inter-domain properties, with the latter explicitly quantifying the energetic coupling between distant functional sites (5, 11, 34). This framework is consistent with the observation that allosteric proteins often partition different activities into distinct domains, such as the ligand- and DNA-binding domains in transcription factors and the effector-binding and catalytic domains in enzymes (35–38). Such thermodynamic approach finds broad applicability across proteins in general, as it has been proposed that all proteins are potentially allosteric (29, 39). Consequently, this raises intriguing questions about the nature of allostery. For instance, do intrinsic connections exist between the intra- and inter-domain properties of a protein, given the highly cooperative nature of allosteric networks? What roles do sequence and structure play in synergistically determining these properties? Furthermore, what are the parameters within the model that are most essential to the accurate description of realistic allosteric systems, especially prediction of activity upon multiple mutations? To answer these questions and deepen our understanding of allosteric regulation, it is essential to parameterize and test the thermodynamic model using comprehensive mutational data, a topic that still requires further exploration (9, 40).

A critical test of any thermodynamic model is whether it can explain the effect of mutations on allosteric signaling. Deep mutational scanning (DMS) analysis has emerged as a powerful function-centric approach over the past decade, which measures the impact of all possible single mutations (41–46). The methodology provides an unbiased way of identifying critical residues for protein allostery and generates extensive data for validating existing computational and theoretical models. Along this line, recent DMS study of four homologous bacterial transcription factors (TFs) in the TetR family (TetR, TtgR, MphR and RolR) revealed that the residues

Z.L., S.R. and Q.C. conceptualized the work. Z.L. and T.G. conducted the research. Z.L. and T.G. analyzed the data. All authors are involved in writing the manuscript.

The authors have no competing interest.

[1]To whom correspondence should be addressed. E-mail: qiangcui@bu.edu

critical for allosteric signaling (hotspots) in these TFs are broadly distributed with no apparent structural link to either the allosteric or the active site (40, 47–50). This contrasts the commonly held view that hotspot residues tend to form well-defined pathways linking the two sites (7, 51–54); the observations also hinted at common molecular rules of allostery in the TetR family of TFs (55, 56). Moreover, systematic analysis of higher order TetR mutants in the background of five noninducible ("dead") mutants revealed remarkable functional plasticity in allosteric regulation (9). Specifically, the loss of inducibility due to the mutations of allostery hotspots could be rescued (restored wildtype-like inducibility) by additional mutations, and different dead mutations exhibit varying numbers of rescuing mutations, which are usually distal and lacking any obvious structural rationale.

While the identification of broadly distributed dead and rescuing mutations in these studies is exciting, such qualitative activity characterization of mutants (inducible and noninducible at a given ligand concentration) prevents a deeper mechanistic understanding of the observation. In fact, mutations exert a graded impact on the allosteric signaling of the TFs; i.e., the expression level of the regulated gene varies among both inducible and noninducible mutants in a continuous, ligand-concentration dependent manner. The nuanced mutational effect on allosteric regulation is linked apparently to the equilibria among different conformational and binding states of the mutant TF, which are determined by the allosteric parameters (intra- and inter-domain properties). Therefore, mapping the mutants onto the parameter space of a biophysical model by exploiting additional experimental data (see below) is crucial for elucidating the observed allosteric phenomena in a comprehensive and physically transparent manner.

In this study, we develop a two-domain statistical thermodynamic model for TetR, in which the protein is generically divided into ligand- and DNA-binding domains (LBD and DBD). Our model incorporates three essential biophysical parameters that capture the intra- and inter-domain properties of the protein, as elaborated in the subsequent section. The model readily rationalizes the myriad ways that mutations can perturb allostery as observed in the aforementioned DMS measurements, by revealing that mutations perturbing intra- and inter-domain properties can lead to similar TF inducibilities at a single ligand concentration. Moreover, the model elucidates the distinct influences of these parameters on the complete induction curve, serving as a diagnostic tool for dissecting the intricate allosteric effects of mutations. We validate the model by accurately describing the induction curves of a comprehensive set of TetR mutants, thereby enabling the quantification of mutational effects and epistatic interactions (57, 58). The insights from this combined theoretical and experimental investigation of TetR allostery are expected to generally apply to other two-domain allosteric systems, such as transcription factors like catabolite activator protein (CAP) (36), receptors like pentameric ligand-gated ion channels (pLGICs) (59, 60), and allosteric enzymes like aspartate transcarbamoylase (ATCase) (37, 38).

Results

Overview of the Two-Domain Thermodynamic Model of Allostery. As shown in Fig. 1*A* and *B* TetR can be generically divided into a LBD and a DBD, disregarding its homodimeric nature (9, 19, 24, 40, 61, 62). In the same vein of the classic allostery models (1, 2), each domain features two (the relaxed/inactive and tense/active) conformations that differ in free energies and binding affinities. For the simplicity and mechanistic clarity of the model, we assume negligible binding affinity of the LBD to the ligand and the DBD to DNA in their inactive conformations and consider competent binding only for the active ones. Each domain must overcome a free energy increase to transition from the inactive to the active conformation (ε_L for LBD and ε_D for DBD). Importantly, the two domains are allosterically coupled, in that there is a free energy penalty γ when both domains adopt the active conformations simultaneously. For example, when the ligand binds to the LBD, selecting its active conformation, it discourages the active conformation of the DBD and therefore DNA binding. This anti-cooperativity establishes the foundation for allosteric regulation in TetR.

Accordingly, there are four possible conformational states of TetR, namely $L_I D_I$, $L_A D_I$, $L_I D_A$ and $L_A D_A$, with L/D and I/A denoting LBD/DBD and inactive/active conformation, respectively (see Fig. 1*B* and the upper four states of Fig. 1*D*). Binding of ligand/operator to the active LBD/DBD further lowers the free energy of the corresponding state in a concentration dependent manner following the standard formulation of binding equilibrium (see the lower three states of Fig. 1*D*). The regulatory mechanism of TetR allostery can then be qualitatively explained by the schematics in Fig. 1*C* and *D*. Without the ligand, wildtype (WT) repressor predominantly binds to the operator sequence, which obstructs the binding of RNA polymerase (RNAP) to the adjacent promoter of the regulated gene (*SI Appendix*, Fig. S1A). In the presence of ligand (inducer) at a sufficiently high concentration, the ligand-bound $L_A D_I$ state (L-$L_A D_I$) has the lowest free energy compared with other possible (DNA-bound) states, thus the repressor predominantly releases the operator upon ligand binding, enabling the expression of downstream genes (Fig. 1*C* and *SI Appendix*, Fig. S1A).

Mutations perturb the three biophysical parameters (ε_D, ε_L and γ) and hence the free energy landscape (Fig. 1*D*), leading to changes in TetR function. The three limiting scenarios are depicted in Fig. 1*D*: 1. when a mutation significantly increases ε_L, it raises the free energy of the L-$L_A D_I$ state relative to the DNA-bound state $L_I D_A$-D, which results in a dead (noninducible) phenotype; 2. when a mutation substantially decreases ε_D, it raises the free energy of the L-$L_A D_I$ state relative to both DNA-bound states ($L_I D_A$-D and L-$L_A D_A$-D), also leading to a dead mutant; 3. mutations that decrease γ greatly lower the free energy of the double-bound state (L-$L_A D_A$-D) relative to L-$L_A D_I$, again leading to the suppression of induction. Therefore, these schemes highlight that hotspot mutations can disrupt allostery by perturbing either intra- (ε_D, ε_L) or inter-domain (γ) properties. Functional readouts like DMS can help distinguish these mechanistic differences at the biophysical level. Notably, mutations mainly perturbing ε_D or ε_L do not have to lie on the structural pathways linking the allosteric and active sites, which explains the broad hotspot distributions observed in the DMS measurement (9, 24, 40, 62). In general, a dead mutation is likely of mixed nature, as long as its effect on the free energy landscape promotes the dominance of the DNA-bound states. Likewise, rescuing mutations may restore WT-like inducibility by modifying intra- and inter-

Fig. 1. Schematic illustration of the two-domain statistical thermodynamic model of TetR allostery. (A) The crystal structure of TetR(B) in complex with minocycline and magnesium (PDB code: 4AC0). The red residues are the hotspots identified in the DMS study (9). (B) Four possible conformations of a two-domain TetR molecule with their corresponding free energies (G). G of the $L_I D_I$ state is set to 0. Blue/green circle (square) denotes the inactive (active) state of LBD/DBD. (C) A simple repression scheme of TetR function. Binding of the ligand (inducer) favors the inactive state of DBD in TetR, which then releases the DNA operator and enables the transcription of the downstream gene. (D) Schematic free energy diagram of the possible binding states of TetR at fixed ligand and operator concentrations. Red, orange and purple arrows show how a mutation can disrupt allostery by 1. increasing ε_L; 2. decreasing ε_D and 3. decreasing γ. The L-$L_A D_A$ state is not explicitly shown in the last column as the doubly-bound L-$L_A D_A$-D state is expected to have a lower free energy. Note that mutations that change the binding affinities of the active LBD/DBD to ligand/operator are not discussed here as we focus on the intrinsic allosteric properties of the TF itself.

domain energetics in various ways, as far as the dominance of the L-$L_A D_I$ state is re-established, rationalizing the broad rescuing mutation distributions. Naturally, the rescuability of a dead mutation depends on which biophysical parameters it perturbs and by how much. At a qualitative level, the distributed nature of dead and rescuing mutations observed in the DMS measurement emerges naturally from the two-domain model (9).

In summary, the interplay of intra- and inter-domain properties that govern the free energy landscape of TetR's conformational and binding states, as elucidated by the two-domain model, aligns well with the unexpected DMS results on a qualitative level. To gain deeper insight into TetR allostery and the model itself, we aim to establish a quantitative framework with the model that accurately captures mutant induction curves in the next section.

System-Level Ramifications of the Two-Domain Model . The discussions in the previous section and Fig. 1 are primarily meant to give an intuitive and qualitative understanding of the two-domain model and rationalization of recent DMS measurements (9, 40). In this section, we further establish a quantitative connection between the model and the induction curves of TetR variants, leveraging the recent success of linking sequence-level perturbations to system-level responses (57, 58, 63–67). An induction curve describes the in vivo expression level of the TetR-regulated gene as a function of inducer concentration.

As described in the last section, the intra-domain parameters (ε_D, ε_L) and inter-domain parameter (γ) of the two-domain model collectively determine the free energy landscape of TetR (Fig. 1). Consequently, a degenerate relationship arises between combinations of parameter values and the induction level at a specific ligand concentration as measured in

the DMS study. To tease apart the allosteric effects of different parameters, we aim to formulate their connection to the full induction curve, which is characterized by (1) the expression level of the TF-regulated gene without ligand (leakiness), (2) gene expression level at saturating ligand concentration (saturation) and (3) the ligand concentration required for half-maximal expression (EC_{50}). As revealed in previous studies, these induction curve properties encode information for the key parameters of an allosteric regulation system, for example, the TF binding affinity to ligand and operator, equilibrium between different conformational states of the TF and the abundance of various essential molecules within the system. The establishment of a quantitative connection between the MWC model and the induction curve has yielded valuable insights into several allosteric systems (33, 57, 58). However, the analogous aspects within a multi-domain thermodynamic model for allostery require further investigation.

Inspired by the pioneering works on transcription with MWC models (57, 58), we derived a quantitative relation between the gene expression level in cells and the three biophysical parameters of the two-domain model (Eq. **1**). Briefly, the ratio of the expression level of a TF-regulated gene to that of an unregulated gene, termed fold change (FC) and bound between 0 and 1, is used to quantify gene expression. The gene expression level on the other hand, is assumed to be proportional to the probability that a promoter in the system is bound by RNA polymerase, p(RNAP). Thus, FC is evaluated as the ratio of p(RNAP) in the presence of the TF to that in the absence of the TF. Intuitively, p(RNAP) can be calculated based on the equilibria among different binding and conformational states of the TF (*SI Appendix*, Fig. S1 and S2), which are determined by the allosteric properties (ε_D, ε_L and γ) and several other parameters. Therefore, FC can be expressed as a function of these parameters, which under

Fig. 2. Schematic illustration of the characteristic effects of perturbations in the three biophysical parameters on the free energy landscape and the corresponding induction curves of a two-domain allosteric system. Panels A, B and C illustrate how changing ε_L, ε_D, and γ alone affects the free energy landscape for the binding states shown in Fig. 2*D* and the induction curve. For the black induction curve in (C), the values of ε_L and ε_D are also adjusted to aid visualization of the negative monotonicity of the gene expression level (fold change) as a function of ligand concentration. The green shade in the middle column separates the DNA-bound states from the rest. In the free energy landscapes shown in the middle column, ligand- or DNA-binding is always assumed when the corresponding domain is in the active conformation.

the assumption that intra- and inter-domain energetics adopt typical values of several $k_B T$, is simplified to Eq. **1**.

$$FC = \left(1 + R^* e^{-\varepsilon_D} \frac{1 + e^{-\varepsilon_L - \gamma}(\frac{c}{K})^2}{1 + e^{-\varepsilon_L}(\frac{c}{K})^2}\right)^{-1} \quad [1]$$

Here, R^* is the rescaled TF copy number in the cell; c is ligand concentration, and K is the dissociation constant of ligand to the repressor with an active LBD. As we focus on understanding how sequence level perturbations are manifested in the allosteric properties of the protein (depicted by ε_D, ε_L and γ), the residues we choose for mutational study here are mostly not in direct contact with either the ligand or the operator (*SI Appendix* for more discussions). Thus, K and R^*, which is determined by the TF copy number in the cell and the affinity of the operator to the repressor with an active DBD, are taken to be constants across mutations in all subsequent discussions (58). Detailed derivations of Eq. **1** are provided in the *SI Appendix*.

Despite the considerable degeneracy in the activity (inducible and noninducible) within the parameter space of the two-domain model (Fig. 1*D*), Eq. **1** demonstrates that mutations affecting distinct biophysical parameters can be discerned based on their characteristic effects on the induction curve.

First, as ε_L and γ have no effect on the leakiness of the induction curve, only mutations that modify ε_D can lead to its changes (Fig. **2** and *SI Appendix* for additional discussions). In addition, these mutations also change the level of saturation (FC value where the induction curve plateaus at large c). Thus, dead mutations that disrupt allostery by decreasing ε_D alone will uniquely feature a noticeably lower leakiness and a lower level of saturation compared with the WT (Fig. 1*D* and Fig. **2**, see also *SI Appendix* and Fig. S3 for more discussions). Second, mutations that solely perturb the other intra-domain property ε_L, a crucial determinant of TetR's ligand detection limit, primarily shift the EC_{50} of the induction curve (see Eq. **1**). As ε_L increases/decreases from the WT value, the induction curve is right/left shifted with leakiness and the

level of saturation remaining unchanged (see the blue and red curves of Fig. 2*A*). However, when ε_L further increases, the induction curve loses the sigmoidal shape, with its sharply varying tail region being the characteristic of a dead mutation that disrupts allostery mainly by increasing ε_L (purple curve of Fig. 2*A*, see *SI Appendix* and Fig. S3 for more discussions).

Finally, in the high concentration limit, Eq. **1** converges to a constant value (see Eq. **2**).

$$\lim_{c \to \infty} FC = \left(1 + R^* e^{-\varepsilon_D - \gamma}\right)^{-1} \quad [2]$$

Hence, mutations affecting γ alone will tune the saturation of a sigmoidal induction curve, which increases/decreases as γ increases/decreases (see red and purple curves of Fig. 2*C*). Therefore, in contrast to the two aforementioned scenarios, dead mutations that disrupt allostery through decreasing γ alone will feature a full sigmoidal induction curve with a low level of saturation and unchanged leakiness compared with the WT (with $\gamma > 0$, see the purple curve of Fig. 2*C*). Furthermore, as γ dictates the inter-domain cooperativity, it thereby controls the sign of the monotonicity of FC as a function of c (*SI Appendix* for detailed proof). Specifically, when $\gamma > / < 0$, there is negative/positive cooperativity between ligand and operator binding, and FC increases/decreases monotonically with c (see Fig. 2*C*). When $\gamma = 0$, however, the bindings of ligand and operator become independent of each other, and c no longer affects FC (*SI Appendix*, Fig. S3).

Having identified the distinctive impacts of the different allosteric parameters of the two-domain model on the induction curve, as outlined in Eq. **1**, we next apply the model to analyze actual experimental induction curves of TetR mutants. The analysis enables us to uncover the underlying biophysical factors contributing to diverse mutational effects. Additionally, it allows us to evaluate the model's validity in capturing TetR allostery by assessing its accuracy in reproducing the experimental data.

Fig. 3. Induction data of 15 TetR mutants and the corresponding parameter estimation results. (A) Shaded blue curves in each plot show the percentiles of the simulated fold-change measurements using the inferred posterior parameters of the mutant. The white data points represent the corresponding experimental induction measurement of 4 or more biological replicates (three replicates for C203V and G102D-HQQ). (B) The inferred parameter values of the 15 mutants. The error bars of ε_L and γ represent the 95^{th} percentile of the Bayesian posterior samples, while the error bar of ε_D is calculated based on the standard error of the mean (SEM) of the corresponding leakiness measurement. The horizontal lines indicate the WT parameter values for reference.

Extensive Induction Curves Fitting of TetR Mutants. With the diagnostic tools established, we mapped mutants to the parameter space of the two-domain model through fitting of their induction curves. We choose 15 mutants for analysis in this section, which contain mutations that span different regions in the sequence and structure of TetR (*SI Appendix*, Fig. S4). Five of the 15 mutants consist of a dead mutation G102D and one of its rescuing mutations (see the third row of Fig. 3*A*), while the other 10 contains the WT, 8 single mutants and a double mutant Q32A-E147G (see the first two rows of Fig. 3*A*). In all cases, fitting of an induction curve is divided into two steps; first, at $c = 0$, Eq. **1** can be rearranged to give

$$\varepsilon_D = -ln\left(\frac{leakiness^{-1} - 1}{R^*}\right), \qquad [3]$$

which enables the direct calculation of ε_D from the leakiness of the induction curve; second, ε_L and γ are inferred based on the remaining induction data using the method of Bayesian inference (58). Briefly, given ε_D calculated in the first step, we select a set of ε_L and γ values from physical ranges of the parameters, based on the probability of observing the induction curve with the parameter values using Monte Carlo sampling. The medians of the selected sets of ε_L and γ values,

which are known as posterior distributions, are reported as the inferred parameter values, and error bars show the 95% credible regions of the posterior distributions (Fig. 3*B* and *SI Appendix*, Fig. S12B). All details are provided in the "*Materials* and *Methods*" section and *SI Appendix*.

As shown in Fig. 3*A*, the fitted curves accurately capture the induction data of all the mutants studied here, enabling the quantification of their biophysical parameters with little uncertainty (see Fig. 3*B*). These results support the general applicability of the two-domain model to describing TetR allostery, as these 15 mutants vary significantly in terms of location of mutations and phenotype. They are one to four mutations away from the WT, and the sites of mutations are distributed across LBD, DBD and the domain interface (*SI Appendix*, Fig. S4). In terms of the phenotype, they fall into different classes as characterized by the FC value at $c = 1000$ nM (FC^{1000}), including dead ($FC^{1000} < 0.1$), enhanced induction ($FC^{1000} > FC_{WT}^{1000}$) and neutral activity (the rest).

Besides inspecting the goodness of fit, a closer examination of the induction curves and the Bayesian inference results reveal additional features of the two-domain model of TetR allostery. First, the apparent binding affinity of TetR to the

ligand (anhydrotetracycline; aTC) and operator (*tetO*2) used in our experiments are estimated to be about 17.4 $k_B T$ and 16.4 $k_B T$, respectively, based on the parameter values inferred for the WT (see *SI Appendix* for detailed calculations); these agree well with the range of reported experimental values ([68]–[73]). Second, as shown in Fig. 3*B*, the perturbations in ε_D are generally small in magnitude ($< 1 \; k_B T$) among the dead mutants, while substantially larger perturbations (up to ~ 5 $k_B T$) are observed for ε_L and γ. The latter observation is in line with the predictions of the model (see the previous section) that two types of qualitatively different induction curves are expected for partially dead mutants (see the first and second rows of Fig. 3*A*). Specifically, the induction curve of P176N-I174K-F177S (PIF) loses the sigmoidal shape, with its sharply varying tail region suggesting a significant increase of ε_L. This is confirmed by the parameter fitting result, which shows that the triple mutations of PIF, located in the core region of LBD (*SI Appendix* and Fig. S4), lead to the largest increase of ε_L among all mutants (4.5 $k_B T$ higher than the WT, see Fig. 3*B*). The induction curves of the other four dead mutants (R49G, D53H, P105M and G143M), however, maintain the sigmoidal shape yet with low levels of saturation. As they all exhibit a higher ε_D (leakiness) than the WT, the low levels of saturation have to result solely from weakened inter-domain couplings (Fig. 3). In other words, these four dead mutations, located mostly at the domain interface (*SI Appendix*, Fig. S4), disrupt allostery primarily through decreasing γ. Indeed, when the γ of these mutants is set to the WT value but keeping their respective ε_D and ε_L parameters unchanged, higher FC^{1000} values than the WT are obtained instead (*SI Appendix*, Fig. S10).

Despite the discussion of two limiting types of changes in the induction curves, it is interesting to observe in Fig. 3*B* that the three biophysical parameters, especially ε_L and γ, are often perturbed together in the mutants. Indeed, although the three parameters are theoretically independent of each other, at the structural level, it is less likely to have a scenario where a mutation perturbs inter-domain coupling (γ) but leaves intra-domain properties unchanged. Lastly, while quantification of G102D from its flat induction curve incurs large uncertainties, (*SI Appendix*, Fig. S3), the diversities of the induction curves and the biophysical parameters of the G102D-rescuing mutants (Fig. 3*B*) provide clear support for the qualitative anticipation from the previous section; i.e., rescuing mutations of a dead mutant may restore inducibility through different combinations of tuning ε_D, ε_L and γ. For example, while Y42M-I57N and K98Q both rescue G102D to achieve similar FC^{1000}, they exert distinct influences on ε_D and ε_L, as reflected in the variations of leakiness and EC_{50} between the corresponding induction curves. In another case, while the induction curves of G102D-T26A and G102D-L146A show comparable values of saturation and leakiness, their different EC_{50} values suggest the different effects of the two rescuing mutations on ε_L, further quantified by the inferred parameter values. The diverse mechanistic origins of the rescuing mutations revealed here provide a rational basis for the broad distributions of such mutations.

In summary, Eq. **1** accurately captures the induction curves of a comprehensive set of TetR mutants using a minimum set of parameters with realistic values (see the section "*Discussion*" and *SI Appendix*, Fig. S4). The two-domain model thus pro- vides a quantitative platform for investigating TetR allostery beyond an intuitive rationalization of the DMS results.

Exploring Epistasis between Mutations

With the 15 mutants mapped to the parameter space of the two-domain model, we now explore the epistatic interactions between the relevant mutations. To do so, we start by assuming additivity in the perturbation of all three biophysical parameters (see Eq. **4** and **5**, where p represents any one of ε_D, ε_L and γ).

$$p^{mut1+mut2} - p^{WT} = \alpha_{1,p}\delta p^{mut1} + \alpha_{2,p}\delta p^{mut2} \qquad [4]$$

$$\delta p^{mut1} = p^{mut1} - p^{WT} \qquad [5]$$

We then evaluate how the induction curves generated by this additive model $\alpha_{1,p} = \alpha_{2,p} = 1$) deviate from the corresponding experimental results, and the magnitude of deviation quantifies the significance of epistasis. Eight mutation combinations were chosen for the analysis, where we pair up C203V and Y132A, the two single mutations that enhance the level of induction relative to the WT, with mutations from different structural regions of TetR.

Although predictions from the additive model are qualitatively correct for 5 of the 8 mutation combinations on phenotypic effects, none of them captures the corresponding induction curve accurately (*SI Appendix*, Fig. S11). This again highlights the parameter space degeneracy of the phenotypes, and that mechanistic specificity is required of a model for making reliable predictions on combined mutants ([74]). For a deeper understanding of epistatic interactions between the mutations queried here, we next directly fit for the three biophysical parameters of the combined mutants using their induction curves, which are then compared with those obtained from the additive model.

As shown in *SI Appendix*, Fig. S12A, the induction data of all eight combined mutants are well captured by the fitted curves, which reaffirms the applicability of the two-domain model. Interestingly, despite the discrepancy between the experimental induction curves and those predicted by the additive model, the fitted parameters of the combined mutants are comparable to those from the additive model in many cases (see *SI Appendix*, Fig. S12B). In this regard, one noticeable example is the mutant C203V-G102D-L146A, for which the result of direct fitting is close to the additive model in terms of both ε_L and γ (with a difference of $\leqslant 0.5 \; k_B T$), while they differ more in ε_D (1.6 $k_B T$). Inspired by such observations, we then seek the minimal modifications to the additive model that can account for epistasis. As C203 is one of the most distant residues from DBD in TetR, we reason that its effect on ε_D may get dominated by mutations much closer to the DNA binding residues like G102D and L146A (*SI Appendix*, Fig. S4). Remarkably, when we quench C203V's contribution to ε_D in the additive model of C203V-G102D-L146A $\alpha_{C203V,\varepsilon_D} = 0$, see Eq. **4**), the predicted induction curve well recapitulates the experimental data (Fig. 4*G*). In another example, when C203V is combined with the triple mutations PIF, which are much closer to the DBD, the fitted parameters of the combined mutant also align well with the additive model except only for ε_D. Here, quenching C203V's contribution to ε_D again leads

Fig. 4. The induction curves for the eight combined TetR mutants from experimental measurement and prediction from the modified additive model. In each plot, black, orange and white points represent the experimental data for mutant 1, mutant 2 and the combined mutant (named mutant 1-mutant 2), specified in the legend and title. The blue band show the 95^{th} percentile of the induction curve prediction from the modified additive model. The modification to the basic additive model in each plot is specified by the 6 weights $\left\{ \alpha_{1,\varepsilon_D}, \alpha_{2,\varepsilon_D}, \alpha_{1,\varepsilon_L}, \alpha_{2,\varepsilon_L}, \alpha_{1,\gamma}, \alpha_{2,\gamma} \right\}$ (see Eq. **4**), which are (A) $\{1, 1, 1, 1, 1, 1\}$; (B) $\{1, 1, 0.5, 1, 1, 1\}$; (C) $\{1, 1, 0.5, 1, 1, 1\}$; (D) $\{1, 1, 0, 1, 1, 1\}$; (E) $\{0, 1, 1, 1, 0, 1\}$; (F) $\{0, 1, 1, 1, 0, 1\}$; (G) $\{0, 1, 1, 1, 1, 1\}$; (H) $\{0, 1, 1, 1, 1, 1\}$.

to good agreement between the additive model and experiment in terms of the induction curve (Fig. 4*H*).

The epistatic effects in other C203V-containing combined mutants can be largely accounted for by modifying the additive model following the same physical reasoning as above. For instance, mutation D53H is near the DNA-binding residues and located at the domain interface, which suggests its dominant role in defining the DBD energetics and inter-domain cooperativity when paired up with C203V (62). Accordingly, quenching C203V's effect on both ε_D and γ in the additive model for mutant C203V-D53H leads to dramatic improvement in the induction curve prediction (*SI Appendix*, Fig. S11F and Fig. 4*F*). Along this line, success is also observed when accounting for the epistasis between C203V and R49G by the same approach (Fig. 4*E*). These examples might also explain why C203V, although being able to enhance the induction of TetR, fails to rescue a range of dead mutations including R49A, D53V, G102D, N129D and G196D (9). More broadly, these observations together indicate that epistasis in combined mutants (e.g., those containing C203V) can be captured by the additive two-domain model with modifications based on physical reasoning.

The epistasis in combined mutants containing Y132A can be understood in a similar manner. Noticeably, the fitted parameter values of Y132A-C203V agree very well with the additive model for ε_D and γ (within a difference of 0.4 $k_B T$), while they differ in ε_L by 3.8 $k_B T$. Interestingly, such discrepancy in ε_L values can be essentially resolved by quenching Y132A's contribution in the additive model (reduced to 0.4 $k_B T$). This likely suggests that the effect of Y132A on ε_L tends to be compromised when combined with other mutations that perturb ε_L. Indeed, for most of the Y132A containing mutants investigated here (Y132A-C203V/PIF/R49G), the accuracy of induction curve prediction by the additive model improves greatly when $\alpha_{Y132A,\varepsilon_L}$ is tuned down (Fig. 4*B − D*, see *SI Appendix* for more discussions).

The only exception to this trend is the mutant Y132A-G102D-T26A, for which we observe strong epistasis between Y132A and the dead-rescue mutation pair G102D-T26A. Although both mutations Y132A and G102D-T26A enhance the allosteric response of TetR (Fig. 3), their combined effects radically change the sign of cooperativity between the two domains (γ), turning the ligand (aTC) from an inducer into a corepressor (Fig. 4*A* and *SI Appendix*, Fig. S12B). Here, large disparity exists between the additive model and the direct fitting result in all three biophysical parameters, which cannot be explained by simple modifications of the additive model.

Discussion

Allostery, a major regulatory mechanism in biology, has attracted intense research interest in the past few decades due to its complexity and implications in biomedicine and protein engineering. A central goal is to develop a quantitative understanding of the phenomena through a physical model that can be tested by comprehensive data. Along this line, one of us has in recent years advanced a function-centric approach to studying protein allostery with DMS, which provides a comprehensive and most direct test of our mechanistic understandings. Specifically, we have shown in an unbiased way that allostery hotspots and dead-rescue mutation pairs in four TetR family TFs are distributed across the protein structures (9, 40). This highlights that modifying the propagation of conformational distortions between the effector and active sites is not the only way to tune allosteric regulation (9).

The rich and surprising observations for the TetR family TFs call for a physical understanding of the underlying allostery mechanism, for which we resort to statistical thermodynamic models. Statistical thermodynamic models have played a central role in shaping our understanding of allostery (1, 2). In particular, Chure et al. have developed the methodology of fitting the MWC model to the induction data of LacI

Fig. 5. Distribution of the 23 investigated TetR mutants in the parameter space of the two-domain model illustrates the correlation between perturbations in ε_L and γ. Color of the contour plot encodes the fold change at $c = 1000$ nM calculated for each point in the two-dimensional space of ε_L and γ with the WT ε_D value. The color within the data point of each mutant is based on the FC^{1000} calculated with its specific ε_D value. The notation of each mutant is abbreviated based on the 1-letter codes of the residues that are mutated. The specific mutations corresponding to each letter code from upper left to lower right are C: C203V; Y: Y132A; GL: G102D-L146A; GT: G102D-T26A; R: R49G; D: D53H; GK: G102D-K98Q; E: E150Y; QE: Q32A-E147G; G: G143M; P: P105M; GYI: G102D-Y42M-I57N; GHQQ: G102D-H44F-Q47S-Q76K; PIF: P176N-I174K-F177S. The least-squares regression line between ε_L and γ is shown with their Pearson correlation coefficient (ρ).

repressors using Bayesian inference (58). This established the connection between sequence-level perturbations to system-level responses, which enables the exploration of mutational effects on transcription within the framework of biophysical models. However, as pointed out in several previous studies, the MWC model is not consistent with the observation that effector-bound TetR crystal structures are closer to the DNA-bound form compared to the apo crystal structures (5, 24, 34). Hence, it's at least difficult to provide a complete picture of TetR allostery with the MWC model. The ensemble allosteric model (EAM) on the other hand, presents a more flexible and detailed domain-specific view of allostery, which can be applied to more complex allosteric systems (e.g., allostery with intrinsically disordered proteins) (5, 34).

Inspired by these pioneering developments and our DMS results, in this work, we propose a two-domain statistical thermodynamic model, the parameter-activity degeneracy of which readily elucidates the distributed allosteric network observed in the DMS result of TetR (Fig. 1). On the other hand, the functional form derived in Eq. **1** establishes the qualitative differences in the impacts of various model parameters on the characteristics of the induction curve (Fig. 2); for example, dead mutations perturbing inter- and intra-domain properties are predicted to cause induction curves with and without a saturating plateau, which are both observed experimentally (Fig. 3). Moreover, Eq. **1** accurately captures the induction data of a diverse set of TetR mutants (Fig. 3, *SI Appendix*, Fig. S4 and S12) in a quantitative manner, enabling their mapping to the model parameter space with high precision.

The success of Eq. **1** allows for a quantitative investigation of epistasis between the characterized mutations. In a previous study of another TF, LacI, using the MWC model, no epistasis was observed between mutations of ligand-binding and DNA-binding residues analyzed therein (58). In the present analysis of TetR, epistasis is observed in all eight queried mutation combinations, which can be largely rationalized through physical reasoning. For example, our results intuitively indicate that the effect of a distant mutation like C203V on DBD energetics

and inter-domain coupling tend to be dominated by mutations much closer to these regions (Fig. 4 and *SI Appendix*, Fig. S11). Such phenomena suggest that these mutations affect allosteric regulation by influencing not only the relative populations of conformations in different ligation states, but also through shifting the dominant conformations themselves (29). On the other hand, epistasis involving Y132A is more complex. For instance, we observe that Y132A's influence on ε_L is diminished when paired up with other mutations in most cases. Further mechanistic understanding, likely from a biochemical perspective, is required to fully explain this phenomenon. Nonetheless, these observations provide a plausible explanation for why C203V and Y132A, although being able to enhance the induction response of TetR individually, could not rescue a range of dead mutations (9).

Our results reveal additional insights into the nature of two-domain allostery as well. Although our model assumes no *a priori* correlation between the intra- and inter-domain properties of the TF, we find that they are always modified together by mutations (Fig. 3 and *SI Appendix*, Fig. S12). This immediately points to the interconnectivity of allosteric networks (19, 24, 62). That is, when a residue is involved in the conformational rearrangements induced by both effector and operator binding, it functions as a bridge through which the two events are coupled. Within such a conceptual framework, it is highly unlikely to observe a mutation that changes γ alone without perturbing the intra-domain energetics.

This point is explicitly illustrated in Fig. 5, which summarizes the locations of all the characterized 23 mutants in the two-dimensional space of ε_L and γ. Here, the color of the contour plot encodes the FC^{1000} value calculated for each (ε_L, γ) combination together with the WT ε_D value, while the color of the specific mutant data points is based on the true FC^{1000} value. Thus, the color of a data point reflects the ε_D of the corresponding mutant relative to the WT; i.e., a brighter/darker color than the surrounding indicates that the mutant features a higher/lower ε_D value than the WT. Since the variations of ε_D among the investigated mutants are

modest compared with ε_L and γ (see Fig. 3B), we'll primarily focus on the latter two in the discussion. Evidently, mutants with the strongest allosteric responses feature small ε_L and large γ, which correspond to the upper left region of Fig. 5. Additional mutations in the background of these mutants lead to weaker allosteric response, moving the mutants to the darker regions of the plot. The interconnected nature of allosteric networks described by the two-domain model, however, dictates that such shifts would take place along the diagonal of the ε_L-γ plane, which describe mutations that modify intra- and inter-domain properties simultaneously. This feature potentially offers a functional advantage of preventing mutants from regions of low allosteric response (corresponding to upper right and lower left of Fig. 5), facilitating efficient adaptation to new effectors during evolution. In the future, it is of interest to examine whether such a negative correlation between ε_L and γ is a generic feature of two-domain allostery.

Our results establish the two-domain model as a flexible and quantitative platform for investigating TetR allostery, adding to the value of statistical thermodynamic models as advocated by the seminal works on MWC molecules (57, 58). We note, however, that the two-domain model is a more generic description of allostery compared with the MWC model.

First, although our model identifies the importance of intra-domain energetics in tuning the free energies of different TF states, it points out the more fundamental role of inter-domain coupling in defining allosteric response (Fig. 2 and *SI Appendix*). A large inter-domain coupling is assumed to be obligatory in the MWC model and required to create the two pre-existing conformations of different ligand and operator affinities in the first place. Various experimental studies as well as our own data reveal that the inter-domain coupling of TetR is amenable to tuning, especially by mutations of residues at the domain interface (24, 62, 75, 76). In some cases, even single mutations can reverse the sign of cooperativity between LBD and DBD, indicating that the dead mutational effects observed in the DMS experiments and elsewhere could well originate from diminishing γ. These considerations necessitate an explicit treatment of γ in a physical model of TetR allostery.

Second, the MWC model, when applied to two-domain systems, contains at least four biophysical parameters (57, 58), is thus more complex than our two-domain model, which requires only three. The simplicity of our model reduces its parameter space degeneracy, enabling high precision in fitting while being able to accurately describe a broad set of induction data. The three parameters also offer an interpretable mechanistic picture of TetR allostery in a physically transparent fashion.

We have focused our experimental and theoretical analyses on the TetR-family TFs, which represent important systems for building allosteric models, due to their broad involvement in many aspects of cell physiology (35, 55). Along this line, a natural extension of the current study is to use the two-domain model as a platform for comparing allostery in different TetR homologs (40). For example, coupling the model with systematic dose-response study of hotspot mutations could reveal the roles of different allosteric hotspots (whether they dictate intra- or inter-domain properties or both). The similarity in key structural features among TetR homologs then enables the comparison of the nature of hotspots at similar structural locations. This could potentially lead to novel insights into the different roles that sequence and structure play in defining

the allostery of a system, which is a fundamental question to address in rational engineering of allosteric proteins.

Although our two-domain model is inspired by and established with the data of TetR family TFs, the mechanistic and modelling insights gained here should be generally applicable to other allosteric systems sharing the two-domain architecture. This includes other transcription factors like catabolite activator protein (CAP), a homodimeric TF with a cyclic AMP (effector) binding domain and a DNA binding domain (36); receptors like pentameric ligand-gated ion channels (pLGICs), which allosterically translate the binding of neurotransmitters to their extracellular domains into the activation of ionotropic pores located in their transmembrane domains (59, 60); and allosteric enzymes like aspartate transcarbamoylase (ATCase), in which binding of effector like ATP to the regulatory domain allosterically alters the catalytic activity of the functional domain (37, 38). Ultimately, we envision that such model provides a minimalist but physically sound framework for integrating different types of experimental data and computational methods (40, 61, 77, 78) for exploring rational engineering and regulation of allostery for biotechnological and biomedical applications.

Materials and Methods

Library Construction. Using a low-copy backbone (SC101 origin of replication) carrying spectinomycin resistance, we constructed a sensor plasmid with TetR(B) (Uniprot P04483). The *tetRb* gene was driven by a variant of promoter apFAB61 and Bba_J61132 RBS (79). On a second reporter plasmid, superfolder GFP (80) was cloned into a high-copy backbone (ColE1 origin of replication) carrying kanamycin resistance. The expression of the superfolder GFP reporter was placed under the control of the ptetO promoter. To control for plasmid copy number, RFP was constitutively expressed with the BBa_J23106 promoter and Plotkin RBS (79) in a divergent orientation to sfGFP.

Library Synthesis. A comprehensive single-mutant TetR library was generated by replacing wildtype residues at positions 2–207 of TetR to all other 19 canonical amino acids (3,914 total mutant sequences). Oligonucleotides encoding each single-point mutation were synthesized as single-stranded Oligo Pools from Twist Bioscience and organized into 6 subpools, spanning six segments of the *tetRb* gene, corresponding to residues 2–39, 40–77, 78–115, 116–153, 154–191, and 192–207 of TetR(B) respectively. Additional sequence diversity was observed in the library due to error rates in the synthesis of single-stranded Oligo Pools, leading to the downstream identification of some double and triple mutant TetR variants. Oligo pools were encoded as a concatemer of the forward priming sequence, a BasI restriction site (5′-GGTCTC), six-base upstream constant region, TetR mutant sequence, six-base downstream constant region, a BsaI site (5′-GAGACC), and the reverse priming sequence. Subpools were resuspended in double-distilled water (ddH_2O) to a final molal concentration of 25 ng/μL and amplified using primers specific to each oligonucleotide subpool with KAPA SYBR FAST qPCR (KAPA Biosystems; 1-ng template). A second PCR amplification was performed with KAPA HiFi (KAPA Biosystems; 1-μL qPCR template, 15 cycles maximum). We amplified corresponding regions of pSC101_TetR_specR with primers

that linearized the backbone, added a BsaI restriction site, and removed the replaced wildtype sequence. Vector backbones were further digested with DpnI, BsaI, and Antarctic phosphatase before library assembly.

We assembled mutant libraries by combining the linearized sensor backbone with each oligo subpool at a molar ratio of 1:5 using Golden Gate Assembly Kit (New England Biolabs; 37 °C for 5 min and 60 °C for 5 min, repeated 30 times). Reactions were dialyzed with water on silica membranes (0.025-μm pores) for 1 h before transformed into DH10B cells (New England Biolabs). Library sizes of at least 100,000 colony-forming units (cfu) were considered successful. DH10B cells containing the reporter pColE1_sfGFP_RFP_kanR were transformed with extracted plasmids to obtain cultures of at least 100,000 cfu. Following co-transformation, the cultures for each subpool were stored as glycerol stocks and kept at -80 °C.

Fluorescence Activated Cell Sorting. The subpool library cultures were seeded from glycerol stocks into 3 mL lysogeny broth (LB) containing 50 μg/mL kanamycin (kan) and 50 μg/mL spectinomycin (spec) and grown for 16 h at 37 °C. Each culture was then back-diluted into two wells of a 96-well plate containing LB kan/spec media using a dilution factor of 1:50 and grown for a period of 5 hours at 37 °C. Following incubation, each well containing saturated library culture was diluted 1:75 into 1× phosphate saline buffer (PBS), and fluorescence intensity was measured on a SH800S Cell Sorter (Sony). We first gated cells to remove debris and doublets and selected for variants constitutively expressing RFP. Using this filter, we then proceeded to draw an additional gate to select for mutants that displayed low fluorescence intensities in the absence of anhydrotetracycline (aTC). This gate allowed us to select for mutants which retained the ability to repress GFP expression in the presence of an inducer, using prior measurements of WT TetR(B) as a reference for selecting repression competent mutants (9). Utilizing these gates, we sorted 500,000 events for each gated population and recovered these cells in 5 mL of LB at 37 °C before adding 50 μg/mL of kan and spec. Following the addition of antibiotics, the cultures were incubated at 37 °C for 16 hours.

Following overnight growth, each culture was back diluted into two wells of a 96-well plate containing LB kan/spec media using a 1:50 dilution factor. Upon reaching an OD600 $\sim$ 0.2, aTC was added to one of the two wells at a final concentration of 1 μM, with the well not receiving aTC serving as the uninduced control population. Cells were then incubated for another 4-hour growth period at 37 °C before being diluted into 1x PBS using a dilution factor of 1:75, and fluorescence intensity was measured on a SH800S Cell Sorter.

We first gated cells to remove debris and doublets and selected for variants constitutively expressing RFP before obtaining the distribution of fluorescence intensities (FITC-A) across the uninduced and induced populations of each of the 6 subpools (*SI Appendix*, Fig. S13). Using the uninduced control population of each subpool as a reference, gates were drawn to capture mutants which displayed no response to the presence of aTC in the induced populations. Using these gates, 500,000 events were sorted from the induced populations of each subpool and cells were subsequently recovered at 37 °C using 5 mL of LB media before being plated onto LB agar plates containing 50 μg/mL kan and spec.

Clonal Screening of Dead Mutants. To screen for functionally deficient variants of TetR, 100 individual colonies were picked from each subpool and grown to saturation in a 96-well plate for 6 hours. Saturated cultures were then diluted 1:50 in LB-kan/spec media and grown in the presence and absence of 1 μM aTC for 6 hours before OD600 and GFP fluorescence (Gain: 40; Excitation: 488/20; Emission: 525/20) were read on a multiplate reader (Synergy HTX, BioTek). Fluorescence was normalized to OD600, and the fold inductions of each clone were calculated by dividing its normalized fluorescence in the presence of inducer by the normalized fluorescence in the absence of inducer. The fold-induction of WT TetR was tested in parallel with these screens to serve as a benchmark for function, and clones which displayed < 50% activity to WT were replated and validated in triplicate before being sent for sanger sequencing (Functional Biosciences).

aTC Dose Response Measurements. TetR mutants identified during clonal screening were reinoculated into 3 mL of LB Kan/Spec media and grown overnight. After an overnight growth period, each culture was added to 4 rows of 12 wells in a 96-well plate containing LB Kan/Spec media using a dilution factor of 1:50. Each mutant was tested across 12 different concentrations of aTC in quadruplicate fashion, with each row having an identical concentration gradient across the 12 wells. The aTC concentration gradient across the 12 wells were 0 nM, 10 nM, 20 nM, 40 nM, 60 nM, 80 nM, 100 nM, 150 nM, 250 nM, 500 nM, 750 nM, and 1μM aTC.

Following reinoculation into 96-well plates, selected mutants were placed on a plate shaker set to 900 RPM and incubated at 37 °C for 5 hours. Following this 5-hour growth period, the plates were removed from the plate shaker and the OD600 and GFP fluorescence (Gain: 40; Excitation: 488/20; Emission: 525/20) of each well was read using a multiplate reader (Synergy HTX, BioTek). Fluorescence was normalized to OD600 for each well. The normalized fluorescence at each concentration was measured for 4 replicates of each mutant (unless otherwise stated). The fluorescence of GFP reporter-only control (no TetR), grown under identical conditions, was measured in the same way. This reporter-only control served as an upper bound to the fluorescence that could be measured in the experiment and was accompanied by a WT TetR-GFP reporter control, used to certify the precision of measurements made at different times.

Synthesis of Combined TetR mutants. To assess the predictive capability of the additive model, eight combined mutants were selected to undergo aTC dose response characterization. These combined mutants were synthesized using clonal DNA fragments from Twist Biosciences coding for the amino acid sequence of each of the selected combined mutants. The gene fragments were individually cloned into sensor plasmid backbones identical to those used in the previous identification of non-functional TetR(B) mutants following the manufacturers protocol for Gibson Assembly (New England Biolabs). The newly constructed plasmids carrying each of the combined mutants were then transformed into E. coli (DH10B) cells containing a GFP reporter plasmid identical to that used in the previous characterization of non-functional TetR(B) mutants. After a 1-hour recovery period, 100 μL of recovered cells were plated onto LB kan/spec agar plates for each of the combination mutants and incubated for 16 hours at 37 °C.

Following overnight growth, individual colonies were picked and inoculated into 3 mL of LB kan/spec media to be grown overnight at 37 °C. Following overnight growth, samples from each culture were submitted for sanger sequencing to confirm the identity of each of the synthesized combined mutants. Following sequence verification, the aTC dose-response behavior of each combined mutant was characterized using an experimental setup consistent with dose-response measurements made for the previously identified TetR(B) mutants.

Model Parameter Estimation. As described above, the estimation of the three biophysical parameters (ε_D, ε_L and γ) for each mutant is divided into two steps. Firstly, as ε_L and γ do not affect FC at $c = 0$ (Eq. **1**), we directly calculate the value of ε_D from the leakiness of an induction curve, which is measured with high precision in our experiments. With the calculated ε_D (ε_D^{WT} is set to 0 as reference), we then fit the full induction curve to obtain the values of ε_L and γ using the method of Bayesian inference. Here, we first construct the statistical model that describes the data generating process based on Eq. **1**, and specify the prior distributions of the relevant parameters. This enables the derivation of the conditional probability of the parameter values of a mutant given its induction data, known as the posterior distribution. The posterior distribution is then sampled using Markov chain Monte Carlo (MCMC), from which we infer the values of ε_L and γ directly. The validity of the statistical model and computational algorithm is fully tested with several metrics, and all details regarding the parameter estimation process briefed here are provided in the *SI Appendix*.

Code and Data Availability. All code and data used in this work can be accessed at the Github repository https://github.com/liuzhbu/Two_Domain_Allostery.

ACKNOWLEDGMENTS. This work is funded by NIH Director's New Innovator Award DP2GM132682 (SR) and Shaw Scientist Award (SR), and R35-GM141930 (QC). Development of the Bayesian inference model was partially supported by grant ML-21-016 from the Dreyfus foundation (QC). Computational resources from the Extreme Science and Engineering Discovery Environment (XSEDE), which is supported by NSF grant number ACI-1548562, are greatly appreciated; part of the computational work was performed on the Shared Computing Cluster which is administered by Boston University's Research Computing Services (URL: https://www.bu.edu/tech/ support/research/).

1. F Bacon, On the nature of allosteric transitions: a plausible model. *J Mol Biol* **12**, 88–118 (1965).
2. DE Koshland Jr, G Némethy, D Filmer, Comparison of experimental binding data and theoretical models in proteins containing subunits. *Biochemistry* **5**, 365–385 (1966).
3. JP Changeux, SJ Edelstein, Allosteric mechanisms of signal transduction. *Science* **308**, 1424–1428 (2005).
4. Q Cui, M Karplus, Allostery and cooperativity revisited. *Protein science* **17**, 1295–1307 (2008).
5. HN Motlagh, JO Wrabl, J Li, VJ Hilser, The ensemble nature of allostery. *Nature* **508**, 331–339 (2014).
6. EW Yu, DE Koshland Jr, Propagating conformational changes over long (and short) distances in proteins. *Proc. Natl. Acad. Sci.* **98**, 9517–9520 (2001).
7. GM Süel, SW Lockless, MA Wall, R Ranganathan, Evolutionarily conserved networks of residues mediate allosteric communication in proteins. *Nat. structural biology* **10**, 59–69 (2003).
8. NV Dokholyan, Controlling allosteric networks in proteins. *Chem. reviews* **116**, 6463–6487 (2016).
9. M Leander, Y Yuan, A Meger, Q Cui, S Raman, Functional plasticity and evolutionary adaptation of allosteric regulation. *Proc. Natl. Acad. Sci.* **117**, 25445–25454 (2020).
10. A Peracchi, A Mozzarelli, Exploring and exploiting allostery: Models, evolution, and drug targeting. *Biochimica et Biophys. Acta (BBA)-Proteins Proteomics* **1814**, 922–933 (2011).
11. SJ Wodak, et al., Allostery in its many disguises: from theory to applications. *Structure* **27**, 566–578 (2019).
12. J Wang, et al., Mapping allosteric communications within individual proteins. *Nat. communications* **11**, 3862 (2020).
13. SW Lockless, R Ranganathan, Evolutionarily conserved pathways of energetic connectivity in protein families. *Science* **286**, 295–299 (1999).
14. MD Daily, JJ Gray, Allosteric communication occurs via networks of tertiary and quaternary motions in proteins. *PLoS computational biology* **5**, e1000293 (2009).
15. GJ Rodriguez, R Yao, O Lichtarge, TG Wensel, Evolution-guided discovery and recoding of allosteric pathway specificity determinants in psychoactive bioamine receptors. *Proc. Natl. Acad. Sci.* **107**, 7787–7792 (2010).
16. J Lee, et al., Surface sites for engineering allosteric control in proteins. *Science* **322**, 438–442 (2008).
17. AS Walker, WP Russ, R Ranganathan, A Schepartz, Rna sectors and allosteric function within the ribosome. *Proc. Natl. Acad. Sci.* **117**, 19879–19887 (2020).
18. M Sevvana, et al., An exclusive α/β code directs allostery in tetr–peptide complexes. *J. molecular biology* **416**, 46–56 (2012).
19. K Takeuchi, M Imai, I Shimada, Conformational equilibrium defines the variable induction of the multidrug-binding transcriptional repressor qacr. *Proc. Natl. Acad. Sci.* **116**, 19963–19972 (2019).
20. S Marzen, HG Garcia, R Phillips, Statistical mechanics of monod–wyman–changeux (mwc) models. *J. molecular biology* **425**, 1433–1460 (2013).
21. JP Changeux, Allostery and the monod-wyman-changeux model after 50 years. *Annu. review biophysics* **41**, 103–133 (2012).
22. J Guo, HX Zhou, Protein allostery and conformational dynamics. *Chem. reviews* **116**, 6503–6515 (2016).
23. O Schueler-Furman, SJ Wodak, Computational approaches to investigating allostery. *Curr. Opin. Struct. Biol.* **41**, 159–171 (2016).
24. SE Reichheld, Z Yu, AR Davidson, The induction of folding cooperativity by ligand binding drives the allosteric response of tetracycline repressor. *Proc. Natl. Acad. Sci.* **106**, 22263–22268 (2009).
25. C Xu, D Tobi, I Bahar, Allosteric changes in protein structure computed by a simple mechanical model: hemoglobin t r2 transition. *J. molecular biology* **333**, 153–168 (2003).
26. L Nierzwicki, et al., Enhanced specificity mutations perturb allosteric signaling in crispr-cas9. *Elife* **10**, e73601 (2021).
27. KW East, et al., Allosteric motions of the crispr–cas9 hnh nuclease probed by nmr and molecular dynamics. *J. Am. Chem. Soc.* **142**, 1348–1358 (2019).
28. J Liu, R Nussinov, Allostery: an overview of its history, concepts, methods, and applications. *PLoS computational biology* **12**, e1004966 (2016).
29. Y Zhang, et al., Intrinsic dynamics is evolutionarily optimized to enable allosteric behavior. *Curr. opinion structural biology* **62**, 14–21 (2020).
30. A Szabo, M Karplus, A mathematical model for structure-function relations in hemoglobin. *J. molecular biology* **72**, 163–197 (1972).
31. C Viappiani, et al., Experimental basis for a new allosteric model for multisubunit proteins. *Proc. Natl. Acad. Sci.* **111**, 12758–12763 (2014).
32. ER Henry, et al., Allosteric control of hemoglobin s fiber formation by oxygen and its relation to the pathophysiology of sickle cell disease. *Proc. Natl. Acad. Sci.* **117**, 15018–15027 (2020).
33. WA Eaton, A retrospective on statistical mechanical models for hemoglobin allostery. *The J. Chem. Phys.* **157** (2022).
34. VJ Hilser, JO Wrabl, HN Motlagh, Structural and energetic basis of allostery. *Annu. review biophysics* **41**, 585–609 (2012).
35. JL Ramos, et al., The tetr family of transcriptional repressors. *Microbiol. molecular biology reviews* **69**, 326–356 (2005).
36. SR Tzeng, CG Kalodimos, Protein activity regulation by conformational entropy. *Nature* **488**, 236–240 (2012).
37. A Velyvis, YR Yang, HK Schachman, LE Kay, A solution nmr study showing that active site ligands and nucleotides directly perturb the allosteric equilibrium in aspartate transcarbamoylase. *Proc. Natl. Acad. Sci.* **104**, 8815–8820 (2007).
38. WN Lipscomb, ER Kantrowitz, Structure and mechanisms of escherichia coli aspartate transcarbamoylase. *Accounts chemical research* **45**, 444–453 (2012).
39. K Gunasekaran, B Ma, R Nussinov, Is allostery an intrinsic property of all dynamic proteins? *Proteins: Struct. Funct. Bioinforma.* **57**, 433–443 (2004).
40. M Leander, Z Liu, Q Cui, S Raman, Deep mutational scanning and machine learning reveal structural and molecular rules governing allosteric hotspots in homologous proteins. *eLife* **11**, e79932 (2022).
41. DM Fowler, et al., High-resolution mapping of protein sequence-function relationships. *Nat. Methods* **7**, 741–746 (2010).
42. DM Fowler, S Fields, Deep mutational scanning: a new style of protein science. *Nat. Methods* **11**, 801–807 (2014).
43. KS Sarkisyan, et al., Local fitness landscape of the green fluorescent protein. *Nature* **533**, 397–401 (2016).
44. JM Flynn, et al., Comprehensive fitness maps of hsp90 show widespread environmental dependence. *eLife* **9** (2019).
45. TN Starr, et al., Deep mutational scanning of sars-cov-2 receptor binding domain reveals constraints on folding and ace2 binding. *Cell* **182**, 1295 – 1310.e20 (2020).
46. P Huss, A Meger, M Leander, K Nishikawa, S Raman, Mapping the functional landscape of the receptor binding domain of t7 bacteriophage by deep mutational scanning. *eLife* **10**, e63775 (2021).
47. DS Tack, et al., The genotype-phenotype landscape of an allosteric protein. *Mol. Syst. Biol.* **17**, e10179 (2021).
48. AJ Faure, et al., Mapping the energetic and allosteric landscapes of protein binding domains. *Nature* **604**, 175 – 183 (2022).
49. EM Jones, et al., Structural and functional characterization of g protein–coupled receptors with deep mutational scanning. *eLife* **9**, e54895 (2020).

50. JW McCormick, MA Russo, S Thompson, A Blevins, KA Reynolds, Structurally distributed surface sites tune allosteric regulation. *eLife* **10**, e68346 (2021).

51. N Ota, DA Agard, Intramolecular signaling pathways revealed by modeling anisotropic thermal diffusion. *J. molecular biology* **351**, 345–54 (2005).

52. D Strickland, K Moffat, TR Sosnick, Light-activated dna binding in a designed allosteric protein. *Proc. Natl. Acad. Sci.* **105**, 10709 – 10714 (2008).

53. KA Reynolds, RN McLaughlin, R Ranganathan, Hot spots for allosteric regulation on protein surfaces. *Cell* **147**, 1564–1575 (2011).

54. BRC Amor, MT Schaub, SN Yaliraki, M Barahona, Prediction of allosteric sites and mediating interactions through bond-to-bond propensities. *Nat. Commun.* **7** (2016).

55. L Cuthbertson, JR Nodwell, The tetr family of regulators. *Microbiol. Mol. Biol. Rev.* **77**, 440 – 475 (2013).

56. K Fukami-Kobayashi, Y Tateno, K Nishikawa, Parallel evolution of ligand specificity between laci/galr family repressors and periplasmic sugar-binding proteins. *Mol. biology evolution* **20**, 267–77 (2003).

57. RD Daber, M Sochor, M Lewis, Thermodynamic analysis of mutant lac repressors. *J. molecular biology* **409**, 76–87 (2011).

58. G Chure, et al., Predictive shifts in free energy couple mutations to their phenotypic conse­quences. *Proc. Natl. Acad. Sci. United States Am.* **116**, 18275 – 18284 (2019).

59. L Sauguet, A Shahsavar, M Delarue, Crystallographic studies of pharmacological sites in pentameric ligand-gated ion channels. *Biochimica et biophysica acta* **1850 3**, 511–523 (2015).

60. H Hu, RJ Howard, U Bastolla, E Lindahl, M Delarue, Structural basis for allosteric transitions of a multidomain pentameric ligand-gated ion channel. *Proc. Natl. Acad. Sci.* **117**, 13437 – 13446 (2020).

61. Y Yuan, J Deng, Q Cui, Molecular dynamics simulations establish the molecular basis for the broad allostery hotspot distributions in the tetracycline repressor. *J. Am. Chem. Soc.* (2022).

62. O Scholz, et al., Activity reversal of tet repressor caused by single amino acid exchanges. *Mol. Microbiol.* **53** (2004).

63. HG Garcia, R Phillips, Quantitative dissection of the simple repression input–output function. *Proc. Natl. Acad. Sci.* **108**, 12173 – 12178 (2011).

64. RC Brewster, et al., The transcription factor titration effect dictates level of gene expression. *Cell* **156**, 1312–1323 (2014).

65. FM Weinert, RC Brewster, M Rydenfelt, R Phillips, WK Kegel, Scaling of gene expression with transcription-factor fugacity. *Phys. review letters* **113**, 258101 (2014).

66. M Rydenfelt, RS Cox, HG Garcia, R Phillips, Statistical mechanical model of coupled tran­scription from multiple promoters due to transcription factor titration. *Phys. review. E, Stat. nonlinear, soft matter physics* **89**, 012702 (2014).

67. M Razo-Mejia, et al., Tuning transcriptional regulation through signaling: A predictive theory of allosteric induction. *Cell systems* **6**, 456 – 469.e10 (2017).

68. O Scholz, P Schubert, M Kintrup, WP Hillen, Tet repressor induction without mg2+. *Biochem­istry* **39**, 10914–10920 (2000).

69. P Schubert, K Pfleiderer, WP Hillen, Tet repressor residues indirectly recognizing anhydrote­tracycline. *Eur. journal biochemistry* **271**, 2144–2152 (2004).

70. S Kędracka-Krok, Z Wasylewski, Kinetics and equilibrium studies of tet repressor–operator interaction. *J. Protein Chem.* **18**, 117–125 (1999).

71. A Kamionka, J Bogdanska-Urbaniak, O Scholz, WP Hillen, Two mutations in the tetracycline repressor change the inducer anhydrotetracycline to a corepressor. *Nucleic acids research* **32**, 842–847 (2004).

72. DS Bolintineanu, et al., Investigation of changes in tetracycline repressor binding upon muta­tions in the tetracycline operator. *J. Chem. Eng. Data* **59**, 3167 – 3176 (2014).

73. D Normanno, et al., Probing the target search of dna-binding proteins in mammalian cells using tetr as model searcher. *Nat. Commun.* **6**, 7357 (2015).

74. X Li, B Lehner, Biophysical ambiguities prevent accurate genetic prediction. *Nat. Commun.* **11**, 4923 (2020).

75. G Müller, et al., Characterization of non-inducible tet repressor mutants suggests conforma­tional changes necessary for induction. *Nat. Struct. Biol.* **2**, 693–703 (1995).

76. B Hecht, G Müller, WP Hillen, Noninducible tet repressor mutations map from the operator binding motif to the c terminus. *J. Bacteriol.* **175**, 1206 – 1210 (1993).

77. J Xie, W Zhang, X Zhu, M Deng, L Lai, Coevolution-based prediction of key allosteric residues for protein function regulation. *eLife* **12** (2022).

78. PD Tonner, AD Pressman, D Ross, Interpretable modeling of genotype–phenotype landscapes with state-of-the-art predictive power. *Proc. Natl. Acad. Sci. United States Am.* **119** (2021).

79. S Kosuri, et al., Composability of regulatory sequences controlling transcription and translation in escherichia coli. *Proc. Natl. Acad. Sci.* **110**, 14024 – 14029 (2013).

80. J Pedelacq, S Cabantous, TH Tran, TC Terwilliger, GS Waldo, Engineering and characterization of a superfolder green fluorescent protein. *Nat. Biotechnol.* **24**, 79–88 (2006).