Cell-free chromatin particles from dying cells promote the induction of an immune response in human lymphocytes

Authors:

Snehal Shabrish[1,2], Gorantla V. Raghuram[1,2], Nimisha Raphael[1,2], Ranjani Narayanan[1,2], Relestina Lopes[1,2], Dipali Kondhalkar[1,2], Naveen Kumar Khare[1,2], Indraneel Mittra[1,2] *

Affiliations:

[1]Translational Research Laboratory, Advanced Centre for Treatment, Research and Education in Cancer, Tata Memorial Centre; Kharghar, Navi Mumbai – 410210, India.

[2]Homi Bhabha National Institute; Anushakti Nagar, Mumbai – 400094, India.

*Corresponding author. Email: imittra@actrec.gov.in

Abstract

It is long established that cell death and immune response are closely related, although the nature of this relationship has remained unclear. We earlier reported that cell-free chromatin particles (cfChPs) released from the billions of cells that die in the body every day to enter into the blood circulation, or those that are released locally from dying cells, can readily enter into healthy cells to induce DNA damage and activate inflammatory cytokines. In this study we investigated whether cfChPs from dying cells might be the missing link between cell death and immune response. We treated human lymphocytes with cfChPs isolated from sera of healthy individuals or with cfChPs released from hypoxia-induced dying lymphocytes. We observed that cfChPs from both sources readily entered into lymphocytes to accumulate in their nuclei within 2 h. This was associated with marked activation of CD69 and release of inflammatory cytokines, as well as of p-STING and cGAS expression. The addition of the STING protein inhibitor H151 to the cfChPs treated cells abolished the release of inflammatory cytokines. These findings lead us to suggest that cfChPs from dying cells are the critical triggers of immune response which act via cGAS-STING pathway.

Introduction

The mechanism that triggers immune response is a central but an unresolved question in immunology, although there is a well-established association between immune response and cell death (Ashida et al., 2011). Cell death results in the release of cellular components which are thought to act as damage-associated molecular patterns (DAMPs) to activate immune response (Akira et al., 2006; Chaplin, 2010; Land, 2015; Rock et al., 2011). Optimal levels of DAMPs in healthy individuals are required to maintain immune homeostasis; however, in pathological conditions such as sepsis, cancer, neurodegenerative disorders, and auto-inflammatory as well as autoimmune diseases, excessive cell death leads to an uncontrolled release of DAMPs that can provoke hyper-inflammatory responses (Denning et al., 2019; Land, 2015; Rock et al., 2011). However, despite intensive research, the precise component of DAMPs that might be critical for mounting immune response continues to remain obscure, hindering the development of effective therapies for life-threatening disorders (Fleischmann et al., 2016; Schaefer, 2014; Ye et al., 2020).

We have previously reported that cell-free chromatin particles (cfChPs) that are released from dying cells are readily internalized by healthy cells wherein they induce DNA damage and inflammation (Indraneel Mittra et al., 2017; Mittra et al., 2015). We have also shown that cfChPs play an important role in disorders associated with hyper-inflammation, such as toxicity from chemo- and radiotherapy, sepsis, ageing, and cancer (Agarwal et al., 2022; Kirolikar et al., 2018; I. Mittra et al., 2017; Mittra Id et al., 2020; Ostwal et al., 2022; Pal et al.,

2022; Pilankar et al., 2022). Since inflammation is a critical component of immune response, these observations led us to hypothesize that cfChPs released from dying cells could be the elusive agents that could link cell death with an inflammatory immune response (Shabrish and Mittra, 2021). In pursuit of this hypothesis, we utilized two experimental models: 1) a culture system wherein peripheral blood mononuclear cells (PBMCs) were exposed to conditioned medium (CM) containing cfChPs released from hypoxia-induced dying PBMCs, and 2) direct treatment of PBMCs with cfChPs isolated from healthy human sera. The immunological effects of cfChPs were investigated by evaluating three integral aspects of an inflammatory immune response viz. 1) immune cell activation characterized by CD69 expression on T cells and NK cells, 2) cellular stress response, and 3) activation of inflammatory cytokines.

Results

cfChPs released from dying cells are readily internalized by PBMCs

As a first step, we revisited our previous finding that cfChPs released from dying cells are readily internalized by mouse fibroblast cells (Indraneel Mittra et al., 2017) and confirmed that cfChPs released from dying PBMCs are also internalized by healthy PBMCs. To demonstrate this, PBMCs were dually fluorescently labelled in their DNA and histones using bromodeoxyuridine (BrdU) and histone-2B-GFP, respectively (see Materials and Methods), and induced to undergo cell death by hypoxia. The CM of the dying PBMCs, when added to

80 fresh PBMCs, showed copious accumulation of fluorescently dually labelled

81 cfChPs in their nuclei at 4 h **(Suppl. Fig. 1).**

Immunogenic effects of CM containing cfChPs

We next investigated the possible immunogenic effects of cfChPs that are

contained in CM by evaluating CD69 expression, activation of cellular stress

markers, and cytokine release by treating fresh PBMCs with hypoxic CM. To

determine the optimum volume of CM required, a dose-response analysis was

initially performed, which showed that maximum CD69 expression on T- and NK-

cells was achieved with 200 µL CM **(Fig. 1A).** All further experiments were

therefore performed using 200 µl CM. The treatment of PBMCs with 200 µL CM

was also associated with marked activation of the inflammatory cytokines IL-1β,

IL-6, TNF-α, IFN-γ, and CXCL9 **(Fig. 1B).** Since stress-related proteins interact

with and regulate signaling intermediates involved in an immune response, we

next investigated whether cfChPs from dying cells would activate the cellular

stress-related markers c-Jun, c-Fos, JunB, FosB, EGR-1, and NF-KB. Activation

of stress markers was evaluated utilizing qRT-PCR in a time course experiment,

which showed upregulation of all six stress markers albeit at different time points

(Suppl. Fig. 2). Of these, we could confirm the expression of FosB at a protein

level by flow cytometry when examined at the maximum time-point of its mRNA

expression **(Fig. 1C)**.

Serum derived cfChPs are readily internalized by PBMCs

We previously reported that cfChPs isolated from human sera are readily

internalized by mouse fibroblast cells (Mittra et al., 2015). Microarray analysis

had suggested that cellular uptake of cfChPs likely occurs via the phagocytosis

pathway (Indraneel Mittra et al., 2017). Other publications have also shown that

nucleosome uptake happens via endocytosis (Wang et al., 2021). Thus, we

speculated, whether serum derived cfChPs, like those contained in hypoxic CM,

might also activate inflammatory cytokines? As a first step, we reconfirmed that

serum derived cfChPs are internalized by PBMCs. We labelled the serum

derived cfChPs in their DNA and histones with Pt Bright 550 (red) and ATTO-488

(green; see Materials and Methods), respectively. When the dually labelled

cfChPs (10 ng) were added to fresh PBMCs, co-localizing red and green

fluorescence signals representing cfChPs were clearly detected in nuclei of fifty

percent of the treated PBMCs by 2 h (Suppl. Fig. 3).

Immunogenic effects of serum derived cfChPs

To investigate whether serum derived cfChPs, like those released from dying

cells, would trigger immune response, we treated PBMCs with serum derived

cfChPs (10 ng) and evaluated their immunogenic potential. The treatment of

PBMCs with cfChPs triggered a marked increase in the expression of CD69 on

T-cells and NK-cells at 24 h (Fig. 2A). A pronounced increase in levels of the

pro-inflammatory cytokines IL-1β, IL-6, TNF-α, IFN-γ, CXCL9, and IFN-β, as well

as the two anti-inflammatory cytokines IL-10 and TGF-β1 (Fig. 2B), was also

detected. cfChPs also activated mRNA expression NF-KB, c-Jun, c-Fos, Jun-B,

FosB, and EGR-1 on human PBMCs, but at different time points **(Suppl. Fig. 4).**

Of these, upregulation of four markers namely, c-Jun, c-Fos, Jun-B and FosB were also found to be upregulated at protein level when analyzed by flow cytometry at the peak time point of mRNA expression **(Fig. 2C)**.

Serum derived cfChPs act via the cGAS-STING pathway

Since the cGAS-STING pathway is activated when DNA is atypically present in the cytoplasm (Motwani et al., 2019)**,** we hypothesized that cGAS-STING might be activated when cfChPs appear in the cytoplasm. We investigated this indirectly by treating PBMCs with cfChPs in the presence of the STING protein inhibitor H151 and demonstrate a significantly reduced production of inflammatory cytokines as estimated by flow cytometry and ELISA **(Fig. 3A)**. qRT-PCR performed 12h after treatment of PBMCs with serum derived cfChPs (10 ng) provided direct evidence of the involvement of cGAS-STING pathway **(Fig. 3B)**. Further direct evidence came from our demonstration that cfChPs treatment significantly upregulated phosphorylated STING protein (p-STING) expression in lymphocytes at 24 h as determined by flow cytometry **(Fig. 3C)**. We further show using immunofluorescence that uptake of cfChPs (labelled in their histone with ATTO-488) markedly increased cGAS expression in PBMCs by 4h **(Fig. 3D)**. cfChPs were labelled only in their histones with ATTO-488 (green) to prevent spectral overlap with cGAS staining. Interestingly, in response to the internalized cfChPs, cGAS was found to shuttle between the cytoplasm and nucleus. At 2h of cfChPs treatment, cGAS translocated from the nucleus to the

cytoplasm, while at 4 h it had translocated back to the nucleus, indicating that

cGAS is directly involved in cfChPs induced immune response. These findings

suggest that cfChPs activate immune response via the cGAS-STING pathway.

Discussion

Several notable observations were made in this study. First, cfChPs released

from dying PBMCs, as well as serum derived cfChPs, rapidly enter immune cells

and accumulate within their nuclei. Second, cfChPs from both sources activate

CD69 expression on T cells and NK cells; cellular stress-related pathways and

inflammatory cytokines. Third, cfChPs-induced immune response is mediated via

the cGAS-STING pathway.

It is currently believed that cell death results in the release of DAMPs, which are

responsible for the activation of an inflammatory immune response (Denning et

al., 2019; Land, 2015; Rock et al., 2011); however, the critical component(s) of

DAMPs that is responsible for mounting immune response continues to remain

elusive (Fleischmann et al., 2016; Schaefer, 2014; Ye et al., 2020). The

composition of DAMPs is complex as they comprise a wide range of molecules

derived from the extracellular matrix as well as nuclear components of dying cells

(Schaefer, 2014)**.** Our results identify that cfChPs might be a critical component

of DAMPs that activate immune response following their internalization by

lymphocytes. Using indirect and direct approaches, we have further

demonstrated that the cfChPs-induced immune response may be mediated via

the cGAS-STING pathway. For the former, we utilized a well-researched

pharmacological inhibitor of STING, H151 (Al-Asmari et al., 2022; Gong et al., 2021; Hu et al., 2022, 2023; Varga et al., 2023), which prevents palmitoylation of STING and thus blocks STING signaling pathway (Haag et al., 2018). We found that H151 could abrogate cfChPs induced inflammatory cytokine release. Since the cGAS-STING pathway is activated when nucleic acids are atypically present in the cytoplasm (Motwani et al., 2019), we suspected that cfChPs activated cGAS-STING pathway while present in cytoplasm. Direct confirmation of involvement of cGAS-STING pathway came from flow cytometry and fluorescent confocal microscopy experiments. We show by flow cytometry that cfChPs treatment upregulates the phosphorylated STING protein. Experiments using fluorescent confocal microscopy demonstrated upregulation of cGAS expression in the nucleus of cfChPs treated PBMCs which appeared to shuttle back and forth between the nucleus and the cytoplasm (Herzner et al., 2021; Liu et al., 2018; Song et al., 2022). These findings suggest that cfChPs-induced immune response activation is mediated via the cGAS-STING pathway. However, further studies are required to evaluate the involvement of the cGAS-STING pathway in the cfChPs-induced immune response.

We have shown, in several studies reported elsewhere, that the various pathological activities of cfChPs can be abrogated by agents that deactivate cfChPs (Agarwal et al., 2022; Kirolikar et al., 2018; I. Mittra et al., 2017; Mittra Id et al., 2020; Ostwal et al., 2022; Pal et al., 2022; Pilankar et al., 2022). Of these, the pro-oxidant combination of the commonly used nutraceuticals resveratrol and copper (R-Cu) holds the most promise for clinical application because of its lack

of toxicity and low cost. We have reported that R-Cu can ameliorate sepsis in a pre-clinical model (Mittra Id et al., 2020) and prevent cytokine storm related death in severe coronavirus disease 2019 in humans (Mittra et al., 2020). For this reason, R-Cu may be a worthy candidate for evaluation in clinical trials involving life-threatening disorders associated with hyper-inflammation, which can include sepsis, cancer, neurodegenerative disorders, and auto-inflammatory as well as autoimmune diseases.

Materials and Methods

Institutional ethics approval

This study was approved by the Institutional Ethics Committee (IEC) of Advanced Centre for Treatment, Research and Education in Cancer (ACTREC), Tata Memorial Centre (TMC), Mumbai (Approval no. 900520). Peripheral blood samples (10 mL) were collected from healthy volunteers after obtaining informed consent as required by the IEC.

Collection of blood samples

Blood samples were used for isolation of PBMCs and cfChPs. To isolate PBMCs, blood samples (10 mL) were collected in heparinized Vacutainer TM tubes (Becton-Dickinson Vacutainer Systems, Franklin Lakes, NJ, USA), whereas to isolate cfChPs, samples were collected in plain Vacutainer tubes (Becton-Dickinson Vacutainer Systems). The volunteers had no history of febrile or other illnesses in the preceding 3 months.

Isolation of PBMCs

PBMCs were isolated using Ficoll-Hypaque according to standard procedures.

Procedure for collecting conditioned medium (CM) from hypoxia-induced

dying PBMCs

ThinCert® Cell Culture Inserts, containing a separating membrane of pore size

400 nm, are designed for dual chamber experiments (Greiner Bio-One,

Kremsmünster, Austria). To collect CM containing cfChPs released from dying

cells, PBMCs (1 x 10^6) were seeded in ThinCert® Cell Culture Inserts containing

1.5 mL Dulbecco's Modified Eagle Medium (DMEM) and placed in a 35-mm 6-

well plate. The plate was incubated in a hypoxia chamber (1% O_2) for 72 h. Post

72 h, sufficient DMEM (~700 μL) was added to the lower chamber such that the

level of media touched the base of the insert placed above. The plates were

incubated at 37°C in a humidified atmosphere of 5% CO_2 for 48 h under

normoxic conditions. Media in the lower chamber allowed cfChPs <400 nm in

size, released from hypoxia-induced dying PBMCs, to seep into the medium in

the lower chamber. Post-incubation, conditioned media from the wells was

pooled and was used for further experiments.

In some experiments, PBMCs were dually fluorescently labelled in their DNA and

histones such that the collected CMs contained dually fluorescently labelled

cfChPs (see below).

Collection of CM containing dually fluorescently labelled cfChPs

CM containing dually fluorescently labelled cfChPs was used in experiments to

detect the uptake of fluorescently dually labelled cfChPs by recipient PBMCs.

PBMCs were fluorescently dually labelled in their DNA with bromodeoxyuridine

(10 µM; Sigma-Aldrich, St. Louis, MO, USA), and in their histones (H2B) with

CellLight® Histone 2BGFP (Thermo Fisher Scientific, Waltham, MA, USA) for 36

h. The detailed labelling procedure has been described by us previously

(Kirolikar et al., 2018). The dually labelled PBMCs were induced to undergo

hypoxic cell death, as described above, such that the collected CM contained

fluorescently dually labelled cfChPs (<400 nm). Two hundred micro-liters of

conditioned media containing cfChPs <400nm that had seeped into the lower

chamber of the insert was applied to isolated PBMCs in a time course

experiment (2h, 4h and 6h). Cells were then washed and processed for

fluorescence microscopy to detect presence of fluorescent signals of BrdU and

histone H2BGFP in the recipient PBMCs.

Treatment of PBMCs with CM containing cfChPs

CM collected from hypoxia-induced dying (unlabeled) PBMCs containing cfChPs

were used in two experimental settings. In the first setting, a time course analysis

was performed wherein PBMCs were treated with varying volumes of CMs (25,

50, 100, 200, 300, 400, and 500 µL) for estimation of CD69 expression (see

below). Optimal CD69 expression was obtained using 200 µL CM. In the second

experiment, PBMCs were treated with 200 µL CM to facilitate the estimation of

stress markers and inflammatory cytokines (see below).

Isolation of cfChPs from human sera

cfChPs were isolated from sera of healthy volunteers according to a protocol

described by us previously (Mittra et al., 2015). Pooled serum (typically from ~5

individuals) was used to isolate cfChPs to maintain inter-experimental

consistency. Isolated cfChPs were quantified in terms of their DNA content using

a Pico-Green quantification assay.

Fluorescent dual labelling of cfChPs

Fluorescently dual-labelled cfChPs were utilized to detect the uptake of

fluorescently dually labelled cfChPs by recipient PBMCs. cfChPs were dually

fluorescently labelled in their DNA and histone H4 with Platinum Bright 550 (red)

and ATTO-TEC 488 (green), respectively, according to the protocol described by

us previously (Mittra et al., 2015).

Treatment of PBMCs with serum derived cfChPs

PBMCs were treated with cfChPs and were analyzed for CD69 expression,

cellular stress markers, and inflammatory cytokine levels. PBMCs (5×10^5) were

plated in 24-well plates containing 1 mL DMEM. Following overnight incubation,

cells were treated with serum derived cfChPs (10 ng equivalent of DNA), with

and without H151 for 24 h in a humidified atmosphere of 5% CO_2. Untreated

PBMCs were utilized as negative controls.

qRT-PCR

qRT-PCR was used for the evaluation of stress markers, STING and cGAS after

the treatment of PBMCs with 1) CM from hypoxia-induced dying PBMCs (200

μL), or 2) serum derived cfChPs (10 ng). The time points for both analyses were:

0, 6, 12, 24, 36, 48, and 72 h and appropriate untreated control PBMCs were

analyzed in parallel. Total RNA was isolated using RNeasy Mini Kit (Qiagen,

Hilden, Germany), and approximately 1 μg of isolated RNA was converted to

cDNA using a RT^2 First Strand Kit (Qiagen, Hilden, Germany). qRT-PCR was

performed using a SYBR Select Master Mix (Applied Biosystems, Waltham, MA,

USA). All samples were assayed on a QuantStudio5 system (Applied

Biosystems) in duplicates. Data were analyzed using a comparative C_T method,

and fold change in mRNA expression was calculated as $2^{\wedge}(-\Delta\Delta C_T)$.

Flow cytometry

Flow cytometry was utilized for the evaluation of 1) CD69 expression, 2)

inflammatory cytokine levels, and 3) stress-related markers and p-STING

following the treatment of PBMCs with either CM (200 μL) or serum derived

cfChPs (10 ng).

CD69 expression

After 24 h incubation with CM or cfChPs, PBMCs were labelled for 20 min with

anti-CD45-BUV395 (HI30), anti-CD3-BV711 (UCHT1), anti-CD56-PE-Cy7

(NCAM) and anti-CD69-APC (FN50) (Becton Dickinson Biosciences, San Jose,

CA, USA.) in dark at room temperature. Samples were analyzed on a FACSAria

III cytometer (Becton Dickinson).

Estimation of cytokine levels

After 48 h treatment either with CM or with cfChPs, cultured cells were

centrifuged at 3,000 rpm for 10 min at 4°C and supernatants were collected and

stored at -80°C for cytokine analysis. A LEGENDplex™ Multi-Analyte Flow assay

kit (BioLegend, San Diego, CA, USA) or ELISA was used as per the

manufacturer's instructions to quantify levels of secreted cytokines in culture

supernatants.

Stress-related markers and pSTING expression

After incubation with cfChPs at the peak time-point as detected by qRT-PCR,

PBMCs were fixed and permeabilized with either 4% formaldehyde and 90%

methanol or cytofix/cytoperm kit (Becton Dickinson) based on the manufacturer's

instructions and were then stained using specific primary antibodies against

respective proteins. For unconjugated antibodies, cells were stained with

appropriate secondary antibody. Cells were acquired on Attune NxT flow

cytometer (ThermoFisher Scientific, USA).

Flow cytometry acquisition

To determine CD69 expression, lymphocytes were gated on forward and side

scatter parameters and further as CD45$^+$ cells **(Suppl. Fig. 5).** T-cells were gated

as CD3$^+$ cells and NK-cells as CD56$^+$CD3$^-$ cells. CD69 expression on the cells

was determined as percentage of positive cells in the respective gated cell

regions. The threshold between negative and positive was defined by the

fluorescence minus one method. At least 10,000 lymphocytes were acquired on

a FACSAria III cytometer (Becton Dickinson). To determine stress markers and

pSTING expression, lymphocytes were gated on forward and side scatter

parameters and specific marker expression on the cells was determined by

increase in Mean fluorescence intensity (MFI) compared to only secondary

antibody staining or only cells (whichever applicable). At least 5,000 lymphocytes

were acquired on Attune NxT flow cytometer (ThermoFisher Scientific, USA). For

the estimation of cytokine levels, sample acquisition was performed as per the

manufacturer's instructions. At least 4,000 cytometric beads were acquired on a

FACSAria III cytometer (Becton Dickinson).

Flow cytometry analysis

Data was analyzed using FlowJo software (Version 10; FlowJo LLC, Ashland,

OR, USA) and cytokine data was analyzed using LEGENDplex Data Analysis

Software (BioLegend).

Enzyme-linked immunosorbent assay (ELISA)

ELISA was used to estimate IL-1β, IFN-β, and CXCL9 levels in the supernatant

media (Krishgen Biosystems, Mumbai, India) as per the manufacturer's

instructions.

Fluorescent confocal microscopy

For co-staining of cfChPs and cGAS protein, cfChPs were labelled only in their

histones with ATTO-488 (green) to prevent spectral overlap with cGAS staining.

PBMCs were treated with labelled cfChPs and incubated for 2 h and 4 h at $37^{o}C$

in a humidified atmosphere of 5% CO_2. Cells were cytospun onto a slide,

immediately fixed with 4% PFA and immuno-stained using rabbit anti-cGAS

primary antibody and appropriate secondary antibody by indirect

immunofluorescence method as described by us earlier (Indraneel Mittra et al.,

2017). Slides were mounted in VectaShield DAPI (Vector labs, USA). For

confocal microscopy, images were acquired using a 63x oil objective on Nikon

AX Confocal Microscope System, (Nikon Corporation, Japan). All experiments

were performed in duplicate, and 50 cells were analysed in each case. Mean

fluorescence intensity (MFI) of images was measured using ImageJ FIJI software

1.5t version (W. Rasband, Bethesda, USA) and results were expressed as mean

± SEM.

Statistical analysis

Statistical analysis was performed using two-tailed unpaired student's t-tests

using GraphPad Prism 8 (GraphPad Software, Boston, MA, USA) and the results

are expressed as mean ± SEM. The following significance thresholds were

applied: * $p<0.05$, ** $p<0.01$, *** $p<0.005$, and **** $p<0.0001$.

357
358

359 **Conflict of Interest:** Authors declare that they have no competing interests.

360 **Data Availability:** All data supporting the findings of this study are available

361 within the article and supplementary materials.

Acknowledgments: The authors thank personnel in the flow cytometry facility at

ACTREC-TMC for their technical support. The authors thank Dr. Rajan Basak for

his guidance in qRT-PCR.

Funding: Department of Atomic Energy, Government of India grant CTCTMC

(IM).

The funder had no role in the preparation, review, or approval of the manuscript

and decision to submit the manuscript for publication.

References

Agarwal A, Khandelwal A, Pal K, Khare NK, Jadhav V, Gurjar M, Punatar S,

Gokarn A, Bonda A, Nayak L, Kannan S, Gota V, Khattry N, Mittra I. 2022. A

novel pro-oxidant combination of resveratrol and copper reduces transplant

related toxicities in patients receiving high dose melphalan for multiple

myeloma (RESCU 001). *PLoS One* **17**:e0262212.

doi:10.1371/journal.pone.0262212

Akira S, Uematsu S, Takeuchi O. 2006. Pathogen recognition and innate

immunity. *Cell.* doi:10.1016/j.cell.2006.02.015

Al-Asmari SS, Rajapakse A, Ullah TR, Pépin G, Croft L V., Gantier MP. 2022.

Pharmacological Targeting of STING-Dependent IL-6 Production in Cancer

Cells. *Front Cell Dev Biol* **9**. doi:10.3389/fcell.2021.709618

Ashida H, Mimuro H, Ogawa M, Kobayashi T, Sanada T, Kim M, Sasakawa C.

2011. Cell death and infection: A double-edged sword for host and pathogen

384  survival. *J Cell Biol.* doi:10.1083/jcb.201108081

385 Chaplin DD. 2010. Overview of the immune response. *J Allergy Clin Immunol*

386  **125**:S3. doi:10.1016/j.jaci.2009.12.980

387 Denning NL, Aziz M, Gurien SD, Wang P. 2019. Damps and nets in sepsis. *Front*

388  *Immunol.* doi:10.3389/fimmu.2019.02536

389 Fleischmann C, Scherag A, Adhikari NKJ, Hartog CS, Tsaganos T, Schlattmann

390  P, Angus DC, Reinhart K. 2016. Assessment of global incidence and

391  mortality of hospital-treated sepsis current estimates and limitations. *Am J*

392  *Respir Crit Care Med* **193**:259–272. doi:10.1164/rccm.201504-0781OC

393 Gong W, Lu L, Zhou Y, Liu J, Ma H, Fu L, Huang S, Zhang Y, Zhang A, Jia Z.

394  2021. The novel STING antagonist H151 ameliorates cisplatin-induced acute

395  kidney injury and mitochondrial dysfunction. *Am J Physiol Physiol* **320**:F608–

396  F616. doi:10.1152/ajprenal.00554.2020

397 Haag SM, Gulen MF, Reymond L, Gibelin A, Abrami L, Decout A, Heymann M,

398  van der Goot FG, Turcatti G, Behrendt R, Ablasser A. 2018. Targeting

399  STING with covalent small-molecule inhibitors. *Nature* **559**:269–273.

400  doi:10.1038/s41586-018-0287-8

401 Herzner A-M, Schlee M, Bartok E. 2021. The many faces of cGAS: how cGAS

402  activation is controlled in the cytosol, the nucleus, and during mitosis. *Signal*

403  *Transduct Target Ther* **6**:260. doi:10.1038/s41392-021-00684-3

Hu S, Gao Y, Gao R, Wang Y, Qu Y, Yang J, Wei X, Zhang F, Ge J. 2022. The selective STING inhibitor H-151 preserves myocardial function and ameliorates cardiac fibrosis in murine myocardial infarction. *Int Immunopharmacol* **107**:108658. doi:10.1016/j.intimp.2022.108658

Hu Z, Zhang F, Brenner M, Jacob A, Wang P. 2023. The protective effect of H151, a novel STING inhibitor, in renal ischemia-reperfusion-induced acute kidney injury. *Am J Physiol Physiol* **324**:F558–F567. doi:10.1152/ajprenal.00004.2023

Kirolikar S, Prasannan P, Raghuram G V., Pancholi N, Saha T, Tidke P, Chaudhari P, Shaikh A, Rane B, Pandey R, Wani H, Khare NK, Siddiqui S, D'souza J, Prasad R, Shinde S, Parab S, Nair NK, Pal K, Mittra I. 2018. Prevention of radiation-induced bystander effects by agents that inactivate cell-free chromatin released from irradiated dying cells. *Cell Death Dis* **9**:1–16. doi:10.1038/s41419-018-1181-x

Land WG. 2015. The role of damage-associated molecular patterns in human diseases: Part I - Promoting inflammation and immunity. *Sultan Qaboos Univ Med J*.

Liu H, Zhang H, Wu X, Ma D, Wu J, Wang L, Jiang Y, Fei Y, Zhu C, Tan R, Jungblut P, Pei G, Dorhoi A, Yan Q, Zhang F, Zheng R, Liu S, Liang H, Liu Z, Yang H, Chen J, Wang P, Tang T, Peng W, Hu Z, Xu Z, Huang X, Wang J, Li H, Zhou Y, Liu F, Yan D, Kaufmann SHE, Chen C, Mao Z, Ge B. 2018.

Nuclear cGAS suppresses DNA repair and promotes tumorigenesis. *Nature*

563:131–136. doi:10.1038/s41586-018-0629-6

Mittra I, de Souza R, Bhadade R, Madke T, Shankpal PD, Joshi M, Qayyumi B,

Bhattacharya A, Gota V, Gupta S, Chaturvedi P, Badwe R. 2020.

Resveratrol and Copper for treatment of severe COVID-19: an observational

study (RESCU 002) (preprint). *medRxiv* 2020.07.21.20151423.

Mittra I, Khare NK, Raghuram GV, Chaubal R, Khambatti F, Gupta D, Gaikwad

A, Prasannan P, Singh Akshita, Iyer A, Singh Ankita, Upadhyay P, Nair NK,

Mishra PK, Dutt A. 2015. Circulating nucleic acids damage DNA of healthy

cells by integrating into their genomes. *J Biosci* **40**:91–111.

doi:10.1007/s12038-015-9508-6

Mittra I., Pal K, Pancholi N, Shaikh A, Rane B, Tidke P, Kirolikar S, Khare NK,

Agrawal K, Nagare H, Nair NK. 2017. Prevention of chemotherapy toxicity by

agents that neutralize or degrade cell-free chromatin. *Ann Oncol* **28**:2119–

2127. doi:10.1093/annonc/mdx318

Mittra Indraneel, Samant U, Sharma S, Raghuram G V, Saha T, Tidke P,

Pancholi N, Gupta D, Prasannan P, Gaikwad A, Gardi N, Chaubal R,

Upadhyay P, Pal K, Rane B, Shaikh A, Salunkhe S, Dutt S, Mishra PK,

Khare NK, Nair NK, Dutt A. 2017. Cell-free chromatin from dying cancer

cells integrate into genomes of bystander healthy cells to induce DNA

damage and inflammation. *Cell Death Discov* **3**:17015.

doi:10.1038/cddiscovery.2017.15

Mittra Id I, Pal K, Pancholi N, Tidke P, Siddiqui S, Rane B, D'souza J, Shaikh A,

Parab S, Shinde S, Jadhav V, Shende S, Raghuram G V. 2020. Cell-free

chromatin particles released from dying host cells are global instigators of

endotoxin sepsis in mice. *PLoS One* **15(3)**:e0229017.

doi:10.1371/journal.pone.0229017

Motwani M, Pesiridis S, Fitzgerald KA. 2019. DNA sensing by the cGAS–STING

pathway in health and disease. *Nat Rev Genet* **20**:657–674.

doi:10.1038/s41576-019-0151-1

Ostwal V, Ramaswamy A, Bhargava P, Srinivas S, Mandavkar S, Chaugule D,

Peelay Z, Baheti A, Tandel H, Jadhav VK, Shinde S, Jadhav S, Gota V,

Mittra I. 2022. A pro-oxidant combination of resveratrol and copper reduces

chemotherapy-related non-haematological toxicities in advanced gastric

cancer: results of a prospective open label phase II single-arm

study (RESCU III study). *Med Oncol* **40**:17. doi:10.1007/s12032-022-01862-

1

Pal K, Raghuram G V., Dsouza J, Shinde S, Jadhav V, Shaikh A, Rane B,

Tandel H, Kondhalkar D, Chaudhary S, Mittra I. 2022. A pro-oxidant

combination of resveratrol and copper down-regulates multiple biological

hallmarks of ageing and neurodegeneration in mice. *Sci Rep* **12**:17209.

doi:10.1038/s41598-022-21388-w

Pilankar A, Singhavi H, Raghuram G V., Siddiqui S, Khare NK, Jadhav V, Tandel

H, Pal K, Bhattacharjee A, Chaturvedi P, Mittra I. 2022. A pro-oxidant

combination of resveratrol and copper down-regulates hallmarks of cancer

and immune checkpoints in patients with advanced oral cancer: Results of

an exploratory study (RESCU 004). *Front Oncol* **12**.

doi:10.3389/fonc.2022.1000957

Rock KL, Lai JJ, Kono H. 2011. Innate and adaptive immune responses to cell

death. *Immunol Rev* **243**:191–205. doi:10.1111/j.1600-065X.2011.01040.x

Schaefer L. 2014. Complexity of danger: The diverse nature of damage-

associated molecular patterns. *J Biol Chem*. doi:10.1074/jbc.R114.619304

Shabrish S, Mittra I. 2021. Cytokine Storm as a Cellular Response to dsDNA

Breaks: A New Proposal. *Front Immunol* **12**.

doi:10.3389/fimmu.2021.622738

Song J-X, Villagomes D, Zhao H, Zhu M. 2022. cGAS in nucleus: The link

between immune response and DNA damage repair. *Front Immunol* **13**.

doi:10.3389/fimmu.2022.1076784

Varga KZ, Gyurina K, Radványi Á, Pál T, Sasi-Szabó L, Yu H, Felszeghy E,

Szabó T, Röszer T. 2023. Stimulator of Interferon Genes (STING) Triggers

Adipocyte Autophagy. *Cells* **12**:2345. doi:10.3390/cells12192345

Wang H, Shan X, Ren M, Shang M, Zhou C. 2021. Nucleosomes enter cells by

clathrin- and caveolin-dependent endocytosis. *Nucleic Acids Res* **49**:12306–

12319. doi:10.1093/nar/gkab1121

Ye Q, Wang B, Mao J. 2020. The pathogenesis and treatment of the 'Cytokine

Storm'' in COVID-19.' *J Infect.* doi:10.1016/j.jinf.2020.03.037

Legends to Figures

Fig. 1. cfChPs released from dying PBMCs into conditioned medium (CM) induce immune response. (A) Representative flow cytometry plots of CM-treated PBMCs. Line graphs represent the expression of CD69 in CM-treated PBMCs through dose-response analyses (left hand panel). Representative flow cytometry plots (middle panel) and box-plot (right hand panel) represent the relative upregulation of CD69 expression on T-cells and NK-cells using 200 µL CM. **(B)** Box-plot represent the relative production of inflammatory cytokines by PBMCs in response to CM (200 µl) as estimated by flow cytometry and ELISA. It is to be noted that CM itself does not contain cytokines. **(C)** Upregulation of stress related marker expression in PBMCs treated with CM. Samples were analyzed by flow cytometry and respective protein expression was analyzed on untreated (blue line) and CM treated (red line) PBMCs. Box-plot represents quantitative analysis (MFI) of stress related markers by PBMCs in response to CM. Box-plot represent median with minimum and maximum data values. Statistical analyses were performed using two-tailed unpaired student's *t*-tests in GraphPad Prism 8. * p<0.05, ** p<0.01, *** p<0.005.

Fig. 2. Serum derived cfChPs activate immune response. (A) Representative flow cytometry plots of cfChPs (10 ng)-treated PBMCs. Line graphs represent the expression of CD69 in a time course experiment (blue line represents untreated cells; red line represents cfChP-treated cells) (left hand panel). Representative flow cytometry plots (middle panel) and box-plot (right hand panel) represent the relative upregulation of CD69 expression on T-cells and NK-cells in response to cfChPs at the peak time-point (24 h). **(B)** Box-plots representing the production of cytokines by PBMCs in response to cfChPs (10 ng). **(C)** Upregulation of stress related marker expression in PBMCs treated with cfChPs. Samples were analyzed by flow cytometry and respective protein expression was analyzed on untreated (blue line) and cfChPs treated (red line) PBMCs (upper panel). Box-plots in the lower panel represent quantitative analysis (MFI) of stress related markers by PBMCs in response to cfChPs (10 ng). Box-plot represent median

with minimum and maximum data values and the data were analyzed using two-
tailed unpaired student's *t*-tests in GraphPad Prism 8. * p<0.05, ** p<0.01, ***
p<0.005, **** p<0.0001.

Fig. 3. Serum derived cfChPs activate immunogenic effects *via* the cGAS-
STING pathway. (A) Box-plots represent data of inflammatory cytokines
secreted by PBMCs in response to cfChPs (10 ng) and their inhibition by prior
treatment with the STING protein inhibitor H151 (50 ng/ml). It is to be noted that
treatment with H151 itself had no effects. **(B)** Upregulation of mRNA expression
of cGAS and STING in human PBMCs following treatment with serum derived
cfChPs (10 ng) detected by qRT-PCR and analyzed using a comparative C_T
method. Line graphs represent the results of a time course experiment calculated
as expression fold [$2^{(-\Delta\Delta C_T)}$]. **(C)** Upregulation of p-STING expression in
PBMCs following treatment with serum derived cfChPs (10 ng) detected by flow
cytometry (blue line represents untreated cells; red line represents cfChPs-
treated cells). Box-plot represent the relative upregulation of p-STING on PBMCs
in response to cfChPs treatment. **(D)** Representative fluorescent confocal
microphotographs showing activation of c-GAS protein in PBMCs in response to
cfChPs treatment. Internalization by PBMCs of cfChPs labelled in their histones
(ATTO-488 green), triggered upregulation of c-GAS (red) protein. Quantitative
analysis (MFI) reveal visible elevation of cGAS expression in the cfChPs- treated
cells by 2 h and 4 h compared to untreated control cells. Results of quantitative
analyses of overall cGAS activation at 2 h and 4 h and the relative presence of
cGAS in the nucleus and the cytoplasm are presented as box-plots (upper and
lower respectively). Results are represented as box-plot indicating median with
minimum and maximum data values and the data were analyzed using two-tailed
unpaired student's *t*-tests in GraphPad Prism 8. * p<0.05, ** p<0.01.

Supplementary Files

Suppl. Fig. 1: Accumulation of cfChPs released from dying PBMCs in the nuclei of healthy PBMCs. PBMCs were dually labelled in their DNA and histone with bromodeoxyuridine (red) and Histone2B-GFP (green), respectively, and induced to undergo hypoxic cell death. The filtered CM (200 µL) containing dually labelled cfChPs (<400 nm) were applied to fresh PBMCs as described in the Materials and Methods section. The upper panel represents the images of dually labelled donor PBMCs; the lower panel represents treated recipient PBMCs. Fluorescent microscopy images were taken at 4 h.

Suppl. Fig. 2: Upregulation of stress-related markers in human PBMCs following treatment with CM (200µl). mRNA expression of six stress-related markers in PBMCs was detected by qRT-PCR and analyzed using a comparative C_T method. Line graphs represent the results of time course analyses calculated as expression fold [$2^{\wedge}(-\Delta\Delta C_T)$].

Suppl. Fig. 3: Accumulation of cfChPs in the nuclei of PBMCs. Serum derived cfChPs were dually labelled in their DNA and histones with Platinum Bright 550 (red) and ATTO-TEC-488 (green), respectively. PBMCs were treated with the dually labelled cfChPs (10 ng) and fluorescent microscopy images were taken at 2 h.

Suppl. Fig. 4: Upregulation of stress-related markers in human PBMCs following treatment with cfChPs (10 ng). mRNA expression of six stress-related markers in PBMCs was detected by qRT-PCR and analyzed using a comparative C_T method. Line graphs represent the results of time course analyses calculated as expression fold [$2^{\wedge}(-\Delta\Delta C_T)$].

Suppl. Fig. 5: Flow cytometry plots representing gating strategy. Gating of lymphocytes on forward/side scatter and further as CD45+ cells, T-cells were identified as CD3+ cells and NK cells as CD56+/CD3- cells.

Figure 1

Figure 2

Figure 3