The Moa the Merrier: Resolving When the Dinornithiformes Went Extinct

Floe Foxon[1],

[1] Folk Zoology Society – PO Box 97014, Pittsburgh, PA 15229, USA

Correspondence: floefoxon@protonmail.com

Abstract

The Moa (Aves: Dinornithiformes) are an extinct group of the ratite clade from New Zealand. The overkill hypothesis asserts that the first New Zealand settlers hunted the Moa to extinction by 1450 CE, whereas the staggered survival hypothesis allows for Moa survival until after Europeans began to arrive on New Zealand. Alleged Moa sightings post-1450 CE may shed light on these competing hypotheses. A dataset of 97 alleged Moa sightings from circa 1675 CE to 1993 CE was constructed, with sightings given subjective quality ratings corresponding to various statistical probabilities. Cumulative probabilities of Moa persistence were calculated with a conservative survival model using these probabilistic sighting-records; a method recently applied to sightings of the Thylacine. Cumulative persistence probability fell sharply after 1390 CE, and across pessimistic and optimistic variations of the model, it was more likely than not that the Moa were extinct by 1770 CE. Probabilistic sighting-record models favour the overkill hypothesis, but give probabilities of Moa survival post-1450 CE that cannot be dismissed entirely. Eyewitness data on Moa sightings are amenable to scientific study, and these methods may be applied to similar animals.

Keywords: Moa; Dinornithiformes; Extinction Dynamics; Citizen Science; Overkill hypothesis; Folk Zoology; Hidden Animals

Introduction

In a recent study in *Science of the Total Environment*, Brook et al. (2023) used a database of alleged Thylacine sighting records from 1910 CE to 2023 CE in a computational extinction dynamics estimator (EDE) to estimate probabilities of persistence of the predatory marsupial Thylacine, *Thylacinus cynocephalus*. Brook et al. (2023) bring a much-needed level of nuance, scientific rigour, and statistical methodology to bear on a topic that is often shrouded in pseudoscience and incredulity; by assigning probabilities to individual sightings rather treating these in binary, Brook et al. transcend the traditional paradigm in zoology of either blind belief in, or arrogant denial of, the existence of hidden animals.

The methods presented by Brook et al. (2023) can be applied to other species whose extinction dates are contested. Roughly 2,000 kilometers east of the Thylacine's native Tasmania lies New Zealand, a land dominated by avifauna (Worthy et al., 2017). The greatest among the New Zealand avifauna were the Dinornithiformes, or Moa: large flightless birds of the ratite clade (Canoville et al., 2022), one species of which, *Dinornis robustus*, reached heights of up to 3.6 meters (Widrig and Field, 2022). Moa lived in relative peace on New Zealand as megaherbivorous browsers (Widrig and Field, 2022), preyed upon only by Haast's eagle, *Harpagornis moorei* (Scofield and Ashwell, 2009), until the arrival of the Austronesian peoples circa 1250–1300 CE (Irwin and Walrond, 2016).

The latest and now generally-accepted view is that, as part of the greater Quaternary 'overkill' (Sandom et al., 2014), an extremely low-density population of first New Zealand settlers wiped out the entire population of Moa on the islands in a period of less than 150 years (Holdaway et al., 2014). This rapid megafaunal extinction is evidenced by high-quality radiocarbon dates on Moa remains from natural and archaeological sites (Perry et al., 2014). Indeed, it is thought that the Moa had relatively low reproduction rates and older ages of sexual maturity (McCulloch, 1982) making them vulnerable to extinction via predation, and a recent collaborative analysis of ancestral sayings suggests that early Maori used the Moa as a metaphor for extinction, e.g. 'lost as the Moa is lost', similar to the English phrase 'dead as a dodo' (Wehi et al., 2018).

The overkill hypothesis is not a view held by all; in particular, critics point to the marked variability in terrain across New Zealand, and question the ability of settlers to eradicate all Moa species across a vast, difficult, and (then) unknown land at such speed (Diamond, 2000). Rather than a rapid war against the Moa, critics opt for 'staggered survival', perhaps until circa 1800 CE (Silverberg, 1973, Chapter 7), by which time Europeans had begun to arrive on the islands.

Entertaining the possibility of Moa persistence post-1450 CE, when did the Moa go extinct? Like the Thylacine studied by Brook et al. (2023), alleged Moa sightings have persisted to as recently as 1993 CE, with reports originating from both Maori and European peoples (Silverberg, 1973, Chapter 7). Indeed, zoologist Bernard Heuvelmans (1986) lists "A surviving species of the Moa family... known to some as *roa-roa*" in his 'Annotated Checklist of Apparently Unknown Animals,' and biologist Roy P. Mackal (1983) dedicated an entire chapter to the possible survival of the New Zealand Moa, among other birds, in his intriguing though not entirely skeptical book 'Searching for Hidden Animals.'

The aim of the present study was to extend the methods of Brook et al. (2023) to the Moa as a test of the 'staggered survival' hypothesis, i.e. that the Moa persisted after 1450 CE, by which time the competing 'overkill' hypothesis asserts that the Moa were extinct.

Material and methods

A new dataset named the New-Zealand Moa Sighting Records Database (NMSRD) was constructed following the Tasmanian Thylacine Sighting Records Database (TTSRD) of Brook et al. (2023). Alleged sightings of Moa in New Zealand were taken manually from Spittle (2010) and were coded in a flat-file format (.xlsx) with one observation per row. The columns in this dataset correspond to unique sighting ID; sighting location; year of observation; month of observation; location name; subjective quality-rating (a score between 1 [lowest] to 5 [highest]); number of human observers; number of Moa; reference; observer name(s); and sighting description.

The approach taken was to adopt the most conservative survival model in Brook et al. (2023); the method using only probabilistic sighting-records and no EDE, which resulted in the lowest persistence probabilities in that study. This approach is justified because the sightings in the NMSRD were generally of lower quality than those in the TTSRD. Following Brook et al. (2023), sightings rated 5 were assigned a probability of 1; 4-rated sightings were assigned a probability of 0.05; 3-rated sightings 0.01; 2-rated sightings 0.005; and 1-rated sightings 0.001. The cumulative probabilities of Moa persistence were then calculated as in Brook et al. (2023) for each year.

For clarity and transparency on the subjective rating system, the following rules were generally applied.

1. Sightings that were quite possibly hoaxes, sightings that were almost certainly cases of misidentification, sightings with reference to the informant's 'father' or 'grandfather' (which may be a mistranslation of 'ancestors'), sightings that referred to 'fresh' looking bones (which may have been well-preserved and therefore deceptively old), and especially those with uncertainty in the number and identities of the observers, the date and location of the observation, &c, were generally rated 1 (the lowest rating).

2. Sightings that were less likely to be hoaxes and less clearly misidentification, especially those with known observer identity/date/location/&c, were generally rated 2.

3. Sightings with circumstantial evidence in support (e.g., bones subsequently found in a location where moa were reportedly hunted recently, reported moa behaviour consistent with other struthious birds or reported appearance consistent with museum specimens unknown to the observer, and tracks of uncertain age observed by an expert) were generally rated 3.

4. Sightings involving direct observation by an expert zoologist or ornithologist were rated 4.

5. Radiocarbon dated Moa specimens described by Higham (1994) from archaeological sites across New Zealand (i.e., Moa remains contemporaneous with people in New Zealand) were rated 5. These were included because the model requires starting values for which Moa persistence was definite.

Confirmed hoaxes were necessarily excluded.

In addition to the most pessimistic survival model in Brook et al. (2023) being used, further conservative measures taken were as follows.

• Related sightings (e.g., multiple sightings made by the same individual) were aggregated into single records in the dataset, which had the effect of decreasing the cumulative probability in a given year.

• To test the sensitivity of the model to the subjective probability assigned to each sighting, a 'pessimistic' model assigning a probability of 0.001 to all sightings rated 1 to 4, an 'optimistic' model assigning a probability of 0.005 to all sightings rated 1 and 2, and a 'super optimisitc' model assigning a probability of 0.01 to all sightings rated 1 to 3, were implemented (all other probabilties kept the same as in the main model).

- Where a range of years was given for a sighting (e.g., 1800–1830 CE) the approximate midpoint was taken (e.g., 1815 CE), except where a preferred year was provided in the source text.

- Chapters/sightings in Spittle (2010) in which the persistence of Moa into recent times was not explicitly asserted or implied by the observer, as well as sightings for which no specific date or range of dates were provided or could be reasonably deduced (e.g. in the "dim past" or "olden times"), were all excluded from the dataset.

Analyses were perfomed in Python version 3.8.16 with the packages Numpy 1.21.5, Pandas 1.5.2, and Matplotlib 3.6.2. The NMSRD and analysis code are made freely available in the online supplemental materials at https://doi.org/10.17605/OSF.IO/E59H6.

Results

The NMSRD consists of 97 alleged sighting records involving a total of over 160 eyewitnesses. The NMSRD contains 66 sightings rated 1, 26 sightings rated 2, five sightings rated 3, zero sightings rated 4, and eight radiocarbon dated Moa specimens rated 5. Sighting dates ranged from circa 1675 CE to 1993 CE.

The cumulative probability of persistence of the Moa over time is plotted in Figure 1, where a value of 1.0 implies the Moa were definitely extant at that time, and a value of 0.0 implies the Moa were definitely extinct at that time. Cumulative persistence probability fell sharply after 1390 CE (the last radiocarbon dated Moa specimen in the dataset). For 1770 CE, the cumulative persistence probability was 0.21 in the main model, 0.09 in the pessimistic model, 0.39 in the optimistic model, and 0.61 in the super optimistic model. Therefore, in all but the most optimistic model, the cumulative persistence probability was < 0.5 circa 1770 CE. This may be interpreted as it being more likely than not that the Moa were extinct circa 1770 CE. Results are practically unchanged by removing the earliest three records from 1675 CE to 1725 CE. All models provided cumulative persistence probabilities ≤ 0.17 circa 1900 CE and ≤ 0.01 circa 2000 CE.

Discussion

Conservatively assuming a low but non-zero probability of Moa persistence associated with each alleged Moa sighting post-1450 CE, it is more likely than not that the Moa were extinct before 1770 CE, when Europeans began arriving on New Zealand. This finding favours the overkill hypothesis. Moa sightings in the period post-1450 CE are likely not numerous or reliable enough to support Moa survival into more recent times. However, if one assumes the most optimistic model (generously assigning a probability of 0.01 to all of the sightings), the extinction of the Moa in New Zealand may have been as late as circa 1850 CE. Indeed, there were very few Maori proverbs relating to Moa extinction before 1800 CE (Wehi et al., 2018), which could suggest persistence of some Moa species up to that time.

This work provides for the first time in the published literature an estimate for the extinction date of the Moa on statistical principles using a probabilistic sighting-record model, as in Brook et al. (2023). Proponents of both the staggered survival and overkill hypotheses may be satisfied with the results of this study, the former because the probability of Moa persistence after 1450 CE is non-zero, and the latter because the Moa most probably became extinct rapidly by circa 1450 CE. This finding is more nuanced than previous studies, and demonstrates that the methods of Brook et al. (2023) are applicable to other controversial animals besides the Thylacine.

Figure 1. Cumulative probability of Moa persistence over time, as inferred from alleged eyewitness Moa sightings in New Zealand. Dots represent Moa sightings from Spittle (2010), combined in each year. Red dots represent the 'pessimistic' model, black dots the main model, blue dots the 'optimistic' model, and green dots the 'super optimistic' model (see Methods). Black crosses represent radiocarbon dated Moa specimens from archaeological sites (Higham, 1994). Results are practically unchanged by removing the earliest 'sightings' circa 1675 CE to 1725 CE.

Important limitations must be noted. First is the contentious assumption in assigning subjective non-zero probabilities to all moa sightings. Given the lack of recent physical material (i.e., specimens), it is likely that many of the sightings in the NMSRD constitue innocent cases of mistaken identity, or else hoaxes. Nickell (2017) notes that most Moa 'sightings' occured around the time when fossil Moa were first described by Richard Owen in 1839 CE (Silverberg, 1973, Chapter 7). This may be evidence of 'expectant attention'; the tendency for observers to 'see' what they anticipate they will see (Radford and Nickell, 2006, Chapter 2). Statistical evidence for expectant attention in the context of other 'maybe' animals has been demonstrated by the author in the context of Lake Champlain (Foxon, 2023b). This said, the model was conservative in other ways.

A further limitation is that the NMSRD is not final and is subject to change. The dataset may contain errors that can be ammended as new historical material is brought to the attention of the author. Readers are encouraged to download the data and code in the online supplemental materials and perform their own tests of the subjective ratings/probabilities and adding/removing records as they feel is appropriate. The dataset is also almost certainly incomplete, in part because some 'sightings' will go unreported for fear of ridicule. According to William Scoble who reported his alleged Moa sighting to the Dominion Museum director R. A. Falla, Scoble did not want to be met "with the ridicule metered out to those who talk of sea serpents... when one has met a moa [sic] so close at hand and over so many years reads so many conflicting statements from Science one must be forgiven for reminding you how grey can be all theory" (Scoble, 1952).

Another limitation is that this model applies generally to the order Dinornithiformes, and does not distinguish between individual Moa species. Indeed, there were inhomogeneities across sightings in morphological

characteristics such as size and plumage (see 'Description' column of the NMSRD dataset), which could either suggest different species/sex/ages of the animals being described, or simply noise from misidentification and hoaxes. It should probably be noted for transparency that the author is a skeptic and so approached these sightings *cum grano salis*.

The reliance in the present study and in Brook et al. (2023) on eyewitness testimony (i.e., circumstantial evidence) to infer the existence of an animal is, in a sense, 'folk-zoological', and represents a more mathematically-grounded implementation of the methods described by Heuvelmans (1982, 1984, 1988) in his seminal works. The Thylacine and Moa are animals of known taxa, believed to have become extinct during historical times, and whose existence in the present is unrecognised in conventional zoology. These are described as 'Category IV' animals in the classificatory system of hidden or unknown animals described by ecologist J. Richard Greenwell (1985). Other Category IV animals, such as the ivory-billed woodpecker, *Campephilus principalis*, are amenable to the same probability-based analysis, and future works may investigate that species and similar animals using these methods.

In conclusion, a probabilistic sighting-record model favours the overkill hypothesis of rapid Moa extinction, but gives a probability of Moa survival post-1450 CE that cannot be dismissed entirely. Eyewitness data on Moa sightings are amenable to meaningful scientific analysis as a form of citizen science, similar to previous studies on sightings of animals in folklore (Brook et al., 2023; Foxon, 2023a,b; Paxton, 2009). Regretfully, the Moa are now extinct. As Silverberg (1973, Chapter 7) wrote, "The mighty *Dinornis*... is no more likely to be seen again than Sinbad's rukh." The Moa have had a tremendous impact on New Zealand culture (Armstrong, 2010), and doubtless will continue to do so.

Acknowledgements

The pun in the title of this article originates from an anonymous comment in the *Otago Witness* column 'Passing Notes' from October 10, 1874.

Fundings

This work was not supported by any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest disclosure

The author declares that they have no financial conflicts of interest in relation to the content of the article.

Data, script, code, and supplementary information availability

Data and code are available online: https://doi.org/10.17605/OSF.IO/E59H6.

References

Armstrong P (2010). Moa Citings. *The Journal of Commonwealth Literature* 45, 325–339. https://doi.org/10.1177/0021989410376799.

Brook BW, SR Sleightholme, CR Campbell, I Jarić, and JC Buettel (2023). Resolving when (and where) the Thylacine went extinct. *Science of The Total Environment* 877, 162878. https://doi.org/10.1016/j.scitotenv.2023.162878. [207][208][209]

Canoville A, A Chinsamy, and D Angst (2022). New Comparative Data on the Long Bone Microstructure of Large Extant and Extinct Flightless Birds. *Diversity* 14. ISSN: 1424-2818. https://doi.org/10.3390/d14040298. [210][211]

Diamond J (2000). Blitzkrieg Against the Moas. *Science* 287, 2170–2171. https://doi.org/10.1126/science.287.5461.2170. [212][213]

Foxon F (2023a). If it's there, could it be a bear? *PeerRef*. https://doi.org/10.1101/2023.01.14.524058. [214]

— (2023b). What's in Lake Champlain? Analysing historic sightings of the cryptid known as "Champ". *The Skeptic*. https://www.skeptic.org.uk/2023/06/whats-in-lake-champlain-analysing-historic-sightings-of-the-cryptid-known-as-champ/. [215][216][217]

Greenwell JR (1985). A Classificatory System for Cryptozoology. *Cryptozoology* 4, 1–14. [218]

Heuvelmans B (1982). What is Cryptozoology? *Cryptozoology* 1, 1–12. [219]

— (1984). The Birth and Early History of Cryptozoology. *Cryptozoology* 3, 1–30. [220]

— (1986). Annotated Checklist of Apparently Unknown Animals With Which Cryptozoology is Concerned. *Cryptozoology* 5, 1–26. [221][222]

— (1988). The Sources and Method of Cryptozoological Research. *Cryptozoology* 7, 1–21. [223]

Higham T (1994). Radiocarbon dating New Zealand prehistory with moa eggshell: some preliminary results. *Quaternary Science Reviews* 13, 163–169. https://doi.org/10.1016/0277-3791(94)90043-4. [224][225]

Holdaway RN, ME Allentoft, C Jacomb, CL Oskam, NR Beavan, and M Bunce (2014). An extremely low-density human population exterminated New Zealand moa. *Nature Communications* 5, 5436. https://doi.org/10.1038/ncomms6436. [226][227][228]

Irwin G and C Walrond (2016). When was New Zealand first settled? *Te Ara - the Encyclopedia of New Zealand*. http://www.TeAra.govt.nz/en/when-was-new-zealand-first-settled. [229][230]

Mackal RP (1983). *Searching for Hidden Animals*. Cadogan Books. [231]

McCulloch B (1982). *No Moa*. Canterbury Museum, Christchurch. [232]

Nickell J (2017). The New Zealand Moa: From Extinct Bird to Cryptid. *Skeptical Inquirer*. https://skepticalinquirer.org/newsletter/the-new-zealand-moa-from-extinct-bird-to-cryptid/. [233][234]

Paxton CGM (2009). The plural of 'anecdote' can be 'data': statistical analysis of viewing distances in reports of unidentified large marine animals 1758–2000. *Journal of Zoology* 279, 381–387. https://doi.org/10.1111/j.1469-7998.2009.00630.x. [235][236][237]

Perry GL, AB Wheeler, JR Wood, and JM Wilmshurst (2014). A high-precision chronology for the rapid extinction of New Zealand moa (Aves, Dinornithiformes). *Quaternary Science Reviews* 105, 126–135. ISSN: 0277-3791. https://doi.org/10.1016/j.quascirev.2014.09.025. [238][239][240]

Radford B and J Nickell (2006). *Lake Monster Mysteries: Investigating the World's Most Elusive Creatures*. University Press of Kentucky. [241][242]

Sandom C, S Faurby, B Sandel, and JC Svenning (2014). Global late Quaternary megafauna extinctions linked to humans, not climate change. *Proceedings of the Royal Society B: Biological Sciences* 281, 20133254. https://doi.org/10.1098/rspb.2013.3254. [243][244][245]

Scoble W (1952). Unpublished letter to Dr R. A. Falla, Director of the Dominion Museum. Dated December 28, 1952. [246][247]

Scofield RP and KWS Ashwell (2009). Rapid somatic expansion causes the brain to lag behind: the case of the brain and behavior of New Zealand's Haast's Eagle (Harpagornis moorei). *Journal of Vertebrate Paleontology* 29, 637–649. https://doi.org/10.1671/039.029.0325. [248][249][250]

Silverberg R (1973). *The Dodo, the Auk and the Oryx: Vanished and Vanishing Creatures*. Puffin Books. [251]

Spittle B (2010). *Moa Sightings Volumes 1–3*. Paua Press Limited, Dunedin. [252]

Wehi PM, MP Cox, T Roa, and H Whaanga (2018). Human perceptions of megafaunal extinction events revealed by linguistic analysis of indigenous oral traditions. *Human Ecology* 46, 461–470. https://doi.org/10.1007/s10745-018-0004-0.

Widrig K and DJ Field (2022). The evolution and fossil record of Palaeognathous birds (Neornithes: Palaeognathae). *Diversity* 14, 105. https://doi.org/10.3390/d14020105.

Worthy TH, VL De Pietri, and RP Scofield (2017). Recent advances in avian palaeobiology in New Zealand with implications for understanding New Zealand's geological, climatic and evolutionary histories. *New Zealand Journal of Zoology* 44, 177–211. https://doi.org/10.1080/03014223.2017.1307235.