Profound seizure suppression and disease modification by targeting JAK1, a key driver of a pro-epileptogenic gene network.

Olivia R. Hoffman[1,2], Anna M. Patterson[1], Emily S. Gohar[1], Emanuel Coleman[3], Jose Ezekial Clemente Espina[4], Barry A. Schoenike[1], Claudia Espinosa-Garcia[4], Felipe Paredes[5], Raymond J. Dingledine[5], Jamie L. Maguire[3], Avtar S. Roopra[1,2]*

[1]Department of Neuroscience, University of Wisconsin-Madison School of Medicine and Public Health, Madison, WI 53705, USA; [2]Molecular and Cellular Pharmacology Graduate Program, University of Wisconsin-Madison School of Medicine and Public Health, Madison, WI 53705, USA; [3]Department of Neuroscience, Tufts University School of Medicine, Boston, MA 02111, USA; ; [4]Cellular and Molecular Biology Graduate Program, University of Wisconsin-Madison School of Medicine and Public Health, Madison, WI 53705, USA [5]Department of Pharmacology & Chemical Biology, Emory University School of Medicine, Atlanta, GA 30322, USA.

ABSTRACT

Epilepsy is the 4th most prevalent neurological disorder with over 50 million cases worldwide[1,2]. While a number of drugs exist to suppress seizures, approximately 1/3 of patients remain drug resistant, and no current treatments are disease modifying[3]. Using network and systems-based approaches, we find that the histone methylase EZH2 suppresses epileptogenesis and slows disease progression, via repression of JAK1 and STAT3 signaling in hippocampal neurons. Pharmacological inhibition of JAK1 with the orally available, FDA-approved drug CP690550 (Tofacitinib)[4,5] profoundly suppresses behavioral and electrographic seizures after the onset of epilepsy across preclinical rodent models of acquired epilepsy. This seizure suppression persists for weeks after drug withdrawal. Identification of an endogenous protective response to status epilepticus in the form of EZH2 induction has highlighted a critical role for the JAK1 kinase and STAT3 in both the initiation and propagation of epilepsy across preclinical rodent models and human disease. Overall, we find that STAT3 is transiently activated after insult, reactivates with spontaneous seizures, and remains targetable for disease modification in chronic epilepsy.

INTRODUCTION

Epilepsy is a highly prevalent neurological disorder with over 50 million cases worldwide[1,2]. Arising from a myriad of etiologies including gene mutations and environmental insults, the unifying characteristic of the epilepsies is the emergence of recurring, spontaneous seizures. Epileptogenesis is the process that links initial insults to the eventual propagation of spontaneous seizures. One of the insults that may initiate epileptogenesis is status epilepticus: an unremitting bout of seizures lasting for more than five minutes, during which the patient does not regain consciousness[6-8]. The first-line treatments for epilepsy are anticonvulsants aimed at controlling seizures. While these medications effectively suppress seizures for some patients, they do not target the processes that orchestrate progression of the disease, and fail to suppress seizures altogether in approximately 1/3 of patients[3]. There is a critical need for treatments that target the progressive decline in seizure threshold that arguably leads to spontaneous seizures.

A poor understanding of the mechanisms underlying epileptogenesis has so far prevented the development of disease biomarkers and translation of antiepileptogenic treatments from pre-clinical rodent models to the clinic. Although others have reported the disease-modifying potential of manipulations to metabolism[9-11], inflammation[12,13], chromatin modification[14], and signal transduction pathways[15], progress has been hindered by a lack of consensus among laboratories employing different models of status epilepticus (SE)[16]. With few exceptions[9,17-20], little effort has been made to identify the principal transcription factors and chromatin modifiers whose function precipitates the gene changes that arise after status epilepticus.

Recently, we applied a systems approach and novel bioinformatic tool[21] to a transcriptomic consortium dataset that incorporated gene expression data across multiple laboratories and models of epilepsy[22]. This allowed for identification of potential mechanisms that span across rodent models of status epilepticus (SE) and human temporal lobe epilepsy (TLE)[23]. With this approach, we identified the histone methylase Enhancer of Zeste Homolog 2 (EZH2) as a potential driver of gene changes in dentate granule cells post-SE that is induced by an order of magnitude across epilepsy models. Transient pharmacological inhibition of EZH2 post-SE worsened

epileptic phenotypes one month later, suggesting that EZH2 induction plays a protective role in
epileptogenesis. In addition to EZH2, our
analysis predicted Signal Transducer
and Activator of Transcription 3 (STAT3)
to be a driver of gene activation across
models of SE[22,23].

Of the studies that have profiled gene
changes in the epileptic brain, nearly all
find a robust activation of inflammatory
pathways[17-20,23-25]. In the brain,
inflammation can be triggered by glial
cells[26], but neurons also exhibit an
innate immune response that has been
associated with hyper-excitability[27]. Anti-
inflammatory drugs have been used with
varying degrees of success to suppress
seizures in human patients[28-31], and
some drugs targeting neuroinflammation
in rodents have shown antiepileptogenic
potential[32-35]. However, the endogenous
mechanisms that regulate neurogenic
inflammation post-SE are poorly
understood.

Herein, we show that EZH2 induction
with SE suppresses pathological
activation of pro-inflammatory JAK/STAT
signaling in neurons, both in animal
models and human disease. EZH2
deletion in neurons leads to hyper-
activation of the JAK/STAT pathway,
robust reductions in seizure threshold
after SE, and an exaggerated epileptic
phenotype. Pharmacological inhibition of
JAK1/3 activity acutely after SE rescues
these phenotypes. Finally, targeting
STAT3 reactivation after the onset of
epilepsy with the JAK inhibitor
CP690550 results in an enduring,
disease-modifying suppression of
spontaneous seizures.

Fig. 1. EZH2 maintains seizure threshold after status epilepticus and mitigates spontaneous seizure frequency and severity. (A) Experimental paradigms for panels B-C and D-E. **(B)** Baseline fluorothyl seizure threshold data for C57BL6J EZH2nWT, nHet, and nHom mice. In naïve mice, deletion of EZH2 in neurons does not alter seizure threshold (one-way ANOVA; nWT:nHet p=0.471, nWT:nHom p=0.134, nHet:nHom p=0.471; n=8-16). **(C)** After induction of SE by KA injection, seizure threshold does not change robustly in EZH2nWT mice (p=0.061). EZH2nHet mice exhibit a 35% reduction in seizure threshold post-SE (p<10^{-4}), with thresholds 25% lower than kainate-injected EZH2nWT mice (p=0.003). EZH2nHom mice show a 25% reduction in seizure threshold post-SE (p<10^{-4}), with thresholds 17% lower than KA EZH2nWT mice (p=0.026) (2-way ANOVA; n=7-14). **(D)** FVB EZH2nWT and nHom mice were injected with KA to induce SE and video recorded from 5-7 weeks post-SE to count and score spontaneous seizure activity. Deletion of an EZH2 allele in neurons leads to a 2.5-fold increase in daily seizure events (unpaired t-test; p<10^{-4}; n=6-10). **(E)** Total observed Racine scale events for EZH2nHom and EZH2nWT mice were quantified as a frequency distribution. 50% of EZH2nWT mice had only Class I seizures, while 50% of EZH2nHom mice had Class I and II seizures (Mann-Whitney test; U:284, p=0.002; n=6-10).

RESULTS

Neuronal EZH2 maintains seizure threshold and protects against exacerbated epileptic phenotypes.

We previously identified the histone methyltransferase EZH2 as a potential driver of gene repression post-SE,
and EZH2 protein was shown to be potently induced over a 10-day window post-SE in rats and mice across
models of epilepsy. EZH2 induction was protective because pharmacological inhibition resulted in accelerated
disease progression[23]. To determine the role of EZH2 in epilepsy, we crossed EZH2 floxed mice[36] with
Syapsin1 Cre drivers[37] to delete EZH2 in mature neurons. Neuronal specificity of Cre expression was
confirmed via crossing Syn1Cre with *flox*-STOP-*flox*-tdTomato mice and immunofluorescence (Fig. S1A).

Immunofluorescence and western blot analysis confirmed successful ablation of neuronal EZH2 post-SE. Fig. S1B-C show diminished induction of EZH2 on day 4 post-SE in mice lacking one EZH2 allele in neurons (EZH2nHet) or lacking both alleles (EZH2nHom). Staining for EZH2 confirmed reduced induction of neuronal EZH2 in EZH2nHet versus wildtype (EZH2nWT) mice (Fig. S1D, quantified in S1E).

We tested the impact of EZH2 deletion on fluorothyl seizure threshold acutely post-SE, as a correlation between fluorothyl seizure threshold[38] and the development of spontaneous seizures has been suggested[39-42]. We measured time to generalized tonic clonic seizure (GTCS) after exposure to the volatile chemoconvulsant fluorothyl in EZH2nWT, EZH2nHet, and EZH2nHom mice aged 4-6 weeks, before and after kainate-induced SE[43,44] (Fig. 1A). Loss of EZH2 alleles in neurons had no impact on seizure threshold in naïve mice (Fig. 1B), and induction of SE did not alter the seizure threshold of EZH2nWT mice on day 4 post-SE (Fig. 1C). In contrast, robust reductions in threshold were detected after SE in EZH2nHet mice (2-way ANOVA; Mean difference:55 ± 8sec, $p<10^{-4}$) and EZH2nHom mice (Mean difference:46 ± 8sec, $p<10^{-4}$). These were significant reductions compared to EZH2nWT mice (EZH2nWT:EZH2nHet: 28%, $p<0.01$; EZH2nWT:EZH2nHom: 20%, $p<0.05$). These data suggest that EZH2 is required to maintain fluorothyl seizure threshold post-SE.

We considered the possibility that EZH2 loss primed the mice to experience more severe responses to kainate injection, and it was this vulnerability to SE driving the observed reduction in mutant seizure threshold. However, we found no differences in time to first class V seizure and no correlation between time to first class V seizure and time to GTCS post-SE for either EZH2nWT (Pearson correlation; R^2:0.016, $p=0.633$), EZH2nHet (R^2:0.261, $p=0.195$), or EZH2nHom mice (R^2:0.152, $p=0.209$) (Fig. S2A-D).

To test the impact of neuronal EZH2 deletion on spontaneous seizures, we monitored behavioral spontaneous seizures in FVB EZH2nWT and EZH2nHom mice by video weeks five to seven post-SE (Fig. 1A). EZH2nHom mice exhibited a 2.5-fold increase in average daily seizure events compared to EZH2nWT mice (unpaired t-test; $p<10^{-4}$) (Fig. 1D). EZH2nHom mice also exhibited more severe Racine scale seizures than EZH2nWT mice (Mann Whitney test; U: 284, $p<0.01$) (Fig. 1E). These data support the hypothesis that neuronal EZH2 induction is a protective response post-SE.

EZH2 tempers induction of the JAK/STAT pathway post-SE in neurons.

To begin to understand the mechanism behind EZH2-mediated protection, we performed bulk RNAseq on whole hippocampal tissue from naïve and epileptic (4d. post-SE) EZH2nWT and EZH2nHom mice. Of the 10556 expressed genes, 6241 were found to be differentially expressed at an adjusted $p<10^{-2}$ (Likelihood Ratio Test) across the four conditions (saline, kainate, EZH2nWT, EZH2nHom). A community-based network approach was used to group genes into clusters. Leiden clustering generated 12 distinct groups (Fig. 2A). The grey, orange, gold, brown, skyblue, and purple clusters were generally repressed after SE, whereas the red, blue, indigo, yellow, and black clusters were generally increased. We focused on the Black cluster because these 439 genes manifested induction with SE in EZH2nWT mice but showed an exaggerated induction with SE in EZH2nHom mice. We reasoned that these would be genes whose expression is induced with SE but tempered in the presence of EZH2 post-SE, and undergo unfettered induction in the absence of EZH2.

To understand the nature of genes comprising the Black cluster, we performed an ontological analysis. The 439 Black cluster genes were compared to gene lists associated with terms in the KEGG, Reactome, Biocarta and Hallmark databases. (Fig. 2B). The ontology network (see Fig. S3A) highlights the innate immune response (NFkB, Interferon Gamma), microglial processes (Complement) and inflammation (IL6 Jak Stat3) amongst others. The term 'IL6 Jak Stat3' was of particular interest given that our previous work highlighted STAT3 as a driver of gene activation across models of epilepsy[23], and others have identified a role for the JAK/STAT pathway in epileptogenesis[27,32,45-48].

To determine which transcription factors and cofactors (hereafter together referred to as 'Factors') control expression of Black cluster genes, we used MAGIC[21]. STAT3 was the highest scoring Factor (Fig. 2D). When projected as a Gene Regulatory Network (GRN) (Fig. 2C) such that an arrow projects from Factor X to Factor Y if the gene encoding Factor Y is a transcriptional target of Factor X, STAT3 stood out as a central node. Thus, STAT3 had the highest in-degree, out-degree and total edge count (Fig. 2E) of any Factor in the

network. Finally, we
assessed the Betweenness
Centrality of the network[49] to
determine which Factors
might be critical
for information flow in the
GRN. STAT3 had by far the
largest Betweenness
Centrality of any Factor (Fig.
2F).

To ascertain whether the
mouse Black cluster has a
counterpart in human
epilepsy, we used whole
transcriptome data from 129
temporal lobectomy
resections collected ante-
mortem from human
temporal lobe epilepsy
(TLE) patients[23,50]. We used
k-means clustering to
generate 10 gene clusters
from the 7658 genes co-
expressed across all
samples (brown, blue,
purple, skyblue, grey,
orange, indigo, green,
yellow, red). To identify the
human TLE cluster most
similar to the mouse Black
cluster, we performed Fisher
exact tests to measure the
degree of overlap between
the 439 Black cluster genes,
and the genes comprising
each cluster of the human
TLE dataset (Fig. 2G). The
human Brown cluster
showed the greatest
overlap, sharing 65 genes in
common with the mouse
Black cluster (Fisher exact;
Odds: 4.79, p=1.19E-20).
We then performed
ontology analysis on the
genes in the human Brown
cluster. The 850 genes were
compared to gene lists
associated with terms in the
KEGG, Reactome, Biocarta
and Hallmark databases.
(Fig. 2H). The ontology

Fig. 2. STAT3 drives hyper-activation of pro-inflammatory gene networks post-SE in EZH2 knockout mice and human temporal lobe epilepsy. (A) Leiden clusters from 6241 differentially expressed genes (fold change >1.2x, adjusted p<0.01) across saline, kainate, EZH2nWT, and EZH2nHom conditions. (B) Ontological analysis of genes in the Black cluster highlights inflammatory processes. (C) MAGIC was used to determine which Factors control the expression of Black cluster genes, and positive hits were projected as a Gene Regulatory Network (GRN). (D) Top 5 Factors driving expression in the Black cluster sorted by –log(adjusted p value). (E, F) Factors were ranked by their in degree, out degree counts, and betweenness centrality. (G) Comparison of 10 human TLE k-means gene clusters to the mouse Black cluster by Fisher Exact test. Plotting -log(adjusted p) versus odds ratio of Fisher tests against the Black cluster shows that the Human Brown cluster is the most similar to the mouse Black cluster (Fisher exact; Odds:4.79, p(adj)=1.2x10[-20]). Dotted line shows p=0.05. (H) Ontological analysis of genes in the human-Brown cluster. The 850 human-Brown cluster genes were compared to gene lists associated with terms in KEGG, Reactome, Biocarta, and Hallmark databases (I) MAGIC GRN for Factors driving the human-Brown cluster. (J, K) Factors were ranked by their in degree, out degree, and betweenness centrality. (L) STAT3 is the top driver predicted by MAGIC to drive expression of genes common to the mouse-Black and human-Brown clusters.

network (see Fig. S3B) highlights inflammation (Cytokine Receptor Interaction/Complement/Innate Immune
System) and interferon signaling (Interferon Gamma Response). Mirroring results of the black cluster, STAT3

221 in the human Brown GRN had the highest in-degree, out-degree, and betweenness centrality of any Factor in
222 the network (Fig. 2J-K).
223
224 MAGIC called STAT3 as the highest scoring Factor driving expression of the 65 genes co-expressed in early
225 rodent epileptogenesis (mouse Black) and frankly epileptic human tissue (human Brown) (Fig. 2L). This data is
226 consistent with the hypothesis that STAT3-mediated disease mechanisms that are activated acutely in a rodent
227 model of epileptogenesis persist in chronic human epilepsy.
228
Transient activation of the JAK/STAT pathway after SE in WT mice has been reported by others[23,46] (and Fig.
S4A-B, S5A-D). To test for a neuronal EZH2-dependent change in JAK/STAT signaling, we performed western
blot analysis on hippocampal lysates from 4d. post-SE WT and EZH2 mutant mice (Fig. 3A-B). On day 4 post-
SE, pY705-STAT3 was induced 2-fold in EZH2nWT mice (p<0.01) compared to naïve controls, 2.9-fold in
EZH2nHet mice (p<10^{-4}), and 4.5-fold in EZH2nHom mice (2-way ANOVA; p<10^{-4}) (Fig. 3C-D). These results
demonstrated an EZH2 gene dose-dependent hyperactivation of STAT3.
235
236 Given the involvement of STAT3 in regulating gene expression across acute rodent SE and chronic human
237 epilepsy, we asked whether STAT3 reactivation occurred in chronic rodent epilepsy. We found that at 14
238 weeks post-SE, epileptic mice showed a 4.7-fold induction of pY705-STAT3 over naïve controls (unpaired t-
239 test; p<0.01) (Fig. 3E-F).

Fig. 3. JAK/STAT pathway components are hyper-activated in neurons of EZH2 knockout mice. EZH2nWT, nHet, and nHom mice were injected with saline or KA to induce SE; hippocampi were harvested on day 4 post-SE. **(A, B)** Representative western blots for pY705-STAT3, total STAT3, and beta-actin on day 4 post-SE. **(C)** pY705-STAT3 levels are unchanged across genotypes in naïve animals (p=0.220). pY705-STAT3 is induced by 2-fold in EZH2nWT mice (p=0.022), 2.9-fold in EZH2nHet mice (p<10^{-4}), and 4.5-fold in EZH2nHom mice post-SE (p<10^{-4}). pY705 levels in EZH2nHom mice post-SE are 2.6-fold greater than in EZH2nWT mice post-SE (p<10^{-4}) (2-way ANOVA; n=7-12). **(D)** Total STAT3 levels on day 4 post-SE. Total STAT3 is induced by 1.2-fold in nHom mice post-SE, compared to nWT mice post-SE (p=0.041). Total STAT3 is induced after seizures in nHet mice by 1.5-fold (p=0.041) (2-way ANOVA; n=7-12). **(E)** Representative western blots for pY705-STAT3, total STAT3, and beta actin in naïve and chronically epileptic mice at 14 weeks post-SE. **(F)** pY705-STAT3 is induced 4.7-fold in epileptic mice (unpaired t-test; p=0.007, n=5-7). **(G)** 20x images of the dentate gyrus (DG) and CA1 pyramidal cell layer (CA1). Hippocampal slices were collected on day 1 post-SE from saline- and KA-injected mice; immunofluorescence was performed to detect nuclear DNA, NeuN, and total STAT3.

To define the cell types in which STAT3 is expressed after seizures, we performed immunofluorescence with DAPI, NeuN, and total STAT3 antibodies on day 1 post-SE hippocampi taken from mice injected with saline or kainic acid (Fig 3G). We made use of the fact that the STAT3 gene is induced with STAT3 activation, and so elevated total STAT3 protein was a proxy for activated STAT3. In both the dentate granule and CA1 pyramidal cell layer, we observed a robust induction of total STAT3 in NeuN-positive cells.

EZH2 targets STAT3 via JAK1 to suppress JAK/STAT signaling.

STAT3 was a central factor in the Black cluster GRN whereas, surprisingly, EZH2 was not. In order to understand how STAT3 hyperactivation could have occurred in the absence of EZH2, shRNA was used to knock down EZH2 in the Neuro2A (N2A) neuronal cell line. We observed robust diminution of EZH2 in knockdown cells (shEZH2) (Fig. 4A, quantified by S6A) (unpaired t-test; 3.7-fold, $p<10^{-4}$). This was associated with a marked increase in pY705-STAT3 levels (unpaired t-test; 2.4-fold $p<10^{-4}$) (quantified by S6B). This result suggests that EZH2 suppresses STAT3 activation in a cell-autonomous manner without circuit level alterations.

To determine whether STAT3-upstream kinases are required for pY705-STAT3 induction in shEZH2 cell lines, we incubated cells with either WP1066 (an analogue of the JAK2 inhibitor AG490[51-53]) or CP690550 citrate (a small molecule inhibitor of JAK1/3 kinase activity[4,5,54]). WP1066 did not impact STAT3 phosphorylation in shEZH2 cells at any concentration tested (Fig. S7A-C). In contrast, 500nM CP690550 citrate resulted in a 2.3-fold (2-

Fig. 4. EZH2 suppresses STAT3 activation via transcriptional repression of JAK1. N2A cells were stably transfected with EZH2 shRNA (shEZH2) or non-targeting shRNA (shNT). (A) Knockdown of EZH2 causes induction of pY705-STAT3 and total STAT3 protein (quantified in Fig. S6). (B,C) pY705-STAT3 activation in shEZH2 cells is inhibited with the JAK inhibitor CP690550 citrate. pY705-STAT3 is induced in vehicle-treated shEZH2 cells by 3.7-fold (p=0.0004). CP690550 treatment reduces pY705-STAT3 levels of KD cells by 2.3-fold (at 0.5µM) and 3.2-fold (p=0.0006) (at 5µM). At 0.5µM CP, pY705 levels in KD cells are reduced to that of untreated control cells (p=0.18) (2-way ANOVA; n=3). (D) JAK1 mRNA expression is induced by 5.1-fold in shEZH2 cells (unpaired t-test; p=0.019; n=6). (E) JAK1 protein is induced 5.5 fold (p=0.0008), and pY705-STAT3 by 6.7-fold (p=0.0004) in the absence of changes to total STAT3 upon transfection with FLAG-JAK1 compared to control transfection (unpaired t-tests; n=3). (F) Schematics depicting EZH2 binding sites on JAK1. (G) H3K27Me3 is enriched at Site 2 at the JAK1 locus. Chromatin IP (ChIP) on N2A cells treated with either vehicle or 10µM UNC1999 for 24 hours, or transfected with expression vectors GFP (control) of cotransfected with GFP-EZH2, SUZ12, and EED (PRC2). Results are graphed as fold H3K27Me3 over input. For JAK1 (Site 2), treatment with UNC1999 reduces H3K27Me3 enrichment by 3.5-fold (p=0.029), and PRC2 overexpression leads to a 2.7-fold increase in H3K27Me3 enrichment (p=0.0006) (unpaired t-tests; n=3). (H) Knockdown of JAK1 suppresses pY705-STAT3 activation in shEZH2 cells. shNT and shEZH2 N2A cells transfected with either non-targeting (NT) or JAK1 siRNA were subjected to western blotting to confirm JAK1 and EZH2 knockdown, and assess pY705-STAT3 levels. (I) Total JAK1 protein is induced in shEZH2 cells transfected with non-targeting siRNA by 1.2-fold (p=0.023), and reduced in both shNT (p=0.0005) and shEZH2 ($p<10^{-4}$) after JAK1 RNAi (2-way ANOVA; n=9-12). (J) pY705-STAT3 is induced in shEZH2:NTsiRNA cells by 1.7-fold (p=0.027) over shNT:NTsiRNA cells. JAK1 RNAi in shEZH2 cells reduces pY705-STAT3 by 2.5-fold (p=0.006), to the levels of shNT:NTsiRNA cells (2-way ANOVA; n=9-12).

way ANOVA; $p<0.01$) reduction of pY705-STAT3 in shEZH2 cells (Fig. 4B,C), rescuing pY705-STAT3 levels to that of vehicle-treated control cells. Even greater reduction occurred in the shEZH2 cells treated with 5μM CP690550 (3.2-fold, $p<10^{-3}$). CP690550 is a JAK inhibitor selective for JAK1 and JAK3, but JAK3 is not expressed in neurons or N2As (data not shown). We therefore tested whether JAK1 was elevated in the shEZH2 cells. Fig. 4D shows a 5.1-fold induction of JAK1 mRNA over control cells (unpaired t-test; $p<0.05$).

To determine whether induction of JAK1 is sufficient to induce STAT3 activation, we overexpressed JAK1 in N2A cells (Fig. 4E). JAK1 shows a 5.6-fold induction upon overexpression (unpaired t-test; $p<10^{-3}$), and pY705-STAT3 is induced by 6.7-fold (unpaired t-test; $p<10^{-3}$) in the absence of changes to total STAT3 (quantified in S8A-B). This suggests that CP690550 works to suppress STAT3 activation primarily through JAK1 downstream of EZH2.

To understand how EZH2 regulates JAK1 activity, we noted three putative EZH2 binding sites in the gene body and promoter region of JAK1 using ENCODE tracks in the UCSC Genome Browser[55] (Fig. 4F, Fig. S9A). We performed chromatin immunoprecipitation with anti-H3K27Me3 antibodies to determine whether EZH2 methylase activity was enriched at any of these sites. In Fig. 4G H3K27Me3 was enriched at site 2. Incubation of cells with UNC1999 (an EZH2 inhibitor[23,56]) for 24 hours resulted in a 3.5-fold loss of signal (unpaired t-test; $p<0.05$). Overexpression of EZH2, SUZ12, and EED (the core components of Polycomb) increased H3K27Me3 signal by 2.7-fold compared to overexpression of GFP as a control (unpaired t-test; $p<10^{-3}$). PAX2 and actin were immunoprecipitated as positive and negative controls for H3K27Me3 enrichment, respectively (Fig. S9B-C); ChIP-PCR results for JAK1 (Site 1), JAK1 (Site 3), JAK2, and STAT3 were also collected (Fig. S9D-G). None of these sites showed both a loss of H3K27Me3 enrichment after UNC1999 treatment and an increase in signal upon overexpression of Polycomb. These data are consistent with the hypothesis that EZH2 targets JAK1 transcription to temper activation of STAT3.

Finally, to corroborate our pharmacological findings, we transfected control and shEZH2 cells with siRNA against JAK1 or a non-targeting transcript, and validated knockdown of EZH2 and JAK1 by western blot (Fig. 4H). Total JAK1 protein is induced 1.2-fold upon EZH2 knockdown (2-way ANOVA; $p<0.05$). JAK1 RNA interference reduced JAK1 protein levels in both control (2-way ANOVA; 1.6-fold, $p<10^{-3}$) and shEZH2 cells (2.4-fold, $p<10^{-4}$) (Fig. 4I). Knockdown of JAK1 expression in shEZH2 cells was sufficient to rescue pY705-STAT3 levels to that of control cells transfected with non-targeting siRNA (2-way ANOVA; 2.0-fold, $p<0.01$) (Fig. 4J). These data support the hypothesis that CP690550 rescues STAT3 hyperactivation primarily through targeting JAK1.

Transient JAK inhibition acutely post-SE rescues epileptic phenotypes in EZH2 knockout mice.

To test whether loss of EZH2-mediated regulation of JAK1 contributed to the exacerbated disease progression observed in EZH2 mutant mice, we treated EZH2nWT and EZH2nHet mice with CP690550 after induction of SE. Because the hyper-induction of pY705-STAT3 in mutant mice occurred already at day 1 post SE (Fig. S10A-C), we injected mice with CP690550 or vehicle at 6, 12, and 24 hours post-SE. A cohort of mice was sacrificed three hours after final injection to assess hippocampal STAT3 activation by western blot to confirm target engagement (Fig. S10D-E). Consistent with the previous result (see Fig. 1B), neuronal deletion of EZH2 did not alter fluorothyl seizure threshold in naïve mice (Fig. 5B). CP690550 treatment raised the 4d. post-SE seizure threshold of EZH2nHet mice by 30 seconds (1-way ANOVA; $p<0.05$), rendering them indistinguishable from epileptic EZH2nWT mice (1-way ANOVA; $p>0.05$) (Fig. 5C). Thus, CP690550 rescued the increased seizure susceptibility phenotype associated with the loss of neuronal EZH2. These data support the hypothesis that EZH2 maintains seizure threshold post-SE by tempering activation of the JAK/STAT pathway.

To test whether transient CP690550 treatment immediately after SE mitigated long-term disease progression, mice were video recorded from weeks five to seven post-SE (Fig. 5A). CP690550 treatment immediately after SE reduced daily seizure frequency by 41% in EZH2nHom mice 5 weeks later (1-way ANOVA; $p<10^{-4}$). EZH2nHom seizure frequency was reduced from 2.6-fold above EZH2nWT to 1.5-fold (1-way ANOVA; $p<0.01$) (Fig. 5D). EZH2nHom mice treated acutely with CP690550 also exhibited a dramatic reduction in seizure

severity from
untreated EZH2nHom
mice (Mann-Whitney
test; U:138.5,
p<0.01), such that
they were
indistinguishable from
WT mice (Fig. 5E).
360
To test whether
CP690550 could
rescue the increased
seizure burden of
EZH2nHom mice if
administered during
the chronic period,
we recorded
spontaneous
seizures for two
weeks post-SE in
EZH2nWT and nHom
mice starting 5 weeks
post SE, followed by
another 2-week
recording period
during which the
EZH2nHom mice
received daily
injections of
CP690550 (Fig. 5A).
Daily CP690550
treatment in the
chronic period
reduced seizure
severity and
frequency by 80% in
EZH2nHom mice (1-
way ANOVA; p<10^{-4})
(Fig. 5F-G). Indeed,
seizure frequency of
EZH2nHom+CP mice
was reduced to below
EZH2nWT levels (1-

Fig. 5. JAK inhibition rescues seizure threshold reductions and spontaneous seizure burden in EZH2nKO mice. (A) (For panels B-C) Naïve seizure threshold was measured in C57 EZH2nWT and nHet mice one week prior to KA injections. (B) Deletion of an EZH2 allele does not alter seizure threshold in naïve mice (unpaired t-test; p=0.246; n=24-29). (C) nHet mice were injected with vehicle or CP690550 at 6, 12, and 24h. post-SE. Seizure threshold was measured 4 days post-SE. Untreated nHet mice show a 35 second reduction in seizure threshold (p=0.007) compared to nWT mice. CP690550 raises seizure thresholds of nHet mice by 30 seconds (p=0.025), extinguishing the difference between treated nHet and nWT mice (p=0.709) (1-way ANOVA; n=7-9). (D) FVB EZH2nWT and nHom mice received CP690550 or no treatment at 1, 6, and 24h. post-SE, and SRS activity was monitored by video from 5-7 weeks post-SE (see panel A). nHom mice treated acutely 6,12 and 24 hours post-SE with CP690550 exhibit a 41% reduction in daily seizures compared to untreated nHom mice (p<10^{-4}); a 34% difference remains between treated nHom mice and nWT mice (p=0.0073) (1-way ANOVA; n=4-10). (E) Racine scale events represented as a frequency distribution. Acute CP690550 treatment reduces seizure severity in nHom mice (Mann Whitney; U:138.5, p=0.009; n=4-10) to nWT levels (U:357, p=0.955). 50% of nWT and treated nHom mice had Class I seizures, while 50% of untreated nHom mice had Class I and II seizures. (F) SRS activity was monitored from 5-7 weeks post-SE, after which nHom mice received daily 10mg/kg CP690550 injections during monitoring from 7-9 weeks post-SE (see panel A). CP treatment reduced daily seizures by 80% in nHom mice (p<10^{-4}); treated nHom mice exhibit daily seizure frequencies 58% lower than nWT mice (p=0.0008) (1-way ANOVA; n=6-10). (G) Total observed Racine scale events represented as a frequency distribution. Daily CP690550 reduces seizure severity in nHom mice. 90th percentile seizure severities of nHom mice before and after treatment: 5.3 and 1, respectively (Wilcoxon matched pairs signed rank; W:-36.00, p=0.007, n=6). 90th percentile severity of WT mice: 2.6 (Mann Whitney test comparing nWT and treated nHom mice; U:119, p=0.042; n=6-10).

way ANOVA; 1.9-fold, p<10^{-3}). Given the robust seizure suppression when administered during the chronic
period, we considered the possibility that CP690550 may simply be an anticonvulsant. To test this, we
measured seizure threshold using both flurothyl and PTZ (pentylenetetrazole)[57,58] in mice pre-treated with
vehicle or CP690550 (15mg/kg, i.p.) thirty minutes prior to testing. This time was chosen because peak plasma
levels are observed thirty minutes after oral administration[59] CP690550 pre-treatment had no effect on flurothyl
or PTZ seizure threshold (Fig. S10F-G).
401
402 **CP690550 profoundly suppresses electrographic and behavioral spontaneous seizures in**
403 **epileptic WT mice.**
404
405 CP690550 treatment during the chronic period reduced seizure frequency in EZH2nHom mice to below WT
406 levels, so we tested its effect on WT epileptic mice. Chronic administration of CP690550 following
407 intrahippocampal injection of kainate (IHKA) in WT epileptic mice significantly diminished spontaneous

recurrent seizures. In a paradigm involving one week of daily vehicle (i.p.) administration followed by an additional week of either vehicle or CP690550 injections (15mg/kg, i.p.) under continuous EEG recording (Fig. 6A), CP690550 treatment during the chronic period profoundly suppressed spontaneous seizures. Notably, 50% of vehicle-treated mice exhibited up to 1 seizure per day, with corresponding daily seizure burdens up to 99 (Fig. 6D-E). In contrast, 50% of CP690550-treated mice displayed up to 0.1 seizures per day, with daily seizure burdens up to 11. CP690550 treatment robustly reduced both spontaneous seizure frequency (Kolmogorov-Smirnov test; D:0.9091, $p<10^{-3}$) and burden (D:0.7083, p<0.01) in epileptic wildtype mice.

Additionally, while all animals experienced seizures during vehicle treatment, 17% of mice experienced no seizures at all during CP690550 treatment (Fig. 6F-G). Animals treated with vehicle throughout the experiment exhibited seizures consistently, contrasting with the profound reduction observed during CP690550 treatment (Fig. 6B, F-G).

Next, we continued exploration of CP690550's antiepileptogenic potential by inclusion of the systemic kainate model of epilepsy, as well as adopting video-based behavioral seizure monitoring. These experimental modifications aimed to generalize the efficacy of CP690550 across epilepsy models, minimize potential confounding effects introduced by headmount implantation, and prioritize clinically relevant seizure endpoints.

Fig 6. CP690550 profoundly suppresses spontaneous seizures in epileptic wildtype mice. **(A)** SE was induced via IHKA, and mice equipped with EEG headmounts. All under continuous EEG recording, vehicle (i.p.) was administered daily for one week followed by one week of daily vehicle or CP690550 treatment (i.p., 15mg/kg). **(B)** Vehicle-treated mice have spontaneous seizures across both weeks of the experiment. **(C)** Example power spectrum, trace, and seizure. **(D-E)** Grouped comparisons for seizure frequency and burden were displayed as a cumulative frequency distributions and violin plots. For cumulative graphs, bin centers are marked. 50% of vehicle-treated mice had up to 1.1 seizures per day on average, while 50% of CP-treated mice had up to 0.1 seizures per day. 50% of untreated mice showed seizure burdens of 99 per day, while 50% of treated mice showed burdens of up to 11 (Kolmogorov-Smirnov test; **Frequency:** D:0.9091, p=0.0003; **Burden:** D:0.7083, p=0.0162; n=5-12). **(F-G)** Heat maps display paired data for seizure frequency and burden. With the exception of one seizure observed on day 10, CP treatment eliminated seizures for the last four days of treatment.

452 introduced by headmount implantation, and prioritize clinically relevant seizure endpoints. Additionally,
453 expansion of the observation period by two weeks after drug washout provided insights into any potential
454 disease-modifying effects of CP690550 (Fig. 7A). Using a different model of epilepsy and method of seizure
455 detection, we showed that CP690550 substantially curtailed seizures.
456
During weeks 11-12 post-SE, 75% of vehicle-treated mice exhibited up to 1.6 seizures per day and seizure
burdens up to 10.7, while 75% of CP690550-treated animals had up to 0.05 seizures per day with seizure
burdens up to 0.5 (Fig. 7B-C) (Kolmogorov-Smirnov test; **Frequency:** D:0.8333, p<0.05, **Burden:** D:0.8333,
p<0.05). In paired analyses, vehicle-treated mice exhibited ongoing spontaneous seizure activity[60] (Fig. 7D,G),
while CP690550 markedly suppressed seizures (Fig. 7E,H). CP690550 eliminated seizures in 60% of mice
during treatment, and 40% of mice across both treatment and washout.

Fig 7. CP690550 profoundly suppresses spontaneous seizures in epileptic wildtype mice. **(A)** SE was induced via systemic kainate (SKA), and behavioral spontaneous seizure activity was monitored by video from weeks 8-14 post-SE. Vehicle (i.p.) was administered daily for three weeks followed by two weeks of daily vehicle or CP690550 treatment (i.p., 15mg/kg). Video recording continued for two weeks after drug washout. **(B-C)** Grouped comparisons for seizure frequency and burden were displayed as a cumulative frequency distributions and violin plots. For cumulative graphs, bin centers are marked. 75% of vehicle-treated mice had up to 1.6 seizures per day with daily burdens up to 10.7, while 75% of CP-treated mice had up to 0.05 seizures per day with daily burdens up to 0.5 (Kolmogorov-Smirnov test; **Frequency:** D:0.8333, p=0.015; **Burden:** D:0.8333, p=0.026; n=6-7). Heat maps display paired data for seizure frequency **(D-E)** and seizure burden **(G-H)**. CP690550 treatment eliminated seizures in 40% of mice during weeks 11-12 post-SE. Seizure frequency **(F)** and seizure burden **(I)** before treatment and after washout for vehicle- and CP690550-treated animals. Vehicle-treated mice show no robust change in seizure frequency (p=0.106) or burden (p=0.115) from before treatment to washout, while CP690550-treated mice exhibit an ongoing 83% reduction in frequency (p=0.005) and 88% reduction in burden (p=0.009) after drug is withdrawn. Compared to vehicle-treated mice during washout, this represents an 87% reduction in frequency (p=0.006) and 81% reduction in burden (p=0.002) (2-way ANOVA; n=5).

Drug washout occurred over the weekend following week 12 post-SE[59]. Mice treated with CP690550 continued to show profound seizure reduction for at least 2 weeks post-washout compared to vehicle-treated mice (2-way ANOVA; **Frequency:** 87% reduction, p<0.01, **Burden:** 81% reduction, p<0.01). This data supports the hypothesis that CP690550 exerts robust disease modification in a preclinical rodent model of epilepsy.

DISCUSSION

We show that CP690550 (Tofacitinib), an FDA approved, orally available drug with no known behavioral side effects, exerts a potent and enduring seizure-suppressive effect when administered transiently across rodent preclinical epilepsy models. We demonstrate that CP690550 shuts down a pro-epileptic gene network centered around STAT3 by inhibition of JAK1, an endogenous gene target of the protective chromatin modifier EZH2. We provide evidence that the STAT3 upstream kinase JAK1 is a novel therapeutic target for seizure suppression and disease modification across epilepsy models.

After status epilepticus, we find EZH2 protein is potently induced in hippocampal pyramidal and dentate granule neurons across models of epilepsy[23]. Previously, we reported that treatment of mice with the SAM-competitive EZH2 inhibitor UNC-1999 exacerbated spontaneous seizures, and here we find that deletion of EZH2 in mature neurons has the same effect. Induction of Polycomb group (PcG) proteins has been studied in association with other brain insults. For example, sublethal ischemic insults lead to induction of PcG proteins in mouse cortical neurons, and *in vitro* overexpression of PRC1 component BMI1 was sufficient to protect against oxygen-glucose deprivation-associated injury[61]. Further, sevoflurane (a volatile anesthetic that exerts positive allosteric modulation on GABA_A receptors[62] and

antagonizes NMDA receptors[63]) mediates the protective induction of EZH2 and H3K27Me3 to attenuate hypoxic-ischemic brain injury, reducing hippocampal autophagy and improving post-injury behavioral

assessments in rats. Intriguingly, these improvements are reversed when rats treated with sevoflurane also receive a dose of the EZH2 inhibitor GSK126[64]. These studies, along with our findings, point to a potentially general protective role for EZH2 in brain insults.

EZH2 deletion in neurons results in large transcriptomic changes that manifest after SE (Fig. 2). The mouse Black gene cluster houses genes whose expression are relatively unchanged in naïve EZH2nWT and EZH2nHom animals, induced after SE in EZH2nWT, and hyper-induced in EZH2nHom mice after SE. This pattern suggests that EZH2 might be responsible for tempering expression of genes in this cluster because its deletion results in their unfettered activation. The Black cluster ontological analysis produces terms associated with innate immunity, neuroinflammation, and the JAK/STAT pathway. Network analysis places STAT3 in the center of the Black GRN, and immunofluorescence confirms induction of STAT3 in neurons post SE. While activation of the JAK/STAT pathway in non-neuronal cells post-SE is well-characterized[26,47,65,66], the role of STAT3 in neurons, especially in epilepsy, has been less studied; thus the mechanisms of neurogenic inflammation post-SE are not clear[45]. Cytokines released from glial cells following seizures increase neuronal excitability[67], and type I IFNs increase excitability in neocortical pyramidal neurons via PKC[68]. Others report that seizure-induced activation of the JAK/STAT pathway reduces the number of gamma-aminobutyric acid (GABA) type A alpha1 receptors expressed in neurons following seizures, influencing GABA(A)R-dependent inhibition[27]; this group has also utilized pharmacological inhibition of STAT3[32] and genetic ablation of the STAT3 SH2 domain[45] to reduce spontaneous seizure frequency in rodent models of epilepsy. Thus, activation of neuroinflammatory pathways, including the JAK/STAT pathway, has been shown to modulate neuronal excitability. However, our work is the first to identify an endogenous protective mechanism induced by SE that works to temper JAK/STAT activation and associated neurogenic inflammation. Our's is also the first to demonstrate an enduring protective effect of targeting this pathway in chronic epilepsy after the onset of spontaneous seizures.

In order to understand the mechanism by which EZH2 suppresses STAT3 activation, and to rule out the possibility that STAT3 hyper-induction in EZH2 mutant mice might be a consequence of worsened disease progression more generally, rather than loss of EZH2 *per se*, we turned to an *in vitro* cell line model. Knocking down EZH2 in N2A cells results in induction of STAT3 Y705 phosphorylation. This activation is not via secretion of soluble factors as swapping media from control and knockdown cells fails to activate STAT3 (data not shown). WP1066, a JAK2 inhibitor[51-53] fails to suppress activated STAT3 in shEZH2 cells whereas CP690550 (a JAK1/3 inhibitor) (Fig. 4B-C) and direct knockdown of JAK1 (Fig. 4H-J) succeeds. Given that JAK3 is not expressed in these cells, we conclude that EZH2 tempers JAK1. Finally, we find that EZH2 directly interacts with the JAK1 locus (Fig. 4G). Increased activation of STAT3 at pY705-STAT3 as a result of EZH2 inactivation has also been observed in a transformed murine cell model of early T cell precursor acute lymphoblastic leukemia; EZH2 inactivation leads to enrichment of documented STAT3 target genes, increased expression of IL6RA, and an exaggerated pY705-STAT3 induction in response to IL6 stimulation[69]. STAT-upstream receptors and kinases are also regulated by EZH2. In prostate cancer EZH2 directly targets the interferon gamma receptor gene INFGR1 to dampen JAK2/STAT1 signaling, and EZH2 depletion restores expression of IFN target genes as well as STAT1 phosphorylation.[70]

Others have identified EZH2 as an activator of JAK/STAT signaling as well as a target of the pathway. When phosphorylated by AKT at serine 21, EZH2 disassociates from Polycomb to enhance STAT3 transcriptional activity through di-methylation of STAT3-K49 in glioblastoma cells[71-73]; we have performed western blotting to assess this posttranslational mark in mouse hippocampi post-SE and find no change in STAT3-K49Me2 after seizures or deletion of EZH2 (data not shown). Thus, this mechanism does not appear to play a role in N2As or the mouse brain. In NK/T-cell lymphoma, JAK3 phosphorylates EZH2 to mediate a switch in its function, promoting the transcriptional upregulation of oncogenes through the interaction of EZH2 and Pol-II[74]. To our knowledge, ours is the first report of EZH2 repressing JAK/STAT activation through JAK1.

Analysis of human TLE samples identifies a cluster of genes strikingly similar to the STAT3-driven mouse Black cluster; the human Brown cluster is a group of genes associated with innate immunity and neuroinflammation. Like the mouse Black cluster, the most influential driver of gene expression is STAT3. This is a possible explanation for why CP690550 suppresses disease whether it is administered shortly after SE (Fig. 5), or after the onset of spontaneous seizures (Fig. 6). The robust overlap between clusters of genes, one from frankly epileptic human samples and another from rodents, suggests a couple of points. Firstly, EZH2-

575 STAT3 mediated mechanisms identified in mice are clinically relevant. Secondly, the EZH2-STAT3 mechanism
576 evoked acutely after insult plays a role later in chronic epilepsy; this is supported by the identification of STAT3
577 as a driver of gene expression, its transient activation acutely post-SE, and its reactivation in chronic rodent
578 epilepsy (see Fig. 3).
579
EZH2nKO mice show both a decreased seizure threshold 4 days post SE as well as exaggerated seizure
burden 5 weeks later compared to WT mice. CP690550 administration for 24 hours post-SE is sufficient to
rescue both these phenotypes. This suggests that STAT3 hyper-induction drives seizure threshold reductions
and spontaneous seizure increases in mutant mice. The observation that CP690550 profoundly suppresses
spontaneous seizures when administered to EZH2nHom mice during the chronic period is consistent with
reactivation of JAK/STAT signaling during the chronic period (Fig. 3E-G, 5F-G, 6C-K). This supports the
possibility that mechanisms involved in disease initiation (e.g., activation of neuroinflammatory pathways)
continue to perpetuate disease chronically. Leveraging the brain's own protective mechanism, we use
CP690550 to buttress EZH2 suppression of JAK1. CP690550 profoundly suppresses seizures in WT epileptic
mice across models of epilepsy. Following cessation of CP690550 treatment we observe robust disease
modification for at least two weeks. Whether this disease-modifying effect is permanent will be the focus of
future studies in the lab.
592
Anti-inflammatory drugs that have been used to suppress seizures in human patients[28-31] or have shown
antiepileptogenic potential[32-35] in rodents pose several drawbacks: some or all are proposed only as
supplemental therapies[28,29], risk worsened disease outcomes[29], show only transient effects[30], or are effective in
only a subset of epilepsy patients[29,31]. We argue that an approach involving multiple animal models and
multiple laboratories to identify drivers of disease increases the likelihood that druggable targets will be viable
across human etiologies. With this approach we identify EZH2 as a protective negative regulator of the
JAK/STAT pathway in epilepsy. Our current data is consistent with the hypothesis that CP690550 mitigates
spontaneous seizures in chronic epilepsy and is disease-modifying. The cross-model approach by which we
identified JAK1 and selected CP690550 suggests this drug may be useful across multiple epilepsies.
602
603 Recently, others find that neuron-specific deletion of the STAT3 SH2 domain plateaus, but does not reduce,
604 seizure frequency in chronic epilepsy[45], whereas CP690550 treatment greatly reduces seizures. The
605 explanation for this difference may be that there is incomplete ablation of STAT3 in their model, or it may
606 suggest that the more robust therapeutic effect of CP690550 is attributable to JAK/STAT pathway suppression
607 in non-neuronal cells (e.g., peripheral immune cells, microglia, etc.) as well as neurons.
608
The JAK/STAT pathway is likely one of many processes orchestrating disease progression. The data
presented here represents the first evidence that EZH2 regulates JAK/STAT function through JAK1/STAT3 in
epileptogenesis to suppress disease progression. Transient treatment with CP690550 not only mitigates
exaggerated disease in EZH2 mutant mice, but is seizure suppressing and chronically disease modifying in
WT mice in two different epilepsy models. Together, these data suggest an ongoing interplay between EZH2
and the JAK/STAT pathway, starting in early epileptogenesis and persisting in chronic epilepsy.
615
616 We conclude that EZH2 tempers the JAK/STAT pathway in epilepsy as a protective response but is ultimately
617 overwhelmed. Buttressing this protective program with the FDA approved, orally available drug CP690550
618 thwarts disease progression and offers a novel therapeutic approach to treating epilepsy.
619
620 **METHODS**
621
622 **Bioinformatics**
623
Reads were mapped to the mm39 build mouse genome and subjected to Likelihood Ratio Tests using
Deseq2[75]. Transcripts per million were calculated and genes were classified as expressed if they had at least 3
TPM in all samples of any one of the 4 conditions (saline:WT, kainate:WT, saline:EZH2nHom,
kainate:EZH2nHom). Deseq2 output was then filtered for only expressed genes and adjusted p-values were
recalculated for the reduced number of genes. Genes with an adjusted p<.01 and 1.2-fold change were
subjected to Leiden clustering with resolution=0.5 using the scanpy Leiden tool[76]. Ontology was performed on
all clusters using Ontomancer (manuscript in preparation). Briefly, Biocarta, Hallmark, KEGG, and Reactome

.gmt files were downloaded from https://www.gsea-msigdb.org/gsea/downloads.jsp. Gene sets associated with
every term in the databases were collected. For each Leiden cluster, Scores (S_t) for each term, t, were
calculated as:

$$S_t = \sqrt{(\log O_t)^2 + (-\log P_t)^2}$$

Where O_t and P_t, are the odds ratio and p values from Fisher exact tests for the overlap of Leiden cluster genes
and genes for each term t. Only terms with odds ratios>1 and p values<0.05 were kept for further analysis. For
each term, a gene list is generated containing the overlap of genes in the cluster and term, G_t. A term x term
similarity matrix was generated where similarity between term i and j is the Fisher exact test log odds ratio $O_{i,j}$
for the intersection of genes in the terms G_i and G_j. This was converted to a network where nodes are terms.
Edge weights were calculated as:

$$weight = (\log (O_{i,j}) / \max \log O)^6$$

Where max log odds is the maximum log odds value across all comparisons. The 6 power was used to separate
nodes for visualization. Networks were generated using the scanpy Leiden tool with resolution=2. Each community
was named after the highest scoring term in that community. Networks were visualized via Python's networkx and
Plotly tools.

MAGIC was performed on all clusters essentially as described[21] but with r=1, and GTFs were ignored. GTFs were
defined as any ENCODE Factors containing the strings "TBP", "POL", "GTF", or "TAF".

Gene Regulatory Network was projected using Cytoscape[77] using MAGIC outputs. Factors were connected to their
targets with edges defined as ChIP signal extracted from the Target Folder generated by MAGIC. Target genes of
Factors that were not themselves Factors were removed to generate the GRN. Network statistics were calculated in
Cytoscape[77].

Analysis of human TLE expression data.

GSE63808[50] from GEO contained 129 transcriptomes from resected hippocampal TLE tissue. Probes with an
Illumina p-value<0.05 in at least 20% of samples were retained in the analysis; the probes were consolidated
to gene symbols by taking the median probe expression value per symbol. These 9076 genes formed the
background list for ontological and MAGIC analyses. Expression values were normalized by the median
expression value in that sample. Genes kept for further analysis exhibited variance greater that the median
coefficient of variance across all samples. The resulting 7658 unique genes were subjected to k-means
clustering using the k-means tools from Sci-kit learn[78] to divide genes into 10 clusters (brown, blue, purple,
skyblue, grey, orange, indigo, green, yellow, red).

Overlap between each of the 10 human TLE clusters with mouse Black cluster genes was measured by Fisher
exact test with the Benjamini-Hochberg correction for multiple comparisons. Ontological and MAGIC analyses
were performed on the highest-scoring human Brown cluster as described in the previous section.

Cell culture.

Growth conditions. Cells were grown in 5% CO_2 at 37°C. Neuro2a cells were grown in MEM (#10-009-CV
Corning, Manassas, VA) with 1.0g/L glucose, 1.5g/L L-Glutamine, and 10% fetal bovine serum (#26140-079
Gibco, Grand Island, NY).

Lentiviral and siRNA knockdown. Stable EZH2 knockdown in Neuro2a cells for western blot and RNA
analysis was achieved using SMARTvector lentiviral delivery of shRNA per the manufacturer instructions.
Puromycin selection began 48 hours after infection and maintained during cell expansion and experimentation.
SMARTvector lentiviral mouse EZH2 mEF1a-TurboRFP shRNA (#V3SM7592-232015353, Dharmacon,
Lafayette, CO) targeted the sequence *ATCGTAGTAAGTACCAATG*, and non-targeting mEF1a-TurboRFP
control particles (#S10-005000-01) were used as an infection control. Transient JAK1 knockdown in Neuro2a

cells was accomplished using ON-TARGETplus Human JAK1 siRNA (#L-003145-00-0005, Dharmacon). Control cells were transfected with the ON-TARGETplus Non-targeting control pool (#D-001810-10-05, Dharmacon). Transfections were performed with RNAiMAX lipofectamine transfection reagent (#13778030, Invitrogen).

PRC2 and JAK1 overexpression. For overexpression of Polycomb, all plasmids were ordered and modified (if necessary) from Addgene (Watertown, MA). Control cells were transfected with pAdtrack-CMV (#16405). To overexpress PRC2, cells were transfected with pCMVHA-hEZH2 (#24230), pCMVHA-EED (#24231), and pCMVHA-SUZ12 (#24232). The EZH2 plasmid was modified in-house to produce an HA-EZH2-GFP fusion protein. To overexpress JAK1, cells were transfected with FLAG-JAK1 (#174572), and controls were transfected with pAdtrack-CMV. All transfections were carried out with Lipofectamine LTX reagent with PLUS reagent (#15338100, Invitrogen).

Animal care

All animal procedures and experiments were performed with approval from the University of Wisconsin-Madison School of Medicine and Public Health Instructional Animal Care and Use Committee, or the Institutional Animal Care and Use Committee of Emory University, as appropriate, and according to NIH national guidelines and policies.

Transgenic mice. Cre-driver and EZH2 floxed mice were ordered from Jackson Laboratories (#003966, #022616, Bar Harbor, ME). Mice were bred on a C57BL6 background. Using the previously described Synapsin1-Cre system[37] $EZH2^{-/-}$; $Syn1Cre^{+/-}$ (EZH2nHom) mice were generated by mating $EZH2^{flox/flox}$; $Syn1Cre^{+/-}$ females with $EZH2^{flox/flox}$ males. $EZH2^{+/-}$; $Syn1Cre^{+/-}$ (EZH2nHet) mice were generated by mating $EZH2^{flox/+}$; $Syn1Cre^{+/-}$ females with $EZH2^{flox/flox}$ and $EZH2^{flox/+}$ males [36]. $EZH2^{+/+}$; $Syn1Cre^{+/-}$ (EZH2nWT) mice were generated from mating pairs containing an $EZH2^{flox/+}$ male. To assess Cre expression, we used a reporter mouse with a lox-P flanked STOP cassette preventing transcription of tdTomato (#007909 Jackson Laboratories) [79]. Reporter males were bred with $Syn1Cre^{+/-}$ females to generate Cre-positive animals and Cre-negative controls.

Repeated low-dose KA model. Male and female C57BL6 and FVB mice were bred and housed under a 12-hour light/dark cycle with access to food and water *ad libitum*. Mice were allowed to reach 4-6 weeks of age (C57BL6) 5-7 weeks of age (FVB) before undergoing experimentation. Kainate injections were performed at the same time of day (~10 A.M.), and mice were returned to home cages before the start of the dark cycle (~5 P.M.). Mice were housed with littermates for the duration of the experiment, with males and females separated at weaning.
We used a repeated low-dose kainate mouse model to induce SE. To begin, mice were weighed and singly housed in observation chambers for the duration of the injections. Mice were injected intraperitoneally (i.p.) with 5.0mg/kg synthetic kainic acid (KA) dissolved in 0.9% saline (#7065, Tocris Bioscience, Bristol, United Kingdom). At twenty-minute intervals, mice were given 5.0mg/kg injections of KA up to the third injection. The dosage was then reduced to 3.75mg/kg. Animals continued to receive 3.75mg/kg injections of KA every twenty minutes until each reached SE. The subsequent KA injection was skipped if an animal experienced two or more Class V seizures within a single twenty-minute interval. Injections resumed for the next round, unless the animal reached SE. During induction of SE, seizures were scored using a modified Racine Scale[80] where I = freezing, behavioral arrest, staring spells; II = head nodding and facial twitches; III = forelimb clonus, whole-body jerks or twitches; IV = rearing; V = rearing and falling; VI = violent running or jumping behavior. Animals were considered to have reached SE after experiencing at least five Class V or VI seizures within a 90-minute window. KA mice were observed for 1-2 hours after SE. Animals were injected with 0.9% saline (s.c.), and soft gel food was provided in home cages. In the days following injection, animals were weighed and injected with 0.9% saline (s.c.) if body weight decreased by more than 0.5g.

Pilocarpine model of status epilepticus. Adult C57BL6 mice (8-12 weeks old, Charles River) were injected with terbutaline (4 mg/kg ip, Sigma T2528) and methylscopolamine (4 mg/kg ip, Sigma S8502) to alleviate respiratory and cardiovascular effects of pilocarpine, then 30 min later received saline or pilocarpine (280 mg/kg i.p. as free base, Sigma P6503) as described[34]. After 1 hour of SE, seizures were interrupted by diazepam (10 mg/kg ip).

IHKA model of status epilepticus. The intrahippocampal kainic acid (IHKA) model was employed to generate chronically epileptic mice as previously described[81]. Kainic acid (100 nL of 20 mM; vIHKA) or vehicle (100 nL of sterile saline; vIHSA) was stereotaxically injected into the hippocampus using the following coordinate (from Bregma): 3.6 mm posterior, -2.8 mm lateral, and 2.8 mm depth. Only mice which experienced status epilepticus and went on to develop spontaneous recurrent seizures were used in the current study.

Fluorothyl and PTZ seizure threshold testing. Male and female C57BL6 mice were bred and housed under a 12-hour light/dark cycle with access to food and water *ad libitum*. Animals were allowed to reach 4-6 weeks of age before undergoing experimentation. Fluorothyl seizure threshold tests were performed at the same time of day (~10 A.M.), and mice were returned to home cages before 5 P.M. Mice were housed with littermates for the duration of the experiment, with males and females separated at weaning. All seizure threshold tests were video recorded, and seizure behavior was scored by a blinded observer. Two tests per mouse were performed: the first at one week prior to kainate injections and the second at four days post-SE.
Seizure threshold tests were performed on male and female EZH2nWT, EZH2nHet, and EZH2nHom mice using 100% bis(2,2,2-trifluoroethyl) ether (#287571, Sigma-Aldrich, St. Louis, MO). All tests were performed in a fume hood. Mice were placed in an airtight 10L. Plexiglas chamber (8.5"X10.75"X6.75"). Fluorothyl was infused into the chamber via a peristaltic pump at a rate of 40μL/minute onto a piece of Whatman filter paper placed at the top of the chamber. The time to generalized tonic clonic seizure (GTCS) was recorded from the start of fluorothyl infusion. Mice reached GTCS when they exhibited a complete loss of postural and motor control. After experiencing a generalized tonic clonic seizure, mice were rapidly removed from the chamber, and fluorothyl infusion was terminated. Mice were transferred to a recovery cage and kept isolated until motor control and normal behavior returned (~10 minutes). Mice were returned to home cages after recovery. PTZ was dissolved at 5mg/mL in 0.9% normal saline, and a single 45mg/kg dose was administered (i.p.). Time to generalized tonic clonic seizure was recorded.

Long-term video monitoring for spontaneous seizures. Male and female FVB mice were bred and housed under a 12-hour light/dark cycle with access to food and water *ad libitum*. Animals were allowed to reach 5-7 weeks of age before undergoing kainate injections. Starting at either week 5 or week 7 post-SE, video recording for spontaneous seizures was performed at the same time of day (~10 A.M.), and mice were returned to home cages before 5 P.M. Mice were singly housed in observation chambers for the duration of recording with access to food and water. During daily CP690550 treatment, mice were weighed and injected daily before recording began. Videos were reviewed and scored by an experimenter blinded to treatment and genotype using the aforementioned modified Racine scale. In the final experiment described by Figure 7, only Racine class III-VI seizures were recorded. For each mouse, daily seizure burden was calculated as: (Total # of seizures * Average seizure duration (s))/(Total recording hours).

CP690550 citrate treatment of KA mice. Following kainate injections, an independent researcher (not the experimenter) randomly assigned mice to drug and vehicle conditions, and prepared both solutions. Drug and vehicle solutions were coded as "A" or "B." CP690550 citrate (tofacitinib citrate) (#4556, Tocris Bioscience, Bristol, United Kingdom) was dissolved in a solution of 70% saline (0.9%)/15% ethanol/15% DMSO. The 70% saline/15% ethanol/15% DMSO solution was administered as vehicle. The independent researcher provided the blinded experimenter with the drug solutions and a list of all mice assigned to group A or B; this researcher did not participate in any seizure threshold testing or spontaneous seizure video scoring. CP690550 or vehicle was delivered to mice intraperitoneally at 6, 12, and 24 hours or 1, 6, and 24 hours post-SE. The drug doses for each of these time points, respectively, were 25mg/kg, 15mg/kg, and 15mg/kg.

Continuous EEG recording and seizure detection.
At the time of stereotaxic IHKA or IHSA injections, mice were implanted with chronic EEG recording headmounts (Pinnacle Technology, cat. #8201) which were affixed to the skull with 4 screws: 2 screws serving as EEG leads placed in the anterior and posterior cortex, 1 as a reference ground, and 1 as the animal ground. The headmounts were secured with dental cement. 24/7 EEG recordings were collected at 4 KHz using a 100X gain preamplifier high pass filtered at 1KHz (Pinnacle Technology, cat. #8292-SE) and tethered turnkey system (Pinnacle Technology, cat. #8200).
Seizures were detected using a custom analysis pipeline developed in-house in python (seizy; https://github.com/neurosimata/seizy) as previously described[81]. Briefly, EEG traces were automatically

detected using a thresholding method from traces that were downsampled (decimated) to 100 Hz and divided
into 5 second segments. Power (2-40 Hz) and line length were used to identify seizures. Seizures were only
included if two consecutive 5 second segments were classified as seizures (minimum 10 seconds length) and
were manually verified by an experienced investigator with expertise in electrographic seizure detection.
Seizure frequency was calculated as the # of seizures per day by measuring total number of seizures a mouse
had over a 24-hour period. The average seizure duration was calculated by taking the mean seizure duration
for each mouse in each recording period. For each mouse, seizure burden was calculated as: (Total # of
seizures * Average seizure duration)/(Total EEG recording hours).
806
807 **Tissue isolation and homogenization**
808
After KA seizure induction and/or treatment with CP690550 citrate, animals were sacrificed by rapid
decapitation. Whole hippocampal hemispheres were harvested and flash-frozen in liquid nitrogen or on dry ice.
Hippocampal tissue was lysed in Radioimmunoprecipitation Assay Buffer (RIPA: 50mM Tris, 150mM NaCl, 1%
nonidet P-40, 0.5% sodium deoxycholate, 0.1% SDS) with mammalian protease inhibitor (1:1000, Sigma or
1860932 ThermoScientific) and phosphatase inhibitor (1:1000, Sigma or 78428 ThermoScientific). Protein from
cell lines was harvested in Triton lysis buffer (3% 5M NaCl, 10% glycerol, 0.3% Triton X-100, 5% 1M Tris
pH8.0) after PBS washes. Tissue was homogenized in lysis buffer by probe sonication (Fisher Scientific, Sonic
Dismembrator, Model 100, Hampton, NH) on power 3 for two rounds, with 10 pulses per round. Supernatants
were isolated by centrifugation and quantified using the DC Protein Assay (Bio-Rad, Hercules, CA) or BCA
assay (ThermoScientific 23227). Protein extracts were stored at -80°C. 5x loading buffer (0.5mM Tris, 10%
SDS, 50% glycerol, 10mM EDTA, and 1% beta-mercaptoethanol) was added to each sample to reach a 1x
final concentration. Extracts in loading buffer were boiled at 95°C for 5 minutes and stored for up to one month
at -20°C until run on an acrylamide gel.
822
823 **Western blotting**
824
Protein extracts in loading buffer were loaded at 20µg per lane and resolved by electrophoresis in hand-poured
acrylamide gels with a 5% acrylamide stacking layer (125mM Tris pH6.8, 5% acrylamide, 0.01% ammonium
persulfate, 0.01% SDS, 0.01% TEMED) and an 8% acrylamide separating layer (375mM Tris pH8.8, 8%
acrylamide, 0.015% ammonium persulfate, 0.015% SDS, 0.08% TEMED). Gels were transferred to polyvinyl
difluoride membranes (PVDF; Millipore, Bedford, MA) using Tris-glycine transfer buffer (20mM Tris, 1.5M
glycine, 20% methanol). Membranes were blocked with 5% bovine serum albumin (Fisher Scientific, Fair
Lawn, NJ) diluted in low-salt Tris-buffered saline (w/w TBST; 20mM Tris pH7.5, 150mM NaCl, 0.1% Tween 20)
with mammalian protease inhibitor (1:2000) and phosphatase inhibitor (1:2000), or Odyssey blocking solution
(LI-COR, 927-60001) for 1 hour at room temperature. Primary antibodies were diluted in the same blocking
buffer and incubated with membranes overnight at 4°C. Antibodies include: EZH2 (1:1000, #5426 Cell
Signaling, Danvers, MA), pY705-STAT3 (1:2000, #9145 Cell Signaling), STAT3 (1:1000, #12640 Cell
Signaling), JAK2 (1:1000, #3230 Cell Signaling), JAK1 (1:1000, #3344S Cell Signaling), STAT1 (1:1000,
#14994 Cell Signaling), pS727-STAT1 (1:1000, #8826 Cell Signaling), STAT2 (1:1000, #72604 Cell Signaling),
pY1007/08-JAK2 (1:500, ab32101, Abcam, Cambridge, UK), Tubulin (1:1000, #T9026 Sigma-Aldrich), and
Actin (1:10,000, #691001 MP Biomedicals). The next day, the membranes were washed three times in 1X
TBST and incubated with horseradish peroxidase-conjugated goat-anti-mouse or -rabbit secondary antibodies
for one hour at room temperature (1:10,000, Invitrogen #31430, #31460, Rockford, IL). Membranes were
subsequently washed three times in TBST, and membranes were developed in SuperSignal West Femto ECL
reagent (ThermoFisher, Waltham, MA). Bands were imaged using a ChemiDoc-It Imaging System (UVP
Vision-Works, Upland, CA) and quantified using UVP Vision Works software or ImageJ. Band intensities were
graphed and analyzed using Prism 9 software (GraphPad Software, La Jolla, CA).
846
847 **RNA extraction and clean-up for whole hippocampal RNAseq**
848
Hippocampi were harvested from saline- and kainate-treated EZH2nWT and EZH2nHom mice on day 4 post-
SE (n=4-5 per treatment/genotype). RNA was extracted from hippocampal tissue with Trizol (#1559602, Life
Technologies, Carlsbad, CA), and concentration was determined by spectrometry. This was followed by a
treatment with DNase (#AM2238, Invitrogen). RNA clean-up was carried with the RNeasy Mini Kit (#74104,

Qiagen, Hilden, Germany). Purified samples were submitted to the Gene Expression core at the University of
Wisconsin-Madison Biotechnology Center for library preparation and sequencing (https://biotech.wisc.edu).
855
Chromatin immunoprecipitation and qPCR
857
4% formaldehyde was used to crosslink proteins to chromatin before neutralization of the reaction in 125mM
glycine, and washes in 1X PBS (pH 7.4). Chromatin was fragmented in Triton-X lysis buffer and 1:100
protease inhibitors; lysates were sonicated and centrifuged to remove cell debris. Protein concentration of the
supernatant was quantified with BioRad Bradford assay reagent (#5000006). Chromatin was incubated
overnight at 4 degrees Celsius with antibodies against H3K27Me3 (#9733 Cell Signaling) or rabbit IgG. The
next day, protein A/G magnetic beads (#88803, ThermoScientific) were blocked in BSA and salmon sperm
DNA while chromatin samples were centrifuged, after which chromatin-antibody complexes were added to the
beads to incubate. Washes were performed in a low-salt ChIP buffer (1% SDS, 1% Triton X-100, 0.4% 500mM
EDTA, 2% 1M Tris pH8.1, 3% 5M NaCl). Decrosslinking buffer (1% SDS, 100mM $NaHCO_3$, 200mM NaCl) was
then added to IgG samples, H3K27Me3 samples, and inputs; decrosslinking reactions were allowed to
incubate overnight in a 65 degree Celsius water bath. Phenol-choloform DNA extraction was performed the
next day. Quantitative PCR was performed with 2x SYBR green mix (#RR420L, Takara Bio, Otsu, Shiga,
Japan) and primers against JAK1, STAT3, JAK2, PAX2, and beta actin (primers listed at the end of the
methods section). Signal was calculated from standard curves generated by serial dilution of an input sample,
and enrichment values were expressed as fold over input. RNA for RT-PCR was extracted with Trizol, and
concentration was determined by spectrometry. PCR reactions were run with primers against JAK1 and
GAPDH. cDNA synthesis was run on the MJ Mini Thermocycler from Bio-Rad, and PCR reactions were run on
the MJ Research DNA Engine Opticon 2 Continuous Fluorescence Detector using Opticon Monitor software.
876
Immunofluorescence
878
Whole brains were harvested on day 1 or 4 post-SE, bisected sagitally, fixed in 4% paraformaldehyde,
dehydrated in a series of graded TBS-Tween/ethanol solutions, embedded in paraffin, sectioned at 5μm and
subjected to immunofluorescence for NeuN, GFAP, total STAT3, and DAPI. Heat induced antigen retrieval was
performed in a 10mM citric acid buffer (pH 6.0). After a blocking step, mouse anti-NeuN (1:200 ab104224
Abcam, or 1:500 MAB377 from Millipore), rabbit anti-STAT3 (1:500 #12640 Cell Signaling), or rabbit anti-EZH2
(1:50 #5246 Cell Signaling) primary antibodies were applied and incubated overnight at 4°C. Slides were
washed with PBS then incubated with goat-anti-mouse or goat-anti-rabbit Alexa Fluor 488 (ab150113 Abcam
or A11005, A11008 from InVitrogen, respectively) and goat-anti-rabbit Alexa Fluor 594 (ab150080 Abcam or
Jackson ImmunoResearch) secondary antibodies at a concentration of 1:200 for one or two hours at room
temperature, followed in some experiments by two-hour incubation at room temperature with Alexa Fluro 488-
conjugated mouse anti-GFAP antibody (1:200) (#53-9892-82 Invitrogen). Nuclear DNA was stained with DAPI
(300 nM) or a DAPI-containing mounting media (#H-1200 Vector Laboratories, Burlingame, CA). Slides were
imaged at 20-40x magnification using a fluorescent microscope (Nikon E600 microscope with fluorescence or
Zeiss Axio Observer A1 epifluorescence microscope). Images were captured with Zeiss Axiovision or Cellsens
software (Olympus Life Science Solutions, Center Valley, PA). Image analysis and processing were performed
in ImageJ (NIH, https://imagej.nih.gov/ij/).
895
Statistical analysis
897
898 For all statistical analysis – with the exception of RNAseq data – FDR < 0.05 corrected for multiple
899 comparisons was considered statistically significant. Data were graphed as mean ± standard deviation (SD) in
900 Prism 9 software (Graphpad Software, La Jolla, CA). Statistical tests were also performed in Prism. Western
901 blot, seizure threshold, and spontaneous seizure data was analyzed by 2-way or 1-way ANOVA with the
902 Benjamini-Hochberg FDR method for multiple comparisons, or by unpaired t-test where appropriate.
903
904 Primer List
905 mouse-JAK1-Site1 (ChIP):
906 (forward) *CTCTGCTCGCTTTCCCGGAGATGC*
907 (reverse) *CTCGGAGGCTGGAGCGTGACTTCT*
908

mouse-JAK1-Site2 (ChIP):
(forward) *TGACTCGCACCCAGCCTGAAATGG*
(reverse) *GCGGCCCAGTCACCCGAAGAAATA*

mouse-JAK1-Site3 (ChIP):
(forward) *TTCGCGGGGCAAGGGAAAGGATTC*
(reverse) *CCACCGCTCAGGGCTCTGGGTTTT*

mouse-JAK2 (ChIP):
(forward) *CGCCGCTGCGGGATGTGAGTGC*
(reverse) *GGAGGCGGGATCGCAGGGAGAACC*

mouse-STAT3 (ChIP):
(forward) *CGGGCGAGCGCTGTACTTCCTA*
(reverse) *AAGCGAGCCGTATCAGGGCATTTA*

mouse-PAX2 (ChIP):
(forward) *GCGGCGCTCAGTCGCGTTCATAG*
(reverse) *CTCCCCGGAGTCTGGGCAGTCCA*

mouse-JAK1 (RT-PCR):
(forward) *GCGCCCAGCCTCGCTCGTC*
(reverse) *AACGCCATGGCATTGCAGTCCTCT*

mouse-GAPDH (RT-PCR):
(forward) *ACGGCAAATTCAACGGCACAGTCA*
(reverse) *TGGGGGCATCGGCAGAAGG*

Funding. Supported by CURE (AR), Lily's Fund (AR), NIH grant 1R01NS108756 (AR, RD) and
R01NS105628, R01NS102937, and R21NS120868 (JM).

Acknowledgements. We thank Asheebo Rojas for providing rat brain tissues in the DFP model.

Conflict of Interest. JM serves on the Scientific Advisory Board and has a Sponsored Research Agreement
with SAGE Therapeutics for work unrelated to this project. AR and RD have no conflicts of interest to declare.

References

1   Hirtz, D. *et al.* How common are the "common" neurologic disorders? *Neurology* **68**, 326 (2007).
https://doi.org/10.1212/01.wnl.0000252807.38124.a3
2   Gavvala, J. R. & Schuele, S. U. New-Onset Seizure in Adults and Adolescents: A Review. *Jama* **316**, 2657 (2016).
https://doi.org/10.1001/jama.2016.18625
3   Sankaraneni, R. & Lachhwani, D. Antiepileptic drugs--a review. *Pediatric Annals* **44**, e36 (2015).
https://doi.org/10.3928/00904481-20150203-10
4   Meyer, D. M. *et al.* Anti-inflammatory activity and neutrophil reductions mediated by the JAK1/JAK3 inhibitor,
CP-690,550, in rat adjuvant-induced arthritis. *Journal of Inflammation (London, England)* **7** (2010).
https://doi.org/10.1186/1476-9255-7-41
5   Dowty, M. E. *et al.* Preclinical to clinical translation of tofacitinib, a Janus kinase inhibitor, in rheumatoid
arthritis. *The Journal of Pharmacology and Experimental Therapeutics* **348**, 165 (2014).
https://doi.org/10.1124/jpet.113.209304
6   French, J. A. *et al.* Characteristics of medial temporal lobe epilepsy: I. Results of history and physical
examination. *Annals of Neurology* **34**, 774 (1993). https://doi.org/10.1002/ana.410340604
7   Trinka, E. *et al.* A definition and classification of status epilepticus--Report of the ILAE Task Force on
Classification of Status Epilepticus. *Epilepsia* **56**, 1515 (2015). https://doi.org/10.1111/epi.13121

8　Hesdorffer, D. C., Logroscino, G., Cascino, G., Annegers, J. F. & Hauser, W. A. Risk of unprovoked seizure after
acute symptomatic seizure: effect of status epilepticus. *Annals of Neurology* **44**, 908 (1998).
https://doi.org/10.1002/ana.410440609
9　Garriga-Canut, M. *et al.* 2-deoxy-D-glucose reduces epilepsy progression by NRSF-CtBP-dependent metabolic
regulation of chromatin structure. *Nature Neuroscience* **9**, 1382-1387 (2006). https://doi.org/10.1038/nn1791
10　Stafstrom, C. E., Roopra, A. & Sutula, T. P. Seizure suppression via glycolysis inhibition with 2-deoxy-D-glucose
(2DG). *Epilepsia* **49 Suppl 8**, 97 (2008). https://doi.org/10.1111/j.1528-1167.2008.01848.x
11　Stafstrom, C. E. *et al.* Anticonvulsant and antiepileptic actions of 2-deoxy-D-glucose in epilepsy models. *Annals of
Neurology* **65**, 435 (2009). https://doi.org/10.1002/ana.21603
12　Vezzani, A., Friedman, A. & Dingledine, R. J. The role of inflammation in epileptogenesis. *Neuropharmacology* **69**,
16 (2013). https://doi.org/10.1016/j.neuropharm.2012.04.004
13　Rojas, A. *et al.* A Novel Second-Generation EP2 Receptor Antagonist Reduces Neuroinflammation and Gliosis
After Status Epilepticus in Rats. *Neurotherapeutics: The Journal of the American Society for Experimental
NeuroTherapeutics* (2021). https://doi.org/10.1007/s13311-020-00969-5
14　Williams-Karnesky, R. L. *et al.* Epigenetic changes induced by adenosine augmentation therapy prevent
epileptogenesis. *Journal of Clinical Investigation* **123**, 3552-3563 (2013). https://doi.org/10.1172/jci65636
15　Liu, G. *et al.* Transient inhibition of TrkB kinase after status epilepticus prevents development of temporal lobe
epilepsy. *Neuron* **79**, 31 (2013). https://doi.org/10.1016/j.neuron.2013.04.027
16　Landis, S. C. *et al.* A call for transparent reporting to optimize the predictive value of preclinical research. *Nature*
490, 187 (2012). https://doi.org/10.1038/nature11556
17　Becker, A. J. *et al.* Correlated stage- and subfield-associated hippocampal gene expression patterns in
experimental and human temporal lobe epilepsy. *The European Journal of Neuroscience* **18**, 2792 (2003).
https://doi.org/10.1111/j.1460-9568.2003.02993.x
18　Borges, K., Shaw, R. & Dingledine, R. Gene expression changes after seizure preconditioning in the three major
hippocampal cell layers. *Neurobiology of Disease* **26**, 66 (2007). https://doi.org/10.1016/j.nbd.2006.12.001
19　Hansen, K. F., Sakamoto, K., Pelz, C., Impey, S. & Obrietan, K. Profiling status epilepticus-induced changes in
hippocampal RNA expression using high-throughput RNA sequencing. *Scientific Reports* **4** (2014).
https://doi.org/10.1038/srep06930
20　McClelland, S. *et al.* The transcription factor NRSF contributes to epileptogenesis by selective repression of a
subset of target genes. *eLife* **3** (2014). https://doi.org/10.7554/eLife.01267
21　Roopra, A. MAGIC: A tool for predicting transcription factors and cofactors driving gene sets using ENCODE data.
PLoS computational biology **16** (2020). https://doi.org/10.1371/journal.pcbi.1007800
22　Dingledine, R. *et al.* Transcriptional profile of hippocampal dentate granule cells in four rat epilepsy models.
Scientific Data **4** (2017). https://doi.org/10.1038/sdata.2017.61
23　Khan, N. *et al.* A systems approach identifies Enhancer of Zeste Homolog 2 (EZH2) as a protective factor in
epilepsy. *Plos One* **14** (2019). https://doi.org/10.1371/journal.pone.0226733
24　Pfisterer, U. *et al.* Identification of epilepsy-associated neuronal subtypes and gene expression underlying
epileptogenesis. *Nature Communications* **11** (2020). https://doi.org/10.1038/s41467-020-18752-7
25　Cid, E. *et al.* Sublayer- and cell-type-specific neurodegenerative transcriptional trajectories in hippocampal
sclerosis. *Cell Reports* **35** (2021). https://doi.org/10.1016/j.celrep.2021.109229
26　Kumar, P. *et al.* Single-cell transcriptomics and surface epitope detection in human brain epileptic lesions
identifies pro-inflammatory signaling. *Nature Neuroscience* **25**, 956 (2022). https://doi.org/10.1038/s41593-022-
01095-5
27　Lund, I. V. *et al.* BDNF Selectively Regulates GABAA Receptor Transcription by Activation of the JAK/STAT
Pathway. *Science Signaling* **1**, ra9 (2008). https://doi.org/10.1126/scisignal.1162396
28　You, S. J., Jung, D. E., Kim, H. D., Lee, H. S. & Kang, H. C. Efficacy and prognosis of a short course of prednisolone
therapy for pediatric epilepsy. *European journal of paediatric neurology: EJPN: official journal of the European
Paediatric Neurology Society* **12**, 314 (2008). https://doi.org/10.1016/j.ejpn.2007.09.003
29　Dubey, D. *et al.* Randomized Placebo-Controlled Trial of Intravenous Immunoglobulin in Autoimmune
LGI1/CASPR2 Epilepsy. *Annals of Neurology* **87**, 313 (2020). https://doi.org/10.1002/ana.25655
30　Gayatri, N. A., Ferrie, C. D. & Cross, H. Corticosteroids including ACTH for childhood epilepsy other than epileptic
spasms. *The Cochrane Database of Systematic Reviews* (2007).
https://doi.org/10.1002/14651858.CD005222.pub2

31   Lai, Y.-C. *et al.* Anakinra usage in febrile infection related epilepsy syndrome: an international cohort. *Annals of*
Clinical and Translational Neurology **7**, 2467 (2020). https://doi.org/10.1002/acn3.51229

32   Grabenstatter, H. L. *et al.* The effect of STAT3 inhibition on status epilepticus and subsequent spontaneous
seizures in the pilocarpine model of acquired epilepsy. *Neurobiology of Disease* **62**, 73 (2014).
https://doi.org/10.1016/j.nbd.2013.09.003

33   Jiang, J. *et al.* Suppressing pro-inflammatory prostaglandin signaling attenuates excitotoxicity-associated
neuronal inflammation and injury. *Neuropharmacology* **149**, 149 (2019).
https://doi.org/10.1016/j.neuropharm.2019.02.011

34   Varvel, N. H. *et al.* Preclinical development of an EP2 antagonist for post-seizure cognitive deficits.
Neuropharmacology **224** (2023). https://doi.org/10.1016/j.neuropharm.2022.109356

35   Rojas, A., Ganesh, T., Lelutiu, N., Gueorguieva, P. & Dingledine, R. Inhibition of the prostaglandin EP2 receptor is
neuroprotective and accelerates functional recovery in a rat model of organophosphorus induced status
epilepticus. *Neuropharmacology* **93**, 15 (2015). https://doi.org/10.1016/j.neuropharm.2015.01.017

36   Shen, X. *et al.* EZH1 mediates methylation on histone H3 lysine 27 and complements EZH2 in maintaining stem
cell identity and executing pluripotency. *Molecular Cell* **32**, 491 (2008).
https://doi.org/10.1016/j.molcel.2008.10.016

37   Zhu, Y. *et al.* Ablation of NF1 function in neurons induces abnormal development of cerebral cortex and reactive
gliosis in the brain. *Genes & Development* **15**, 859 (2001). https://doi.org/10.1101/gad.862101

38   Basu, T. *et al.* Histone deacetylase inhibitors restore normal hippocampal synaptic plasticity and seizure
threshold in a mouse model of Tuberous Sclerosis Complex. *Scientific Reports* **9** (2019).
https://doi.org/10.1038/s41598-019-41744-7

39   Cilio, M. R. *et al.* Anticonvulsant action and long-term effects of gabapentin in the immature brain.
Neuropharmacology **40**, 139 (2001). https://doi.org/10.1016/s0028-3908(00)00103-9

40   Bierkamper, G. G. & Cenedella, R. J. Induction of chronic epileptiform activity in the rat by an inhibitor of
cholesterol synthesis, U18666A. *Brain Research* **150**, 343 (1978). https://doi.org/10.1016/0006-8993(78)90285-8

41   Liu, Z. & Holmes, G. L. Basic fibroblast growth factor is highly neuroprotective against seizure-induced long-term
behavioural deficits. *Neuroscience* **76**, 1129 (1997). https://doi.org/10.1016/s0306-4522(96)00412-5

42   Applegate, C. D. & Tecott, L. H. Global increases in seizure susceptibility in mice lacking 5-HT2C receptors: a
behavioral analysis. *Experimental Neurology* **154**, 522 (1998). https://doi.org/10.1006/exnr.1998.6901

43   Rho, J. M., Kim, D. W., Robbins, C. A., Anderson, G. D. & Schwartzkroin, P. A. Age-dependent differences in
flurothyl seizure sensitivity in mice treated with a ketogenic diet. *Epilepsy Research* **37**, 233 (1999).
https://doi.org/10.1016/s0920-1211(99)00068-6

44   Samoriski, G. M. & Applegate, C. D. Repeated generalized seizures induce time-dependent changes in the
behavioral seizure response independent of continued seizure induction. *The Journal of Neuroscience: The*
Official Journal of the Society for Neuroscience **17**, 5581 (1997).

45   Tipton, A. E. *et al.* Selective Neuronal Knockout of STAT3 Function Inhibits Epilepsy Progression, Improves
Cognition, and Restores Dysregulated Gene Networks in a Temporal Lobe Epilepsy Model. *Annals of Neurology*
(2023). https://doi.org/10.1002/ana.26644

46   Ahmed, M. M. *et al.* Altered Protein Profiles During Epileptogenesis in the Pilocarpine Mouse Model of Temporal
Lobe Epilepsy. *Frontiers in Neurology* **12** (2021). https://doi.org/10.3389/fneur.2021.654606

47   Vishwakarma, S., Singh, S. & Singh, T. G. Pharmacological modulation of cytokines correlating
neuroinflammatory cascades in epileptogenesis. *Molecular Biology Reports* **49**, 1437 (2022).
https://doi.org/10.1007/s11033-021-06896-8

48   Martín-Suárez, S., Cortes, J. M. & Bonifazi, P. Blockage of STAT3 during epileptogenesis prevents GABAergic loss
and imprinting of the epileptic state. *Brain: A Journal of Neurology* (2023).
https://doi.org/10.1093/brain/awad055

49   Freeman, L. C. Centrality in social networks conceptual clarification. *Social Networks* **1**, 215-239 (1978).

50   Johnson, M. R. *et al.* Systems genetics identifies Sestrin 3 as a regulator of a proconvulsant gene network in
human epileptic hippocampus. *Nature Communications* **6** (2015). https://doi.org/10.1038/ncomms7031

51   Kupferman, M. E. *et al.* Therapeutic suppression of constitutive and inducible JAK\STAT activation in head and
neck squamous cell carcinoma. *Journal of Experimental Therapeutics & Oncology* **8**, 117 (2009).

52 Hussain, S. F. *et al.* A Novel Small Molecule Inhibitor of Signal Transducers and Activators of Transcription 3
Reverses Immune Tolerance in Malignant Glioma Patients. *Cancer Research* **67**, 9630 (2007).
https://doi.org/10.1158/0008-5472.can-07-1243
53 Iwamaru, A. *et al.* A novel inhibitor of the STAT3 pathway induces apoptosis in malignant glioma cells both in
vitro and in vivo. *Oncogene* **26**, 2435 (2007). https://doi.org/10.1038/sj.onc.1210031
54 Changelian, P. S. *et al.* Prevention of organ allograft rejection by a specific Janus kinase 3 inhibitor. *Science (New
York, N.Y.)* **302**, 875 (2003). https://doi.org/10.1126/science.1087061
55 *UCSC Genome Browser*, <https://genome.ucsc.edu/index.html> (2022).
56 Konze, K. D. *et al.* An orally bioavailable chemical probe of the Lysine Methyltransferases EZH2 and EZH1. *ACS
chemical biology* **8**, 1324 (2013). https://doi.org/10.1021/cb400133j
57 Rocha, L. Subchronic treatment with antiepileptic drugs modifies pentylenetetrazol-induced seizures in mice: Its
correlation with benzodiazepine receptor binding. *Neuropsychiatric Disease and Treatment* **4**, 619 (2008).
https://doi.org/10.2147/ndt.s2118
58 Baradaran Rahimi, V., Askari, V. R., Hosseini, M., Yousefsani, B. S. & Sadeghnia, H. R. Anticonvulsant Activity of
Viola tricolor against Seizures Induced by Pentylenetetrazol and Maximal Electroshock in Mice. *Iranian Journal of
Medical Sciences* **44**, 220 (2019).
59 Cohen, S., Zwillich, S. H., Chow, V., Labadie, R. R. & Wilkinson, B. Co-administration of the JAK inhibitor CP-
690,550 and methotrexate is well tolerated in patients with rheumatoid arthritis without need for dose
adjustment. *British Journal of Clinical Pharmacology* **69**, 143 (2010). https://doi.org/10.1111/j.1365-
2125.2009.03570.x
60 Maguire, J. Epileptogenesis: More Than Just the Latent Period. *Epilepsy Currents* **16**, 31 (2016).
https://doi.org/10.5698/1535-7597-16.1.31
61 Stapels, M. *et al.* Polycomb Group Proteins as Epigenetic Mediators of Neuroprotection in Ischemic Tolerance.
Science Signaling **3**, ra15 (2010). https://doi.org/10.1126/scisignal.2000502
62 Jenkins, A., Franks, N. P. & Lieb, W. R. Effects of temperature and volatile anesthetics on GABA(A) receptors.
Anesthesiology **90**, 484 (1999). https://doi.org/10.1097/00000542-199902000-00024
63 Brosnan, R. J. & Thiesen, R. Increased NMDA receptor inhibition at an increased Sevoflurane MAC. *BMC
anesthesiology* **12** (2012). https://doi.org/10.1186/1471-2253-12-9
64 Xue, H. *et al.* Sevoflurane post-conditioning alleviates neonatal rat hypoxic-ischemic cerebral injury via Ezh2-
regulated autophagy. *Drug Design, Development and Therapy* **13**, 1691 (2019).
https://doi.org/10.2147/dddt.s197325
65 Choi, J.-S. *et al.* Upregulation of gp130 and differential activation of STAT and p42/44 MAPK in the rat
hippocampus following kainic acid-induced seizures. *Molecular Brain Research* **119**, 10 (2003).
https://doi.org/10.1016/j.molbrainres.2003.08.010
66 Wang, H. *et al.* LncRNA-UCA1 inhibits the astrocyte activation in the temporal lobe epilepsy via regulating the
JAK/STAT signaling pathway. *Journal of Cellular Biochemistry* **121**, 4261 (2020).
https://doi.org/10.1002/jcb.29634
67 Iori, V., Frigerio, F. & Vezzani, A. Modulation of neuronal excitability by immune mediators in epilepsy. *Current
Opinion in Pharmacology* **26**, 118 (2016). https://doi.org/10.1016/j.coph.2015.11.002
68 Reetz, O., Stadler, K. & Strauss, U. Protein kinase C activation mediates interferon-β-induced neuronal
excitability changes in neocortical pyramidal neurons. *Journal of Neuroinflammation* **11** (2014).
https://doi.org/10.1186/s12974-014-0185-4
69 Danis, E. *et al.* Ezh2 Controls an Early Hematopoietic Program and Growth and Survival Signaling in Early T Cell
Precursor Acute Lymphoblastic Leukemia. *Cell Reports* **14**, 1953 (2016).
https://doi.org/10.1016/j.celrep.2016.01.064
70 Wee, Z. N. *et al.* EZH2-mediated inactivation of IFN-γ-JAK-STAT1 signaling is an effective therapeutic target in
MYC-driven prostate cancer. *Cell Reports* **8**, 204 (2014). https://doi.org/10.1016/j.celrep.2014.05.045
71 Kim, E. *et al.* Phosphorylation of EZH2 activates STAT3 signaling via STAT3 methylation and promotes
tumorigenicity of glioblastoma stem-like cells. *Cancer Cell* **23**, 839 (2013).
https://doi.org/10.1016/j.ccr.2013.04.008
72 Dasgupta, M., Dermawan, J. K. T., Willard, B. & Stark, G. R. STAT3-driven transcription depends upon the
dimethylation of K49 by EZH2. *Proceedings of the National Academy of Sciences of the United States of America*
112, 3985 (2015). https://doi.org/10.1073/pnas.1503152112

73   Cha, T.-L. *et al.* Akt-mediated phosphorylation of EZH2 suppresses methylation of lysine 27 in histone H3.
Science (New York, N.Y.) **310**, 306 (2005). https://doi.org/10.1126/science.1118947

74   Yan, J. *et al.* EZH2 phosphorylation by JAK3 mediates a switch to noncanonical function in natural killer/T-cell
lymphoma. *Blood* **128**, 948 (2016). https://doi.org/10.1182/blood-2016-01-690701

75   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with
DESeq2. *Genome Biology* **15** (2014). https://doi.org/10.1186/s13059-014-0550-8

76   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data analysis. *Genome*
Biology **19** (2018). https://doi.org/10.1186/s13059-017-1382-0

77   Assenov, Y., Ramírez, F., Schelhorn, S.-E., Lengauer, T. & Albrecht, M. Computing topological parameters of
biological networks. *Bioinformatics (Oxford, England)* **24**, 282 (2008).
https://doi.org/10.1093/bioinformatics/btm554

78   Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *Journal of Machine Learning Research* **12**, 2825
(2011).

79   Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization system for the whole mouse
brain. *Nature Neuroscience* **13**, 133 (2010). https://doi.org/10.1038/nn.2467

80   Racine, R. J. Modification of seizure activity by electrical stimulation. II. Motor seizure. *Electroencephalography*
and Clinical Neurophysiology **32**, 281 (1972). https://doi.org/10.1016/0013-4694(72)90177-0

81   Basu, T. *et al.* Hypothalamic-pituitary-adrenal axis dysfunction worsens epilepsy outcomes and increases SUDEP
risk. (Neuroscience, 2022).

1139

1140

1141

1142

SUPPLEMENTAL FIGURES

S1. Synapsin1-Cre expression and recombination occur in neurons, and conditional knockout diminishes EZH2 induction post-SE.

S2. Neuronal ablation of EZH2 does not alter latency to 1st Class V seizure during kainate injections.

S3. Ontology networks for the mouse-Black and human-Brown clusters.

S4. Transient induction of STAT3 in wildtype animals is observed across models of SE.

S5. Other STAT family proteins are hyper-induced in EZH2nKO mice.

S6. Western blot quantifications in shNT and shEZH2 N2A cells.

S7. JAK2 activity is not required for hyper-activation of STAT3 in the absence of EZH2.

S8. Western blot quantifications in N2A cells transfected with control plasmids or overexpressing JAK1.

S9. H3K27Me3 ChIP-PCR at JAK1 (Sites 1 and 3), JAK2, and STAT3.

S10. CP treatment window selection and target engagement.

Fig. S1. Synapsin1-Cre expression and recombination occur in neurons, and conditional knockout diminishes EZH2 induction post-SE. (A) Syn1Cre[+/-] driver mice were bred with TdTm reporter mice. Immunofluorescence was performed on hippocampal sections, probing with the neuronal marker NeuN. TdTm colocalizes with NeuN. **(B)** Representative western blot probed for EZH2 in EZH2nWT, EZH2nHet, and EZH2nHom mice. **(C)** Quantification of western blot in panel B (2-way ANOVA; n=5-7). **(D)** (Left) Merged immunofluorescence images of the dentate gyrus labeling DAPI (blue), EZH2 (red), and (right) EZH2 binary images generated after auto-thresholding. Images were captured at 20x magnification in KA-injected EZH2nWT and EZH2nHet mice 4 days post-SE (n=1). **(E)** Quantification of binary images in panel D as a cumulative frequency distribution (Kolmogorov-Smirvnov test; D:0.239, p<10[-4]).

Fig. S2. Neuronal ablation of EZH2 does not alter latency to 1st Class V seizure during kainate injections. (A) Deletion of one or both EZH2 alleles does not alter time to first class V seizure during induction of SE with kainate (1-way ANOVA; n=8-18). **(B-D)** There is no correlation between time to first class V seizure and time to generalized tonic clonic seizure post-SE during fluorothyl testing (Pearson correlation; **nWT**: R^2:0.016, p=0.633, **nHet**: R^2:0.261, p=0.195, **nHom**: R^2:0.152, p=0.209; n=8-15).

Fig. S3. Ontology networks for the mouse-Black and human-Brown clusters. (A) The 439 mouse-Black cluster genes and **(B)** 850 human-Brown cluster genes were compared to gene lists associated with terms in KEGG, Reactome, Biocarta, and Hallmark databases; terms with an odds ratio>1 and p<0.05 across all databases were sorted into ten communities via a network approach. Community names are the ontological term with the highest odds ratio in the community.

Fig S4. Transient induction of STAT3 in wildtype animals is observed across models of SE. **(A)** Hippocampi were collected from wildtype control and pilo-SE rats at 12, 24, 48, and 72h. post-SE and protein lysates were run on western blots for pY705-STAT3, total STAT3, and GAPDH. **(B)** pY705-STAT3 is robustly induced up to 48h. post-SE, but begins to diminish by 72h. post-SE (1-way ANOVA; n=4-8).

Fig S5. Other STAT family proteins are hyper-induced in EZH2nKO mice. **(A)** Representative western blots for total STAT2, total STAT1, phospho-STAT1 (S727) and beta-actin on day 4 post-SE. For all proteins quantified in panels B-D, deletion of an EZH2 allele in naïve animals does not alter protein levels. **(B)** Total STAT2 is robustly induced after SE in EZH2nHom mice only by 3.4-fold (p=0.0008), 2.6-fold greater than EZH2nWT mice post-SE (p=0.001) (2-way ANOVA; n=3). **(C)** Total STAT1 protein is induced by 3-fold (p=0.001) in EZH2nWT mice post-SE, and 5.7-fold (p<10⁻⁴) in EZH2nHom mice post-SE. Total STAT1 is hyper-induced in EZH2nHom mice following seizures by 1.5-fold (p=0.007) compared to EZH2nWT mice (2-way ANOVA; n=9). **(D)** pS727-STAT1 is not robustly altered across the four conditions, but trends toward a 2-fold (p=0.148) induction in EZH2nHom mice post-SE (2-way ANOVA; n=3).

Fig. S6. Western blot quantifications in shNT and shEZH2 N2A cells. **(A)** Protein was harvested from shNT and shEZH2 cells for western blot analysis. A 3.7-fold reduction in EZH2 protein in the knockdown cells was confirmed by western blot **(A)** (unpaired t-test; p<10⁻⁴; n=3). We observe a 2.4-fold induction of pY705-STAT3 **(B)** in shEZH2 cells (unpaired t-test; p<10⁻⁴; n=5-6). **(C)** Total STAT3 protein was induced by 2.1-fold in shEZH2 cells (unpaired t-test; p=0.004; n=6).

Fig. S7. JAK2 activity is not required for hyper-activation of STAT3 in the absence of EZH2. **(A)** shNT and shEZH2 cells were treated with the JAK2/STAT3 inhibitor WP1066 to a final concentration of 0 (DMSO), 3, or 10μM for one hour. Protein was harvested and subjected to western blot analysis. **(B)** Quantification of western blots in panel A. pY705-STAT3 is induced in vehicle-treated shEZH2 cells by 3-fold (p=0.005). WP1066 does not alter pY705 levels in control or knockdown cells at any concentration (2-way ANOVA; n=3). **(C)** WP1066 reduces pY1007/08-JAK2 in shNT cells by 1.4-fold for both concentrations (p=0.014), but no change was observed in knockdown cells (2-way ANOVA; n=3).

Fig. S8. Western blot quantifications in N2A cells transfected with control plasmids or overexpressing JAK1. N2A cells were transfected with either Adtrack plasmid (Con) or FLAG-JAK1 (JAK1 OE), and samples were subjected to western blotting. **(A)** A 5.5-fold induction of JAK1 upon overexpression was confirmed (unpaired t-test; p=0.0008; n=3). **(B)** pY705-STAT3 levels were induced by 6.7-fold in JAK1 OE cells (p=0.0004) in the absence of changes to total STAT3 (unpaired t-test; p=0.0004; n=3).

Fig. S9. H3K27Me3 ChIP-PCR at JAK1 (Sites 1 and 3), JAK2, and STAT3. **(A)** Image from the UCSC Genome Browser depicting putative EZH2 binding sites 1 and 2 on JAK1. Schematics representing all EZH2 binding sites on JAK1, JAK2, and STAT3. **(B)** Positive control PAX2. H3K27Me3 is enriched in both vehicle-treated (p=10⁻⁴) and GFP-transfected (p=0.008) N2A cells. Treatment with 10mM UNC1999 reduces H3K27Me3 enrichment (p=0.004), and overexpression of PRC2 increases H3K27Me3 enrichment (p=0.009) (1-way ANOVA; n=3-5). **(C)** Negative control Actin. H3K27Me3 is not enriched unless PRC2 is overexpressed (p=0.0116) (1-way ANOVA; n=3). **(D-G)** Parallel ChIP-qPCR experiments performed with antibodies against H3K27Me3, wherein N2A cells were treated with either vehicle or 10mM UNC1999 for 24 hours, or transfected with expression vectors for GFP-EZH2, SUZ12, and EED or GFP. qPCR was performed. Results were graphed as fold H3K27Me3 over input. At JAK1, Site 1 **(D)**, treatment with 10mM UNC1999 leads to a trends towards reduced H3K27Me3 enrichment (p=0.079), and overexpression of PRC2 increases H3K27Me3 enrichment (p=0.010) (unpaired t-test; n=3-4). At JAK1, Site 3 **(E)**, treatment with 10mM UNC1999 does not alter H3K27Me3 enrichment, and overexpression of PRC2 leads to a trend towards increased H3K27Me3 enrichment (p=0.059) (unpaired t-test; n=3-4). At JAK2 **(F)**, treatment with 10mM UNC1999 does not alter H3K27Me3 enrichment, and overexpression of PRC2 leads to a trend towards increased H3K27Me3 enrichment (p=0.080) (unpaired t-test; n=3-4). At STAT3 **(G)**, treatment with 10mM UNC1999 reduces H3K27Me3 enrichment (p=0.023), but overexpression of PRC2 does not alter H3K27Me3 enrichment (unpaired t-test; n=3-4).

Fig. S10. CP treatment window selection and target engagement. **(A-C)** Hippocampi from saline- or kainate-injected EZH2nWT and EZH2nHet mice were harvested on day 1 post-SE to quantify pY705-STAT3 induction. **(B)** All kainate animals exhibit robust induction of pY705 on day 1 post-SE ($p < 10^{-4}$), with EZH2nHet mice showing a 1.4-fold greater induction of pY705 over EZH2nWT mice ($p < 10^{-4}$) (2-way ANOVA; n=3). **(C)** Total STAT3 is not altered by EZH2 deletion or kainate injection (2-way ANOVA; n=3). **(D)** After induction of SE and CP treatment at 6, 12, and 24h. post-SE, a cohort of mice was sacrificed 3 hours after the last injection, and hippocampi were extracted to confirm drug target engagement via western blot. **(E)** Protein extracts from treated and untreated animals were run alongside extracts from naïve mice (lanes 1,4) to confirm the pY705 induction typically observed on day 1 post-SE. Drug-treated animals exhibited a 2.7-fold reduction in pY705 compared to vehicle-treated animals (unpaired t-test; p=0.004; n=3). **(F-G)** Naïve wildtype mice were pre-treated with CP690550 or vehicle (i.p., 15mg/kg) 30 minutes prior to threshold testing with flurothyl or PTZ (i.p., 45mg/kg). Time to generalized tonic clonic seizure was recorded. Results were analyzed by unpaired t-test. CP690550 pre-treatment does not alter **(F)** flurothyl (unpaired t-test; p=0.544; n=4-5) or **(G)** PTZ seizure threshold (unpaired t-test; p=0.236; n=5).