Profound seizure suppression and disease modification by targeting JAK1, a key driver of a pro-epileptogenic gene network.

Olivia R. Hoffman[1,2], Anna M. Patterson[1], Emily S. Gohar[1], Emanuel Coleman[3], Jose Ezekial Clemente Espina[4], Barry A. Schoenike[1], Claudia Espinosa-Garcia[4], Felipe Paredes[5], Raymond J. Dingledine[5], Jamie L. Maguire[3], Avtar S. Roopra[1,2]*

[1]Department of Neuroscience, University of Wisconsin-Madison School of Medicine and Public Health, Madison, WI 53705, USA; [2]Molecular and Cellular Pharmacology Graduate Program, University of Wisconsin-Madison School of Medicine and Public Health, Madison, WI 53705, USA; [3]Department of Neuroscience, Tufts University School of Medicine, Boston, MA 02111, USA; ; [4]Cellular and Molecular Biology Graduate Program, University of Wisconsin-Madison School of Medicine and Public Health, Madison, WI 53705, USA [5]Department of Pharmacology & Chemical Biology, Emory University School of Medicine, Atlanta, GA 30322, USA.

* asroopra@wisc.edu

ABSTRACT

Epilepsy is the 4th most prevalent neurological disorder with over 50 million cases worldwide[1,2]. While a number of drugs exist to suppress seizures, approximately 1/3 of patients remain drug resistant, and no current treatments are disease modifying[3]. Using network and systems-based approaches, we find that the histone methylase EZH2 suppresses epileptogenesis and slows disease progression, via repression of JAK1 and STAT3 signaling in hippocampal neurons. Pharmacological inhibition of JAK1 with the orally available, FDA-approved drug CP690550 (Tofacitinib)[4,5] profoundly suppresses behavioral and electrographic seizures after the onset of epilepsy across preclinical rodent models of acquired epilepsy. This seizure suppression persists for weeks after drug withdrawal. Identification of an endogenous protective response to status epilepticus in the form of EZH2 induction has highlighted a critical role for the JAK1 kinase and STAT3 in both the initiation and propagation of epilepsy across preclinical rodent models and human disease. Overall, we find that STAT3 is transiently activated after insult, reactivates with spontaneous seizures, and remains targetable for disease modification in chronic epilepsy.

INTRODUCTION

Epilepsy is a highly prevalent neurological disorder with over 50 million cases worldwide[1,2]. Arising from a myriad of etiologies including gene mutations and environmental insults, the unifying characteristic of the epilepsies is the emergence of recurring, spontaneous seizures. Epileptogenesis is the process that links initial insults to the eventual propagation of spontaneous seizures. One of the insults that may initiate epileptogenesis is status epilepticus: an unremitting bout of seizures lasting for more than five minutes, during which the patient does not regain consciousness[6-8]. The first-line treatments for epilepsy are anticonvulsants aimed at controlling seizures. While these medications effectively suppress seizures for some patients, they do not target the processes that orchestrate progression of the disease, and fail to suppress seizures altogether in approximately 1/3 of patients[3]. There is a critical need for treatments that target the progressive decline in seizure threshold that arguably leads to spontaneous seizures.

A poor understanding of the mechanisms underlying epileptogenesis has so far prevented the development of disease biomarkers and translation of antiepileptogenic treatments from pre-clinical rodent models to the clinic. Although others have reported the disease-modifying potential of manipulations to metabolism[9-11], inflammation[12,13], chromatin modification[14], and signal transduction pathways[15], progress has been hindered by a lack of consensus among laboratories employing different models of status epilepticus (SE)[16]. With few exceptions[9,17-20], little effort has been made to identify the principal transcription factors and chromatin modifiers whose function precipitates the gene changes that arise after status epilepticus.

Recently, we applied a systems approach and novel bioinformatic tool[21] to a transcriptomic consortium dataset that incorporated gene expression data across multiple laboratories and models of epilepsy[22]. This allowed for identification of potential mechanisms that span across rodent models of status epilepticus (SE) and human temporal lobe epilepsy (TLE)[23]. With this approach, we identified the histone methylase Enhancer of Zeste

Homolog 2 (EZH2) as a potential driver of gene changes in dentate granule cells post-SE that is induced by an order of magnitude across epilepsy models. Transient pharmacological inhibition of EZH2 post-SE worsened epileptic phenotypes one month later, suggesting that EZH2 induction plays a protective role in epileptogenesis. In addition to EZH2, our analysis predicted Signal Transducer and Activator of Transcription 3 (STAT3) to be a driver of gene activation across models of SE[22,23].

Of the studies that have profiled gene changes in the epileptic brain, nearly all find a robust activation of inflammatory pathways[17-20,23-25]. In the brain, inflammation can be triggered by glial cells[26], but neurons also exhibit an innate immune response that has been associated with hyper-excitability[27]. Anti-inflammatory drugs have been used with varying degrees of success to suppress seizures in human patients[28-31], and some drugs targeting neuroinflammation in rodents have shown antiepileptogenic potential[32-35]. However, the endogenous mechanisms that regulate neurogenic inflammation post-SE are poorly understood.

Herein, we show that EZH2 induction with SE suppresses pathological activation of pro-inflammatory JAK/STAT signaling in neurons, both in animal models and human disease. EZH2 deletion in neurons leads to hyper-activation of the JAK/STAT pathway, robust reductions in seizure threshold after SE, and an exaggerated epileptic phenotype. Pharmacological inhibition of JAK1/3 activity acutely after SE rescues these phenotypes. Finally, targeting STAT3 reactivation after the onset of epilepsy with the JAK inhibitor CP690550 results in an enduring, disease-modifying suppression of spontaneous seizures.

Fig. 1. EZH2 maintains seizure threshold after status epilepticus and mitigates spontaneous seizure frequency and severity. (A) Experimental paradigms for panels B-C and D-E. (B) Baseline fluorothyl seizure threshold data for C57BL6J EZH2nWT, nHet, and nHom mice. In naïve mice, deletion of EZH2 in neurons does not alter seizure threshold (one-way ANOVA; nWT:nHet p=0.471, nWT:nHom p=0.134, nHet:nHom p=0.471; n=8-16). (C) After induction of SE by KA injection, seizure threshold does not change robustly in EZH2nWT mice (p=0.061). EZH2nHet mice exhibit a 35% reduction in seizure threshold post-SE (p<10⁻⁴), with thresholds 25% lower than kainate-injected EZH2nWT mice (p=0.003). EZH2nHom mice show a 25% reduction in seizure threshold post-SE (p<10⁻⁴), with thresholds 17% lower than KA EZH2nWT mice (p=0.026) (2-way ANOVA; n=7-14). (D) FVB EZH2nWT and nHom mice were injected with KA to induce SE and video recorded from 5-7 weeks post-SE to count and score spontaneous seizure activity. Deletion of an EZH2 allele in neurons leads to a 2.5-fold increase in daily seizure events (unpaired t-test; p<10⁻⁴; n=6-10). (E) Total observed Racine scale events for EZH2nHom and EZH2nWT mice were quantified as a frequency distribution. 50% of EZH2nWT mice had only Class I seizures, while 50% of EZH2nHom mice had Class I and II seizures (Mann-Whitney test; U:284, p=0.002; n=6-10).

RESULTS

Neuronal EZH2 maintains seizure threshold and protects against exacerbated epileptic phenotypes.

We previously identified the histone methyltransferase EZH2 as a potential driver of gene repression post-SE, and EZH2 protein was shown to be potently induced over a 10-day window post-SE in rats and mice across models of epilepsy. EZH2 induction was protective because pharmacological inhibition resulted in accelerated disease progression[23]. To determine the role of EZH2 in epilepsy, we crossed EZH2 floxed mice[36] with

Syapsin1 Cre drivers[37] to delete EZH2 in mature neurons. Neuronal specificity of Cre expression was
confirmed via crossing Syn1Cre with *flox*-STOP-*flox*-tdTomato mice and immunofluorescence (Fig. S1A).

Immunofluorescence and western blot analysis confirmed successful ablation of neuronal EZH2 post-SE. Fig.
S1B-C show diminished induction of EZH2 on day 4 post-SE in mice lacking one EZH2 allele in neurons
(EZH2nHet) or lacking both alleles (EZH2nHom). Staining for EZH2 confirmed reduced induction of neuronal
EZH2 in EZH2nHet versus wildtype (EZH2nWT) mice (Fig. S1D, quantified in S1E).

We tested the impact of EZH2 deletion on fluorothyl seizure threshold acutely post-SE, as a correlation
between fluorothyl seizure threshold[38] and the development of spontaneous seizures has been suggested[39-42].
We measured time to generalized tonic clonic seizure (GTCS) after exposure to the volatile chemoconvulsant
fluorothyl in EZH2nWT, EZH2nHet, and EZH2nHom mice aged 4-6 weeks, before and after kainate-induced
SE[43,44] (Fig. 1A). Loss of EZH2 alleles in neurons had no impact on seizure threshold in naïve mice (Fig. 1B),
and induction of SE did not alter the seizure threshold of EZH2nWT mice on day 4 post-SE (Fig. 1C). In
contrast, robust reductions in threshold were detected after SE in EZH2nHet mice (2-way ANOVA; Mean
difference:55±8sec, $p<10^{-4}$) and EZH2nHom mice (Mean difference:46±8sec, $p<10^{-4}$). These were significant
reductions compared to EZH2nWT mice (EZH2nWT:EZH2nHet: 28%, $p<0.01$; EZH2nWT:EZH2nHom: 20%,
$p<0.05$). These data suggest that EZH2 is required to maintain fluorothyl seizure threshold post-SE.

We considered the possibility that EZH2 loss primed the mice to experience more severe responses to kainate
injection, and it was this vulnerability to SE driving the observed reduction in mutant seizure threshold.
However, we found no differences in time to first class V seizure and no correlation between time to first class
V seizure and time to GTCS post-SE for either EZH2nWT (Pearson correlation; R^2:0.016, $p=0.633$), EZH2nHet
(R^2:0.261, $p=0.195$), or EZH2nHom mice (R^2:0.152, $p=0.209$) (Fig. S2A-D).

To test the impact of neuronal EZH2 deletion on spontaneous seizures, we monitored behavioral spontaneous
seizures in FVB EZH2nWT and EZH2nHom mice by video weeks five to seven post-SE (Fig. 1A). EZH2nHom
mice exhibited a 2.5-fold increase in average daily seizure events compared to EZH2nWT mice (unpaired t-
test; $p<10^{-4}$) (Fig. 1D). EZH2nHom mice also exhibited more severe Racine scale seizures than EZH2nWT
mice (Mann Whitney test; U: 284, $p<0.01$) (Fig. 1E). These data support the hypothesis that neuronal EZH2
induction is a protective response post-SE.

EZH2 tempers induction of the JAK/STAT pathway post-SE in neurons.

To begin to understand the mechanism behind EZH2-mediated protection, we performed bulk RNAseq on
whole hippocampal tissue from naïve and epileptic (4d. post-SE) EZH2nWT and EZH2nHom mice. Of the
10556 expressed genes, 6241 were found to be differentially expressed at an adjusted $p<10^{-2}$ (Likelihood Ratio
Test) across the four conditions (saline, kainate, EZH2nWT, EZH2nHom). A community-based network
approach was used to group genes into clusters. Leiden clustering generated 12 distinct groups (Fig. 2A). The
grey, orange, gold, brown, skyblue, and purple clusters were generally repressed after SE, whereas the red,
blue, indigo, yellow, and black clusters were generally increased. We focused on the Black cluster because
these 439 genes manifested induction with SE in EZH2nWT mice but showed an exaggerated induction with
SE in EZH2nHom mice. We reasoned that these would be genes whose expression is induced with SE but
tempered in the presence of EZH2 post-SE, and undergo unfettered induction in the absence of EZH2.

To understand the nature of genes comprising the Black cluster, we performed an ontological analysis. The
439 Black cluster genes were compared to gene lists associated with terms in the KEGG, Reactome, Biocarta
and Hallmark databases. (Fig. 2B). The ontology network (see Fig. S3A) highlights the innate immune
response (NFkB, Interferon Gamma), microglial processes (Complement) and inflammation (IL6 Jak Stat3)
amongst others. The term 'IL6 Jak Stat3' was of particular interest given that our previous work highlighted
STAT3 as a driver of gene activation across models of epilepsy[23], and others have identified a role for the
JAK/STAT pathway in epileptogenesis[27,32,45-48].

To determine which transcription factors and cofactors (hereafter together referred to as 'Factors') control
expression of Black cluster genes, we used MAGIC[21]. STAT3 was the highest scoring Factor (Fig. 2D). When
projected as a Gene Regulatory Network (GRN) (Fig. 2C) such that an arrow projects from Factor X to Factor

Y if the gene encoding
Factor Y is a transcriptional
target of Factor X, STAT3
stood out as a central node.
Thus, STAT3 had the
highest in-degree, out-
degree and total edge count
(Fig. 2E) of any Factor in the
network. Finally, we
assessed the Betweenness
Centrality of the network[49] to
determine which Factors
might be critical
for information flow in the
GRN. STAT3 had by far the
largest Betweenness
Centrality of any Factor (Fig.
2F).

To ascertain whether the
mouse Black cluster has a
counterpart in human
epilepsy, we used whole
transcriptome data from 129
temporal lobectomy
resections collected ante-
mortem from human
temporal lobe epilepsy
(TLE) patients[23,50]. We used
k-means clustering to
generate 10 gene clusters
from the 7658 genes co-
expressed across all
samples (brown, blue,
purple, skyblue, grey,
orange, indigo, green,
yellow, red). To identify the
human TLE cluster most
similar to the mouse Black
cluster, we performed Fisher
exact tests to measure the
degree of overlap between
the 439 Black cluster genes,
and the genes comprising
each cluster of the human
TLE dataset (Fig. 2G). The
human Brown cluster
showed the greatest
overlap, sharing 65 genes in
common with the mouse
Black cluster (Fisher exact;
Odds: 4.79, p=1.19E-20).
We then performed

Fig. 2. STAT3 drives hyper-activation of pro-inflammatory gene networks post-SE in EZH2 knockout mice and human temporal lobe epilepsy. (A) Leiden clusters from 6241 differentially expressed genes (fold change >1.2x, adjusted p<0.01) across saline, kainate, EZH2nWT, and EZH2nHom conditions. (B) Ontological analysis of genes in the Black cluster highlights inflammatory processes. (C) MAGIC was used to determine which Factors control the expression of Black cluster genes, and positive hits were projected as a Gene Regulatory Network (GRN). (D) Top 5 Factors driving expression in the Black cluster sorted by –log(adjusted p value). (E, F) Factors were ranked by their in degree, out degree counts, and betweenness centrality. (G) Comparison of 10 human TLE k-means gene clusters to the mouse Black cluster by Fisher Exact test. Plotting -log(adjusted p) versus odds ratio of Fisher tests against the Black cluster shows that the Human Brown cluster is the most similar to the mouse Black cluster (Fisher exact; Odds:4.79, p(adj)=1.2x10[-20]). Dotted line shows p=0.05. (H) Ontological analysis of genes in the human-Brown cluster. The 850 human-Brown cluster genes were compared to gene lists associated with terms in KEGG, Reactome, Biocarta, and Hallmark databases (I) MAGIC GRN for Factors driving the human-Brown cluster. (J, K) Factors were ranked by their in degree, out degree, and betweenness centrality. (L) STAT3 is the top driver predicted by MAGIC to drive expression of genes common to the mouse-Black and human-Brown clusters.

ontological analysis on the genes in the human Brown cluster. The 850 genes were compared to gene lists
associated with terms in the KEGG, Reactome, Biocarta and Hallmark databases. (Fig. 2H). The ontology
network (see Fig. S3B) highlights inflammation (Cytokine Receptor Interaction/Complement/Innate Immune

System) and interferon signaling (Interferon Gamma Response). Mirroring results of the black cluster, STAT3
in the human Brown GRN had the highest in-degree, out-degree, and betweenness centrality of any Factor in
the network (Fig. 2J-K).
226
227 MAGIC called STAT3 as the highest scoring Factor driving expression of the 65 genes co-expressed in early
228 rodent epileptogenesis (mouse Black) and frankly epileptic human tissue (human Brown) (Fig. 2L). This data is
229 consistent with the hypothesis that STAT3-mediated disease mechanisms that are activated acutely in a rodent
230 model of epileptogenesis persist in chronic human epilepsy.
231
Transient activation of the JAK/STAT pathway after SE in WT mice has been reported by others[23,46] (and Fig.
S4A-B, S5A-D). To test for a neuronal EZH2-dependent change in JAK/STAT signaling, we performed western
blot analysis on hippocampal lysates from 4d. post-SE WT and EZH2 mutant mice (Fig. 3A-B). On day 4 post-
SE, pY705-STAT3 was induced 2-fold in EZH2nWT mice (p<0.01) compared to naïve controls, 2.9-fold in
EZH2nHet mice (p<10^{-4}), and 4.5-fold in EZH2nHom mice (2-way ANOVA; p<10^{-4}) (Fig. 3C-D). These results
demonstrated an EZH2 gene dose-dependent hyperactivation of STAT3.
238
239 Given the involvement of STAT3 in regulating gene expression across acute rodent SE and chronic human
240 epilepsy, we asked whether STAT3 reactivation occurred in chronic rodent epilepsy. We found that at 14
241 weeks post-SE, epileptic mice showed a 4.7-fold induction of pY705-STAT3 over naïve controls (unpaired t-

Fig. 3. JAK/STAT pathway components are hyper-activated in neurons of EZH2 knockout mice. EZH2nWT, nHet, and nHom mice were injected with saline or KA to induce SE; hippocampi were harvested on day 4 post-SE. **(A, B)** Representative western blots for pY705-STAT3, total STAT3, and beta-actin on day 4 post-SE. **(C)** pY705-STAT3 levels are unchanged across genotypes in naïve animals (p=0.220). pY705-STAT3 is induced by 2-fold in EZH2nWT mice (p=0.022), 2.9-fold in EZH2nHet mice (p<10^{-4}), and 4.5-fold in EZH2nHom mice post-SE (p<10^{-4}). pY705 levels in EZH2nHom mice post-SE are 2.6-fold greater than in EZH2nWT mice post-SE (p<10^{-4}) (2-way ANOVA; n=7-12). **(D)** Total STAT3 levels on day 4 post-SE. Total STAT3 is induced by 1.2-fold in nHom mice post-SE, compared to nWT mice post-SE (p=0.041). Total STAT3 is induced after seizures in nHet mice by 1.5-fold (p=0.041) (2-way ANOVA; n=7-12). **(E)** Representative western blots for pY705-STAT3, total STAT3, and beta actin in naïve and chronically epileptic mice at 14 weeks post-SE. **(F)** pY705-STAT3 is induced 4.7-fold in epileptic mice (unpaired t-test; p=0.007, n=5-7). **(G)** 20x images of the dentate gyrus (DG) and CA1 pyramidal cell layer (CA1). Hippocampal slices were collected on day 1 post-SE from saline- and KA-injected mice; immunofluorescence was performed to detect nuclear DNA, NeuN, and total STAT3.

test; p<0.01) (Fig. 3E-F).

To define the cell types in which STAT3 is expressed after seizures, we performed immunofluorescence with
DAPI, NeuN, and total STAT3 antibodies on day 1 post-SE hippocampi taken from mice injected with saline or
kainic acid (Fig 3G). We made use of the fact that the STAT3 gene is induced with STAT3 activation, and so
elevated total STAT3 protein was a proxy for activated STAT3. In both the dentate granule and CA1 pyramidal
cell layer, we observed a robust induction of total STAT3 in NeuN-positive cells.

250 EZH2 targets STAT3 via JAK1 to suppress JAK/STAT signaling.

252 STAT3 was a central factor in the Black cluster GRN whereas, surprisingly, EZH2 was not. In order to
253 understand how STAT3 hyperactivation could have occurred in the absence of EZH2, shRNA was used to
254 knock down EZH2 in the Neuro2A (N2A) neuronal cell line. We observed robust diminution of EZH2 in
255 knockdown cells (shEZH2) (Fig. 4A, quantified by S6A) (unpaired t-test; 3.7-fold, $p<10^{-4}$). This was associated

with a marked increase in pY705-STAT3 levels (unpaired t-test; 2.4-fold $p<10^{-4}$) (quantified by S6B). This result suggests that EZH2 suppresses STAT3 activation in a cell-autonomous manner without circuit level alterations.

To determine whether STAT3-upstream kinases are required for pY705-STAT3 induction in shEZH2 cell lines, we incubated cells with either WP1066 (an analogue of the JAK2 inhibitor AG490[51-53]) or CP690550 citrate (a small molecule inhibitor of JAK1/3 kinase activity[4,5,54]). WP1066 did not impact STAT3 phosphorylation in shEZH2 cells

Fig. 4. EZH2 suppresses STAT3 activation via transcriptional repression of JAK1. N2A cells were stably transfected with EZH2 shRNA (shEZH2) or non-targeting shRNA (shNT). **(A)** Knockdown of EZH2 causes induction of pY705-STAT3 and total STAT3 protein (quantified in Fig. S6). **(B,C)** pY705-STAT3 activation in shEZH2 cells is inhibited with the JAK inhibitor CP690550 citrate. pY705-STAT3 is induced in vehicle-treated shEZH2 cells by 3.7-fold (p=0.0004). CP690550 treatment reduces pY705-STAT3 levels of KD cells by 2.3-fold (at 0.5µM) and 3.2-fold (p=0.0006) (at 5µM). At 0.5µM CP, pY705 levels in KD cells are reduced to that of untreated control cells (p=0.18) (2-way ANOVA; n=3). **(D)** JAK1 mRNA expression is induced by 5.1-fold in shEZH2 cells (unpaired t-test; p=0.019; n=6). **(E)** JAK1 protein is induced 5.5 fold (p=0.0008), and pY705-STAT3 by 6.7-fold (p=0.0004) in the absence of changes to total STAT3 upon transfection with FLAG-JAK1 compared to control transfection (unpaired t-tests; n=3). **(F)** Schematics depicting EZH2 binding sites on JAK1. **(G)** H3K27Me3 is enriched at Site 2 at the JAK1 locus. Chromatin IP (ChIP) on N2A cells treated with either vehicle or 10µM UNC1999 for 24 hours, or transfected with expression vectors GFP (control) of cotransfected with GFP-EZH2, SUZ12, and EED (PRC2). Results are graphed as fold H3K27Me3 over input. For JAK1 (Site 2), treatment with UNC1999 reduces H3K27Me3 enrichment by 3.5-fold (p=0.029), and PRC2 overexpression leads to a 2.7-fold increase in H3K27Me3 enrichment (p=0.0006) (unpaired t-tests; n=3). **(H)** Knockdown of JAK1 suppresses pY705-STAT3 activation in shEZH2 cells. shNT and shEZH2 N2A cells transfected with either non-targeting (NT) or JAK1 siRNA were subjected to western blotting to confirm JAK1 and EZH2 knockdown, and assess pY705-STAT3 levels. **(I)** Total JAK1 protein is induced in shEZH2 cells transfected with non-targeting siRNA by 1.2-fold (p=0.023), and reduced in both shNT (p=0.0005) and shEZH2 ($p<10^{-4}$) after JAK1 RNAi (2-way ANOVA; n=9-12). **(J)** pY705-STAT3 is induced in shEZH2:NTsiRNA cells by 1.7-fold (p=0.027) over shNT:NTsiRNA cells. JAK1 RNAi in shEZH2 cells reduces pY705-STAT3 by 2.5-fold (p=0.006), to the levels of shNT:NTsiRNA cells (2-way ANOVA; n=9-12).

at any concentration tested (Fig. S7A-C). In contrast, 500nM CP690550 citrate resulted in a 2.3-fold (2-way
ANOVA; $p<0.01$) reduction of pY705-STAT3 in shEZH2 cells (Fig. 4B,C), rescuing pY705-STAT3 levels to that
of vehicle-treated control cells. Even greater reduction occurred in the shEZH2 cells treated with 5µM
CP690550 (3.2-fold, $p<10^{-3}$). CP690550 is a JAK inhibitor selective for JAK1 and JAK3, but JAK3 is not
expressed in neurons or N2As (data not shown). We therefore tested whether JAK1 was elevated in the
shEZH2 cells. Fig. 4D shows a 5.1-fold induction of JAK1 mRNA over control cells (unpaired t-test; $p<0.05$).
305

306   To determine whether induction of JAK1 is sufficient to induce STAT3 activation, we overexpressed JAK1 in
307   N2A cells (Fig. 4E). JAK1 shows a 5.6-fold induction upon overexpression (unpaired t-test; $p<10^{-3}$), and
308   pY705-STAT3 is induced by 6.7-fold (unpaired t-test; $p<10^{-3}$) in the absence of changes to total STAT3
309   (quantified in S8A-B). This suggests that CP690550 works to suppress STAT3 activation primarily through
310   JAK1 downstream of EZH2.
311

To understand how EZH2 regulates JAK1 activity, we noted three putative EZH2 binding sites in the gene body
and promoter region of JAK1 using ENCODE tracks in the UCSC Genome Browser[55] (Fig. 4F, Fig. S9A). We
performed chromatin immunoprecipitation with anti-H3K27Me3 antibodies to determine whether EZH2
methylase activity was enriched at any of these sites. In Fig. 4G H3K27Me3 was enriched at site 2. Incubation
of cells with UNC1999 (an EZH2 inhibitor[23,56]) for 24 hours resulted in a 3.5-fold loss of signal (unpaired t-test;
$p<0.05$). Overexpression of EZH2, SUZ12, and EED (the core components of Polycomb) increased
H3K27Me3 signal by 2.7-fold compared to overexpression of GFP as a control (unpaired t-test; $p<10^{-3}$). PAX2
and actin were immunoprecipitated as positive and negative controls for H3K27Me3 enrichment, respectively
(Fig. S9B-C); ChIP-PCR results for JAK1 (Site 1), JAK1 (Site 3), JAK2, and STAT3 were also collected (Fig.
S9D-G). None of these sites showed both a loss of H3K27Me3 enrichment after UNC1999 treatment and an
increase in signal upon overexpression of Polycomb. These data are consistent with the hypothesis that EZH2
targets JAK1 transcription to temper activation of STAT3.
324

Finally, to corroborate our pharmacological findings, we transfected control and shEZH2 cells with siRNA
against JAK1 or a non-targeting transcript, and validated knockdown of EZH2 and JAK1 by western blot (Fig.
4H). Total JAK1 protein is induced 1.2-fold upon EZH2 knockdown (2-way ANOVA; $p<0.05$). JAK1 RNA
interference reduced JAK1 protein levels in both control (2-way ANOVA; 1.6-fold, $p<10^{-3}$) and shEZH2 cells
(2.4-fold, $p<10^{-4}$) (Fig. 4I). Knockdown of JAK1 expression in shEZH2 cells was sufficient to rescue pY705-
STAT3 levels to that of control cells transfected with non-targeting siRNA (2-way ANOVA; 2.0-fold, $p<0.01$)
(Fig. 4J). These data support the hypothesis that CP690550 rescues STAT3 hyperactivation primarily through
targeting JAK1.
333

334   **Transient JAK inhibition acutely post-SE rescues epileptic phenotypes in EZH2 knockout**
335   **mice.**
336

To test whether loss of EZH2-mediated regulation of JAK1 contributed to the exacerbated disease progression
observed in EZH2 mutant mice, we treated EZH2nWT and EZH2nHet mice with CP690550 after
induction of SE. Because the hyper-induction of pY705-STAT3 in mutant mice occurred already at day 1 post
SE (Fig. S10A-C), we injected mice with CP690550 or vehicle at 6, 12, and 24 hours post-SE. A cohort of mice
was sacrificed three hours after final injection to assess hippocampal STAT3 activation by western blot to
confirm target engagement (Fig. S10D-E). Consistent with the previous result (see Fig. 1B), neuronal deletion
of EZH2 did not alter fluorothyl seizure threshold in naïve mice (Fig. 5B). CP690550 treatment raised the 4d.
post-SE seizure threshold of EZH2nHet mice by 30 seconds (1-way ANOVA; $p<0.05$), rendering them
indistinguishable from epileptic EZH2nWT mice (1-way ANOVA; $p>0.05$) (Fig. 5C). Thus, CP690550 rescued
the increased seizure susceptibility phenotype associated with the loss of neuronal EZH2. These data support
the hypothesis that EZH2 maintains seizure threshold post-SE by tempering activation of the JAK/STAT
pathway.
349

To test whether transient CP690550 treatment immediately after SE mitigated long-term disease progression,
mice were video recorded from weeks five to seven post-SE (Fig. 5A). CP690550 treatment immediately after
SE reduced daily seizure frequency by 41% in EZH2nHom mice 5 weeks later (1-way ANOVA; $p<10^{-4}$).
EZH2nHom seizure frequency was reduced from 2.6-fold above EZH2nWT to 1.5-fold (1-way ANOVA; $p<0.01$)
(Fig. 5D). EZH2nHom mice treated acutely with CP690550 also exhibited a dramatic reduction in seizure

severity from
untreated EZH2nHom
mice (Mann-Whitney
test; U:138.5,
p<0.01), such that
they were
indistinguishable from
WT mice (Fig. 5E).
363
To test whether
CP690550 could
rescue the increased
seizure burden of
EZH2nHom mice if
administered during
the chronic period,
we recorded
spontaneous
seizures for two
weeks post-SE in
EZH2nWT and nHom
mice starting 5 weeks
post SE, followed by
another 2-week
recording period
during which the
EZH2nHom mice
received daily
injections of
CP690550 (Fig. 5A).
Daily CP690550
treatment in the
chronic period
reduced seizure
severity and
frequency by 80% in
EZH2nHom mice (1-
way ANOVA; p<10^{-4})
(Fig. 5F-G). Indeed,
seizure frequency of
EZH2nHom+CP mice
was reduced to below
EZH2nWT levels (1-

Fig. 5. JAK inhibition rescues seizure threshold reductions and spontaneous seizure burden in EZH2nKO mice. (**A**) (For panels B-C) Naïve seizure threshold was measured in C57 EZH2nWT and nHet mice one week prior to KA injections. (**B**) Deletion of an EZH2 allele does not alter seizure threshold in naïve mice (unpaired t-test; p=0.246; n=24-29). (**C**) nHet mice were injected with vehicle or CP690550 at 6, 12, and 24h. post-SE. Seizure threshold was measured 4 days post-SE. Untreated nHet mice show a 35 second reduction in seizure threshold (p=0.007) compared to nWT mice. CP690550 raises seizure thresholds of nHet mice by 30 seconds (p=0.025), extinguishing the difference between treated nHet and nWT mice (p=0.709) (1-way ANOVA; n=7-9). (**D**) FVB EZH2nWT and nHom mice received CP690550 or no treatment at 1, 6, and 24h. post-SE, and SRS activity was monitored by video from 5-7 weeks post-SE (see panel A). nHom mice treated acutely 6,12 and 24 hours post-SE with CP690550 exhibit a 41% reduction in daily seizures compared to untreated nHom mice (p<10^{-4}); a 34% difference remains between treated nHom mice and nWT mice (p=0.0073) (1-way ANOVA; n=4-10). (**E**) Racine scale events represented as a frequency distribution. Acute CP690550 treatment reduces seizure severity in nHom mice (Mann Whitney; U:138.5, p=0.009; n=4-10) to nWT levels (U:357, p=0.955). 50% of nWT and treated nHom mice had Class I seizures, while 50% of untreated nHom mice had Class I and II seizures. (**F**) SRS activity was monitored from 5-7 weeks post-SE, after which nHom mice received daily 10mg/kg CP690550 injections during monitoring from 7-9 weeks post-SE (see panel A). CP treatment reduced daily seizures by 80% in nHom mice (p<10^{-4}); treated nHom mice exhibit daily seizure frequencies 58% lower than nWT mice (p=0.0008) (1-way ANOVA; n=6-10). (**G**) Total observed Racine scale events represented as a frequency distribution. Daily CP690550 reduces seizure severity in nHom mice. 90th percentile seizure severities of nHom mice before and after treatment: 5.3 and 1, respectively (Wilcoxon matched pairs signed rank; W:-36.00, p=0.007, n=6). 90th percentile severity of WT mice: 2.6 (Mann Whitney test comparing nWT and treated nHom mice; U:119, p=0.042; n=6-10).

way ANOVA; 1.9-fold, p<10^{-3}). Given the robust seizure suppression when administered during the chronic
period, we considered the possibility that CP690550 may simply be an anticonvulsant. To test this, we
measured seizure threshold using both flurothyl and PTZ (pentylenetetrazole)[57,58] in mice pre-treated with
vehicle or CP690550 (15mg/kg, i.p.) thirty minutes prior to testing. This time was chosen because peak plasma
levels are observed thirty minutes after oral administration[59] CP690550 pre-treatment had no effect on flurothyl
or PTZ seizure threshold (Fig. S10F-G).
404
405 **CP690550 profoundly suppresses electrographic and behavioral spontaneous seizures in**
406 **epileptic WT mice.**
407
408 CP690550 treatment during the chronic period reduced seizure frequency in EZH2nHom mice to below WT
409 levels, so we tested its effect on WT epileptic mice. Chronic administration of CP690550 following
410 intrahippocampal injection of kainate (IHKA) in WT epileptic mice significantly diminished spontaneous

recurrent seizures. In a paradigm involving one week of daily vehicle (i.p.) administration followed by an additional week of either vehicle or CP690550 injections (15mg/kg, i.p.) under continuous EEG recording (Fig. 6A), CP690550 treatment during the chronic period profoundly suppressed spontaneous seizures. Notably, 50% of vehicle-treated mice exhibited up to 1 seizure per day, with corresponding daily seizure burdens up to 99 (Fig. 6D-E). In contrast, 50% of CP690550-treated mice displayed up to 0.1 seizures per day, with daily seizure burdens up to 11. CP690550 treatment robustly reduced both spontaneous seizure frequency (Kolmogorov-Smirnov test; D:0.9091, p<10^{-3}) and burden (D:0.7083, p<0.01) in epileptic wildtype mice.

Additionally, while all animals experienced seizures during vehicle treatment, 17% of mice experienced no seizures at all during CP690550 treatment (Fig. 6F-G). Animals treated with vehicle throughout the experiment exhibited seizures consistently, contrasting with the profound reduction observed during CP690550 treatment (Fig. 6B, F-G).

Next, we continued exploration of CP690550's antiepileptogenic potential by inclusion of the systemic kainate model of epilepsy, as well as adopting video-based behavioral seizure monitoring. These experimental modifications aimed to generalize the efficacy of CP690550 across epilepsy models, minimize potential confounding effects

Fig 6. CP690550 profoundly suppresses spontaneous seizures in epileptic wildtype mice. **(A)** SE was induced via IHKA, and mice equipped with EEG headmounts. All under continuous EEG recording, vehicle (i.p.) was administered daily for one week followed by one week of daily vehicle or CP690550 treatment (i.p., 15mg/kg). **(B)** Vehicle-treated mice have spontaneous seizures across both weeks of the experiment. **(C)** Example power spectrum, trace, and seizure. **(D-E)** Grouped comparisons for seizure frequency and burden were displayed as a cumulative frequency distributions and violin plots. For cumulative graphs, bin centers are marked. 50% of vehicle-treated mice had up to 1.1 seizures per day on average, while 50% of CP-treated mice had up to 0.1 seizures per day. 50% of untreated mice showed seizure burdens of 99 per day, while 50% of treated mice showed burdens of up to 11 (Kolmogorov-Smirnov test; **Frequency:** D:0.9091, p=0.0003; **Burden:** D:0.7083, p=0.0162; n=5-12). **(F-G)** Heat maps display paired data for seizure frequency and burden. With the exception of one seizure observed on day 10, CP treatment eliminated seizures for the last four days of treatment.

introduced by headmount implantation, and prioritize clinically relevant seizure endpoints. Additionally, expansion of the observation period by two weeks after drug washout provided insights into any potential disease-modifying effects of CP690550 (Fig. 7A). Using a different model of epilepsy and method of seizure detection, we showed that CP690550 substantially curtailed seizures.

During weeks 11-12 post-SE, 75% of vehicle-treated mice exhibited up to 1.6 seizures per day and seizure burdens up to 10.7, while 75% of CP690550-treated animals had up to 0.05 seizures per day with seizure burdens up to 0.5 (Fig. 7B-C) (Kolmogorov-Smirnov test; **Frequency:** D:0.8333, p<0.05, **Burden:** D:0.8333, p<0.05). In paired analyses, vehicle-treated mice exhibited ongoing spontaneous seizure activity[60] (Fig. 7D,G), while CP690550 markedly suppressed seizures (Fig. 7E,H). CP690550 eliminated seizures in 60% of mice during treatment, and 40% of mice across both treatment and washout.

Fig 7. CP690550 profoundly suppresses spontaneous seizures in epileptic wildtype mice. **(A)** SE was induced via systemic kainate (SKA), and behavioral spontaneous seizure activity was monitored by video from weeks 8-14 post-SE. Vehicle (i.p.) was administered daily for three weeks followed by two weeks of daily vehicle or CP690550 treatment (i.p., 15mg/kg). Video recording continued for two weeks after drug washout. **(B-C)** Grouped comparisons for seizure frequency and burden were displayed as a cumulative frequency distributions and violin plots. For cumulative graphs, bin centers are marked. 75% of vehicle-treated mice had up to 1.6 seizures per day with daily burdens up to 10.7, while 75% of CP-treated mice had up to 0.05 seizures per day with daily burdens up to 0.5 (Kolmogorov-Smirnov test; **Frequency:** D:0.8333, p=0.015; **Burden:** D:0.8333, p=0.026; n=6-7). Heat maps display paired data for seizure frequency **(D-E)** and seizure burden **(G-H)**. CP690550 treatment eliminated seizures in 40% of mice during weeks 11-12 post-SE. Seizure frequency **(F)** and seizure burden **(I)** before treatment and after washout for vehicle- and CP690550-treated animals. Vehicle-treated mice show no robust change in seizure frequency (p=0.106) or burden (p=0.115) from before treatment to washout, while CP690550-treated mice exhibit an ongoing 83% reduction in frequency (p=0.005) and 88% reduction in burden (p=0.009) after drug is withdrawn. Compared to vehicle-treated mice during washout, this represents an 87% reduction in frequency (p=0.006) and 81% reduction in burden (p=0.002) (2-way ANOVA; n=5).

Drug washout occurred over the weekend following week 12 post-SE[59]. Mice treated with CP690550 continued to show profound seizure reduction for at least 2 weeks post-washout compared to vehicle-treated mice (2-way ANOVA; **Frequency:** 87% reduction, p<0.01, **Burden:** 81% reduction, p<0.01). This data supports the hypothesis that CP690550 exerts robust disease modification in a preclinical rodent model of epilepsy.

DISCUSSION

We show that CP690550 (Tofacitinib), an FDA approved, orally available drug with no known behavioral side effects, exerts a potent and enduring seizure-suppressive effect when administered transiently across rodent preclinical epilepsy models. We demonstrate that CP690550 shuts down a pro-epileptic gene network centered around STAT3 by inhibition of JAK1, an endogenous gene target of the protective chromatin modifier EZH2. We provide evidence that the STAT3 upstream kinase JAK1 is a novel therapeutic target for seizure suppression and disease modification across epilepsy models.

After status epilepticus, we find EZH2 protein is potently induced in hippocampal pyramidal and dentate granule neurons across models of epilepsy[23]. Previously, we reported that treatment of mice with the SAM-competitive EZH2 inhibitor UNC-1999 exacerbated spontaneous seizures, and here we find that deletion of EZH2 in mature neurons has the same effect. Induction of Polycomb group (PcG) proteins has been studied in association with other brain insults. For example, sublethal ischemic insults lead to induction of PcG proteins in mouse cortical neurons, and *in vitro* overexpression of PRC1 component BMI1 was sufficient to protect against oxygen-glucose deprivation-associated injury[61]. Further, sevoflurane (a volatile anesthetic that exerts positive allosteric modulation on GABA_A receptors[62] and

antagonizes NMDA receptors[63]) mediates the protective induction of EZH2 and H3K27Me3 to attenuate hypoxic-ischemic brain injury, reducing hippocampal autophagy and improving post-injury behavioral

assessments in rats. Intriguingly, these improvements are reversed when rats treated with sevoflurane also receive a dose of the EZH2 inhibitor GSK126[64]. These studies, along with our findings, point to a potentially general protective role for EZH2 in brain insults.

EZH2 deletion in neurons results in large transcriptomic changes that manifest after SE (Fig. 2). The mouse Black gene cluster houses genes whose expression are relatively unchanged in naïve EZH2nWT and EZH2nHom animals, induced after SE in EZH2nWT, and hyper-induced in EZH2nHom mice after SE. This pattern suggests that EZH2 might be responsible for tempering expression of genes in this cluster because its deletion results in their unfettered activation. The Black cluster ontological analysis produces terms associated with innate immunity, neuroinflammation, and the JAK/STAT pathway. Network analysis places STAT3 in the center of the Black GRN, and immunofluorescence confirms induction of STAT3 in neurons post SE. While activation of the JAK/STAT pathway in non-neuronal cells post-SE is well-characterized[26,47,65,66], the role of STAT3 in neurons, especially in epilepsy, has been less studied; thus the mechanisms of neurogenic inflammation post-SE are not clear[45]. Cytokines released from glial cells following seizures increase neuronal excitability[67], and type I IFNs increase excitability in neocortical pyramidal neurons via PKC[68]. Others report that seizure-induced activation of the JAK/STAT pathway reduces the number of gamma-aminobutyric acid (GABA) type A alpha1 receptors expressed in neurons following seizures, influencing GABA(A)R-dependent inhibition[27]; this group has also utilized pharmacological inhibition of STAT3[32] and genetic ablation of the STAT3 SH2 domain[45] to reduce spontaneous seizure frequency in rodent models of epilepsy. Thus, activation of neuroinflammatory pathways, including the JAK/STAT pathway, has been shown to modulate neuronal excitability. However, our work is the first to identify an endogenous protective mechanism induced by SE that works to temper JAK/STAT activation and associated neurogenic inflammation. Our's is also the first to demonstrate an enduring protective effect of targeting this pathway in chronic epilepsy after the onset of spontaneous seizures.

In order to understand the mechanism by which EZH2 suppresses STAT3 activation, and to rule out the possibility that STAT3 hyper-induction in EZH2 mutant mice might be a consequence of worsened disease progression more generally, rather than loss of EZH2 *per se*, we turned to an *in vitro* cell line model. Knocking down EZH2 in N2A cells results in induction of STAT3 Y705 phosphorylation. This activation is not via secretion of soluble factors as swapping media from control and knockdown cells fails to activate STAT3 (data not shown). WP1066, a JAK2 inhibitor[51-53] fails to suppress activated STAT3 in shEZH2 cells whereas CP690550 (a JAK1/3 inhibitor) (Fig. 4B-C) and direct knockdown of JAK1 (Fig. 4H-J) succeeds. Given that JAK3 is not expressed in these cells, we conclude that EZH2 tempers JAK1. Finally, we find that EZH2 directly interacts with the JAK1 locus (Fig. 4G). Increased activation of STAT3 at pY705-STAT3 as a result of EZH2 inactivation has also been observed in a transformed murine cell model of early T cell precursor acute lymphoblastic leukemia; EZH2 inactivation leads to enrichment of documented STAT3 target genes, increased expression of IL6RA, and an exaggerated pY705-STAT3 induction in response to IL6 stimulation[69]. STAT-upstream receptors and kinases are also regulated by EZH2. In prostate cancer EZH2 directly targets the interferon gamma receptor gene INFGR1 to dampen JAK2/STAT1 signaling, and EZH2 depletion restores expression of IFN target genes as well as STAT1 phosphorylation.[70]

Others have identified EZH2 as an activator of JAK/STAT signaling as well as a target of the pathway. When phosphorylated by AKT at serine 21, EZH2 disassociates from Polycomb to enhance STAT3 transcriptional activity through di-methylation of STAT3-K49 in glioblastoma cells[71-73]; we have performed western blotting to assess this posttranslational mark in mouse hippocampi post-SE and find no change in STAT3-K49Me2 after seizures or deletion of EZH2 (data not shown). Thus, this mechanism does not appear to play a role in N2As or the mouse brain. In NK/T-cell lymphoma, JAK3 phosphorylates EZH2 to mediate a switch in its function, promoting the transcriptional upregulation of oncogenes through the interaction of EZH2 and Pol-II[74]. To our knowledge, ours is the first report of EZH2 repressing JAK/STAT activation through JAK1.

Analysis of human TLE samples identifies a cluster of genes strikingly similar to the STAT3-driven mouse Black cluster; the human Brown cluster is a group of genes associated with innate immunity and neuroinflammation. Like the mouse Black cluster, the most influential driver of gene expression is STAT3. This is a possible explanation for why CP690550 suppresses disease whether it is administered shortly after SE (Fig. 5), or after the onset of spontaneous seizures (Fig. 6). The robust overlap between clusters of genes, one from frankly epileptic human samples and another from rodents, suggests a couple of points. Firstly, EZH2-

578 STAT3 mediated mechanisms identified in mice are clinically relevant. Secondly, the EZH2-STAT3 mechanism
579 evoked acutely after insult plays a role later in chronic epilepsy; this is supported by the identification of STAT3
580 as a driver of gene expression, its transient activation acutely post-SE, and its reactivation in chronic rodent
581 epilepsy (see Fig. 3).
582
EZH2nKO mice show both a decreased seizure threshold 4 days post SE as well as exaggerated seizure
burden 5 weeks later compared to WT mice. CP690550 administration for 24 hours post-SE is sufficient to
rescue both these phenotypes. This suggests that STAT3 hyper-induction drives seizure threshold reductions
and spontaneous seizure increases in mutant mice. The observation that CP690550 profoundly suppresses
spontaneous seizures when administered to EZH2nHom mice during the chronic period is consistent with
reactivation of JAK/STAT signaling during the chronic period (Fig. 3E-G, 5F-G, 6C-K). This supports the
possibility that mechanisms involved in disease initiation (e.g., activation of neuroinflammatory pathways)
continue to perpetuate disease chronically. Leveraging the brain's own protective mechanism, we use
CP690550 to buttress EZH2 suppression of JAK1. CP690550 profoundly suppresses seizures in WT epileptic
mice across models of epilepsy. Following cessation of CP690550 treatment we observe robust disease
modification for at least two weeks. Whether this disease-modifying effect is permanent will be the focus of
future studies in the lab.
595
Anti-inflammatory drugs that have been used to suppress seizures in human patients[28-31] or have shown
antiepileptogenic potential[32-35] in rodents pose several drawbacks: some or all are proposed only as
supplemental therapies[28,29], risk worsened disease outcomes[29], show only transient effects[30], or are effective in
only a subset of epilepsy patients[29,31]. We argue that an approach involving multiple animal models and
multiple laboratories to identify drivers of disease increases the likelihood that druggable targets will be viable
across human etiologies. With this approach we identify EZH2 as a protective negative regulator of the
JAK/STAT pathway in epilepsy. Our current data is consistent with the hypothesis that CP690550 mitigates
spontaneous seizures in chronic epilepsy and is disease-modifying. The cross-model approach by which we
identified JAK1 and selected CP690550 suggests this drug may be useful across multiple epilepsies.
605
606 Recently, others find that neuron-specific deletion of the STAT3 SH2 domain plateaus, but does not reduce,
607 seizure frequency in chronic epilepsy[45], whereas CP690550 treatment greatly reduces seizures. The
608 explanation for this difference may be that there is incomplete ablation of STAT3 in their model, or it may
609 suggest that the more robust therapeutic effect of CP690550 is attributable to JAK/STAT pathway suppression
610 in non-neuronal cells (e.g., peripheral immune cells, microglia, etc.) as well as neurons.
611
The JAK/STAT pathway is likely one of many processes orchestrating disease progression. The data
presented here represents the first evidence that EZH2 regulates JAK/STAT function through JAK1/STAT3 in
epileptogenesis to suppress disease progression. Transient treatment with CP690550 not only mitigates
exaggerated disease in EZH2 mutant mice, but is seizure suppressing and chronically disease modifying in
WT mice in two different epilepsy models. Together, these data suggest an ongoing interplay between EZH2
and the JAK/STAT pathway, starting in early epileptogenesis and persisting in chronic epilepsy.
618
619 We conclude that EZH2 tempers the JAK/STAT pathway in epilepsy as a protective response but is ultimately
620 overwhelmed. Buttressing this protective program with the FDA approved, orally available drug CP690550
621 thwarts disease progression and offers a novel therapeutic approach to treating epilepsy.
622
623 # METHODS
624
625 ## Bioinformatics
626
Reads were mapped to the mm39 build mouse genome and subjected to Likelihood Ratio Tests using
Deseq2[75]. Transcripts per million were calculated and genes were classified as expressed if they had at least 3
TPM in all samples of any one of the 4 conditions (saline:WT, kainate:WT, saline:EZH2nHom,
kainate:EZH2nHom). Deseq2 output was then filtered for only expressed genes and adjusted p-values were
recalculated for the reduced number of genes. Genes with an adjusted $p<.01$ and 1.2-fold change were
subjected to Leiden clustering with resolution=0.5 using the scanpy Leiden tool[76]. Ontology was performed on
all clusters using Ontomancer (manuscript in preparation). Briefly, Biocarta, Hallmark, KEGG, and Reactome

.gmt files were downloaded from https://www.gsea-msigdb.org/gsea/downloads.jsp. Gene sets associated with every term in the databases were collected. For each Leiden cluster, Scores (S_t) for each term, t, were calculated as:

$$S_t = \sqrt{(\log O_t)^2 + (-\log P_t)^2}$$

Where O_t and P_t, are the odds ratio and p values from Fisher exact tests for the overlap of Leiden cluster genes and genes for each term t. Only terms with odds ratios>1 and p values<0.05 were kept for further analysis. For each term, a gene list is generated containing the overlap of genes in the cluster and term, G_t. A term x term similarity matrix was generated where similarity between term i and j is the Fisher exact test log odds ratio $O_{i,j}$ for the intersection of genes in the terms G_i and G_j. This was converted to a network where nodes are terms. Edge weights were calculated as:

$$weight = (\log(O_{i,j})/\max \log O)^6$$

Where max log odds is the maximum log odds value across all comparisons. The 6 power was used to separate nodes for visualization. Networks were generated using the scanpy Leiden tool with resolution=2. Each community was named after the highest scoring term in that community. Networks were visualized via Python's networkx and Plotly tools.

MAGIC was performed on all clusters essentially as described[21] but with r=1, and GTFs were ignored. GTFs were defined as any ENCODE Factors containing the strings "TBP", "POL", "GTF", or "TAF".

Gene Regulatory Network was projected using Cytoscape[77] using MAGIC outputs. Factors were connected to their targets with edges defined as ChIP signal extracted from the Target Folder generated by MAGIC. Target genes of Factors that were not themselves Factors were removed to generate the GRN. Network statistics were calculated in Cytoscape[77].

Analysis of human TLE expression data.

GSE63808[50] from GEO contained 129 transcriptomes from resected hippocampal TLE tissue. Probes with an Illumina p-value<0.05 in at least 20% of samples were retained in the analysis; the probes were consolidated to gene symbols by taking the median probe expression value per symbol. These 9076 genes formed the background list for ontological and MAGIC analyses. Expression values were normalized by the median expression value in that sample. Genes kept for further analysis exhibited variance greater that the median coefficient of variance across all samples. The resulting 7658 unique genes were subjected to k-means clustering using the k-means tools from Sci-kit learn[78] to divide genes into 10 clusters (brown, blue, purple, skyblue, grey, orange, indigo, green, yellow, red).

Overlap between each of the 10 human TLE clusters with mouse Black cluster genes was measured by Fisher exact test with the Benjamini-Hochberg correction for multiple comparisons. Ontological and MAGIC analyses were performed on the highest-scoring human Brown cluster as described in the previous section.

Cell culture conditions, lentiviral knockdown, and PRC2 overexpression.

Cells were grown in 5% CO_2 at 37°C. Neuro2a cells were grown in MEM (#10-009-CV Corning, Manassas, VA) with 1.0g/L glucose, 1.5g/L L-Glutamine, and 10% fetal bovine serum (#26140-079 Gibco, Grand Island, NY). Stable EZH2 knockdown in Neuro2a cells for western blot and RNA analysis was achieved using SMARTvector lentiviral delivery of shRNA per the manufacturer instructions. Puromycin selection began 48 hours after infection and maintained during cell expansion and experimentation. SMARTvector lentiviral mouse EZH2 mEF1a-TurboRFP shRNA (#V3SM7592-232015353, Dharmacon, Lafayette, CO) targeted the sequence *ATCGTAGTAAGTACCAATG*, and non-targeting mEF1a-TurboRFP control particles (#S10-005000-01) were used as an infection control.

688 For overexpression of Polycomb, all plasmids were ordered and modified (if necessary) from Addgene
689 (Watertown, MA). Control cells were transfected with pAdtrack-CMV (#16405). To overexpress PRC2, cells
690 were transfected with pCMVHA-hEZH2 (#24230), pCMVHA-EED (#24231), and pCMVHA-SUZ12 (#24232).
691 The EZH2 plasmid was modified in-house to produce an HA-EZH2-GFP fusion protein. All transfections were
692 carried out with Lipofectamine 3000 transfection reagent (#L3000015, Invitrogen).
693

Animal care

All animal procedures and experiments were performed with approval from the University of Wisconsin-
Madison School of Medicine and Public Health Instructional Animal Care and Use Committee, or the
Institutional Animal Care and Use Committee of Emory University, as appropriate, and according to NIH
national guidelines and policies.

Transgenic mice. Cre-driver and EZH2 floxed mice were ordered from Jackson Laboratories
(#003966, #022616, Bar Harbor, ME). Mice were bred on a C57BL6 background. Using the previously
described Synapsin1-Cre system [37] $EZH2^{+/-}$; $Syn1Cre^{+/-}$ (EZH2nHet) mice were generated by mating $EZH2^{flox/+}$;
$Syn1Cre^{+/-}$ females with $EZH2^{flox/flox}$ and $EZH2^{flox/+}$ males [36]. $EZH2^{+/+}$; $Syn1Cre^{+/-}$ (EZH2nWT) mice were
generated from mating pairs containing an $EZH2^{flox/+}$ male. To assess Cre expression, we used a reporter
mouse with a lox-P flanked STOP cassette preventing transcription of tdTomato (#007909 Jackson
Laboratories) [79]. Reporter males were bred with $Syn1Cre^{+/-}$ females to generate Cre-positive animals and Cre-
negative controls.

Repeated low-dose KA model. Male and female C57BL6 and FVB mice were bred and housed under
a 12-hour light/dark cycle with access to food and water *ad libitum*. C57BL6 mice were allowed to reach 4-6
weeks of age, and FVB animals 5-7 weeks of age, before undergoing experimentation. Kainate injections were
performed at the same time of day (~10 A.M.), and mice were returned to home cages before the start of the
dark cycle (~5 P.M.). Mice were housed with littermates for the duration of the experiment, with males and
females separated at weaning.
We used a repeated low-dose kainate mouse model to induce SE. To begin, mice were weighed and
singly housed in observation chambers for the duration of the injections. Mice were injected intraperitoneally
(i.p.) with 5.0mg/kg synthetic kainic acid (KA) dissolved in 0.9% saline (#7065, Tocris Bioscience, Bristol,
United Kingdom). At twenty-minute intervals, mice were given 5.0mg/kg injections of KA up to the third
injection. The dosage was then reduced to 3.75mg/kg. Animals continued to receive 3.75mg/kg injections of
KA every twenty minutes until each reached SE. The subsequent KA injection was skipped if an animal
experienced two or more Class V seizures within a single twenty-minute interval. Injections resumed for the
next round, unless the animal reached SE. During induction of SE, seizures were scored using a modified
Racine Scale [80] where I = freezing, behavioral arrest, staring spells; II = head nodding and facial twitches; III =
forelimb clonus, whole-body jerks or twitches; IV = rearing; V = rearing and falling; VI = violent running or
jumping behavior. Animals were considered to have reached SE after experiencing at least five Class V or VI
seizures within a 90-minute window. KA mice were observed for 1-2 hours after SE. Animals were injected with
0.9% saline (s.c.), and soft gel food was provided in home cages. In the days following injection, animals were
weighed and injected with 0.9% saline (s.c.) if body weight decreased by more than 0.5g.

Pilocarpine model of status epilepticus. Adult C57Bl/6 mice (8-12 weeks old, Charles River) were
injected with terbutaline (4 mg/kg ip, Sigma T2528) and methylscopolamine (4 mg/kg ip, Sigma S8502) to
alleviate respiratory and cardiovascular effects of pilocarpine, then 30 min later received saline or pilocarpine
(280 mg/kg ip as free base, Sigma P6503) as described[34]. After 1 h of SE, seizures were interrupted by
diazepam (10 mg/kg ip).

IHKA model of status epilepticus. The intrahippocampal kainic acid (IHKA) model was employed to
generate chronically epileptic mice as previously described[81]. Kainic acid (100 nL of 20 mM; vIHKA) or vehicle
(100 nL of sterile saline; vIHSA) was stereotaxically injected into the hippocampus using the following
coordinate (from Bregma): 3.6 mm posterior, -2.8 mm lateral, and 2.8 mm depth. Only mice which experienced
status epilepticus and went on to develop spontaneous recurrent seizures were used in the current study.
742

Fluorothyl and PTZ seizure threshold testing. Male and female C57BL6 mice were bred and housed under a 12-hour light/dark cycle with access to food and water *ad libitum*. Animals were allowed to reach 4-6 weeks of age before undergoing experimentation. Fluorothyl seizure threshold tests were performed at the same time of day (~10 A.M.), and mice were returned to home cages before 5 P.M. Mice were housed with littermates for the duration of the experiment, with males and females separated at weaning. All seizure threshold tests were video recorded, and seizure behavior was scored by a blinded observer. Two tests per mouse were performed: the first at one week prior to kainate injections and the second at four days post-SE.

Seizure threshold tests were performed on male and female EZH2nWT wildtypes and EZH2nHet mutant mice using 100% bis(2,2,2-trifluoroethyl) ether (#287571, Sigma-Aldrich, St. Louis, MO). All tests were performed in a fume hood. Mice were placed in an airtight 10L. Plexiglas chamber (8.5"X10.75"X6.75"). Fluorothyl was infused into the chamber via a peristaltic pump at a rate of 40μL/minute onto a piece of Whatman filter paper placed at the top of the chamber. The time to generalized tonic clonic seizure (GTCS) was recorded from the start of fluorothyl infusion. Mice reached GTCS when they exhibited a complete loss of postural and motor control. After experiencing a generalized tonic clonic seizure, mice were rapidly removed from the chamber, and fluorothyl infusion was terminated. Mice were transferred to a recovery cage and kept isolated until motor control and normal behavior returned (~10 minutes). Mice were returned to home cages after recovery. PTZ was dissolved at 5mg/mL in 0.9% normal saline, and a single 45mg/kg dose was administered (i.p.). Time to generalized tonic clonic seizure was recorded.

Long-term video monitoring for spontaneous seizures. Male and female FVB mice were bred and housed under a 12-hour light/dark cycle with access to food and water *ad libitum*. Animals were allowed to reach 5-7 weeks of age before undergoing experimentation. Video recording was performed at the same time of day (~10 A.M.), and mice were returned to home cages before 5 P.M. Mice were singly housed in chambers for the duration of recording with access to food and water. During daily CP690550 treatment, mice were weighed and injected daily before recording began. Videos were reviewed and scored by an experimenter blinded to treatment and genotype using the aforementioned modified Racine scale. In the final experiment described by Figure 7, only Racine class III-VI seizures were recorded. For each mouse, daily seizure burden was calculated as: (Total # of seizures * Average seizure duration (s))/(Total recording hours).

CP690550 citrate treatment of KA mice. Following kainate injections, an independent researcher (not the experimenter) randomly assigned mice to drug and vehicle conditions, and prepared both solutions. Drug and vehicle solutions were coded as "A" or "B." CP690550 citrate (tofacitinib citrate) (#4556, Tocris Bioscience, Bristol, United Kingdom) was dissolved in a mixture of 70% saline (0.9%)/15% ethanol/15% DMSO. The 70% saline/15% ethanol/15% DMSO solution was administered as vehicle. The independent researcher provided the blinded experimenter with the drug solutions and a list of all mice assigned to group A or B; this researcher did not participate in any seizure threshold testing or kainate injections. CP690550 or vehicle was delivered to mice intraperitoneally at 6, 12, and 24 hours post-SE. The drug doses for each of these time points, respectively, were 25mg/kg, 15mg/kg, and 15mg/kg.

Continuous EEG recording and seizure detection.
At the time of stereotaxic IHKA or IHSA injections, mice were implanted with chronic EEG recording headmounts (Pinnacle Technology, cat. #8201) which was affixed to the skull with 4 screws: 2 screws serving as EEG leads placed in the anterior and posterior cortex, 1 as a reference ground, and 1 as the animal ground. The headmounts were secured with dental cement. 24/7 EEG recordings were collected at 4 KHz using a 100X gain preamplifier high pass filtered at 1KHz (Pinnacle Technology, cat. #8292-SE) and tethered turnkey system (Pinnacle Technology, cat. #8200).

Seizures were detected using a custom analysis pipeline developed in-house in python (seizy; https://github.com/neurosimata/seizy) as previously described[81]. Briefly, EEG traces were automatically detected using a thresholding method from traces that were downsampled (decimated) to 100 Hz and divided into 5 second segments. Power (2-40 Hz) and line length were used to identify seizures. Seizures were only included if two consecutive 5 second segments were classified as seizures (minimum 10 seconds length) and were manually verified by an experienced investigator with expertise in electrographic seizure detection. Seizure frequency was calculated as the # of seizures per day by measuring total number of seizures a mouse had over a 24-hour period. The average seizure duration was calculated by taking the mean seizure duration for each mouse in each recording period. For each mouse, seizure burden was calculated as: (Total # of seizures * Average seizure duration)/(Total EEG recording hours).

Tissue isolation and homogenization

After KA seizure induction and/or treatment with CP690550 citrate, animals were sacrificed by rapid decapitation. Whole hippocampal hemispheres were harvested and flash-frozen in liquid nitrogen or on dry ice. Single hemispheres were lysed in Radioimmunoprecipitation Assay Buffer (RIPA: 50mM Tris, 150mM NaCl, 1% nonidet P-40, 0.5% sodium deoxycholate, 0.1% SDS) with mammalian protease inhibitor (1:1000, Sigma or 1860932 ThermoScientific) and phosphatase inhibitor (1:1000, Sigma or 78428 ThermoScientific). Protein from cell lines was harvested in Triton lysis buffer (3% 5M NaCl, 10% glycerol, 0.3% Triton X-100, 5% 1M Tris pH8.0) after PBS washes. Tissue was homogenized in lysis buffer by probe sonication (Fisher Scientific, Sonic Dismembrator, Model 100, Hampton, NH) on power 3 for two rounds, with 10 pulses per round. Supernatants were isolated by centrifugation and quantified using the DC Protein Assay (Bio-Rad, Hercules, CA) or BCA assay (ThermoScientific 23227). Protein extracts were stored at -80°C. 5x loading buffer (0.5mM Tris, 10% SDS, 50% glycerol, 10mM EDTA, and 1% beta-mercaptoethanol) was added to each sample to reach a 1x final concentration. Extracts in loading buffer were boiled at 95°C for 5 minutes and stored for up to one month at -20°C until run on an acrylamide gel.

Western blotting

Protein extracts in loading buffer were loaded at 20µg per lane and resolved by electrophoresis in hand-poured acrylamide gels with a 5% acrylamide stacking layer (125mM Tris pH6.8, 5% acrylamide, 0.01% ammonium persulfate, 0.01% SDS, 0.01% TEMED) and an 8% acrylamide separating layer (375mM Tris pH8.8, 8% acrylamide, 0.015% ammonium persulfate, 0.015% SDS, 0.08% TEMED). Gels were transferred to polyvinyl difluoride membranes (PVDF; Millipore, Bedford, MA) using Tris-glycine transfer buffer (20mM Tris, 1.5M glycine, 20% methanol). Membranes were blocked with 5% bovine serum albumin (Fisher Scientific, Fair Lawn, NJ) diluted in low-salt Tris-buffered saline (w/w TBST; 20mM Tris pH7.5, 150mM NaCl, 0.1% Tween 20) with mammalian protease inhibitor (1:2000) and phosphatase inhibitor (1:2000), or Odyssey blocking solution (LI-COR, 927-60001) for 1 hour at room temperature. Primary antibodies were diluted in the same blocking buffer and incubated with membranes overnight at 4°C. Antibodies include: EZH2 (1:1000, #5426 Cell Signaling, Danvers, MA), pY705-STAT3 (1:2000, #9145S Cell Signaling), STAT3 (1:1000, #12640 Cell Signaling), JAK2 (1:1000, #3230 Cell Signaling), JAK1 (1:1000, #3344S Cell Signaling), pY1034/35-JAK1 (1:1000, #74129S Cell Signaling), pY1007/08-JAK2 (1:500, ab32101, Abcam, Cambridge, UK), and Actin (1:10,000, #691001 MP Biomedicals) or GAPDH (1:10,000; Millipore, CB1001). The next day, the membranes were washed three times in 1X TBST and incubated with horseradish peroxidase-conjugated goat-anti-mouse or -rabbit secondary antibodies for one hour at room temperature (1:10,000, Invitrogen #31430, #31460, Rockford, IL). Membranes were subsequently washed three times in TBST, and membranes were developed in SuperSignal West Femto ECL reagent (ThermoFisher, Waltham, MA). Bands were imaged using a ChemiDoc-It Imaging System (UVP Vision-Works, Upland, CA) and quantified using UVP Vision Works software or ImageJ. Band intensities were graphed and analyzed using Prism 9 software (GraphPad Software, La Jolla, CA).

RNA extraction and clean-up for whole hippocampal RNAseq

Hippocampi were harvested from saline- and kainate-treated EZH2nWT and EZH2nHom mice on day 4 post-SE (n=5 per treatment/genotype). RNA was extracted from hippocampal tissue with Trizol (#1559602, Life Technologies, Carlsbad, CA), and concentration was determined by spectrometry. This was followed by a treatment with DNase (#AM2238, Invitrogen). RNA clean-up was carried with the RNeasy Mini Kit (#74104, Qiagen, Hilden, Germany). Purified samples were submitted to the Gene Expression core at the University of Wisconsin-Madison Biotechnology Center for library preparation and sequencing (https://biotech.wisc.edu).

Chromatin immunoprecipitation and qPCR.

4% formaldehyde was used to crosslink proteins to chromatin before neutralization of the reaction in 125mM glycine, and washes in 1X PBS (pH 7.4). Chromatin was fragmented in Triton-X lysis buffer and 1:100 protease inhibitors; lysates were sonicated and centrifuged to remove cell debris. Protein concentration of the

supernatant was quantified with BioRad Bradford assay reagent (#5000006). Chromatin was incubated overnight at 4 degrees Celsius with antibodies against H3K27Me3 (#9733 Cell Signaling) or rabbit IgG. The next day, protein A/G magnetic beads (#88803, ThermoScientific) were blocked in BSA and salmon sperm DNA while chromatin samples were centrifuged, after which chromatin-antibody complexes were added to the beads to incubate. Washes were performed in a low-salt ChIP buffer (1% SDS, 1% Triton X-100, 0.4% 500mM EDTA, 2% 1M Tris pH8.1, 3% 5M NaCl). Decrosslinking buffer (1% SDS, 100mM $NaHCO_3$, 200mM NaCl) was then added to IgG samples, H3K27Me3 samples, and inputs; decrosslinking reactions were allowed to incubate overnight in a 65 degree Celsius water bath. Phenol-choloform DNA extraction was performed the next day. Quantitative PCR was performed with 2x SYBR green mix (#RR420L, Takara Bio, Otsu, Shiga, Japan) and primers against JAK1, STAT3, JAK2, PAX2, Hoxc10, and beta actin (primers listed at the end of the methods section). Signal was calculated from standard curves generated by serial dilution of an input sample, and enrichment values were expressed as fold over input. RNA for RT-PCR was extracted with Trizol, and concentration was determined by spectrometry. PCR reactions were run with primers against JAK1 and GAPDH. cDNA synthesis was run on the MJ Mini Thermocycler from Bio-Rad, and PCR reactions were run on the MJ Research DNA Engine Opticon 2 Continuous Fluorescence Detector using Opticon Monitor software.

Immunofluorescence

Whole brains were harvested on day 3 or 4 post-SE, bisected sagitally, fixed in 4% paraformaldehyde, dehydrated in a series of graded TBS-Tween/ethanol solutions, embedded in paraffin, sectioned at 5μm and subjected to immunofluorescence for EZH2, H3K27me3, NeuN, GFAP, total STAT3, and DAPI. Heat induced antigen retrieval was performed in a 10mM citric acid buffer (pH 6.0). After a blocking step, rabbit anti-EZH2 (1:50) (#5246 Cell Signaling), rabbit anti-H3K27me3 (1:300) (#39155 Active Motif), mouse anti-NeuN (1:200 ab104224 Abcam, or 1:500 MAB377 from Millipore) primary antibodies were applied and incubated overnight at 4°C. Slides were washed with PBS then incubated with goat-anti-mouse or goat-anti-rabbit Alexa Fluor 488 (ab150113 Abcam or A11005, A11008 from InVitrogen, respectively) and goat-anti-rabbit Alexa Fluor 594 (ab150080 Abcam or Jackson ImmunoResearch) secondary antibodies at a concentration of 1:200 for one or two hours at room temperature, followed in some experiments by two-hour incubation at room temperature with Alexa Fluro 488-conjugated mouse anti-GFAP antibody (1:200) (#53-9892-82 Invitrogen). Nuclear DNA was stained with DAPI (300 nM) or a DAPI-containing mounting media (#H-1200 Vector Laboratories, Burlingame, CA). Slides were imaged at 20-40x magnification using a fluorescent microscope (Nikon E600 microscope with fluorescence or Zeiss Axio Observer A1 epifluorescence microscope). Images were captured with Zeiss Axiovision or Cellsens software (Olympus Life Science Solutions, Center Valley, PA). Image analysis and quantification was performed in ImageJ (NIH, https://imagej.nih.gov/ij/). EZH2 antibody specificity was confirmed before usage, as shown in Fig. S1A.

Statistical analysis

For all statistical analysis – with the exception of RNAseq data – FDR < 0.05 corrected for multiple comparisons was considered statistically significant. Data were graphed as mean $\pm$ standard deviation (SD) in Prism 9 software (Graphpad Software, La Jolla, CA). Statistical tests were also performed in Prism. Western blot, seizure threshold, and spontaneous seizure data was analyzed by 2-way or 1-way ANOVA with the Benjamini-Hochberg FDR method for multiple comparisons, or by unpaired t-test where appropriate.

Primer List
mouse-JAK1-Site1 (ChIP):
(forward) *CTCTGCTCGCTTTCCCGGAGATGC*
(reverse) *CTCGGAGGCTGGAGCGTGACTTCT*

mouse-JAK1-Site2 (ChIP):
(forward) *TGACTCGCACCCAGCCTGAAATGG*
(reverse) *GCGGCCCAGTCACCCGAAGAAATA*

mouse-JAK1-Site3 (ChIP):
(forward) *TTCGCGGGGCAAGGGAAAGGATTC*
(reverse) *CCACCGCTCAGGGCTCTGGGTTTT*

mouse-JAK2 (ChIP):
(forward) *CGCCGCTGCGGGATGTGAGTGC*
(reverse) *GGAGGCGGGATCGCAGGGAGAACC*

mouse-STAT3 (ChIP):
(forward) *CGGGCGAGCGCTGTACTTCCTA*
(reverse) *AAGCGAGCCGTATCAGGGCATTTA*

mouse-PAX2 (ChIP):
(forward) *GCGGCGCTCAGTCGCGTTCATAG*
(reverse) *CTCCCCGGAGTCTGGGCAGTCCA*

mouse-Hoxc10 (ChIP):
(forward) *GCGGGGGCCCTCTGCCTAAT*
(reverse) *CCCGGCCCTGGGCTCACACT*

mouse-JAK1 (RT-PCR):
(forward) *GCGCCCAGCCTCGCTCGTC*
(reverse) *AACGCCATGGCATTGCAGTCCTCT*

mouse-GAPDH (RT-PCR):
(forward) *ACGGCAAATTCAACGGCACAGTCA*
(reverse) *TGGGGGCATCGGCAGAAGG*

Funding. Supported by CURE (AR), Lily's Fund (AR), NIH grant 1R01NS108756 (AR, RD) and R01NS105628, R01NS102937, and R21NS120868 (JM).

Acknowledgements. We thank Asheebo Rojas for providing rat brain tissues in the DFP model.

Conflict of Interest. JM serves on the Scientific Advisory Board and has a Sponsored Research Agreement with SAGE Therapeutics for work unrelated to this project. AR and RD have no conflicts of interest to declare.

References

Hirtz, D. *et al.* How common are the "common" neurologic disorders? *Neurology* **68**, 326 (2007). https://doi.org/10.1212/01.wnl.0000252807.38124.a3

Gavvala, J. R. & Schuele, S. U. New-Onset Seizure in Adults and Adolescents: A Review. *Jama* **316**, 2657 (2016). https://doi.org/10.1001/jama.2016.18625

Sankaraneni, R. & Lachhwani, D. Antiepileptic drugs--a review. *Pediatric Annals* **44**, e36 (2015). https://doi.org/10.3928/00904481-20150203-10

Meyer, D. M. *et al.* Anti-inflammatory activity and neutrophil reductions mediated by the JAK1/JAK3 inhibitor, CP-690,550, in rat adjuvant-induced arthritis. *Journal of Inflammation (London, England)* **7** (2010). https://doi.org/10.1186/1476-9255-7-41

Dowty, M. E. *et al.* Preclinical to clinical translation of tofacitinib, a Janus kinase inhibitor, in rheumatoid arthritis. *The Journal of Pharmacology and Experimental Therapeutics* **348**, 165 (2014). https://doi.org/10.1124/jpet.113.209304

French, J. A. *et al.* Characteristics of medial temporal lobe epilepsy: I. Results of history and physical examination. *Annals of Neurology* **34**, 774 (1993). https://doi.org/10.1002/ana.410340604

Trinka, E. *et al.* A definition and classification of status epilepticus--Report of the ILAE Task Force on Classification of Status Epilepticus. *Epilepsia* **56**, 1515 (2015). https://doi.org/10.1111/epi.13121

Hesdorffer, D. C., Logroscino, G., Cascino, G., Annegers, J. F. & Hauser, W. A. Risk of unprovoked seizure after acute symptomatic seizure: effect of status epilepticus. *Annals of Neurology* **44**, 908 (1998). https://doi.org/10.1002/ana.410440609

9   Garriga-Canut, M. *et al.* 2-deoxy-D-glucose reduces epilepsy progression by NRSF-CtBP-dependent metabolic
regulation of chromatin structure. *Nature Neuroscience* **9**, 1382-1387 (2006). https://doi.org/10.1038/nn1791
10   Stafstrom, C. E., Roopra, A. & Sutula, T. P. Seizure suppression via glycolysis inhibition with 2-deoxy-D-glucose
(2DG). *Epilepsia* **49 Suppl 8**, 97 (2008). https://doi.org/10.1111/j.1528-1167.2008.01848.x
11   Stafstrom, C. E. *et al.* Anticonvulsant and antiepileptic actions of 2-deoxy-D-glucose in epilepsy models. *Annals of
Neurology* **65**, 435 (2009). https://doi.org/10.1002/ana.21603
12   Vezzani, A., Friedman, A. & Dingledine, R. J. The role of inflammation in epileptogenesis. *Neuropharmacology* **69**,
16 (2013). https://doi.org/10.1016/j.neuropharm.2012.04.004
13   Rojas, A. *et al.* A Novel Second-Generation EP2 Receptor Antagonist Reduces Neuroinflammation and Gliosis
After Status Epilepticus in Rats. *Neurotherapeutics: The Journal of the American Society for Experimental
NeuroTherapeutics* (2021). https://doi.org/10.1007/s13311-020-00969-5
14   Williams-Karnesky, R. L. *et al.* Epigenetic changes induced by adenosine augmentation therapy prevent
epileptogenesis. *Journal of Clinical Investigation* **123**, 3552-3563 (2013). https://doi.org/10.1172/jci65636
15   Liu, G. *et al.* Transient inhibition of TrkB kinase after status epilepticus prevents development of temporal lobe
epilepsy. *Neuron* **79**, 31 (2013). https://doi.org/10.1016/j.neuron.2013.04.027
16   Landis, S. C. *et al.* A call for transparent reporting to optimize the predictive value of preclinical research. *Nature
490, 187 (2012). https://doi.org/10.1038/nature11556
17   Becker, A. J. *et al.* Correlated stage- and subfield-associated hippocampal gene expression patterns in
experimental and human temporal lobe epilepsy. *The European Journal of Neuroscience* **18**, 2792 (2003).
https://doi.org/10.1111/j.1460-9568.2003.02993.x
18   Borges, K., Shaw, R. & Dingledine, R. Gene expression changes after seizure preconditioning in the three major
hippocampal cell layers. *Neurobiology of Disease* **26**, 66 (2007). https://doi.org/10.1016/j.nbd.2006.12.001
19   Hansen, K. F., Sakamoto, K., Pelz, C., Impey, S. & Obrietan, K. Profiling status epilepticus-induced changes in
hippocampal RNA expression using high-throughput RNA sequencing. *Scientific Reports* **4** (2014).
https://doi.org/10.1038/srep06930
20   McClelland, S. *et al.* The transcription factor NRSF contributes to epileptogenesis by selective repression of a
subset of target genes. *eLife* **3** (2014). https://doi.org/10.7554/eLife.01267
21   Roopra, A. MAGIC: A tool for predicting transcription factors and cofactors driving gene sets using ENCODE data.
PLoS computational biology **16** (2020). https://doi.org/10.1371/journal.pcbi.1007800
22   Dingledine, R. *et al.* Transcriptional profile of hippocampal dentate granule cells in four rat epilepsy models.
Scientific Data **4** (2017). https://doi.org/10.1038/sdata.2017.61
23   Khan, N. *et al.* A systems approach identifies Enhancer of Zeste Homolog 2 (EZH2) as a protective factor in
epilepsy. *Plos One* **14** (2019). https://doi.org/10.1371/journal.pone.0226733
24   Pfisterer, U. *et al.* Identification of epilepsy-associated neuronal subtypes and gene expression underlying
epileptogenesis. *Nature Communications* **11** (2020). https://doi.org/10.1038/s41467-020-18752-7
25   Cid, E. *et al.* Sublayer- and cell-type-specific neurodegenerative transcriptional trajectories in hippocampal
sclerosis. *Cell Reports* **35** (2021). https://doi.org/10.1016/j.celrep.2021.109229
26   Kumar, P. *et al.* Single-cell transcriptomics and surface epitope detection in human brain epileptic lesions
identifies pro-inflammatory signaling. *Nature Neuroscience* **25**, 956 (2022). https://doi.org/10.1038/s41593-022-
01095-5
27   Lund, I. V. *et al.* BDNF Selectively Regulates GABAA Receptor Transcription by Activation of the JAK/STAT
Pathway. *Science Signaling* **1**, ra9 (2008). https://doi.org/10.1126/scisignal.1162396
28   You, S. J., Jung, D. E., Kim, H. D., Lee, H. S. & Kang, H. C. Efficacy and prognosis of a short course of prednisolone
therapy for pediatric epilepsy. *European journal of paediatric neurology: EJPN: official journal of the European
Paediatric Neurology Society* **12**, 314 (2008). https://doi.org/10.1016/j.ejpn.2007.09.003
29   Dubey, D. *et al.* Randomized Placebo-Controlled Trial of Intravenous Immunoglobulin in Autoimmune
LGI1/CASPR2 Epilepsy. *Annals of Neurology* **87**, 313 (2020). https://doi.org/10.1002/ana.25655
30   Gayatri, N. A., Ferrie, C. D. & Cross, H. Corticosteroids including ACTH for childhood epilepsy other than epileptic
spasms. *The Cochrane Database of Systematic Reviews* (2007).
https://doi.org/10.1002/14651858.CD005222.pub2
31   Lai, Y.-C. *et al.* Anakinra usage in febrile infection related epilepsy syndrome: an international cohort. *Annals of
Clinical and Translational Neurology* **7**, 2467 (2020). https://doi.org/10.1002/acn3.51229

32 Grabenstatter, H. L. *et al.* The effect of STAT3 inhibition on status epilepticus and subsequent spontaneous
seizures in the pilocarpine model of acquired epilepsy. *Neurobiology of Disease* **62**, 73 (2014).
https://doi.org/10.1016/j.nbd.2013.09.003
33 Jiang, J. *et al.* Suppressing pro-inflammatory prostaglandin signaling attenuates excitotoxicity-associated
neuronal inflammation and injury. *Neuropharmacology* **149**, 149 (2019).
https://doi.org/10.1016/j.neuropharm.2019.02.011
34 Varvel, N. H. *et al.* Preclinical development of an EP2 antagonist for post-seizure cognitive deficits.
Neuropharmacology **224** (2023). https://doi.org/10.1016/j.neuropharm.2022.109356
35 Rojas, A., Ganesh, T., Lelutiu, N., Gueorguieva, P. & Dingledine, R. Inhibition of the prostaglandin EP2 receptor is
neuroprotective and accelerates functional recovery in a rat model of organophosphorus induced status
epilepticus. *Neuropharmacology* **93**, 15 (2015). https://doi.org/10.1016/j.neuropharm.2015.01.017
36 Shen, X. *et al.* EZH1 mediates methylation on histone H3 lysine 27 and complements EZH2 in maintaining stem
cell identity and executing pluripotency. *Molecular Cell* **32**, 491 (2008).
https://doi.org/10.1016/j.molcel.2008.10.016
37 Zhu, Y. *et al.* Ablation of NF1 function in neurons induces abnormal development of cerebral cortex and reactive
gliosis in the brain. *Genes & Development* **15**, 859 (2001). https://doi.org/10.1101/gad.862101
38 Basu, T. *et al.* Histone deacetylase inhibitors restore normal hippocampal synaptic plasticity and seizure
threshold in a mouse model of Tuberous Sclerosis Complex. *Scientific Reports* **9** (2019).
https://doi.org/10.1038/s41598-019-41744-7
39 Cilio, M. R. *et al.* Anticonvulsant action and long-term effects of gabapentin in the immature brain.
Neuropharmacology **40**, 139 (2001). https://doi.org/10.1016/s0028-3908(00)00103-9
40 Bierkamper, G. G. & Cenedella, R. J. Induction of chronic epileptiform activity in the rat by an inhibitor of
cholesterol synthesis, U18666A. *Brain Research* **150**, 343 (1978). https://doi.org/10.1016/0006-8993(78)90285-8
41 Liu, Z. & Holmes, G. L. Basic fibroblast growth factor is highly neuroprotective against seizure-induced long-term
behavioural deficits. *Neuroscience* **76**, 1129 (1997). https://doi.org/10.1016/s0306-4522(96)00412-5
42 Applegate, C. D. & Tecott, L. H. Global increases in seizure susceptibility in mice lacking 5-HT2C receptors: a
behavioral analysis. *Experimental Neurology* **154**, 522 (1998). https://doi.org/10.1006/exnr.1998.6901
43 Rho, J. M., Kim, D. W., Robbins, C. A., Anderson, G. D. & Schwartzkroin, P. A. Age-dependent differences in
flurothyl seizure sensitivity in mice treated with a ketogenic diet. *Epilepsy Research* **37**, 233 (1999).
https://doi.org/10.1016/s0920-1211(99)00068-6
44 Samoriski, G. M. & Applegate, C. D. Repeated generalized seizures induce time-dependent changes in the
behavioral seizure response independent of continued seizure induction. *The Journal of Neuroscience: The
Official Journal of the Society for Neuroscience* **17**, 5581 (1997).
45 Tipton, A. E. *et al.* Selective Neuronal Knockout of STAT3 Function Inhibits Epilepsy Progression, Improves
Cognition, and Restores Dysregulated Gene Networks in a Temporal Lobe Epilepsy Model. *Annals of Neurology*
(2023). https://doi.org/10.1002/ana.26644
46 Ahmed, M. M. *et al.* Altered Protein Profiles During Epileptogenesis in the Pilocarpine Mouse Model of Temporal
Lobe Epilepsy. *Frontiers in Neurology* **12** (2021). https://doi.org/10.3389/fneur.2021.654606
47 Vishwakarma, S., Singh, S. & Singh, T. G. Pharmacological modulation of cytokines correlating
neuroinflammatory cascades in epileptogenesis. *Molecular Biology Reports* **49**, 1437 (2022).
https://doi.org/10.1007/s11033-021-06896-8
48 Martín-Suárez, S., Cortes, J. M. & Bonifazi, P. Blockage of STAT3 during epileptogenesis prevents GABAergic loss
and imprinting of the epileptic state. *Brain: A Journal of Neurology* (2023).
https://doi.org/10.1093/brain/awad055
49 Freeman, L. C. Centrality in social networks conceptual clarification. *Social Networks* **1**, 215-239 (1978).
50 Johnson, M. R. *et al.* Systems genetics identifies Sestrin 3 as a regulator of a proconvulsant gene network in
human epileptic hippocampus. *Nature Communications* **6** (2015). https://doi.org/10.1038/ncomms7031
51 Kupferman, M. E. *et al.* Therapeutic suppression of constitutive and inducible JAK\STAT activation in head and
neck squamous cell carcinoma. *Journal of Experimental Therapeutics & Oncology* **8**, 117 (2009).
52 Hussain, S. F. *et al.* A Novel Small Molecule Inhibitor of Signal Transducers and Activators of Transcription 3
Reverses Immune Tolerance in Malignant Glioma Patients. *Cancer Research* **67**, 9630 (2007).
https://doi.org/10.1158/0008-5472.can-07-1243

| 1068 | 53 | Iwamaru, A. *et al.* A novel inhibitor of the STAT3 pathway induces apoptosis in malignant glioma cells both in vitro and in vivo. *Oncogene* **26**, 2435 (2007). https://doi.org/10.1038/sj.onc.1210031 |

53   Iwamaru, A. *et al.* A novel inhibitor of the STAT3 pathway induces apoptosis in malignant glioma cells both in
vitro and in vivo. *Oncogene* **26**, 2435 (2007). https://doi.org/10.1038/sj.onc.1210031

54   Changelian, P. S. *et al.* Prevention of organ allograft rejection by a specific Janus kinase 3 inhibitor. *Science (New*
York, N.Y.) **302**, 875 (2003). https://doi.org/10.1126/science.1087061

55   *UCSC Genome Browser*, <https://genome.ucsc.edu/index.html> (2022).

56   Konze, K. D. *et al.* An orally bioavailable chemical probe of the Lysine Methyltransferases EZH2 and EZH1. *ACS*
chemical biology **8**, 1324 (2013). https://doi.org/10.1021/cb400133j

57   Rocha, L. Subchronic treatment with antiepileptic drugs modifies pentylenetetrazol-induced seizures in mice: Its
correlation with benzodiazepine receptor binding. *Neuropsychiatric Disease and Treatment* **4**, 619 (2008).
https://doi.org/10.2147/ndt.s2118

58   Baradaran Rahimi, V., Askari, V. R., Hosseini, M., Yousefsani, B. S. & Sadeghnia, H. R. Anticonvulsant Activity of
Viola tricolor against Seizures Induced by Pentylenetetrazol and Maximal Electroshock in Mice. *Iranian Journal of*
Medical Sciences **44**, 220 (2019).

59   Cohen, S., Zwillich, S. H., Chow, V., Labadie, R. R. & Wilkinson, B. Co-administration of the JAK inhibitor CP-
690,550 and methotrexate is well tolerated in patients with rheumatoid arthritis without need for dose
adjustment. *British Journal of Clinical Pharmacology* **69**, 143 (2010). https://doi.org/10.1111/j.1365-
2125.2009.03570.x

60   Maguire, J. Epileptogenesis: More Than Just the Latent Period. *Epilepsy Currents* **16**, 31 (2016).
https://doi.org/10.5698/1535-7597-16.1.31

61   Stapels, M. *et al.* Polycomb Group Proteins as Epigenetic Mediators of Neuroprotection in Ischemic Tolerance.
Science Signaling **3**, ra15 (2010). https://doi.org/10.1126/scisignal.2000502

62   Jenkins, A., Franks, N. P. & Lieb, W. R. Effects of temperature and volatile anesthetics on GABA(A) receptors.
Anesthesiology **90**, 484 (1999). https://doi.org/10.1097/00000542-199902000-00024

63   Brosnan, R. J. & Thiesen, R. Increased NMDA receptor inhibition at an increased Sevoflurane MAC. *BMC*
anesthesiology **12** (2012). https://doi.org/10.1186/1471-2253-12-9

64   Xue, H. *et al.* Sevoflurane post-conditioning alleviates neonatal rat hypoxic-ischemic cerebral injury via Ezh2-
regulated autophagy. *Drug Design, Development and Therapy* **13**, 1691 (2019).
https://doi.org/10.2147/dddt.s197325

65   Choi, J.-S. *et al.* Upregulation of gp130 and differential activation of STAT and p42/44 MAPK in the rat
hippocampus following kainic acid-induced seizures. *Molecular Brain Research* **119**, 10 (2003).
https://doi.org/10.1016/j.molbrainres.2003.08.010

66   Wang, H. *et al.* LncRNA-UCA1 inhibits the astrocyte activation in the temporal lobe epilepsy via regulating the
JAK/STAT signaling pathway. *Journal of Cellular Biochemistry* **121**, 4261 (2020).
https://doi.org/10.1002/jcb.29634

67   Iori, V., Frigerio, F. & Vezzani, A. Modulation of neuronal excitability by immune mediators in epilepsy. *Current*
Opinion in Pharmacology **26**, 118 (2016). https://doi.org/10.1016/j.coph.2015.11.002

68   Reetz, O., Stadler, K. & Strauss, U. Protein kinase C activation mediates interferon-β-induced neuronal
excitability changes in neocortical pyramidal neurons. *Journal of Neuroinflammation* **11** (2014).
https://doi.org/10.1186/s12974-014-0185-4

69   Danis, E. *et al.* Ezh2 Controls an Early Hematopoietic Program and Growth and Survival Signaling in Early T Cell
Precursor Acute Lymphoblastic Leukemia. *Cell Reports* **14**, 1953 (2016).
https://doi.org/10.1016/j.celrep.2016.01.064

70   Wee, Z. N. *et al.* EZH2-mediated inactivation of IFN-γ-JAK-STAT1 signaling is an effective therapeutic target in
MYC-driven prostate cancer. *Cell Reports* **8**, 204 (2014). https://doi.org/10.1016/j.celrep.2014.05.045

71   Kim, E. *et al.* Phosphorylation of EZH2 activates STAT3 signaling via STAT3 methylation and promotes
tumorigenicity of glioblastoma stem-like cells. *Cancer Cell* **23**, 839 (2013).
https://doi.org/10.1016/j.ccr.2013.04.008

72   Dasgupta, M., Dermawan, J. K. T., Willard, B. & Stark, G. R. STAT3-driven transcription depends upon the
dimethylation of K49 by EZH2. *Proceedings of the National Academy of Sciences of the United States of America*
112, 3985 (2015). https://doi.org/10.1073/pnas.1503152112

73   Cha, T.-L. *et al.* Akt-mediated phosphorylation of EZH2 suppresses methylation of lysine 27 in histone H3.
Science (New York, N.Y.) **310**, 306 (2005). https://doi.org/10.1126/science.1118947

74   Yan, J. *et al.* EZH2 phosphorylation by JAK3 mediates a switch to noncanonical function in natural killer/T-cell
lymphoma. *Blood* **128**, 948 (2016). https://doi.org/10.1182/blood-2016-01-690701
75   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with
DESeq2. *Genome Biology* **15** (2014). https://doi.org/10.1186/s13059-014-0550-8
76   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data analysis. *Genome*
Biology **19** (2018). https://doi.org/10.1186/s13059-017-1382-0
77   Assenov, Y., Ramírez, F., Schelhorn, S.-E., Lengauer, T. & Albrecht, M. Computing topological parameters of
biological networks. *Bioinformatics (Oxford, England)* **24**, 282 (2008).
https://doi.org/10.1093/bioinformatics/btm554
78   Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *Journal of Machine Learning Research* **12**, 2825
(2011).
79   Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization system for the whole mouse
brain. *Nature Neuroscience* **13**, 133 (2010). https://doi.org/10.1038/nn.2467
80   Racine, R. J. Modification of seizure activity by electrical stimulation. II. Motor seizure. *Electroencephalography*
and Clinical Neurophysiology **32**, 281 (1972). https://doi.org/10.1016/0013-4694(72)90177-0
81   Basu, T. *et al.* Hypothalamic-pituitary-adrenal axis dysfunction worsens epilepsy outcomes and increases SUDEP
risk. (Neuroscience, 2022).

1137
1138
1139
1140

SUPPLEMENTAL FIGURES

S1. Synapsin1-Cre expression and recombination occur in neurons, and conditional knockout diminishes EZH2 induction post-SE.

S2. Neuronal ablation of EZH2 does not alter latency to 1st Class V seizure during kainate injections.

S3. Ontology networks for the mouse-Black and human-Brown clusters.

S4. Transient induction of STAT3 in wildtype animals is observed across models of SE.

S5. Other STAT family proteins are hyper-induced in EZH2nKO mice.

S6. Western blot quantifications in shNT and shEZH2 N2A cells.

S7. JAK2 activity is not required for hyper-activation of STAT3 in the absence of EZH2.

S8. Western blot quantifications in N2A cells transfected with control plasmids or overexpressing JAK1.

S9. H3K27Me3 ChIP-PCR at JAK1 (Sites 1 and 3), JAK2, and STAT3.

S10. CP treatment window selection and target engagement.