IBAS: Interaction-bridged association studies discovering novel genes
underlying complex traits

Pathum Kossinna[1,#], Senitha Kumarapeli[2,3], Qingrun Zhang[1,3,4,*]

[1]Department of Biochemistry & Molecular Biology, University of Calgary, Calgary, Alberta,
T2N 1N4, Canada

[2]Schulich School of Medicine & Dentistry, University of Western Ontario, London, Ontario,
N6A 5C1, Canada

[3]Department of Mathematics and Statistics, University of Calgary, Calgary, Alberta, T2N
1N4, Canada

[4]Alberta Children's Hospital Research Institute, University of Calgary, Calgary, Alberta, T2N
1N4, Canada

* Corresponding author: qingrun.zhang@ucalgary.ca

Abstract
The contribution of genetic variants to a complex phenotype may be mediated by various
forms of complicated interactions. Currently, the discovery of genetic variants underlying
interaction is limited, partly due to that the real interaction patterns are diverse and unknown,
whereas exhaustively examining all potential combinations confers the risk of overfitting and
instability. We propose IBAS, Interaction-Bridged Association Study, a new model using
statistical learning techniques to extract representations of interaction patterns in
transcriptome data, which act as a mediator for the next genotype-phenotype association test.
Using simulated perturbation experiments, it is demonstrated that IBAS is more robust to
noise than similar mediation-based protocols replying on single-genes, i.e., transcriptome-
wide association studies (TWAS). By applying IBAS to real genotype-phenotype and
expression data, we reported additional genes underlying complex traits as well as their
biological annotations. IBAS unlocks the power of integrating gene-gene interactions in
association mapping without concerning overfitting and instability.

Keywords
Genotype-phenotype association study/Stability model/Transcriptome/Gene-gene interactions

Introduction

Discovering genetic basis of complex traits is a long-standing theme in genetics and genomics (Glazier *et al*, 2002). Towards this end, genotype-phenotype association mapping has emerged as an essential tool over the last 20 years (Klein *et al*, 2005; Young *et al*, 2019). In recent times, while identifying genes statistically associated with diseases remains a primary goal, researchers have gradually shifted their focus to their underlying mechanisms through various means of interpretations (Sheng *et al*, 2021). Towards this aim, the availability of multi-scale -omics data especially transcriptome allows researchers to develop sophisticated statistical methods to improve both the discovery of genes and their interpretations (Gamazon *et al*, 2015; Zeng *et al*, 2017; Xie *et al*, 2017; Brandes *et al*, 2020; Okada *et al*, 2016; Xu *et al*, 2017). Among many biologically sensible hypothetical mechanisms, it is intuitive to expect that the variation of gene-gene interaction networks would play a critical role mediating genetic effects to phenotype. Methods such as iPath (Su *et al*, 2021) take this premise and study it at the transcriptomic level, investigating perturbations of given pathways at the sample level and the effect on clinical phenotypes. However, despite gene-gene interactions having been studied for a century (Bateson William *et al*, 1909; Fisher, 1919), no consensus has been reached in understanding the genetic basis of variation of interactions and its implications to phenotypic changes.

As far as our understanding, the lack of statistical genetic models studying the genetic basis of gene-gene interactions is due to the astronomic number of potential combinations, which leads to serious problem of instability. Indeed, studying the interaction patterns associated with phenotypes is already a $p \gg n$ problem, suffering from statistical instability as well as computational burden (Fang *et al*, 2019; Zhang *et al*, 2014). By adding an extra layer of the genetic basis of variation of interaction patterns, it only adds substantial risk of instability, potentially leading to many results that may not be robust to perturbation of input data.

Transcriptome-Wide Association Study (TWAS) is a popular model to conduct association mapping utilizing transcriptomes (Gamazon *et al*, 2015; Gusev *et al*, 2016). Although it only focuses on single genes, the in-depth characterization of TWAS by our group (Cao *et al*, 2021a, 2022) offered a unique insight of developing models incorporating interactions. The mainstream format of TWAS is a two-step protocol: first, one can form an expression prediction model using (usually cis) genotypes in a reference dataset (e.g.: GTEx (Consortium *et al*, 2015)). The predicted expression is called Genetically Regulated eXpression, or GReX. Then, in the second step, in the main dataset for association mapping (that does not contain expressions), one can predict expressions using the available genotypes and finally associate the predicted expression to phenotype.

Our recent works showed this interpretation of "predicting expressions" was not the essence of TWAS, but rather, misleading. First, theoretical power analysis showed that TWAS could be more powerful than a hypothetical scenario in which expression data is available in the main GWAS dataset (Cao *et al*, 2021a). Additionally, TWAS could be underpowered compared to GWAS when the expression heritability is low (Cao *et al*, 2021a). Both results question the interpretation of the prediction of expression in TWAS. As such, we proposed to interpret the "prediction" step as a selection of genetic variants directed by expression. From the perspective of Machine Learning, the first step in TWAS is essentially feature selection and the second, feature aggregation. With this interpretation in mind, one may disregard GReX and instead conduct feature selection and aggregation independently. As a result, methodological research focusing on optimal combinations of feature selection and aggregation approaches could be flexibly conducted. Indeed, novel methods splitting these

two steps developed by us (Cao *et al*, 2021b, 2022) and independently by others (Tang *et al*,
2021) showed higher power than standard TWAS.

The above unique insight into TWAS has opened a door of conducting association mapping
mediated by any "data-bridge" by replacing the gene expression and its predicted value with
another entity for feature selection and a form of feature aggregation (Cao *et al*, 2022).
Indeed, another development in our group has provided a useful tool leveraging brain images
(He *et al*, 2023) in a similar manner.

In this work, we designed an unconventional framework by forming a mediator representing
interactions computed using gene expressions through statistical learning techniques. We
then used it to facilitate the discovery of the genetic basis of variations of interactions and
their consequence to phenotypic changes. More specifically, starting from a gene expression
matrix of all genes in a prespecified pathway, we extract representative statistics such as PCA
(Hotelling, 1933), t-SNE (der Maaten & Hinton, 2008) or UMAP (McInnes *et al*, 2018) to
form the objective function for feature selection. Then, following our previous work (Cao *et*
al, 2021b, 2022; He *et al*, 2022), a kernel machine (Wu *et al*, 2010) is employed for feature
aggregation aimed at association test. The outcomes are single genes whose genetic variants
alter the variation of gene-gene interactions in the focal pathway, which also impact the
phenotypic changes. This method is named "Interaction Bridged Association Study", or
IBAS. Importantly, such representative statistics contributed by many genes in the pathway is
more robust to noise than single genes alone, therefore intuitively could be more stable
mediators than single genes in a typical TWAS protocol. Indeed, perturbation experiments
using GTEx data show that IBAS is more stable than standard TWAS protocols that rely on
single genes. We further applied IBAS to the seven diseases of the Wellcome Trust Case
Control Consortium, or WTCCC (The Wellcome Trust Case Control Consortium, 2007) and
discovered novel genes underlying pathways and diseases.

Results
The IBAS Framework
Step 1 of IBAS utilizes a reference dataset where gene expression and genotype data are
available. Known biological pathways (or even user specified collection of genes known to
contain interactions) are then used to extract the expression data. The extracted subset of gene
expressions (corresponding to a particular pathway) then undergoes dimensionality reduction
(using methods such as PCA, t-SNE or UMAP) to produce interaction components ("data-
bridges") (**Fig 1A**). These components are expected to capture interactions at the
transcriptome level. In Step 2, each component is then associated with individual *cis*-variants
corresponding to the genes in the pathway (**Fig 1B**). The coefficients obtained for each
variant indicate their contribution to the interactions captured by the corresponding
interaction component. Step 3 of IBAS involves a GWAS dataset where genotype data is
assessed together with a phenotype of interest. Variants in the GWAS dataset are then
aggregated to the gene level, weighted by their contribution to interactions (obtained in Step
2 depicted in **Fig 1B**) and tested for association with phenotypes using the Sequence Kernel
Association Test, or SKAT (Wu *et al*, 2010) (**Fig 1C**). This results in the identification of
genes associated with the phenotype of interest through the mediation of the pathway being
considered.

The natural cascade of genetic variants affecting phenotype may be depicted in **Fig 1D,**
upper panel: from variants to gene expression, to pathways and finally phenotype. TWAS
captures the effects of variants on gene expression, and then gene expression on phenotype,
ignoring the contribution of interactions at the pathway level (**Fig 1D, middle panel**). IBAS

on the other hand, captures the effects of cis-variants on the interactions at the pathway level
(where it could be argued that the gene-level effect itself is present) and identifies association
with the phenotype (**Fig 1D, lower panel**). Therefore, IBAS reveals the genetic basis of
complex traits from a compensatory angle.

IBAS is more stable than single gene-mediated TWAS
To test whether interaction-based protocol is more stable than single-gene based ones,
perturbation experiments were conducted to compare the stability of IBAS to PrediXcan with
respect to different reference data sets (**Fig 2**). We utilized real data from the GTEx
Consortium for building the data-bridge and investigate the effects on the Type 1 Diabetes
(T1D) cohort from the WTCCC dataset (along with the other 6 cohorts in the **Appendix Fig**
S1-S6 to ensure this stability holds across different genetic architectures). To simulate
various signal-to-noise ratios, expression data in the GTEx dataset was perturbed by adding
uniform noise of varying ranges ($= 0.1, 0.25$ and 0.5) relative to the expression levels of
each gene (**Materials & Methods**) to produce 10 replicates each for every noise-level.
PrediXcan elastic-net models were trained on these data sets, and IBAS weights were
obtained for dimensionality reduced data-bridges of the same. These models, and weights
were then used for association analysis in WTCCC T1D data. In the case of IBAS, for each
dimensionality reduction method, weights were combined for all components by mean for
coefficient-based and Fisher combined p-values for the p-value based method (we call this
combination 'meta'-weights) (**Materials & Methods**).

IBAS-based methods showed clear advantages over PrediXcan in stability of discoveries. **Fig**
2A showcases that IBAS has lower variance in p-values over a wide range of noise levels for
genes identified as significant ($p < 0.05$) in at least one replicate. PCA-based IBAS
demonstrates a significantly lower variability in p-values. It is also notable that weighting by
p-value, rather than by coefficient across all dimensionality reduction methods reduces
variability of p-values. We also look at the recurrence percentage (**Fig 2B**) of genes that were
significant in at least one replicate and uncover a similar pattern. PrediXcan performs poorly
indicating that a gene uncovered in one replicate may not be uncovered again if there was
even a slight perturbation in the reference data. IBAS is significantly more stable – with
PCA-based, p-value weighted IBAS performing the best (**Fig 2A**).

IBAS discovers novel genes through the mediation of interactions
IBAS uncovered significant discoveries across the 7 diseases of the WTCCC case-control
cohort; a number of which were previously identified in the DisGeNET database (Piñero *et*
al, 2017) with an additional number of discoveries that were novel (**Fig 3**). Most discoveries
were for the Rheumatoid Arthritis (RA) and T1D cohorts both of which also contain the most
annotations in the DisGeNET database being autoimmune diseases that have been
extensively studied. However, IBAS was also able to uncover significantly associated genes
for Bipolar Disorder (BD), Coronary Artery Disease (CAD), Hypertension (HT) and Type-2
Diabetes (T2D). Furthermore, the discovery of numerous novel genes is expected and
welcomed since most association studies incorporate purely genetic information (or, in the
case of TWAS, transcriptomic information of a single gene).

IBAS shows contrastive results compared to other gene-disease association tools (**Fig 4**).
PrediXcan (Gamazon *et al*, 2015) and kTWAS (Cao *et al*, 2021b) are used for this
comparison since they represent analyses based on the typical TWAS framework and a
kernel-based framework similar to IBAS but without the interaction effects, respectively.
Results from IBAS are once again based on the meta-weights (mean for coefficient based and
Fisher combined p-values for p-value based across all dimensionalities reduced components).

kTWAS and PrediXcan were found to report fewer results in most diseases, but with a level
of overlap between them that suggests that while kTWAS does gain more power due to the
use of the kernel-based test for feature aggregation as shown previously (Cao *et al*, 2021b),
they extract similar information from transcriptomic data. IBAS on the other hand,
incorporates interactions from gene expression and thus is expected to discover genes based
on their association with these interactions. The high number of genes discovered in T1D and
RA by both kTWAS and PrediXcan further support the theory previously mentioned that
there may be a high marginal effect from genes associated with these diseases. It is also
noteworthy that IBAS uncovers several genes associated with HT: one of which, *CLOCK*,
has known associations with HT in the DisGeNET database (**Fig 3**). This further strengthens
the hypothesis that IBAS is able to effectively uncover genes that may be associated with
disease through their interactions at the pathway level.

Biological annotation of results
IBAS distinguishes itself from previous annotations in the WTCCC cohort by identifying
genes that may have a small marginal effect but contribute to disease phenotype mediated
through relevant pathways (**Fig 5**). In particular, observing the patterns in diseases where
many significant genes are identified such as RA and T1D, previously annotated diseases are
identified across multiple pathways – indicating that their strong marginal effect may have
contributed to their significance. This could explain the discovery of a large number of genes
in the HLA region being identified as significant for these auto-immune disorders (Serrano *et*
al, 2006) as well as *TNF* (Serrano *et al*, 2006) and *ITPR3* (Huang *et al*, 2010). The weaker
marginal effects of genes may explain the lower number of genes identified to be associated
with the other diseases since most methods do not account for pathway-level interactions as
IBAS does.

IBAS identifies a key mechanism suggesting the mediation of metabolism related pathways
and genes in BD. 4 out of the 5 novel genes identified by IBAS are associated with various
metabolic processes (GPI Gene - GeneCards | G6PI Protein | G6PI Antibody; Yang *et al*,
2019; Dupuy *et al*, 2015; Zempleni *et al*, 2014). The role of metabolism in psychiatric
disorders has long been investigated (Zuccoli *et al*, 2017) with a clear association established
between dysfunctional metabolism and psychiatric disorders including BD (Rosso *et al*,
2015).

HLCS, a gene known for its role in biotin metabolism (Zempleni & Kuroishi, 2012) in the
body, is found to be highly associated with BD through the biotin metabolism pathway
(Kanehisa Laboratories; Kanehisa & Goto, 2000) by IBAS indicating a potential pathway that
could mediate the development of BD. Biotin or Vitamin B7 is known to be vital to brain
function as it is a crucial component in the process of glucose metabolism (Kennedy, 2016).
Furthermore, *HLCS* is highly specific to neurons (both inhibitory and excitatory) in the brain
according to the Human Protein Atlas (Karlsson *et al*, 2021; The Human Protein Atlas).
Changes in the excitatory and inhibitory (E/I) synaptic balance have been previously
associated with bipolar disorder in animal models (Lee *et al*, 2018) and the specificity of
HLCS as well as its importance in metabolism make it a potential mediator in the emergence
of the BD phenotype. While no studies have been conducted on the direct association of
HLCS on BD, one study identified increased methylation at CpG sites in *HLCS* in borderline
personality disorder patients in their genome-wide association study but failed to find the
same in a validation analysis (Teschler *et al*, 2013). Increased methylation could indeed be a
potential mechanism of repression of *HLCS* expression, resulting in dysfunctional biotin-
dependant metabolic pathways which in turn could create an imbalance in the activation of
E/I neurons contributing to manic episodes.

In the case of other significant associations, IBAS hints at potential biologically pivotal
candidates in disease etiology. *VARS2*, the most significant gene identified with association to
CAD (excluding all previously annotated genes in DisGeNET) has been identified to cause
heart failure in zebrafish models when knocked-out (Kayvanpour *et al*, 2022). Another
significant gene, *ENPP3*, has known cardiac-specific function and is downstream of *NKX2-5*,
a transcription factor associated with congenital heart disease (Barth *et al*, 2010). *PPAT*,
identified to be significant in the HT phenotype, was concluded to be an important gene in
the control of BP rhythm in rats (Murata *et al*, 2020). A significant gene (*ACSL5*) identified
in T2D, has been studied to be the primary target of the most strongly associated variant of
T2D in the *TCF7L2* gene (Xia *et al*, 2016), and is important in the metabolism of fatty acids.

Discussion
Stability and validations
We expect that the development of IBAS is a significant progress towards the goal of
developing statistically stable models involving interactions. This assumption is because
IBAS does not aim to explore all combinations of interactions and therefore does not run the
risk of overfitting at this stage. Indeed, we have revealed that the performance of IBAS is
stable with respect to the alternation of input expression data (**Fig 2**). Additionally, as t-SNE
and UMAP can extract nonlinear components incorporating interactions flexibly, its
performance should be also stable to alternations of the prior knowledge of membership of
pathways.

More rigorous validations are needed to quantitatively support the above assumptions. First,
numerical simulations with known genetic architectures must be conducted to assess the
theoretical properties of IBAS. This is not a trivial task as there is very limited literature
reporting patterns of gene-gene interactions that involve many genes in a pathway. During the
development of IBAS, we have carried out simulations following the three non-linear genetic
architectures, i.e., epistatic, compensatory, and heterogenous models. These non-linear
models are better accepted by the field and have been used in the previous publications in our
group (Cao *et al*, 2021b, 2022; Li *et al*, 2020). However, these architectures are designed to
mimic interactions of two genes and are therefore not suitable for the purpose of verifying
IBAS. Instead, we opted to use perturbations of real data in our simulations since there was a
lack of literature and benchmarking surrounding the simulation of realistic pathway-level
interactions.

Association v.s. Causality
It is worth noting that, although IBAS used PCA/t-SNE/UMAP represented gene-gene
interactions to mediate the association test, we are not claiming the causality of "genetics ⇒
interaction patterns ⇒ phenotype". Instead, there could be all kinds of causality models, for
instance, the pleiotropic model in which genetics causes changes in both interaction patterns
and phenotype. Here, the key insight is that we used the interaction patterns to
methodologically mediate the discovery, despite the real biological causality relationship is
unknown. The same issue applies to many existing works leveraging multi-scale omics to
identify association between genetics and phenotype (Gamazon *et al*, 2015; Zeng *et al*, 2017;
Xie *et al*, 2017, 2016; Cao *et al*, 2021b, 2022; Brandes *et al*, 2020; Okada *et al*, 2016; Xu *et*
al, 2017).

Materials and Methods
IBAS Implementation (I): Biological pathways
Biological pathways refer to known interactions among genes, proteins and metabolites (van
Iersel *et al*, 2008). These interactions are often curated from literature by various projects

such as Kyoto Encyclopedia of Genes and Genomes (KEGG) (Kanehisa *et al*, 2017, 2023;
Kanehisa & Goto, 2000), Reactome (Joshi-Tope *et al*, 2005) or WikiPathways (Kelder *et al*,
2012). These databases provide access to both downloading and visualizing these pathways
and their known interactions. This work utilizes the KEGG database through the use of the
KEGGREST (Tenenbaum & Maintainer, 2022) R package. However, IBAS could be applied
in a similar manner to any of the pathway databases or even other databases containing
known biological interactions such as the Gene Ontology Resource (Consortium, 2019).

IBAS Implementation (II): Dimensionality Reduction
The downloaded pathways provide context for sub-setting the reference data, but we must
turn to dimensionality reduction to uncover the complex interactions within the pathways.
Here, we take advantage of three methods often used in the analysis of genomics data: PCA
(Hotelling, 1933), t-distributed Stochastic Neighbour Embedding (t-SNE) (der Maaten &
Hinton, 2008) and Uniform Manifold Approximation and Projection (UMAP) (McInnes *et al*,
2018). These methods are based on the concept of identifying a low-dimensional
representation of points in a high-dimensional space. We use these methods to reduce the
dimensionality of the pathways to a maximum of 10 components in the case of PCA and
UMAP, and 2 in the case of t-SNE. In instances where a pathway has less than 10 genes, we
set the number of components to be generated as the number of genes.

PCA utilizes the principle of iteratively identifying the combinations of components of the
higher-dimensional space that maximize captured variance. These components are identified
in such a way that they are uncorrelated with all previous PCs resulting in orthogonal
combinations of the original data (Lever *et al*, 2017). These components have been identified
in various fields to capture variance and identify differences in subpopulations of the data
(Reich *et al*, 2008; Jollife & Cadima, 2016).

t-SNE was originally designed for the purpose of visualizing high-dimensional data in a low-
dimensional space (typically 2-dimensional). t-SNE is now often used in the visualization of
transcriptomic data, particularly in RNA-Seq (Kobak & Berens, 2019). t-SNE is based upon
the concept of minimizing the probability distribution of points in a low-dimensional space
close to their probability distribution in a high-dimensional space. Consider a set of points x_i
where $i \in \{1, \ldots, n\}$ and $x_i \in \Re^p$ and their embedding in a lower-dimensional space y_i
where $y_i \in \Re^d$ and $d \ll p$.

The conditional probability between two distinct points x_i and x_j in the higher-dimensional
space is defined as:

$$300 \quad p_{j|i} = \frac{\exp{-\frac{\left\lVert x_i - x_j \right\rVert^2}{2\sigma_i^2}}}{\sum_{k \neq i} \exp{-\frac{\left\lVert x_k - x_j \right\rVert^2}{2\sigma_i^2}}} \tag{1}$$

301 and $p_{i|i} = 0$. We then define the joint probability of the two points x_i and x_j as:

$$302 \quad p_{ij} = \frac{p_{j|i} + p_{i|j}}{2n} \tag{2}$$

303 The joint probability of the same points in the lower dimension is then modelled using a
304 Student t-distribution with one degree of freedom as:

$$305 \quad q_{ij} = \frac{\left(1 + \left\lVert y_i - y_j \right\rVert^2\right)^{-1}}{\sum_{k \neq l}\left(1 + \left\lVert y_k - y_l \right\rVert^2\right)^{-1}}. \tag{3}$$

The Kullback-Leibler (KL) divergence between P and Q distributions is then used as the cost
function and optimized via gradient descent to obtain the optimal embedding.

UMAP uses concepts similar to t-SNE, to achieve a low-dimensional embedding. The
mathematics behind UMAP are fairly complex (McInnes *et al*, 2018), and for the purposes of
this paper are left out and instead, a high level explanation provided. UMAP begins by
constructing what is termed a "fuzzy simplicial complex" which is a type of weighted graph
between the points in the high-dimensional space with the edges denoting the likelihood that
two points are connected. To determine this connectivity (similar to **Equation 1**), UMAP
extends a radius determined by the distance to the nth nearest neighbour. Then, the likelihood
is reduced based on the distance from the point. It is also required that each point is
connected to at least its closest neighbour. The last step involves constructing a similar
network to the first, but this time in the lower dimensional space, with the use of cross-
entropy as the cost function compared with the KL-divergence used by t-SNE.

IBAS Implementation (III): Interaction Directed Feature Selection
Once a pathway has undergone dimensionality reduction, it is then in a form suitable for
association analysis to "select" relevant genetic variants (here we reference feature selection
with regards to feature prioritization). Using the same reference dataset, cis-variants are
tested for association with each individual component of the interaction components. Each
component is analyzed individually as it is expected that different components would capture
a different set of interactions among the gene expression in the pathway. While any number
of methods could be used for this association analysis, we use the linear mixed model (Kang
et al, 2008, 2010) which accounts for population stratification by the use of a kinship matrix
(which is also called a Genomic Relationship Matrix (GRM) in some literature).

The linear mixed model is defined as:

$$\boldsymbol{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{u}_g + \boldsymbol{\epsilon} \tag{4}$$

where $\boldsymbol{y}$ denotes the vector of phenotypes (interaction component in IBAS), $\mathbf{X}$ is the matrix
containing the intercept, variant to be tested and other covariates, $\boldsymbol{u}_g \sim \boldsymbol{N}(\mathbf{0}, \sigma_g^2 \mathbf{K})$ denotes the
random effects and $\boldsymbol{\epsilon} \sim N(\mathbf{0}, \sigma_e^2 \mathbf{I})$. Here $\mathbf{K}$ denotes the kinship matrix which provides a
measure of the degree of relatedness of the individuals in the sample.

This association is carried out for all cis-variants of the genes in the pathway. While it is
possible that trans-variants also do influence the expression of the genes in the pathway, such
testing is extremely resource-intensive and would suffer from a large multiple-testing
problem (Goeman & Solari, 2014). Typically, cis-variants are considered as the variants
within 1Mb of the start and end of the gene and we used this definition for our analysis as
well.

Genotype-Phenotype Association Analysis via Feature Aggregation
Finally, gene-level testing is conducted using the Sequence Kernel Association Test (SKAT).
This test combines the effect of multiple variants in a kernel-based score test to test
phenotypic association of the aggregate:

$$Q = \boldsymbol{y}^T \mathbf{K} \boldsymbol{y} \tag{5}$$

where y is the vector of phenotypes and $\mathbf{K}$ is a kernel function based on the centralized
genotype matrix of the GWAS dataset, $\mathbf{G}$. While multiple kernels may be defined for the use
in the score test (Wu *et al*, 2010), we use the modified kernel previously used in our work
(Cao *et al*, 2021b, 2022):

$\mathbf{K_W} = \mathbf{GWG}^T$ (6)

where $\mathbf{W} = \text{diag}(\alpha_1, \ldots, \alpha_i, \ldots, \alpha_p)$. α_i here, denotes the coefficients of association derived
from **Equation 4** for the reference data. These coefficients of association may be selected as
the direct coefficients from the linear mixed model (i.e. $\alpha_i = |\beta_i|$; coefficient-based
weighting) or as the p-values associated with each coefficient from the same (i.e. $\alpha_i = -\log p_i$; p-value based weighting). This is then used in **Equation 5** to calculate Q which
follows a mixture of chi-square distributions (Davies, 1980) (**Fig 1C**).
356

While weights associated with individual components of each dimensionality reduction
method could be used to identify genes effecting phenotypes through various interactions, a
'meta'-weight was also considered for each pathway. In this case, the weights for each
variant were calculated as the mean of the coefficients in the case of coefficient-based
weighting, and in the case of p-value based weighting, the Fisher combined p-values. These
'meta'-weights could be assumed as the average contribution of each variant on the major
interactions in the pathway under study.

Thus, the outputs of IBAS are genes associated with phenotypes of interest, mediated by the
interaction effects of the pathway considered. Since a single pathway may contain up to
hundreds of genes, a multiple hypothesis testing correction in the form of a Benjamini-
Hochberg correction (Benjamini & Hochberg, 1995) is applied to reduce the risk of false
positives.

Other Methods compared to IBAS
TWAS: PrediXcan
PrediXcan (Gamazon *et al*, 2015) trains an elastic-net (Zou & Hastie, 2005) predictive model
of gene expression levels on reference data in a similar manner to IBAS as described before.
Considering the expression of gene j to be denoted as T_j, the PrediXcan model can be stated:

$T_j = \sum_k w_k \mathbf{X}_k + \epsilon$ (7)

where X_k denote the (*cis*-)variants of gene j. Once the model is trained, it is used to predict
the Genetically Regulated eXpression (GReX) in a GWAS dataset. Finally, the predicted
expressions are associated with the phenotypes of interest using either linear, logistic or Cox
regression models or other non-parametric methods (such as testing for Spearman's
correlation) (Gamazon *et al*, 2015).

kernel-based TWAS (kTWAS)
kernel-based TWAS (kTWAS) (Cao *et al*, 2021b) was the first data bridge model proposed
by our group. Following the first steps of PrediXcan, an elastic-net model is trained on
reference data. However, instead of using this model to predict gene expression in the GWAS
dataset, the weights from the elastic-net model, $\mathbf{W}_{kTWAS} = \text{diag}(\gamma_1, \ldots, \gamma_i, \ldots \gamma_p)$, are used in
SKAT in the form of a weighted kernel matrix for the score test. Similar to IBAS, these
weights may be derived from the elastic-net model used in PrediXcan as $\gamma_i = w_i$ or using
weights derived from p-values associated with each term. Following the results in the original
paper, we use the coefficient-based analysis for comparison. Essentially, the kernel matrix
$\mathbf{K_w}$ is constructed as:

$\mathbf{K_w} = \mathbf{GW}_{kTWAS}\mathbf{G}^T$ (8)

and the score statistic Q is defined as:

$Q = \boldsymbol{y}^T \mathbf{K_w} \boldsymbol{y}$ (9)

which is distributed as a mixture of chi-squared distributions (**Fig 1C**) (Davies, 1980).

Real Data Analysis: Reference Data
Publicly available, normalized, filtered expression data of "Whole Blood" tissue from the
GTEx Consortium (Consortium *et al*, 2015) which contains samples from 670 individuals
and for 20315 transcripts was used as the reference expression data. This data was then
corrected for PEER factors (Stegle *et al*, 2010), the top 5 genotype principal components, age
and gender as suggested by the GTEx Portal. This expression data was then used for
dimensionality reduction by subsetting for individual pathways.

Genotype data for 838 individuals and 43,066,422 SNPs were present in the raw VCF data.
Since IBAS works best if there is a high degree of overlap between the GWAS and reference
datasets when considering genotypes, imputation was carried out using SHAPEIT4
(Delaneau *et al*, 2019) for phasing and IMPUTE5 (Rubinacci *et al*, 2020) for the final
imputation based on data from the 1000 Genome Project (Consortium & others, 2015). This
results in 88,863,451 SNPs for the imputed data. The data was then filtered to include only
individuals who had expression data available and general QC (MAF < 0.05, deviation from
HWE $p < 10^{-8}$ and relatedness > 0.025) was applied, resulting in 6,187,683 SNPs for 553
individuals. This data was converted to the {tped, tfam} file format and JAWAMix5 was
used to convert it to hdf5 (Long *et al*, 2013) for out-of-core association analysis with the
interaction components obtained from the expression data.

Real Data Analysis: GWAS Data Containing Genotype and Phenotype
The Wellcome Trust Case Control Consortium (The Wellcome Trust Case Control
Consortium, 2007) contains genotype data for 13,241 cases and 2,938 controls (individual
sample sizes are provided in **Appendix Table S1**) across 7 diseases: bipolar disease (BD),
coronary artery disease (CAD), Crohn's disease (CD), rheumatoid arthritis (RA), type 1
diabetes (T1D), type 2 diabetes (T2D) and hypertension (HT). The original dataset only
contained 393,272 variants since this data was obtained using the Affymetrix GeneChip
500K arrays and thus imputation (Rubinacci *et al*, 2020) was conducted as in GTEx genotype
data to produce 73,162,888 variants.

Simulations Generating Perturbated Data
Simulations were carried out by adding perturbations to gene expression data from the Whole
Blood tissue in GTEx (Consortium *et al*, 2015) to evaluate robustness to changes in reference
data. Each expression value was regenerated using a uniform distribution $U \sim \big(x(1 +$
$r), x(1 - r)\big)$ where x is the current expression value and $r = \{0.1, 0.25, 0.5\}$. This approach
attempts to hold the same expression patterns within pathways while adding individual noise
levels to genes as could be present due to various factors such as sampling time and
sequencing batch effects. The varying levels of r, adds increasing levels of noise and 10
replicates are generated at each level. These 30 individual expression datasets are then used
in conjunction with unperturbed genotype data in generating data-bridges for IBAS and in
training PrediXcan models for association testing with WTCCC cohorts.

Acknowledgements
Funding:
NSERC Discovery grant RGPIN-2018-05147 (QZ)

New Frontiers in Research Fund NFRFE-2018-00748 (QZ)

VPR Catalyst Grant (QZ)

Alberta Innovates LevMax Bridge Funds (222300769) (QZ)

Author contributions:
Designed the study: PK, QZ
Implemented the tool: PK
Analyzed data: PK, SK, QZ
Wrote the manuscript: PK and QZ

Competing interests: The authors declare that they have no conflict of interest.

Data and materials availability:
GTEx gene expression: https://gtexportal.org/home/datasets

GTEx whole genome sequencing data: https://www.ncbi.nlm.nih.gov/projects/gap/cgi-
bin/study.cgi?study_id=phs000424.v9.p2

WTCCC genotype: https://www.wtccc.org.uk/

IBAS source code available at: https://github.com/QingrunZhangLab/IBAS

References
Barth JL, Clark CD, Fresco VM, Knoll EP, Lee B, Argraves WS & Lee KH (2010)
Jarid2 is among a set of genes differentially regulated by Nkx2.5 during outflow
tract morphogenesis. *Developmental Dynamics* 239: 2024–2033

Bateson William, Mendel Gregor & Leighton Arthur G. (1909) Mendel's principles of
heredity, by W. Bateson Cambridge [Eng.], University Press, 1909

Benjamini Y & Hochberg Y (1995) Controlling the False Discovery Rate: A Practical
and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical*
Society: Series B (Methodological) 57: 289–300

Brandes N, Linial N & Linial M (2020) Pwas: Proteome-wide association study. In
International Conference on Research in Computational Molecular Biology pp
237–239.

Cao C, Ding B, Li Q, Kwok D, Wu J & Long Q (2021a) Power analysis of
transcriptome-wide association study: Implications for practical protocol choice.
PLoS Genet 17: e1009405

Cao C, Kossinna P, Kwok D, Li Q, He J, Su L, Guo X, Zhang Q & Long Q (2022)
Disentangling genetic feature selection and aggregation in transcriptome-wide
association studies. *Genetics* 220: iyab216

Cao C, Kwok D, Edie S, Li Q, Ding B, Kossinna P, Campbell S, Wu J, Greenberg M
& Long Q (2021b) kTWAS: integrating kernel machine with transcriptome-wide
association studies improves statistical power and reveals novel genes. *Brief*
Bioinform 22: bbaa270

Consortium 1000 Genomes Project & others (2015) A global reference for human
genetic variation. *Nature* 526: 68

Consortium GO (2019) The gene ontology resource: 20 years and still GOing strong.
Nucleic Acids Res 47: D330–D338

Consortium Gte, Ardlie KG, Deluca DS, Segrè A V, Sullivan TJ, Young TR, Gelfand
ET, Trowbridge CA, Maller JB, Tukiainen T, *et al* (2015) The Genotype-Tissue

Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science (1979)* 348: 648–660

Davies RB (1980) The Distribution of a Linear Combination of χ2 Random Variables. *J R Stat Soc Ser C Appl Stat* 29: 323–333

Delaneau O, Zagury J-F, Robinson MR, Marchini JL & Dermitzakis ET (2019) Accurate, scalable and integrative haplotype estimation. *Nat Commun* 10: 1–10

Dupuy F, Tabariès S, Andrzejewski S, Dong Z, Blagih J, Annis MG, Omeroglu A, Gao D, Leung S, Amir E, *et al* (2015) PDK1-dependent metabolic reprogramming dictates metastatic potential in breast cancer. *Cell Metab* 22: 577–589

Fang G, Wang W, Paunic V, Heydari H, Costanzo M, Liu X, Liu X, VanderSluis B, Oately B, Steinbach M, *et al* (2019) Discovering genetic interactions bridging pathways in genome-wide association studies. *Nat Commun* 10: 1–18

Fisher RA (1919) XV.—The correlation between relatives on the supposition of Mendelian inheritance. *Earth Environ Sci Trans R Soc Edinb* 52: 399–433

Gamazon ER, Wheeler HE, Shah KP, Mozaffari S V, Aquino-Michaels K, Carroll RJ, Eyler AE, Denny JC, Nicolae DL, Cox NJ, *et al* (2015) A gene-based association method for mapping traits using reference transcriptome data. *Nat Genet* 47: 1091–1098

Glazier AM, Nadeau JH & Aitman TJ (2002) Finding genes that underlie complex traits. *Science (1979)* 298: 2345–2349

Goeman JJ & Solari A (2014) Multiple hypothesis testing in genomics. *Stat Med* 33: 1946–1978

GPI Gene - GeneCards | G6PI Protein | G6PI Antibody

Gusev A, Ko A, Shi H, Bhatia G, Chung W, Penninx BWJH, Jansen R, De Geus EJC, Boomsma DI, Wright FA, *et al* (2016) Integrative approaches for large-scale transcriptome-wide association studies. *Nat Genet* 48: 245–252

He J, Antonyan L, Zhu H, Li Q, Enoma D, Zhang W, Liu A, Cao B, MacDonald ME, Arnold PD, *et al* (2023) A statistical method for image-mediated association studies discovers genes and pathways associated with four brain disorders. *bioRxiv*: 2023.06.16.545326

He J, Li Q, Cao C, Zhu H, Shang K, Liu A, Cao B, McDonald E, Arnold P & Long Q (2022) IMAS: A novel statistical method for image-mediated association studies – application to the UK Biobank images discovers image and genetic variants associated with four brain disorders

Hotelling H (1933) Analysis of a complex of statistical variables into principal components. *J Educ Psychol* 24: 417

Huang YC, Lin YJ, Chang JS, Chen SY, Wan L, Sheu JC, Lai CH, Lin CW, Liu SP, Chen CP, *et al* (2010) Single nucleotide polymorphism rs2229634 in the ITPR3 gene is associated with the risk of developing coronary artery aneurysm in children with Kawasaki disease. *Int J Immunogenet* 37: 439–443

van Iersel MP, Kelder T, Pico AR, Hanspers K, Coort S, Conklin BR & Evelo C
(2008) Presenting and exploring biological pathways with PathVisio. *BMC*
Bioinformatics 9: 1–9

Jollife IT & Cadima J (2016) Principal component analysis: a review and recent
developments. *Philosophical Transactions of the Royal Society A: Mathematical,*
Physical and Engineering Sciences 374

Joshi-Tope G, Gillespie M, Vastrik I, D'Eustachio P, Schmidt E, de Bono B, Jassal B,
Gopinath GR, Wu GR, Matthews L, *et al* (2005) Reactome: a knowledgebase of
biological pathways. *Nucleic Acids Res* 33: D428–D432

Kanehisa Laboratories KEGG PATHWAY: hsa00780.

Kanehisa M, Furumichi M, Sato Y, Kawashima M & Ishiguro-Watanabe M (2023)
KEGG for taxonomy-based analysis of pathways and genomes. *Nucleic Acids Res*
51: D587–D592

Kanehisa M, Furumichi M, Tanabe M, Sato Y & Morishima K (2017) KEGG: new
perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res* 45:
D353–D361

Kanehisa M & Goto S (2000) KEGG: Kyoto Encyclopedia of Genes and Genomes.
Nucleic Acids Res 28: 27–30

Kang HM, Sul JH, Service SK, Zaitlen NA, Kong S, Freimer NB, Sabatti C & Eskin E
(2010) Variance component model to account for sample structure in genome-
wide association studies. *Nat Genet* 42: 348–354

Kang HM, Zaitlen NA, Wade CM, Kirby A, Heckerman D, Daly MJ & Eskin E (2008)
Efficient control of population structure in model organism association mapping.
Genetics 178: 1709–1723

Karlsson M, Zhang C, Méar L, Zhong W, Digre A, Katona B, Sjöstedt E, Butler L,
Odeberg J, Dusart P, *et al* (2021) A single–cell type transcriptomics map of
human tissues. *Sci Adv* 7

Kayvanpour E, Wisdom M, Lackner MK, Sedaghat-Hamedani F, Boeckel JN, Müller
M, Eghbalian R, Dudek J, Doroudgar S, Maack C, *et al* (2022) VARS2 Depletion
Leads to Activation of the Integrated Stress Response and Disruptions in
Mitochondrial Fatty Acid Oxidation. *Int J Mol Sci* 23: 7327

Kelder T, Van Iersel MP, Hanspers K, Kutmon M, Conklin BR, Evelo CT & Pico AR
(2012) WikiPathways: building research communities on biological pathways.
Nucleic Acids Res 40: D1301–D1307

Kennedy DO (2016) B Vitamins and the Brain: Mechanisms, Dose and Efficacy—A
Review. *Nutrients* 8: 68

Klein RJ, Zeiss C, Chew EY, Tsai J-Y, Sackler RS, Haynes C, Henning AK,
SanGiovanni JP, Mane SM, Mayne ST, *et al* (2005) Complement factor H
polymorphism in age-related macular degeneration. *Science (1979)* 308: 385–389

Kobak D & Berens P (2019) The art of using t-SNE for single-cell transcriptomics.
Nat Commun 10: 1–14

Lee Y, Zhang Y, Kim S & Han K (2018) Excitatory and inhibitory synaptic dysfunction in mania: an emerging hypothesis from animal model studies. *Experimental & Molecular Medicine 2018 50:4* 50: 1–11

Lever J, Krzywinski M & Altman N (2017) Points of Significance: Principal component analysis. *Nat Methods* 14: 641–642

Li Q, Cao C, Perera D, He J, Chen X, Azeem F, Howe A, Au B, Yan J & Long Q (2020) Statistical model integrating interactions into genotype-phenotype association mapping: An application to reveal 3D-genetic basis underlying Autism. *bioRxiv*: 2020.07.27.222364 doi:10.1101/2020.07.27.222364 [PREPRINT]

Long Q, Zhang Q, Vilhjalmsson BJ, Forai P, Seren Ü & Nordborg M (2013) JAWAMix5: an out-of-core HDF5-based java implementation of whole-genome association studies using mixed models. *Bioinformatics* 29: 1220–1222

der Maaten L & Hinton G (2008) Visualizing data using t-SNE. *Journal of machine learning research* 9

McInnes L, Healy J & Melville J (2018) Umap: Uniform manifold approximation and projection for dimension reduction. *arXiv preprint arXiv:180203426*

Murata Y, Ueno T, Tanaka S, Kobayashi H, Okamura M, Hemmi S, Fuke Y, Matsumoto Y, Abe M & Fukuda N (2020) Identification of Clock Genes Related to Hypertension in Kidney From Spontaneously Hypertensive Rats. *Am J Hypertens* 33: 1136–1145

Okada H, Ebhardt HA, Vonesch SC, Aebersold R & Hafen E (2016) Proteome-wide association studies identify biochemical modules associated with a wing-size phenotype in Drosophila melanogaster. *Nat Commun* 7: 1–11

Piñero J, Bravo Á, Queralt-Rosinach N, Gutiérrez-Sacristán A, Deu-Pons J, Centeno E, García-García J, Sanz F & Furlong LI (2017) DisGeNET: A comprehensive platform integrating information on human disease-associated genes and variants. *Nucleic Acids Res* 45: D833–D839

Reich D, Price AL & Patterson N (2008) Principal component analysis of genetic data. *Nature Genetics 2008 40:5* 40: 491–492

Rosso G, Cattaneo A, Zanardini R, Gennarelli M, Maina G & Bocchio-Chiavetto L (2015) Glucose metabolism alterations in patients with bipolar disorder. *J Affect Disord* 184: 293–298

Rubinacci S, Delaneau O & Marchini J (2020) Genotype imputation using the positional burrows wheeler transform. *PLoS Genet* 16: e1009049

Serrano NC, Millan P & Páez MC (2006) Non-HLA associations with autoimmune diseases. *Autoimmun Rev* 5: 209–214

Sheng X, Guan Y, Ma Z, Wu J, Liu H, Qiu C, Vitale S, Miao Z, Seasock MJ, Palmer M, *et al* (2021) Mapping the genetic architecture of human traits to cell types in the kidney identifies mechanisms of disease and potential treatments. *Nat Genet* 53: 1322–1333

601   Stegle O, Parts L, Durbin R & Winn J (2010) A Bayesian framework to account for
602   complex non-genetic factors in gene expression levels greatly increases power in
603   eQTL studies. *PLoS Comput Biol* 6: e1000770

604   Su K, Yu Q, Shen R, Sun SY, Moreno CS, Li X & Qin ZS (2021) Pan-cancer analysis
605   of pathway-based gene expression pattern at the individual level reveals
606   biomarkers of clinical prognosis. *Cell Reports Methods* 1: 100050

607   Tang S, Buchman AS, De Jager PL, Bennett DA, Epstein MP & Yang J (2021) Novel
608   Variance-Component TWAS method for studying complex human diseases with
609   applications to Alzheimer's dementia. *PLoS Genet* 17: e1009482

610   Tenenbaum D & Maintainer BP (2022) KEGGREST: Client-side REST access to the
611   Kyoto Encyclopedia of Genes and Genomes (KEGG). [PREPRINT]

612   Teschler S, Bartkuhn M, Künzel N, Schmidt C, Kiehl S, Dammann G & Dammann R
613   (2013) Aberrant Methylation of Gene Associated CpG Sites Occurs in Borderline
614   Personality Disorder. *PLoS One* 8: e84180

615   The Human Protein Atlas Single cell type - HLCS.

616   The Wellcome Trust Case Control Consortium (2007) Genome-wide association study
617   of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 447:
618   661–678

619   Wu MC, Kraft P, Epstein MP, Taylor DM, Chanock SJ, Hunter DJ & Lin X (2010)
620   Powerful SNP-set analysis for case-control genome-wide association studies. *The
621   American Journal of Human Genetics* 86: 929–942

622   Xia Q, Chesi A, Manduchi E, Johnston BT, Lu S, Leonard ME, Parlin UW, Rappaport
623   EF, Huang P, Wells AD, *et al* (2016) The type 2 diabetes presumed causal variant
624   within TCF7L2 resides in an element that controls the expression of ACSL5.
625   *Diabetologia* 59: 2360–2368

626   Xie R, Quitadamo A, Cheng J & Shi X (2016) A predictive model of gene expression
627   using a deep learning framework. In *2016 IEEE International Conference on
628   Bioinformatics and Biomedicine (BIBM)* pp 676–681.

629   Xie R, Wen J, Quitadamo A, Cheng J & Shi X (2017) A deep auto-encoder model for
630   gene expression prediction. *BMC Genomics* 18: 39–49

631   Xu Z, Wu C, Pan W, Initiative ADN & others (2017) Imaging-wide association study:
632   integrating imaging endophenotypes in GWAS. *Neuroimage* 159: 159–169

633   Yang JS, Hsu JW, Park SY, Lee SY, Li J, Bai M, Alves C, Tseng W, Michelet X, Ho
634   IC, *et al* (2019) ALDH7A1 inhibits the intracellular transport pathways during
635   hypoxia and starvation to promote cellular energy homeostasis. *Nat Commun* 10

636   Young AI, Benonisdottir S, Przeworski M & Kong A (2019) Deconstructing the
637   sources of genotype-phenotype associations in humans. *Science (1979)* 365:
638   1396–1400

Zempleni J & Kuroishi T (2012) Biotin. *Advances in Nutrition* 3: 213–214

Zempleni J, Liu D, Teixeira Camara D & Cordonier EL (2014) Novel roles of
holocarboxylase synthetase in gene regulation and intermediary metabolism. *Nutr*
Rev 72: 369

Zeng P, Zhou X & Huang S (2017) Prediction of gene expression with cis-SNPs using
mixed models and regularization methods. *BMC Genomics* 18: 1–11

Zhang Q, Long Q & Ott J (2014) AprioriGWAS, a new pattern mining strategy for
detecting genetic variants associated with disease through interaction effects.
PLoS Comput Biol 10: e1003627

Zou H & Hastie T (2005) Regularization and variable selection via the elastic net. *J R*
Stat Soc Series B Stat Methodol 67: 301–320

Zuccoli GS, Saia-Cereda VM, Nascimento JM & Martins-de-Souza D (2017) The
energy metabolism dysfunction in psychiatric disorders postmortem brains: Focus
on proteomic evidence. *Front Neurosci* 11: 277593

Figures

Figure 1 – Overview of IBAS Framework.

A) Gene expression matrices of a reference dataset (such as GTEx) are subset for known
biological pathways (from sources such as KEGG, GO, etc.) and undergo dimensionality
reduction (using methods such as t-SNE and UMAP) to produce interaction components
(“data-bridges”). **B)** Individual interaction components are then associated with *cis*-variants

of the genes in the pathway. The coefficients (or significance) of association are interpreted
as the contribution of the variants towards the interaction effects at the transcriptome level.
C) Genotypes in a GWAS dataset are then tested for association with a phenotype of interest,
aggregated at the gene level using the Sequence Kernel Association Test, weighted by the
interaction contributions obtained in (B). **D)** The natural cascade of genetic effects affecting
phenotype are modelled to various extents by existing methods. TWAS methods capture the
effect of variants on individual genes' expression and ignore the contributions of interactions
at the pathway level on phenotype while IBAS captures the effect of variants on these
interactions (where the individual gene contribution may be also present).

     Figure 2 – Simulated Perturbations reveal robustness of IBAS to noise in reference
data.

Simulations were conducted by perturbing gene expression values of each gene by percentage
ranges (0.1, 0.25 and 0.5) in the reference data. 10 perturbed expression datasets (replicates)
were generated for each of the noise levels and variability of results for case-control
association with the Wellcome Trust Case Control Consortium Type 1 Diabetes cohort was
evaluated (other cohorts were tested and available in **Appendix Fig S1-S6**) using both IBAS
and PrediXcan. PCA, t-SNE and UMAP dimensionality reduction methods as well as p-value
based (p) and coefficient based (c) interaction association weights were tested in the case of
IBAS. **A)** Indicates the variance of observed p-values in genes identified as significant
(   ) in at least one replicate while **B)** indicates the recurrent significance of genes
identified as significant in at least one replicate.

Figure 3 – Evaluation of genes associated with disease uncovered by IBAS.

Genes identified as associated with each of the 7 diseases (columns) in the WTCCC dataset
(with GTEx used as reference) across both coefficient-based and p-value based (rows) for the
PCA data-bridge were annotated as "Rediscovered" if found as previously associated with the
same disease in the DisGeNET database. Remaining genes are annotated as "Novel".
Evaluation of t-SNE and UMAP can be found in **Appendix Fig S7-S8**.

Figure 4 – Comparison of IBAS results with kTWAS and PrediXcan.

Genes identified by each of the IBAS methods (PCA, t-SNE and UMAP) as well as PrediXcan and kTWAS were compared to identify overlap and uniqueness of discoveries across the 7 diseases of the WTCCC dataset.

Figure 5 – IBAS identifies associations with marginal effects as well as interaction-
driven effects.

Comparison of significance of previously discovered ("Rediscovered") and "Novel" genes
associated with the 6 disease (no significant genes associated with CD using IBAS-PCA)
cohorts of the WTCCC dataset. Combined P-values (y-axis) indicate Fisher combined p-
values of genes identified across multiple pathways; while the x-axis denotes the number of
pathways mediated through which the gene was identified in as associated with the
phenotype.