Haplotype Function Score improves biological interpretation and cross-ancestry polygenic prediction of human complex traits

Weichen Song[1,2✉], Yongyong Shi[2,3✉] and Guan Ning Lin[1✉]

1 Shanghai Mental Health Center, Shanghai Jiaotong University School of Medicine, School of Bioengineering, Shanghai Jiaotong University, Shanghai, China
2 Bio-X Institutes, Key Laboratory for the Genetics of Developmental and Neuropsychiatric Disorders (Ministry of Education), Collaborative Innovation Center for Brain Science, Shanghai Jiao Tong University, Shanghai, China
3 Biomedical Sciences Institute of Qingdao University (Qingdao Branch of SJTU Bio-X Institutes), Qingdao University, Qingdao, China

✉Correspondence:
Dr. Weichen Song, goubegou@sjtu.edu.cn
Prof. Yongyong Shi, shiyongyong@gmail.com
Prof. Guan Ning Lin, nickgnlin@sjtu.edu.cn

Haplotype Function Score improves biological interpretation

and cross-ancestry polygenic prediction of complex traits

Abstract

We propose a new framework for human genetic association studies: at each locus, a
deep learning model (in this study, Sei) is used to calculate the functional genomic
activity score for two haplotypes per individual. This score, defined as the Haplotype
Function Score (HFS), replaces the original genotype in association studies. Applying
the HFS framework to 14 complex traits in the UK Biobank, we identified 3,619
independent HFS-trait associations with a significance of $p<5\times10^{-8}$. Fine-mapping
revealed 2,699 causal associations, corresponding to a median increase of 63 causal
findings per trait compared with SNP-based analysis. HFS-based enrichment analysis
uncovered 727 pathway-trait associations and 153 tissue-trait associations with strong
biological interpretability, including "circadian pathway-chronotype" and
"arachidonic acid-intelligence". Lastly, we applied LASSO regression to integrate
HFS prediction score with SNP-based polygenic risk scores, which showed an
improvement of 16.1% to 39.8% in cross-ancestry polygenic prediction. We
concluded that HFS is a promising strategy for understanding the genetic basis of
human complex traits.

Introduction

Genome-wide association studies (GWAS) have witnessed remarkable advancements
over recent years, both in terms of sample size and genetic discovery. However, the
elucidation of downstream mechanisms and subsequent applications still face certain
limitations(Visscher et al., 2017). One caveat is that the statistical power of GWAS on
a variant relies on its population frequency(Li et al., 2020; Null et al., 2022; Zhou et
al., 2022), whereas most variants with large effect size are rare(Zeng et al., 2021),
leading to insufficient discoveries. Moreover, Linkage Disequilibrium (LD) among
neighboring variants can significantly inflate false positive results(Nowbandegani et
al., 2022). The variability of LD structure among different populations further
compounds the challenges associated with training predictive models and discovering
causal genes. Lastly, most trait-relevant variants reside in non-coding
regions(Watanabe et al., 2019), which lack direct functional annotations as coding
variants. The prevalent approach to addressing this issue is to annotate each variant
based on its location within functionally significant regions(Finucane et al., 2015;
Grotzinger et al., 2022; Iotchkova et al., 2019; Weissbrod et al., 2020; Zheng et al.,
2022), such as transcription factor binding sites or enhancers. While this strategy has
considerably advanced the analysis, it's not optimal, as a variant's placement within a
functionally important region does not inherently signify that the variant has
substantial functional impacts.

The central dogma, proposing that DNA alterations' effects on phenotype are mediated via RNA and protein changes, offers a novel strategy to address these challenges. More precisely, by replacing the original genotypes in association studies with the aggregated impact of variants on transcription or functional genomics, the central dogma ensures the preservation of the majority of genetic information. This 'aggregated impact' offers several benefits for GWAS analysis: it provides direct biological interpretations, bypasses the effects of LD and population genetic history, and amalgamates information from both common and rare variants. One successful implementation of this strategy is Polygenic Transcriptome Risk Scores (PTRS)(Hu et al., 2022; Liang et al., 2022), which employ genetically determined transcription levels rather than genotypes to predict complex trait, and achieved remarkable portability. Nonetheless, the accuracy of imputing transcription levels from genotypes, given the sample size of currently available cohorts (such as the Genotype-Tissue Expression project, GTEx(Aguet et al., 2020)), remains limited (R2 around 0.1 for most genes)(Barbeira et al., 2018). Thus, the performance of PTRS is yet to reach its optimal potential.

Following the success of PTRS, we made one step forward to utilize functional genomics in this strategy. Compared with transcription levels, predicting genetically-determined functional genomic levels has achieved much higher accuracy by multiple recent deep learning (DL) studies(Avsec et al., 2021; Chen et al., 2022; Kelley, 2020; Yan et al., 2021; Zhou et al., 2018). These DL models utilize segments of the human reference genome as training samples, substantially increasing the sample size. Furthermore, functional genomics serve as a mediator between DNA and transcription, thus lessening the influence of non-genic factors such as the environment. Given these advancements, we propose that using the outputs of one of the state-of-the-art DL models, Sei(Chen et al., 2022), as the 'aggregated impact' in this novel strategy could effectively address the challenges aforementioned. Sei accepts a DNA sequence and computes multiple sequence class scores that represent different facets of the functional genomic activities of that sequence. This score integrates impacts from all variants, even those as rare as singletons, into one continuous variable, and is, in theory, unaffected by LD. In line with this notion, a recent similar strategy called cistrome-wide association study (CWAS) integrated variant-chromatin activity and variant-phenotype association to boost power of genetic study of cancer(Baca et al., 2022).

In this study, we present an analytical framework founded on this strategy (Figure 1) and implement it on complex traits in the UK Biobank to pinpoint causal loci and genes, decipher biological mechanisms, and devise cross-ancestry prediction models. We segmented the human reference genome into multiple 4,096 bp loci, generated DNA sequences for each locus for two haplotypes per individual, and employed Sei to compute the functional genomic activities of these sequences. We designated this activity score as the Haplotype Function Score (HFS) and analyzed the association between the HFS and each trait. Our findings confirm that the HFS framework offers a unique improvement in the biological interpretation and polygenic prediction of complex traits compared to classic SNP-based methods, thereby demonstrating its

103   value in genetic association studies.

Result

Overview of genome-wide HFS

We used the HFS framework to analyze imputed genotype data from the UK Biobank
(Figure 1). We segmented the human genome (hg38) into 617,378 discrete, non-
overlapping loci, each 4,096 base pairs long. Of these, 590,959 loci carried at least
one non-reference haplotype in the UKB cohort (see Method and Table S1). After
quality control, these loci contained approximately 1.2 billion haplotypes, with a
median count of 819 per loci (Figure S1). We then employed the deep learning
framework, sei(Chen et al., 2022), to compute sequence class scores for each
haplotype. In its sequence-mode, sei accepts DNA sequences in fasta format and
produces multiple distinct sequence class scores, 39 of which were included in our
study (Method). Our analysis identified significant variation in sequence class scores
across different loci. In fact, 49.7% of loci housed haplotypes whose sequence class
(as defined by the maximum of the 39 sequence class scores) differed from the
reference haplotype sequence class. Using the reference sequence class as a
benchmark, we noted that 16.8% of loci showed a difference between the maximum
and minimum haplotype scores that surpassed the score of the reference haplotype.
Moreover, the correlation between sequence class scores of adjacent loci was low,
with a median R^2 value of 0.013 (Figure S2), effectively reducing the impact of LD in
association studies. Further evaluation indicated that this low LD was led by two
factors: integration of rare variant impacts and segmentation. Firstly, excluding rare
variants from HFS caused the LD raised to median=0.14 (Method; Figure S2C).
Secondly, median LD of SNPs from adjacent loci was 0.06, which was significantly
higher than HFS LD (paired Wilcoxon $p=1.76\times10^{-5}$) but significantly lower than HFS
LD without rare variants (paired Wilcoxon $p<2.2\times10^{-16}$).

Expanding on the sequence class scores, we defined HFS for each locus.
Specifically, we computed the mean sequence class score of two haplotypes per
individual, reflecting an additive model. We selected the score corresponded to the
sequence class of reference sequence as the HFS of the corresponding locus, and its
association with each trait was computed using a generalized linear model. Simulation
analysis revealed that when a non-reference sequence class score was associated the
trait, reference class score could still capture median 70% of HFS-trait association R^2.
We applied this framework to 14 polygenic traits in the UKB British ancestry training
set (n=350,587; Table S2 and Method), identifying 16,597 significant HFS-trait
associations at a threshold of $p<5\times10^{-8}$ (n=15 for insomnia, n=7,573 for height; Table
S2), equating to roughly 3,619 independent associations. The most significant
associations were between the "promotor" score of chr7:121327898-121331994
(WNT16) and bone mineral density (BMD; regression beta=-0.02, $p<10^{-300}$), and the
"promotor" score of chr9:4760952-4765048 (AK3) and platelet count (beta=3.20,

$p=2.79\times10^{-262}$; Table S3).

When comparing HFS association with the standard SNP-based GWAS on the
same data, we found that 98% of significant HFS loci also harbored a significant SNP.
There were a few cases (n=0~5) where significant HFS loci did not harbored even
marginal SNP association (GWAS p>0.01), which were due to the lack of common
SNP in these loci. HFS association p value was higher than GWAS p value in 95 % of
significant loci, suggested that HFS did not improve power to detect marginal effect.
The genomic control inflation factor (λ_{GC}) for the HFS association test varied between
0.99 for asthma and 1.50 for height, closely resembling the SNP GWAS (Pearson
Correlation Coefficient [PCC]=0.91, paired t-test p=0.16; Method and Figure S3). We
concluded that HFS-based association tests had adequate power and do not introduce
additional p-value inflation.

Fine-mapping based on HFS
Based on these data, we applied SUSIE to fine-map the causal loci that were
associated with each of the 14 traits. We divided hg38 genome into 1,361 independent
blocks as defined by MacDonald et al.(MacDonald et al., 2022), and applied SUSIE
to loci HFS in each of these blocks (number of loci per block=4~2392). As shown in
Figure 2 and Table S4, we identified a total of 2,699 causal loci-trait associations at
the threshold of posterior inclusion probability (PIP)>0.95, hereafter referred to as
"causal loci". Compared with SNP-based functionally-aware fine-mapping methods
(PolyFun(Weissbrod et al., 2020) and SbayesRC(Zheng et al., 2022)), HFS-based
SUSIE detected -11 to 334 more causal signals (median=63, Table S5) for each trait.
We cautioned that these methods use summary statistics as input and are by nature
less sensitive than individual data-based methods. Yet, we suggested that such impact
would be mild, since we used in-sample LD reference (from UKB European sample).

Among these causal loci, only 22% were also lead loci in association analysis

(loci with the lowest p-value in 200kb region), and 58% had association p-value > 5×

10^{-8}. In line with previous SNP-based analysis(Weissbrod et al., 2020), this result
highlighted the importance of using causal signals instead of lead signals in post-
GWAS analysis. We found 67 causal loci showing pleiotropic effects on at least two
independent traits, including "CTCF-Cohesin" score of chr9:89596537-89600633 that
was associated with age at menarche, body mass index (BMI) and height (PIP>0.97;
Table S4). We also found that rare variants played an important role in the good find-
mapping performance of HFS: when variants with MAF<0.01 were removed, 55.3%
of the causal signals would be missed in HFS+SUSIE analysis.

When looking at the reference sequence class of loci, those with functional
importance were more likely to be causal loci, including "Promoter" (Odds Ratio

[OR]=2.33, $p=1.41\times10^{-14}$) and "Bivalent stem cell enhancer" (OR=2.22, $p=1.11\times10^{-8}$)

and "Transcribed region 1" (OR=1.71, $p=1.581\times10^{-10}$, Figure 2). Such functional

enrichment was even higher for pleiotropic loci ("Promoter": OR=7.20, $p=3.35\times10^{-5}$).

We also observed trait-specific patterns of such sequence class enrichment, such as
"CEBPB binding site" (Insomnia: OR=5.25, p=0.01) and "FOXA1/AR/ESR1 binding
site" (intelligence: OR=4.69, p=0.01, Figure 2 and Table S6). These results
demonstrated the expected functional patterns of causal loci, and indicated that HFS-
based fine-mapping was biologically interpretable and reliable.
Despite the functional enrichment, we applied several secondary analyses to
verify the reliability of HFS-based SUSIE result. Firstly, we took causal SNP fine-
mapped by PolyFun(Weissbrod et al., 2020) as positive control, and find that
compared with genomic region-matched control loci, causal loci were significantly

enriched for causal SNP (OR=1.33 to 5.08, Fisher test p=0.12 to $4.72{\times}10^{-52}$, Table S5).

Secondly, we calculated the heritability tagged by causal loci and PolyFun causal SNP
in independent test set (defined as the R^2 of linear regression; Method), and found that
causal loci tagged 38% to 251% more heritability than causal SNP (median=151%;
Table S5). This was not an artifact of larger number of causal loci, since the Akaike
information criterion (AIC) was similar between causal loci and causal SNP (paired t-
test p=0.36; Table S5). Thirdly, for traits with sufficient causal loci coverage, we also
applied Linkage Disequilibrium Score regression (LDSC) on independent GWAS
summary statistic to evaluate heritability enrichment in causal loci. On average,
causal loci showed 124-fold enrichment of heritability, significantly larger than
genomic region-matched control loci (124-fold vs 101-fold; p=0.0002, Method and
Figure S4). Lastly, we applied simulation analysis and found that HFS+SUSIE
showed similar advantages over SNP-based methods as in real data, with high
accuracy and low false positive rate (Supplementary materials).
We further applied a sliding-window analysis (step=2048 bp, Method) to test
whether HFS-based result is robust against the choice of sequence interval. 29.4% of
causal loci (PIP>0.95) in the original analysis were still causal in sliding window
analysis. 31.1% and 29.3% of causal loci whose 5' and 3' overlapping locus had
PIP>0.95 in sliding window analysis, respectively, while themselves were no longer
causal. Besides, HFS+SUSIE was also robust when the predefined number of causal
loci (L=2 to 10) was changed, and the number of detected loci were not changed.
Lastly, removing insertion and deletion would reveal 9% more significant association
($p{<}5{\times}10^{-8}$) but 4.7% less causal association (PIP>0.95), and slightly increased
inflation factor (Wilcoxon p=0.0001, Figure S4). Taken together, HFS-based SUSIE is
a powerful and robust strategy for individual data-based genetic fine-mapping.
217

Biological interpretation based on HFS
Pinpointing causal loci of complex traits provides the opportunity of analyzing the
biological mechanism of them. Thus, based on the HFS-based fine-mapping result, we
applied a linear regression model to analyze the underlying pathways, cell types and
tissues of each complex trait. For each locus, we annotated its relevance to a pathway
by combined SNP to Gene (CS2G) strategy(Gazal et al., 2022), and regressed the PIP
against this annotation, with a set of baseline annotations included as covariates,
similar to the LDSC framework(Finucane et al., 2018) (Method). After p-value

correction and recurrent pathway removal (Method), we detected a total of 727
pathway-trait associations (Figure 3A and Table S7). The most significant associations
were "megakaryocyte differentiation" with platelet count ($p=2.26\times10^{-34}$), "Insulin-like
growth factor receptor signaling pathway", "Endochondral ossification" with height
($p=4.95\times10^{-33}$ and 1.17×10^{-27}), "PD-1 signaling" with allergic disease ($p=5.55\times10^{-25}$),

and "major histocompatibility complex pathway" with asthma ($p=1.22\times10^{-23}$). In fact,
asthma and allergic disease were predominantly associated with more than 80
immune-related pathways. These associations were all in line with existing
knowledge of trait mechanism, and extended the understanding of their genetic basis.
For example, PD-1 has recently been suggested as potential targets of allergic diseases
like atopic dermatitis(Morales et al., 2021), but such association has not been
highlighted by previous genetic association studies.
For other traits, the most significant associations also replicated known
mechanisms, such as "osteoblast differentiation", "Wnt ligand biogenesis and
trafficking" with BMD ($p=4.59\times10^{-13}$ and 2.78×10^{-12}); "circadian pathway" with

chronotype ($p=4.25\times10^{-12}$); "calcium regulated exocytosis of neurotransmitter",

"Arachidonic acid metabolism" with intelligence ($p=5.52\times10^{-7}$ and 2.78×10^{-6});

"GPCR pathway" and "adipogenesis" with BMI ($p=4.97\times10^{-10}$ and 2.02×10^{-7}) and

"physiological cardiac muscle hypertrophy" with systolic blood pressure ($p=6.32\times10^{-11}$
). We also highlighted less significant association which provided novel insights,

such as "synaptic vesicle docking" and "neuron migration" with chronotype ($p=4.00\times$

10^{-7} and 4.55×10^{-7}), "Prostaglandins synthesis" with insomnia ($p=5.30\times10^{-9}$),

"behavioral response to cocaine" with alcohol intake ($p=3.39\times10^{-8}$) and "roof of

mouth development" and "glycoside metabolism" with FVC ($p=2.19\times10^{-12}$ and $5.73\times$

10^{-11}).
For cell type and tissue analysis (Figure 3B and Table S8), we applied the same
linear model to evaluate whether causal loci enriched in active chromatin regions of
each cell type (Method). We found 153 biologically interpretable associations with

complex traits. For example, fetal megakaryocyte ($p=5.67\times10^{-22}$) and child spleen

($p=2.15\times10^{-13}$) were found to be key cell type and tissue of platelet count. Systolic

blood pressure was significantly associated with multiple heart and artery tissues and

fetal cardiomyocyte ($p<1.63 \times 10^{-5}$), whereas allergic disease was associated with

multiple immune cells including natural killer, Treg and B cells ($p<4.79\times10^{-16}$). For

brain-related traits, we found 21 significant associations, 14 of which were from
central nervous system. For example, adult Hippocampus and cingulate gyrus were

both linked to alcohol intake, smoking and insomnia ($p<1.11 \times 10^{-5}$), whereas

chronotype was associated with embryonic brain germinal matrix ($p<8.68\times10^{-6}$) and

intelligence with embryonic neuron derived stem cell ($p<6.89\times10^{-7}$).

We also applied other modified strategies for this task but did not get satisfying
result. For example, using cS2G to link locus to gene lists specifically expressed in
each cell type suffered from scRNA dataset batch effect, whereas linear mix model
was less sensitive than standard linear model (Supplementary Materials).

Taken together, our result suggests that fine-mapping results based on HFS could
pinpoint the causal pathways, cell types and tissues underlying complex traits, and is
valuable for the biological interpretation of genetic association study.

Highlighted genes for complex traits

Enhanced power of fine-mapping and biological enrichment could reveal novel key
genes for trait mechanism study. Below we integrated fine-mapping result and their
functional annotation in several case studies to find causal signals and trait-relevant
genes in regions not resolved by previous genetic association studies.

In our study, platelet count had large number of causal loci (Figure 2) which
showed significant functional enrichment (Figure 3). To find key loci and genes
underlying platelet count, we focused on causal loci that overlapped with active
regions in "fetal megakaryocyte" and "child spleen tissue", and applied cS2G(Gazal
et al., 2022) to link them to two key pathways ("megakaryocyte differentiation" and
"platelet morphogenesis", Method and Figure 4A). We chose these annotations based
on p-value in biological enrichment analysis in Figure 3. A total of 25 loci were
highlighted (Figure 4A), which were recurrently linked to well-known platelet-
regulating genes like MEF2C, SH2B3, FLI1, RUNX1, THPO and NFE2. Among
them we noticed a less-studied gene RBBP5, a target of key transcriptome factor
MEF2C during megakaryopoiesis(Kong et al., 2019). Specifically, in 1q32.1 region,
HFS+SUSIE identified two loci with PIP>0.9 (Figure 4B). SNP-based association
also found significant association in this region, but SNP fine-mapping(Weissbrod et
al., 2020) could not resolve this signal and only found seven signals between PIP=0.1
to 0.5. This was unlikely a statistical inflation, since HFS-based association test p-
value was actually higher than SNP-based one (Figure S5). One of the causal loci,
chr1:47401806-47405902 (PIP=1), overlapped with spleen active chromatin and
harbored a cCRE in megakaryocyte, and was linked to RBBP5 and three other genes

(Table S9). RBBP5 is known to be involved in megakaryocyte differentiation during megakaryopoiesis and was regulated by MEF2C(Kong et al., 2019), but previous genetic association studies provided little evidence for its association with platelet count.

The major histocompatibility complex (MHC) region has long been a challenge of genetic association study due to its long-range LD, and is often excluded in fine-mapping tools. However, many disorders like Schizophrenia(Sekar et al., 2016) and immune diseases(Nawijn et al., 2011) are robustly associated with MHC region. In our HFS-based fine-mapping of asthma, we found 15 loci within MHC region had PIP>0.95, 11 of which overlapped with active chromatin regions in Treg or natural killer cells (Figure 4C and Table S10). This result showed good discrimination between causal and non-causal loci: despite these 15 likely causal loci, only six loci had PIP between 0.25 to 0.95. Since MHC region harbored a large number of genes, these causal loci were linked to as much as 105 potential target genes, which hindered the discovery of true targets. We further filtered them based on the involvement in pathway "TNFR2-NFKB pathway" and "innate lymphocyte [ILC] development", since these pathways were most significantly associated with asthma (Figure 3), even after excluding MHC region (p=2.57×10^{-13} and 1.39×10^{-17}). We found five genes (LTA, LTB, TNF, PSMB8 and PSMB9) that were predicted to be regulated by five causal loci overlapped with active chromatin regions (Figure 4C), which could be considered as potential key genes for further validation.

Similarly, we fine-mapped MHC region for other allergic diseases (Figure S6 and Table S11) and found potential key genes including HLA family and AGER. We also highlighted other gene-trait association not previously emphasized by GWAS, including GATA4 and NPPA (cardiac muscle hypertrophy) with SBP, ALOX5 (Arachidonic acid metabolism) with intelligence and CRY1 (circadian pathway) with chronotype, as further discussed in Table S12 to S14 and supplementary information.

On the other hand, HFS perform worse than SNP-based fine-mapping on exonic regions. Taking height as an example, PolyFun detected 125 causal SNPs (PIP>0.95) in the exonic regions, but only 16% (20) of loci that harbored them also reached PIP>0. 5 (11 reached PIP>0.95) in HFS+SUSIE analysis. Among the 105 loci that missed such signals (HFS PIP<0.5), 12 had a nearby loci (within 10kb) showing HFS PIP>0.95, which likely reflected false positive led by LD. Thus, SNP-based analysis should be prioritized over HFS in coding regions.

HFS-based polygenic prediction

Lastly, we analyzed the potentiality of HFS in polygenic prediction accuracy. Compared with state-of-the-art SNP-based polygenic risk score (PRS) algorithm LDAK-BOLT(Q. Zhang et al., 2021), HFS-based PRS (weighted by SUSIE posterior effect size) reached 47% to 90% of R^2 in independent European test set (meta-analyzed proportion=75.6%, 95% confidence interval=75.3%~75.8%, Figure S7). The gap between performance of HFS-based and SNP-based PRS reflected the fact that HFS only captured (the majority of) functional genomic alterations and missed the information of amino acid sequence and post-translational modification. We thus

proposed that integrating information from HFS and SNP could provide better
performance. Specifically, in the large European training set we trained SNP PRS
model by LDAK. Then, in a small tuning sample of target ancestry, we calculated per-
locus HFS prediction score of height (sum of HFS within this block, weighted by
SUSIE posterior effect size), then used machine learning to integrate them with
LDAK PRS into a final polygenic prediction score, hereafter referred to as
"HFS+LDAK". To choose the proper machine learning tools to achieve this goal, in
British European test set we applied LASSO, ridge regression and elastic net and
compared the result (Figure 5B). They gave comparable result with only difference of
R^2 around 0.01, and all of them were profoundly better than simple linear regression.
We chose LASSO as the algorithm in the formal analysis.

Using height as a representative trait, we first estimated the proportion of
variance captured by top loci, and found that HFS of loci with PIP>0.4 (n=5,101)
captured roughly 80% of variance explained by all genome-wide loci (n=1,200,024
corresponded to sling-window strategy; Figure 5A). We then calculated HFS+LDAK
in non-British European (NBE), South Asian (SAS), East Asian (EAS) and African
(AFR) population in UK Biobank, and observed 17.5%, 16.1%, 17.2% and 39.8%

improvement over LDAK alone (p=3.21×10^{-16}, 0.0001, 0.002 and 0.001, respectively.

Figure 5C). As a comparison, we integrated LDAK with PolyFun-pred(Weissbrod et
al., 2022) and SbayesRC(Zheng et al., 2022) using Polypred framework(Weissbrod et
al., 2022), but did not observe significant improvement over LDAK alone (difference
in R^2<0.01, p=0.001~0.07, Figure 5C). Since PolyFun-pred+BOLT-LMM has been
shown to significantly outperformed BOLT-LMM alone(Weissbrod et al., 2022), we
reasoned that the improvement of LDAK over BOLT-LMM might have attenuated the
improvement brought about by PolyFun-pred, making it difficult to reach significance
threshold. Taken together, we concluded that HFS could bring about mild but
significant improvement to classic SNP-based PRS in the task of cross-ancestry
polygenic prediction.

Discussion

In this study, we designed the new HFS framework for genetic association analysis
and demonstrated that it could improve classic SNP-based analysis in terms of causal
loci and gene identification, biological interpretation and polygenic prediction. We
suggest that HFS is a promising strategy for future genetic studies, but more
progresses in algorithm and computation and data resources is still desired.

Compared with SNP, HFS has several compelling features. For instance, LD
between adjacent HFS is much lower than SNP, which enhances the precision of
statistical fine-mapping. For those false-positive variants caused by LD, they are
expected to make little impacts on functional genomics, thus their HFS would be
close to reference and would not influence downstream analysis significantly. In line
with these advantages, we showed that HFS-based fine-mapping had high statistical
power, and downstream enrichment analysis was capable of revealing biologically
interpretable mechanisms. As a typical example, our findings of enrichment of

intelligence-associated loci in arachidonic acid metabolism pathway is in line with the
well-known role of polyunsaturated fatty acid in neurodevelopment(Helland et al.,
2003). Nonetheless, previous GWAS provided little evidence on this association.
Secondly, HFS could integrate effects of all variants within a locus, regardless of their
population frequency. Thus, HFS could capture information from rare variants
overlooked by classic association study and improve polygenic prediction, as shown
by our result. In fact, HFS framework could directly extend to whole-genomes
sequencing data and capture all mutations as rare as singleton, making one step
forward to fill in the "missing heritability".
Despite its potential, the current HFS framework carries several drawbacks and
necessitates significant enhancements. A key limitation is the substantial
computational cost. In this study, the transformation phase of the genotype-haplotype-
sequence for UK Biobank SNP data required hundreds of thousands of CPU core
hours. This computation cost would increase exponentially when analyzing whole-
genome sequencing (WGS) data or employing a sliding-window strategy. A potential
solution could involve developing a new algorithm that bypasses the variant calling
stage and directly generates DNA sequences per locus from raw sequencing or SNP
array data. For the sequence-to-HFS step, Sei(Chen et al., 2022) required about 1.8
GPU hours per one million sequences. Intriguingly, the majority of Sei's output is
unused in the HFS framework, since Sei predicts over twenty thousand functional
genomic features, while the HFS only represents one of their integrated scores. Future
development of novel deep learning models that predict functional genomics in a
manner more fitting to the HFS framework could considerably reduce computation
costs. Lastly, it's currently unfeasible to incorporate all genome-wide HFS into a
single LASSO model. This limitation forced us to first integrate HFS into pre-locus
score, which inevitably sacrificed the accuracy.
Another hurdle arises in integrating HFS with other genomic features.
Intrinsically, HFS captures only the variant effect mediated by functional genomics,
while a genetic variant might also influence amino acids, post-transcriptional
modifications (PTMs)(Park et al., 2021), and 3D chromosomal structures(Zhou, 2022).
Therefore, HFS alone cannot wholly replace SNP without any loss, as our results
demonstrate that the HFS-based prediction model captured approximately 70% of the
variance explainable by the SNP-based prediction model. One potential solution is to
extend the concept of HFS, applying deep learning to quantify the genetically
determined values of PTMs, protein biochemical properties(Pejaver et al., 2020), and
protein and chromosomal structures, potentially employing AlphaFold(Jumper et al.,
2021)-derived features(Liu et al., 2022). Analyzing HFS in conjunction with these
multi-modal function scores could provide a comprehensive depiction of the genetic
architecture of complex traits. However, the colossal computational cost is currently
prohibitive. As a compromise, we simply performed joint analysis of HFS with SNP
PRS in our prediction model analysis. This approach is far from optimal, as it led to
only moderate improvement and did not enhance fine-mapping and biological
enrichment analysis.
The challenge of using sequence-based deep learning (DL) models in HFS

applications is further compounded by their difficulty in predicting variations between individuals. Recent studies(Huang et al., 2023; Sasse et al., 2023) indicate that DL models, trained on the reference human genome, demonstrate limited accuracy in predicting gene expression levels across different individuals. This limitation is likely due to the models' inability to account for long-range regulatory patterns, which are crucial for understanding the impact of variants on gene expression and vary across genes. In contrast, our study leveraged sequence-determined functional genomic profiles in association studies, which mitigates this issue to an extent. For instance, although sei cannot identify the specific gene regulated by a given input sequence, it can predict changes in the sequence's functional activity. Future improvements in DL models' ability to predict interindividual differences could be achieved by incorporating cross-individual data in the training process. An example of such data is the EN-TEX(Rozowsky et al., 2023) dataset, which aligns functional genomic peaks with the specific individuals and haplotypes they correspond to.

In summary, our results demonstrate that incorporating HFS to represent genetically-determined functional genomic activities in genetic association studies offers robust improvements in both the biological interpretation and polygenic prediction of complex traits. Thus, the application of the HFS framework in future genetic association studies holds considerable promise.

Method

Sample description

This study analyzed UK Biobank data, with application ID 84436. We only included participants with array imputed genotype data in bgen format that passed UKB quality control, and removed related individuals. We randomly selected 350,587 self-identified British ancestry Caucasians as training sample. The remaining participants were grouped according to their ancestry, where Non-British European, South Asian, East Asian and African groups serve as test samples.

All phenotypes analyzed (Table S1) were collected from UKB table browser, which came from self-report or physical measurement. Phenotypes were first adjusted by age, sex, top ten principal components, Townsend index and genotype array quality metrics by linear regression. We then applied inverse-normal transformation on the residuals. Binary phenotypes were adjusted in the same way except by generalized linear regression.

Genotype data processing

We first segmented hg38 genome into 4,096 bp loci. To do so, we downloaded chromatin state annotation of 222 human tissues at different developmental stage (embryo, newborn, adult) from epimap(Boix et al., 2021) database. For each tissue, all chromosomal regions annotated as "transcription start site (TSS), transcription region (TX), enhancer, promoter" in at least half of the samples were marked as active regions. The union of active regions across all tissues was taken, and regions annotated as genomic gaps (centromere, ambiguous base pairs, etc.) in the Hg38 genome were removed. Then, for this series of active regions, if the length is less than 4096bp, the locus is defined as a 4096bp area centered around the active region. If the

length is greater than 4096bp, 4096bp length loci are gradually delineated from the midpoint outward. Finally, non-overlapping 4096bp blocks were used to cover the remaining genomic regions. This resulted in about 617,378 genomic regions in total. In the sliding-window analysis, all these blocks were shifted 2048 bp towards 5' end, generating another 617,378 blocks. We repeated the fine-mapping analysis and applied polygenic analysis on these combined blocks, using height as a representative trait.

For each of the loci, we obtained ID of variants within this locus by bedtools(Quinlan and Hall, 2010), then extracted genotypes from UKB .bgen file by bgenix, finally used Plink(Purcell et al., 2007) to remove all variants with INFO<0.8, Hardy-Weinberg p<10^{-6}, allele count<10 or missing rate >10%, and removed individual that missed more than 10% of retained variants in this locus. The output vcf file was liftover to hg38 by Crossmap(Zhao et al., 2014) and phased by SHAPEIT4(Delaneau et al., 2019). Phased vcf was transformed to .haps format by Plink, which in turn gave rise to two files: a vcf file containing information of each haplotype, and an n x 2 matrix in plain text that recorded the id of two haplotypes per individual.

HFS calculation

There has been several DL models that predict functional genomic profiles based on DNA sequence(Avsec et al., 2021; Chen et al., 2022; Kelley, 2020; Yan et al., 2021; Zhou et al., 2018). Among them, we chose sei(Chen et al., 2022) to calculate HFS for the following reasons: 1) the required input length (4,096bp) is moderate. 2) it represents 21,906 functional genomic tracks, more comprehensive than other models. 3) it integrated information of the entire sequence, not only the few bp at the center. For each haplotype at each locus, we generated its corresponding DNA sequence by bcftools(Danecek et al., 2021) consensus option. At each locus, the start point of each sequence was matched to the start point of reference sequence. When insertion variants made the sequence longer than 4,096bp, we discarded base pairs at the 3' end. Likewise, with deletion variants, we added N to the 3' end. We applied sei to predict 21,906 functional genomic tracks for each sequence, without normalizing for histone mark (divided each track score by the sum of histone mark score) as suggested by the sei author. We then used the projection matrix provided by sei to calculate forty sequence class scores, which could be regards as the weighted sum of these tracks and represented different aspect of functional genomic activities. We discarded the last score (heterochromatin 6 [centromere]), since its proportion is too low and is functionally trivial, leading to 39 scores per haplotype.

On each individual, we derived from each sequence class score the mean of two haplotypes, corresponding to additive model. For HFS Linkage Disequilibrium calculation, we extracted the mean value of sequence class score corresponding to reference sequence class of adjacent loci, and calculate R^2 value between them. The sequence class score of the reference sequence class was defined as the HFS for this locus, and was used for downstream trait association analysis.

HFS-trait association

For each locus, we calculated the association between trait-specific HFS and adjusted,

normalized trait value by linear regression, without any covariates (this is because all
selected covariates have been adjusted at the normalization step). For uniformity, we

set the significance threshold at $p<5\times10^{-8}$, even if it was over-stringent for n=590,959

loci. Among significant associations, we defined an independent association as the
locus with the lowest p value in the 200kb regions. As a positive control, we applied
quantitative and binary GWAS with REGENIE(Mbatchou et al., 2021), using default
settings and the same British training sample. The main difference is that we used raw
trait values in REGENIE, and provided the same covariates. We calculated the
genomic control inflation factor, $\lambda_{GC,}$ as the median of X^2 statistics, separately for
HFS association test and GWAS (only those SNPs in hapmap3(Altshuler et al., 2010)
project were calculated). We compared the λ_{GC} between HFS and SNP by Pearson
correlation analysis and paired t-test.
Fine-mapping analysis
We divided hg38 genome into 1,361 independent blocks as defined by MacDonald et
al.(MacDonald et al., 2022), and applied SUSIE to HFS of all loci within each block,
separately for each trait (parameters: maximum number of causal signal=10,
coverage=0.95). We subtracted reference HFS value for each locus prior to analysis,
such that homozygous reference haplotype corresponded to HFS=0. To avoid
influence of sei prediction noise, we rounded the HFS value at two decimals. This is
due to the fact that even if a variant actually makes no impact on functional genomics,
sei would still output a value that are close to but not equal to reference sequence
class score. Rounding procedure would set such HFS to zero and remove the random
value from sei. Loci whose HFS had PIP>0.95 were defined as causal loci, and loci
that had causal association with multiple traits were defined as pleiotropic loci. As a
positive control, we applied PolyFun(Weissbrod et al., 2020) and SbayesRC(Zheng et
al., 2022) on the GWAS summary statistics by REGENIE on the same training set,
and extracted the reported PIP to define causal SNP.
To analyze the functional characteristics of causal loci, we first defined the
sequence class of each locus by the maximum sequence class score of reference
haplotype. We then tested whether each sequence class contained excess causal loci of
each trait by Fisher test. For each causal locus, we also defined a "control" locus as
the nearest locus that matched the p-value of this causal locus, and tested whether
causal loci carried more PolyFun causal SNP than control loci by Fisher test.
Furthermore, For traits whose causal loci covered >0.1% of genome-wide SNP, we
applied LDSC(Finucane et al., 2015) to quantify the heritability enrichment in causal
and control loci, and compare their difference by jackknife method. To avoid winner's
curse, we used external GWAS summary statistics for this analysis(Mikaelsdottir et al.,
2021; Yengo et al., 2022). As an alternative method to quantify the heritability
captured by causal loci, we ran multivariate linear regression in independent British
test set where HFS of causal loci were independent variables and trait value were
dependent variable, and calculated the R^2 and AIC. We applied the same analysis on
causal SNP, and compared AIC between HFS and SNP multivariate regression.
Functional enrichment analysis

Similar to the idea of LDSC(Finucane et al., 2015), we first generated a series of
baseline annotation of each locus, then tested whether locus PIP was associated with
functional annotations after controlling the impact of these baseline annotations.
Specifically, we defined the following baseline annotations:
1) number of haplotypes, range of HFS distribution of all haplotypes (scaled by
reference HFS), and 39 sequence class score of reference haplotype.
2) genomic regions of conserved base, high Phastcons score(Siepel et al., 2005) in
mammals, primates and vertebrate, exon, intron, untranslated regions at 3' and 5' and
200bp flanking regions of TSS. We used bedtools *intersect -f 0.1* option to annotate
each locus by these annotations.
3) maximum B statistics(McVicker et al., 2009), minimum allele age and
$ASMC_{avg}$(Palamara et al., 2018) of all variants within this locus.
Type two and three annotations were directly obtained from LDSC(Finucane et
al., 2015) baseline annotations. We did not include annotations related to functional
genomics, since 39 sequence class score were used to capture functional genomic
characteristics. Conditioned on these baseline annotations, we analyzed the
enrichment of PIP in the following functional annotations:
1) Biological pathways: We downloaded all pathways from MsigDB(Subramanian et
al., 2005) C2: canonical pathways category (including Reactome(Fabregat et al.,
2018), PID(Schaefer et al., 2009), Biocarta and Wikipathway) and C6: Gene
ontology(Ashburner et al., 2000) (biological process) category. We retained only

pathways with >5 genes and <500 genes. We generated a gene × pathway binary

matrix and applied hierarchical clustering so that similar pathways were placed close
to each other. We sequentially compared adjacent pathways, and removed the smaller
one if the fraction of overlap > 30%. A total of 3,219 pathways were retained. We then
linked each locus to these pathways by cS2G(Gazal et al., 2022) strategy. Specifically,
a locus L would be annotated as 1 for pathway P only if L contained a SNP that was
link to P with cS2G score>0.5.
2) Tissue-specific chromatin activity: We downloaded chromHMM(Ernst and Kellis,
2012) chromatin state annotation for 833 samples from epimap(Boix et al., 2021), and
grouped them according to developmental stages and second-level tissue types. For
each group, all chromosomal regions annotated as "transcription start site (TSS),
transcription region (TX), enhancer, promoter" in at least half of the samples were
marked as active regions. We used bedtools *intersect -f 0.1* option to annotate whether
each locus was active in each tissue.
3) cell type-specific open chromatin regions: we downloaded scATAC-seq peak data
from Zhang et al.(K. Zhang et al., 2021), and annotated each locus by bedtools
intersect -f 0.1 option.
We applied multivariate linear regression of PIP against baseline annotations +
one of the functional annotations. Regression coefficient > 0 and Bonferroni-adjusted
regression p value<0.05 were used as significance threshold. From the final results,
we manually removed those pathways and cell types that reached significance
threshold in more than half of the traits, since these pathways likely reflected

unrecognized confounders.

Polygenic prediction

We used the posterior effect size estimated by SUSIE on sliding window strategy
(doubling the number of loci) as weights, and calculated the weighted sum of HFS as
the polygenic risk score of each trait, and calculated R^2 in independent British test
sample with simple linear regression. As a positive control, we applied LDAK-
BOLT(Q. Zhang et al., 2021) algorithm on the SNP array data (about seven hundred
thousand variants) with tenfold cross-validation and max iteration=200 in the same
training sample, and calculated SNP-based PRS with the output SNP weights.
Normalized trait values were analyzed, without any covariates provided. Array data
were filtered by Plink with option --geno 0.1 --hwe 1e-6 --mac 100 --maf 0.01 --mind
0.1.

To train the refined model that predict height, we first calculated per-block HFS-
based prediction score of height as the weighted sum of HFS within this block. Then,
within each target ancestry group (non-British European (NBE), South Asian (SAS),
East Asian (EA) and African (AFR) participants in UK Biobank), we randomly
selected half as tuning sample and half as test sample. In the tuning sample, we
applied LASSO regression that included both LDAK PRS and genome-wide per-
block HFS score (1,361 in total). The choice of LASSO regression was based on a
comparison on British European test set (Figure 5B), where LASSO, ridge and elastic
net gave similar results and LASSO was relatively better. In the tuning sample of
target ancestry, LASSO estimated the weights to combine per-block HFS score and
LDAK PRS. We calculated the final prediction score in the test sample using these
weights, and evaluated its prediction by linear regression R^2. Since the outcome
(height) has already been adjusted and standardized, no covariates were included in
this step. Additionally, we applied PolyFun-pred(Weissbrod et al., 2022) and
SbayesRC(Zheng et al., 2022) to the summary statistics of height (calculated by
REGENIE in the same training sample), and integrated their effect size with LDAK
weight in the tuning sample using Polypred(Weissbrod et al., 2022) method. PRS for
LDAK, LDAK+PolyFun and LDAK+SbayesRC were calculated by plink *score*
option, excluding variants with INFO<0.8, Hardy-Weinberg p<10^{-6}, allele count<2 or
missing rate >10% in the target test set.

Statistical analysis

All p-values were two-sided and adjusted by Bonferroni unless otherwise specified.
For group comparison, we used Fisher test for count data and paired t-test for
continuous data. For R^2 of PRS comparison, we applied r2redux(Momin et al., 2023)
R package to estimate 95% confidence interval and its p-value for the difference of R^2.

Availability of data and materials

635 This work was based on UK Biobank data from project 84436. Code for this analysis
636 is available at https://github.com/WeiCSong/HFS

Acknowledgement

This work was supported by grants from the 2030 Science and Technology Innovation

Key Program of Ministry of Science and Technology of China (No.
2022ZD020910001), the National Natural Science Foundation of China (No.
81971292, 82150610506) and the Natural Science Foundation of Shanghai (No.
21ZR1428600), the Medical-Engineering Cross Foundation of Shanghai Jiao Tong
University (No. YG2022ZD026).

Authors' contributions

645 W.S designed the study, collected and analyzed the data and wrote the manuscript.
646 G.N.L and Y.S supervised the study. All authors read, revise and approved the
647 manuscript.

Competing interests

649 The authors declare that they have no competing interests.

Reference

Aguet F, Barbeira AN, Bonazzola R, Brown A, Castel SE, Jo B, Kasela S, Kim-
Hellmuth S, Liang Y, Oliva M, Flynn ED, Parsana P, Fresard L, Gamazon ER,
Hamel AR, He Y, Hormozdiari F, Mohammadi P, Muñoz-Aguirre M, Park YS,
Saha A, Segrè A V., Strober BJ, Wen X, Wucher V, Ardlie KG, Battle A, Brown
CD, Cox N, Das S, Dermitzakis ET, Engelhardt BE, Garrido-Martín D, Gay NR,
Getz GA, Guigó R, Handsaker RE, Hoffman PJ, Im HK, Kashin S, Kwong A,
Lappalainen T, Li Xiao, MacArthur DG, Montgomery SB, Rouhana JM,
Stephens M, Stranger BE, Todres E, Viñuela A, Wang G, Zou Y, Anand S,
Gabriel S, Graubert A, Hadley K, Huang KH, Meier SR, Nedzel JL, Nguyen DT,
Balliu B, Conrad DF, Cotter DJ, deGoede OM, Einson J, Eskin E, Eulalio TY,
Ferraro NM, Gloudemans MJ, Hou L, Kellis M, Li Xin, Mangul S, Nachun DC,
Nobel AB, Park Y, Rao AS, Reverter F, Sabatti C, Skol AD, Teran NA, Wright
F, Ferreira PG, Li G, Melé M, Yeger-Lotem E, Barcus ME, Bradbury D, Krubit
T, McLean JA, Qi L, Robinson K, Roche N V., Smith AM, Sobin L, Tabor DE,
Undale A, Bridge J, Brigham LE, Foster BA, Gillard BM, Hasz R, Hunter M,
Johns C, Johnson M, Karasik E, Kopen G, Leinweber WF, McDonald A, Moser
MT, Myer K, Ramsey KD, Roe B, Shad S, Thomas JA, Walters G, Washington
M, Wheeler J, Jewell SD, Rohrer DC, Valley DR, Davis DA, Mash DC, Branton
PA, Sobin L, Barker LK, Gardiner HM, Mosavel M, Siminoff LA, Flicek P,
Haeussler M, Juettemann T, Kent WJ, Lee CM, Powell CC, Rosenbloom KR,
Ruffier M, Sheppard D, Taylor K, Trevanion SJ, Zerbino DR, Abell NS, Akey J,
Chen L, Demanelis K, Doherty JA, Feinberg AP, Hansen KD, Hickey PF, Hou L,
Jasmine F, Jiang L, Kaul R, Kellis M, Kibriya MG, Li JB, Li Q, Lin S, Linder
SE, Pierce BL, Rizzardi LF, Smith KS, Snyder M, Stamatoyannopoulos J, Tang
H, Wang M, Branton PA, Carithers LJ, Guan P, Koester SE, Little AR, Moore
HM, Nierras CR, Rao AK, Vaught JB, Volpi S. 2020. The GTEx Consortium
atlas of genetic regulatory effects across human tissues. *Science (80-)* **369**:1318–
1330. doi:10.1126/SCIENCE.AAZ1776
Altshuler DM, Gibbs RA, Peltonen L, Schaffner SF, Yu F, Dermitzakis E, Bonnen PE,

De Bakker PIW, Deloukas P, Gabriel SB, Gwilliam R, Hunt S, Inouye M, Jia X,
Aarno Palotie, Parkin M, Whittaker P, Chang K, Hawes A, Lewis LR, Ren Y,
Wheeler D, Muzny DM, Barnes C, Darvishi K, Hurles M, Korn JM, Kristiansson
K, Lee C, McCarroll SA, Nemesh J, Keinan A, Montgomery SB, Samuela
Pollack, Price AL, Soranzo N, Gonzaga-Jauregui C, Anttila V, Brodeur W, Daly
MJ, Leslie S, McVean G, Moutsianas L, Nguyen H, Zhang Q, Ghori MJR,
McGinnis R, McLaren W, Pollack S, Takeuchi F, Grossman SR, Shlyakhter I,
Hostetter EB, Sabeti PC, Adebamowo CA, Foster MW, Gordon DR, Licinio J,
Manca MC, Marshall PA, Matsuda I, Ngare D, Wang VO, Reddy D, Rotimi CN,
Royal CD, Sharp RR, Zeng C, Brooks DB, McEwen JE. 2010. Integrating
common and rare genetic variation in diverse human populations. *Nature*
467:52–58. doi:10.1038/nature09298
Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP,
Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis
A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G.
2000. Gene Ontology: tool for the unification of biology. *Nat Genet* **25**:25–29.
doi:10.1038/75556
Avsec Ž, Agarwal V, Visentin D, Ledsam JR, Grabska-Barwinska A, Taylor KR,
Assael Y, Jumper J, Kohli P, Kelley DR. 2021. Effective gene expression
prediction from sequence by integrating long-range interactions. *Nat Methods*
18:1196. doi:10.1038/S41592-021-01252-X
Baca SC, Singler C, Zacharia S, Seo JH, Morova T, Hach F, Ding Y, Schwarz T,
Huang CCF, Anderson J, Fay AP, Kalita C, Groha S, Pomerantz MM, Wang V,
Linder S, Sweeney CJ, Zwart W, Lack NA, Pasaniuc B, Takeda DY, Gusev A,
Freedman ML. 2022. Genetic determinants of chromatin reveal prostate cancer
risk mediated by context-dependent gene regulation. *Nat Genet* **54**:1364.
doi:10.1038/S41588-022-01168-Y
Barbeira AN, Dickinson SP, Bonazzola R, Zheng J, Wheeler HE, Torres JM,
Torstenson ES, Shah KP, Garcia T, Edwards TL, Stahl EA, Huckins LM, Aguet
F, Ardlie KG, Cummings BB, Gelfand ET, Getz G, Hadley K, Handsaker RE,
Huang KH, Kashin S, Karczewski KJ, Lek M, Li Xiao, MacArthur DG, Nedzel
JL, Nguyen DT, Noble MS, Segrè A V., Trowbridge CA, Tukiainen T, Abell NS,
Balliu B, Barshir R, Basha O, Battle A, Bogu GK, Brown A, Brown CD, Castel
SE, Chen LS, Chiang C, Conrad DF, Damani FN, Davis JR, Delaneau O,
Dermitzakis ET, Engelhardt BE, Eskin E, Ferreira PG, Frésard L, Gamazon ER,
Garrido-Martín D, Gewirtz ADH, Gliner G, Gloudemans MJ, Guigo R, Hall IM,
Han B, He Y, Hormozdiari F, Howald C, Jo B, Kang EY, Kim Y, Kim-Hellmuth
S, Lappalainen T, Li G, Li Xin, Liu B, Mangul S, McCarthy MI, McDowell IC,
Mohammadi P, Monlong J, Montgomery SB, Muñoz-Aguirre M, Ndungu AW,
Nobel AB, Oliva M, Ongen H, Palowitch JJ, Panousis N, Papasaikas P, Park YS,
Parsana P, Payne AJ, Peterson CB, Quan J, Reverter F, Sabatti C, Saha A,
Sammeth M, Scott AJ, Shabalin AA, Sodaei R, Stephens M, Stranger BE,
Strober BJ, Sul JH, Tsang EK, Urbut S, Van De Bunt M, Wang G, Wen X,
Wright FA, Xi HS, Yeger-Lotem E, Zappala Z, Zaugg JB, Zhou YH, Akey JM,

Bates D, Chan J, Claussnitzer M, Demanelis K, Diegel M, Doherty JA, Feinberg AP, Fernando MS, Halow J, Hansen KD, Haugen E, Hickey PF, Hou L, Jasmine F, Jian R, Jiang L, Johnson A, Kaul R, Kellis M, Kibriya MG, Lee K, Li JB, Li Q, Lin J, Lin S, Linder S, Linke C, Liu Y, Maurano MT, Molinie B, Nelson J, Neri FJ, Park Y, Pierce BL, Rinaldi NJ, Rizzardi LF, Sandstrom R, Skol A, Smith KS, Snyder MP, Stamatoyannopoulos J, Tang H, Wang L, Wang M, Van Wittenberghe N, Wu F, Zhang R, Nierras CR, Branton PA, Carithers LJ, Guan P, Moore HM, Rao A, Vaught JB, Gould SE, Lockart NC, Martin C, Struewing JP, Volpi S, Addington AM, Koester SE, Little AR, Brigham LE, Hasz R, Hunter M, Johns C, Johnson M, Kopen G, Leinweber WF, Lonsdale JT, McDonald A, Mestichelli B, Myer K, Roe B, Salvatore M, Shad S, Thomas JA, Walters G, Washington M, Wheeler J, Bridge J, Foster BA, Gillard BM, Karasik E, Kumar R, Miklos M, Moser MT, Jewell SD, Montroy RG, Rohrer DC, Valley DR, Davis DA, Mash DC, Undale AH, Smith AM, Tabor DE, Roche N V., McLean JA, Vatanian N, Robinson KL, Sobin L, Barcus ME, Valentino KM, Qi L, Hunter S, Hariharan P, Singh S, Um KS, Matose T, Tomaszewski MM, Barker LK, Mosavel M, Siminoff LA, Traino HM, Flicek P, Juettemann T, Ruffier M, Sheppard D, Taylor K, Trevanion SJ, Zerbino DR, Craft B, Goldman M, Haeussler M, Kent WJ, Lee CM, Paten B, Rosenbloom KR, Vivian J, Zhu J, Nicolae DL, Cox NJ, Im HK. 2018. Exploring the phenotypic consequences of tissue specific gene expression variation inferred from GWAS summary statistics. *Nat Commun* **9**:1–20. doi:10.1038/s41467-018-03621-1

Boix CA, James BT, Park YP, Meuleman W, Kellis M. 2021. Regulatory genomic circuitry of human disease loci by integrative epigenomics. *Nature* **590**:300–307. doi:10.1038/s41586-020-03145-z

Chen KM, Wong AK, Troyanskaya OG, Zhou J. 2022. A sequence-based global map of regulatory activity for deciphering human genetics. *Nat Genet* **54**:940–949. doi:10.1038/s41588-022-01102-2

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA, Davies RM. 2021. Twelve years of SAMtools and BCFtools. *Gigascience* **10**:1–4. doi:10.1093/GIGASCIENCE/GIAB008

Delaneau O, Zagury JF, Robinson MR, Marchini JL, Dermitzakis ET. 2019. Accurate, scalable and integrative haplotype estimation. *Nat Commun 2019 101* **10**:1–10. doi:10.1038/s41467-019-13225-y

Ernst J, Kellis M. 2012. ChromHMM: Automating chromatin-state discovery and characterization. *Nat Methods*. doi:10.1038/nmeth.1906

Fabregat A, Jupe S, Matthews L, Sidiropoulos K, Gillespie M, Garapati P, Haw R, Jassal B, Korninger F, May B, Milacic M, Roca CD, Rothfels K, Sevilla C, Shamovsky V, Shorser S, Varusai T, Viteri G, Weiser J, Wu G, Stein L, Hermjakob H, D'Eustachio P. 2018. The Reactome Pathway Knowledgebase. *Nucleic Acids Res* **46**:D649–D655. doi:10.1093/nar/gkx1132

Finucane HK, Bulik-Sullivan B, Gusev A, Trynka G, Reshef Y, Loh P-R, Anttila V, Xu H, Zang C, Farh K, Ripke S, Day FR, Purcell S, Stahl E, Lindstrom S, Perry JRB, Okada Y, Raychaudhuri S, Daly MJ, Patterson N, Neale BM, Price AL,

Patterson N, Neale BM, Price AL. 2015. Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nat Genet* **47**:1228–1235. doi:10.1038/ng.3404

Finucane HK, Reshef YA, Anttila V, Slowikowski K, Gusev A, Byrnes A, Gazal S, Loh P-RR, Lareau C, Shoresh N, Genovese G, Saunders A, Macosko E, Pollack S, Perry JRBB, Buenrostro JD, Bernstein BE, Raychaudhuri S, McCarroll S, Neale BM, Price AL. 2018. Heritability enrichment of specifically expressed genes identifies disease-relevant tissues and cell types. *Nat Genet* **50**:621–629. doi:10.1038/s41588-018-0081-4

Gazal S, Weissbrod O, Hormozdiari F, Dey KK, Nasser J, Jagadeesh KA, Weiner DJ, Shi H, Fulco CP, O'Connor LJ, Pasaniuc B, Engreitz JM, Price AL. 2022. Combining SNP-to-gene linking strategies to identify disease genes and assess disease omnigenicity. *Nat Genet* **54**:827–836. doi:10.1038/s41588-022-01087-y

Grotzinger AD, de la Fuente J, Davies G, Nivard MG, Tucker-Drob EM. 2022. Transcriptome-wide and stratified genomic structural equation modeling identify neurobiological pathways shared across diverse cognitive traits. *Nat Commun 2022 131* **13**:1–15. doi:10.1038/s41467-022-33724-9

Helland IB, Smith L, Saarem K, Saugstad OD, Drevon CA. 2003. Maternal Supplementation With Very-Long-Chain n-3 Fatty Acids During Pregnancy and Lactation Augments Children's IQ at 4 Years of Age. *Pediatrics* **111**:e39–e44. doi:10.1542/PEDS.111.1.E39

Hu X, Qiao D, Kim W, Moll M, Balte PP, Lange LA, Bartz TM, Kumar R, Li X, Yu B, Cade BE, Laurie CA, Sofer T, Ruczinski I, Nickerson DA, Muzny DM, Metcalf GA, Doddapaneni H, Gabriel S, Gupta N, Dugan-Perez S, Cupples LA, Loehr LR, Jain D, Rotter JI, Wilson JG, Psaty BM, Fornage M, Morrison AC, Vasan RS, Washko G, Rich SS, O'Connor GT, Bleecker E, Kaplan RC, Kalhan R, Redline S, Gharib SA, Meyers D, Ortega V, Dupuis J, London SJ, Lappalainen T, Oelsner EC, Silverman EK, Barr RG, Thornton TA, Wheeler HE, Cho MH, Im HK, Manichaikul A. 2022. Polygenic transcriptome risk scores for COPD and lung function improve cross-ethnic portability of prediction in the NHLBI TOPMed program. *Am J Hum Genet* **0**. doi:10.1016/J.AJHG.2022.03.007

Huang C, Shuai R, Baokar P, Chung R, Rastogi R, Kathail P, Ioannidis N. 2023. Personal transcriptome variation is poorly explained by current genomic deep learning models. *bioRxiv* 2023.06.30.547100. doi:10.1101/2023.06.30.547100

Iotchkova V, Ritchie GRS, Geihs M, Morganella S, Min JL, Walter K, Timpson NJ, Dunham I, Birney E, Soranzo N. 2019. GARFIELD classifies disease-relevant genomic features through integration of functional annotations with association signals. *Nat Genet* **51**:343–353. doi:10.1038/s41588-018-0322-6

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW,

Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein structure
prediction with AlphaFold. *Nat 2021 5967873* **596**:583–589.
doi:10.1038/s41586-021-03819-2
Kelley DR. 2020. Cross-species regulatory sequence activity prediction. *PLoS*
Comput Biol **16**. doi:10.1371/JOURNAL.PCBI.1008050
Kong X, Ma L, Chen E, Shaw CA, Edelstein LC. 2019. Identification of the
Regulatory Elements and Target Genes of Megakaryopoietic Transcription
Factor MEF2C. *Thromb Haemost* **119**:716. doi:10.1055/S-0039-1678694
Li Xihao, Li Z, Zhou H, Gaynor SM, Liu Yaowu, Chen H, Sun R, Dey R, Arnett DK,
Aslibekyan S, Ballantyne CM, Bielak LF, Blangero J, Boerwinkle E, Bowden
DW, Broome JG, Conomos MP, Correa A, Cupples LA, Curran JE, Freedman BI,
Guo X, Hindy G, Irvin MR, Kardia SLR, Kathiresan S, Khan AT, Kooperberg
CL, Laurie CC, Liu XS, Mahaney MC, Manichaikul AW, Martin LW, Mathias
RA, McGarvey ST, Mitchell BD, Montasser ME, Moore JE, Morrison AC,
O'Connell JR, Palmer ND, Pampana A, Peralta JM, Peyser PA, Psaty BM,
Redline S, Rice KM, Rich SS, Smith JA, Tiwari HK, Tsai MY, Vasan RS, Wang
FF, Weeks DE, Weng Z, Wilson JG, Yanek LR, Abe N, Abecasis GR, Aguet F,
Albert C, Almasy L, Alonso A, Ament S, Anderson P, Anugu P, Applebaum-
Bowden D, Ardlie K, Arking D, Arnett DK, Ashley-Koch A, Aslibekyan S,
Assimes T, Auer P, Avramopoulos D, Barnard J, Barnes K, Barr RG, Barron-
Casella E, Barwick L, Beaty T, Beck G, Becker D, Becker L, Beer R,
Beitelshees A, Benjamin E, Benos T, Bezerra M, Bielak LF, Bis J, Blackwell T,
Blangero J, Boerwinkle E, Bowden DW, Bowler R, Brody J, Broeckel U,
Broome JG, Bunting K, Burchard E, Bustamante C, Buth E, Cade B, Cardwell J,
Carey V, Carty C, Casaburi R, Casella J, Castaldi P, Chaffin M, Chang C, Chang
YC, Chasman D, Chavan S, Chen BJ, Chen WM, Chen YDI, Cho M, Choi SH,
Chuang LM, Chung M, Chung RH, Clish C, Comhair S, Conomos MP, Cornell
E, Correa A, Crandall C, Crapo J, Cupples LA, Curran JE, Curtis J, Custer B,
Damcott C, Darbar D, Das S, David S, Davis C, Daya M, de Andrade M, Fuentes
L de las, DeBaun M, Deka R, DeMeo D, Devine S, Duan Q, Duggirala R, Durda
JP, Dutcher S, Eaton C, Ekunwe L, El Boueiz A, Ellinor P, Emery L, Erzurum S,
Farber C, Fingerlin T, Flickinger M, Fornage M, Franceschini N, Frazar C, Fu M,
Fullerton SM, Fulton L, Gabriel S, Gan W, Gao S, Gao Y, Gass M, Gelb B,
Geng X (Priscilla), Geraci M, Germer S, Gerszten R, Ghosh A, Gibbs R,
Gignoux C, Gladwin M, Glahn D, Gogarten S, Gong DW, Goring H, Graw S,
Grine D, Gu CC, Guan Y, Guo X, Gupta N, Haessler J, Hall M, Harris D,
Hawley NL, He J, Heckbert S, Hernandez R, Herrington D, Hersh C, Hidalgo B,
Hixson J, Hobbs B, Hokanson J, Hong E, Hoth K, Hsiung C (Agnes), Hung YJ,
Huston H, Hwu CM, Irvin MR, Jackson R, Jain D, Jaquish C, Jhun MA, Johnsen
J, Johnson A, Johnson C, Johnston R, Jones K, Kang HM, Kaplan R, Kardia
SLR, Kathiresan S, Kelly S, Kenny E, Kessler M, Khan AT, Kim W, Kinney G,
Konkle B, Kooperberg CL, Kramer H, Lange C, Lange E, Lange L, Laurie CC,
Laurie C, LeBoff M, Lee J, Lee SS, Lee WJ, LeFaive J, Levine D, Levy D,
Lewis J, Li Xiaohui, Li Y, Lin Henry, Lin Honghuang, Lin KH, Lin X, Liu S,

Liu Yongmei, Liu Yu, Loos RJF, Lubitz S, Lunetta K, Luo J, Mahaney MC,
Make B, Manichaikul AW, Manson JA, Margolin L, Martin LW, Mathai S,
Mathias RA, May S, McArdle P, McDonald ML, McFarland S, McGarvey ST,
McGoldrick D, McHugh C, Mei H, Mestroni L, Meyers DA, Mikulla J, Min N,
Minear M, Minster RL, Mitchell BD, Moll M, Montasser ME, Montgomery C,
Moscati A, Musani S, Mwasongwe S, Mychaleckyj JC, Nadkarni G, Naik R,
Naseri T, Natarajan P, Nekhai S, Nelson SC, Neltner B, Nickerson D, North K,
O'Connell JR, O'Connor T, Ochs-Balcom H, Paik D, Palmer ND, Pankow J,
Papanicolaou G, Parsa A, Peralta JM, Perez M, Perry J, Peters U, Peyser PA,
Phillips LS, Pollin T, Post W, Becker JP, Boorgula MP, Preuss M, Psaty BM,
Qasba P, Qiao D, Qin Z, Rafaels N, Raffield L, Vasan RS, Rao DC, Rasmussen-
Torvik L, Ratan A, Redline S, Reed R, Regan E, Reiner A, Reupena MS, Rice
KM, Rich SS, Roden D, Roselli C, Rotter JI, Ruczinski I, Russell P, Ruuska S,
Ryan K, Sabino EC, Saleheen D, Salimi S, Salzberg S, Sandow K, Sankaran VG,
Scheller C, Schmidt E, Schwander K, Schwartz D, Sciurba F, Seidman C,
Seidman J, Sheehan V, Sherman SL, Shetty Amol, Shetty Aniket, Sheu WHH,
Shoemaker MB, Silver B, Silverman E, Smith JA, Smith J, Smith N, Smith T,
Smoller S, Snively B, Snyder M, Sofer T, Sotoodehnia N, Stilp AM, Storm G,
Streeten E, Su JL, Sung YJ, Sylvia J, Szpiro A, Sztalryd C, Taliun D, Tang H,
Taub M, Taylor KD, Taylor M, Taylor S, Telen M, Thornton TA, Threlkeld M,
Tinker L, Tirschwell D, Tishkoff S, Tiwari HK, Tong C, Tracy R, Tsai MY,
Vaidya D, Van Den Berg D, VandeHaar P, Vrieze S, Walker T, Wallace R,
Walts A, Wang FF, Wang H, Watson K, Weeks DE, Weir B, Weiss S, Weng LC,
Wessel J, Willer CJ, Williams K, Williams LK, Wilson C, Wilson JG, Wong Q,
Wu J, Xu H, Yanek LR, Yang I, Yang R, Zaghloul N, Zekavat M, Zhang Y,
Zhao SX, Zhao W, Zhi D, Zhou X, Zhu X, Zody M, Zoellner S, Abdalla M,
Abecasis GR, Arnett DK, Aslibekyan S, Assimes T, Atkinson E, Beitelshees A,
Bielak LF, Bis J, Bodea C, Boerwinkle E, Bowden DW, Brody J, Cade B,
Carlson J, Chang IS, Chen YDI, Chun S, Chung RH, Conomos MP, Correa A,
Cupples LA, Damcott C, de Vries P, Do R, Elliott A, Fu M, Ganna A, Gong DW,
Graham S, Haas M, Haring B, He J, Heckbert S, Himes B, Hixson J, Irvin MR,
Jain D, Jarvik G, Jhun MA, Jiang J, Jun G, Kalyani R, Kardia SLR, Kathiresan S,
Khera A, Klarin D, Kooperberg CL, Kral B, Lange L, Laurie CC, Laurie C,
Lemaitre R, Li Z, Li Xihao, Lin X, Mahaney MC, Manichaikul AW, Martin LW,
Mathias RA, Mathur R, McGarvey ST, McHugh C, McLenithan J, Mikulla J,
Mitchell BD, Montasser ME, Moran A, Morrison AC, Nakao T, Natarajan P,
Nickerson D, North K, O'Connell JR, O'Donnell C, Palmer ND, Pampana A,
Patel A, Peloso GM, Perry J, Peters U, Peyser PA, Pirruccello J, Pollin T, Psaty
BM, Redline S, Reed R, Reiner A, Rich SS, Rosenthal S, Rotter JI, Schoenberg J,
Selvaraj MS, Sheu WHH, Smith JA, Sofer T, Stilp AM, Sunyaev SR, Surakka I,
Sztalryd C, Tang H, Taylor KD, Tsai MY, Uddin MM, Urbut S, Verbanck M,
Von Holle A, Wang H, Wang FF, Wiggins K, Willer CJ, Wilson JG, Wolford B,
Xu H, Yanek LR, Zaghloul N, Zekavat M, Zhang J, Neale BM, Sunyaev SR,
Abecasis GR, Rotter JI, Willer CJ, Peloso GM, Natarajan P, Lin X. 2020.

Dynamic incorporation of multiple in silico functional annotations empowers rare variant association analysis of large whole-genome sequencing studies at scale. *Nat Genet* **52**:969–983. doi:10.1038/s41588-020-0676-4

Liang Y, Pividori M, Manichaikul A, Palmer AA, Cox NJ, Wheeler HE, Im HK. 2022. Polygenic transcriptome risk scores (PTRS) can improve portability of polygenic risk scores across ancestries. *Genome Biol* **23**:1–18. doi:10.1186/S13059-021-02591-W/FIGURES/5

Liu Z, Pan W, Li W, Zhen X, Liang J, Cai W, Xu F, Yuan K, Lin GN. 2022. Evaluation of the Effectiveness of Derived Features of AlphaFold2 on Single-Sequence Protein Binding Site Prediction. *Biology (Basel)* **11**:1454. doi:10.3390/BIOLOGY11101454/S1

MacDonald JW, Harrison T, Bammler TK, Mancuso N, Lindström S. 2022. An updated map of GRCh38 linkage disequilibrium blocks based on European ancestry data. *bioRxiv* 2022.03.04.483057. doi:10.1101/2022.03.04.483057

Mbatchou J, Barnard L, Backman J, Marcketta A, Kosmicki JA, Ziyatdinov A, Benner C, O'Dushlaine C, Barber M, Boutkov B, Habegger L, Ferreira M, Baras A, Reid J, Abecasis G, Maxwell E, Marchini J. 2021. Computationally efficient whole-genome regression for quantitative and binary traits. *Nat Genet* **53**:1097–1103. doi:10.1038/s41588-021-00870-7

McVicker G, Gordon D, Davis C, Green P. 2009. Widespread genomic signatures of natural selection in hominid evolution. *PLoS Genet* **5**. doi:10.1371/journal.pgen.1000471

Mikaelsdottir E, Thorleifsson G, Stefansdottir L, Halldorsson G, Sigurdsson JK, Lund SH, Tragante V, Melsted P, Rognvaldsson S, Norland K, Helgadottir A, Magnusson MK, Ragnarsson GB, Kristinsson SY, Reykdal S, Vidarsson B, Gudmundsdottir IJ, Olafsson I, Onundarson PT, Sigurdardottir O, Sigurdsson EL, Grondal G, Geirsson AJ, Geirsson G, Gudmundsson J, Holm H, Saevarsdottir S, Jonsdottir I, Thorgeirsson G, Gudbjartsson DF, Thorsteinsdottir U, Rafnar T, Stefansson K. 2021. Genetic variants associated with platelet count are predictive of human disease and physiological markers. *Commun Biol* **4**. doi:10.1038/S42003-021-02642-9

Momin MM, Lee S, Wray NR, Lee SH. 2023. Significance tests for R2 of out-of-sample prediction using polygenic scores. *Am J Hum Genet* **0**. doi:10.1016/J.AJHG.2023.01.004

Morales MAG, Montero⎕vargas JM, Vizuet⎕de⎕rueda JC, Teran LM. 2021. New Insights into the Role of PD-1 and Its Ligands in Allergic Disease. *Int J Mol Sci* **22**:11898. doi:10.3390/IJMS222111898

Nawijn MC, Piavaux BJA, Jeurink P V., Gras R, Reinders MA, Stearns T, Foote S, Hylkema MN, Groot PC, Korstanje R, Van Oosterhout AJM. 2011. Identification of the Mhc Region as an Asthma Susceptibility Locus in Recombinant Congenic Mice. *Am J Respir Cell Mol Biol* **45**:295. doi:10.1165/RCMB.2009-0369OC

Nowbandegani PS, Wohns AW, Ballard JL, Lander ES, Bloemendal A, Neale BM, O'Connor LJ. 2022. Extremely sparse models of linkage disequilibrium in

ancestrally diverse association studies. *bioRxiv* 2022.09.06.506858.
doi:10.1101/2022.09.06.506858
Null M, Dupuis J, Sheinidashtegol P, Layer RM, Gignoux CR, Hendricks AE. 2022.
RAREsim: A simulation method for very rare genetic variants. *Am J Hum Genet*
0. doi:10.1016/J.AJHG.2022.02.009
Palamara PF, Terhorst J, Song YS, Price AL. 2018. High-throughput inference of
pairwise coalescence times identifies signals of selection and enriched disease
heritability. *Nat Genet* **50**:1311–1317. doi:10.1038/s41588-018-0177-x
Park CY, Zhou J, Wong AK, Chen KM, Theesfeld CL, Darnell RB, Troyanskaya OG.
2021. Genome-wide landscape of RNA-binding protein target site dysregulation
reveals a major impact on psychiatric disorder risk. *Nat Genet* **53**:166.
doi:10.1038/S41588-020-00761-3
Pejaver V, Urresti J, Lugo-Martinez J, Pagel KA, Lin GN, Nam HJ, Mort M, Cooper
DN, Sebat J, Iakoucheva LM, Mooney SD, Radivojac P. 2020. Inferring the
molecular and phenotypic impact of amino acid variants with MutPred2. *Nat*
Commun **11**:1–13. doi:10.1038/s41467-020-19669-x
Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J,
Sklar P, De Bakker PIW, Daly MJ, Sham PC. 2007. PLINK: A tool set for
whole-genome association and population-based linkage analyses. *Am J Hum*
Genet **81**:559–575. doi:10.1086/519795
Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**:841–842.
doi:10.1093/BIOINFORMATICS/BTQ033
Rozowsky J, Gao J, Borsari B, Yang YT, Galeev T, Gürsoy G, Epstein CB, Xiong K,
Xu J, Li T, Liu J, Yu K, Berthel A, Chen Z, Navarro F, Sun MS, Wright J,
Chang J, Cameron CJF, Shoresh N, Gaskell E, Drenkow J, Adrian J, Aganezov S,
Aguet F, Balderrama-Gutierrez G, Banskota S, Corona GB, Chee S, Chhetri SB,
Cortez Martins GC, Danyko C, Davis CA, Farid D, Farrell NP, Gabdank I, Gofin
Y, Gorkin DU, Gu M, Hecht V, Hitz BC, Issner R, Jiang Y, Kirsche M, Kong X,
Lam BR, Li S, Li B, Li X, Lin KZ, Luo R, Mackiewicz M, Meng R, Moore JE,
Mudge J, Nelson N, Nusbaum C, Popov I, Pratt HE, Qiu Y, Ramakrishnan S,
Raymond J, Salichos L, Scavelli A, Schreiber JM, Sedlazeck FJ, See LH,
Sherman RM, Shi X, Shi M, Sloan CA, Strattan JS, Tan Z, Tanaka FY, Vlasova
A, Wang J, Werner J, Williams B, Xu M, Yan C, Yu L, Zaleski C, Zhang J,
Ardlie K, Cherry JM, Mendenhall EM, Noble WS, Weng Z, Levine ME, Dobin
A, Wold B, Mortazavi A, Ren B, Gillis J, Myers RM, Snyder MP, Choudhary J,
Milosavljevic A, Schatz MC, Bernstein BE, Guigó R, Gingeras TR, Gerstein M.
2023. The EN-TEx resource of multi-tissue personal epigenomes & variant-
impact models. *Cell* **186**:1493-1511.e40. doi:10.1016/J.CELL.2023.02.018
Sasse A, Ng B, Spiro AE, Tasaki S, Bennett DA, Gaiteri C, De Jager PL, Chikina M,
Mostafavi S, Allen PG. 2023. Benchmarking of deep neural networks for
predicting personal gene expression from DNA sequence highlights
shortcomings. *bioRxiv* 2023.03.16.532969. doi:10.1101/2023.03.16.532969
Schaefer CF, Anthony K, Krupa S, Buchoff J, Day M, Hannay T, Buetow KH. 2009.

PID: the Pathway Interaction Database. *Nucleic Acids Res* **37**:D674.
doi:10.1093/NAR/GKN653

Sekar A, Bialas AR, Rivera H de, Davis A, Hammond TR, Kamitaki N, Tooley K,
Presumey J, Baum M, Doren V Van, Genovese G, Rose SA, Handsaker RE,
Consortium SWG of the PG, Daly MJ, Carroll MC, Stevens B, McCarroll SA.
2016. Schizophrenia risk from complex variation of complement component 4.
Nature **530**:177. doi:10.1038/NATURE16549

Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K, Clawson H,
Spieth J, Hillier LDW, Richards S, Weinstock GM, Wilson RK, Gibbs RA, Kent
WJ, Miller W, Haussler D. 2005. Evolutionarily conserved elements in
vertebrate, insect, worm, and yeast genomes. *Genome Res* **15**:1034–1050.
doi:10.1101/GR.3715005

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA,
Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. 2005. Gene set
enrichment analysis: A knowledge-based approach for interpreting genome-wide
expression profiles. *Proc Natl Acad Sci U S A* **102**:15545–15550.
doi:10.1073/PNAS.0506580102/SUPPL_FILE/06580FIG7.JPG

Visscher PM, Wray NR, Zhang Q, Sklar P, McCarthy MI, Brown MA, Yang J. 2017.
10 Years of GWAS Discovery: Biology, Function, and Translation. *Am J Hum*
Genet **101**:5–22. doi:10.1016/J.AJHG.2017.06.005

Wang G, Sarkar A, Carbonetto P, Stephens M. 2020. A simple new approach to
variable selection in regression, with application to genetic fine mapping. *J R*
Stat Soc Ser B (Statistical Methodol **82**:1273–1300. doi:10.1111/RSSB.12388

Watanabe K, Stringer S, Frei O, Umićević Mirkov M, de Leeuw C, Polderman TJC,
van der Sluis S, Andreassen OA, Neale BM, Posthuma D. 2019. A global
overview of pleiotropy and genetic architecture in complex traits. *Nat Genet*
51:1339–1348. doi:10.1038/s41588-019-0481-0

Weissbrod O, Hormozdiari F, Benner C, Cui R, Ulirsch J, Gazal S, Schoech AP, van
de Geijn B, Reshef Y, Márquez-Luna C, O'Connor L, Pirinen M, Finucane HK,
Price AL. 2020. Functionally informed fine-mapping and polygenic localization
of complex trait heritability. *Nat Genet* **52**:1355–1363. doi:10.1038/s41588-020-
00735-5

Weissbrod O, Kanai M, Shi H, Gazal S, Peyrot WJ, Khera A V., Okada Y, Matsuda K,
Yamanashi Y, Furukawa Y, Morisaki T, Murakami Y, Kamatani Y, Muto K,
Nagai A, Obara W, Yamaji K, Takahashi K, Asai S, Takahashi Y, Suzuki T,
Sinozaki N, Yamaguchi H, Minami S, Murayama S, Yoshimori K, Nagayama S,
Obata D, Higashiyama M, Masumoto A, Koretsune Y, Martin AR, Finucane HK,
Price AL. 2022. Leveraging fine-mapping and multipopulation training data to
improve cross-population polygenic risk scores. *Nat Genet* **54**:450–458.
doi:10.1038/s41588-022-01036-9

Yan J, Qiu Y, Ribeiro dos Santos AM, Yin Y, Li YE, Vinckier N, Nariai N, Benaglio
P, Raman A, Li X, Fan S, Chiou J, Chen F, Frazer KA, Gaulton KJ, Sander M,
Taipale J, Ren B. 2021. Systematic analysis of binding of transcription factors to
noncoding variants. *Nature* **591**:147–151. doi:10.1038/s41586-021-03211-0

Yang J, Bakshi A, Zhu Z, Hemani G, Vinkhuyzen AAE, Lee SH, Robinson MR,
1033   Perry JRB, Nolte IM, Van Vliet-Ostaptchouk J V., Snieder H, Esko T, Milani L,
1034   Mägi R, Metspalu A, Hamsten A, Magnusson PKE, Pedersen NL, Ingelsson E,
1035   Soranzo N, Keller MC, Wray NR, Goddard ME, Visscher PM. 2015. Genetic
1036   variance estimation with imputed variants finds negligible missing heritability
1037   for human height and body mass index. *Nat Genet* **47**:1114–1120.
1038   doi:10.1038/ng.3390

1039   Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, Madden PA,
1040   Heath AC, Martin NG, Montgomery GW, Goddard ME, Visscher PM. 2010.
1041   Common SNPs explain a large proportion of the heritability for human height.
1042   *Nat Genet* **42**:565–569. doi:10.1038/ng.608

1043   Yengo L, Vedantam S, Marouli E, Sidorenko J, Bartell E, Sakaue S, Graff M, Eliasen
1044   AU, Jiang Y, Raghavan S, Miao J, Arias JD, Graham SE, Mukamel RE,
1045   Spracklen CN, Yin X, Chen SH, Ferreira T, Highland HH, Ji Y, Karaderi T, Lin
1046   K, Lüll K, Malden DE, Medina-Gomez C, Machado M, Moore A, Rüeger S, Sim
1047   X, Vrieze S, Ahluwalia TS, Akiyama M, Allison MA, Alvarez M, Andersen MK,
1048   Ani A, Appadurai V, Arbeeva L, Bhaskar S, Bielak LF, Bollepalli S,
1049   Bonnycastle LL, Bork-Jensen J, Bradfield JP, Bradford Y, Braund PS, Brody JA,
1050   Burgdorf KS, Cade BE, Cai H, Cai Q, Campbell A, Cañadas-Garre M, Catamo E,
1051   Chai JF, Chai X, Chang LC, Chang YC, Chen CH, Chesi A, Choi SH, Chung
1052   RH, Cocca M, Concas MP, Couture C, Cuellar-Partida G, Danning R, Daw EW,
1053   Degenhard F, Delgado GE, Delitala A, Demirkan A, Deng X, Devineni P, Dietl
1054   A, Dimitriou M, Dimitrov L, Dorajoo R, Ekici AB, Engmann JE, Fairhurst-
1055   Hunter Z, Farmaki AE, Faul JD, Fernandez-Lopez JC, Forer L, Francescatto M,
1056   Freitag-Wolf S, Fuchsberger C, Galesloot TE, Gao Y, Gao Z, Geller F,
1057   Giannakopoulou O, Giulianini F, Gjesing AP, Goel A, Gordon SD, Gorski M,
1058   Grove J, Guo X, Gustafsson S, Haessler J, Hansen TF, Havulinna AS, Haworth
1059   SJ, He J, Heard-Costa N, Hebbar P, Hindy G, Ho YLA, Hofer E, Holliday E,
1060   Horn K, Hornsby WE, Hottenga JJ, Huang H, Huang J, Huerta-Chagoya A,
1061   Huffman JE, Hung YJ, Huo S, Hwang MY, Iha H, Ikeda DD, Isono M, Jackson
1062   AU, Jäger S, Jansen IE, Johansson I, Jonas JB, Jonsson A, Jørgensen T, Kalafati
1063   IP, Kanai M, Kanoni S, Kårhus LL, Kasturiratne A, Katsuya T, Kawaguchi T,
1064   Kember RL, Kentistou KA, Kim HN, Kim YJ, Kleber ME, Knol MJ, Kurbasic A,
1065   Lauzon M, Le P, Lea R, Lee JY, Leonard HL, Li SA, Li Xiaohui, Li Xiaoyin,
1066   Liang J, Lin H, Lin SY, Liu Jun, Liu X, Lo KS, Long J, Lores-Motta L, Luan J,
1067   Lyssenko V, Lyytikäinen LP, Mahajan A, Mamakou V, Mangino M,
1068   Manichaikul A, Marten J, Mattheisen M, Mavarani L, McDaid AF, Meidtner K,
1069   Melendez TL, Mercader JM, Milaneschi Y, Miller JE, Millwood IY, Mishra PP,
1070   Mitchell RE, Møllehave LT, Morgan A, Mucha S, Munz M, Nakatochi M,
1071   Nelson CP, Nethander M, Nho CW, Nielsen AA, Nolte IM, Nongmaithem SS,
1072   Noordam R, Ntalla I, Nutile T, Pandit A, Christofidou P, Pärna K, Pauper M,
1073   Petersen ERB, Petersen L V., Pitkänen N, Polašek O, Poveda A, Preuss MH,
1074   Pyarajan S, Raffield LM, Rakugi H, Ramirez J, Rasheed A, Raven D, Rayner
1075   NW, Riveros C, Rohde R, Ruggiero D, Ruotsalainen SE, Ryan KA, Sabater-

Lleal M, Saxena R, Scholz M, Sendamarai A, Shen B, Shi J, Shin JH, Sidore C,
Sitlani CM, Slieker RC, Smit RAJ, Smith A V., Smith JA, Smyth LJ, Southam L,
Steinthorsdottir V, Sun L, Takeuchi F, Tallapragada DSP, Taylor KD, Tayo BO,
Tcheandjieu C, Terzikhan N, Tesolin P, Teumer A, Theusch E, Thompson DJ,
Thorleifsson G, Timmers PRHJ, Trompet S, Turman C, Vaccargiu S, van der
Laan SW, van der Most PJ, van Klinken JB, van Setten J, Verma SS, Verweij N,
Veturi Y, Wang CA, Wang C, Wang L, Wang Z, Warren HR, Bin Wei W,
Wickremasinghe AR, Wielscher M, Wiggins KL, Winsvold BS, Wong A, Wu Y,
Wuttke M, Xia R, Xie T, Yamamoto K, Yang Jingyun, Yao J, Young H, Yousri
NA, Yu L, Zeng L, Zhang W, Zhang X, Zhao JH, Zhao W, Zhou W,
Zimmermann ME, Zoledziewska M, Adair LS, Adams HHH, Aguilar-Salinas
CA, Al-Mulla F, Arnett DK, Asselbergs FW, Åsvold BO, Attia J, Banas B,
Bandinelli S, Bennett DA, Bergler T, Bharadwaj D, Biino G, Bisgaard H,
Boerwinkle E, Böger CA, Bønnelykke K, Boomsma DI, Børglum AD, Borja JB,
Bouchard C, Bowden DW, Brandslund I, Brumpton B, Buring JE, Caulfield MJ,
Chambers JC, Chandak GR, Chanock SJ, Chaturvedi N, Chen YDI, Chen Z,
Cheng CY, Christophersen IE, Ciullo M, Cole JW, Collins FS, Cooper RS, Cruz
M, Cucca F, Cupples LA, Cutler MJ, Damrauer SM, Dantoft TM, de Borst GJ,
de Groot LCPGM, De Jager PL, de Kleijn DPV, Janaka de Silva H, Dedoussis G
V., den Hollander AI, Du S, Easton DF, Elders PJM, Eliassen AH, Ellinor PT,
Elmståhl S, Erdmann J, Evans MK, Fatkin D, Feenstra B, Feitosa MF, Ferrucci L,
Ford I, Fornage M, Franke A, Franks PW, Freedman BI, Gasparini P, Gieger C,
Girotto G, Goddard ME, Golightly YM, Gonzalez-Villalpando C, Gordon-
Larsen P, Grallert H, Grant SFA, Grarup N, Griffiths L, Gudnason V, Haiman C,
Hakonarson H, Hansen T, Hartman CA, Hattersley AT, Hayward C, Heckbert
SR, Heng CK, Hengstenberg C, Hewitt AW, Hishigaki H, Hoyng CB, Huang PL,
Huang W, Hunt SC, Hveem K, Hyppönen E, Iacono WG, Ichihara S, Ikram MA,
Isasi CR, Jackson RD, Jarvelin MR, Jin ZB, Jöckel KH, Joshi PK, Jousilahti P,
Jukema JW, Kähönen M, Kamatani Y, Kang KD, Kaprio J, Kardia SLR, Karpe F,
Kato N, Kee F, Kessler T, Khera A V., Khor CC, Kiemeney LALM, Kim BJ,
Kim EK, Kim HL, Kirchhof P, Kivimaki M, Koh WP, Koistinen HA, Kolovou
GD, Kooner JS, Kooperberg C, Köttgen A, Kovacs P, Kraaijeveld A, Kraft P,
Krauss RM, Kumari M, Kutalik Z, Laakso M, Lange LA, Langenberg C, Launer
LJ, Le Marchand L, Lee H, Lee NR, Lehtimäki T, Li H, Li L, Lieb W, Lin X,
Lind L, Linneberg A, Liu CT, Liu Jianjun, Loeffler M, London B, Lubitz SA,
Lye SJ, Mackey DA, Mägi R, Magnusson PKE, Marcus GM, Vidal PM, Martin
NG, März W, Matsuda F, McGarrah RW, McGue M, McKnight AJ, Medland SE,
Mellström D, Metspalu A, Mitchell BD, Mitchell P, Mook-Kanamori DO,
Morris AD, Mucci LA, Munroe PB, Nalls MA, Nazarian S, Nelson AE, Neville
MJ, Newton-Cheh C, Nielsen CS, Nöthen MM, Ohlsson C, Oldehinkel AJ,
Orozco L, Pahkala K, Pajukanta P, Palmer CNA, Parra EJ, Pattaro C, Pedersen O,
Pennell CE, Penninx BWJH, Perusse L, Peters A, Peyser PA, Porteous DJ,
Posthuma D, Power C, Pramstaller PP, Province MA, Qi Q, Qu J, Rader DJ,
Raitakari OT, Ralhan S, Rallidis LS, Rao DC, Redline S, Reilly DF, Reiner AP,

Rhee SY, Ridker PM, Rienstra M, Ripatti S, Ritchie MD, Roden DM, Rosendaal
FR, Rotter JI, Rudan I, Rutters F, Sabanayagam C, Saleheen D, Salomaa V,
Samani NJ, Sanghera DK, Sattar N, Schmidt B, Schmidt H, Schmidt R, Schulze
MB, Schunkert H, Scott LJ, Scott RJ, Sever P, Shiroma EJ, Shoemaker MB, Shu
XO, Simonsick EM, Sims M, Singh JR, Singleton AB, Sinner MF, Smith JG,
Snieder H, Spector TD, Stampfer MJ, Stark KJ, Strachan DP, 't Hart LM, Tabara
Y, Tang H, Tardif JC, Thanaraj TA, Timpson NJ, Tönjes A, Tremblay A, Tuomi
T, Tuomilehto J, Tusié-Luna MT, Uitterlinden AG, van Dam RM, van der Harst
P, Van der Velde N, van Duijn CM, van Schoor NM, Vitart V, Völker U,
Vollenweider P, Völzke H, Wacher-Rodarte NH, Walker M, Wang YX,
Wareham NJ, Watanabe RM, Watkins H, Weir DR, Werge TM, Widen E,
Wilkens LR, Willemsen G, Willett WC, Wilson JF, Wong TY, Woo JT, Wright
AF, Wu JY, Xu H, Yajnik CS, Yokota M, Yuan JM, Zeggini E, Zemel BS,
Zheng W, Zhu X, Zmuda JM, Zonderman AB, Zwart JA, Partida GC, Sun Y,
Croteau-Chonka D, Vonk JM, Chanock S, Le Marchand L, Chasman DI, Cho
YS, Heid IM, McCarthy MI, Ng MCY, O'Donnell CJ, Rivadeneira F,
Thorsteinsdottir U, Sun Y V., Tai ES, Boehnke M, Deloukas P, Justice AE,
Lindgren CM, Loos RJF, Mohlke KL, North KE, Stefansson K, Walters RG,
Winkler TW, Young KL, Loh PR, Yang Jian, Esko T, Assimes TL, Auton A,
Abecasis GR, Willer CJ, Locke AE, Berndt SI, Lettre G, Frayling TM, Okada Y,
Wood AR, Visscher PM, Hirschhorn JN. 2022. A saturated map of common
genetic variants associated with human height. *Nature* **610**:704–712.
doi:10.1038/s41586-022-05275-y
Zeng J, Xue A, Jiang L, Lloyd-Jones LR, Wu Y, Wang H, Zheng Z, Yengo L,
Kemper KE, Goddard ME, Wray NR, Visscher PM, Yang J. 2021. Widespread
signatures of natural selection across human complex traits and functional
genomic categories. *Nat Commun* **12**:752527. doi:10.1038/s41467-021-21446-3
Zhang K, Hocker JD, Miller M, Wang A, Preissl S, Correspondence BR, Hou X,
Chiou J, Poirion OB, Qiu Y, Li YE, Gaulton KJ, Ren B. 2021. A single-cell atlas
of chromatin accessibility in the human genome. *Cell* **184**:1–17.
doi:10.1016/j.cell.2021.10.024
Zhang Q, Privé F, Vilhjálmsson B, Speed D. 2021. Improved genetic prediction of
complex traits from individual-level data or summary statistics. *Nat Commun*
12:1–9. doi:10.1038/s41467-021-24485-y
Zhao H, Sun Z, Wang J, Huang H, Kocher JP, Wang L. 2014. CrossMap: a versatile
tool for coordinate conversion between genome assemblies. *Bioinformatics*
30:1006–1007. doi:10.1093/BIOINFORMATICS/BTT730
Zheng Z, Liu S, Sidorenko J, Yengo L, Turley P, Ani A, Wang R, Nolte IM, Snieder
H, Study LC, Yang J, Wray NR, Goddard ME, Visscher PM, Zeng J. 2022.
Leveraging functional genomic annotations and genome coverage to improve
polygenic prediction of complex traits within and between ancestries. *bioRxiv*
2022.10.12.510418. doi:10.1101/2022.10.12.510418
Zhou J. 2022. Sequence-based modeling of three-dimensional genome architecture
from kilobase to chromosome scale. *Nat Genet* **54**:725–734.

doi:10.1038/s41588-022-01065-4

Zhou J, Theesfeld CL, Yao K, Chen KM, Wong AK, Troyanskaya OG. 2018. Deep

learning sequence-based ab initio prediction of variant effects on expression and

disease risk. *Nat Genet* **50**:1171–1179. doi:10.1038/s41588-018-0160-6

Zhou W, Bi W, Zhao Z, Dey KK, Jagadeesh KA, Karczewski KJ, Daly MJ, Neale

BM, Lee S. 2022. SAIGE-GENE+ improves the efficiency and accuracy of set-

based rare variant association tests. *Nat Genet* **54**:1466–1469.

doi:10.1038/s41588-022-01178-w

1172

1173

Figure 1 Flowchart of the study. Ind: individual.

Figure 2 Fine-mapping result summary. Grey bar plots indicated the number of loci
with posterior inclusion probability (PIP)>0.95 in HFS+SUSIE (causal loci). Black
bar plots indicated number of SNP with PIP>0.95 in PolyFun or SbayesRC analysis
(the larger number was shown). Each grid of heatmap showed the Odds ratio of each
sequence class loci being causal loci for each trait. "All_OR" indicated Odds ratio for
pooling all traits together. Enh: enhancer. TF: transcription factor binding site.

Figure 3 Biological enrichment analysis based on HFS fine-mapping. X-axis
indicated t statistics of the analyzed term in a multivariate linear regression (Method).
Cell: single-cell ATAC peak for 222 cell types from Zhang et al(K. Zhang et al., 2021).
Tissue: active chromatin regions of 222 tissues from epimap(Boix et al., 2021). For
each trait, we showed the most significant term plus one or two terms with high
biological interpretation that also passed significance threshold. Full enrichment result
was shown in Table S7 and S8.

Figure 4 HFS linked trait to causal genes. A: Target genes of causal loci identified
by HFS+SUSIE for platelet count. Only genes that showed functional convergence
were shown. B: Regional plot for RBBP5. HFS: loci PIP calculated by HFS+SUSIE.
SNP: SNP PIP calculated by PolyFun. cCRE: credible cis-regulation elements. C:
Regional plot of MHC region for asthma. Thickened curve linked highlighted causal
loci to its target genes predicted by cS2G(Gazal et al., 2022).

Figure 5 HFS-based polygenic prediction. A: Prediction R^2 of HFS-based polygenic
risk score (PRS) using different threshold of PIP. allSNP: SNP-based PRS calculated
by LDAK-BOLT(Q. Zhang et al., 2021). n: number of features included in the
corresponding PRS. B: Prediction R^2 of per-block HFS score in British European test
set by different methods. EN: elastic net. C: Prediction R^2 of different tools in non-
British European (NBE), South Asian (SAS), East Asian (EAS) and African (AFR)
groups in UK Biobank.

1201

1202