Efficient assembly of plant genomes: A case study with

evolutionary implications in *Ranunculus* (Ranunculaceae)

Karbstein, Kevin[1,2*], Choudhary, Nancy[3], Xie, Ting[4], Tomasello, Salvatore[1], Wagner,

Natascha D.[1], Barke, Birthe H.[1], Paetzold, Claudia[5], Bradican, John P.[1], Preick,

Michaela[6], Himmelbach, Axel[7], Stein, Nils[7,8], Papantonis, Argyris[4], Irisarri, Iker[9], de

Vries, Jan[10,11,12], Pucker, Boas[3], & Hörandl, Elvira[1*]

[1] *University of Göttingen, Albrecht-von-Haller Institute for Plant Sciences, Department of*

Systematics, Biodiversity and Evolution of Plants (with Herbarium), Göttingen, Germany

[2] *Max Planck Institute for Biogeochemistry, Department of Biogeochemical Integration, Jena,*

Germany

[3] *TU Braunschweig, Institute of Plant Biology, Braunschweig, Germany*

[4] *University Medical Center Göttingen, Institute of Pathology, Göttingen, Germany*

[5] *Senckenberg Naturhistorische Sammlungen, Dresden, Germany*

[6] *University of Potsdam, Institute for Biochemistry and Biology, Potsdam, Germany*

[7] *Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), Seeland, Germany*

[8] *Center of integrated Breeding Research (CiBreed), Department of Crop Sciences,*

Göttingen, Germany

[9] *Museo Nacional de Ciencias Naturales (MNCN-CSIC), Department of Biodiversity and*

Evolutionary Biology, Madrid, Spain

[10] *University of Göttingen, Institute for Microbiology and Genetics, Department of Applied*

Bioinformatics, Göttingen, Germany

[11] *University of Göttingen, Campus Institute Data Science (CIDAS), Göttingen, Germany*

[12] *University of Göttingen, Göttingen Center for Molecular Biosciences (GZMB), Department*

of Applied Bioinformatics, Göttingen, Germany

ORCID

Kevin Karbstein 0000-0003-1424-6557

Nancy Choudhary 0000-0002-2562-7905

Ting Xie 0000-0002-6648-7697

Salvatore Tomasello 0000-0001-5201-1156

Natascha D. Wagner 0000-0001-6623-7623

Birthe H. Barke 0000-0001-5379-6055

Claudia Paetzold 0000-0002-4128-6645

John P. Bradican 0000-0001-5650-6172

Axel Himmelbach 0000-0001-7338-0946

Nils Stein 0000-0003-3011-8731

Argyris Papantonis 0000-0001-7551-1073

Iker Irisarri 0000-0002-3628-1137

Jan de Vries 0000-0003-3507-5195

Boras Pucker 0000-0002-3321-7471

Elvira Hörandl 0000-0002-7600-1128

*   corresponding authors: *email: kkarb@bgc-jena.mpg.de or kevin.karbstein@tu-*
ilmenau.de, elvira.hoerandl@biologie.uni-goettingen.de

Running Head

Strategies to assemble non-model plant genome sequences

Abstract

Abstract

Currently, it is still a challenge - in terms of laboratory effort and cost, as well as assembly

quality - to unravel the sequence of large and complex genomes from non-model/crop plants.

This often hampers the study of evolutionarily intricate species groups. The species-rich

genus *Ranunculus* (Ranunculaceae) is an important angiosperm group for the study of

polyploidy, apomixis, reticulate evolution, and biogeography. However, neither

mitochondrial, nor high-quality nuclear genome sequences are available. This limits

phylogenomic, functional, and taxonomic analyses thus far. Here, we tested Illumina short-

read, Oxford Nanopore Technology (ONT) or PacBio/HiFi long-read, and hybrid-read

assembly strategies. We used the diploid progenitor species *R. cassubicifolius* (*R. auricomus*

complex), and selected the best assemblies in terms of completeness, contiguity, and quality

scores. We first assembled the plastome (156 kbp, 85 genes) and mitogenome (1.18 Mbp, 40

genes) sequences using Illumina and Illumina-PacBio-hybrid strategies, respectively. We also

present an updated plastome and the first mitogenome phylogeny of Ranunculaceae, including

studies of gene loss (e.g., *infA*, *ycf15*, or *rps*) with evolutionary implications. For the nuclear

genome, we favored a PacBio-based assembly three-times polished with filtered reads and

subsequently scaffolded into 8 pseudochromosomes by chromatin conformation data (Hi-C)

as the representative sequence. We obtained a haploid genome sequence with 2.69 Gbp,

94.1% complete BUSCO 'embryophyta_odb10' genes found, and 31,322 annotated genes.

The genomic information presented here will improve phylogenomic analyses in this species

complex, and will enable advanced functional, evolutionary, and biogeographic analyses for

the genus and beyond Ranunculaceae in the future.

Keywords

gene evolution, large plant genomes, plastome, mitogenome, nuclear genome, Illumina vs.

Nanopore vs. PacBio sequencing, *de novo* assembly strategies, *Ranunculus auricomus,*

Ranunculaceae

Significance Statement

The genus *Ranunculus* is a model system in flowering plants for polyploidy, apomixis,

evolution, and biogeography research. We present the first nuclear and mitochondrial genome

sequence, and the plastid evolution of Ranunculaceae. Using Illumina, ONT, and PacBio data,

we developed an efficient assembly strategy that can be applied to other non-model plants.

Results presented here are useful for improving population genomic and phylogenomic

analyses, and enable better functional analyses at species, genus and family level.

Resource Article

Introduction

Over the past decade, the scientific community has experienced a massive expansion in accessible sequencing technologies. For constructing genome assemblies, short-reads (e.g., Weisenfeld et al., 2018; Prjibelski et al., 2020), long-reads (e.g., Koren et al., 2017; Cheng et al., 2021), or a combination of both (hybrid-read; e.g., Di Genova et al., 2021; Gatter et al., 2021; Zimin and Salzberg, 2022) have frequently been used. Nevertheless, biologists have moved beyond short-read (100-250 bp) Illumina sequence data to perform genome assemblies. Short reads often produce incomplete assemblies, and lack the contiguity required to resolve repetitive regions that plague large genomes, leading to highly fragmented and incomplete *de novo* genome assemblies (Gatter et al., 2021; Rhie et al., 2021; Zimin and Salzberg, 2022). Today, long-read technologies, such as nanopore sequencing (Oxford Nanopore Technologies, ONT) or single molecule real time sequencing (SMRT; Pacific Biosciences), can bridge such gaps and improve the quality of genome assemblies. High-fidelity (HiFi) reads derived from SMRT sequencing are typically of high per-base quality (>99.5%) and up to 25 kbp in length. In contrast, ONT sequencing just crossed the 99% per-base accuracy threshold, read lengths are limited only by the input DNA fragment size thus reaching >1 Mbp, and can be easily performed in any laboratory (reviewed in Pucker et al., 2022).

HiFi or ONT sequencing, combined with proximity information obtained by chromosome conformation capture sequencing (Hi-C), has started to become prevalent for chromosome-scale, high-quality model but also non-model plant genome sequences (e.g., Mascher et al., 2017; Xie et al., 2020; Hoencamp et al., 2021; Qu et al., 2023). The quality and completeness of the assembly can be improved by polishing strategies using available, quality-filtered Illumina, ONT, or PacBio reads (e.g., Zimin and Salzberg, 2020; Dmitriev et al., 2021; Chang et al., 2023). The combination of low- to medium-sequencing coverage (that is, the proportion of the total assembly that is supported by the raw reads) long-read data with highly accurate, often already available Illumina sequences in hybrid assemblies is particularly attractive for improving genome assemblies when resources are limited. This is usually the case when working with non-model plants.

Producing "high-quality" genome sequences, that is genome assemblies characterized by > 90% completeness with respect to expected genome size, > 1-10 Mbp contiguity (N_{50}),

and > 90% recovery rate of conserved protein-coding genes (BUSCO; see assembly standards

of the Earth Biogenome Project; Table S1 in Lawniczak et al., 2022), represent an important

target for research groups. However, some metrics are rather adapted to animals. For example,

flowering plants have undergone many ancient or recent polyploidization events (Van De

Peer et al., 2017; Leebens-Mack et al., 2019; Hörandl, 2022) and are characterized by quite

large, repetitive genomes with duplicated sequence content. Consequently, a BUSCO

'embryophyta_odb10' score of > 90% single-copy genes can often not be met because genes

are often duplicated in the genome (Dmitriev et al., 2021) or is not informative for quality

assessment in non-coding, repetitive DNA regions (e.g., LTR assembly index; Mokhtar et al.,

2023). Irrespective of chosen metrics, efficient strategies in terms of laboratory effort, cost,

and assembly quality are needed to reliably reconstruct large and complex high-quality

genome sequences for non-model organisms. This often hampers the study of evolutionary,

biogeographic, and ecological relationships within groups characterized by intricate

evolutionary processes such as hybridization, polyploidy, and/or apomixis (Ennos et al., 2005;

Karbstein et al., 2024).

Ranunculus represents a cosmopolitan genus with approximately 600 species, of

which about 40% are polyploid (Hörandl et al., 2005) and have rather large genomes (ranging

from 1.8 to 24.6 Gbp; Leitch et al., 2019). The *Ranunculus auricomus* complex is one of the

largest polyploid apomictic plant species groups in Eurasia, with more than 840 described

apomictic taxa (Hörandl and Raab-Straube, 2015). The complex is a model system for

understanding the evolution and biogeography of young flowering plant groups formed by

reticulate evolution and allopolyploidy (Paun et al., 2006; Hörandl et al., 2009; Pellino et al.,

2013; Hodač et al., 2018; Karbstein et al., 2020b, 2020a, 2021, 2022; Tomasello et al., 2020;

Karbstein, 2021; Bradican et al., 2023, 2024; Hodač & Karbstein et al., 2023). It has also been

frequently studied for the molecular, cytogenetic, environmental, and evolutionary

mechanisms of apomixis (i.e., the asexual reproduction via seeds; Hörandl et al., 2024).

Apomixis is relevant for agricultural applications (e.g., Schmidt, 2020), and it is often

associated with hybridization and/or polyploidy, which are considered major evolutionary

forces in plant evolution. At least 30% of flowering plant species are of recent polyploid

origin, while > 25% of extant plant species are known to be involved in interspecific

hybridization or introgression events (Mallet, 2005; Soltis et al., 2015; Van De Peer et al.,

2017; Leebens-Mack et al., 2019; Hörandl, 2022). Consequently, improving (phylo)genomic

analyses for di- and polyploids, and observing regulatory mechanisms of apomixis (e.g.,

Brukhin et al., 2019) with *R. auricomus* reference genome sequences will help develop a

better understanding of plant evolution and biodiversity, and is therefore of general interest in
biological research. Organellar genome reconstructions are also valuable targets for studying
nuclear-organelle compatibility and the chimerism of cyto-nuclear complexes (Sloan, 2013;
Møller et al., 2021). Little is known, for instance, about plant mitochondrial genomes,
although mitochondrial-nuclear compatibility is important for speciation, evolution of
crossing barriers, and emergence of cytoplasmic male sterility (Postel and Touzet, 2020;
Møller et al., 2021). Elucidating cyto-nuclear complexes may be important for studying a
variety of evolutionary processes. For instance, plastid information may provide additional
insights into phylogenetic relationships (Tomasello et al., 2024), maternal progenitors of
hybrids (Karbstein et al., 2022), or the ability of hybridization (Sobanski et al., 2019; Postel
and Touzet, 2020), or mitochondrial information into plant sex evolution and determination
(e.g., male sterility; (Havird et al., 2015; Park et al., 2021; Hörandl, 2024).

The *R. auricomus* complex contains five newly circumscribed, predominantly diploid
sexual species (Karbstein et al., 2020b), which gave rise to the >800 described allopolyploid
apomictic taxa within the last ca. 800 kyr (Tomasello et al., 2020; Karbstein et al., 2022). The
diploid genome size of the *R. auricomus* complex (2n = 16) ranges between 6.12 and 6.30 pg
DNA based on DNA image cytometry (Hörandl and Greilhuber, 2002), recalculated as 5.998-
6.174 Gbp (Leitch et al., 2019). Therefore, *R. auricomus* genomes are large even at the
diploid level. Increasing proportions of repetitive regions are responsible for genome size
inflation (Bennetzen, 2005; Peška et al., 2019; Guo et al., 2021; Jayakodi et al., 2023), which
could also be relevant for *R. auricomus* genomes (Hörandl et al., 2009; Hodač et al., 2018;
Karbstein et al., 2022). However, so far, high-quality reference genome sequences for the
species complex, but also for the entire genus *Ranunculus* are missing. In Ranunculaceae, the
only other reference genome sequences are available from diploid *Aquilegia* and *Coptis*
species (Xie et al., 2020; Liu et al., 2021; Mokhtar et al., 2023), but these genomes are much
smaller (ca. 1 Gbp; Leitch et al., 2019), and no polyploid species complexes are known in
these genera (TROPICOS database v3.4.2, http://www.tropicos.org; CCDB database v1.66,
https://taux.evolseq.net/CCDB_web; Rice et al., 2015).

The evolutionary relationships among the diploid sexual progenitors of *R. auricomus*
have recently been clarified (Karbstein et al., 2020b; Tomasello et al., 2020). A diploid
Ranunculus reference genome sequence would be a major step forward in improving the
reconstruction of the species by facilitating read mapping to orthologous loci, allele phasing,
and subgenome assignments in hybrids; it would be the starting point for more sophisticated
approaches such as reference-guided genome assemblies (e.g., Eaton and Overcast, 2020),

polyploid phasing used for multilabeled species trees (e.g., Dauphin et al., 2018), or
allopolyploid network reconstructions (e.g., Jones, 2017; Šlenker et al., 2021). In addition, the
annotation of thousands of nuclear genes would also help enable sophisticated structural and
functional analyses in the species complex. New developments allow the incorporation of
external hints such as RNA-seq data or the annotation of homologous genes through
knowledge transfer from other species (Pucker, 2024). Tools are also needed for
automatically extracting and comparing group-specific annotations across genome sequences
from databases, which are essential for evolutionary studies. In *Ranunculus*, and in
Ranunculaceae, such large-scale studies are missing. Annotated high-quality genome
sequences are essential for functional genomics in the context of apomixis, adaptation, and
gene expression (e.g., Syngelaki et al., 2021), and for studying loci under selection and
mutation-selection dynamics (e.g., Del Amparo et al., 2021) that could so far only be
performed on (incomplete) transcriptomes (Pellino et al., 2013; Paetzold et al., 2022).

To generate plastid, mitochondrial, and high-quality nuclear genome assemblies of a
diploid species of the *R. auricomus* complex, we applied different assembly strategies using
short Illumina reads and long ONT or PacBio reads under low to medium sequencing
coverage, with the best nuclear-genome assembly being scaffolded by Hi-C reads. This
represents a typical scenario when working with non-model plant groups on a limited budget.
We thus focused on applying different approaches based on the available sequencing data. To
study organellar gene evolution in detail, we also downloaded and analyzed all available
sequences from NCBI.

Methods

Plant Material

In the present study, analyses were performed on diploid sexual individuals (2n = 16)
of *Ranunculus cassubicifolius* (*R. auricomus* complex; taxonomic treatment follows Karbstein
et al., 2020b). Chromosome numbers, measurements of ploidy and mode of reproduction are
published in Hörandl and Greilhuber (2002) under '*R. carpaticola*', and in Karbstein et al.
(2020b, 2021). Fresh leaf material was harvested in 2019 from individual EH8483/10 (NCBI
BioSample SAMN27753242), sampled in Revúca, Slovakia (48°41'39"N, 20°07'42"E;
Hörandl and Greilhuber, 2002; under '*R. carpaticola*'). High-molecular-weight DNA was
extracted immediately after harvest and used for Illumina DNA sequencing. The same
population was already used for transcriptome (Pellino et al., 2013) and population genetic

analyses (Paun et al., 2006; Hodač et al., 2018). In autumn 2021, we harvested fresh, young

leaf material for ONT sequencing, after 12-hour dark-adaptation at room temperature in the

early morning from another conspecific individual collected nearby (LH040/02,

SAMN27753242; 48°40'38"N, 20°06'24"E; Figure 1; Karbstein et al., 2020b) because the

previously used individual died. In 2023, we used individuals from the same population for

PacBio and Hi-C sequencing, LH040/06 and LH040/05+06 (SAMN44350262), respectively,

because the previously harvested individual did not sprout again. All plants were cultivated at

the Old Botanical Garden of the University of Göttingen.

Figure 1. (a) Habitat of *Ranunculus cassubicifolius* s.l. (previously listed as '*R. carpaticola*')

population LH040 in a forest near Liptov, Slovakia (4th May 2018, L. Hodač & K. Spitzer;

details in Karbstein et al., 2020b). (b) Shown are the typical undissected basal leaves *in situ*

and a typical *R. cassubicifolius* flower from a different population (LH009) and (c) the scan of

a representative herbarium specimen from LH040. Field images: L. Hodač.

Illumina DNA extraction and sequencing

For Illumina short-read sequencing, genomic DNA (gDNA) was directly extracted

from ca. 200 mg of fresh leaf material from a single *R. cassubicifolius* s.l. individual

(EH8483/10) using liquid nitrogen and the DNeasy PowerPlant Pro Kit (Qiagen, Hilden,

Germany) according to the manufacturer's instructions. DNA size and concentration were

measured with the Qubit Fluorometer 2.0 and the Qubit dsDNA HS Assay Kit (ThermoFisher

Scientific, Waltham, USA), and the Genomic DNA ScreenTape device (Agilent
Technologies, Santa Clara, USA). The extraction yielded 16 ng/µl of DNA with a median
fragment length of 19 kb.

Library preparation followed the 10x Chromium Controller Genomics Kit (10x
Genomics, Pleasanton, USA) according to the manufacturer's instructions. Library quality
was assessed by a qPCR based on the NEBNext Library Quant Kit (New England Biolabs
Inc., Ipswich, USA). Sequencing with standard Illumina primers was performed on a NextSeq
500/550 instrument (Illumina, San Diego, USA) with 2 x 150 bp reads.

We converted the raw output into FASTQ files using SUPERNOVA v2.1.1 (10x
Genomics, San Francisco, USA; Marks et al., 2017; Weisenfeld et al., 2018) and
BLC2FASTQ v2.20 (Illumina, San Diego, USA), resulting in 51 Gb raw FASTQ files. We
conducted polyX-tail and adapter trimming, and quality filtering of Illumina raw reads using
ATRIA v3.1.2 (Chuan et al., 2021) with default settings, and BBTools/BBMap v39.01
('repair.sh'; (Bushnell, 2022) to reorder filtered paired-end reads.

Nanopore DNA extraction and sequencing

For ONT, gDNA extraction was mainly performed according to the Qiagen protocol
for the isolation of gDNA from plants and fungi, and the manufacturer's instructions of the
Qiagen Genomic DNA Buffer Set using Genomic Tip 20/G (see details in Text S1). Several
protocol modifications were applied to increase DNA yield and to optimize DNA quality (see
also Vaillancourt and Buell, 2019; Li et al., 2020). Three gDNA extractions were performed
in parallel from a single *R. cassubicifolius* s.l. individual (LH040). Fresh leaf material (ca. 75
mg) was frozen in liquid nitrogen and ground with a mortar and pestle, and transferred into a
2.0 ml Eppendorf tube containing 400 µl API Qiagen extraction buffer (Qiagen DNeasy Plant
Mini Kit).

The mixture was incubated at 40°C for 1.5 hours with gentle agitation. Then, 4 µl
Qiagen RNase A (100 mg/ml) was added, and the mixture was incubated again at 37°C for 30
minutes, followed by the addition of 18 µl Qiagen Proteinase K (20 mg/ml), and incubation at
50°C for 2 h with gentle agitation. The mixture was centrifuged at 12,000 x g for 20 min, and
ca. 400 µl of the lysate was transferred with wide-bore tips to a buffer QBT-equilibrated
Genomic Tip. DNA was washed with 5 x 1 ml QC Buffer, and eluted with 0.8 ml 55°C pre-
warmed QF Buffer into a new 2.0 ml Eppendorf tube. DNA was precipitated by adding 70%
room-temperature isopropanol and inverting the tube 20 times, followed by incubation at
room temperature for 15 min. Using a glass rod, the DNA was rolled up and placed into a new

1.5 ml Eppendorf tube containing ca. 0.8 ml ice-cold 80% ethanol. The DNA was air dried,

resuspended in 200 µl autoclaved ddH$_2$O, and all three extractions were pooled into a single

tube, stored at 4 °C until library preparation.

DNA concentration was assessed using the Qubit Fluorometer 3.0 and the Qubit

dsDNA HS Assay Kit, and the DNA purity was checked with the NanoDrop 2000

Spectrophotometer (ThermoFisher Scientific, Waltham, USA; Text S1). DNA fragment

length distribution and RNA absence were checked on a 2% agarose gel via electrophoresis

(50 V, 90 min) and the Quick-Load 1 kb DNA Ladder (New England Biolabs Inc., Ipswich,

USA; Text S1, Fig. S1). After several dilution steps, we obtained a final DNA yield of 61.2

µg (51.0 ng/µl * 1200 µl ddH$_2$O).

Library preparation was conducted using the ONT Ligation Sequencing Kit SQK-

LSK110 optimized for high throughput and long reads (ONT, Oxford, UK), and applicable

for singleplex gDNA sequencing. We adjusted the DNA concentration to 1000 ng in 47 µl

(ca. 21.5 µl/ng) and followed the manufacturer's instructions for library preparation (protocol

vGDE_9108_v110_revL_10Nov2020, available at community.nanoporetech.com) with a few

modifications. Incubation times were increased up to 15 min, the concentration of ethanol

wash buffer was increased to 80%, and DNA fragments > 3 kbp were enriched using the

buffers provided with the kit. Active pores ranged from ca. 800-1500. Flow cell 1 was loaded

with freshly prepared libraries, flow cell 2 with one freshly prepared and one library stored at

-80°C for two days, and flow cells 3-6 with two libraries stored at -80°C for 1-2 weeks.

Freezing of gDNA had no apparent effect on N$_{50}$ fragment size (see also Results &

Discussion and flow cell reports on FigShare). We used a MinION 101B device, the ONT

software MinKNOW v21.11.7, 21.11.9, and 22.03.5 on a local Linux system, and six R9.4.1

flow cells following the manufacturer's instructions for priming and loading. After 1-2 days

of run time, the old library was exhausted and replaced with a second library using the

supplied flow cell wash kit (protocol vWFC_9120_v1_revB_08Dec2020). Sequencing was

run for 3-5 days per flow cell until the majority of pores were inactive (> 99%).

Reports per flow cell can be accessed via FigShare. Basecalling was performed with

the   ONT   software   GUPPY   v6.0.1   specifying   the   configuration   file

"dna_r9.4.1_450bps_fast.cfg" for fast base-calling (in preruns, the difference of median per-

base quality between 'fast' and the substantially slower 'high accuracy' was quite small, i.e.,

97 vs. 98%, see FastQC files on Figshare). After that, NanoFilt v2.2.0 (De Coster et al., 2018)

was applied to remove the first 50 bp of the reads that were usually of low quality.

PacBio DNA extraction and sequencing

High-molecular-weight gDNA was isolated from fresh leaf material of a single *R. cassubicifolius* s.l. individual (LH040) with the NucleoBond HMW DNA kit (Macherey Nagel, Germany), following the manufacturer's instructions. The integrity of the DNA was determined by the Femto Pulse System (Agilent Technologies Inc, CA, USA). The DNA concentration was measured with the Qubit dsDNA High Sensitivity assay kit (Thermo Fisher Scientific, MA, USA). For the construction of a HiFi library, 7.5 µg HMW DNA was fragmented (speed 29) using the Megaruptor 3 device (Diagenode SA, Seraing, Belgium).

In total, two HiFi libraries were constructed and sequenced into one Revio 25M SMRT cell. Libraries were generated according to the 'Procedure & Checklist - Preparing whole genome and metagenome libraries using SMRTbell® prep kit 3.0' manual (102-166-600, Pacific Biosciences of California Inc., USA). For size fractionation, a SageELF (Sage Science, USA) device was employed. The final sequencing library size (16.2 kb) was measured using the Femto Pulse System (Agilent Technologies Inc, USA). Polymerase-bound SMRTbell complexes were formed according to standard manufacturer's protocols. Sequencing (HiFi CCS) was conducted using the Pacific Biosciences Revio instrument (24 h movie time, loading concentration 230 pM, 2 h pre-extension time, diffusion loading, mean insert length according to SMRT link raw data report: 18.0 kb) following standard manufacturer's protocols. The Revio 25M SMRT cell generated 82 Gb HiFi CCS data. All steps were done at IPK Gatersleben.

HiFiAdapterFilt v2.0.0 (Sim et al., 2022) was run with default settings to convert the HiFi reads from BAM to FASTQ format, and to remove reads with remnant PacBio adapter sequences. The quality of the sequenced reads was checked by FastQC v0.11.4 (Andrews, 2010) (reports on FigShare).

Chromosome conformation capture (Hi-C) DNA extraction and sequencing

The Hi-C sequencing library was generated from fresh leaf material of two *R. cassubicifolius* s.l. individuals (LH040) with the *DpnII* enzyme, essentially following the method described in Padmarasu et al. (2019). The library was sequenced (paired-end: 2 x 111 cycles, two indexing reads: 8 cycles, v1.5 chemistry) using the NovaSeq6000 instrument (Illumina, Inc., San Diego, USA) according to standard manufacturer's protocols. All steps were performed at IPK Gatersleben.

Genomic Data Analysis

The plastid genome. The plastome sequence was assembled using Illumina short-read data and the software GetOrganelle v1.7.5.3 (Jin et al., 2020) as recommended by Freudenthal et al. (2020), specifying the 'embplant_pt' database, k-mer size range between 45 and 121, and as seed the *Ranunculus repens* plastome (154,247 bp, GenBank accession number NC_036976; Dann et al., 2017). The average embplant_pt base sequencing depth was 560x. The plastome was annotated using CPGAVAS2 (Shi et al., 2019) with default settings and 2,544 reference plastomes, and visualized with CPGView (Liu et al., 2023). We checked and edited the annotation results with Geneious R11 2023.1.2 (Kearse et al., 2012). Results were also visually inspected using Bandage v0.9.0 (Wick et al., 2015). The final annotated plastome sequence (Fig. 2) was deposited in GenBank (NC_077490). We applied MAFFT v7.490 to align and investigate detailed differences between *R. cassubicifolius* and available *Ranunculus* plastome sequences. The plastid raw data (SRA: PRJNA826743) and the annotated plastome (GenBank: NC_077490) are deposited in NCBI.

The mitochondrial genome. Two alternative assembly strategies were followed. First, we used GetOrganelle with the same settings as described above, except that we specified "embplant_mt" and as seed used the closest available mitogenome (*Hepatica maxima*, GenBank accession number NC_053368.1; Park and Park, 2020). The average embplant_mt base sequencing depth (i.e., the average number of reads per position) of Illumina reads was estimated at 92x. To improve the fragmented mitogenome assembly, we first mapped ONT reads against the contigs derived from GetOrganelle and extracted the mapped reads using SAMtools v1.9 ('faidx' and 'view'; Danecek et al., 2021), MINIMAP2 v2.27 ('-ax map-ont'; Li, 2018), and BEDTools v2.29.1 ('bamtofastq'; Quinlan and Hall, 2010). We applied MaSuRCA v4.0.9 (Zimin and Salzberg, 2022) to perform a hybrid assembly based on extracted ONT reads and "embplant_mt" hitting Illumina reads derived from the GetOrganelle assembly process.

Then, we retained an ONT-based mitogenome sequence in 18 contigs and a total length of 351,681 bp. We aligned contigs in Geneious to merge overlapping regions, which resulted in 4 contigs, onto which Illumina and ONT reads were mapped using minimap2 ('-ax sr' and '-ax map-ont', respectively). Alignments were inspected with the Integrative Genomics Viewer (IGV) v2.16.1 (Thorvaldsdottir et al., 2013) and Geneious to ensure that previous gaps were spanned by raw reads (see alignments and information concerning contig borders on FigShare). Using Geneious, we inferred a mean Illumina depth of 62x (2.87% sites

with no coverage) and ONT depth of 3640x (9.88% sites with no coverage), respectively.
Remaining gaps were closed manually based on mapping results of Illumina and ONT reads.
The assembly was annotated using MitoHiFi pipeline v3.2.1 (Uliano-Silva et al., 2023)
including gene annotation with *H. maxima* as reference, and visualized with PMGmap (Zhang
et al., 2024; http://www.1kmpg.cn/pmgmap) web browser tool. We ran RepeatModeler2
v2.0.5 to additionally check for dispersed repeats.

In a second assembly strategy, we used PacBio HiFi reads, which were subjected to
the MitoHiFi all-in-one pipeline to assemble, annotate, and draw the mitogenome based on
the PacBio HiFi reads. The pipeline represents a similar strategy as described before by first
mapping length-filtered HiFi reads to the reference mitogenome, followed by assembly using
Hifiasm and annotation with MitoFinder (Allio et al., 2020). The pipeline resulted in a very
small mitogenome of about 125 kb (about 10% of the expected length). Therefore, we
performed a hybrid assembly strategy with the HiFi reads as described above. This resulted in
40 raw contigs with 1,913,766 bp summarized into 9 contigs that were aimed to be arranged
like the ones in the ONT hybrid mitogenome. We inferred a mean Illumina sequencing depth
of 269x (1.96% sites with no coverage) and HiFi depth of 3324x (0.12% sites with no
coverage). Finally, the mitogenome sequences were retained in a single contig based on ONT-
and PacBio-hybrid assemblies. Due to substantial intragenomic rearrangements, we
performed progressiveMauve (2015-02-25; Darling et al., 2010) instead of MAFFT in
Geneious to investigate sequence differences between both assemblies, and between *R.*
cassubicifolius and available Ranunculaceae assemblies.

Geneious was used to inspect and edit the annotation results. The resulting GeneBank
file was converted into a feature table file for GenBank submission using the GB2SEQUIN
tool (Lehwark and Greiner, 2019) within the MPI-MP CHLOROBOX webserver. Finally, we
selected the PacBio-based mitogenome because of its (4x) higher sequencing depth of
mapped Illumina reads and 1.5x and 82x fewer Illumina and long-read sites without coverage,
although the gene content (but not the gene block arrangement, see FigShare) was the same
compared to the ONT-based mitogenome. The annotated mitogenome is illustrated in Figure
3. The mitochondrial raw data (SRA: PRJNA831351) and annotated mitogenome (GenBank:
PP657143) are deposited in NCBI.

Phylogenomics and organellar gene evolution within Ranunculaceae. To study
gene evolution within Ranunculaceae, we downloaded all available organellar genome
sequences from NCBI using a custom Python script 'findGenome.py'

(https://github.com/KK260/NCBI-Genome-Tools/). Applying the following specifications

"complete genome" AND "Ranunculaceae" AND "chloroplast"/ "mitochondrion", we found

1161 plastome and 16 mitogenome sequences on NCBI that were initially filtered by

duplicate sequence removal, a maximum of 1 individual per species and 2 species per genus,

and selection of the latest RefSeq 'NC' records. Taxon names were checked for synonyms

using information from the Catalogue of Life on GBIF. This yielded 308 plastomes including

6 outgroup taxa of Berberidaceae and Papaveraceae (295 taxa), and 10 mitogenomes

including 1 outgroup taxon of Papaveraceae (none available for Berberidaceae, 10 taxa) (see

Table S1).

Plastome sequences were aligned using MAFFT with mode 'FFT-NS-2' (Katoh and

Standley, 2013), following the deletion of sites with less than 50%, 70%, and 90% of

available samples using Goalign v0.3.7 (Lemoine and Gascuel, 2021). We then ran

ModelTest-NG v0.1.7 (Darriba et al., 2020), which determined 'TVM+I+G4' (unfiltered),

'TIM1+I+G4' (min50), and 'GTR+I+G4' (min70, min90) as best-fit evolutionary models for

the datasets. Maximum likelihood trees were inferred with RAxML-NG v1.2.1 (Kozlov et al.,

2019) specifying the estimated sequence model and 100 pseudoreplicates of non-parametric

bootstrapping, and transfer bootstrap expectation (TBE) metric. To further assess branch

support,   Quartet   Sampling   (QS)   v1.3.1   (Pease   et   al.,   2018;

https://www.github.com/fephyfofum/quartetsampling) scores were calculated with default

settings   and   visualized   using   the   supplied   R   script

(https://github.com/ShuiyinLIU/QS_visualization). Following the approach of Karbstein et al.

(2020b), the final phylogenetic tree was selected according to the highest mean TBE support

and optimal QS scores (QC=1, QD=0, QI=1) using the R package PHYTOOLS v2.0-3

(Revell, 2012, 2024; https://github.com/liamrevell/phytools). Finally, we chose 'min90' (126

kbp) with the lowest number of gaps excluding poorly assembled non-coding DNA regions,

and the highest TBE and QS values in the phylogeny as the optimal alignment (see details in

Figs. S2-S4 and Text S2, and alignments on FigShare). Branches of all tribes except

Ranunculeae were collapsed with FigTree v1.4.4 (Rambaut, 2014), gene gain and loss events

per genus were mapped onto the tree (Fig. 4a).

The mitogenome phylogeny was built based on coding regions due to substantial

genomic rearrangements and size differences present among sequences. Using a custom

Python script 'assembleGenes.py' (https://github.com/KK260/NCBI-Genome-Tools/), we run

Astral v5.7.8 with 100 bootstraps based on collapsed gene trees of low support previously

inferred by RAxML_NG with 100 bootstraps, and mapped gene features onto the resulting

tree. In Ranunculaceae, the loss of genes per group derived from the feature table of

'assembleGenome.py' was manually verified by mapping extracted genes from closely

related Ranunculaceae species to the sequence of interest in Geneious, and edited if necessary.

In *R. cassubicifolius*, we aimed to find lost organellar genes by mapping extracted genes from

closely related *Ranunculus* species to the nuclear genome sequence trees (Fig. 4b).

The nuclear genome. We applied different genome assembly approaches with default

settings and a haploid genome size of 3.2 Gbp and RAM of at least 1.5 TB, unless otherwise

specified: (i) using Illumina short-reads, we performed analyses with Supernova

(incorporation of all reads, acceptance of extreme coverages) and SPAdes v3.13.2 (Prjibelski

et al., 2020); (ii) using Illumina short-reads and ONT or PacBio/HiFi long-reads, we run

SPAdes (untrusted Supernova contigs for gap filling), MaSuRCA v4.1.0 (Zimin et al., 2017),

MuCHSALSA v0.02 (Gatter et al., 2021) and WENGAN (M) v0.2 (Di Genova et al., 2021);

(iii a) using ONT long-reads, we performed analyses with Canu v2.1.1 (Koren et al., 2017)

and Flye v2.9.1 (Kolmogorov et al., 2019) option '--nano-raw'), followed by the best

polishing strategy of Dmitriev et al. (2021) of two-times via filtered ONT reads using Racon

v1.4.3 (Vaser et al., 2017) and Medaka v0.7.1 (https://github.com/nanoporetech/medaka), and

once using Illumina reads via the POLCA function (Zimin and Salzberg, 2020); (iii b) using

HiFi long-reads, we run Canu, Flye, and Hifiasm v0.19.8 (Cheng et al., 2021), followed by

the previous polishing strategy except for the use of polishCLR v1.1.0 (Chang et al., 2023)

instead of Medaka. To ensure a fair comparison between assemblies based on ONT or PacBio

reads within the same tool, we downsampled PacBio reads to 16x coverage using reformat.sh

from BBTools and reran all analyses.

Statistics for genome assemblies were calculated using QUAST v5.0.1 (Gurevich et

al., 2013), and are summarized in Table 1 (reports on FigShare). Assembly completeness in

terms the expected single-copy genes in inherent to land plants was estimated by

'Benchmarking Universal Single-copy Orthologs' (BUSCO v5.0.0; Waterhouse et al., 2018;

Manni et al., 2021) with the 'embryophyta_odb10' 2020-09-10 dataset containing 60 species

and 1375 core genes. The assembly with the highest completeness, largest contig size, and

best BUSCO scores (Table 1, Fig. 3, reports on FigShare) was selected for scaffolding with

Hi-C data and genome annotation. Filtered Hi-C reads were mapped to the HiFi polished

contigs using Juicer v1.6 (Durand et al., 2016) with default parameters. Paired reads mapping

to different contigs were used for the Hi-C-associated scaffolding. We applied the 3D-DNA

pipeline v170123 (Dudchenko et al., 2017) to order and orient the clustered contigs with sizes

> 15,000 bp to produce chromosome-level scaffolds. Misassembles were identified and fixed

manually using assembling mode in the Juicebox software, which was manually corrected

based on the neighboring interactions. The resulting Hi-C contact map is shown in Fig. S5. To

fulfill NCBI quality standards, we screened for non-plant contigs using FCS-GX v0.5.4

(Astashyn et al., 2024;https://github.com/ncbi/fcs) and for mitochondrial contigs using seqkit

v2.8.2 (Shen et al., 2016) removing 552 short contigs in total. The functional annotation was

performed using InterProScan5 (Jones et al., 2014) and eggNOG-mapper (Cantalapiedra et

al., 2021) using the Funannotate "annotate" function.

The genome assembly was then subjected to the identification of tandem repeats (TRs)

and transposable elements (TEs). Tandem Repeats Finder v4.10.1 (Benson, 1999) using

parameters 'Match=2, Mismatch=5, Delta=7, PM=80, PI=10, Minscore=5, and

MaxPeriod=2000', and EDTA v2.2.1. The EDTA pipeline integrates the results of multiple

high-performing tools: LTR-FINDER (Xu and Wang, 2007; Ou and Jiang, 2019),

LTR_retriever (Ou and Jiang, 2018), Generic Repeat Finder (Shi and Liang, 2019), TIR-

Learner (Xiong et al., 2014; Su et al., 2019), HelitronScanner (Xiong et al., 2014; Zhang et

al., 2022), and TEsorter, to achieve accurate and efficient TE identification and classification.

Given the limited availability of RNA-seq reads from *R. cassubicifolius*, we used a

multi-species RNA-seq mapping approach. This strategy included RNA reads from multiple

Ranunculus species (37 individuals, Table S1), retrieved using prefetch and fasterq-dump

from NCBI SRA toolkit v3.1.0 (https://github.com/ncbi/sra-tools). RNA data was mapped

onto the genome using HISAT2 v2.2.1 (Kim et al., 2019).

Structural annotation of protein-coding genes used a combination of three approaches.

(i) First, homology-based gene prediction using GeMoMa v1.9 (Keilwagen et al., 2016,

2018), using annotations from three Ranunculaceae and eleven Ranunculales species (Table

S2). The coding genes from these 14 species were aligned to the *R. cassubicifolius* genome

sequence using MMseqs2 v15-6f452 (Steinegger and Söding, 2017) following the default

GeMoMa setting. GeMoMa is based on the conservation of amino acid sequences and intron

position conservation among closely related species. To refine intron boundaries, RNA-seq

data from five recently published diploid *R. auricomus* transcriptomes (Paetzold et al., 2022)

were incorporated. The resulting fourteen gene annotation sets were merged and filtered using

the GeMoMa Annotation Filter (GAF) with parameters: f="start=='M' and stop=='*' and

(isNaN(score) or score/aa>=1.50 or (score/aa>0.8 and avgCov>50)) and avgCov>0.0 and

(iAA>=0.5 or pAA>=0.5) and (evidence>1 or tpc>=0.8)" and atf="tie==1 and

sumWeight>4".

(ii) Second, we used the BRAKER3 pipeline (Gabriel et al., 2024), which uses ab

initio gene prediction using AUGUSTUS v3.5.0 (Stanke et al., 2006, 2008) and GeneMark-

ETP v1.02 (Brůna et al., 2024), which is corrected by extrinsic evidence provided by RNA-

seq and protein homology. For protein evidence, we used 'Viridiplantae' library (Kuznetsov

et al., 2023) and annotated NCBI genomes of Ranunculaceae species (*Aquilegia caerulea*

(GCA_002738505.1), *Thalictrum thalictroides* (GCA_013358455.1), and *Coptis chinensis*

(GCA_015680905.1)). The resulting BRAKER3 annotation was filtered using GeMoMa

Annotation Filter (GAF) with parameters: f="start=='M' and stop=='*' and avgCov>0.0 and

tpc>0.5" and atf="tie==1 or sumWeight>1".

(iii) Third, we used Funannotate v1.8.15 (https://github.com/nextgenusfs/funannotate;

Palmer and Stajich, 2022), initially developed for fungi but now capable of handling larger

genomes. Repetitive contigs were cleaned from the genome using minimap2 v2.26 and simple

repeats were masked using TANTAN ('-s arabidopsis'; Frith, 2011). Approximately 8.04%

(217 Mbp) of the genome sequence was masked. Funannotate's gene prediction involved

three steps: train, predict, and update. In all steps, the parameter '--max_intronlen 1000000'

was used. The filtered genome and the transcriptome sequences were used to train the

annotation process by a genome-guided transcriptome assembly using Trimmomatic v0.39

(Bolger et al., 2014), Trinity v2.8.5 (Grabherr et al., 2011), HISAT2, kallisto v0.46.1 (Bray et

al., 2016), and PASA v2.5.3 (Haas et al., 2008) to filter, normalize, and cluster (genome-

guided) transcriptomic data. The sorted alignments, transcripts, UniProt protein library, and

transcriptome annotations were the input for the gene prediction process performed by

AUGUSTUS v3.5.0, GlimmerHMM (Majoros et al., 2004), and SNAP (Korf, 2004), and

results were summarized with EVidenceModeler v1.1.1 (Haas et al., 2008). Additional

parameters '--busco_db eudicots_odb10 --organism other --busco_seed_species arabidopsis'

were used in this step.

The annotations resulting from GeMoMa, BRAKER3, and Funannotate were

compared based on BUSCO completeness scores. Based on these comparisons, the GeMoMa

and BRAKER3 gene predictions were merged using GeMoMa Annotation Filter (GAF) with

the following parameters: f="start=='M' and stop=='*' and avgCov>0 and tpc>0 and aa >=50

and ((isNaN(score) and (tie==1 or isNaN(tie) and tpc>0.5)) or (tie>0.5 or tpc>0.5))" and atf=

" tie==1 and tpc>0.5 and (isNaN(score) or sumWeight>2)". Gene models in disagreement

with RNA evidence data were fixed manually. The annotation quality was evaluated using

BUSCO v5.6.1 with the embryophyta_odb10 database.

545    The nuclear genome sequences and annotated protein-coding genes (Fig. 6) are

546 deposited in NCBI under SAMN44350262 in the Bioproject PRJNA831351 (until publication

547 in FigShare). Non-coding RNA features, including rRNAs, tRNAs, and short ncRNAs were

548 identified using Infernal v1.1.5 (Nawrocki and Eddy, 2013) cmscan, which scanned

549 covariance models from the Rfam database v15 (Ontiveros-Palacios et al., 2024).

550 Additionally, tRNAscan-SE v2.0.12 (Chan et al., 2021) and barrnap v0.9 (available at

551 https://github.com/tseemann/barrnap) were used to enhance the detection of tRNAs and

552 rRNAs, respectively.

553    All data analyses were performed on the HPC cluster of the GWDG (Göttingen,

554 Germany), except for the genome annotation which was run on the de.NBI cloud.

Results and Discussion

More than 504 million filtered Illumina reads (150 bp) were generated of *R. cassubicifolius*, resulting in 70.7 Gbp raw data and genome coverage of 22x (mean base accuracy of 99.97%), assuming a haploid genome size of 3.2 Gbp. In addition, the final ONT long-read dataset contains 51.6 Gbp and 5.83 million reads (N_{50}=22.6 kbp), yielding a coverage of 16x (mean base accuracy of 96.62%). The final HiFi (PacBio) long-read dataset harbors 79.1 Gbp and 4.40 million reads (N_{50}=17.9 kbp), resulting in a coverage of 25x (mean base accuracy of 99.98%).

Plastome features and gene evolution within *Ranunculus* and Ranunculaceae. Based on short Illumina reads, we retained a circular plastid genome sequence with 156,233 bp in two states caused by the differing orientations of the inverted repeat (IR) regions ("flip-flop"; Stein et al., 1986; Walker et al., 2015). The plastome shows a typical tetrapartite structure of a small single-copy region (SSC, length: 19,826 bp) and a large single-copy region (LSC, length: 85,629 bp) separated by two IR regions (length: 2 x 25,389 bp). The plastome contains 85 (78 unique) protein-coding genes, 37 (30 unique) tRNAs, and 8 (4 unique) rRNAs, resulting in a coding content of 53.2%. We also observed 35 SSR, 32 long tandem, and 34 dispersed repeat motifs (2.2% repeat content). For example, in comparison with the closely related *R. sceleratus* (nested within the sister clade to sect. *Auricomus*) and the more distantly related *R. repens* (nested within a different subgenus within Ranunculaceae), *R. cassubicifolius* shows a similar number of unique protein-coding genes (78/77/78), tRNAs (30/28/29), and rRNAs (4/3/4). The plastome size of *R. cassubicifolius* was similar to that of *R. sceleratus* (+96 bp, 156,329 bp), but larger compared to *R. repens* (-1,986 bp; 154,247 bp), mainly due to additional mono-, oligo-, or polynucleotide insertions between different protein-coding genes (e.g., see positions at 15,754 bp or 126,944 bp in alignment file(s) on FigShare).

Within Ranunculaceae, the plastome sequence of *R. cassubicifolius* is quite average, considering the longest (167 kbp; *Helleborus atrorubens*) and shortest (150 kbp; *Myosurus apetalus*) plastomes. A clear phylogenetic trend of size evolution in Ranunculaceae is not recognized (Fig. 4). With similar gene contents, length variation can be mainly attributed to non-coding regions (Fig. 4a, and feature tables and alignments on FigShare). However, Long et al. (2024) observed in *Myosurus* that gene length and number can also be responsible for smaller plastomes, possibly reflecting adaptation to new environments, or an effect of annual life cycles, as known from nuclear genomes (e.g., Lunkova and Ivanov, 2022). In general,

compared to other flowering plant lineages, we detected usual size variation and no substantial plastome rearrangement among investigated plastome sequences of *Ranunculus* and Ranunculaceae, typical for an obligate autotrophic lifestyle (Petersen et al., 2018; Konupková et al., 2024).

The maximum likelihood plastome phylogeny with 292 taxa revealed a fully resolved and well-supported phylogeny of Ranunculaceae (TBE>85; Figs. 4 and S4, Text S3). Low branch support occurs within species-rich genera such as *Anemone, Hepatica,* or *Aconitum* and is probably due to fast radiation and the slow evolutionary rates of plastomes (Wang et al., 2016; Zhai et al., 2019). Almost all subfamilies or tribes are monophyletic, and branching is widely consistent with previous studies on plastome sequences (35 taxa) or four nuclear and plastid loci (105 taxa) (Wang et al., 2009; Zhai et al., 2019). Within *Ranunculeae*, *Oxygraphis*, *Ficaria*, *Myosuru*s*, and *Ceratocephala* represent close relatives of *Ranunculus*, in agreement with Emadzade et al. (2010). Compared to Zhai et al. (2019) and Ji et al. (2023), the here presented plastome phylogeny is the most recent and taxon-rich to date of Ranunculaceae. The plastome phylogeny of *Ranunculus* widely fits the one presented in Ji et al. (2023), that is, an Eurasian-American (clade 1), European-Asian clade including *R.* sect. *Auricomus* (clade 2), and the predominant Asian clade including aquatic taxa (clade 3). The phylogeny is also largely congruent with combined plastid-nuclear marker trees in Hörandl and Emadzade (2012), but based on a smaller taxon sampling and therefore not representative for this big genus. Further plastome sequencing efforts would allow a more taxon-rich and -balanced phylogenomic reconstruction of Ranunculaceae.

Regarding gene evolution within Ranunculaceae and particularly *Ranunculus*, we frequently did not detect some specific genes (*infA*, *pbf1*, *rpl32*, *rps16*, and *ycf15*) suggesting a potential loss (Fig. 4a; see feature tables on FigShare). Gene transfers from the chloroplast to the nucleus have been reported for *infA* but also for *accD, rpl22, rpl20, rpl32, rpl23, rps7, rps16, ycf1*, and *ycf2* in seed plants (Park et al., 2015; Zhou et al., 2023). In our study, the plastid gene *rpl32* was lost in the common ancestor of *Caltheae, Asteropyreae*, *Cimifugeae,* and *Nigelleae*, and *Delphinieae* and reappeared in *Callianthemeae* and all remaining subfamilies/tribes. The ribosomal gene *rpl32* is involved in protein biosynthesis and plastid function. Its putative loss and functional transfer to the nuclear genome have been observed in other clades within Ranunculaceae (Park et al., 2015; Zhai et al., 2019; Park and Park, 2020). The gene *rpl32* may be involved in rapid adaptation to novel environments, such as in *Myosurus* adapting from mesophytic to wet habitats during climate change over the last 5 Ma

(Long et al., 2024). In *R. cassubicifolius, rpl32* shows a typical length of 162 bp and is not

reduced as in *Myosurus* (138 bp), or in alpine *Clematis alpina* (69 bp).

The plastid gene *infA* is present in the outgroups and all tribes of Ranunculaceae but

appears to have been lost several times (22x in *Ranunculus*), including *R. cassubicifolius*.

Independent losses and transfers to the nuclear genome are already documented for various

angiosperm lineages, making *infA* one of the most mobile plastome genes in plants (Millen et

al., 2001). In the case of *R. cassubicifolius*, we were even able to confirm the loss of *infA* in

the plastome sequence and its transfer to the nuclear genome (two presumably functional,

size-reduced (ca. 150-160 bp) copies exist in chromosomes 1 and 4), a result not previously

reported in any other plant. The gene *infA* encodes the translation initiation factor 1 important

for ribosome assembly, translation initiation, and thus cell viability (Cummings and Hershey,

1994). Transfer to the nucleus may reduce the metabolic burden of the chloroplast, enhance

the functional integration and regulation of plastid- and nucleus-encoded proteins, or even

escape from the accumulation of deleterious mutations in asexual organellar genomes

(Muller's Ratchet; Millen et al., 2001). Studies have shown that *infA* plays a crucial role in

adaptation to cold shocks (*Escherichia coli*) or to different temperatures (slipper orchids;

Giangrossi et al., 2007; Hu et al., 2022). The transfers of *infA* into the nuclear genome of *R.*

cassubicifolius could thus, even more, promote the adaptation of sexual diploids, and

particularly of allopolyploid apomicts distributed from Arctic to Mediterranean climates

(Karbstein et al., 2020b, 2021, 2022; Hodač & Karbstein et al., 2023).

In addition, the gene *ycf15* appears to have been lost early in the evolution of

Ranunculaceae (e.g., *Hydrastideae*, *Caltheae*), and regained several times in the plastome also

within *Ranunculus* (12x in *Ranunculus)* including *R. cassubicifolius*. The gene *ycf15* probably

originated at the roots of the angiosperms and has been pseudogenized or lost many times,

and compared to other (pseudo)genes like *infA* or *rpl32*, no transfer to the nuclear genome

was observed across many angiosperm lineages (Shi et al., 2013). For the genus *Ranunculus*,

the complete loss of this gene had been assumed (Shi et al., 2013) but the recent study by Kim

et al. (2023) for *R. austro-oreganus* and *R. occidentalis,* and the results presented here

contradict this assumption. Although the gene is only 54 bp long in *R. cassubicifolius* and

thus apparently degraded (e.g., compare to 198 bp in *R. austro-oreganus*), and seemingly

without essential function to photosynthesis and survival of plants (but see its application for

species delimitation; Shi et al., 2013; Li et al., 2021), we found two copies of *ycf15* in the IRA

and IRB regions with regular ATG start and TGA stop codons, suggesting that it is functional

in *R. cassubicifolius*. However, the functions of *ycf15* and the co-transcribed large plastid

gene *ycf2* are still unclear, although *ycf2* and *accD* influence plastid structure and
competitiveness and thus may play a role in uniparental vs. biparental inheritance of plants
that is known to shape the ability of hybridization (Drescher et al., 2000; Shi et al., 2013; Liu
et al., 2017; Sobanski et al., 2019; Postel and Touzet, 2020). Its role in the hybridization of
sexual progenitors that produced the hundreds of allopolyploid *R. auricomus* nothotaxa
(Karbstein et al., 2022) needs further investigation.

Other genes frequently known to be transferred into the nuclear genome like *accD,*
yc1, or *ycf2* remain in the plastomes of Ranunculaceae. Noteworthy, we observed that
annotations of (pseudo)genes are sometimes missing (e.g., *clpp*, *psbG*, *rpl22*, or *rpl32*) or
renamed (*pbsN* vs. *pbf1*), which complicates plastome feature comparisons and requires
additional manual work such as mapping gene sequences from closely related species to the
plastome sequence of interest. Therefore, we advocate for a more careful annotation of
plastome sequences through cross-species alignments of homologous genes.

Figure 2. (a) The annotated plastome sequence of *Ranunculus cassubicifolius* (LH040). The
plastome assembly is a 156,233 bp-long single circular contig. It contains 85 (78 unique)
protein-coding genes, 37 (30 unique) tRNAs, and 8 (4 unique) rRNAs, as shown in the outer
circle. Genes and t/rRNAs on the inside and outside of the circle are transcribed in the
clockwise and counterclockwise directions, respectively (gene codon usage bias in brackets).
The first inner circle represents GC content (%), and the second inner circle the plastome
structure, i.e., a small single-copy region (SSC) and a large single-copy region (LSC)
separated by two inverted repeat (IRA/B) regions. The third inner circle reflects repeat
content, i.e., microsatellites (1-6 bp per repeat), tandem repeats (>6 bp per repeat), and
dispersed repeats (transposable elements or satellite DNA; reports on FigShare). See Shi et al.
(2019) and http://www.1kmpg.cn/cpgview/ for more details.

Mitogenome features and gene evolution within *Ranunculus* and Ranunculaceae.

Based on the Illumina-PacBio hybrid assembly, we assembled a mitochondrial *R.*

cassubicifolius genome sequence of 1,183,731 bp. It contains 40 (40 unique) protein-coding

genes, 77 (18 unique) tRNAs, and 7 (3 unique) rRNAs, resulting in a coding content of 8.0%.

This is consistent with observations in angiosperms (Skippington et al., 2015; Dong et al.,

2018; Park and Park, 2020; Jiang et al., 2023). In comparison with *Hepatica maxima* (the

closest available reference within Ranunculaceae, see Fig. S4a; Zhai et al., 2019), both

mitogenomes show similar numbers of unique protein-coding genes (40/39), t-RNAs (18/18),

and rRNAs (3/3), but the *R. cassubicifolius* mitogenome is slightly longer (+61,185 bp). In

Ranunculaceae, variation in mitogenome size is high, ranging from 0.207 Mbp in *Isopyrum*

anemonoides to 1.184 Mbp in our study species *R. cassubicifolius*. In angiosperms,

mitogenomes can even range from 66 kbp to 11.7 Mbp (Skippington et al., 2015; Putintseva

et al., 2020). The mitogenome of *R. cassubicifolius* is therefore relatively small, but one of the

largest known for Ranunculaceae.

Size differences in the mitogenomes of Ranunculaceae can only be explained to a

limited extent by variation in repeat content. In *R. cassubicifolius*, many short (187) and long

(168) tandem repeats and a few dispersed repeats were detected, leading to 2.1% repeat

content. This is lower compared to *H. maxima* (6.9%) and substantially lower (28x) than the

smallest known mitogenome of *I. anemonoides* (0.207 Mbp, 56%). In contrast to plastomes

(and animal mitogenomes), plant mitogenomes are recombinationally active, can exist as

populations of alternative structures even within individuals, and are considered to be

organized in noncircular molecules (Sloan, 2013; Møller et al., 2021). Sequence differences

despite similar gene content are thus mainly attributable to sequence recombination and

rearrangements in non-coding regions, supported by whole genome alignment results in the

present study (Fig. S5a). Significant structural differences were also found between the

PacBio and ONT mitogenome assemblies (Fig. S5b), which might stem from the fact that

different individuals were sequenced rather than from the use of different sequencing

technologies (see Table 1). In addition, despite reads bridging contig boundaries, assembly

results may encourage the presence of at least 3 chromosomes instead of a single circular

molecule in the active cell state (Skippington et al., 2015; Feng et al., 2021).

The coalescent-based mitogene phylogeny of Ranunculaceae was well-supported

(FBP>85%; Fig. 4b), marking the first comprehensive one for the family and one of the first

for plants (e.g., Lin et al., 2022). Like in the plastome tree, low branch support occurs

especially within fast radiating, species-rich genera such as *Pulsatilla* (*Anemoneae*).

Branching of tribes is also widely consistent with the here presented and previously described

plastome phylogenies (Fig. 4a; Wang et al., 2009; Zhai et al., 2019), that is, *Thalictroideae*,

Delphinieae, *Ranunculeae*, and *Anemoneae*. Regarding mitochondrial gene evolution within

Ranunculaceae and particularly *Ranunculus*, we observed that some genes (*rps1*, *rps2*, *rps10*,

rps11, *rps12*, *rps14*, *rps19*, or *sdh3*) were frequently lost (Fig. 4b; see raw feature table on

FigShare). In flowering plants, gene loss and transfer to the nuclear genome have been

frequently observed for *rps*, *sdh3,* and *sdh4* (Adams et al., 2000, 2001; Bi et al., 2020).

In our study, the mitochondrial genes *rps10* and *rps14* appeared to be lost from

Ranunculeae to *Anemoneae* and putatively transferred to the nuclear genome (no reference

genome sequence available). The *rps* genes encode small ribosomal subunit proteins and are

thus essential for mitochondrial ribosome function, as well as vegetative and reproductive

plant development (Adams et al., 2000, 2001; Robles and Quesada, 2017). These two genes

were shown to be under evolutionary positive selection in certain plant groups (e.g., *rps1*,

rps10, rps14 in legumes: Bi et al., 2020; *rps13* in rosids: Liu and Adams, 2008). In addition,

rps1, *rps2*, *rps11*, and *rps19* were not found in *I. anemonoides*; *rps2* and *rps12* in *Aconitum*

kusnezoffii; and *rps2*, *rps11*, and *rps14* in *A. carmichaelii*. In plant evolution, the majority of

mitochondrial genes have already been transferred to the nuclear genome. This enhances

regulatory host control of these genes and integration with nuclear-encoded mitochondrial

proteins, and has important consequences for ATP synthesis, metabolism, and abiotic cold or

heat and biotic stress responses (Adams et al., 2000; Bonen and Calixte, 2006; Woodson and

Chory, 2008; Saha et al., 2017; Fakih et al., 2023).

Interestingly, only in *R. cassubicifolius,* all ribosomal genes are present. In contrast to

our obligate self-incompatible species of interest (Hörandl, 2008; Karbstein et al., 2020a), all

other genera examined, such as *Anemone*, *Pulsatilla*, and partially *Aconitum*, are known to be

largely self-compatible (Lindell, 1998; Zhigang et al., 2006; Liao et al., 2009). Selfing (and

clonal reproduction), in the absence of deleterious mutation accumulation in the mitogenome,

is known to increase the transfer of mitochondrial genes to the nuclear genome and provides

an adaptive advantage due to fixation of co-adapted gene combinations (Brandvain and Wade,

2009). Therefore, it would be worth investigating whether self-compatible, apomictic

polyploids retain as many *rps* genes in their mitogenomes as their sexual diploid *R. auricomus*

progenitors, such as *R. cassubicifolius*. However, also the putative transfer of all *rps10* and

rps14 genes to the nuclear genome within the fast radiating, species-rich clade Anemoneae

(e.g., only *Anemone* and *Clematis* with >100 and >400 accepted species on gbif.org) needs

further investigation. In Ranunculaceae, the lost and putatively transferred *rps* genes are

known to be regulated under temperature heat and cold stress (Alafari and Abd-Elgawad,
2021; Fakih et al., 2023), and in the investigated taxa it could play a crucial role in adaptation
to a broad range of subtropical to subarctic and lowland to mountainous habitats in central
and (south)eastern Asia. Similar to its common forest understory ancestor, *R. cassubicifolius*
grows in temperate habitats in humid forests or along stream-sides (Tomasello et al., 2020;
Karbstein et al., 2021), facing low selective environmental pressure, which, along with its
reproductive traits, may explain the retention of all *rps* genes in its mitogenome sequence.
Whether this pattern is consistent across the 600 *Ranunculus* species with worldwide
distribution in various environments, needs further investigation.

The protein-coding gene *sdh3* was detected neither in *R. cassubicifolius* nor *I.*
anemonoides (see raw feature table on FigShare). It encodes the subunit 3 of the succinate
dehydrogenase and plays a role in electron transfer in complex II of the respiratory chain, and
is thus important for energy production (Gebert et al., 2011). Ancient and recent events of
*sdh*3 loss with and without transfer to the nuclear genome have been observed in several
flowering plant lineages (>40x; (Adams et al., 2001; Choi et al., 2019; Huang et al., 2019).
Surprisingly, transfer of *sdh3* to the nuclear genome could not be confirmed in *R.*
cassubicifolius, and for *I. anemonoides* no reference genome sequence is available. Loss of
sdh3 might reflect optimization of cellular efficiency, and that the gene is not essential in the
investigated plant lineages.
770
771

Figure 3. The mitogenome sequence of *Ranunculus cassubicifolius* (LH040). The mitogenome is 1,183,731 bp in size represented in a single, circular contig, and contains 40 (40 unique) protein-coding genes, 77 (unique 18) tRNAs, and 7 (unique 3) rRNAs, as shown in the outer circle. Genes and t/rRNAs on the inside and outside of the circle are transcribed in the clockwise and counterclockwise directions, respectively. The first inner circle represents GC content (%). The second inner circle reflects repeat content, i.e., short tandem repeats/microsatellites (1-6 bp per repeat), long tandem repeats (>6 bp per repeat), and dispersed repeats detected by RepeatModeler2 (reports on FigShare). See (Zhang et al., 2024) and http://www.1kmpg.cn/pmgmap for more details.

(a)

Anemoneae (118)
Adonideae (6)

Ranunculeae (29)

-infA
99
R. bungei SE-AS OR625573
R. bungei SE-AS NC_060613
R. bungei SE-AS MK253468
R. bungei SE-AS OR625572
R. trichophyllus NH OR625582
-ycf15
99
R. trichophyllus NH OR625577
-infA
R. mongolicus C-AS OR625576
-infA
R. sceleratus NH MK253452
99
99
R. cassubicifolius CE-EU NC_077490
R. monophyllus EU+AS OR625578
84
R. polyrhizos EU+AS OR625579
-infA
R. brotherusii SE-AS OR625580
-ycf15
R. ternatus SE-AS NC_081908

clade III
clade II

R. silerifolius E-AS ON462450
R. cantoniensis E-AS NC_045920
-infA
-ycf15
R. chinensis E-AS NC_079824
R. macranthus N-AM NC_008796
97
R. occidentalis NW-AM NC_031651
R. austro-oreganus NW-AM KX639503
92
-ycf15 -ycf15
R. reptans NH NC_036977
95
-infA
R. flammula NH KY562595
-ycf15
R. japonicus SE-AS MZ169045
R. repens EU+AS NC_036976

clade I

C. falcata EU+AS MK253464
C. testiculata EU+AS NC_069854
92
C. orthoceras EU+AS PP155435
-infA
M. minimus NH PP155438
M. apetalus NW-AM PP155437
99
-infA
F. verna EU+AS PP155436
O. kamchatica NH NC_041538

Helleboreae (2)
Callianthemeae (2)
-rpl32
Delphinieae (56)
Nigelleae (2)
Cimicifugeae (14)
-ycf15
Asteropyreae (2)
Caltheae (1)
Thalictroideae (33)
-ycf15
Adonideae (16)
Coptideae (6)
Glaucidieae (1)
Hydrastideae (1)

Papaveraceae (3)
Berberidaceae (3)

4.0

Figure 4. Phylogenetic relationships and gene evolution within Ranunculaceae. (a)

Maximum-likelihood phylogeny based on 308 plastome sequences (295 taxa) and the min90

alignment (126 kbp) of the plant family Ranunculaceae. Branches of tribes except for

Ranunculeae were collapsed, and gained or lost genes are shown in different colored gray

bars along the tree. All branches received full (1) transfer bootstrap expectation (TBE) values,

unless otherwise shown. See also Quartet Sampling (QS) metrics in Fig. S2, the complete

phylogeny in Fig. S4a,b, and more technical details in Text S3. (b) Coalescent-based

phylogeny of 10 mitogenome sequences and 42 genes of Ranunculaceae. All branches

received full (100%) Felsenstein Bootstrap Proportions (FBP) values, unless otherwise

shown. All NCBI species names were checked by gbif.org, and synonyms were replaced by

accepted names (synonyms in brackets). The color of clades corresponds to tribe/subfamily

names (tribal classification follows Wang et al. (2016) and Zhai et al. (2019). *Ranunculus*

cassubicifolius is highlighted in bold with an arrow. Abbreviations for native distribution

ranges of taxa: EU = Europe, AS = Asia, AM = America, NH = Northern Hemisphere, C =

Central, E = Eastern, N= Northern, S = Southern, and W = Western.

Comparison of nuclear genome assembly strategies.

800
This study presents a systematic and comprehensive comparison of assembly
strategies for large non-model plant genomes based on short- and long-read sequencing data.
Related studies have mainly focused on model plants with high coverage and few tested tools
(e.g., Mascher et al., 2021), or non-model plants with small-sized genomes (e.g., Murigneux
et al., 2020). Here, the 18 genome assembly strategies yielded substantial differences
concerning completeness, continuity, and quality (Table 1, Fig. 5). SUPERNOVA and
SPAdes assemblies based on Illumina reads delivered mixed results in terms of completeness
(86/17%), with too many (6.92/1.23 million contigs), short (N_{50}: 2/3 kbp) contigs. BUSCO
genes were also highly fragmented (F: 36/28%), with low levels of complete single-copy (S:
60/20%) and duplicated (D: 11/2%) genes, and many missing genes (M: 34/10%). This is in
line with studies showing highly fragmented, gappy, low-quality assemblies of large genomes
when only short Illumina reads are available (Gatter et al., 2021; Rhie et al., 2021; Zimin and
Salzberg, 2022; Hotaling et al., 2023). Especially in *R. cassubicifolius* with 86% repetitive
regions, short reads with moderate coverage cannot resolve these long and complex patterns.

Hybrid assemblies of SPAdes, MaSuRCA, and WENGAN based on Illumina and
ONT or PacBio reads performed slightly better in terms of contiguity, with fewer but larger
contigs (368-1.18/0.530-6.57 million contigs, with N_{50}: 5-21/5-6 kbp), but not in terms of
genome completeness (0.16-13/14-38%) and BUSCO completeness as many genes were often
missing (M: 7.7-99.6 / 7.5-90%, F: 0.2-24 / 9-23 %, D: 0-2.4 / 0-2.4%, S: 0.2-66 / 1-67%).
SPAdes produced the best assemblies in terms of completeness, contig size, and BUSCO
scores, despite being developed and tested for very small microbial genomes (<0.1 Gbp;
Prjibelski et al., 2020). MaSuRCA and WENGAN (MuCHSALSA, results not shown)
performed worse. Many hybrid assemblers are considered by developers to be particularly
useful under low long-read coverage (<10-15x) or large/complex genomes (e.g, Zimin et al.,
2017; Di Genova et al., 2021; Gatter et al., 2021). However, this cannot be confirmed in our
study. These tools have been tested on model species like *Arabidopsis thaliana* with small
genomes (0.135 Gbp) and very low levels of TE (ca. 21%, reviewed in Quesneville, 2020) or
in human data with similar genome size (3.2 Gbp) to *R. cassubicifolius* but also lower levels
of repeat content (ca. 50%, reviewed in Liao et al., 2023), or they need Illumina data at higher
coverage (>30-40x) than we provide to build initial robust reads for subsequent long read
mapping, which may explain their poorer performance here.

Compared to already discussed strategies, the ONT or PacBio-based assemblies
conducted with Canu, Flye, and Hifiasm were generally superior in terms of completeness

(25-158%), contig number (1,453- 65,232), median contig size (N_{50}: 0.037-20.0 Mbp), and

quality as the majority of BUSCO genes were found (M: 0.2-29%, F: 0.3-7.1%, D: 7.9-88%,

S: 11-76 %). This is also consistent with the recent review by Hotaling et al. (2023), which

concludes that long reads allow for more complete and contiguous genome assemblies of

higher quality, particularly for large repeat-rich genomes. However, we also observed that

PacBio delivered better assemblies than ONT using Canu and Flye. PacBio produced the

longer (N_{50}: 617/853 vs. 37-103 kbp) and fewer contigs (16,262/12,506 vs. 30,612/64,618).

The lower base accuracy of ONT compared to PacBio sequence data (97 vs. 98%), despite a

slightly larger median sequence length (23 vs. 18 kbp), can explain this observation (Hotaling

et al., 2023). It is important to note that the ONT read N_{50} length of the runs in this study was

ca. 20 kbp, while other studies have achieved runs with more than twice this value when using

fresh libraries based on different DNA extraction protocols (Hakim et al., 2024; Horz et al.,

2024; Nowak et al., 2024). Using the latest ONT flow cell generation and fresh DNA extracts

from different kits would probably deliver much longer, higher quality reads and thus

improved assemblies (see Text S5 for further discussion). However, the PacBio-based

assemblies generated by Canu and Flye were far too large considering the expected haploid

genome size (158/153 vs. 25/112%) and characterized by too many duplicated BUSCO genes

(70/88 vs. 8/14%). This suggests uncollapsed heterozygous regions in the assembly.

However, Hifiasm – both using the full PacBio 25x coverage or the 16x downsampled

set – outperformed Canu and Flye by being closest to expected genome size (101/113 vs.

158/153%), median contig length (2,890/20,000 vs. 617/853 kbp) and contig number

(3449/1453 vs. 16,262/12,506), and runtime (2-3 vs. 7-14 days), but all delivered similar

BUSCO gene completeness (99/99% vs. 99/99%). This agrees with recent benchmark

analyses on large and complex eukaryotic and synthetic genomes (Cheng et al., 2021; Yu et

al., 2024). Interestingly, our polishing strategy slightly improved assembly in terms of

missing and fragmented BUSCO scores (mean M: 8.41 vs. 3.67%, and F: 2.51 vs. 1.16%),

and contig number of >1 Mbp (mean 21.9 vs. 19.8 Mbp). Percentage of single-copy BUSCO

genes improved significantly only for ONT-Canu assemblies (56 vs. 66%). After polishing,

genome completeness (mean 118 vs. 117%) and N_{50} contig length (mean 4.13 vs. 4.13 Mbp)

remained constant. We generally observed that polishing improved BUSCO statistics more

for lower-quality assemblies rather than for good assemblies. Thus, sequencing errors,

misassemblies, and duplications can be improved by polishing, but existing contigs are not

significantly enlarged. The need for polishing thus depends on the obtained assembly quality,

and is particularly attractive when working under low coverage scenarios (<30x). In addition,

25x compared to downsampled 16x coverage PacBio data delivered assemblies of
substantially fewer and larger contigs (mean 26,873 vs. 21,268, and 0.832 vs. 1.83 Mbp),
more single-copy BUSCO genes (mean 37.6 vs. 31.7%) and less duplicated (mean 40 vs.
46%), fragmented (mean 5.23 vs. 5.63%) and missing ones (mean 8.4 vs. 16%), but had
similar genome completeness (mean 105 vs 106%) across all tested hybrid and long-read
strategies (Tables 1, S4). Coverage much below 25x is therefore not recommended for either
strategy.

The PacBio-based Hifiasm assembly polished twice by PacBio and once more by
Illumina reads delivered the best results among all tested strategies (100% completeness,
1,319 contigs, 20 Mbp N_{50}, and 99% complete BUSCO genes). This represents the best
assembly strategy for the large and complex *R. cassubicifolius* nuclear genome with low to
moderate sequencing depths, and agrees with findings in smaller non-model plant assemblies
(e.g., Dmitriev et al., 2021).

The nuclear genome features.
Based on the best PacBio + Hi-C assembly, we obtained a nuclear *R. cassubicifolius*
genome sequence of 2,691,150,139 bp. Approximately 98.4% (2.65 Gbp) of the scaffolds
were anchored to form 8 pseudochromosomes based on Hi-C data. This high-quality,
chromosome-level assembly has an N_{50} of 355 Mbp, with the longest pseudochromosome
spanning 447 Mbp (Fig. 6, Table 2). BUSCO analysis revealed that 94.1% (78.8% single
copy, 15.3% duplicated) of the BUSCO embyophyta_odb10 genes were present, indicating
the completeness of the assembly. The BUSCO score is slightly lower than before Hi-C
scaffolding (99%). The Hi-C scaffolding procedure automatically removes contigs below
1,500 bp, but all contigs prior to scaffolding were far above this threshold, so contig removal
does not explain the reduced BUSCO gene score. More likely are misassemblies or incorrect
gene annotations based on the original assembly. The retained genome sequences excel the
recommended minimum standard of the Earth Biogenome Project in terms of continuity
(scaffolds >10 Mbp), functional completeness (>90% BUSCO genes), and chromosome status
(>80% contigs assigned to chromosomes; https://www.earthbiogenome.org/report-on-
assembly-standards, Lawniczak et al., 2022).

The genome sequence contains 35,482 (31,322 unique) protein-coding genes with an
average gene length of 3,548 bp and an average coding sequence (CDS) length of 1,169 bp.
The gene annotation showed a high level of BUSCO completeness (94.5% using
embryophyta_odb10). We predicted 984 tRNA, 99 spliceosomal RNA (snRNA), and 6,448

rRNA (3,013 5S rRNA, 1,077 5.8S rRNA, 1,154 18S rRNA, and 1,204 28S rRNAs) gene

models. Further annotation details are available on Github. In comparison with *Aquilegia*

kansuensis (293 Mbp; Xie et al., 2020), *Coptis chinensis* (959 Mbp; Chen et al., 2021), and

Coptis teeta (932 Mbp; Wang et al., 2024) as the only available references within

Ranunculaceae, the *R. cassubicifolius* nuclear genome is up to 10-fold larger in size (+2.42

Gbp / +1.75 Gbp /+1.78 Gbp) but has similar numbers of chromosomes (8 vs. 7/9/9).

Ranunculus, however, has mostly the long, strongly winding chromosome type (R-type) of

the family, while *Aquilegia* and *Coptis* have small, simply curved chromosome types (T-type

like *Thalictrum*) (Tamura, 1993). Within *Ranunculus*, our target species shows an ancestral

karyotype with large chromosomes (4 metacentric and 4 submetacentric, Baltisberger and

Hörandl, 2016). Comparing all available genome sequences, *Coptis teeta* shows the most

genes (43,979 in Hap1 /46,311 in Hap2), followed by *C. chinensis* (34,109), *R. cassubicifolius*

(31,322), and *A. kansuensis* (25,571), suggesting that despite different gene annotation

pipelines, gene content and genome size are probably decoupled (Michael, 2014; Li et al.,

2017). In angiosperms, haploid genome size varies ca. 2400-fold from 0.063 to 149 Gbp

(median = 0.6 / mean = 5.7 Gbp, Dodsworth et al., 2015). The nuclear genome of *R.*

cassubicifolius is larger than most flowering plants, and one of the largest assembled yet for

Ranunculaceae. Within diploid *Ranunculus* species, haploid genome sizes measured with

karyological methods were estimated to range from 1.86 to 8.43 Gbp (Leitch et al., 2019). *R.*

cassubicifolius thus represents a typical, middle-sized genome reference for the genus.

The low proportion of duplicated BUSCO genes (15%) hints at no recent

polyploidization event. In ancestral angiosperms, a baseline of 12,000 – 14,000 protein-

coding genes is postulated (Sterck et al., 2007). In Liu et al. (2021) and He et al. (2022),

ancient whole genome duplication (WGD) events were detected in Beberidaceae and

Papaveraceae as close outgroups to Ranunculaceae, and also shared and lineage-specific ones

(e.g., *Coptis chinensis* with ca. 39,500 genes, Coptideae) at the base of Ranunculaceae.

Ranunculus is nested within a far later branching clade (Ranunculeae) within Ranunculaceae,

but *R. cassubicifolius* shows only slightly higher gene number. Either no further shared WGD

event occurred and gene number remained quite conserved in the several million years of

Ranunculaceae evolution, or re-diploidization masked another, more recent WGD event.

Interestingly, the *A. kansuensis* (Thalictroideae) genome with only ca. 25,000 genes is placed

shortly after *Coptis* and before *Ranunculus* in the Ranunculaceae phylogeny. Whether this re-

diploidisation represents a lineage-specific or general trend followed by subsequent WGD

events needs to be tested using higher quality genome assemblies and additional analysis tools

(e.g., Sun et al., 2022b).

Almost 86% of the *R. cassubicifolius* assembly consists of repetitive elements (Table

4), among the highest in flowering plants (85% in *Zea mays*; Schnable et al., 2009; Wang et

al., 2021). Long terminal repeat (LTR) retrotransposons are most abundant, making up to ca.

77% of the genome. Additionally, DNA transposons, and long interspersed nuclear elements

(LINEs) and short interspersed nuclear elements (SINEs) retrotransposons account for 5.7%,

0.81%, and 0.2% of the genome, respectively. In angiosperms, genome size correlates well

with repetitive element content and particularly with LTR-retrotransposon content but not

with polyploidization (Li et al., 2017; Wang et al., 2021). Size differences between

Ranunculaceae genomes can be largely explained by variation in LTR-retrotransposon

contents. In *R. cassubicifolius*, LTR-RTs account for 77.6% of the genome whereas they

constitute ca. 36% of the genome of *Aquilegia kansuensis* (Xie et al., 2020). In *Coptis teeta,*

LTR-RTs contribute to ca. 36% of genome (Wang et al., 2024), and in *Coptis chinensis*, it is

50% (Chen et al., 2021). Interestingly, LTR retrotransposon bursts have recently been shown

to have occurred during the last three million years (Pleistocene) characterized by severe

climatic variability, potentially promoting adaptation through novel genomic functions and

altered expression patterns (Wang et al., 2021). The progenitor species of the *R. auricomus*

complex also speciated by vicariance processes due to range contraction and expansion during

cold and warm phases 830-580 kya in Europe (Tomasello et al., 2020). Therefore, climatic

variations and the need to adapt to them may be related to the high LTR content in *R.*

cassubicifolius.

Given the overall high completeness of our assembly (94.1% of BUSCO genes), we

think it could represent a very important resource for exploring the genetic background of

reproductive mode shifts from sex to apomixis in *R. cassubicifolius* and beyond. The

apomixis candidate genes under selection in apomictic hybrids (*ASY1, XRI1*, and *APC1*)

recognized by Paetzold et al., 2022) based on RNA-seq data were also detected in the genome

of *R. cassubicifolius* at pseudochromosomes 2 and 5. However, the apomixis candidate gene

MSP1 was not found and needs further investigation. The annotated genes will be also an

important resource for comprehensive analysis of loci under selection in the genus (e.g., via

dN/dS ratios; (Pellino et al., 2013), which could give insights into general genome evolution,

adaptation, and speciation.

Table 1. QUAST statistics for different genome assembly strategies of *R. cassubicifolius* (LH040) (see also reports on Figshare). SUPERNOVA, and partially SPAdes, MaSuRCA, and WENGAN used Illumina short-reads ('Illumina'). Canu, Flye, and Hifiasm, and partially SPAdes, MaSuRCA, and WENGAN used Oxford Nanopore (ONT) or PacBio long-reads. Coverage is given per read type. Assembly polishing included the following steps: polishing via filtered ONT reads using Racon and Medaka / via filtered PacBio reads using Racon and polishCLR, respectively, and via filtered Illumina reads using POLCA (MaSuRCA pipeline). The best assembly in terms of completeness and contig length are highlighted in bold. N_{50} describes the shortest contig length required to summarize at least half (50%) of the bases of the entire assembly; L_{50} is defined as the minimum number of contigs that encompass half (50%) of the bases of the entire assembly. See Table S3 for comparisons with not downsampled PacBio reads. - means no results produced.

Metric	SUPERNOVA (Illumina22x)	SPAdes (Illumina22x)	SPAdes (Illumina22x + ONT16x/ PacBio16x)	MaSuRCA (Illumina22x + ONT16x/ PacBio16x)	WENGAN (Illumina22x + ONT16x/ PacBio16x)	Canu (ONT16x/ PacBio16x)	Canu (ONT16x/ PacBio16x) +3x Polish	Flye (ONT16x/ PacBio16x)	Flye (ONT16x/ PacBio16x) +3x Polish	HIFIASM (PacBio25x/ PacBio16x)	HIFIASM (PacBio25x) +3x Polish
Total assembly length, Mbp	2,76 (86%)	0,557 (17%)	0.408 (13%)/ 0.464 (14%)	0.076 (2.4%)/ -	0.005 (0.16%)/ 1.21 (38%)	0.80 (25%)/ 5.05 (158%)	0.81 (25%)/ 5.04 (157%)	3.58 (112%)/ 4.90 (153%)	3.45 (108%)/ 4.90 (153%)	3.22 (101%)/ 3.62 (113%)	**3.21** **(100%)**
No. contigs	6,920,808	1,231,317	1,184,036/ 529,574	38,650/ -	368/ 6,565,544	30,612/ 16,262	32,959/ 15,711	65,232/ 12,514	54,310/ 12,356	1,453/ 3,449	**1,319**
No. contigs > 1 Mbp	562,716	122,407	110,594/ 108,266	12,799/ -	368/ 5,046	30,612/ 16,262	31,072/ 15,711	64,618/ 12,506	53,572/ 12,356	1,453/ 3,449	**1,319**
Largest contig, Mbp	0.039	0.069	0.096/ 0.112	0.145/ -	0.095/ 0.009	0.540/ 14.3	0.542/ 14.3	0.976/ 6.57	0.974/ 6.57	84/ 32	**84**
GC, %	40	39	39/ 39	40/ -	40 / 39	40/ 42	40/ 42	43/ 42	43/ 42	42/ 42	**42**
N_{50}, Mbp	0.002	0.003	0.005/ 0.005	0.010/ -	0.021/ 0.006	0.037/ 0.617	0.038/ 0.618	0.103/ 0.853	0.104/ 0.852	20.0/ 2.89	**20.0**
L_{50}	259,069	29,900	23,910/ 23,533	1,507/ -	76/ 28,344	6,935/ 2,374	6,937/ 2,366	10,305/ 1728	9,889/ 1727	45/ 350	**45**

977 **Figure 5.** BUSCO assessments for different genome assembly strategies of *Ranunculus cassubicifolius* (LH040). SUPERNOVA, and partially
978 SPAdes, MaSuRCA, and WENGAN used Illumina short-reads (SR). Canu, Flye, and HIFIASM, and partially SPAdes, MaSuRCA, and
979 WENGAN used ONT/PacBio long-reads alone (LR) or in combination with short reads (Hybrid). The best assembly in terms of complete
980 BUSCO genes (C+D) is highlighted with a black square (i.e., 'HIFIASM (PacBio) + Polishing'). Analyses are based on 1,614 BUSCO groups
981 searched. See the Materials & Methods section, and Table 1 and Table S4 (Excel Sheet) for more details.
982
983

a: GC content

b: Gene density

Repeat content

c: DNA TE density

d: LTR TE density

e: LTR-classification:
  Unknown
  Gypsy
  Copia

f: Tandem repeats density

g: Chromosome relationships

Figure 6. Circos plot of the chromosome-scaffolded nuclear genome of *Ranunculus cassubicifolius* (LH040). The genome is 2.69 Gbp in size,

has 8 pseudochromosomes and 31,322 annotated genes. Illustrated are (from the outer to the inner circle): Chromosome number, chromosome

size in Mbp, GC content (a), protein-coding gene density (b), DNA transposon element density (c), Long terminal repeat (LTR) retrotransposon

density (d), percentage of unknown, Gypsy, and copia LTR-type TEs (e), tandem repeat density (f), and major interchromosomal synteny (g).

Table 2. Assembly statistics of the annotated plastome, mitogenome, and chromosome-scaffolded nuclear genome of *Ranunculus cassubicifolius*
(LH040). Statistics were calculated with Geneious and R. See Table 4 for a detailed repeat content of the nuclear genome, Materials and
Methods, and Results section for more details.

Metric	Plastome	Mitogenome	High-Quality Nuclear Genome
Assembly Strategy	Illumina SR (GetOrganelle)	Best strategy: Illumina (filtered by GetOrganelle) assembled with previously mapped HiFi (MaSuRCA)	Best strategy: PacBio (HIFIASM) + 3x Polishing [HiFi (Racon), HiFi (polishCLR), Illumina (POLCA)] + Hi-C chromosome-level scaffolding
Assembled Genome Size	156,233 bp	1,183,731 bp	2,691,150,139 bp
Number of Contigs	1 (circular)	1 (circular)	8 (pseudochromosomes)
N_{50} of Contigs	-	-	355 Mbp
Largest Contig	156,233 bp	1,183,731 bp	447 Mbp
BUSCO Genes (S+D/S)	-	-	94.1%/78.8%
Protein Coding Genes	85 (78 unique)	40 (40 unique)	35,482 (31,322 unique)
tRNA Genes	37 (28 unique)	77 (unique 18)	984
rRNA Genes	8 (4 unique)	7 (unique 3)	6,448
GC-Content (%)	37.9%	43.5%	42.1%
Repeat Content (%)	2.15%	2.13%	86.0%
Gene Content (%)	53.2%	7.99%	4.1%

995   **Table 4.** Repeat content of the chromosome-scaffolded nuclear genome of *Ranunculus*

996   *cassubicifolius* (LH040). Long Interspersed Nuclear Elements (LINEs), Short Interspersed

997   Nuclear Elements (SINEs), Long Terminal Repeats (LTRs), and Terminal Inverted Repeats

998   (TIR). A detailed explanation of repeat classes can be found in Wicker et al. (2007).

999   Annotation reports are deposited in Github.

	Total Sequences	Total Length	
	320	2,691,150,139	
Class	**Count**	**bpMasked**	**%masked**
Class I (retrotransposons)			
LINE			
L1	31,842	21,678,526	0.81%
LTR			
Copia	147,394	80,810,425	3.00%
Gypsy	1,105,626	654,391,560	24.32%
unknown	2,075,881	1,353,935,235	50.32%
SINE			
tRNA	309	114,573	0.00%
unknown	347	444,295	0.02%%
Class II (DNA transposons)			
TIR			
CACTA	85,851	32,242,719	1.20%
Mutator	134,007	38,209,632	1.42%
PIF_Harbinger	86,910	30,275,684	1.13%
Tc1_Mariner	32,184	12,038,336	0.45%
hAT	133,119	40,436,252	1.50%
nonTIR			
helitron	123,174	38,735,549	1.44%
rDNA			
45S	191	42,204	0.00%
Repeat Fragment	44,541	11,822,897	0.44%
Satellite DNA	186	83,887	0.00%
total interspersed	4,001,562	2,315,261,774	86.04%
Total	**4,001,562**	**2,315,261,774**	**86.04%**

1000

Conclusions

In this study, we assembled the complete plastome, mitogenome, and large nuclear genome sequences of *R. cassubicifolius,* a diploid sexual progenitor species within the *Ranunculus auricomus* complex, using different combinations of Illumina short-reads and ONT and PacBio long-reads under low to medium sequencing depth scenarios. We generated a plastid and the first intricate mitogenome sequences for *Ranunculus* based on Illumina and Illumina + PacBio data, respectively. An updated plastome phylogeny and the first mitogenome phylogeny of Ranunculaceae indicated frequent gene losses (e.g., *infA*, *ycf15*, and *rps* genes), potentially in the context of environmental adaptation and reproductive features. To assemble the large and complex nuclear genome, we performed 18 assemblies using short-, long-, and hybrid-read strategies, followed by gap filling and/or polishing with various state-of-the-art bioinformatic tools. The best result in terms of contiguity and completeness was achieved by Hifiasm using PacBio data, polished twice with filtered PacBio reads (Racon, polishCLR) and once with Illumina reads (POLCA), representing the first high-quality, chromosome-scaffolded genome sequence within *Ranunculus* and the fourth within Ranunculaceae (although the two currently available genome sequences on NCBI are below 1 Gbp genome size; Xie et al., 2020b; Chen et al., 2021). The large *R. cassubicifolius* genome with 2.69 Gb size, containing 31,322 annotated genes and 86% repetitive elements, represents a typical medium-sized plant genome and is probably representative for *Ranunculus*.

The obtained results and in particular the new nuclear genome assembly are useful for (i) improving read mapping (see Text S4), orthologous gene/locus assembly, and allele phasing of subgenomic reduced-representation sequencing (RRS) (e.g., RAD-Seq, target enrichment), transcriptome (RNA-seq), and whole genome (re)sequencing data (e.g., WGS, WGR), (ii) as a basis for understanding cyto-nuclear compatibility for both plastid-nuclear and mitochondrial-nuclear interactions; (iii) as diploid reference for the analysis of the more complex allo- and autopolyploid nuclear genomes within the *Ranunculus,* (iii) as reference for further studies of genome evolution, loci under selection, adaptation, and speciation, as well as (iv) for improving functional analyses (e.g., gene expression studies) related to apomixis, adaptation, and many other traits.

Data Availability. The basic data supporting the findings are available within the manuscript and Supporting Information. Raw sequencing data and the assembled sequences are deposited at the National Center for Biotechnology Information (NCBI) database under the BioProject accession numbers: PRJNA826743 for Illumina raw reads (https://www.ncbi.nlm.nih.gov/bioproject?LinkName=biosample_bioproject&from_uid=27582367), PRJNA831351 for filtered ONT, PacBio, and Hi-C reads, filtered ONT and PacBio mitochondrial reads, and filtered mitochondrial Illumina reads (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA831351/); SAMN44350262 the assembled and annotated genome (https://www.ncbi.nlm.nih.gov/datasets/genome/SAMN44350262/); NC_077490 for the assembled and annotated plastome (https://www.ncbi.nlm.nih.gov/search/all/?term=NC_077490), PP657143 for the assembled and annotated mitogenome (https://www.ncbi.nlm.nih.gov/search/all/?term=PP657143). The non-coding genes and transposable elements annotation are available on Github at https://github.com/NancyChoudhary28/Ranunculus-genomics. We deposited software assembly reports, statistical results, and genome alignments on FigShare (https://doi.org/10.6084/m9.figshare.20488908; private link: https://figshare.com/s/b7fba31a6f2444123906), and the used Python scripts on GitHub at https://github.com/KK260/NCBI-Genome-Tools.

Author Contributions

Funding

This work was financially supported by the Lindemann-foundation of the University of Göttingen to E.H., and the German Research Foundation (DFG) grant Ho4395/10-2 to E.H. within the priority program SPP 1991 "Taxon-Omics: New Approaches for Discovering and Naming Biodiversity". Work in the lab of JdV is supported by the European Research Council through funding under the European Union's Horizon 2020 research and innovation program (Grant Agreement No. 852725; ERC-StG "TerreStriAL") and the DFG via SPP 2237 "MAdLand" (422691801, 528076711). AP, JdV, and EH are part of the GRK 2984/1 "EvoReSt" financed by DFG. I.I. was supported by the Spanish Ministry of Science and

1065   Innovation (MCIN/AEI) and the European Social Fund Plus (ESF+) (grant RYC2022-
1066   038245-I).

1067

Acknowledgments

We acknowledge the excellent technical assistance of Susanne König, Sylvia Swetik, and

Manuela Knauft during PacBio and Hi-C library preparation and DNA sequencing, Jennifer

Krüger for laboratory help during ONT DNA sequencing, Silvia Friedrichs and Gabriele

Ließmann for garden work, Ladislav Hodač for material sampling and field image supply, and

Magnus Nordborg and Robin Burns for bioinformatic advice. This work was supported by the

BMBF-funded de.NBI Cloud within the German Network for Bioinformatics Infrastructure

(de.NBI) (031A532B, 031A533A, 031A533B, 031A534A, 031A535A, 031A537A,

031A537B, 031A537C, 031A537D, 031A538A).

1077

Supplementary Material

1079   The Supplementary Material for this article can be found attached to this article.

1080

References

Adams, K. L., Daley, D. O., Qiu, Y.-L., Whelan, J., and Palmer, J. D. (2000). Repeated, recent and diverse transfers of a mitochondrial gene to the nucleus in flowering plants. *Nature* 408, 354–357. doi: 10.1038/35042567

Adams, K. L., Rosenblueth, M., Qiu, Y.-L., and Palmer, J. D. (2001). Multiple losses and transfers to the nucleus of two mitochondrial succinate dehydrogenase genes during angiosperm evolution. *Genetics* 158, 1289–1300. doi: 10.1093/genetics/158.3.1289

Alafari, H. A., and Abd-Elgawad, M. E. (2021). Differential expression gene/protein contribute to heat stress-responsive in Tetraena propinqua in Saudi Arabia. *Saudi J Biol Sci* 28, 5017–5027. doi: 10.1016/j.sjbs.2021.05.016

Allio, R., Schomaker□Bastos, A., Romiguier, J., Prosdocimi, F., Nabholz, B., and Delsuc, F. (2020). MitoFinder: Efficient automated large□scale extraction of mitogenomic data in target enrichment phylogenomics. *Mol Ecol Resour* 20, 892–905. doi: 10.1111/1755-0998.13160

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data.

Astashyn, A., Tvedte, E. S., Sweeney, D., Sapojnikov, V., Bouk, N., Joukov, V., et al. (2024). Rapid and sensitive detection of genome contamination at scale with FCS-GX. *Genome Biol* 25, 60. doi: 10.1186/s13059-024-03198-7

Baltisberger, M., and Hörandl, E. (2016). Karyotype evolution supports the molecular phylogeny in the genus Ranunculus (Ranunculaceae). *Perspect Plant Ecol Evol Syst* 18, 1–14. doi: 10.1016/j.ppees.2015.11.001

Bennetzen, J. L. (2005). Mechanisms of recent genome size variation in flowering plants. *Ann Bot* 95, 127–132. doi: 10.1093/aob/mci008

Bi, C., Lu, N., Xu, Y., He, C., and Lu, Z. (2020). Characterization and analysis of the mitochondrial genome of common bean (Phaseolus vulgaris) by comparative genomic approaches. *Int J Mol Sci* 21, 3778. doi: 10.3390/ijms21113778

Bolger, A. M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30, 2114–2120. doi: 10.1093/bioinformatics/btu170

Bonen, L., and Calixte, S. (2006). Comparative analysis of bacterial-origin genes for plant mitochondrial ribosomal proteins. *Mol Biol Evol* 23, 701–712. doi: 10.1093/molbev/msj080

Bradican, J. P., Tomasello, S., Boscutti, F., Karbstein, K., and Hörandl, E. (2023). Phylogenomics of Southern European Taxa in the Ranunculus auricomus Species Complex: The Apple Doesn't Fall Far from the Tree. *Plants* 12, 3664. doi: 10.3390/plants12213664

Bradican, J. P., Tomasello, S., Vollmer, J., and Hörandl, E. (2024). Converging forms: an examination of sub-Arctic, circumarctic, and Central Asian Ranunculus auricomus agg. populations. *Front Plant Sci* 15, 1–10. doi: 10.3389/fpls.2024.1415059

Brandvain, Y., and Wade, M. J. (2009). The functional transfer of genes from the mitochondria to the nucleus: The effects of selection, mutation, population size and rate of self-fertilization. *Genetics* 182, 1129–1139. doi: 10.1534/genetics.108.100024

Bray, N. L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* 34, 525–527. doi: 10.1038/nbt.3519

Brukhin, V., Osadtchiy, J. V., Florez-Rueda, A. M., Smetanin, D., Bakin, E., Nobre, M. S., et al. (2019). The *Boechera* genus as a resource for apomixis research. *Front Plant Sci* 10, 1–19. doi: 10.3389/fpls.2019.00392

Brůna, T., Lomsadze, A., and Borodovsky, M. (2024). GeneMark-ETP significantly improves the accuracy of automatic annotation of large eukaryotic genomes. *Genome Res* 34, 757–768. doi: 10.1101/gr.278373.123

Bushnell, B. (2022). BBTools software package. Available at: http://sourceforgenet/
projects/bbmap
Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P., and Huerta-Cepas, J. (2021).
eggNOG-mapper v2: functional annotation, orthology assignments, and domain
prediction at the metagenomic scale. *Mol Biol Evol* 38, 5825–5829. doi:
10.1093/molbev/msab293
Chan, P. P., Lin, B. Y., Mak, A. J., and Lowe, T. M. (2021). tRNAscan-SE 2.0: improved
detection and functional classification of transfer RNA genes. *Nucleic Acids Res* 49,
9077–9096. doi: 10.1093/nar/gkab688
Chang, J., Stahlke, A. R., Chudalayandi, S., Rosen, B. D., Childers, A. K., and Severin, A. J.
(2023). polishCLR: A nextflow workflow for polishing PacBio CLR genome assemblies.
Genome Biol Evol 15, 1–5. doi: 10.1093/gbe/evad020
Chen, D., Pan, Y., Wang, Y., Cui, Y.-Z., Zhang, Y.-J., Mo, R., et al. (2021). The
chromosome-level reference genome of Coptis chinensis provides insights into genomic
evolution and berberine biosynthesis. *Hortic Res* 8, 121. doi: 10.1038/s41438-021-
00559-2
Cheng, H., Concepcion, G. T., Feng, X., Zhang, H., and Li, H. (2021). Haplotype-resolved de
novo assembly using phased assembly graphs with hifiasm. *Nat Methods* 18, 170–175.
doi: 10.1038/s41592-020-01056-5
Choi, I.-S., Schwarz, E. N., Ruhlman, T. A., Khiyami, M. A., Sabir, J. S. M., Hajarah, N. H.,
et al. (2019). Fluctuations in Fabaceae mitochondrial genome size and content are both
ancient and recent. *BMC Plant Biol* 19, 448. doi: 10.1186/s12870-019-2064-8
Chuan, J., Zhou, A., Hale, L. R., He, M., and Li, X. (2021). Atria: an ultra-fast and accurate
trimmer for adapter and quality trimming. *GigaByte* 2021, 1–18. doi:
10.46471/gigabyte.31
Cummings, H. S., and Hershey, J. W. B. (1994). Translation initiation factor IF1 is essential
for cell viability in Escherichia coli. *J Bacteriol* 176, 198–205. doi:
10.1128/jb.176.1.198-205.1994
Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., et al. (2021).
Twelve years of SAMtools and BCFtools. *Gigascience* 10, 1–4. doi:
10.1093/gigascience/giab008
Dann, M., Bellot, S., Schepella, S., Schaefer, H., and Tellier, A. (2017). Mutation rates in
seeds and seed-banking influence substitution rates across the angiosperm phylogeny.
bioRxiv, 156398. doi: 10.1101/156398
Darling, A. E., Mau, B., and Perna, N. T. (2010). progressiveMauve: multiple genome
alignment with gene gain, loss and rearrangement. *PLoS One* 5, e11147. doi:
10.1371/journal.pone.0011147
Darriba, D., Posada, D., Kozlov, A. M., Stamatakis, A., Morel, B., and Flouri, T. (2020).
ModelTest-NG: a new and scalable tool for the selection of DNA and protein
evolutionary models. *Mol Biol Evol* 37, 291–294. doi: 10.1093/molbev/msz189
Dauphin, B., Grant, J. R., Farrar, D. R., and Rothfels, C. J. (2018). Rapid allopolyploid
radiation of moonwort ferns (*Botrychium*; Ophioglossaceae) revealed by PacBio
sequencing of homologous and homeologous nuclear regions. *Mol Phylogenet Evol* 120,
342–353. doi: 10.1016/j.ympev.2017.11.025
De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M., and Van Broeckhoven, C. (2018).
NanoPack: visualizing and processing long-read sequencing data. *Bioinformatics* 34,
2666–2669. doi: 10.1093/bioinformatics/bty149
Del Amparo, R., Branco, C., Arenas, J., Vicens, A., and Arenas, M. (2021). Analysis of
selection in protein-coding sequences accounting for common biases. *Brief Bioinform*
22, 1–10. doi: 10.1093/bib/bbaa431

Di Genova, A., Buena-Atienza, E., Ossowski, S., and Sagot, M.-F. (2021). Efficient hybrid de
novo assembly of human genomes with WENGAN. *Nat Biotechnol* 39, 422–430. doi:
10.1038/s41587-020-00747-w

Dmitriev, A. A., Pushkova, E. N., Novakovskiy, R. O., Beniaminov, A. D., Rozhmina, T. A.,
Zhuchenko, A. A., et al. (2021). Genome sequencing of fiber flax cultivar Atlant using
Oxford Nanopore and Illumina platforms. *Front Genet* 11, 1–7. doi:
10.3389/fgene.2020.590282

Dodsworth, S., Leitch, A. R., and Leitch, I. J. (2015). Genome size diversity in angiosperms
and its influence on gene space. *Curr Opin Genet Dev* 35, 73–78. doi:
10.1016/j.gde.2015.10.006

Dong, S., Zhao, C., Chen, F., Liu, Y., Zhang, S., Wu, H., et al. (2018). The complete
mitochondrial genome of the early flowering plant *Nymphaea colorata* is highly
repetitive with low recombination. *BMC Genomics* 19, 1–12. doi: 10.1186/s12864-018-
4991-4

Drescher, A., Ruf, S., Calsa, T., Carrer, H., and Bock, R. (2000). The two largest chloroplast
genome☐encoded open reading frames of higher plants are essential genes. *The Plant
Journal* 22, 97–104. doi: 10.1046/j.1365-313x.2000.00722.x

Dudchenko, O., Batra, S. S., Omer, A. D., Nyquist, S. K., Hoeger, M., Durand, N. C., et al.
(2017). De novo assembly of the Aedes aegypti genome using Hi-C yields chromosome-
length scaffolds. *Science (1979)* 356, 92–95. doi: 10.1126/science.aal3327

Durand, N. C., Shamim, M. S., Machol, I., Rao, S. S. P., Huntley, M. H., Lander, E. S., et al.
(2016). Juicer provides a one-click system for analyzing loop-resolution Hi-C
experiments. *Cell Syst* 3, 95–98. doi: 10.1016/j.cels.2016.07.002

Eaton, D. A. R., and Overcast, I. (2020). ipyrad: Interactive assembly and analysis of RADseq
datasets. *Bioinformatics* 36, 2592–2594. doi: 10.1093/bioinformatics/btz966

Emadzade, K., Lehnebach, C., Lockhart, P., and Hörandl, E. (2010). A molecular phylogeny,
morphology and classification of genera of Ranunculeae (Ranunculaceae). *Taxon* 59,
809–828. doi: 10.1002/tax.593011

Ennos, R., French, G., and Hollingsworth, P. (2005). Conserving taxonomic complexity.
Trends Ecol Evol 20, 164–168. doi: 10.1016/j.tree.2005.01.012

Fakih, Z., Plourde, M. B., and Germain, H. (2023). Differential participation of plant
ribosomal proteins from the small ribosomal subunit in protein translation under stress.
Biomolecules 13, 1160. doi: 10.3390/biom13071160

Feng, Y., Xiang, X., Akhter, D., Pan, R., Fu, Z., and Jin, X. (2021). Mitochondrial
phylogenomics of Fagales provides insights intoplant mitogenomemosaic evolution.
Front Plant Sci 12, 1–13. doi: 10.3389/fpls.2021.762195

Freudenthal, J. A., Pfaff, S., Terhoeven, N., Korte, A., Ankenbrand, M. J., and Förster, F.
(2020). A systematic comparison of chloroplast genome assembly tools. *Genome Biol*
21, 254. doi: 10.1186/s13059-020-02153-6

Frith, M. C. (2011). A new repeat-masking method enables specific detection of homologous
sequences. *Nucleic Acids Res* 39, e23–e23. doi: 10.1093/nar/gkq1212

Gabriel, L., Brůna, T., Hoff, K. J., Ebel, M., Lomsadze, A., Borodovsky, M., et al. (2024).
BRAKER3: Fully automated genome annotation using RNA-seq and protein evidence
with GeneMark-ETP, AUGUSTUS, and TSEBRA. *Genome Res* 34, 769–777. doi:
10.1101/gr.278090.123

Gatter, T., von Löhneysen, S., Fallmann, J., Drozdova, P., Hartmann, T., and Stadler, P. F.
(2021). LazyB: fast and cheap genome assembly. *Algorithms for Molecular Biology* 16,
8. doi: 10.1186/s13015-021-00186-5

Gebert, N., Gebert, M., Oeljeklaus, S., von der Malsburg, K., Stroud, D. A., Kulawiak, B., et
al. (2011). Dual function of Sdh3 in the respiratory chain and TIM22 protein Translocase

of the mitochondrial inner membrane. *Mol Cell* 44, 811–818. doi:
10.1016/j.molcel.2011.09.025

Giangrossi, M., Brandi, A., Giuliodori, A. M., Gualerzi, C. O., and Pon, C. L. (2007).
Cold□shock□induced de novo transcription and translation of infA and role of IF1
during cold adaptation. *Mol Microbiol* 64, 807–821. doi: 10.1111/j.1365-
2958.2007.05699.x

Grabherr, M. G.□;, Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., et al.
(2011). Full-length transcriptome assembly from RNA-Seq data without a reference
genome. *Nat Biotechnol* 29, 644–652. doi: 10.1038/nbt.1883

Guo, X., Fang, D., Sahu, S. K., Yang, S., Guang, X., Folk, R., et al. (2021). Chloranthus
genome provides insights into the early diversification of angiosperms. *Nat Commun* 12,
6930. doi: 10.1038/s41467-021-26922-4

Gurevich, A., Saveliev, V., Vyahhi, N., and Tesler, G. (2013). QUAST: quality assessment
tool for genome assemblies. *Bioinformatics* 29, 1072–1075. doi:
10.1093/bioinformatics/btt086

Haas, B. J., Salzberg, S. L., Zhu, W., Pertea, M., Allen, J. E., Orvis, J., et al. (2008).
Automated eukaryotic gene structure annotation using EVidenceModeler and the
program to assemble spliced alignments. *Genome Biol* 9, R7. doi: 10.1186/gb-2008-9-1-
r7

Hakim, S. E., Choudhary, N., Malhotra, K., Peng, J., Arafa, A., Bültemeier, A., et al. (2024).
Phylogenomics and metabolic engineering reveal a conserved gene cluster in Solanaceae
plants for withanolide biosynthesis. *bioRxiv*, 1–44. doi: 10.1101/2024.09.27.614867

Havird, J. C., Hall, M. D., and Dowling, D. K. (2015). The evolution of sex: A new
hypothesis based on mitochondrial mutational erosion. *BioEssays* 37, 951–958. doi:
10.1002/bies.201500057

He, J., Lyu, R., Luo, Y., Xiao, J., Xie, L., Wen, J., et al. (2022). A phylotranscriptome study
using silica gel-dried leaf tissues produces an updated robust phylogeny of
Ranunculaceae. *Mol Phylogenet Evol* 174, 107545. doi: 10.1016/j.ympev.2022.107545

Hodač, L. & Karbstein, K., Tomasello, S., Wäldchen, J., Bradican, J. P., and Hörandl, E.
(2023). Geometric morphometric versus genomic patterns in a large polyploid plant
species complex. *Biology (Basel)* 12, 418. doi: 10.3390/biology12030418

Hodač, L., Barke, B. H., and Hörandl, E. (2018). Mendelian segregation of leaf phenotypes in
experimental F2 hybrids elucidates origin of morphological diversity of the apomictic
Ranunculus auricomus complex. *Taxon* 67, 1082–1092. doi: 10.12705/676.6

Hoencamp, C., Dudchenko, O., Elbatsh, A. M. O., Brahmachari, S., Raaijmakers, J. A., van
Schaik, T., et al. (2021). 3D genomics across the tree of life reveals condensin II as a
determinant of architecture type. *Science (1979)* 372, 984–989. doi:
10.1126/science.abe2218

Hörandl, E. (2008). Evolutionary Implications of self□Compatibility and reproductive fitness
in the apomictic Ranunculus auricomus polyploid complex (Ranunculaceae). *Int J Plant
Sci* 169, 1219–1228. doi: 10.1086/591980

Hörandl, E. (2022). Novel approaches for species concepts and delimitation in polyploids and
hybrids. *Plants* 11, 204. doi: 10.3390/plants11020204

Hörandl, E. (2024). Apomixis and the paradox of sex in plants. *Ann Bot*, 1–18. doi:
10.1093/aob/mcae044

Hörandl, E., and Emadzade, K. (2012). Evolutionary classification□: A case study on the
diverse plant genus Ranunculus L. (Ranunculaceae). *Perspect Plant Ecol Evol Syst* 14,
310–324. doi: 10.1016/j.ppees. 2012.04.001

Hörandl, E., and Greilhuber, J. (2002). Diploid and autotetraploid sexuals and their
relationships to apomicts in the *Ranunculus cassubicus* group: insights from DNA

content and isozyme variation. *Plant Systematics and Evolution* 234, 85–100. doi: 10.1007/s00606-002-0209-x

Hörandl, E., Greilhuber, J., Klímová, K., Paun, O., Temsch, E., Emadzade, K., et al. (2009). Reticulate evolution and taxonomic concepts in the *Ranunculus auricomus* complex (Ranunculaceae): insights from analysis of morphological, karyological and molecular data. *Taxon* 58, 1194–1215. doi: 10.1002/tax.584012

Hörandl, E., Hojsgaard, D., Caperta, A., Conceição, S. I. R., Róis, A. S., Dickinson, T., et al. (2024). Apomixis in systematics, evolution and phylogenetics of angiosperms: Current developments and prospects. *CRC Crit Rev Plant Sci* 0, 1–43. doi: 10.1080/07352689.2024.2396259

Hörandl, E., Paun, O., Johansson, J. T., Lehnebach, C., Armstrong, T., Chen, L., et al. (2005). Phylogenetic relationships and evolutionary traits in Ranunculus s.l. (Ranunculaceae) inferred from ITS sequence analysis. *Mol Phylogenet Evol* 36, 305–327. doi: 10.1016/j.ympev.2005.02.009

Hörandl, E., and Raab-Straube, E. (2015). Ranunculaceae. *Euro+Med Plantbase - the information resource for Euro-Mediterranean plant diversity.* Available at: http://ww2.bgbm.org/EuroPlusMed

Horz, J. M., Wolff, K., Friedhoff, R., and Pucker, B. (2024). Genome sequence of the medicinal and ornamental plant Digitalis purpurea reveals the molecular basis of flower color variation. *bioRxiv*, 1–27.

Hotaling, S., Wilcox, E. R., Heckenhauer, J., Stewart, R. J., and Frandsen, P. B. (2023). Highly accurate long reads are crucial for realizing the potential of biodiversity genomics. *BMC Genomics* 24, 117. doi: 10.1186/s12864-023-09193-9

Hu, C., Jiao, Z., Deng, X., Tu, X., Lu, A., Xie, C., et al. (2022). The ecological adaptation of the unparalleled plastome character evolution in slipper orchids. *Front Plant Sci* 13, 1–13. doi: 10.3389/fpls.2022.1075098

Huang, S., Braun, H., Gawryluk, R. M. R., and Millar, A. H. (2019). Mitochondrial complex II of plants: subunit composition, assembly, and function in respiration and signaling. *The Plant Journal* 98, 405–417. doi: 10.1111/tpj.14227

Jayakodi, M., Golicz, A. A., Kreplak, J., Fechete, L. I., Angra, D., Bednář, P., et al. (2023). The giant diploid faba genome unlocks variation in a global protein crop. *Nature* 615, 652–659. doi: 10.1038/s41586-023-05791-5

Ji, J., Luo, Y., Pei, L., Li, M., Xiao, J., Li, W., et al. (2023). Complete plastid genomes of nine species of Ranunculeae (Ranunculaceae) and their phylogenetic inferences. *Genes (Basel)* 14, 2140. doi: 10.3390/genes14122140

Jiang, M., Ni, Y., Li, J., and Liu, C. (2023). Characterisation of the complete mitochondrial genome of Taraxacum mongolicum revealed five repeat-mediated recombinations. *Plant Cell Rep* 42, 775–789. doi: 10.1007/s00299-023-02994-y

Jin, J.-J., Yu, W.-B., Yang, J.-B., Song, Y., DePamphilis, C. W., Yi, T.-S., et al. (2020). GetOrganelle: a fast and versatile toolkit for accurate de novo assembly of organelle genomes. *Genome Biol* 21, 241. doi: 10.1186/s13059-020-02154-5

Jones, G. (2017). Bayesian phylogenetic analysis for diploid and allotetraploid species networks. *bioRxiv*, 1–15. doi: 10.1101/129361

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., et al. (2014). InterProScan 5: genome-scale protein function classification. *Bioinformatics* 30, 1236–1240. doi: 10.1093/bioinformatics/btu031

Karbstein, K. (2021). Untying Gordian knots: The evolution and biogeography of the large European apomictic polyploid *Ranunculus auricomus* plant complex. University of Göttingen. doi: 21.11130/00-1735-0000-0008-5946-6

Karbstein, K., Kösters, L., Hodač, L., Hofmann, M., Hörandl, E., Tomasello, S., et al. (2024). Species delimitation 4.0: integrative taxonomy meets artificial intelligence. *Trends Ecol Evol* 39, 771–784. doi: 10.1016/j.tree.2023.11.002

Karbstein, K., Rahmsdorf, E., Tomasello, S., Hodač, L., and Hörandl, E. (2020a). Breeding system of diploid sexuals within the *Ranunculus auricomus* complex and its role in a geographical parthenogenesis scenario. *Ecol Evol* 10, 14435–14450. doi: 10.1002/ece3.7073

Karbstein, K., Tomasello, S., Hodač, L., Dunkel, F. G., Daubert, M., and Hörandl, E. (2020b). Phylogenomics supported by geometric morphometrics reveals delimitation of sexual species within the polyploid apomictic *Ranunculus auricomus* complex (Ranunculaceae). *Taxon* 69, 1191–1220. doi: 10.1002/tax.12365

Karbstein, K., Tomasello, S., Hodač, L., Lorberg, E., Daubert, M., and Hörandl, E. (2021). Moving beyond assumptions: Polyploidy and environmental effects explain a geographical parthenogenesis scenario in European plants. *Mol Ecol* 30, 2659–2675. doi: 10.1111/mec.15919

Karbstein, K., Tomasello, S., Hodač, L., Wagner, N., Marinček, P., Barke, B. H., et al. (2022). Untying Gordian knots: Unraveling reticulate polyploid plant evolution by genomic data using the large *Ranunculus auricomus* species complex. *New Phytologist* 235, 2081–2098. doi: 10.1111/nph.18284

Katoh, K., and Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 30, 772–780. doi: 10.1093/molbev/mst010

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., et al. (2012). Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28, 1647–1649. doi: 10.1093/bioinformatics/bts199

Keilwagen, J., Hartung, F., Paulini, M., Twardziok, S. O., and Grau, J. (2018). Combining RNA-seq data and homology-based gene prediction for plants, animals and fungi. *BMC Bioinformatics* 19, 189. doi: 10.1186/s12859-018-2203-5

Keilwagen, J., Wenk, M., Erickson, J. L., Schattat, M. H., Grau, J., and Hartung, F. (2016). Using intron position conservation for homology-based gene prediction. *Nucleic Acids Res* 44, e89–e89. doi: 10.1093/nar/gkw092

Kim, D., Paggi, J. M., Park, C., Bennett, C., and Salzberg, S. L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 37, 907–915. doi: 10.1038/s41587-019-0201-4

Kolmogorov, M., Yuan, J., Lin, Y., and Pevzner, P. A. (2019). Assembly of long, error-prone reads using repeat graphs. *Nat Biotechnol* 37, 540–546. doi: 10.1038/s41587-019-0072-8

Konupková, A., Tomečková, L., Záhonová, K., Oborník, M., and Füssy, Z. (2024). "Easier lost than found? What we know about plastid genome reduction," in *Endosymbiotic Organelle Acquisition*, (Cham: Springer International Publishing), 147–181. doi: 10.1007/978-3-031-57446-7_5

Koren, S., Walenz, B. P., Berlin, K., Miller, J. R., Bergman, N. H., and Phillippy, A. M. (2017). Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res* 27, 722–736. doi: 10.1101/gr.215087.116

Korf, I. (2004). Gene finding in novel genomes. *BMC Bioinformatics* 5, 1–9. doi: 10.1186/1471-2105-5-59

Kozlov, A. M., Darriba, D., Flouri, T., Morel, B., and Stamatakis, A. (2019). RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* 35, 4453–4455. doi: 10.1093/bioinformatics/btz305

Kuznetsov, D., Tegenfeldt, F., Manni, M., Seppey, M., Berkeley, M., Kriventseva, E. V, et al.
(2023). OrthoDB v11: annotation of orthologs in the widest sampling of organismal
diversity. *Nucleic Acids Res* 51, D445–D451. doi: 10.1093/nar/gkac998

Lawniczak, M. K. N., Durbin, R., Flicek, P., Lindblad-Toh, K., Wei, X., Archibald, J. M., et
al. (2022). Standards recommendations for the Earth BioGenome Project. *Proceedings of*
the National Academy of Sciences 119, 1–8. doi: 10.1073/pnas.2115639118

Leebens-Mack, J. H., Barker, M. S., Carpenter, E. J., Deyholos, M. K., Gitzendanner, M. A.,
Graham, S. W., et al. (2019). One thousand plant transcriptomes and the phylogenomics
of green plants. *Nature* 574, 679–685. doi: 10.1038/s41586-019-1693-2

Lehwark, P., and Greiner, S. (2019). GB2sequin - A file converter preparing custom GenBank
files for database submission. *Genomics* 111, 759–761. doi:
10.1016/j.ygeno.2018.05.003

Leitch, I., Johnston, E., Pellicer, J., Hidalgo, O., and Bennet, M. (2019). Plant DNA C-values
database (release 7.1, Apr 2019). Available at: http://data.kew.org/cvalues/

Lemoine, F., and Gascuel, O. (2021). Gotree/Goalign: toolkit and Go API to facilitate the
development of phylogenetic workflows. *NAR Genom Bioinform* 3, 1–5. doi:
10.1093/nargab/lqab075

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34,
3094–3100. doi: 10.1093/bioinformatics/bty191

Li, S.-F., Su, T., Cheng, G.-Q., Wang, B.-X., Li, X., Deng, C.-L., et al. (2017). Chromosome
evolution in connection with repetitive sequences and epigenetics in plants. *Genes*
(Basel) 8, 290. doi: 10.3390/genes8100290

Li, X., Li, Y., Sylvester, S. P., Zang, M., El□Kassaby, Y. A., and Fang, Y. (2021).
Evolutionary patterns of nucleotide substitution rates in plastid genomes of Quercus.
Ecol Evol 11, 13401–13414. doi: 10.1002/ece3.8063

Li, Z., Parris, S., and Saski, C. A. (2020). A simple plant high-molecular-weight DNA
extraction method suitable for single-molecule technologies. *Plant Methods* 16, 38. doi:
10.1186/s13007-020-00579-4

Liao, W.-J., Hu, Y., Zhu, B.-R., Zhao, X.-Q., Zeng, Y.-F., and Zhang, D.-Y. (2009). Female
reproductive success decreases with display size in monkshood, Aconitum kusnezoffii
(Ranunculaceae). *Ann Bot* 104, 1405–1412. doi: 10.1093/aob/mcp237

Liao, X., Zhu, W., Zhou, J., Li, H., Xu, X., Zhang, B., et al. (2023). Repetitive DNA sequence
detection and its role in the human genome. *Commun Biol* 6, 954. doi: 10.1038/s42003-
023-05322-y

Lin, Q., Braukmann, T. W. A., Soto Gomez, M., Mayer, J. L. S., Pinheiro, F., Merckx, V. S.
F. T., et al. (2022). Mitochondrial genomic data are effective at placing
mycoheterotrophic lineages in plant phylogeny. *New Phytologist* 236, 1908–1921. doi:
10.1111/nph.18335

Lindell, T. (1998). Breeding systems and crossing experiments in Anemone patens and in the
Anemome pulsatilla group (Ranunculaceae). *Nord J Bot* 18, 549–561. doi:
10.1111/j.1756-1051.1998.tb01535.x

Liu, L.-X., Li, R., Worth, J. R. P., Li, X., Li, P., Cameron, K. M., et al. (2017). The complete
chloroplast genome of Chinese bayberry (*Morella rubra*, Myricaceae): Implications for
understanding the evolution of Fagales. *Front Plant Sci* 8, 1–15. doi:
10.3389/fpls.2017.00968

Liu, S., Ni, Y., Li, J., Zhang, X., Yang, H., Chen, H., et al. (2023). CPGView: A package for
visualizing detailed chloroplast genome structures. *Mol Ecol Resour* 23, 694–704. doi:
10.1111/1755-0998.13729

Liu, S.-L., and Adams, K. (2008). Molecular adaptation and expression evolution following
duplication of genes for organellar ribosomal protein S13 in rosids. *BMC Evol Biol* 8, 25.
doi: 10.1186/1471-2148-8-25
Liu, Y., Wang, B., Shu, S., Li, Z., Song, C., Liu, D., et al. (2021). Analysis of the Coptis
chinensis genome reveals the diversification of protoberberine-type alkaloids. *Nat*
Commun 12, 3276. doi: 10.1038/s41467-021-23611-0
Long, J., He, W.-C., Peng, H.-W., Erst, A. S., Wang, W., and Xiang, K.-L. (2024).
Comparative plastome analysis of the sister genera Ceratocephala and Myosurus
(Ranunculaceae) reveals signals of adaptive evolution to arid and aquatic environments.
BMC Plant Biol 24, 202. doi: 10.1186/s12870-024-04891-2
Lunkova, N. F., and Ivanov, V. B. (2022). Features of the genome correlated with the
annuality of plant. *Russian Journal of Plant Physiology* 69, 87. doi:
10.1134/S1021443722050132
Majoros, W. H., Pertea, M., and Salzberg, S. L. (2004). TigrScan and GlimmerHMM: two
open source ab initio eukaryotic gene-finders. *Bioinformatics* 20, 2878–2879. doi:
10.1093/bioinformatics/bth315
Mallet, J. (2005). Hybridization as an invasion of the genome. *Trends Ecol Evol* 20, 229–237.
doi: 10.1016/j.tree.2005.02.010
Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., and Zdobnov, E. M. (2021). BUSCO
update: Novel and streamlined workflows along with broader and deeper phylogenetic
coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Mol Biol Evol* 38,
4647–4654. doi: 10.1093/molbev/msab199
Marks, P., Garcia, S., Martinez, A., Belhocine, K., Bernate, J., Bharadwaj, R., et al. (2017).
Resolving the full spectrum of human genome variation using linked-reads. *bioRxiv*, 1–
43. doi: 10.1101/230946
Mascher, M., Gundlach, H., Himmelbach, A., Beier, S., Twardziok, S. O., Wicker, T., et al.
(2017). A chromosome conformation capture ordered sequence of the barley genome.
Nature 544, 427–433. doi: 10.1038/nature22043
Mascher, M., Wicker, T., Jenkins, J., Plott, C., Lux, T., Koh, C. S., et al. (2021). Long-read
sequence assembly: a technical evaluation in barley. *Plant Cell* 33, 1888–1906. doi:
10.1093/plcell/koab077
Michael, T. P. (2014). Plant genome size variation: bloating and purging DNA. *Brief Funct*
Genomics 13, 308–317. doi: 10.1093/bfgp/elu005
Millen, R. S., Olmstead, R. G., Adams, K. L., Palmer, J. D., Lao, N. T., Heggie, L., et al.
(2001). Many parallel losses of infA from chloroplast DNA during angiosperm evolution
with multiple independent transfers to the nucleus. *Plant Cell* 13, 645–658. doi:
10.1105/tpc.13.3.645
Mokhtar, M. M., Abd-Elhalim, H. M., and El Allali, A. (2023). A large-scale assessment of
the quality of plant genome assemblies using the LTR assembly index. *AoB Plants* 15,
1–8. doi: 10.1093/aobpla/plad015
Møller, I. M., Rasmusson, A. G., and Van Aken, O. (2021). Plant mitochondria – past, present
and future. *The Plant Journal* 108, 912–959. doi: 10.1111/tpj.15495
Murigneux, V., Rai, S. K., Furtado, A., Bruxner, T. J. C., Tian, W., Harliwong, I., et al.
(2020). Comparison of long-read methods for sequencing and assembly of a plant
genome. *Gigascience* 9, 1–11. doi: 10.1093/gigascience/giaa146
Nawrocki, E. P., and Eddy, S. R. (2013). Infernal 1.1: 100-fold faster RNA homology
searches. *Bioinformatics* 29, 2933–2935. doi: 10.1093/bioinformatics/btt509
Nowak, M. S., Harder, B., Meckoni, S. N., Friedhoff, R., Wolff, K., and Pucker, B. (2024).
Genome sequence and RNA-seq analysis reveal genetic basis of flower coloration in the
giant water lily Victoria cruziana. *bioRxiv*, 1–23. doi: 10.1101/2024.06.15.599162

Ontiveros-Palacios, N., Cooke, E., Nawrocki, E. P., Triebel, S., Marz, M., Rivas, E., et al.
(2024). Rfam 15: RNA families database in 2025. *Nucleic Acids Res*, 1–10. doi:
10.1093/nar/gkae1023
Ou, S., and Jiang, N. (2018). LTR_retriever: A highly accurate and sensitive program for
identification of long terminal repeat retrotransposons. *Plant Physiol* 176, 1410–1422.
doi: 10.1104/pp.17.01310
Ou, S., and Jiang, N. (2019). LTR_FINDER_parallel: parallelization of LTR_FINDER
enabling rapid identification of long terminal repeat retrotransposons. *Mob DNA* 10, 48.
doi: 10.1186/s13100-019-0193-0
Padmarasu, S., Himmelbach, A., Mascher, M., and Stein, N. (2019). In situ Hi-C for plants:
An improved method to detect long-range chromatin interactions. *Methods in Molecular*
Biology 1933, 441–472. doi: 10.1007/978-1-4939-9045-0_28
Paetzold, C., Barke, B. H., and Hörandl, E. (2022). Evolution of transcriptomes in early-
generation hybrids of the apomictic *Ranunculus auricomus complex* (Ranunculaceae).
Int J Mol Sci 23, 13881. doi: 10.3390/ijms232213881
Palmer, J. M., and Stajich, J. E. (2022). Funannotate (Version 1.8.15) [Computer software].
Park, H.-S., Lee, W. K., Lee, S.-C., Lee, H. O., Joh, H. J., Park, J. Y., et al. (2021).
Inheritance of chloroplast and mitochondrial genomes in cucumber revealed by four
reciprocal F1 hybrid combinations. *Sci Rep* 11, 2506. doi: 10.1038/s41598-021-81988-w
Park, S., Jansen, R. K., and Park, S. (2015). Complete plastome sequence of *Thalictrum*
coreanum (Ranunculaceae) and transfer of the rpl32 gene to the nucleus in the ancestor
of the subfamily Thalictroideae. *BMC Plant Biol* 15, 40. doi: 10.1186/s12870-015-0432-
6
Park, S., and Park, S. (2020). Large-scale phylogenomics reveals ancient introgression in
Asian Hepatica and new insights into the origin of the insular endemic *Hepatica maxima*.
Sci Rep 10, 16288. doi: 10.1038/s41598-020-73397-2
Paun, O., Stuessy, T. F., and Hörandl, E. (2006). The role of hybridization, polyploidization
and glaciation in the origin and evolution of the apomictic *Ranunculus cassubicus*
complex. *New Phytologist* 171, 223–236. doi: 10.1111/j.1469-8137.2006.01738.x
Pease, J. B., Brown, J. W., Walker, J. F., Hinchliff, C. E., and Smith, S. A. (2018). Quartet
Sampling distinguishes lack of support from conflicting support in the green plant tree of
life. *Am J Bot* 105, 385–403. doi: 10.1002/ajb2.1016
Pellino, M., Hojsgaard, D., Schmutzer, T., Scholz, U., Hörandl, E., Vogel, H., et al. (2013).
Asexual genome evolution in the apomictic *Ranunculus auricomus* complex: examining
the effects of hybridization and mutation accumulation. *Mol Ecol* 22, 5908–5921. doi:
10.1111/mec.12533
Peška, V., Mandáková, T., Ihradská, V., and Fajkus, J. (2019). Comparative dissection of
three giant genomes: Allium cepa, Allium sativum, and Allium ursinum. *Int J Mol Sci*
20, 733. doi: 10.3390/ijms20030733
Petersen, G., Zervas, A., Pedersen, H. Æ., and Seberg, O. (2018). Genome reports:
Contractedgenes and dwarfed plastome in mycoheterotrophic Sciaphila thaidanica
(Triuridaceae, Pandanales). *Genome Biol Evol* 10, 976–981. doi: 10.1093/gbe/evy064
Postel, Z., and Touzet, P. (2020). Cytonuclear genetic incompatibilities in plant speciation.
Plants 9, 487. doi: 10.3390/plants9040487
Prjibelski, A., Antipov, D., Meleshko, D., Lapidus, A., and Korobeynikov, A. (2020). Using
SPAdes de novo assembler. *Curr Protoc Bioinformatics* 70, 1–29. doi: 10.1002/cpbi.102
Pucker, B. (2024). Functional annotation - How to tackle the bottleneck in plant genomics.
Preprints.org, 1–15. doi: 10.20944/preprints202402.0645.v1

Pucker, B., Irisarri, I., de Vries, J., and Xu, B. (2022). Plant genome sequence assembly in the era of long reads: Progress, challenges and future directions. *Quantitative Plant Biology* 3, e5. doi: 10.1017/qpb.2021.18

Putintseva, Y. A., Bondar, E. I., Simonov, E. P., Sharov, V. V., Oreshkova, N. V., Kuzmin, D. A., et al. (2020). Siberian larch (Larix sibirica Ledeb.) mitochondrial genome assembled using both short and long nucleotide sequence reads is currently the largest known mitogenome. *BMC Genomics* 21, 654. doi: 10.1186/s12864-020-07061-4

Qu, Z., Wang, W., and Adelson, D. L. (2023). Chromosomal level genome assembly of medicinal plant Sophora flavescens. *Sci Data* 10, 572. doi: 10.1038/s41597-023-02490-8

Quesneville, H. (2020). Twenty years of transposable element analysis in the Arabidopsis thaliana genome. *Mob DNA* 11, 28. doi: 10.1186/s13100-020-00223-x

Quinlan, A. R., and Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* 26, 841–842. doi: 10.1093/bioinformatics/btq033

Rambaut, A. (2014). Figtree, a graphical viewer of phylogenetic trees. Available at: http://tree.bio.ed.ac.uk/software/figtree

Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology (and other things). *Methods Ecol Evol* 3, 217–223. doi: 10.1111/j.2041-210X.2011.00169.x

Revell, L. J. (2024). phytools 2.0: an updated R ecosystem for phylogenetic comparative methods (and other things). *PeerJ* 12, e16505. doi: 10.7717/peerj.16505

Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., et al. (2021). Towards complete and error-free genome assemblies of all vertebrate species. *Nature* 592, 737–746. doi: 10.1038/s41586-021-03451-0

Rice, A., Glick, L., Abadi, S., Einhorn, M., Kopelman, N. M., Salman-Minkov, A., et al. (2015). The Chromosome Counts Database (CCDB) - a community resource of plant chromosome numbers. *New Phytologist* 206, 19–26. doi: 10.1111/nph.13191

Robles, P., and Quesada, V. (2017). Emerging roles of mitochondrial ribosomal proteins in plant development. *Int J Mol Sci* 18, 2595. doi: 10.3390/ijms18122595

Saha, A., Das, S., Moin, M., Dutta, M., Bakshi, A., Madhav, M. S., et al. (2017). Genome-wide identification and comprehensive expression profiling of ribosomal protein small subunit (RPS) genes and their comparative analysis with the large subunit (RPL) genes in rice. *Front Plant Sci* 8, 1–21. doi: 10.3389/fpls.2017.01553

Schmidt, A. (2020). Controlling apomixis: Shared features and distinct characteristics of gene regulation. *Genes (Basel)* 11, 329. doi: 10.3390/genes11030329

Schnable, P. S., Ware, D., Fulton, R. S., Stein, J. C., Wei, F., Pasternak, S., et al. (2009). The B73 maize genome: complexity, diversity, and dynamics. *Science (1979)* 326, 1112–1115. doi: 10.1126/science.1178534

Shen, W., Le, S., Li, Y., and Hu, F. (2016). SeqKit: A cross-platform and ultrafast toolkit for fasta/q file manipulation. *PLoS One* 11, e0163962. doi: 10.1371/journal.pone.0163962

Shi, C., Liu, Y., Huang, H., Xia, E.-H., Zhang, H.-B., and Gao, L.-Z. (2013). Contradiction between plastid gene transcription and function due to complex posttranscriptional splicing: an exemplary study of *ycf15* function and evolution in angiosperms. *PLoS One* 8, e59620. doi: 10.1371/journal.pone.0059620

Shi, J., and Liang, C. (2019). Generic Repeat Finder: A high-sensitivity tool for genome-wide de novo repeat detection. *Plant Physiol* 180, 1803–1815. doi: 10.1104/pp.19.00386

Shi, L., Chen, H., Jiang, M., Wang, L., Wu, X., Huang, L., et al. (2019). CPGAVAS2, an integrated plastome sequence annotator and analyzer. *Nucleic Acids Res* 47, W65–W73. doi: 10.1093/nar/gkz345

Sim, S. B., Corpuz, R. L., Simmonds, T. J., and Geib, S. M. (2022). HiFiAdapterFilt, a memory efficient read processing pipeline, prevents occurrence of adapter sequence in

PacBio HiFi reads and their negative impacts on genome assembly. *BMC Genomics* 23, 157. doi: 10.1186/s12864-022-08375-1

Skippington, E., Barkman, T. J., Rice, D. W., and Palmer, J. D. (2015). Miniaturized mitogenome of the parasitic plant Viscum scurruloideum is extremely divergent and dynamic and has lost all nad genes. *Proceedings of the National Academy of Sciences* 112, E3515–E3524. doi: 10.1073/pnas.1504491112

Šlenker, M., Kantor, A., Marhold, K., Schmickl, R., Mandáková, T., Lysak, M. A., et al. (2021). Allele sorting as a novel approach to resolving the origin of allotetraploids using Hyb-Seq data: A case study of the Balkan Mountain endemic *Cardamine barbaraeoides*. *Front Plant Sci* 12, 1–22. doi: 10.3389/fpls.2021.659275

Sloan, D. B. (2013). One ring to rule them all? Genome sequencing provides new insights into the 'master circle' model of plant mitochondrial DNA structure. *New Phytologist* 200, 978–985. doi: 10.1111/nph.12395

Sobanski, J., Giavalisco, P., Fischer, A., Kreiner, J. M., Walther, D., Schöttler, M. A., et al. (2019). Chloroplast competition is controlled by lipid biosynthesis in evening primroses. *Proc Natl Acad Sci U S A* 116, 5665–5674. doi: 10.1073/pnas.1811661116

Soltis, P. S., Marchant, D. B., Van de Peer, Y., and Soltis, D. E. (2015). Polyploidy and genome evolution in plants. *Curr Opin Genet Dev* 35, 119–125. doi: 10.1016/j.gde.2015.11.003

Stanke, M., Diekhans, M., Baertsch, R., and Haussler, D. (2008). Using native and syntenically mapped cDNA alignments to improve de novo gene finding. *Bioinformatics* 24, 637–644. doi: 10.1093/bioinformatics/btn013

Stanke, M., Schöffmann, O., Morgenstern, B., and Waack, S. (2006). Gene prediction in eukaryotes with a generalized hidden Markov model that uses hints from external sources. *BMC Bioinformatics* 7, 62. doi: 10.1186/1471-2105-7-62

Stein, D. B., Palmer, J. D., and Thompson, W. F. (1986). Structural evolution and flip-flop recombination of chloroplast DNA in the fern genus *Osmunda*. *Curr Genet* 10, 835–841. doi: 10.1007/BF00418530

Steinegger, M., and Söding, J. (2017). MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nat Biotechnol* 35, 1026–1028. doi: 10.1038/nbt.3988

Sterck, L., Rombauts, S., Vandepoele, K., Rouze, P., and Vandepeer, Y. (2007). How many genes are there in plants (… and why are they there)? *Curr Opin Plant Biol* 10, 199–203. doi: 10.1016/j.pbi.2007.01.004

Su, W., Gu, X., and Peterson, T. (2019). TIR-Learner, a new ensemble method for TIR transposable element annotation, provides evidence for abundant new transposable elements in the maize genome. *Mol Plant* 12, 447–460. doi: 10.1016/j.molp.2019.02.008

Sun, P., Jiao, B., Yang, Y., Shan, L., Li, T., Li, X., et al. (2022). WGDI: A user-friendly toolkit for evolutionary analyses of whole-genome duplications and ancestral karyotypes. *Mol Plant* 15, 1841–1851. doi: 10.1016/j.molp.2022.10.018

Syngelaki, E., Paetzold, C., and Hörandl, E. (2021). Gene expression profiles suggest a better cold acclimation of polyploids in the Alpine species *Ranunculus kuepferi* (Ranunculaceae). *Genes (Basel)* 12, 1818. doi: 10.3390/genes12111818

Tamura, M. (1993). "Ranunculaceae," in *The Families and Genera of Vascular Plants. 2. Flowering Plants. Dicotyledons, Magnoliid, Hamamelid, and Caryophyllid Families*, eds. K. Kubitzki, J. G. Rohwer, and V. Bittrich, 563–583.

Thorvaldsdottir, H., Robinson, J. T., and Mesirov, J. P. (2013). Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform* 14, 178–192. doi: 10.1093/bib/bbs017

Tomasello, S., Karbstein, K., Hodač, L., Pätzold, C., and Hörandl, E. (2020). Phylogenomics
unravels Quaternary vicariance and allopatric speciation patterns in temperate☐montane
plant species: a case study on the *Ranunculus auricomus* species complex. *Mol Ecol* 29,
2031–2049. doi: 10.1111/mec.15458
Tomasello, S., Manzo, E., and Karbstein, K. (2024). Comparative plastome assembly of the
yellow ironweed (Verbesina alternifolia) using Nanopore and Illumina reads. *Front*
Plant Sci 15, 1–10. doi: 10.3389/fpls.2024.1429494
Uliano-Silva, M., Ferreira, J. G. R. N., Krasheninnikova, K., Blaxter, M., Mieszkowska, N.,
Hall, N., et al. (2023). MitoHiFi: a python pipeline for mitochondrial genome assembly
from PacBio high fidelity reads. *BMC Bioinformatics* 24, 288. doi: 10.1186/s12859-023-
05385-y
Vaillancourt, B., and Buell, C. R. (2019). High molecular weight DNA isolation method from
diverse plant species for use with Oxford Nanopore sequencing. *bioRxiv*, 1–13. doi:
10.1101/783159
Van De Peer, Y., Mizrachi, E., and Marchal, K. (2017). The evolutionary significance of
polyploidy. *Nat Rev Genet* 18, 411–424. doi: 10.1038/nrg.2017.26
Vaser, R., Sović, I., Nagarajan, N., and Šikić, M. (2017). Fast and accurate de novo genome
assembly from long uncorrected reads. *Genome Res* 27, 737–746. doi:
10.1101/gr.214270.116
Walker, J. F., Jansen, R. K., Zanis, M. J., and Emery, N. C. (2015). Sources of inversion
variation in the small single copy (SSC) region of chloroplast genomes. *Am J Bot* 102,
1751–1752. doi: 10.3732/ajb.1500299
Wang, D., Zheng, Z., Li, Y., Hu, H., Wang, Z., Du, X., et al. (2021). Which factors contribute
most to genome size variation within angiosperms? *Ecol Evol* 11, 2660–2668. doi:
10.1002/ece3.7222
Wang, W., Lin, L., Xiang, X.-G., Ortiz, R. D. C., Liu, Y., Xiang, K.-L., et al. (2016). The rise
of angiosperm-dominated herbaceous floras: Insights from Ranunculaceae. *Sci Rep* 6,
27259. doi: 10.1038/srep27259
Wang, W., Lu, A.-M., Ren, Y., Endress, M. E., and Chen, Z.-D. (2009). Phylogeny and
classification of Ranunculales: Evidence from four molecular loci and morphological
data. *Perspect Plant Ecol Evol Syst* 11, 81–110. doi: 10.1016/j.ppees.2009.01.001
Wang, Y., Liu, Y., Miao, K., Hou, L., Guo, X., and Ji, Y. (2024). A haplotype-resolved
genome assembly of Coptis teeta, an endangered plant of significant medicinal value. *Sci*
Data 11, 1012. doi: 10.1038/s41597-024-03861-5
Waterhouse, R. M., Seppey, M., Simão, F. A., Manni, M., Ioannidis, P., Klioutchnikov, G., et
al. (2018). BUSCO applications from quality assessments to gene prediction and
phylogenomics. *Mol Biol Evol* 35, 543–548. doi: 10.1093/molbev/msx319
Weisenfeld, N. I., Kumar, V., Shah, P., Church, D. M., and Jaffe, D. B. (2018). Corrigendum:
Direct determination of diploid genome sequences. *Genome Res* 28, 606.1-606.1. doi:
10.1101/gr.235812.118
Wick, R. R., Schultz, M. B., Zobel, J., and Holt, K. E. (2015). Bandage: interactive
visualization of de novo genome assemblies. *Bioinformatics* 31, 3350–3352. doi:
10.1093/bioinformatics/btv383
Wicker, T., Sabot, F., Hua-Van, A., Bennetzen, J. L., Capy, P., Chalhoub, B., et al. (2007). A
unified classification system for eukaryotic transposable elements. *Nat Rev Genet* 8,
973–982. doi: 10.1038/nrg2165
Woodson, J. D., and Chory, J. (2008). Coordination of gene expression between organellar
and nuclear genomes. *Nat Rev Genet* 9, 383–395. doi: 10.1038/nrg2348

Xie, J., Zhao, H., Li, K., Zhang, R., Jiang, Y., Wang, M., et al. (2020). A chromosome-scale
reference genome of *Aquilegia oxysepala var. kansuensis*. *Hortic Res* 7, 113. doi:
10.1038/s41438-020-0328-y
Xiong, W., He, L., Lai, J., Dooner, H. K., and Du, C. (2014). HelitronScanner uncovers a
large overlooked cache of Helitron transposons in many plant genomes. *Proceedings of*
the National Academy of Sciences 111, 10263–10268. doi: 10.1073/pnas.1410068111
Xu, Z., and Wang, H. (2007). LTR_FINDER: an efficient tool for the prediction of full-length
LTR retrotransposons. *Nucleic Acids Res* 35, W265–W268. doi: 10.1093/nar/gkm286
Yu, W., Luo, H., Yang, J., Zhang, S., Jiang, H., Zhao, X., et al. (2024). Comprehensive
assessment of 11 de novo HiFi assemblers on complex eukaryotic genomes and
metagenomes. *Genome Res* 34, 326–340. doi: 10.1101/gr.278232.123
Zhai, W., Duan, X., Zhang, R., Guo, C., Li, L., Xu, G., et al. (2019). Chloroplast genomic
data provide new and robust insights into the phylogeny and evolution of the
Ranunculaceae. *Mol Phylogenet Evol* 135, 12–21. doi: 10.1016/j.ympev.2019.02.024
Zhang, R.-G., Li, G.-Y., Wang, X.-L., Dainat, J., Wang, Z.-X., Ou, S., et al. (2022). TEsorter:
An accurate and fast method to classify LTR-retrotransposons in plant genomes. *Hortic*
Res 9, 0–3. doi: 10.1093/hr/uhac017
Zhang, X., Chen, H., Ni, Y., Wu, B., Li, J., Burzyński, A., et al. (2024). Plant mitochondrial
genome map (PMGmap): A software tool for the comprehensive visualization of coding,
noncoding and genome features of plant mitochondrial genomes. *Mol Ecol Resour*, 1–15.
doi: 10.1111/1755-0998.13952
Zhigang, Z., Guozhen, D., Xianhui, Z., Mantang, W., and Qingji, R. (2006). Variations with
altitude in reproductive traits and resource allocation of three Tibetan species of
Ranunculaceae. *Aust J Bot* 54, 691. doi: 10.1071/BT05015
Zhou, W., Armijos, C. E., Lee, C., Lu, R., Wang, J., Ruhlman, T. A., et al. (2023). Plastid
genome assembly using long read data. *Mol Ecol Resour* 23, 1442–1457. doi:
10.1111/1755-0998.13787
Zimin, A. V., Puiu, D., Luo, M.-C., Zhu, T., Koren, S., Marçais, G., et al. (2017). Hybrid
assembly of the large and highly repetitive genome of *Aegilops tauschii*, a progenitor of
bread wheat, with the MaSuRCA mega-reads algorithm. *Genome Res* 27, 787–792. doi:
10.1101/gr.213405.116
Zimin, A. V., and Salzberg, S. L. (2020). The genome polishing tool POLCA makes fast and
accurate corrections in genome assemblies. *PLoS Comput Biol* 16, e1007981. doi:
10.1371/journal.pcbi.1007981
Zimin, A. V., and Salzberg, S. L. (2022). The SAMBA tool uses long reads to improve the
contiguity of genome assemblies. *PLoS Comput Biol* 18, 1–11. doi:
10.1371/journal.pcbi.1009860
1708