1 **Green algae CO₂ capture is powered by alternative electron pathways of photosynthesis**
2

Authors: Gilles Peltier[1], Carolyne Stoffel[2], Justin Findinier[2], Sai Kiran Madireddi[2], Ousmane
Dao[1], Virginie Epting[1], Amélie Morin[1], Arthur Grossman[2,3], Yonghua Li-Beisson[1], Adrien
Burlacot[2,3*]

Affiliations:
[1] Aix Marseille Univ, CEA, CNRS, Institut de Biosciences et Biotechnologies Aix-Marseille,
CEA Cadarache, 13108 Saint Paul-lez-Durance, France
[2] Department of Plant Biology, The Carnegie Institution for Science, Stanford, CA, 94305
USA
[3] Department of Biology, Stanford University, Stanford, CA, 94305, USA
12
13 [*]corresponding author: A.B., aburlacot@carnegiescience.edu
14
15 **ORCID IDs:** 0000-0001-7434-6416 (A.B.), 0000-0002-4078-6250 (C.S.), 0000-0001-9019-
16 8900 (J.F.), 0000-0002-1820-1033 (S.M.), 0000-0002-7040-5770 (O.D), 0000-0003-3318-
17 8778 (A.M.), 0000-0003-1064-1816 (Y.L.-B.), 0000-0002-3747-5881(A.G.), 0000-0002-
18 2226-3931 (G.P.)

19
20

21 **Author contributions:** A.B. led and coordinated the research, A.B. and G.P. designed
22 research; A.B., C.S., J.F., O.D., S.M., V.E., A.M. and G.P. performed research; A.B. and G.P.
23 contributed new reagents/analytic tools; A.B. analyzed data and made all figures; A.B. and
24 G.P. wrote the paper with inputs from Y. L.-B. and A.G.

25
26 **One sentence summary:** Quantifying the contribution of the main electron flow pathways to
27 algal CO₂ fixation.
28

Abstract

On Earth, microalgae contribute to about half of global net photosynthesis. During

photosynthesis, sunlight is converted into chemical energy (ATP and NADPH) used by

metabolism to convert CO_2 into biomass. Alternative electron pathways of photosynthesis

have been proposed to generate additional ATP that is required for sustaining CO_2 fixation,

but the relative importance of each pathway remains elusive. Here, we dissect and quantify

the contribution of cyclic, pseudo-cyclic and chloroplast to mitochondria electron flows for

their ability to sustain net photosynthesis in the microalga *Chlamydomonas reinhardtii*. We

show that each pathway has the potential to energize substantial CO_2 fixation, can compensate

each other, and that the additional energy requirement to fix CO_2 is more than 3 times higher

than previous estimations. We further show that all pathways have very different efficiencies

at energizing CO_2 fixation, with the chloroplast- mitochondria interaction being the most

efficient, thus laying bioenergetic foundations for biotechnological improvement of CO_2

capture.

Introduction

Oxygenic photosynthesis is the main process responsible for the carbon input into ecosystems on Earth, with net CO_2 fixation by photosynthesis representing more than ten times anthropic CO_2 emissions (IPCC, 2021). During photosynthesis, carbon fixation is powered by the combined action of photosystems (PS) I and II, establishing a linear electron flow (LEF) that produces reducing power (NADPH) and a trans-thylakoidal proton motive force (*pmf*) used to generate ATP. Both ATP and NADPH fuel the CO_2 fixation reactions of the Calvin-Benson-Bassham (CBB) cycle and the further processing of the glyceraldehyde-3-phosphate (G3P) to regenerated ribulose-1-5 bisphosphate and produce building blocks used for the elaboration of biomass (Johnson and Alric, 2013). However, given the insufficiency of the LEF to generate an adequate *pmf* and the conversion efficiency of the chloroplastic ATP-synthase (ATPase), it has been long recognized that the LEF only generates 85% of the theoretical ATP requirements of the CBB cycle (Allen, 2003).

Cyclic electron flow, which recycles reducing equivalents around PSI (CEF) and pseudo-cyclic electron flow (PCEF), which reduces molecular O_2 at the acceptor side of PSI (**Fig. 1**), were proposed to contribute to the production of extra ATP during photosynthesis (Arnon, 1959, 1984). In most angiosperms and mosses, CEF entails two main pathways, one involving the proton gradient generation (PGR) 5 (Munekage *et al.*, 2002) and PGR-Like 1 proteins (DalCorso *et al.*, 2008) and another involving the plastidial NDH-1 complex (Joe□t *et al.*, 2001; Munekage *et al.*, 2004). In green microalgae, the plastidial NDH-1 complex is absent (Peltier *et al.*, 2016) and it was reported that two CEF pathways operate, one controlled by PGRL1 and PGR5 (Tolleter *et al.*, 2011; Johnson *et al.*, 2014) and the other involving the plastidial NAD(P)H dehydrogenase Nda2 (Jans *et al.*, 2008; Desplats *et al.*, 2009; Saroussi *et al.*, 2016). Since the Nda2 pathway is mostly active during nitrogen deprivation (Peltier and Schmidt, 1991; Saroussi *et al.*, 2016), hereafter CEF refers to the PGRL1/PGR5-controled CEF. PCEF involves Flavodiiron proteins (FLVs) that catalyse O_2 photoreduction at the PSI acceptor side (**Fig. 1**) in cyanobacteria(Helman *et al.*, 2003; Allahverdiyeva *et al.*, 2013), green microalgae (Chaux *et al.*, 2017), mosses (Gerotto *et al.*, 2016), liverwort (Shimakawa *et al.*, 2017) and gymnosperms (Ilík *et al.*, 2017).

Another mechanism that can supply ATP to photosynthesis involves metabolic cooperation between chloroplasts and mitochondria (Raghavendra and Padmasree, 2003). Reducing power generated within the chloroplast is transferred to the mitochondria by metabolic shuttles where it is converted into ATP by the mitochondrial respiratory chain and

might be shuttled back to the cytosol and the chloroplast (**Fig. 1**). Such a cooperation, called here Chloroplast to Mitochondria Electron Flow (CMEF) was first evidenced from the restoration of photoautotrophic growth in a mutant of the green microalgae *Chlamydomonas reinhardtii* (hereafter *Chlamydomonas*) deficient in the plastidial ATPase (Lemaire *et al.*, 1988). Similar cooperation was later suggested from the study of *Chlamydomonas* mutants affected in mitochondrial respiration (Cardol *et al.*, 2003) or in CEF (Dang *et al.*, 2014) as well as in the moss *Physcomytrum patens* (Mellon *et al.*, 2021), and in diatoms (Bailleul *et al.*, 2015).

It is generally considered that both PCEF and CMEF act essentially as electron valves, operating either transiently in the case of PCEF to alleviate excess electrons generated by photosynthesis under light fluctuating regimes (Alric and Johnson, 2017; Chaux *et al.*, 2017; Alboresi *et al.*, 2019) or more continuously for CMEF under high light stress (Kaye *et al.*, 2019). Although in *Chlamydomonas*, CMEF, PCEF and CEF have been recently shown to all contribute to supply energy to the CO_2 concentrating mechanism (CCM) (Burlacot *et al.*, 2022), evidence of their role in powering the CBB cycle and downstream carbon metabolism processes are scarce.

In this work, we show that each pathway (CEF, PCEF and CMEF) has the capacity to sustain net CO_2 fixation to varying degrees and can compensate for the absence of the other pathways, whereas CO_2 fixation is completely abolished in the absence of all three pathways. Moreover, we introduce a new metric: the electron quantum yield of net photosynthesis for each pathway and use it to show that CMEF is by far the most efficient at energizing CO_2 fixation. We further discuss our data in relation to the relative contribution and efficiency of the different alternative pathways to the net primary productivity of terrestrial and oceanic ecosystems.

Results

Net photosynthesis fully relies on mitochondrial respiration when PCEF and CEF are impaired. To test whether PCEF and CEF are required for maximal net photosynthesis (V_{Max}), we first measured photosynthetic O_2 exchange rates using Membrane Inlet Mass Spectrometry (Burlacot *et al.*, 2020) in air-grown *Chlamydomonas* mutants impaired in both CEF and PCEF (*pgrl1 flvB-1 and 3*) and in control strains (WT2 and 3)(Burlacot *et al.*, 2022). Since the *pgrl1* and *flvB* mutants are affected in the functioning of the CO_2 concentration mechanism (CCM) (Burlacot *et al.*, 2022), photosynthesis measurements were performed in the presence of a saturating concentration of inorganic carbon (C_i) when the CCM is inactive. As previously reported (Burlacot *et al.*, 2022), *pgrl1 flvB* double mutants can perform photosynthesis at maximum rates comparable to the single *pgrl1* or *flvB* mutants and to the parental and sibling control strains (**Fig. 2; Sup Figs. S1 A, S2, S3**), therefore showing that neither CEF nor PCEF are essential to reach V_{Max}. To test a possible contribution of mitochondrial respiration to photosynthesis, *pgrl1 flvB* double mutants were treated with myxothiazol (Myxo) and salicylhydroxamic acid (SHAM), inhibitors of mitochondrial electron transport, respectively acting on complex III and the alternative oxidases (AOX) (**Fig. 1**)(Cournac *et al.*, 2002). While the addition of both inhibitors had minimal impact (about 10-20% decrease) on the V_{Max} of control strains (**Fig. 2 A, C, Sup Figs. S1 A, S2, S3**), they completely abolished photosynthesis of *pgrl1 flvB* double mutants (**Fig. 2 B, C, Sup Figs. S1, S2**).

In *Chlamydomonas*, the cellular location of mitochondria depends on the CO_2 supply, with the mitochondria located at the cell periphery in low CO_2-grown cells and mostly inside the chloroplast cup in high CO_2 (Geraghty and Spalding, 1996) (**Fig. 2E, F**). Since this location might affect CMEF (Burlacot and Peltier, 2023), we sought to examine the involvement of the mitochondrial sub-cellular localization on the role of CMEF. Similar to air-grown cells, mitochondrial inhibitors fully stopped net photosynthesis on 2% CO_2-grown *pgrl1 flvB* mutants while having a marginal effect (about 10-20% decrease) on the control strains (**Fig. 2D, Sup Fig. S4**). To control the location of the mitochondria in the different strains, we visualized the spatial distribution of an expressed, mitochondria-targeted fluorophore (Clover). There was similar re-localization of mitochondria depending on the CO_2 supply during growth (at periphery under very low CO_2) of both the single and double mutants and a control strain (**Fig. 2E, F, Sup Fig. S5**). We conclude from this experiment that in the absence of both CEF and PCEF, near wild type levels of photosynthesis can be

achieved that fully rely on CMEF, the role of CMEF being independent of the spatial distribution of the mitochondria.

CEF, PCEF and CMEF each have the potential to sustain most net photosynthesis. We quantified the potential of CEF, PCEF and CMEF to energize net photosynthesis, by measuring the effect of mitochondrial inhibitors on photosynthesis rates in single and double *pgrl1* and *flvB* mutants (**Fig. 3 A, B, C, D**). Inhibition of respiration decreased net photosynthesis by 40-50% in both the *flvB* and *pgrl1* single mutants (**Fig. 3 A, B, Sup Figs. S1 A, S2**), showing that both CEF and PCEF have the potential to energize 50-60% of V_{Max} on their own (**Fig. 3 E, Sup Fig. S1A, B**). Inhibition of respiration decreased the V_{Max} of all control strains by 10-20% (**Fig. S1 A, S2, S3**) showing that combined activities of CEF and PCEF can power approximately 85% of photosynthesis (**Fig. 3 E**). Based on the effects of SHAM or myxothiazol individually, on V_{Max} in *pgrl1 flvB* mutants (**Fig. 3C, D, E, Sup Figs. S1 A, S2**), we conclude that while the complex III-IV pathway of CMEF (**Fig. 1**) can power most net photosynthesis on its own (**Fig. 3D**), the AOX pathway (**Fig. 1**) can only sustain approximately 45% of V_{Max} (**Fig. 3E**). Because the amount of the terminal oxidases involved in CMEF does not differ in the *pgrl1 flvB* mutants and their control strains (**Sup Fig. S6**), these values likely reflect wild type capacities. We also quantified net photosynthesis based on inorganic carbon (C_i) uptake (**Sup Figs. S7, S8**). On all strains tested, the inhibitory effects on V_{max} measured as net C_i uptake was similar to that observed on net O_2 production (**Sup Figs. S7**). We conclude from these experiments that CEF, PCEF and CMEF can sustain photosynthetic C_i uptake on their own but that only the complex III-IV pathway of CMEF is efficient enough to reach near maximal photosynthesis levels.

CEF, PCEF and CMEF show different efficiencies in sustaining net photosynthesis. Mechanistically, for each electron flowing once through the different pathways (CEF, PCEF, the AOX, or the complex III-IV pathway of CMEF), the total number of protons translocated across the thylakoid and/or the inner mitochondrial membranes, highly differ (Burlacot, 2023). To quantify the potential of the different pathways in energizing net photosynthesis, we introduce a new metric, the electron (e^-) quantum yield of a pathway defined by the following equation:

$$e^- quantum\ yield = \frac{e^-(net\ photosynthesis)}{e^-(net\ photosynthesis) + e^-(alternative\ pathway)}$$

Where e⁻ (net photosynthesis) is the flux of photosynthetic electrons directly used for net

photosynthesis and e⁻ (alternative pathway) is the flux of photosynthetic electrons through a

given alternative pathway under conditions where only this alternative pathway is active.

Therefore, the e- quantum yield of an alternative pathway, which is determined when only

this alternative pathway is operating, characterizes the efficiency of electrons passing through

it in powering net photosynthesis. Note that the e⁻ quantum yield, thus does not reflect the rate

or the importance of the pathway considered but rather its capacity to produce ATP for each

electron. We used this equation and experiments from **Fig. 3 A-E** to quantify the e⁻ quantum

yield of each pathway (**Fig. 3 F, Sup Fig. S9**). The e⁻quantum yield of PCEF was estimated at

0.65, whereas that of CMEF at 0.85 (**Fig. 3F**), indicating a better efficiency of CMEF in

driving CO_2 fixation as compared to PCEF. In the absence of an established method to

quantifying CEF in physiologically relevant conditions (Burlacot *et al.*, 2022), CEF quantum

efficiency was estimated to be around 0.55 by hypothesizing that an electron passing once

through CEF is 1.5 time less efficient at generating a *pmf* than if it would pass through PCEF

(Kramer and Evans, 2011; Burlacot, 2023) (**see Materials and Methods**). In the case of

CMEF, the e⁻ quantum yield of the AOX pathway was lower than that of the complex III-IV

pathway (**Fig. 3F**), likely reflecting the proton translocating capacity of each pathway (**Fig.**

1)(Burlacot, 2023). Using the e⁻ quantum yields of each pathway, we then compared the flux

of electrons required to power the same amount of net photosynthesis relative to PCEF (**Fig.**

3G). Strikingly, theoretical values were in accord with values experimentally determined by

the e⁻ quantum yields (**Fig. 3 G**). We conclude from these experiments that CMEF pathways

are more efficient than PCEF and CEF at energizing CO_2 fixation and that this difference

arises from a higher H^+/e^- transfer capacity likely leading to a higher capacity to generate

ATP.

Estimating the respective contributions of CEF, PCEF and CMEF during steady state

photosynthesis and growth. We then sought to estimate (*i*) the electron flux via each

pathway expressed as an additional electron flow relative to net photosynthesis (*%-LEF*) and

(*ii*) their contribution to energize net photosynthesis. We used two different approaches, one

based on the effect of mutations, the other based on the effect of respiratory inhibitors. As

both approaches are based on pathway inhibition, they are potentially biased by the high level

of redundancy existing between the different AEFs and the existence of compensatory

mechanisms (Cardol *et al.*, 2009; Dang *et al.*, 2014; Burlacot *et al.*, 2022). The two

approaches differ, however, in the way compensatory mechanisms likely contribute, mutants

being more subject to compensation than short-terms effects of inhibitors. In the mutant-based

approach, PCEF was quantified by subtracting O_2 uptake rates in the *flvB* mutant from its

control parental strain (allowing the calculation of $PCEF_{mut}$ and $CMEF_{mut}$) (**Sup Fig. S10**). In

the inhibitor-based approach, CMEF was quantified by subtracting O_2 uptake rates between

SHAM plus Myxo treated and untreated control strains (allowing the calculation of $PCEF_{inhib}$

and $CMEF_{inhib}$) (**Sup Fig. S10**). The two methods differ somewhat in their estimates of PCEF

and CMEF (**Fig. 4 A, B, Sup Fig. S11**), with a higher CMEF contribution being estimated

using the mutant method. On average, PCEF and CMEF represent a flux of between 10-20*%-*

LEF and 5-10*%-LEF*, respectively (**Fig. 4 A**). Thanks to the respective e⁻ quantum yield of

each pathway, we can estimate that 10-30% and 30-60% of CO_2 fixation is sustained by

PCEF and CMEF respectively (**Fig. 4 B**). Since CEF sustains the remaining net

photosynthesis, we estimated that 25-30% of CO_2 fixation is powered by CEF (**Fig. 4 B**),

which corresponds to a CEF of about 25*%-LEF* assuming an e⁻ quantum yield of CEF of 0.55

(**Fig. 4 A**).

To further assess the contribution of CEF, PCEF and CMEF over the long term,

growth tests were performed on the different strains in the absence or presence of the

respiration inhibitor Myxo. While Myxo has no effect on the growth of control strains and

single *flvB* or *pgrl1* mutants in moderate light intensities (**Fig. 4 C**), growth of *pgrl1 flvB* was

impaired in the presence of the inhibitor (**Fig. 4 C**). Interestingly, at low light intensity (30

μmol photons m⁻² s⁻¹), the effect of Myxo on the double mutant was absent (**Sup Fig. 12 A**),

suggesting that the AOX pathway of CMEF has the potential to fully power net

photosynthesis in low light conditions. Therefore, whereas CEF, PCEF, and CMEF all

contribute to energizing photosynthesis, CMEF, although representing a relatively small flux,

has the potential to fuel CO_2 fixation and biomass accumulation thanks to its high e⁻ quantum

yield.

Discussion

Since the first description of CEF by Arnon in 1959 (Arnon, 1959), the question was raised as

to whether LEF could sustain the ATP requirement of photosynthetic CO_2 fixation (Arnon,

1959), and it was proposed that photosynthetic electron flows, other than LEF, are involved in

balancing the ATP/NADPH ratio to sustain CCB cycle function (Arnon, 1959; Allen, 2003).

Since then, several mechanisms of CEF, PCEF and CMEF have been uncovered (Munekage

et al., 2002; Helman *et al.*, 2003; Munekage *et al.*, 2004; Tomaz *et al.*, 2010) as well as some

compensations between those different mechanisms (Cardol *et al.*, 2009; Petroutsos *et al.*,

2009; Dang *et al.*, 2014; Storti *et al.*, 2019; Storti *et al.*, 2020; Burlacot *et al.*, 2022), but the

importance and relative efficiency of each mechanism remain mostly unknown. Here we have

shown that in *Chlamydomonas* PGRL1-controlled CEF, FLV-mediated PCEF, and both the

AOX and complex III/IV/cytochrome oxidase pathways of CMEF can all supply additional

energy to complement LEF for CO_2 fixation. The e^- quantum yields of CEF, PCEF and

CMEF greatly differ; CMEF is the most electron efficient and contributes importantly to the

extra ATP requirements of CO_2 fixation in continuous, saturating light. Note, that in spite of

the ongoing controversy on how PGRL1 is involved in CEF (Nawrocki *et al.*, 2019), a clear

integration is observed between PGRL1-controled CEF, PCEF and CMEF in which these

different pathways can compensate each other in powering photosynthesis. Importantly, we

observed similar effects on net O_2 and CO_2 exchange (**Sup Fig. S7**), thus showing that all

effects measured on net photosynthesis correspond to changes in CO_2 fixation and eventually

biomass production and growth (**Fig. 4 C**).

Some compensation between CEF, PCEF, and CMEF has been previously reported; a

Chlamydomonas PGRL1-deficient strain exhibits increased CMEF and PCEF (Dang *et al.*,

2014) and mutants of complexes I and III show enhanced rates of CEF (Cardol *et al.*, 2009).

We observed an elevated CMEF in the absence of FLV (**Fig. 4 A, B**) with a higher level of

terminal oxidases associated with CMEF (**Sup Fig. 6**), corroborating cross-compensation

among all three mechanisms of electron flow. In diatoms, while CMEF is considered to be a

major driver of CO_2 fixation, inhibition of CMEF only leads to a ~20% decrease of the PSII

yield (Bailleul *et al.*, 2015), which might also reflect compensation by CEF. We also

conclude, based on the fact that no CO_2 fixation could be detected in the double *pgrl1 flv*

mutants treated with inhibitors of mitochondrial respiration, that neither the NDA2-mediated

CEF, nor O_2 reduction mediated by the plastid terminal oxidase or Mehler reactions have the

potential by themselves to sustain a measurable amount of net photosynthesis, which is in

agreement with previous studies concluding that these pathway represent a small fraction of

alternative electron flow (Alric *et al.*, 2010; Houille-Vernes *et al.*, 2011; Nawrocki *et al.*,

2019)(**Fig. 2 B**).

The quantitative contribution of each pathway in the wild-type strain in the absence of

perturbation of any AEFs is a challenging task to which the approaches we have developed

here provide an imperfect answer. Indeed, the quantification of each pathway we have

proposed relies on the assumption that inhibiting one pathway (CMEF or PCEF) does not

affect the flow through the others, assumption which is most likely not verified since

compensation between different AEFs mechanisms have been previously documented (Cardol

et al., 2009; Dang *et al.*, 2014; Burlacot *et al.*, 2022). It should therefore be considered as a first attempt to evaluate the respective contributions of each pathway until an integrative quantification method is available. Here, the mutant method likely overestimates CMEF and the inhibitor method likely overestimates PCEF, both approaches likely overestimating CEF. It is clear however, that no single pathway sustains net photosynthesis by itself (**Sup Fig. S11 B**) which might reflect a rather balanced contribution of the different pathways to CO_2 fixation in the WT. In any case, it seems highly likely that, considering the high redundancy of AEFs and the ability of each of them to sustain CO_2 fixation at a high rate, a dynamic mix of the different pathways may be involved in different environmental conditions, each pathway being more or less advantageous in specific conditions (Burlacot, 2023).

The e^- quantum yield, a new tool to evaluate the efficiency of electron transport pathways.

In support to LEF, a combination of PGRL1-controled CEF and PCEF cannot sustain more than 80% of maximal CO_2 fixation (**Fig. 3 E**) whereas each individual flux can help sustain 60% of it (**Fig. 3 F**). This suggests the existence of an intrinsic limitation of the flow of electrons through the photosynthetic electron transport chain, which is likely due to a *pmf*-induced photosynthetic control occurring at the cytochrome b_6f level (Johnson and Berry, 2021; Malone *et al.*, 2021); an elevated *pmf* eliciting a block in electron flow across the b_6f complex. Engaging an electron pathway with a high e^- quantum yield, such as CMEF, might be a way to avoid this limitation since less electrons would be needed to power the same level of CO_2 fixation. Such a strategy seems to be used by diatoms (Bailleul *et al.*, 2015) which lacks FLVs.

Throughout the green lineage, the nature and activity of alternative electron flows are quite different. For example, angiosperms lack FLV-dependent PCEF (Peltier *et al.*, 2010), harbour a NDH-1-dependent CEF pathway (Peltier *et al.*, 2016) and have a weakly active CMEF (Raghavendra and Padmasree, 2003). Since the NDH-1 dependent CEF translocates $4H^+/e^-$ (Strand *et al.*, 2017), its e^- quantum yield is close to that of CMEF, strongly suggesting that in angiosperms the NDH-1 complex can compensate when the CMEF capacity is diminished. The latter is illustrated by the poor growth of *Arabidopsis thaliana* mutants impaired in both CEF pathways (PGR and NDH-1) (Munekage *et al.*, 2004). Because of the number of key proteins and biochemical reactions involved in CMEF, its high e^- quantum yield may not be as rapidly accessible as CEF or PCEF during environmental fluctuations. While under light limiting conditions a high e^- quantum yield pathway might be advantageous, a less efficient pathway would contribute to the safe dissipation of excess

absorbed light energy and diminished reactive oxygen production at saturating light intensities. Overall, it remains to be determined to what extent harbouring a palette of alternative electron flows with varying quantum yields and likely different time responses, is critical for sustaining photosynthetic performance under dynamic conditions in nature.

Note here that while modulation of the capacity of membranes to transform *pmf* into ATP may modulate the absolute e$^-$ quantum yield of each pathway, the relative e$^-$ quantum yield of CEF and PCEF would not change if they are embedded in the same thylakoid system. Similarly, even in conditions where the mitochondrial respiration would be partially or fully uncoupled, the e$^-$ quantum yield of CMEF would be at least as good as PCEF due to the generation of a *pmf* across the thylakoid membranes (Burlacot, 2023). Hence, if there is no electron transport chain dedicated to a single pathway, electrons passing through CMEF and PCEF should always be more efficient than CEF in powering net photosynthesis.

Additional energy cost of photosynthesis.

Allen previously estimated that a PCEF representing only 16%*-LEF* would be sufficient, in addition to LEF, to satisfy the ATP and NADPH requirement of the CBB cycle (Allen, 2002). Our conclusions are strikingly different since we estimate that if only PCEF was active, it would have to represent 54%*-LEF* to sustain CO_2 fixation. This discrepancy may reflect the fact that the actual metabolic cost of CO_2 fixation exceeds the ATP requirement of the CBB cycle. This may result from the involvement of additional ATP-consuming metabolic pathways or transporters, or to the presence of proton leakages from energized membranes. In addition, in natural environments, either terrestrial or aquatic, CO_2 fixation is further limited by the concentration of atmospheric CO_2, which can trigger photorespiration in C_3 plants or induction of the carbon concentration mechanisms in C_4 plants and algae. All these mechanisms would require additional energy for their activities (Osmond, 1981; Takabayashi *et al.*, 2005; Walker *et al.*, 2014; Burlacot *et al.*, 2022), making alternative electron pathways of photosynthesis even more critical for sustaining plant and algal CO_2 capture.

Acknowledgements

The authors acknowledge the constructive comments on the manuscript from Dr. Joe Berry, Dr. Jennifer Johnson, Evan Saldivar, Leron Perez and technical support from Michel Philibert and Emmanuel Capra. This work was supported by the Carnegie Institution for Science (AB, AG), and the ANR grants OTOLHYD n°ANR-18-CE05-0029 (GP) and AlgalCCM n°ANR-

335   22-CE44-0023-01 (GP, AM). This work was also in part supported by DOE award DE-
336   SC0019417 to AG.

Figure legend:

Figure 1. Electron transport pathways operating during oxygenic photosynthesis and
their role in ATP generation. Linear electron flow (LEF) operates from PSII to PSI and
reduces $NADP^+$ to NADPH using electrons from water. It also generates a *pmf* across the
thylakoid membrane composed of a proton gradient (ΔpH) and electrical gradient ($\Delta\Psi$); the
pmf is used by the ATPase to generate ATP. Cyclic electron flow (CEF) and pseudo cyclic
electron flow (PCEF) both contribute to the generation of the *pmf* but do not produce reducing
equivalents since the e^- used for PCEF reduce O_2 to water. Chloroplast to mitochondria
electron flow (CMEF) encompasses multiple metabolic pathways by which reductants
generated in the chloroplast is shuttled to the mitochondria and converted into ATP by the
mitochondrial electron transport chain. NADPH and ATP generated by these different
pathways are eventually used for transforming CO_2 into biomass, which reflects net
photosynthesis. Other abbreviations: AOX, alternative oxidase; ATP, adenosine triphosphate;
CBB, Calvin Benson Bassham; FDX, ferredoxin; FLVs, flavodiiron proteins; FNR,
ferredoxin-$NADP^+$ reductase; Myxo, myxothiazol; NADPH, reduced form of nicotinamide
adenine dinucleotide phosphate; PS, photosystem; PGR5/L1, proton gradient generation
5/like-1; SHAM, salicyl hydroxamic acid.

Figure 2. Dependence of net photosynthesis on CEF, PCEF and CMEF. A, B, O_2
exchange rates upon a dark to light transition in WT2 (**A**) and *pgrl1 flvB-1* (**B**). At t=6min,
mitochondrial respiration inhibitors myxothiazol ($2.5\,\mu M$) and salicylhydroxamic acid
(SHAM, $400\,\mu M$) were added to the reaction. Shown are representative traces of $n=3$
biologically independent samples. **C and D**, Net O_2 evolution in *pgrl1 flvB-1* and *pgrl1 flvB-*3
and their control strains (WT2 and WT3) in the absence (control) or presence (Myxo &
SHAM) of myxothiazol ($2.5\,\mu M$) and salicylhydroxamic acid (SHAM, $400\,\mu M$) in cells
grown in air (**C**) or in 2% CO_2-enriched air (**D**). Values are taken after 9 min of illumination
upon a dark to light transitions. Bars show the average and dots show individual replicates
($n=3$ biologically independent samples). Typical experiments used to extract these data
are shown in **Sup. Fig. S2**. **E** and **F**, Cross section images showing subcellular localization of
mitochondria-localized Clover (yellow fluorescence) in *pgrl1 flvB-1* and a control strain
(WT2), grown in air (**E**) or 2% CO_2-enriched air (**F**). Images are representative of 2
independent experiments. Scale bar: 10 µm. Brightfield, Clover fluorescence, Chlorophyll
autofluorescence and Clover/Chlorophyll merge signals are shown. Images are representative
of 2 experiments. Scale bar: 10 µm.

Figure 3. Relative efficiencies of CEF, PCEF and CMEF in energizing net
photosynthesis. A-D, Net O_2 evolution measured during a dark to light transition in *flvB* (**A**),
pgrl1 (**B**), *pgrl1 flvB-2* (**C**, **D**), treated (gray dots) or not treated (black dots) with both
myxothiazol (Myxo, $2.5\,\mu M$) and salicylhydroxamic acid (SHAM, $400\,\mu M$) (**A, B**), or with
only one inhibitor (**C,D**). Shown are representative traces of 3 biological replicates. **E**,
Relative V_{max} measured when a given set of alternative electron pathways are active after
treatment with myxothiazol (Myxo, $2.5\,\mu M$), salicylhydroxamic acid (SHAM, $400\,\mu M$) or

both inhibitors; CEF (blue, *flvB* treated with Myxo and SHAM), PCEF (green, *pgrl1* treated
with Myxo and SHAM), CEF&PCEF (dark, control strains treated with Myxo and SHAM),
CMEF via AOX (purple, *pgrl1 flvB* treated with Myxo), CMEF via III/IV (red, *pgrl1 flvB*
treated with SHAM). **F,** Electron quantum yield of PCEF (green), CMEF via AOX (purple),
CMEF via III/IV (red), CMEF (pink) and CEF (dotted line). Similar electron quantum yields
were also determined for PCEF in 2% CO_2-grown cells and a slightly lower quantum yield for
CMEF (**Sup Fig. S9 D**). **G**, Electron flux relative to PCEF required to power similar levels of
net photosynthesis when relying on a single pathway. Data shown are individual
extrapolations based on measured electron quantum yields. In **E, F** and **G,** bars show the
average and dots show individual replicates ($n\,≥3$ biologically independent sample).
391

Figure 4. Contribution of CEF, PCEF and CMEF to the electron flow, net
photosynthesis and growth. A Electron flow through PCEF, CEF and CMEF relative to LEF
were determined in the wildtype strain CC-4533 by two different methods, one based on the
use of inhibitors (inhib) the other on mutants (mut). Data used for these calculations are the
same as for **Fig. 2 C** and **Sup Fig. 2**. **B** Proportion of net photosynthesis powered by each
electron flow (PCEF, CEF and CMEF) in the control strain CC-4533 as measured by the
inhibition (inhib) or mutant (mut) method. CC-4533 was chosen because it is the control
strain for *flvB* and is therefore the only strain where both methods could be used. Data used
for these calculations are the same as for **A**. In **A** and **B,** bars show the average and dots show
individual replicates ($n=3$ biologically independent sample). **C**, Growth tests
for *flvB* and *pgrl1* mutants, their corresponding control strains (CC-4533 and 137AH,
respectively) and double mutants (*pgrl1 flvB-1 and pgrl1 flv-3*) and their corresponding
control strains (WT2 and 3). Cells were spotted on plates containing minimal medium at pH
7.2 and grown under continuous illumination (60 µmol photon m^{-2} s^{-1}) in the absence (upper
panel) or presence (lower panel) of 2.5µM Myxothiazol (Myxo) in air enriched at 2% CO_2.
Spots shown are representative of 10 independent experiments.

Material and Methods

Strains and growing conditions. *Chlamydomonas flvB* and *pgrl1* single mutants and their respective parental strains CC-4533 (for *flvB*), 137AH (for *pgrl1*) where previously described(Tolleter *et al.*, 2011; Chaux *et al.*, 2017). *flvB pgrl1*-1, -2 and -3 double mutant and their respective sibling strains WT2 and WT3 (used here as control strains) have been described previously(Burlacot *et al.*, 2022) and are all from the progeny of a crossing of *flvB* and *pgrl1* performed in(Burlacot *et al.*, 2022). Because *pgrl1 flvB* double mutants were obtained by crossing the two single mutants which have different genetic backgrounds (as described in (Burlacot *et al.*, 2022)), the control strains used are siblings that resulted from the cross and that contained wild-type copies and levels of the FLVB and PGRL1 proteins. *pgrl1-2* is a mutant impaired in accumulation of the PGRL1 protein originating from crossing *flvB* and *pgrl1* performed in (Burlacot *et al.*, 2022). All strains were grown photo-autotrophically in 125-mL flasks with cotton stoppers at 25 °C in a buffered minimal medium (20 mM MOPS pH 7.2) under constant illumination (80 µmol photons m^{-2} s^{-1}) either under ambient CO_2 or 2% CO_2 in air. Amongst *pgrl1 flvB* double mutants and their control strains already described (Burlacot *et al.*, 2022), *pgrl1 flvB*-1 and *pgrl1 flvB*-3 and WT2 and 3, respectively, were used for physiological measurements since their maximal photosynthesis was representative of other double mutants and control strains (Burlacot *et al.*, 2022). All strains used in this study are available under the same names used here (except for *flvB* which is available as *flvB-21*) at the Chlamydomonas Resource Center (www.chlamycollection.org). Spectrum of the growing light used are shown in **Sup Fig. 13**.

Gas exchange measurements. Gas exchange rates were measured using membrane inlet mass spectrometry (MIMS). Cells grown in air enriched or not at 2% CO_2 were collected during the exponential phase by centrifugation at 450 g for 3 min and resuspended in 1.5 mL of fresh buffered minimal medium (pH 7.2) at 30 µg chlorophyll mL^{-1}. HCO_3^- was subsequently added to the cell suspension (10 mM final concentration). The cell suspension was then placed in the MIMS reaction vessel, 2 mL of ^{18}O-enriched O_2 (97% ^{18}O Eurisop, France) were bubbled in the suspension. After closing the vessel, gas exchange was recorded during a dark to light transition (2,000 µmol photons m^{-2} s^{-1}; green LEDs picked at 525nm; Luminus reference PT-121-G-L11-MPK). Light intensity was set to be saturating (**Sup Fig. 14**) using weakly absorbed green light for better homogeneity. Calculations of gross and net photosynthesis were done using the MIMS analysis software as previously described (Burlacot *et al.*, 2020). In some experiments salicyl hydroxamic acid (SHAM, 400 µM final

concentration) and/or myxothizol (Myxo, 2.5 µM final concentration) were added 1 min

before the beginning of the measurements, except for the data presented in **Fig. 2A**, **B** where

inhibitors were added during the measurement. Note here that SHAM has been reported to not

inhibit the plastid terminal oxidase (Cournac *et al.*, 2000). All Net O_2, Net CO_2, e⁻ quantum

yield, e⁻ flow and Net photosynthesis sustained shown as histograms in **Figs. 2, 3, 4** and **Sup.**

Figs. 1, 7, 9, 11 are based on measurements after 9 minutes of illumination.

Clover Expressing cells. 137AH, WT2, *pgrl1*, *flvB*, *pgrl1flvB*-1 and -2 expressing the

mitochondria targeted fluorophore Clover (mito-Clover) were generated as described in

(Harmon *et al.*, 2022). Briefly, cells were transformed with a linearized plasmid encoding

mito-Clover and a hygromycin-resistance cassette using the Max Efficiency Reagent

(Invitrogen). Resistant clones to hygromycin were screened for Clover fluorescence using a

M1000 Tecan plate reader (Excitation: 488/7 nm; Emission: 515/10 nm) and mitochondrial

signal was checked by confocal microscopy. Selected clones were grown at 80 µmol photons

m^{-2} s^{-1} in 25 mL minimal medium in 125 mL flasks, in the presence of either normal or 2%

CO_2-enriched air until they reached a density of ~10 µg chlorophyll mL^{-1}. An aliquot of each

culture was then briefly concentrated by centrifugation at 1,500g and cells layered on a 18-

well microscope slide. Images were acquired using a TCS SP8 confocal laser-scanning

microscope (Leica) and a ×63 objective. Acquisition settings were as follows: Clover

excitation 488 nm, emission 500-547 nm; Chlorophyll excitation 488 nm, emission 694-733

nm. For homogeneity and clarity, signal intensities of each channel were adjusted using the

Fiji software.

Growth tests. The different *Chlamydomonas* strains were cultivated at air level of CO_2 under

moderate light (80 µmol photons m^{-2} s^{-1}) at 25 °C. Cells were collected during exponential

growth and diluted in fresh minimal medium to 0.125, 0.625 or 2.5☐10^6 cells mL^{-1}. Seven-

microlitre drops were spotted on 2% Agar plates of minimal medium (buffered with 20mM

MOPS pH☐7.2) with or without myxothiazol (2.5µM final concentration) and exposed to air

enriched with 2% CO_2. Homogeneous light was supplied by LED panels (see its spectrum in

Sup Fig. 13 B), the different light intensities were obtained with various neutral filters.

Temperature was maintained at 25☐°C at the level of plates by means of fans.

Electron quantum yield calculation. The e⁻ quantum yield of a given alternative electron

pathway is defined as the proportion of electrons generated downstream PSI that are used for

net photosynthesis when only this alternative electron pathway is active, following this

equation:

$$\text{(1)} \quad e^- \, quantum \, yield = \frac{e^-(net\,photosynthesis)}{e^-(net\,photosynthesis) + e^-(alternative\,pathway)}$$

Where e⁻(net photosynthesis) is the flux of photosynthetic electrons directly used as reductants for net photosynthesis and e⁻(alternative pathway) is the flux of photosynthetic electrons through an alternative pathway. Because for PCEF and CMEF the sum of e⁻(alternative pathway) and e⁻(net photosynthesis) equals gross photosynthesis, electron quantum yields of PCEF and CMEF were quantified using gross and net O_2 evolution in mutants harbouring: only PCEF (*pgrl1* and *pgrl1-2* mutants treated with Myxo and SHAM), or only the AOX or the complex III/IV pathway of CMEF (*pgrl1 flvB*-1 and 3 mutants treated with Myxo or SHAM, respectively). The electron quantum yield was then calculated for each pathway using the following equation:

$$\text{(2)} \quad e^- \, quantum \, yield = \frac{net\,O_2\,evolution}{gross\,O_2\,evolution}$$

The theoretical electron quantum yield of CEF was calculated based on the assumption that CEF translocates 1.5 less protons per electron across the thylakoid membranes than PCEF (Kramer and Evans, 2011; Burlacot, 2023) and that this is the driving force leading to net photosynthesis energization. Note that this later assumption is well supported by the independent results from **Fig. 3 G** showing that the relative electron quantum yield of CMEF pathways and PCEF corresponds to their relative capacity at translocating protons across membranes. Hence, for the same net photosynthesis powered, CEF should be 1.5 times higher than PCEF, and the electron quantum yields of CEF could be written as:

$$\text{(3)} \quad e^- \, quantum \, yield \, (CEF) = \frac{e^-(net\,in\,pgrl1\,Myxo\,SHAM)}{e^-(net\,in\,pgrl1\,Myxo\,SHAM) + e^-(PCEF) \times 1.5}$$

Where e⁻(net in *pgrl1* Myxo SHAM) is the net photosynthesis measured in the *pgrl1* mutant treated with Myxo and SHAM (hence having only PCEF active) and e⁻ (PCEF) being the O_2 uptake rate in the same strain and condition.

Based on equation (1), we can rewrite e⁻(PCEF) as:

$$\text{(4)} \quad e^-(PCEF) = \frac{(1 - e^- quantum\,yield(PCEF))}{e^- quantum\,yield(PCEF)} \times e^-(net\,in\,pgrl1\,Myxo\,SHAM)$$

Hence, combining (3) and (4):

$$\text{(5)} \quad e^- \, quantum \, yield \, (CEF) = \frac{e^- quantum\,yield\,(PCEF)}{1.5 - \frac{e^- quantum\,yield\,(PCEF)}{2}}$$

The electron quantum yield of CEF shown in **Fig. 3** is using equation (5) with the average electron quantum yield of PCEF. Note that the quantifications of the e^- quantum yield in *Chlamydomonas* assumes that removing two pathways (elimination of CEF and CMEF in the *pgrl1* mutant treated with inhibitors of respiration, and CEF and PCEF in the *pgrl1 flvB* double mutants) makes CO_2 fixation dependent only on the third pathway. This dependency is supported by the finding that the *pgrl1 flvB* double mutant treated with inhibitors of respiration are unable to support any CO_2 fixation, which also support the hypothesis that the contribution of other pathways, such as the NDA2-dependent CEF, is marginal.

Calculation of electron flows via each pathway and their contribution to net photosynthesis in wildtype strains. The electron flow through PCEF and CMEF were deduced on wildtype strain using two independent methods based on measurements of gross O_2 uptake made in **Sup. Fig. 2 and 3**. In the so called "mutant" method, CMEF was quantified as being the gross O_2 uptake measured in the *flvB* mutant. Then, PCEF was quantified by subtracting the CMEF to the O_2 uptake of CC-4533 (**Sup. Fig. S10**). This method was only used for CC-4533 since it is the only control strain for the *flvB* mutant. In the so called "inhibitor" method, PCEF was quantified as being the gross O_2 uptake remaining in control strains treated with SHAM and Myxo. Then, CMEF was quantified by subtracting the PCEF to the O_2 uptake in the same untreated control strain (**Sup. Fig. S10**). The calculated electron flows of PCEF and CMEF were then used for each strain to quantify the net photosynthesis powered by each of them following the equations derived from (1):

$$net\ photoynthesis\ (AEF) = e^-(AEF) \times \frac{e^-quantum\ yield\ (AEF)}{1 - e^-quantum\ yield\ (AEF)}$$

Where net photosynthesis (AEF) is the net photosynthesis sustained by a specific alternative electron flow, e^-(AEF) being the electron flow through this AEF and e^- quantum yield (AEF) being the electron quantum yield of this alternative electron flow.

The total of net photosynthesis powered in a strain by PCEF and CMEF was then compared to the measured net photosynthesis in this same strain to quantify the "missing" net photosynthesis that had to be powered by CEF. Then the electron flow through CEF (e^-(CEF)) was calculated by the following equation derived from (1):

$$e^-(CEF) = missing\ net\ \times \frac{1 - e^-quantum\ yield\ (CEF)}{e^-quantum\ yield\ (CEF)}$$

Where "missing net" is the calculated net photosynthesis that is not powered by PCEF and

CMEF and e⁻ quantum yield (CEF) being the electron quantum yield of CEF.

Protein extraction and Immunodetection. Cells were harvested by centrifugation at 1,000g

for 5 minutes. Total cellular protein was extracted by heating the cell pellet at 100°C for one

minute using an extraction aqueous solution composed of 1% SDS, 100 mM DTT, 1mM

Aminocaproic acid, and 1mM benzamidine hydrochloride buffered with 60mM Tris (pH 6.8).

The resulting cell lysate was then centrifuged at 10,000 g for 5 minutes to obtain the

supernatant. The supernatant was subsequently incubated with acetone (80% final

concentration) at -20°C for 1 hour. The protein pellet was collected by centrifugation at 5,000

g for 5 minutes and stored in aqueous solution containing 1% SDS, 5 mM DTT and buffered

with 60mM Tris (pH 6.8). Protein content was estimated (Microplate BCA protein assay Kit-

reducing agent compatible, Thermo Scientific) with bovine serum albumin as the standard.

Protein extracts (10 μg protein) were loaded on a 12% SDS-PAGE gel and run for 1h at

150V in Tris-glycine SDS buffer and transferred onto a nitrocellulose membrane using semi-

dry transfer (Transblot turbo, Biorad). Immunoblot detection was performed using antibodies

raised against FlvB (1:1,000)(Chaux *et al.*, 2017), and Pgrl1 (1:5,000)(Tolleter *et al.*, 2011).

Other antibodies against AOX1 (1:10,000) (AS06 152), COXIIB (1:10,000) (AS06 151) were

obtained from Agrisera (https://www.agrisera.com/). An anti-rabbit HRP-conjugated antibody

(1:50,000) (AS09 602) was used as a secondary antibody for immunodetection. Precision Plus

Protein Dual Color Standards (Biorad) was used as the molecular weight marker. Uncropped

and unprocessed scans of the blots shown in this study are visible in source data.

Data Availability. Genes studied in this article can be found on https://phytozome-

next.jgi.doe.gov/ under the loci Cre12.g531900 (FLVA), Cre16.g691800 (FLVB),

Cre07.g340200 (PGRL1), Cre09.g395950 (AOX1), Cre03.g169550 (AOX2) and

Cre01.g049500 (COXIIB).

References

Alboresi A, Storti M, Morosinotto T (2019) Balancing protection and efficiency in the
regulation of photosynthetic electron transport across plant evolution. New Phytol.
221: 105-109 10.1111/nph.15372

Allahverdiyeva Y, Mustila H, Ermakova M, Bersanini L, Richaud P, Ajlani G, Battchikova N, Cournac L, Aro EM (2013) Flavodiiron proteins Flv1 and Flv3 enable cyanobacterial growth and photosynthesis under fluctuating light. Proc. Natl. Acad. Sci. U.S.A. **110:** 4111-4116 10.1073/pnas.1221194110

Allen JF (2002) Photosynthesis of ATP—Electrons, Proton Pumps, Rotors, and Poise. Cell **110:** 273-276 https://doi.org/10.1016/S0092-8674(02)00870-X

Allen JF (2003) Cyclic, pseudocyclic and noncyclic photophosphorylation: new links in the chain. Trends Plant Sci. **8:** 15-19 10.1016/s1360-1385(02)00006-7

Alric J, Johnson X (2017) Alternative electron transport pathways in photosynthesis: a confluence of regulation. Curr. Op. Plant. Biol. **37:** 78-86 https://doi.org/10.1016/j.pbi.2017.03.014

Alric J, Lavergne J, Rappaport F (2010) Redox and ATP control of photosynthetic cyclic electron flow in *Chlamydomonas reinhardtii* (I) aerobic conditions. Biochim. Biophys. Acta **1797:** 44-51

Arnon DI (1959) Conversion of light into chemical energy in photosynthesis. Nature **184:** 10-21 10.1038/184010a0

Arnon DI (1984) The discovery of photosynthetic phosphorylation. Trends Plant Sci **9:** 258-262 https://doi.org/10.1016/0968-0004(84)90159-2

Bailleul B, Berne N, Murik O, Petroutsos D, Prihoda J, Tanaka A, Villanova V, Bligny R, Flori S, Falconet D, Krieger-Liszkay A, Santabarbara S, Rappaport F, Joliot P, Tirichine L, Falkowski PG, Cardol P, Bowler C, Finazzi G (2015) Energetic coupling between plastids and mitochondria drives CO_2 assimilation in diatoms. Nature **524:** 366 10.1038/nature14599

Burlacot A (2023) Quantifying the roles of algal photosynthetic electron pathways: a milestone towards photosynthetic robustness. New Phytol. **n/a**https://doi.org/10.1111/nph.19328

Burlacot A, Burlacot F, Li-Beisson Y, Peltier G (2020) Membrane Inlet Mass Spectrometry: A Powerful Tool for Algal Research. Front. Plant. Sci. **11**10.3389/fpls.2020.01302

Burlacot A, Dao O, Auroy P, Cuiné S, Li-Beisson Y, Peltier G (2022) Alternative photosynthesis pathways drive the algal CO_2 concentrating mechanism. Nature https://doi.org/10.1038/s41586-022-04662-9

Burlacot A, Peltier G (2023) Energy crosstalk between photosynthesis and the algal CCM. Trends Plant Sci. https://doi.org/10.1016/j.tplants.2023.03.018

Cardol P, Alric J, Girard-Bascou J, Franck F, Wollman FA, Finazzi G (2009) Impaired respiration discloses the physiological significance of state transitions in *Chlamydomonas*. **106:** 15979-15984 10.1073/pnas.0908111106

Cardol P, Gloire G, Havaux M, Remacle C, Matagne R, Franck F (2003) Photosynthesis and state transitions in mitochondrial mutants of *Chlamydomonas reinhardtii* affected in respiration. Plant Physiol. **133:** 2010-2020 10.1104/pp.103.028076

Chaux F, Burlacot A, Mekhalfi M, Auroy P, Blangy S, Richaud P, Peltier G (2017) Flavodiiron proteins promote fast and transient O_2 photoreduction in Chlamydomonas. Plant Physiol. **174:** 1825-1836 10.1104/pp.17.00421

Cournac L, Latouche G, Cerovic Z, Redding K, Ravenel J, Peltier G (2002) *In vivo* interactions between photosynthesis, mitorespiration, and chlororespiration in *Chlamydomonas reinhardtii*. Plant Physiol. **129:** 1921-1928 10.1104/pp.001636

Cournac L, Redding K, Ravenel J, Rumeau D, Josse E-M, Kuntz M, Peltier G (2000) Electron flow between photosystem II and oxygen in chloroplasts of photosystem I-deficient algae is mediated by a quinol oxidase involved in chlororespiration. **275:** 17256-17262

DalCorso G, Pesaresi P, Masiero S, Aseeva E, Nemann DS, Finazzi G, Joliot P, Barbato R, Leister D (2008) A complex containing PGRL1 and PGR5 is involved in the switch between linear and cyclic electron flow in *Arabidopsis*. Cell **132:** 273-285 10.1016/j.cell.2007.12.028

Dang KV, Plet J, Tolleter D, Jokel M, Cuine S, Carrier P, Auroy P, Richaud P, Johnson X, Alric J, Allahverdiyeva Y, Peltier G (2014) Combined increases in mitochondrial cooperation and oxygen photoreduction compensate for deficiency in cyclic electron flow in *Chlamydomonas reinhardtii*. Plant Cell **26:** 3036-3050 10.1105/tpc.114.126375

Desplats C, Mus F, Cuine S, Billon E, Cournac L, Peltier G (2009) Characterization of Nda2, a plastoquinone-reducing type II NAD(P)H dehydrogenase in *Chlamydomonas* chloroplasts. J. Biol. Chem. **284:** 4148-4157 10.1074/jbc.M804546200

Geraghty AM, Spalding MH (1996) Molecular and structural changes in *Chlamydomonas* under limiting CO_2 (a possible mitochondrial role in adaptation). **111:** 1339-1347 10.1104/pp.111.4.1339

Gerotto C, Alboresi A, Meneghesso A, Jokel M, Suorsa M, Aro E-M, Morosinotto T (2016) Flavodiiron proteins act as safety valve for electrons in Physcomitrella patens. Proc. Nat. Acad. Sci. U.S.A. **113:** 12322-12327 10.1073/pnas.1606685113

Harmon J, Findinier J, Ishii NT, Herbig M, Isozaki A, Grossman A, Goda K (2022) Intelligent image-activated sorting of *Chlamydomonas reinhardtii* by mitochondrial localization. Cytometry Part A **101:** 1027-1034 https://doi.org/10.1002/cyto.a.24661

Helman Y, Tchernov D, Reinhold L, Shibata M, Ogawa T, Schwarz R, Ohad I, Kaplan A (2003) Genes encoding a-type flavoproteins are essential for photoreduction of O_2 in cyanobacteria. Curr. Biol. **13:** 230-235

Houille-Vernes L, Rappaport F, Wollman F-A, Alric J, Johnson X (2011) Plastid terminal oxidase 2 (PTOX2) is the major oxidase involved in chlororespiration in *Chlamydomonas*. Proc. Nat. Acad. Sci. U. S. A. **108:** 20820-20825 10.1073/pnas.1110518109

Ilík P, Pavlovič A, Kouřil R, Alboresi A, Morosinotto T, Allahverdiyeva Y, Aro E-M, Yamamoto H, Shikanai T (2017) Alternative electron transport mediated by flavodiiron proteins is operational in organisms from cyanobacteria up to gymnosperms. New Phytol. **214:** 967-972 10.1111/nph.14536

IPCC (2021) Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA

Jans F, Mignolet E, Houyoux PA, Cardol P, Ghysels B, Cuine S, Cournac L, Peltier G, Remacle C, Franck F (2008) A type II NAD(P) H dehydrogenase mediates light-independent plastoquinone reduction in the chloroplast of *Chlamydomonas*. Proc. Natl. Acad. Sci. U.S.A. **105:** 20546-20551 10.1073/pnas.0806896105

Joët T, Cournac L, Horvath EM, Medgyesy P, Peltier G (2001) Increased sensitivity of photosynthesis to antimycin A induced by inactivation of the chloroplast ndhB gene. evidence for a participation of the NADH-dehydrogenase complex to cyclic electron flow around photosystem I. Plant Physiol. **125:** 1919-1929 10.1104/pp.125.4.1919

Johnson JE, Berry JA (2021) The role of Cytochrome b6f in the control of steady-state photosynthesis: a conceptual and quantitative model. Phot. Res. **148:** 101-136 10.1007/s11120-021-00840-4

Johnson X, Alric J (2013) Central carbon metabolism and electron transport in *Chlamydomonas reinhardtii*: metabolic constraints for carbon partitioning between oil and starch. Eukaryotic Cell **12:** 776-793 10.1128/ec.00318-12

Johnson X, Steinbeck J, Dent RM, Takahashi H, Richaud P, Ozawa SI, Houille-Vernes L, Petroutsos D, Rappaport F, Grossman AR, Niyogi KK, Hippler M, Alric J (2014) Proton Gradient Regulation 5-mediated cyclic electron flow under ATP- or redox-limited conditions: A study of Δ*ATPase pgr5* and Δ*rbcL pgr5* mutants in the green alga *Chlamydomonas reinhardtii*. Plant Physiol. **165:** 438-452 10.1104/pp.113.233593

Kaye Y, Huang W, Clowez S, Saroussi S, Idoine A, Sanz-Luque E, Grossman AR (2019) The mitochondrial alternative oxidase from *Chlamydomonas reinhardtii* enables survival in high light. J. Biol. Chem. **294:** 1380-1395 10.1074/jbc.RA118.004667

Kramer DM, Evans JR (2011) The importance of energy balance in improving photosynthetic productivity. Plant Physiol. **155:** 70-78 10.1104/pp.110.166652

Lemaire C, Wollman FA, Bennoun P (1988) Restoration of phototrophic growth in a mutant of *Chlamydomonas reinhardtii* in which the chloroplast *atpB* gene of the ATP synthase has a deletion: an example of mitochondria-dependent photosynthesis. Proc. Natl. Acad. Sci. USA **85:** 1344-1348

Malone LA, Proctor MS, Hitchcock A, Hunter CN, Johnson MP (2021) Cytochrome b6f – Orchestrator of photosynthetic electron transfer. Biochim. Biophys. Acta **1862:** 148380 https://doi.org/10.1016/j.bbabio.2021.148380

Mellon M, Storti M, Vera-Vives AM, Kramer DM, Alboresi A, Morosinotto T (2021) Inactivation of mitochondrial complex I stimulates chloroplast ATPase in Physcomitrium patens. Plant Physiol. **187:** 931-946 10.1093/plphys/kiab276

Munekage Y, Hashimoto M, Miyake C, Tomizawa K-I, Endo T, Tasaka M, Shikanai T (2004) Cyclic electron flow around photosystem I is essential for photosynthesis. Nature **429:** 579-582 10.1038/nature02598

Munekage Y, Hojo M, Meurer J, Endo T, Tasaka M, Shikanai T (2002) PGR5 is involved in cyclic electron flow around photosystem I and is essential for photoprotection in *Arabidopsis*. Cell **110:** 361-371 https://doi.org/10.1016/S0092-8674(02)00867-X

Nawrocki WJ, Bailleul B, Picot D, Cardol P, Rappaport F, Wollman FA, Joliot P (2019) The mechanism of cyclic electron flow. Biochim. Biophys. Acta Bioenerg. **1860:** 433-438 https://doi.org/10.1016/j.bbabio.2018.12.005

Osmond CB (1981) Photorespiration and photoinhibition: Some implications for the energetics of photosynthesis. Biochim. Biophys. Acta **639:** 77-98 https://doi.org/10.1016/0304-4173(81)90006-9

Peltier G, Aro EM, Shikanai T (2016) NDH-1 and NDH-2 plastoquinone reductases in oxygenic photosynthesis. Ann. Rev. Plant Biol. **67:** 55-80 10.1146/annurev-arplant-043014-114752

Peltier G, Schmidt GW (1991) Chlororespiration - an adaptation to nitrogen deficiency in *Chlamydomonas reinhardtii*. Proc. Natl. Acad. Sci. USA **88:** 4791-4795 10.1073/pnas.88.11.4791

Peltier G, Tolleter D, Billon E, Cournac L (2010) Auxiliary electron transport pathways in chloroplasts of microalgae. Photosynth. Res. **106:** 19-31

Petroutsos D, Terauchi AM, Busch A, Hirschmann I, Merchant SS, Finazzi G, Hippler M (2009) PGRL1 participates in iron-induced remodeling of the photosynthetic apparatus and in energy metabolism in *Chlamydomonas reinhardtii*. J. Biol. Chem. **284:** 32770-32781 10.1074/jbc.M109.050468

Raghavendra AS, Padmasree K (2003) Beneficial interactions of mitochondrial metabolism with photosynthetic carbon assimilation. Trends Plant Sci. **8:** 546-553 https://doi.org/10.1016/j.tplants.2003.09.015

Saroussi SI, Wittkopp TM, Grossman AR (2016) The Type II NADPH dehydrogenase facilitates cyclic electron flow, energy-dependent quenching, and chlororespiratory

metabolism during acclimation of *Chlamydomonas reinhardtii* to nitrogen deprivation Plant Physiol. **170:** 1975-1988 10.1104/pp.15.02014

Shimakawa G, Ishizaki K, Tsukamoto S, Tanaka M, Sejima T, Miyake C (2017) The Liverwort, Marchantia, drives alternative electron flow using a flavodiiron protein to protect PSI. Plant Physiol. **173:** 1636-1647 10.1104/pp.16.01038

Storti M, Alboresi A, Gerotto C, Aro E-M, Finazzi G, Morosinotto T (2019) Role of cyclic and pseudo-cyclic electron transport in response to dynamic light changes in *Physcomitrella patens*. Plant Cell Environ. **42:** 1590-1602 10.1111/pce.13493

Storti M, Segalla A, Mellon M, Alboresi A, Morosinotto T (2020) Regulation of electron transport is essential for photosystem I stability and plant growth. New Phytol. **228:** 1316-1326 https://doi.org/10.1111/nph.16643

Strand DD, Fisher N, Kramer DM (2017) The higher plant plastid NAD(P)H dehydrogenase-like complex (NDH) is a high efficiency proton pump that increases ATP production by cyclic electron flow. J. Biol. Chem. **292:** 11850-11860 10.1074/jbc.M116.770792

Takabayashi A, Kishine M, Asada K, Endo T, Sato F (2005) Differential use of two cyclic electron flows around photosystem I for driving CO_2-concentration mechanism in C4 photosynthesis. Proc. Natl. Acad. Sci. USA **102:** 16898-16903 10.1073/pnas.0507095102

Tolleter D, Ghysels B, Alric J, Petroutsos D, Tolstygina I, Krawietz D, Happe T, Auroy P, Adriano JM, Beyly A, Cuine S, Plet J, Reiter IM, Genty B, Cournac L, Hippler M, Peltier G (2011) Control of hydrogen photoproduction by the proton gradient generated by cyclic electron flow in *Chlamydomonas reinhardtii*. Plant Cell **23:** 2619-2630 10.1105/tpc.111.086876

Tomaz T, Bagard M, Pracharoenwattana I, Lindén P, Lee CP, Carroll AJ, Ströher E, Smith SM, Gardeström P, Millar AH (2010) Mitochondrial malate dehydrogenase lowers leaf respiration and alters photorespiration and plant growth in *Arabidopsis*. Plant Physiol. **154:** 1143-1157 10.1104/pp.110.161612

Walker BJ, Strand DD, Kramer DM, Cousins AB (2014) The response of cyclic electron flow around photosystem I to changes in photorespiration and nitrate assimilation Plant Physiol. **165:** 453-462 10.1104/pp.114.238238

Figure 1. Electron transport pathways operating during oxygenic photosynthesis and their role in ATP generation. Linear electron flow (LEF) operates from PSII to PSI and reduces NADP$^+$ to NADPH using electrons from water. It also generates a *pmf* across the thylakoid membrane composed of a proton gradient (ΔpH) and electrical gradient ($\Delta\Psi$); the *pmf* is used by the ATPase to generate ATP. Cyclic electron flow (CEF) and pseudo cyclic electron flow (PCEF) both contribute to the generation of the *pmf* but do not produce reducing equivalents since the e$^-$ used for PCEF reduce O_2 to water. Chloroplast to mitochondria electron flow (CMEF) encompasses multiple metabolic pathways by which reductants generated in the chloroplast is shuttled to the mitochondria and converted into ATP by the mitochondrial electron transport chain. NADPH and ATP generated by these different pathways are eventually used for transforming CO_2 into biomass, which reflects net photosynthesis. Other abbreviations: AOX, alternative oxidase; ATP, adenosine triphosphate; CBB, Calvin Benson Bassham; FDX, ferredoxin; FLVs, flavodiiron proteins; FNR, ferredoxin-NADP$^+$ reductase; Myxo, myxothiazol; NADPH, reduced form of nicotinamide adenine dinucleotide phosphate; PS, photosystem; PGR5/L1, proton gradient generation 5/like-1; SHAM, salicyl hydroxamic acid.

Figure 2. Dependence of net photosynthesis on CEF, PCEF and CMEF. A, B, O_2 exchange rates upon a dark to light transition in WT2 (**A**) and *pgrl1 flvB-1* (**B**). At t=6min, mitochondrial respiration inhibitors myxothiazol (2.5 μM) and salicylhydroxamic acid (SHAM, 400 μM) were added to the reaction. Shown are representative traces of *n* = 3 biologically independent samples. **C and D**, Net O_2 evolution in *pgrl1 flvB-1* and *pgrl1 flvB*-3 and their control strains (WT2 and WT3) in the absence (control) or presence (Myxo & SHAM) of myxothiazol (2.5 μM) and salicylhydroxamic acid (SHAM, 400 μM) in cells grown in air (**C**) or in 2% CO_2-enriched air (**D**). Values are taken after 9 min of illumination upon a dark to light transitions. Bars show the average and dots show individual replicates (*n* = 3 biologically independent samples). Typical experiments used to extract these data are shown in **Sup. Fig. S2**. **E and F**, Cross section images showing subcellular localization of mitochondria-localized Clover (yellow fluorescence) in *pgrl1 flvB-1* and a control strain (WT2), grown in air (**E**) or 2% CO_2-enriched air (**F**). Images are representative of 2 independent experiments. Scale bar: 10 µm. Brightfield, Clover fluorescence, Chlorophyll autofluorescence and Clover/Chlorophyll merge signals are shown. Images are representative of 2 experiments. Scale bar: 10 µm.

Figure 3. Relative efficiencies of CEF, PCEF and CMEF in energizing net photosynthesis. A-D, Net O_2 evolution measured during a dark to light transition in *flvB* (**A**), *pgrl1* (B), *pgrl1 flvB*-2 (**C, D**), treated (gray dots) or not treated (black dots) with both myxothiazol (Myxo, 2.5 μM) and salicylhydroxamic acid (SHAM, 400 μM) (**A, B**), or with only one inhibitor (**C,D**). Shown are representative traces of 3 biological replicates. **E**, Relative V_{max} measured when a given set of alternative electron pathways are active after treatment with myxothiazol (Myxo, 2.5 μM), salicylhydroxamic acid (SHAM, 400 μM) or both inhibitors; CEF (blue, *flvB* treated with Myxo and SHAM), PCEF (green, *pgrl1* treated with Myxo and SHAM), CEF&PCEF (dark, control strains treated with Myxo and SHAM), CMEF via AOX (purple, *pgrl1 flvB* treated with Myxo), CMEF via III/IV (red, *pgrl1 flvB* treated with SHAM). **F,** Electron quantum yield of PCEF (green), CMEF via AOX (purple), CMEF via III/IV (red), CMEF (pink) and CEF (dotted line). Similar electron quantum yields were also determined for PCEF in 2% CO_2-grown cells and a slightly lower quantum yield for CMEF (**Sup Fig. S9 D**). **G**, Electron flux relative to PCEF required to power similar levels of net photosynthesis when relying on a single pathway. Data shown are individual extrapolations based on measured electron quantum yields. In **E, F** and **G,** bars show the average and dots show individual replicates ($n \geq 3$ biologically independent sample).

Figure 4. Contribution of CEF, PCEF and CMEF to the electron flow, net photosynthesis and growth. A Electron flow through PCEF, CEF and CMEF relative to LEF were determined in the wildtype strain CC-4533 by two different methods, one based on the use of inhibitors (inhib) the other on mutants (mut). Data used for these calculations are the same as for **Fig. 2 C** and **Sup Fig. 2**. **B** Proportion of net photosynthesis powered by each electron flow (PCEF, CEF and CMEF) in the control strain CC-4533 as measured by the inhibition (inhib) or mutant (mut) method. CC-4533 was chosen because it is the control strain for *flvB* and is therefore the only strain where both methods could be used. Data used for these calculations are the same as for **A**. In **A** and **B,** bars show the average and dots show individual replicates (*n*=3 biologically independent sample). **C**, Growth tests for *flvB* and *pgrl1* mutants, their corresponding control strains (CC-4533 and 137AH, respectively) and double mutants (*pgrl1 flvB-1 and pgrl1 flv-3*) and their corresponding control strains (WT2 and 3). Cells were spotted on plates containing minimal medium at pH 7.2 and grown under continuous illumination (60 µmol photon m^{-2} s^{-1}) in the absence (upper panel) or presence (lower panel) of 2.5µM Myxothiazol (Myxo) in air enriched at 2% CO_2. Spots shown are representative of 10 independent experiments.