The landscape of antibiotic resistance genes in the nasopharynx of preterm infants: Prolonged signature of hospitalization and effects by antibiotics

Achal Dhariwal[1]†, Polona Rajar[1,2]†, Gabriela Salvadori[1], Heidi Aarø Åmdal[1], Dag Berild[3,4], Ola Didrik Saugstad[5], Drude Fugelseth[2,4], Gorm Greisen[6], Ulf Dahle[7], Kirsti Haaland[2], Fernanda Cristina Petersen[1]*

[1] Institute of Oral Biology, Faculty of Dentistry, University of Oslo, Oslo, Norway.

[2] Department of Neonatal Intensive Care, Division of Pediatric and Adolescent Medicine, Oslo University Hospital Ullevål, Oslo, Norway.

[3] Department of Infectious Diseases, Oslo University Hospital, Oslo, Norway.

[4] Institute of Clinical Medicine, Faculty of Medicine, Oslo University, Oslo, Norway.

[5] Department of Pediatric Research, Oslo University Hospital, University of Oslo, Oslo, Norway.

[6] Department of Neonatology, Copenhagen University Hospital Rigshospitalet, Copenhagen, Denmark.

[7] Centre for Antimicrobial Resistance, Norwegian Institute of Public Health, Oslo, Norway

†Shared First authorship

*Correspondence: Fernanda Cristina Petersen

Institute of Oral Biology

Faculty of Dentistry,

University of Oslo, Oslo, Norway.

Tel: +47-22840312

E-mail: f.c.petersen@odont.uio.no

ABSTRACT

Respiratory pathogens, commonly colonizing nasopharynx, are among the leading causes of death due to antimicrobial resistance. Yet, antibiotic resistance determinants within nasopharyngeal microbial communities remain poorly understood. Utilizing shotgun metagenomics, we investigated the nasopharynx resistome development in preterm infants, assessed early antibiotic impact on its trajectory, and explored its association with clinical covariates. Our findings revealed widespread nasopharyngeal carriage of antibiotic resistance genes (ARGs) with resistomes undergoing transient changes, including increased ARG diversity, abundance, and composition alterations due to early antibiotic exposure. ARGs associated with the critical nosocomial pathogen *Serratia marcescens* persisted up to 8-10 months of age, representing a long-lasting hospitalization signature. The nasopharyngeal resistome strongly correlated with microbiome composition, with inter-individual differences and postnatal age explaining most of the variation. Our report on the collateral effects of antibiotics and prolonged hospitalization underscores the urgency of further studies focused on this relatively unexplored reservoir of pathogens and ARGs.

Keywords: respiratory resistome, metagenomics, nasopharynx, microbiome, preterm infants, antibiotics, antimicrobial resistance

INTRODUCTION

Antimicrobial resistant infections are a major global health threat. Among these, respiratory infections account for more than 1.5 million deaths a year, surpassing the number of deaths caused by resistant infections in any other part of the human body niche [1]. Main pathogens include *Streptococcus pneumoniae, Staphylococcus aureus, Klebsiella pneumoniae,* and *Haemophilus influenzae*, all generally present in the nasopharynx of healthy populations. These pathogens share common niches with commensals, which in turn provide colonization resistance against pathogens. However, commensals can also serve as reservoirs of antibiotic resistance genes (ARGs) that can readily be transferred to pathogens [2-4].

Recent studies have highlighted the potential negative impact of antibiotic treatment on the nasopharynx microbiome and its association with an increased risk of developing asthma and respiratory infections in infants [5-7]. These studies have significantly advanced our understanding of the ecological impact of antibiotics on respiratory health and disease. However, as these studies rely on 16S rRNA sequencing technologies, they do not offer a high level of microbial taxonomic resolution nor provide information on the overall gene composition, including the presence of ARGs within microbial communities, referred to as the resistome. In our previous study, we developed an optimization strategy for DNA extraction and library preparation that helped to circumvent the challenges posed by low bacterial biomass and high host DNA content in nasopharynx samples from preterm infants [8]. In the present study, we applied this optimized protocol to nasopharyngeal aspirate samples to characterize the resistome development in 36 preterm infants from birth to 6 months corrected age (8-10 months chronological age). By analyzing ~1.85 terabytes (Tb) of metagenomic DNA data generated from deep shotgun metagenomic sequencing, we obtained the most comprehensive view to date of the characteristics and dynamics of resistome development in the respiratory tract of preterm infants. Our hypothesis was that early exposure to broad-spectrum antibiotics for suspected early onset sepsis (sEONS) treatment in preterm infants may lead to prolonged compositional alterations in the respiratory resistome. We also explored the extensive metadata collected to gain insights into additional clinical covariates that may influence resistome development.

RESULTS

Distribution and characterization of ARGs in the preterm infant nasopharyngeal microbiome

In total, we found 369 ARGs belonging to 15 ARG classes conferring resistance through 5

distinct mechanisms. ARGs associated with multidrug efflux pumps comprised the majority

of the resistome with an average of 27% from T1 to T5 and 17.8% at T6 (six months

corrected age) of the total relative abundance. There was a median of 18 (mean: 24) unique

ARGs ranging from 1 to 155 per analyzed sample. On average, the maximum number of

ARGs were detected after the discontinuation of treatment (T2) (mean: 40) in most infants

receiving early antibiotic treatment. In the naive group, the corresponding mean value at T2

was 19. Only 13 ARGs were found to be both highly prevalent (detected in >40% of all

samples) and highly abundant, with a mean relative abundance of 65.2% (SD: 32%) across all

preterm infant nasopharyngeal resistome. These core set of resistance genes included

multidrug resistance genes (*acrB, oqxB, mexI),* macrolide-lincosamide-streptogramin (MLS)

resistance genes (*mel, pmrA, rlmA(II)*), two beta-lactam resistance genes (*blaZ, SST-1),*

fluoroquinolone resistance genes (*patA, patB),* tetracycline resistance genes (*tet(41), tetM*),

and the aminoglycoside resistance gene *AAC(6')-Ic*.

In both antibiotic-treated and naive groups, the most abundant class of ARG was multidrug

(mean: 27.95%; SD: 21.96%), followed by fluoroquinolone (mean: 17.45%; SD: 20.28%),

beta-lactam (mean: 15.35%; SD: 17.05%), tetracycline (mean: 14.33%; SD: 10.93%), MLS

(mean: 13.18%; SD: 15.43%) and aminoglycoside (mean: 8.95%; SD: 12.94%) (Fig. 1A). In

terms of prevalence, aminoglycoside (76/82) and beta-lactam (75/82) were the most common

classes present in the samples from antibiotic-treated group, while multidrug (92/99) and

beta-lactam (91/99) were the most common classes in the naive infants. When classified

based on resistance mechanism, antibiotic efflux (mean: 59.17%; SD: 21.54%) was the most

abundant category in both the groups across all time points (Supplementary Fig. 4). At

baseline (T1), the ARG composition of the antibiotic-treated group was found to be more

heterogeneous (higher inter-individual variability) and diverse compared to that of naive

infants (Fig. 1B).

The health risks of ARGs were assessed using the recently proposed ARG-ranker list, in

which Rank IV comprises ARGs not associated with humans, Rank III includes human-

associated ARGs found in non-mobile elements, Rank II encompasses mobile ARGs capable

of transferring existing resistance to pathogens, and Rank I comprises ARGs already present

in pathogens, thus posing the highest risk to human health [9]. We detected homologs to

potential high-risk ARGs (rank I & II) in the nasopharyngeal samples at multiple time points

across infants in both antibiotic-treated and naive groups (Supplementary Fig. 5). We also

observed a few high-risk and clinically relevant ARGs that appeared directly after the early

antibiotic treatment at T2. For instance, high-risk ARGs specific to the antibiotics encoding

for aminoglycoside-modifying enzymes (*AAC(3)-II*) along with extended-spectrum TEM

beta-lactamase (*bla*$_{TEM-1}$) were detected in two infants. These ARGs co-occurred with other

non-targeted high-risk (rank II) ARGs such as *qnrS*, *floR* and *aadA*. Previous studies have

shown the co-carriage of these ARGs on multidrug resistance plasmids in Enterobacter

species [10]. Also, other clinically relevant Extended Spectrum Beta-Lactamase (ESBL)

genes belonging to the SHV beta-lactamase (rank II) family were detected in two other

infants following antibiotic exposure. In one patient, the plasmid-mediated AmpC-type β-

lactamase (ACT-beta lactamase), known to confer resistance against all classes of beta-

lactam antibiotics such as penicillins, cephalosporins and carbapenems was detected only

after the treatment cessation (T2). These ARGs were mainly observed in early antibiotics

exposed infants whose mothers had also received antibiotics during pregnancy (prenatal).

Nonetheless, they did not persist in nasopharyngeal samples collected after discharge at 6

months corrected age (T6). Amongst ARGs found in Gram-positive pathogens, the

Staphylococcal methicillin-resistance gene i.e., *mecA* was detected in 44 samples from almost

all (14 of 15) antibiotic-treated infants and 10 of the 21 naive infants.

Impact of early antibiotics on preterm infant nasopharyngeal resistome

Firstly, we investigated the total ARG abundance (quantified as the sum of RPKM) and ARG

α-diversity (quantified by the Shannon index) across sampling time points. In both the

antibiotic-naive and treated groups, we found that total ARG abundance and diversity

increased from T1 to T3-T4, before descending at T5 to the 6 months corrected age (T6). In

treated infants, total ARG abundance and diversity increased more rapidly upon antibiotic

treatment, whereas the changes were generally more gradual over time points in naive infants

(Fig. 2A, B). To identify the direct effects of early antibiotic treatment on the total ARG

abundance and diversity, we used a generalized linear mixed model with individual set as

random effect while correcting for age (DOL). Early antibiotics were significantly associated

with increase in both total ARG abundance (LME model: *p.adj* = 0.01) and Shannon

diversity (*p.adj* = 0.01) after treatment (T2) as compared to the baseline (T1) samples in

preterm infant nasopharyngeal resistome. Moreover, the significant effect of antibiotics on

the increase in total ARG abundance was observed until T3 (*p.adj* = 0.001). Nonetheless,

such effects were short-lived, as we did not identify significant differences at any of the

following time points compared to the baseline (T1) in antibiotic-treated group. In the naive

group of infants, no significant differences appeared in total ARG abundance and α-diversity

between any timepoints during the first 6 months (corrected) of life while correcting for age.

β-diversity analysis revealed that the overall resistome composition did significantly differ

between the two groups (PERMANOVA: R^2 = 8.4%, *p.adj* = 0.02) at the baseline (T1).

Infants that received early antibiotic treatment had more dissimilar (dispersed) resistome

composition than the naive infants (Fig. 3A). Further, we observed that nasopharyngeal

samples were clustering more effectively according to birth weight group (R^2 = 16.5%, *p.adj*

= 0.01; Supplementary Fig. 6A). No significant difference in the baseline resistome

composition between antibiotic-treated and naive infants (R^2 = 3.9%, *p.adj* = 0.26) was found

while correcting for birth weight group. However, a minor yet significant compositional

difference in the resistome was still observed between the two groups (R^2 = 6.4%, *p.adj* =

0.02; adjusted for BW group: R^2 = 5.9%, *p.adj* = 0.03) directly after the cessation of early

antibiotic treatment (T2) (Fig. 3B). Moreover, the compositional changes due to early

antibiotics at T2 were more discernible in infants whose mothers had received prenatal

antibiotics during pregnancy (prenatal + early (*n = 8*) vs naive (*n = 20*): R^2 = 10.1%, *p.adj* =

0.009; Supplementary Fig. 6B). Following treatment, the resistome normalized rapidly as no

significant differences in composition between the naive and treated groups were detected

from T3 until 6 months of corrected age (T6) (Fig. 3C).

Clinical covariates associated with the development of preterm infant nasopharyngeal

resistome composition

The associations between resistome composition and clinical factors were analyzed using

PERMANOVA-tests (all *p.adj* ≤ 0.001) and principal component analysis (PCA) across all

nasopharyngeal samples from preterm birth until 6 months corrected age, using centered log-

ratio (CLR) transformed data (Supplementary Fig. 2). The PCA was explained by two main

principal components, describing 18.7% and 17.6% of the variation, respectively. Besides

inter-individual variation (PERMANOVA (univariate): R^2 = 37.8%, *p.adj* = 0.001), age

was the other significant covariate found to be associated with overall resistome composition

in the overarching cohort. A moderate yet statistically significant effect was observed with

GA (R^2 = 4%, *p.adj* = 0.001) and BW group (R^2 = 4.1%, *p.adj* = 0.001), which are highly

correlated variables (as indicated by the chi-square test: $p < 2.2e\text{-}16$). Mode of delivery, sex,

and intrapartum antibiotic treatment did not appear to shape the overall nasopharyngeal

resistome composition (Supplementary Table 3). Covariates that differed between the
antibiotic-treated and naive infants at baseline and that also had a significant association with
the overall resistome composition (GA and prenatal antibiotics ($R^2 = 4.5\%$, $p.adj = 0.001$))
were used for correction in the subsequent multivariate analysis.

Multivariate PERMANOVA analysis showed that the largest effect on overall resistome
composition remained attributed to inter-individual variation ($R^2 = 30.13\%$, $p.adj = 0.001$),
followed by PMA ($R^2 = 9.8\%$, $p.adj = 0.001$) and the chronological age group (sampling
time points) ($R^2 = 8.4\%$, $p.adj = 0.001$). Although we observed a high degree of variation
between samples with no obvious discrete spatial clustering in relation to these variables
upon visual inspection (Fig. 4A-C). We found no significant influence of early antibiotics (R^2
$= 0.6\%$, $p.adj = 0.339$) and its duration ($R^2 = 1.3\%$, $p.adj = 0.007$) after adjusting for
confounding variables. However, we observed a significant effect of prenatal antibiotics (R^2
$= 3.6\%$, $p.adj = 0.001$) and its duration (categorized) ($R^2 = 4.4\%$, $p.adj = 0.001$) in shaping
the overall nasopharyngeal resistome composition in preterm birth. To account for the
likelihood of prenatal antibiotics with early antibiotics exposure in our analyses (Chi-squared
test, $p < 0.0003$), we further categorized our early antibiotic-treated group based on whether
the mothers had also received antibiotics during pregnancy (prenatal + early, *n = 9*) or not
(only early, *n = 6*). Post-hoc analysis revealed significant differences in the overall resistome
composition of infants having exposure to both prenatal and early life antibiotics (prenatal +
early) compared to infants receiving either only early or naive (no prenatal + no early) (*p.adj*
$= 0.001$; Fig. 4D). No difference in overall resistome composition was observed between the
only early antibiotics and naive groups (*p.adj* = 0.06), underlining the relevance of prenatal
antibiotic exposure in shaping the overall resistome composition of preterm infants during the
first 6 months (corrected) of life.

Unsupervised clustering classifies nasopharyngeal samples into distinct clusters based
on composition differences
Next, we determined whether the nasopharyngeal samples were organized into clusters (i.e.,
resistotypes) according to their ARG abundance profiles using an unsupervised method, i.e.,
DMM models. This way, we identified three distinct clusters (or resistotypes) that were
driven by specific ARGs. The majority of our samples (~87%) were classified into two main
resistotypes, R1 and R2. The first resistotype (R1) was characterized by a predominance of
fluoroquinolone resistance genes (*patA, patB*) and *rlmA(II)* (Supplementary Fig. 7). On the

other hand, resistotype R2 was driven mainly by relatively higher abundance of the beta-
lactam resistance gene *SST-1,* aminoglycoside resistance gene *AAC(6')-Ic*, tetracycline
resistance gene *tet(41)* and the multidrug resistance gene *mexl*. Resistotype R3 was enriched
with the blaZ β-lactamase resistance gene. The identified resistotypes varied in their overall
ARG composition (β-diversity), as shown in a PCA plot (Fig. 5A). Furthermore,
PERMANOVA analysis showed that resistotypes were more strongly (larger effect size) and
significantly associated with the resistome than other clinical covariates or early-life factors
in preterm infants during the first 6 months (corrected) of life ($R^2 = 20.4\%$, *p.adj* = 0.001).

Nasopharyngeal resistome composition of preterm infants is linked to microbial
composition and carries hospitalization signatures
To investigate the impact of microbial composition on the resistome composition, we
taxonomically profiled the infant nasopharyngeal samples using MetaPhlAn3 (manuscript
under review, Rajar et al., 2024). We noticed that microbial community types, as determined
by DMM models, were highly linked to the resistotypes (chi-square test: *p* < 0.0001).
Resistotype R2 was very specific to the *Serratia* community type. Resistotype R1 was
associated with community types mainly driven by *Streptococcus* and *Gemella*, while
Resistotype R3 was enriched in community types predominated by *Staphylococcus* and
Streptococcus (Fig. 5B). These community types were also significantly associated with the
overall composition of the resistome (PERMANOVA: $R^2 = 13.5\%$, *p.adj* = 0.001; Fig. 5C).
Additionally, results from Procrustes analysis showed a strong and significant association
between the microbial taxa and ARG abundance profiles (PROTEST: correlation coefficient
(r) = 0.87, *p* = 0.001, m^2 = 0.24), underlining that bacterial community composition
structured the resistome composition in the nasopharyngeal microbiome of preterm infants.
However, the correlation was more robust in the antibiotic-naive infants (PROTEST: r =
0.94, *p* = 0.001, m^2 = 0.12) as compared to antibiotic-treated infants (PROTEST: r = 0.85, *p* =
0.001, m^2 = 0.27; Fig. 6A, B).

Next, we performed pairwise Spearman's correlation analysis between ARG and microbial
species abundances to predict the origin or potential microbial host of ARGs in
nasopharyngeal samples. The strongest positive correlation (r ≥ 0.8, *p.adj* < 0.05) was found
between the *SST-1, AAC(6')-Ic, tet(41)* ARGs and *Serratia marcescens/nematodiphila* (Fig.
6C). It is worth noting that during our study, a clinically detected outbreak of *S. marcescens*
occurred in the NICU a few months after infant enrollment had commenced. When present at

T1, the carriage of these ARGs persisted in 24 out of 26 preterm infants across both naïve and

treated groups, extending long after hospital discharge as observed at ~8 to 10 months

chronological age (i.e., 6 months corrected age (T6)) (Fig. 6D). Occasional time points in

which ARGs associated with *S. marcescens* showed variations from T1 were observed in few

of the samples. Whether these are true variations is difficult to ascertain as metagenomic

studies can only detect DNA sequences that are present in specific samples and to a depth

that is possible to be captured by present analytical approaches.

The highly prevalent ARGs like *patA, patB and rlmA(II)* within Resistotype 1 were observed

to exhibit a strong co-occurrence with *Streptococcus mitis/oralis* and *Gemella*

haemolysans/sanguinis (Fig. 6C)*. Staphylococcus aureus* and *Staphylococcus epidermidis*

correlated only with *blaZ* beta-lactam ARG. These associations are in accordance with

information in the literature and AMR gene databases on the known microbial hosts of these

ARGs, highlighting the potential of such an analytical approach to predict the origin of ARG

in metagenomes. However, to reduce the bias due to false ranking of features with many zero

values, this analysis is only applicable on the features (ARGs or taxa) present in high

prevalence across samples.

Lastly, we also examined whether some of these strongly associated bacterial species with

ARGs would also directly correlate with our overall resistome outcomes. We plotted the

relative abundance of each species versus the total ARG abundance (sum of RPKM) and the

number of unique ARGs from all samples. Our analysis revealed that only the relative

abundance of *Streptococcus mitis* (rmcorr correlation coefficient (r^{rm} = 0.43, *p.adj* = 5.82e-

08) showed a moderately linear association with the total ARG abundance while adjusting for

inter-individual variability (Supplementary Fig. 8). Additionally, we observed a small but

significant association between the relative abundance of *S. mitis* and the number of unique

ARGs (r^{rm} = 0.19, *p.adj* = 0.02). To statistically assess whether *S. mitis* relative abundance

explains the dispersion of resistomes across samples, we conducted the PERMANOVA test

and found that *S. mitis* significantly explains 8.5% (*p.adj* = 0.001) of the variation in the

overall resistome composition. Only *S. marcescens* (R^2 = 13.9%, *p.adj* = 0.001) accounts for

a higher proportion of the total variance in the resistome composition.

DISCUSSION

The nasopharynx represents a highly accessible microbial community and also serves as a
crucial diagnostic window in combating respiratory infections and AMR [11]. It is also the
primary ecological niche for common respiratory pathogens [12]. To date, one of the closest
attempts at an in-depth investigation of ARGs in the respiratory tract using a culture-free
approach identified ARGs in 86% of the samples. However, this was based on the
microbiomes of the anterior nares and from an adult population (Human Microbiome Project)
[13], which are different from the groups and collection sample sites used in our study. In a
recent study focusing on infants, ARGs were detected in 64% of nasopharynx samples [14].
Samples in this study were enriched for streptococci, thus precluding general conclusions on
the extent of ARG presence in the natural microbiome. In our study, ARGs were present in
95% of the preterm infant samples from the nasopharynx, regardless of whether they were
exposed to antibiotics or not. Although our study differs in several aspects from the ones
above, the combined findings indicate that the respiratory tract is likely an important
reservoir of ARGs and an integral component of human respiratory microbiomes. For
preterm infants, the detection of ARGs in the nasopharynx during the first few hours of life
might occur due to microbiome colonization from multiple sources such as transgenerational
transfer from mothers and the environment [15, 16]. Procedures common in neonatal
intensive care, such as intubation, feeding tube insertion and suction catheters, may also
disrupt anatomical barriers in the respiratory system, potentially altering the patterns of
microbial colonization [17]. In addition, preterm infants typically necessitate extended
hospital stays after birth compared to term infants, with a greater likelihood of requiring care
in NICUs. This most likely contributes to an increased risk for respiratory tract colonization
with drug-resistant microorganisms, as recently reported in relation to the gut resistome [18].

We found that ARGs conferring multidrug resistance via efflux pumps encompassed a large
proportion of the nasopharyngeal resistome in the preterm infants. Similar findings have been
previously reported in the gut and oral resistome of preterm and term infants by other
metagenomic studies [7, 15, 19, 20]. In addition, we detected many of the same
aminoglycoside-modifying enzymes, tetracycline protection proteins, and beta-lactamase
ARGs identified previously in the nasopharynx of South African infants, even though the
samples in the South African study were culture-enriched for streptococci and were from a
country with higher AMR challenges than the origin of samples used in the present study.
Such similarities in resistome composition may, at least in part, be explained by the large
proportion of streptococci found in the natural microbial communities of the infant

nasopharynx. Together, these findings suggest the existence of a core infant nasopharyngeal
resistome.

Staphylococcus aureus is one of the leading pathogens causing neonatal sepsis and one of the
priority drug-resistant pathogens by the WHO [21]. We found a strong correlation between
the abundance of *S. aureus/epidermidis* and the PC1-beta-lactamase (*blaZ*) gene. This gene is
prevalent in the chromosomes and mobile plasmids of Staphylococci [22], thus representing a
potential high-risk ARG. Another strong correlation was found between *Serratia* species and
specific resistance genes such as *SST-1*, which confers resistance against cephalosporins.
Other resistance genes correlated with this group included aminoglycoside (*AAC(6')-Ic)* and
tetracycline (*tet(41)*) resistance genes. Many clinical and environmental isolates of *Serratia*
spp., especially *S. marcescens strains*, have been reported to harbor these multiple ARGs on
their chromosomes or plasmids [23-26]. This holds particular relevance, given that *S.*
marcescens is linked to nosocomial infections and hospital outbreaks, notably in NICUs [26,
27]. Multidrug resistant *Serratia spp.* are listed among the most critical bacteria by the WHO
for which innovative new treatments are urgently needed [28].

Our results showed a significant effect of early broad-spectrum antibiotic regimen
(ampicillin□+□gentamicin) on the total abundance, diversity, and composition of ARGs in
the nasopharynx of preterm infants. These effects on the resistome were short-lived and only
observed directly after the cessation of antibiotic treatment (T2). In a recent study
investigating the impact of antibiotics used in the first week of life on gut resistome, Reyman
et al. reported that three other antibiotic combinations (penicillin + gentamicin, co-amoxiclav
+ gentamicin, and amoxicillin + cefotaxime) also had a transient impact on the infant
resistome [29]. As the effects of antibiotics on microbiome and resistome composition are
unique to each individual and dependent on the antibiotics used and the body site, caution
should be taken when extending the interpretation of such general findings. Interestingly, the
transient effects of ampicillin□+□gentamicin on the nasopharyngeal resistome were more
discernible in preterm infants whose mothers were exposed to antibiotics during pregnancy
before delivery (prenatal antibiotics). Moreover, we also observed a more profound effect of
prenatal antibiotic exposure on the nasopharyngeal resistome of preterm infants at T2 (after
cessation of early antibiotics), which persisted until the last sampling time point before
discharge from the NICU (T4-T5), but not at 6 months corrected age. Although very few
studies have focused on the association between prenatal antibiotic exposure and the
resistome, a recent study based on the gut indicates a possible correlation of prenatal

antibiotics with increased abundance of ARGs, at least when used 5 to 6 weeks before birth [7]. Additionally, in concordance with other studies on the infant gut [15, 30], our data showed no significant impact of intrapartum antibiotic therapy on the overall nasopharyngeal resistome composition. The nasopharyngeal samples of preterm infants at 6 months corrected age were still clustered according to the birth weight group (similar to T1, Supplementary Fig. 6C), suggesting a strong but transient influence of other environmental factors, such as antibiotic exposures. Nonetheless, there is a paucity of relevant studies on nasopharynx that can corroborate our findings, as they are mainly restricted to the infant gut resistome [15, 16, 18, 29-31].

One of the strengths of this study is that it presents the most comprehensive information to date on the characteristics and dynamics of resistome development in the nasopharynx, including results for the first 8-10 months of life (6 months corrected age) – a critical period of airway development in infants. In addition, the naive group was enrolled at the same time as our treated group from the same hospital, thus removing potential confounding differences in enrollment years or sampling duration. However, the small number of samples at each time point and the inter-individual response to antibiotic exposure may be insufficient to unravel significant associations between antibiotic regimen and resistome development. Moreover, the baseline samples for the antibiotic-treated group ideally would represent samples taken before treatment. However, the initiation of antibiotic treatment for sEONS and the clinical stabilization of the preterm infants were a high priority that needed to precede nasopharynx sample collection. To partially address this issue, we included only preterm infants with samples collected no later than 24 hours after antibiotic initiation. Even so, this choice may have resulted in the selection of more clinically stable infants, which could potentially limit the generalizability of our results. Despite this approach, the differences in resistome changes between baseline and after treatment found in our study may underestimate the true impact of antibiotics on the resistome. We rigorously adjusted for such confounding factors and inter-individual variation using multivariable methods or matched analysis to ensure the reliability of our observations as a consequence of early antibiotics. Nevertheless, we cannot isolate early antibiotic effects from other adverse pre-life events that coincide with premature birth, such as prenatal and antepartum maternally administered antibiotics. Furthermore, for low biomass samples such as those used in our study, sensitive approaches such as deep sequencing are necessary to obtain meaningful microbiome information. This increases the risk of detecting potential DNA contaminants. We mitigated these challenges by

implementing rigorous measures spanning from sample collection to data analysis. This
included the concurrent collection of samples from treated and naive groups, stringent
protection measures during sample collection, incorporation of negative and positive controls,
and meticulous *in-silico* identification and removal of possible contaminants using DNA
sequencing data.

In conclusion, our findings suggest that early-life treatment with broad-spectrum antibiotics
for sEONS can cause perturbations in the nasopharyngeal resistome of preterm infants.
However, the overall perturbations were transient, as we did not observe a significant
alteration in the diversity, abundance, composition, or carriage persistence of potentially
high-risk ARGs in treated preterm infants compared to controls at 6 months of corrected age.
Despite this, such findings hold significance, considering the vulnerability of preterm infants
in their first days of life. The individual variability in the distribution of high-risk ARGs and
the changes in the resistome following antibiotic treatment highlight the potential benefits of
developing resistome profiling tools for point-of-care use, which could aid in informing and
tailoring antibiotic treatment decisions [32]. In addition, prolonged carriage of ARGs
associated with nosocomial *S. marcescens* was noted. Future multicenter studies with long-
term follow-up are needed to assess the ecological side-effects of hospitalization and different
antibiotic regimens, including alternatives to broad-spectrum antimicrobial therapy on infant
respiratory resistome development. Understanding the development of the respiratory
resistome and the impact of antibiotic therapies is relevant for striking a balance between the
conflicting goals of minimizing the risk of serious infectious complications to preterm birth
on one hand, and minimizing the selection of organisms with ARGs on the other. In general,
the disproportionate contribution of respiratory pathogens to drug-resistant deaths at all ages
highlights the importance of future studies focusing on this yet relatively unexplored
reservoir of pathogens and ARGs.

METHODS

Study design and cohort characteristics
The study is a prospective observational cohort study. Between July 2019 and January 2021,
we included preterm infants born (or transferred within 48 hours) at the NICU (Oslo
University Hospital, Ullevål) who were between 28+0 and 31+6 weeks of gestational age
(GA). In total, 66 preterm infants were enrolled in the study, and followed from birth until six

months corrected age (i.e., adjusted for prematurity) during their stay in the NICU and after

discharge. Five of the infants were lost to follow-up. Extensive metadata and covariates from

infants and mothers, including mode of delivery, gestational age (GA) at birth, postmenstrual

age (PMA), sex, birth weight (BW), postnatal infant antibiotics usage (duration, type, days),

maternal antibiotic usage, feeding regimen during hospitalization and after discharge, and

other demographic characteristics, were collected through medical records. This data was

stored in a secured server at the Services for Sensitive Data (TSD), University of Oslo.

In our cohort of 66 preterm infants, 33 infants with sEONS received broad-spectrum

antibiotics, consisting of a combination of ampicillin (a beta-lactam antibiotic) and

gentamicin (an aminoglycoside antibiotic), within 24 hours after birth. Due to the

circumstances of premature birth and the instability of the infants, baseline samples could not

be collected before antibiotic therapy initiation. Instead, the first nasopharyngeal aspirate

samples (baseline) were obtained within 48 hours after the initiation of early antibiotic

therapy. To investigate temporal changes in the resistome in response to antibiotics and

control for potential baseline differences as confounding factors, this paper included only

infants who had nasopharyngeal samples collected on the same day as the initiation of early

antibiotic treatment ($n = 15/33$) and antibiotic-naive preterm infants as controls ($n = 21$). The

average early antibiotic treatment duration was 5 days (SD: 2 days). The sampling time point

T2 corresponds to the first available sample collected (except in one infant) after the

cessation of early antibiotic treatment. In addition to early antibiotics, one of the included

infants received subsequent antibiotics initiated > 72 hours after birth and before discharge

from NICU and two other infants received antibiotic therapy between discharge from NICU

and 6 months corrected age. A timeline of sample collection and antibiotic administration is

represented in Supplementary Fig. 1.

Infants exposed to early antibiotics had significantly lower GA and BW, and their mothers

had a more prolonged period of ruptured membranes before delivery compared to antibiotic-

naive infants. None of the mothers of naive infants received antibiotics during pregnancy

(prenatal antibiotics) before onset of delivery. The baseline characteristics of the infants

stratified by antibiotic exposure are presented in Table 1.

Sample collection

Nasopharyngeal aspirate samples were collected from infants at six distinct time points

spanning from birth to 6 months corrected age (T1-T6). The days of life (DOL) at which

samples were collected for each time point varied as follows: T1 (DOL: 0-3), T2 (DOL: 4-9),

T3 (DOL: 12-19), T4 (DOL: 20-31), T5 (DOL: 32-76) and T6 at 6 months corrected age

(DOL: 229-288). Additional samples were taken if the antibiotic therapy was initiated or

terminated more than 48 hours before/after a predefined sampling time. As a result, in two

infants, there was more than one sample available for the defined time points. To facilitate

time point-wise analysis, only one sample was included for each of the time points

(Supplementary Table 2 & Supplementary Fig. 1). In cases where the sample could not be

collected at the scheduled time points (such as with an unstable infant), a new sample was

obtained as soon as possible. Infants who were transferred to another hospital before

discharged to home adhered to the study protocol. When the infants reached six months of

corrected age, they were either invited to an outpatient visit at Ullevål hospital or their local

hospital, or they were visited at home for sample collection. Nasopharynx aspirates were

collected as previously described [8], then stored and cryopreserved in 20% glycerol (2ml) at

-80°C until further processing. In total, we collected and extracted metagenomic DNA from

198 nasopharyngeal samples obtained from 36 preterm infants, from the day of birth until 6

months corrected age. Detailed information on sample exclusion and inclusion statistics at

different steps from sampling to whole metagenomic sequencing (WMS) is described in

Supplementary Table 1.

Metagenomic DNA extraction, quantification, and sequencing

Metagenomic DNA was extracted by first thawing the aspirate samples (2 ml) on ice,

pelleting them by centrifugation at 10,000 g for 10 minutes at 4°C, and then using MolYsis™

for host DNA depletion (Molzym, Bremen, Germany), following the manufacturer's protocol

with some modifications, as previously described [8]. After depletion of human DNA, fresh

pellets were spiked with 20 µl of ZymoBIOMICS Spike-in Control II (catalog number:

D6321 & D6321-10), and bacterial DNA was extracted using the MasterPure™ Gram

Positive DNA Purification Kit (Epicentre, Madison, WI, USA), according to the

manufacturer's recommendations. This kit employs a chemical lysis method for hard-to-

lyse bacteria that has been shown to outperform mechanical lysis methods for reliable

recovery of DNA from low biomass preterm nasopharyngeal samples [8]. One glycerol

sample from each batch used for the cryopreservation of nasopharyngeal samples (n = 5;

sampling blanks) and one reagent blank per extraction batch (n = 5; DNA extraction blanks)

were extracted alongside the cohort samples using the same protocol, serving as negative

controls. As positive controls, six aliquots (20 µL each) of Spike-in Control II Low Microbial

Load (ZymoBIOMICS™, Catalog Nos. D6321 & D6321-10), extracted in the same manner,

were included. The ZymoBIOMICS Spike-in Control II comprises of three bacterial species

not found in the human microbiome (Truepera radiovictrix, Imtechella halotolerans, and

Allobacillus halotolerans). Finally, the DNA pellet was eluted in 35 µl of TE buffer and

stored at -80°C until further use.

The total DNA concentration was quantified using Qubit dsDNA HS assay kits in a Qubit 4.0

Fluorometer (Invitrogen, Thermo Fisher Scientific, USA). Real-time quantitative polymerase

chain reaction (RT-qPCR) assays were used to quantify human (Zymo E2005) and microbial

DNA (Zymo E2006) in all extracted samples using Zymo Femto^{TM} Quantification kits

(ZymoBIOMICS™). The RT-qPCR results for bacterial DNA (16S rRNA) were also used to

exclude samples with low or negligible bacterial load before sequencing. This was

accomplished by subtracting the average yield of bacterial DNA from the positive spike-in

controls from the yields of patient samples. Samples that exhibited a negative yield in RT-

qPCR ($n = 4$) were excluded from further processing (Supplementary Table 4). The DNA

yield from negative controls was extremely low or undetected ($n = 10$; mean = 0.00043

ng/µL), as measured by RT-qPCR. These samples were included in sequencing runs for in-

silico identification and removal of contaminant signals, alongside the spike-in positive

controls.

Metagenomic sequencing libraries were constructed using the Nextera DNA Flex library

preparation kit (Illumina Inc., CA, USA). All samples were subjected to paired-end

sequencing at 2×150 bp on an Illumina NovaSeq S4 high-output platform at the Norwegian

Sequencing Centre (Oslo, Norway). In total, 6.14 billion raw sequencing reads were obtained

from the nasopharyngeal samples of preterm infants, with a mean sample read counts of

33.97 million (M) ranging from 4.19 to 69.25 M reads. Conversely, negative control samples

exhibited an average read count of 0.15 million (M), with a range of 24,113 to 355,861 reads

(Supplementary Table 2).

Controlling for contaminant in low microbial biomass samples

To mitigate biases due to contamination—a critical consideration in low microbial biomass

microbiome research [33]—we implemented several recommendations as follows: (1)

samples from infants in both groups were collected concurrently to ensure temporal

consistency, (2) standard protection measures were strictly adhered to during sample

collection, including the use of clean suits, disposable gloves, and face masks, (3) both

sampling and DNA extraction blanks were included as negative controls, (4) the DNA

extraction and library preparation processes were conducted in a meticulously cleaned

environment, with personnel wearing laboratory coats and disposable gloves, (5) longitudinal

samples from each preterm infant were processed in a single extraction run, (6) spike-in using

non-human bacterial species were used as positive controls. The *in-silico* identification and

removal of possible contaminants using DNA sequencing data is described below.

540   Bioinformatics processing

The raw metagenomic sequencing data was pre-processed using our in-house bioinformatics

pipelines using a high-performance computing cluster inside a secured environment, i.e.,

TSD at the University of Oslo. In brief, the Nextera adaptor sequences and low-quality reads

were filtered using Trim Galore (v.0.6.1) with default parameters [34]. Next, the human DNA

contaminant sequences were identified and removed by mapping the quality-trimmed reads

against the human reference genome (GRCh38) using Bowtie2 (v.2.3.4.2) [35] (non-default

parameters: q -N 1 -k 1 --fr --end-to-end --phred33 --very-sensitive –no-discordant) along

with SAMtools (v.1.9) [36] and BEDTools (v.2.27.1) [37]. On average, 46.71% (range: 1.09 -

97.69%) of total quality-filtered reads were identified as belonging to the human genome.

After removal of the human-associated metagenomic reads, a total of 2.98 billion remaining

high-quality clean reads across all nasopharyngeal samples with number of reads ranging

from 0.76 to 59.9 M per sample, and with median of 16.4 M reads were subjected to

resistome profiling. More detailed results regarding the overall sequence preprocessing are

presented in Supplementary Table 2. The sequence quality reports on raw and processed

reads were generated using FastQC (v.0.11.9) [38] and summarized using MultiQC (v.1.7)

[39].

558   Resistome and microbiome profiling

To characterize the presence of ARGs, cleaned high-quality reads for each sample were

mapped against the nucleotide_fasta_protein_homolog_model from the Comprehensive

Antibiotic Resistance Database (CARD) (v.3.2.2) [22] using Bowtie2 under parameter –very-

sensitive-local. The mapped reads from each sample were then filtered, sorted, and indexed

using SAMtools. The number of reads mapped to each ARG was calculated using SAMtools

idxstats and BEDTools. We considered ARGs with a gene fraction of at least 80% (meaning,

80% or more of the nucleotides in the ARG sequence were covered by at least one read) as

positively detected in a sample. Additionally, we excluded ARGs from subsequent
downstream analysis if they could not be confidently attributed to resistance based solely on a
short-read marker, such as those linked to resistance through mutations, as detailed by
D'Souza et al. [40]. The mapped read counts were normalized for bacterial sequence
abundances and gene lengths by calculating reads per kilobase of reference gene per million
bacterial reads (RPKM) for each CARD reference sequence. The relative abundance of
ARGs for each sample was computed by dividing the RPKM by the total sum of the RPKM
for each sample. For keeping the acyclic hierarchical annotation structure for accurate
resistome profiling at higher functional levels, ARGs were manually re-annotated based on
the drug class to which they confer resistance. ARGs belonging to macrolides, lincosamides
and streptogramins were congregated into the MLS class. While ARGs belonging to
carbapenem, cephamycin, cephalosporin, penem, penam and monobactam were congregated
into the Beta-lactam class.

Rarefaction analysis was performed using Rarefaction Analyzer [41] to assess whether the
samples had sufficient sequencing depth for the comprehensive characterization of the
nasopharyngeal resistome composition in preterm infants (Supplementary Fig. 3). This
analysis involved the random subsampling of sequencing reads at proportions varying from
0% to 100% of the total sequencing depth. The subsampled metagenomes were not utilized
for any further downstream analysis. Ten samples without identified ARGs were excluded
from downstream analysis (Supplementary Table 1). These samples were collected from 9
different infants at various time points and exhibited varying numbers of clean metagenomic
reads, as detailed in Supplementary Table 2. Collectively, we assessed the resistome of the
nasopharyngeal microbiome in a total of 181 samples collected from 36 preterm infants.

To profile the nasopharyngeal microbiota composition of preterm infants, we utilized
MetaPhlAn 3.0 software [42] (Supplementary Table 5). Since *Allobacillus halotolerans*, a
species included in the ZymoBIOMICS spike-in, was not initially represented in the default
MetaPhlAn database (*mpa v30 CHOCOPhlAn 201901*), we took an additional step to
incorporate its clade-specific marker genes into the database. Next, for the *in-silico*
identification of possible contaminants, we employed both the prevalence and frequency-
based (combined) methods of the decontam (v.1.16.0) R package [43] with the default
threshold (p-score = 0.1) on the raw microbiome and resistome count abundance tables.
Contaminants identification was performed independently within each batch of glycerol and

DNA extraction kit used for processing the nasopharyngeal samples from preterm infants. In
total, 28 microbial species, including the three ZymoBIOMICS spike-in species, and 4 ARGs
identified as contaminants, were filtered out from abundance tables prior to any downstream
analysis (Supplementary Table 6).

Statistical analysis and data visualization
All statistical analyses were conducted in R (v.4.2.1) within RStudio (v.2022.07.2+576) [44,
45]. The ARG abundance, annotation table, and metadata file were compiled into a single
data object using the phyloseq (v.1.40.0) package [46]. Figures, unless stated otherwise, were
created using the ggplot2 (v.3.4.0) package [47] and further edited using Adobe Illustrator
(v.16.0.0). α- and β-diversity analysis was performed using the vegan (v.2.6.2) [48] and
phyloseq R packages. α-diversity was calculated using the Shannon diversity metric, and
differences over time points in groups were assessed using a linear mixed-effects (LME)
model with infants set as a random effect while correcting for age. The overall resistome
composition of nasopharyngeal samples was visualized using principal component analysis
(PCA) ordination plots. The ordinates were based on the Aitchison distance with centered
log-ratio (CLR) transformed count abundance data. The variance explained by clinical
covariates in the overall resistome composition by clinical covariates was statistically
evaluated primarily with the Permutational multivariate analysis of variance
(PERMANOVA) test using the *adonis2* function (vegan package) with 999 permutations.
PERMANOVA is a powerful test used to analyze the effects of both categorical and/or
continuously distributed explanatory variables. It can accommodate complex study designs,
including multiple confounding factors as present in our study. However, it is not particularly
sensitive to heterogeneity in dispersion between the groups. To test the homogeneity of
multivariate dispersion, the permutational analysis of multivariate dispersion (PERMDISP)
test was used. In case of heterogeneity, the analysis of similarities (ANOSIM) test using the
anosim function (vegan package) was also employed to statistically re-assess the univariate
association of variables with the overall resistome composition. In the multivariable,
temporal PERMANOVA analyses, only the variables that individually showed a significant
association with resistome composition were included.

To identify the resistotypes, the Dirichlet multinomial mixture (DMM) model approach was
utilized using the DirichletMultinomial (v.1.38.0) R package [49]. Procrustes analysis was

performed to determine the association between microbiota and resistome composition based
on PCA using ARG and species relative abundance profiles. The symmetric Procrustes
correlation coefficients between the ordinations and P-values (*p*) were obtained through the
protest function from the vegan package. Spearman's correlation was applied to resistome
and microbiome relative abundance profiles for paired samples. To avoid the potential bias
introduced by Spearman's rank when ranking zero values, we removed ARGs and species
that were present in less than half of the samples. The correlation coefficient and
corresponding adjusted P-values (*p.adj*) were calculated using the *associate* function from the
microbiome (v.1.18.0) package [50]. Associations between individual taxa and overall
resistome outcomes were calculated using repeated measure correlation analysis from the
rmcorr (v.0.5.4) package while controlling for inter-individual variation [51]. Heatmaps were
created using the *pheatmap* function from the pheatmap R package (v.1.0.12) [52]. For
simple, independent comparisons of group differences, we used a one-way analysis of
variance test, Wilcoxon rank-sum test, Kruskal-Wallis test, or chi-square test as appropriate,
and we considered $p < 0.05$ to be significant. For all analyses involving multiple
comparisons, we applied the Benjamini-Hochberg (BH) method to correct for multiple
testing. The statistical analysis scheme used in the study is shown in Supplementary Fig. 2.
648

Ethics statement
Premature infant metadata and all nasopharynx aspirate samples used in this study were
collected at the Neonatal Intensive Care Unit (NICU) at Oslo University Hospital, Ullevål,
Norway, as part of the Born in the Twilight of Antibiotic project. The study received
approval from the Regional Committee for Medical and Health Research Ethics – South East,
Norway (2018/1381 REKD), and followed the principles of the Declaration of Helsinki
guidelines. Infants were enrolled in the study after obtaining written informed consent from
their parents.
657

DATA AVAILABILITY
The sequencing data (after filtering of human DNA) for all the samples that supports the
findings of this studies have been made available at NCBI SRA under BioProject ID:
PRJNA1009231
(https://dataview.ncbi.nlm.nih.gov/object/PRJNA1009231?reviewer=811m19u88kg6no8nnf2
eafjk99). Source data underlying the main as well as supplementary figures are provided in
this paper.

CODE AVAILABILITY

The software packages used in this manuscript are free and open source. No custom code was used in generating this manuscript. The analysis scripts used here are available from the authors upon reasonable request.

REFERENCES

1. Murray, C., et al., *& Naghavi, M.(2022). Global burden of bacterial antimicrobial*
resistance in 2019: a systematic analysis. The Lancet. **399**(10325): p. 629-655.
2. Von Wintersdorff, C.J., et al., *Dissemination of antimicrobial resistance in microbial*
ecosystems through horizontal gene transfer. Frontiers in microbiology, 2016. **7**: p.
173.
3. Kilian, M., et al., *Parallel evolution of Streptococcus pneumoniae and Streptococcus*
mitis to pathogenic and mutualistic lifestyles. MBio, 2014. **5**(4): p. e01490-14.
4. Pailhoriès, H., et al., *Antibiotic resistance in chronic respiratory diseases: from*
susceptibility testing to the resistome. European Respiratory Review, 2022. **31**(164).
5. Teo, S.M., et al., *The infant nasopharyngeal microbiome impacts severity of lower*
respiratory infection and risk of asthma development. Cell host & microbe, 2015.
17(5): p. 704-715.
6. Toivonen, L., et al., *Antibiotic treatments during infancy, changes in nasal microbiota,*
and asthma development: population-based cohort study. Clinical Infectious
Diseases, 2021. **72**(9): p. 1546-1554.
7. Li, X., et al., *The infant gut resistome associates with E. coli, environmental*
exposures, gut microbiome maturity, and asthma-associated bacterial composition.
Cell Host & Microbe, 2021. **29**(6): p. 975-987. e4.
8. Rajar, P., et al., *Microbial DNA extraction of high-host content and low biomass*
samples: Optimized protocol for nasopharynx metagenomic studies. Frontiers in
Microbiology, 2022. **13**.
9. Zhang, A.-N., et al., *An omics-based framework for assessing the health risk of*
antimicrobial resistance genes. Nature communications, 2021. **12**(1): p. 4765.
10. Long, Y., et al., *High Carriage Rate of the Multiple Resistant Plasmids Harboring*
Quinolone Resistance Genes in Enterobacter spp. Isolated from Healthy Individuals.
Antibiotics, 2022. **11**(1): p. 15.
11. Flynn, M. and J. Dooley, *The microbiome of the nasopharynx.* Journal of Medical
Microbiology, 2021. **70**(6).
12. Man, W.H., W.A. de Steenhuijsen Piters, and D. Bogaert, *The microbiota of the*
respiratory tract: gatekeeper to respiratory health. Nature Reviews Microbiology,
2017. **15**(5): p. 259-270.
13. Consortium, H.M.P., *Structure, function and diversity of the healthy human*
microbiome. Nature, 2012. **486**(7402): p. 207-214.
14. Manenzhe, R.I., et al., *Longitudinal changes in the nasopharyngeal resistome of*
South African infants using shotgun metagenomic sequencing. PloS one, 2020. **15**(4):
p. e0231887.
15. Parnanen, K., et al., *Maternal gut and breast milk microbiota affect infant gut*
antibiotic resistome and mobile genetic elements. Nat Commun, 2018. **9**(1): p. 3891.
16. Yassour, M., et al., *Natural history of the infant gut microbiome and impact of*
antibiotic treatment on bacterial strain diversity and stability. Science translational
medicine, 2016. **8**(343): p. 343ra81-343ra81.
17. Parker, L.A., et al., *Neonatal feeding tube colonization and the potential effect on*
infant health: a review. Frontiers in Nutrition, 2022. **9**: p. 775014.
18. Gasparrini, A.J., et al., *Persistent metagenomic signatures of early-life hospitalization*
and antibiotic treatment in the infant gut microbiota and resistome. Nature
microbiology, 2019. **4**(12): p. 2285-2297.

19.   Rahman, S.F., et al., *Machine learning leveraging genomes from metagenomes*
identifies influential antibiotic resistance genes in the infant gut microbiome.
MSystems, 2018. **3**(1): p. e00123-17.

20.   Sukumar, S., et al., *The development of the oral resistome during the first decade of*
life. 2022.

21.   Tacconelli, E., *Global Priority List of Antibiotic-Resistant Bacteria to Guide Research,*
Discovery, and Development. 2017.

22.   Alcock, B.P., et al., *CARD 2020: antibiotic resistome surveillance with the*
comprehensive antibiotic resistance database. Nucleic acids research, 2020. **48**(D1):
p. D517-D525.

23.   Thompson, S.A., et al., *Novel tetracycline resistance determinant isolated from an*
environmental strain of Serratia marcescens. Applied and environmental
microbiology, 2007. **73**(7): p. 2199-2206.

24.   Matsumura, N., S. Minami, and S. Mitsuhashi, *Sequences of homologous β-*
lactamases from clinical isolates of Serratia marcescens with different substrate
specificities. Antimicrobial agents and chemotherapy, 1998. **42**(1): p. 176-179.

25.   Shaw, K., et al., *Characterization of the chromosomal aac (6')-Ic gene from Serratia*
marcescens. Antimicrobial agents and chemotherapy, 1992. **36**(7): p. 1447-1455.

26.   Piccirilli, A., et al., *Molecular Characterization by Whole-Genome Sequencing of*
Clinical and Environmental Serratia marcescens Strains Isolated during an Outbreak
in a Neonatal Intensive Care Unit (NICU). Diagnostics, 2022. **12**(9): p. 2180.

27.   Cristina, M.L., M. Sartini, and A.M. Spagnolo, *Serratia marcescens infections in*
neonatal intensive care units (NICUs). International journal of environmental
research and public health, 2019. **16**(4): p. 610.

28.   WHO, I., *Global priority list of antibiotic-resistant bacteria to guide research,*
discovery, and development of new antibiotics. Who, 2017. **2017**: p. 1-7.

29.   Reyman, M., et al., *Effects of early-life antibiotics on the developing infant gut*
microbiome and resistome: a randomized trial. Nat Commun, 2022. **13**(1): p. 893.

30.   Lebeaux, R.M., et al., *The infant gut resistome is associated with E. coli and early-life*
exposures. BMC Microbiol, 2021. **21**(1): p. 201.

31.   Gibson, M.K., et al., *Developmental dynamics of the preterm infant gut microbiota*
and antibiotic resistome. Nature Microbiology, 2016. **1**: p. 16024.

32.   Leggett, R.M., et al., *Rapid MinION profiling of preterm microbiota and antimicrobial-*
resistant pathogens. Nature Microbiology, 2020. **5**(3): p. 430-442.

33.   Eisenhofer, R., et al., *Contamination in low microbial biomass microbiome studies:*
issues and recommendations. Trends in microbiology, 2019. **27**(2): p. 105-117.

34.   Krueger, F., *Trim Galore: a wrapper tool around Cutadapt and FastQC.* Trim Galore,
2012.

35.   Langmead, B. and S.L. Salzberg, *Fast gapped-read alignment with Bowtie 2.* Nature
methods, 2012. **9**(4): p. 357-359.

36.   Li, H., et al., *The sequence alignment/map format and SAMtools.* bioinformatics,
2009. **25**(16): p. 2078-2079.

37.   Quinlan, A.R. and I.M. Hall, *BEDTools: a flexible suite of utilities for comparing*
genomic features. Bioinformatics, 2010. **26**(6): p. 841-842.

38.   Andrews, S., *FastQC: a quality control tool for high throughput sequence data.* 2010,
Babraham Bioinformatics, Babraham Institute, Cambridge, United Kingdom.

39.   Ewels, P., et al., *MultiQC: summarize analysis results for multiple tools and samples in*
765   *a single report.* Bioinformatics, 2016. **32**(19): p. 3047-3048.

766   40.   D'Souza, A.W., et al., *Destination shapes antibiotic resistance gene acquisitions,*
767   *abundance increases, and diversity changes in Dutch travelers.* Genome medicine,
768   2021. **13**: p. 1-21.

769   41.   Doster, E., et al., *MEGARes 2.0: a database for classification of antimicrobial drug,*
770   *biocide and metal resistance determinants in metagenomic sequence data.* Nucleic
771   acids research, 2020. **48**(D1): p. D561-D569.

772   42.   Beghini, F., et al., *Integrating taxonomic, functional, and strain-level profiling of*
773   *diverse microbial communities with bioBakery 3.* elife, 2021. **10**: p. e65088.

774   43.   Davis, N.M., et al., *Simple statistical identification and removal of contaminant*
775   *sequences in marker-gene and metagenomics data.* Microbiome, 2018. **6**: p. 1-14.

776   44.   Allaire, J., *RStudio: integrated development environment for R.* Boston, MA, 2012.
777   **770**(394): p. 165-171.

778   45.   R Core Team, R. and R.C. Team, *R: a language and environment for statistical*
779   *computing. R Foundation for Statistical Computing; 2020.* 2021.

780   46.   McMurdie, P.J. and S. Holmes, *phyloseq: an R package for reproducible interactive*
781   *analysis and graphics of microbiome census data.* PloS one, 2013. **8**(4): p. e61217.

782   47.   Wickham, H., *ggplot2.* Wiley interdisciplinary reviews: computational statistics, 2011.
783   **3**(2): p. 180-185.

784   48.   Oksanen, J., et al., *Vegan: Community Ecology Package (R Package Version 2.6-2).*
785   *2022.*

786   49.   Holmes, I., K. Harris, and C. Quince, *Dirichlet multinomial mixtures: generative*
787   *models for microbial metagenomics.* PloS one, 2012. **7**(2): p. e30126.

788   50.   Lahti, L. and S. Shetty, *Introduction to the microbiome R package.* 2018.

789   51.   Bakdash, J.Z. and L.R. Marusich, *Repeated measures correlation.* Frontiers in
790   psychology, 2017. **8**: p. 456.

791   52.   Kolde, R. and M.R. Kolde, *Package 'pheatmap'.* R package, 2018. **1**(10).

ACKNOWLEDGEMENTS

We would like to thank the parents of participating infants, and the clinical staff at the

Neonatal Intensive Care units at Ullevål hospital for their assistance with sample collection.

The sequencing service for this work was provided by the Norwegian Sequencing Centre

(www.sequencing.uio.no) and the computations were performed on resources provided by

Sigma2 - the National Infrastructure for High Performance Computing and Data Storage in

Norway.

AUTHOR CONTRIBUTIONS

FCP conceptualized and designed the study, acquired funding, and contributed to data

interpretation. AD conducted bioinformatics and statistical analysis, performed data

interpretation, and wrote the manuscript with input from FCP. PR and KH were responsible

for patient inclusion, sampling, and metadata analysis. PR, GS, and HAA conducted sample

processing, metagenomic DNA extraction, and library preparation for sequencing. DB, ODS,

DF, GG, UD, and KH contributed to study design, provided critical revisions on the

manuscript. All authors contributed to manuscript review and provided valuable feedback

during the revision process.

FUNDING

This work was financed by the Norwegian Research Council (NFR) (project numbers:

273833 and 299159), the Olav Thon Foundation (project number: 421258), and the Faculty

of Dentistry at the University of Oslo.

COMPETING INTERESTS

The authors declare that they have no financial or non-financial competing interests.

 FIGURE LEGENDS

Fig. 1: Resistome composition in the nasopharynx microbiome of preterm infants.

Stacked bar plots showing the relative abundance of (A) antibiotic resistance genes (ARGs)

clustered at the Class level for the naive and antibiotic-treated groups over time; and (B) the

top 30 most abundant ARGs identified across samples at baseline (T1) in early antibiotic-

treated (left) and naive (right) groups. All other ARGs are grouped into the "Others"

category. MLS = macrolide-lincosamide-streptogramin. Source data are provided as a Source

Data file.

Fig. 2: Impact of early antibiotic treatment on nasopharynx total ARG abundance and

ARG α-diversity Boxplots showing the (A) total ARG abundance (sum of RPKM) and (B)

ARG Shannon diversity per sample, stratified by antibiotic treatment and time points. The

lower and upper horizontal box lines correspond to the first and third quartiles, with the line

in the middle of the boxes representing the median. Whiskers extending from the boxes

represent the range of data points within 1.5 times the interquartile range below the first

quartile and within 1.5 times the interquartile range above the third quartile. (Statistical

significance: LME mixed effect model. Adjusted p-values (p): $*p < .05$, $**p < .01$ and $***p <$

$.001$). RPKM = reads per kilobase of reference gene per million bacterial reads. Source data

are provided as a Source Data file.

Fig. 3: β-diversity PCA plots of overall nasopharyngeal resistome composition stratified

per time point. Principal component analysis (PCA) based on Euclidean distances of CLR-

transformed ARG abundances (*i.e.,* Aitchison dissimilarity) between samples, visualizing the

overall nasopharyngeal resistome composition stratified for antibiotic-treated infants and

naive at time point (A) T1 (baseline) and (B) T2 (immediately after cessation of early

antibiotic therapy). Each data point represents the resistome composition of one sample. The

ellipses represent the 95% confidence interval around the centroid of resistome composition

for each group. Statistical significance (adjusted p-values) and effect size (R^2) of the

differences in beta diversity were assessed using PERMANOVA. In panel (C), the results of

the PERMANOVA-tests for the later time points are presented. Data points of early

antibiotic-treated infants are colored red and those of naive infants are blue. PERMANOVA

= permutational multivariate analysis of variance; *p.adj* = adjusted p-values. Source data are

provided as a Source Data file.

Fig. 4: Factors influencing the overall nasopharyngeal resistome composition in the
preterm infants. PCA visualizations of Aitchison dissimilarities between all the
nasopharyngeal samples from birth until 6 months corrected age at ARG level. Each data
point corresponds to a sample, colored by (A) Infant, (B) PMA, (C) Time points and (D)
Prenatal and post-natal antibiotic exposure and shaped according to early antibiotic exposure
or naive groups. Effect sizes (R^2) and corresponding adjusted p-values (*p.adj*, corrected using
the Benjamini-Hochberg method) are calculated using PERMANOVA-tests (as shown in the
bottom right of each plot). Percentage refers to the percentage of variance explained by the
principal component. PMA = postmenstrual age; PC1 = principal component 1; PC2 =
principal component 2; PERMANOVA = permutational multivariate analysis of variance;
p.adj = adjusted p-values. Source data are provided as a Source Data file.

Fig. 5: Association between resistotypes, community types, and ARG abundance
profiles. The PCA plot showing the association between overall ARG abundance profile with
respect to their (A) resistotypes and (C) community types identified among all the
nasopharyngeal samples using the DMM models. R^2 was obtained using the PERMANOVA
test. (B) Samples proportions for each resistotype shown as a function of community type.
DMM = dirichlet multinomial mixture; Strep = *Streptococcus* ; Staph = *Staphylococcus*.
Source data are provided as a Source Data file.

Fig. 6: The resistome variation in nasopharynx is explained by microbiome variation.
The ARG and microbial abundance profiles were correlated with each other for antibiotic-
treated (A) and naive (B) groups using Procrustes analysis. The length of line connecting two
points denotes the degree of dissimilarity or distance between the microbiome (filled square)
and resistome (filled circle) composition of the same sample. The line and points are colored
based on time points in both groups. The correlation coefficients and significance were
calculated using the *protest* function in vegan package. (C) Heatmap depicts the Spearman's
pairwise correlation coefficients (*r*) between individual ARGs and microbial species from
nasopharyngeal samples. Hierarchical clustering using Euclidean distance was performed on
both rows and columns. Correlation coefficients (r) are shown only where adjusted p-
value $<$ 0.05. (D) Timeline of ARGs associated with *S. marcescens* in the nasopharynx
samples of each preterm infant across time points. Each ARG is represented by a different
color and shape. Source data are provided as a Source Data file.

Supplementary Fig. 1: Sample collection and antibiotic exposure for preterm infants in
this study. Sampling timeline of 198 nasopharyngeal samples collected from 36 preterm
infants over first six months (corrected age), together with information regarding the time of
early antibiotic exposure (red and blue colored circles), exposure to subsequent antibiotic
treatment (green colored circles) and included/excluded samples. The x-axis represents the
days of life (i.e., corrected age) and the y-axis represents the infants. NP = nasopharyngeal;
AB□=□antibiotics.

Supplementary Fig. 2: Statistical analysis design used in the study. ARGs = antibiotic
resistance genes; PERMANOVA = permutational multivariate analysis of variance;
PERMDISP = permutational analysis of multivariate dispersion; ANOSIM = analysis of
similarities.

Supplementary Fig. 3: Rarefaction curve for the number of unique ARGs identified at
subsampled sequencing depth. The red line indicates the median sequencing depth for all
samples, while the gray shading represents one standard deviation. Source data are provided
as a Source Data file.

Supplementary Fig. 4: Mean normalized relative abundances of different ARG mechanisms
found in the nasopharynx of early antibiotic-treated and naive preterm infants at different
time points (T1-T6). **Source data are provided as a Source Data file.**

Supplementary Fig. 5: Heatmap showing the log10-transformed RPKM abundance of
potential high-risk antibiotic resistance genes (ARGs) identified in the nasopharynx of all
preterm infants across samples. The x-axis represents the names of samples, which are
arranged by treatment group and time points. RPKM = reads per kilobase of reference gene
per million bacterial reads. Source data are provided as a Source Data file.

Supplementary Fig. 6: PCA visualizations of beta diversity analysis using the Aitchison
distances at time points: (A) T1, (B) T2 and (C) T6 (6 month corrected age). Each point
corresponds to a sample colored based on birth weight group (A, C) and prenatal and
postnatal antibiotic exposure (B). Ellipses indicate 95% confidence intervals (CI). Adjusted
p-values and effect size (R^2) are based on PERMANOVA test. CLR = centered-log ratio;
BW= Birth Weight. Source data are provided as a Source Data file.

Supplementary Fig. 7: Box plots of the relative abundances of the main ARG contributors
are shown for each Resistotype cluster identified across all the preterm infants'
nasopharyngeal resistomes using DMM models. The X-axis displays the three identified
clusters, and the Y-axis represents the relative abundance of the ARGs. DMM = dirichlet
multinomial mixture. Source data are provided as a Source Data file.

Supplementary Fig. 8: Scatterplots showing the association between correlated species and
overall resistome outcomes across all infant samples from birth to 6 months corrected age.
The total ARG abundance in RPKM was positively correlated with the relative abundance of
Streptococcus mitis (A) and non-linearly negatively associated with the relative abundance of
Serratia marcescens (C). Plots (B) and (D) depict the relationship between the number of
unique ARGs and the relative abundance of *Streptococcus mitis* and *Serratia marcescens*,
respectively. Plot (A) and (C) are colored by the relative abundance of *Streptococcus mitis*,
while plots (B) and (D) are colored by the relative abundance of *Serratia marcescens*. RPKM
= reads per kilobase of reference gene per million bacterial reads. Source data are provided as
a Source Data file.

Supplementary Fig. 9: Stacked bar plots showing the relative abundance of top 30 most
abundant bacterial species identified across samples in the antibiotic-treated (top) and naive
(bottom) groups over sampling time points (T1-T6). All other low abundant species are
merged into the "Others" category.

Fig. 1

A

B

Fig. 2

Fig. 3

Time points	R^2 (%)	p.adj
T3	3.4%	0.345
T4	4.4%	0.148
T5	5.1%	0.371
T6	3.6%	0.355

Fig. 4

Fig. 5

A

PCA2 (17.6%) vs PCA1 (18.7%)

Resistotypes
- R1
- R2
- R3

Group
- ■ Early AB
- ● Naive

R^2 = 20.04%
$p.adj$ = 0.001

B

Sample Proportions vs Resistotypes (R1, R2, R3)

Community types
- 1 (Strep)
- 2 (Strep_Staph)
- 3 (Serratia)
- 4 (Strep_Gemella)

C

PCA2 (17.6%) vs PCA1 (18.7%)

Community types
- 1
- 2
- 3
- 4

Group
- ■ Early AB
- ● Naive

R^2 = 13.5%
$p.adj$ = 0.001

Fig. 6