The landscape of antibiotic resistance genes in the nasopharynx of preterm

infants: Prolonged signature of hospitalization and effects by antibiotics

Achal Dhariwal[1]†, Polona Rajar[1,2]†, Gabriela Salvadori[1], Heidi Aarø Åmdal[1], Dag Berild[3,4], Ola

Didrik Saugstad[5], Drude Fugelseth[2,4], Gorm Greisen[6], Ulf Dahle[7], Kirsti Haaland[2], Fernanda

Cristina Petersen[1]*

[1] Institute of Oral Biology, Faculty of Dentistry, University of Oslo, Oslo, Norway.

[2] Department of Neonatal Intensive Care, Division of Pediatric and Adolescent Medicine, Oslo

University Hospital Ullevål, Oslo, Norway.

[3] Department of Infectious Diseases, Oslo University Hospital, Oslo, Norway.

[4] Institute of Clinical Medicine, Faculty of Medicine, Oslo University, Oslo, Norway.

[5] Department of Pediatric Research, Oslo University Hospital, University of Oslo, Oslo, Norway.

[6] Department of Neonatology, Copenhagen University Hospital Rigshospitalet, Copenhagen,

Denmark.

[7] Centre for Antimicrobial Resistance, Norwegian Institute of Public Health, Oslo, Norway

†Shared First authorship

*Correspondence: Fernanda Cristina Petersen

Institute of Oral Biology

Faculty of Dentistry,

University of Oslo, Oslo, Norway.

Tel: +47-22840312

E-mail: f.c.petersen@odont.uio.no

ABSTRACT

Respiratory pathogens, commonly colonizing nasopharynx, are among the leading causes of death due to antimicrobial resistance. Yet, antibiotic resistance determinants within nasopharyngeal microbial communities remain poorly understood. Utilizing shotgun metagenomics, we investigated the nasopharynx resistome development in preterm infants, assessed early antibiotic impact on its trajectory, and explored its association with clinical covariates. Our findings revealed widespread nasopharyngeal carriage of antibiotic resistance genes (ARGs) with resistomes undergoing transient changes, including increased ARG diversity, abundance, and composition alterations due to early antibiotic exposure. ARGs associated with the critical nosocomial pathogen *Serratia marcescens* persisted (at 8-10 months of age) for months following discharge, signifying hospitalization signature. The nasopharyngeal resistome strongly correlated with microbiome composition, with inter-individual differences and postnatal age explaining most of the variation. The collateral effects of antibiotics and hospitalization reported here highlights the importance of future studies focusing on this yet rather unexplored reservoir of pathogens and ARGs.

Keywords: respiratory resistome, metagenomics, nasopharynx, microbiome, preterm infants, antibiotics, antimicrobial resistance

INTRODUCTION

Antimicrobial resistant infections are one of the biggest global health threats. Among these, respiratory infections account for more than 1.5 million deaths a year, surpassing the number of deaths by resistant infections in any other human body niches [1]. Main pathogens include *Streptococcus pneumoniae, Staphylococcus aureus, Klebsiella pneumoniae, Enterococcus faecalis* and *Haemophilus influenzae*, all generally present in the nasopharynx of healthy populations. These pathogens share common niches with commensals, which in turn provide colonization resistance against pathogens. However, commensals can also serve as reservoirs of antibiotic resistance genes (ARGs) that can readily be transferred to pathogens [2-4].

Recent studies have highlighted the potential negative impact of antibiotic treatment on the nasopharynx microbiome and its association with an increased risk of developing asthma and respiratory infections in infants [5-7]. These studies have significantly advanced our understanding of the ecological impact of antibiotics on respiratory health and disease. However, as these studies rely on 16S rRNA sequencing technologies, they do not offer a high level of microbial taxonomic resolution and lack information on the overall gene composition, including the presence of ARGs within microbial communities (referred to as the resistome). In our previous study, we developed an optimization strategy for DNA extraction and library preparation that helped to circumvent the challenges posed by low biomass and high host DNA content in nasopharynx samples of preterm infants [8]. In this study, we applied the optimized protocol to nasopharyngeal aspirate samples to characterize the development of the resistome in 36 preterm infants from birth up to 6 months corrected age (8-10 months of chronological age). By analyzing ~1.85 terabytes (Tb) of metagenomic DNA data generated from deep shotgun sequencing, we obtained the most comprehensive information to date on the characteristics and dynamics of resistome development in the respiratory tract of preterm infants. Our hypothesis was that early exposure to broad-spectrum antibiotics for treatment of suspected early onset sepsis (sEONS) in preterm infants may promote prolonged compositional alterations in the respiratory resistome. We also explored the extensive metadata collected to gain insights into additional clinical covariates that may be associated with resistome development.

METHODS

Study design and cohort characteristics

The study is a prospective observational cohort study. Between July 2019 and January 2021, we included preterm infants born (or transferred within 48 hours) at the NICU (Oslo University Hospital, Ullevål) who were between 28+0 and 31+6 weeks of gestational age (GA). In total, 66 preterm infants were enrolled in the study, and followed from birth until six months corrected age (i.e., adjusted for prematurity) during their stay in the NICU and after discharge. Five of the infants were lost to follow-up. Extensive metadata and covariates from infants and mothers, including mode of delivery, gestational age (GA) at birth, postmenstrual age (PMA), gender, birth weight (BW), postnatal infant antibiotics usage (duration, type, days), maternal antibiotic usage, feeding regimen during hospitalization and after discharge, and other demographic characteristics, were collected through medical records. This data was stored in secured server at the Services for Sensitive Data (TSD), University of Oslo.

In our cohort of 66 infants, 33 infants with sEONS received broad-spectrum antibiotics, consisting of a combination of ampicillin (a beta-lactam antibiotic) and gentamicin (an aminoglycoside antibiotic), within 24 hours after birth. Due to the circumstances of premature birth and the instability of the infants, baseline samples could not be collected before antibiotic therapy initiation. Instead, the first nasopharyngeal aspirate samples (baseline) were obtained within 48 hours after the initiation of early antibiotic therapy. The average early antibiotic treatment duration was 5 days (SD: 2 days). The sampling time point T2 corresponds to the first available sample collected (except in one infant) after the cessation of early antibiotic treatment. In addition to early antibiotics, one of the included infants received subsequent antibiotics initiated > 72 hours after birth and before discharge from NICU and two other infants received antibiotic therapy between discharge from NICU and 6 months corrected age. A timeline of sample collection and antibiotic administration is represented in Supplementary Fig. 1.

To investigate temporal changes in the resistome in response to antibiotics and control for potential baseline differences as confounding factors, this study included only infants who had nasopharyngeal samples collected on the same day as the initiation of early antibiotic treatment (*n = 15*) and antibiotic-naive infants as controls (*n = 21*). Infants exposed to early antibiotics had significantly lower GA and BW, and their mothers had a more prolonged period of ruptured

membranes before delivery compared to antibiotic-naive infants. None of the mothers of naive

infants received antibiotics during pregnancy (prenatal antibiotics) before onset of delivery. The

baseline characteristics of the infants stratified by antibiotic exposure are presented in Table 1.

Sample collection

Nasopharyngeal aspirate samples were collected from infants at six distinct time points spanning

from birth to 6 months corrected age (T1-T6). The days of life (DOL) at which samples were

collected for each time point varied as follows: T1 (DOL: 0-3), T2 (DOL: 4-9), T3 (DOL: 12-19),

T4 (DOL: 20-31), T5 (DOL: 32-76) and T6 at 6 months corrected age (DOL: 229-288). Additional

samples were taken if the antibiotic therapy was initiated or terminated more than 48 hours

before/after a predefined sampling time. As a result, in two infants, there was more than one sample

available for the defined time points. To facilitate time point-wise analysis, only one sample was

included for each of the time points (Supplementary Table 1 & Supplementary Fig. 1). In cases

where the sample could not be collected at the scheduled time points (such as with an unstable

infant), a new sample was obtained as soon as possible. Infants who were transferred to another

hospital before discharged to home adhered to the study protocol. When the infants reached six

months of corrected age, they were either invited to an outpatient visit at Ullevål or their local

hospital or visited at home for sample collection. Nasopharynx aspirates were collected as

previously described [8] and stored at -80°C until further processing. In total, we collected and

extracted metagenomic DNA from 198 nasopharyngeal samples obtained from 36 preterm infants,

from day of birth until 6 months corrected age. Detailed information on sample exclusion and

inclusion statistics at different steps from sampling to whole metagenomic sequencing (WMS) is

described in Supplementary Table 1.

Metagenomic DNA extraction and sequencing

Metagenomic DNA was extracted by first thawing the aspirate samples (2 ml) on ice, pelleting

them by centrifugation at 10,000 g for 10 minutes at 4°C, and then using MolYsis™ for host DNA

depletion (Molzym, Bremen, Germany), following the manufacturer's protocol with some

modifications, as previously described [8]. After depletion of human DNA, fresh pellets were

spiked with 20 µl of ZymoBIOMICS Spike-in Control II (catalog number: D6321 & D6321-10),

and bacterial DNA was extracted using the MasterPure™ Gram Positive DNA Purification Kit

(Epicentre, Madison, WI, USA), according to the manufacturer's recommendations. The bacterial

DNA was eluted in 35 µl of TE buffer and stored at -80°C until further use. Final DNA

concentration was quantified using Qubit dsDNA HS assay kits in a Qubit 4.0 Fluorometer

(Invitrogen, Thermo Fisher Scientific, USA). Metagenomic sequencing libraries were constructed

using the Nextera DNA Flex library preparation kit (Illumina Inc., CA, USA). All samples were

subjected to paired-end sequencing at 2 × 150 bp on an Illumina NovaSeq S4 high-output platform

at the Norwegian Sequencing Centre (Oslo, Norway). In total, 6.14 billion raw sequencing reads

were obtained with a mean value of sample reads counts of 33.97 million (M) ranging from 4.19

to 69.25 M reads (Supplementary Table 2).

Controls for low microbial biomass

The study followed recent recommendations for low microbial biomass microbiome research [9].

This involved the following steps: (1) collecting samples from the included infants in the two

groups within the same time period, (2) using standard protection measures during sample

collection (such as clean suits, disposable gloves, and face masks), (3) including sampling and

DNA extraction blank controls (negative controls), (4) conducting DNA extraction and library

preparation while wearing laboratory coats and disposable gloves in a carefully cleaned

environment , and (5) using unique barcodes in library preparation. The negative controls had no

or extremely low DNA concentrations to proceed to library preparation as determined by RT-PCR

using the Femto Bacterial DNA Quantification Kit from Zymo and were therefore excluded from

further analysis ($n =10$; mean = 0.00043 ng per µL). Additionally, spike-in controls ($n = 6$) were

extracted and used as positive controls, and longitudinal samples from each preterm infant were

extracted within the same run.

Bioinformatics processing

The raw metagenomic sequencing data was pre-processed using our in-house bioinformatics

pipelines using a high-performance computing cluster inside a secured environment, i.e., TSD at

the University of Oslo. In brief, the Nextera adaptor sequences and low-quality reads were filtered

using Trim Galore (v.0.6.1) with default parameters [10]. Next, the human DNA contaminant

sequences were identified and removed by mapping the quality-trimmed reads against the human

reference genome (GRCh38) using Bowtie2 (v.2.3.4.2) [11] (non-default parameters: *q -N 1 -k 1*

--fr --end-to-end --phred33 --very-sensitive –no-discordant) along with SAMtools (v.1.9) [12] and

BEDTools (v.2.27.1) [13]. On average, 46.71% (range: 1.09 - 97.69%) of total quality-filtered

reads were identified as belonging to the human genome. After removal of the human-associated

metagenomic reads, a total of 2.98 billion remaining high-quality clean reads across all samples

with number of reads ranging from 0.76 to 59.9 M per sample, and with median of 16.4 M reads

were subjected to resistome profiling. More detailed results regarding the overall sequence

preprocessing are presented in Supplementary Table 2. The sequence quality reports on raw and

processed reads were generated using FastQC (v.0.11.9) [14] and summarized using MultiQC

(v.1.7) [15].

Resistome profiling

To characterize the presence of ARGs, cleaned high-quality reads for each sample were mapped

against the *nucleotide_fasta_protein_homolog_model* from the Comprehensive Antibiotic

Resistance Database (CARD) (v.3.2.2) [16] using Bowtie2 under parameter *–very-sensitive-local*.

The mapped reads from each sample were then filtered, sorted, and indexed using SAMtools. The

number of reads mapped to each ARG was calculated using SAMtools *idxstats* and BEDTools.

ARGs with a coverage of at least 80% from the alignment were retained for further downstream

analyses. The mapped read counts were normalized for bacterial sequence abundances and gene

lengths by calculating reads per kilobase of reference gene per million bacterial reads (RPKM) for

each CARD reference sequence. The relative abundance of ARGs for each sample was computed

by dividing the RPKM by the total sum of the RPKM for each sample. For keeping the acyclic

hierarchical annotation structure for accurate resistome profiling at higher functional levels, ARGs

were manually re-annotated based on the drug class to which they confer resistance. ARGs

belonging to macrolides, lincosamides and streptogramins were congregated into the MLS class.

While ARGs belonging to carbapenem, cephamycin, cephalosporin, penem, penam and

monobactam were congregated into the Beta-lactam class.

Rarefaction analysis was performed using Rarefaction Analyzer [17] to assess whether the samples

had sufficient sequencing depth for the comprehensive characterization of the nasopharyngeal

resistome composition in preterm infants (Supplementary Fig. 3). This analysis involved the

random subsampling of sequencing reads at proportions varying from 0% to 100% of the total

sequencing depth. The subsampled metagenomes were not utilized for any further downstream

analysis. Ten samples were excluded from downstream analysis as no reads were assigned to

ARGs. Collectively, we assessed the resistome of the nasopharyngeal microbiome in a total of 181

samples collected from 36 preterm infants.

Statistical analysis and data visualization

All statistical analyses were conducted in R (v.4.2.1) within RStudio (v.2022.07.2+576) [18, 19].

The ARG abundance, annotation table, and metadata file were compiled into a single data object

using the phyloseq (v.1.40.0) package [20]. Figures, unless stated otherwise, were created using

the ggplot2 (v.3.4.0) package [21] and further edited using Adobe Illustrator (v.16.0.0). α- and β-

diversity analysis was performed using the vegan (v.2.6.2) [22] and phyloseq R packages. α-

diversity was calculated using the Shannon diversity metric, and differences over time points in

groups were assessed using a linear mixed-effects (LME) model with infants set as a random effect

while correcting for age. The overall resistome composition of nasopharyngeal samples was

visualized using principal component analysis (PCA) ordination plots. The ordinates were based

on the Euclidean dissimilarity matrix of centered log-ratio (CLR) transformed relative abundance

data (i.e., Aitchison distance). Associations of clinical covariates with the dispersion of sample

resistome compositions were statistically evaluated with the Permutational multivariate analysis

of variance (PERMANOVA) test using the *adonis2* function (vegan package) with 999

permutations. To test the homogeneity of multivariate dispersion, the permutational analysis of

multivariate dispersion (PERMDISP) test was used. In case of heterogeneity, the analysis of

similarities (ANOSIM) test using the *anosim* function (vegan package) was further used to

statistically re-assess the association of variables with the overall resistome composition. In the

multivariable, temporal analyses, only the variables that individually showed a significant

association with resistome composition were included.

To identify the resistotypes, the Dirichlet multinomial mixture (DMM) model approach was

utilized using the DirichletMultinomial (v.1.38.0) R package [23]. Procrustes analysis was

performed to examine the effect of underlying microbiota on the resistome. The ARG RPKM

abundance matrix and the species-level count per million (CPM) abundance matrix were Hellinger-transformed and ordinated using PCoA (Principal coordinates analysis) on Bray–Curtis dissimilarities. The symmetric Procrustes correlation coefficients between the ordinations and P-values were obtained through the *protest* function from the vegan package. Spearman's correlation was applied to resistome and microbiome relative abundance profiles for paired samples. To avoid the potential bias introduced by Spearman's rank when ranking zero values, we removed ARGs and species that were present in less than half of the samples. The correlation coefficient and corresponding adjusted p-values were calculated using the associate function from the microbiome (v.1.18.0) package [24]. Associations between individual taxa and overall resistome outcomes were calculated using repeated measure correlation analysis from the rmcorr (v.0.5.4) package while controlling for inter-individual variation [25]. Heatmaps were created using the *pheatmap* function from the pheatmap R package (v.1.0.12) [26]. For simple, independent comparisons of group differences, we used a one-way analysis of variance test, Wilcoxon rank-sum test, Kruskal-Wallis test, or chi-square test as appropriate, and we considered $p < 0.05$ to be significant. For all analyses involving multiple comparisons, we applied the Benjamini-Hochberg (BH) method to correct for multiple testing. The statistical analysis scheme used in the study is shown in Supplementary Fig. 2.

Ethics statement

Premature infant metadata and all nasopharynx aspirate samples used in this study were collected at the Neonatal Intensive Care Unit (NICU) at Oslo University Hospital, Ullevål, Norway, as part of the Born in the Twilight of Antibiotic project. The study received approval from the Regional Committee for Medical and Health Research Ethics – South East, Norway (2018/1381 REKD), and followed the principles of the Declaration of Helsinki guidelines. Infants were enrolled in the study after obtaining written informed consent from their parents.

RESULTS

Distribution and characterization of ARGs in the preterm infant nasopharyngeal microbiome

In total, we found 373 ARGs belonging to 15 ARG classes conferring resistance via 5 distinct mechanisms. Multidrug efflux mediated ARGs comprised the majority of the resistome with an average of 27% at T1 to T5 and 17.8% at T6 (six months corrected age) of the total relative abundance. There was a median of 18 (mean: 24) unique ARGs ranging from 1 to 157 per analyzed sample. On average, maximum number of ARGs were detected after the discontinuation of treatment (T2) (mean: 40) in most infants receiving early antibiotics treatment. In the non-treated group, the corresponding mean value at T2 was 19. Only 13 ARGs were both most prevalent (detected in > 40% of all samples) and highly abundant with mean relative abundance of 64.8% (SD: 32.06%) across all preterm infant nasopharyngeal resistome. These core set of resistance genes included multidrug resistance genes (*acrB, oqxB, mexI),* macrolide-lincosamide-streptogramin (MLS) resistance genes (*mel, pmrA, rlmA(II)*), two beta-lactam resistance genes (*blaZ, SST-1),* fluoroquinolone resistance genes (*patA, patB),* tetracycline resistance genes (*tet(41), tetM*), and the aminoglycoside resistance gene *AAC(6')-Ic*.

In both antibiotic-treated and naive group, the most abundant class of ARG was multidrug (mean: 27.85%; SD: 21.90%), followed by fluoroquinolone (mean: 17.41%; SD: 20.18%), beta-lactam (mean: 15.27%; SD 16.99%), tetracycline (mean: 14.70%; SD: 10.75%), MLS (mean: 13.06%; SD: 15.29%) and aminoglycoside (mean: 8.77%; SD: 12.48%) (Fig. 1A). In terms of prevalence, aminoglycoside (76/82) and beta-lactam (75/82) were the most common class present in the samples from antibiotic-treated group, while multidrug (92/99) and beta-lactam (91/99) were the most common class in the naive infants. When classified based on resistance mechanism, antibiotic efflux (mean: 59.56%; SD: 20.97%) was the most abundant category in both the groups across all time points (Supplementary Fig. 3). At baseline (T1), the ARG composition of antibiotic-treated group was found to be more heterogeneous (higher inter-individual variability) and diverse compared to naive infants (Fig. 1B).

Homology to ARGs posing major current threat for public health, including those identified by recent omics-based framework [27], were identified in nasopharyngeal samples of our cohort. Most of these potential high-risk ARGs (rank I) were detected at multiple time points across infants in both antibiotic-treated and naive group (Supplementary Fig. 4). We also observed a few high risk and clinically relevant ARGs that appeared directly after the early antibiotic treatment at T2.

For instance, high-risk ARGs specific to the antibiotics encoding for aminoglycoside-modifying

enzymes (*AAC(3)-II*) along with extended-spectrum TEM beta-lactamase (*bla*$_{TEM-1}$) were detected

in two infants. These ARGs co-occurred with other non-targeted high-risk ARGs such as *qnrS*,

floR and *aadA* (rank II). Previous studies have shown the co-carriage of these ARGs on multidrug

resistance plasmids in Enterobacter species [28]. Also, other clinically relevant Extended Spectrum

Beta-Lactamase (ESBL) genes belonging to SHV beta-lactamase (rank II) family were detected in

two other infants following antibiotic exposure. In one patient, the plasmid-mediated AmpC-type

β-lactamase (ACT-beta lactamase), known to confer resistance against all classes of beta-lactam

antibiotics such as penicillins, cephalosporins and carbapenems was detected only after the

treatment cessation (T2). These high-risk ARGs were observed mainly in early antibiotics exposed

infants whose mothers had also received antibiotics during pregnancy (prenatal). Nonetheless, they

did not persist in nasopharyngeal samples collected after discharge at the corrected age of 6

months. Amongst ARGs that are found in Gram-positive pathogens, the *Staphylococcus aureus*

methicillin-resistance gene i.e., *mecA* was detected in 44 samples from almost all (14/15)

antibiotic-treated infants and 10 of the 21 naive infants.

Impact of early antibiotics on preterm infant nasopharyngeal resistome

Firstly, we investigated the total ARG abundance (quantified as the sum of RPKM) and ARG α-

diversity (quantified by Shannon index) across sampling time points. In both the antibiotic-naive

and treated groups, we found that total ARG abundance and diversity increased from T1 to T3-T4,

before descending at T5 until 6 month corrected age (T6). In treated infants, total ARG abundance

and diversity increased more rapidly upon antibiotic treatment, whereas the changes were

generally more gradual over time points in naive infants (Fig. 2A, B). To identify the direct effects

of early antibiotic treatment on the total ARG abundance and diversity, we used a generalized

linear mixed model with individual set as random effect while correcting for age (DOL). Early

antibiotics were significantly associated with increase in both total ARG abundance (LME model:

p.adj = 0.008) and Shannon diversity (*p.adj* = 0.008) after treatment (T2) as compared to baseline

(T1) samples in preterm infant nasopharyngeal resistome. Moreover, the significant effect of

antibiotics on increase in total ARG abundance was observed until T3 (*p.adj* = 0.001).

Nonetheless, such effects were short-lived, as we did not identify significant differences at any of

the following timepoints compared to baseline (T1) in antibiotic-treated group. In naive group of

infants, no significant differences appeared in total ARG abundance and α-diversity between any timepoints during the first 6 months of life while correcting for age.

β-diversity analysis revealed that the overall resistome composition did significantly differ between the two groups (PERMANOVA: $R^2 = 8.2\%$, *p.adj* = 0.02) at baseline (T1). Infants that received early antibiotic treatment had more dissimilar (dispersed) resistome composition than the naive infants (Fig. 3A). Further, we observed that nasopharyngeal samples were clustering more effectively according to birth weight group ($R^2 = 16.4\%$, *p.adj* = 0.012; Supplementary Fig. 5A). No significant difference in baseline resistome composition between antibiotic-treated and naive infants ($R^2 = 0.03$, *p.adj* = 0.319) was found while correcting for birth weight group. However, a minor yet significant compositional difference in resistome was still observed between the two groups ($R^2 = 5.9\%$, *p.adj* = 0.03) directly after the cessation of early antibiotic treatment (T2) (Fig. 3B). Moreover, the compositional changes due to early antibiotics at T2 were more discernible in infants whose mothers had received prenatal antibiotics during pregnancy (prenatal + early (*n = 8*) vs naive (*n = 20*): $R^2 = 10.09\%$, *p.adj* = 0.006; Supplementary Fig. 5B). Following treatment, resistome normalized rapidly as no significant differences in composition between naive and treated group were detected from T3 until 6 months of corrected age (T6) (Fig. 3C).

Clinical covariates associated with preterm infant nasal resistome composition development
The associations between resistome composition and clinical factors were analyzed using PERMANOVA-tests (all *p.adj* ≤ 0.001) and principal component analysis (PCA) across all nasopharyngeal samples from preterm birth until 6 months corrected age, using centered log-ratio (CLR) transformed relative abundance data (Supplementary Fig. 2). The PCA was explained by two main principal components, describing 18.5% and 17.7% of the variation, respectively. Besides inter-individual variation (PERMANOVA (univariate): $R^2 = 37.7\%$, *p.adj* = 0.001), age was the other significant covariate found to be associated with overall resistome composition in the overarching cohort. A moderate yet statistically significant effect was observed with GA (R^2 4%, *p.adj* = 0.001) and BW group (R^2 4.1%, *p.adj* = 0.001), which are highly correlated variables (as indicated by chi-squared test: $p < 2.2e\text{-}16$). Mode of delivery, gender, and intrapartum antibiotic treatment did not appear to shape the overall nasopharyngeal resistome composition (Supplementary Table 3). Covariates that differed between the antibiotic-treated and naive infants

at baseline and that also had a significant association with the overall resistome composition (GA

and prenatal antibiotics (R^2 4.5%, *p.adj* = 0.001)) were used for correction in the subsequent

multivariate analysis.

Multivariate PERMANOVA analysis showed that the largest effect on overall resistome

composition remained attributed to inter-individual variation (R^2 30.03%), followed by PMA (R^2

9.7%) and chronological age group (sampling time points) (R^2 8.4%, all *p.adj* = 0.001). Although

we observed a high degree of variation between samples with no obvious discrete spatial clustering

in relation to these variables upon visual inspection (Fig. 4A-C). We found no significant influence

of early antibiotics (R^2 = 0.6%, *p.adj* = 0.31) and its duration (R^2 = 1.2%, *p.adj* = 0.009) after

adjusting for confounding variables. However, we observed a significant effect of prenatal

antibiotics (R^2 = 3.6%, *p.adj* = 0.001) and its duration (categorized) (R^2 = 4.4%, *p.adj* = 0.001) in

shaping the overall nasopharyngeal resistome composition in preterm birth. To account for the

likelihood of prenatal antibiotics with early antibiotics exposure in our analyses (Chi-squared test,

p < 0.0003), we further categorized our early antibiotic-treated group based on whether the mothers

had also received antibiotics during pregnancy (prenatal + early, *n = 9*) or not (only early, *n = 6*).

Post-hoc analysis revealed significant differences in the overall resistome composition of infants

having exposure to both prenatal and early life antibiotics (prenatal + early) compared to infants

receiving either only early or naive (no prenatal + no early) (*p.adj* = 0.001; Fig. 4D). No difference

in overall resistome composition was observed between the only early antibiotics and naive group

(*p.adj* = 0.14), underlining the relevance of prenatal antibiotic exposure in shaping the overall

resistome composition of preterm infants during the first six months of life (corrected).

Unsupervised clustering classifies nasopharyngeal samples into distinct clusters based on

composition differences

Next, we determined whether the nasopharyngeal samples were organized into clusters (i.e.,

resistotypes) according to their ARG abundance profiles using an unsupervised method, i.e., DMM

models. This way, we identified three distinct clusters (or resistotypes) that were driven by specific

ARGs. Majority of our samples (~90%) were classified into two main resistotypes, R1 and R2. No

significant difference in the sequencing depth, was detected between the two Resistotypes

(pairwise Wilcox test: *p* = 0.14), suggesting that sequencing depth was not driving their separation.

The first resistotype (R1) was characterized by a predominance of fluoroquinolone resistance

genes (*patA, patB*) and *rlmA(II)* (Supplementary Fig. 6). On the other hand, resistotype R2 was

driven mainly by relatively higher abundance of the beta-lactam resistance gene *SST-1* and the

tetracycline resistance gene *tet(41)*. While resistotype R3 was enriched with the blaZ β-lactamase

resistance gene. The identified resistotypes did vary in their overall ARG composition (β-

diversity), as shown in a PCA plot (Fig. 5A). Further, PERMANOVA analysis showed that

resistotypes were more strongly (larger effect size) and significantly associated with the resistome

than other clinical covariates or early-life factors in preterm infants during the first 6 months of

life (R^2 = 19.05%, *p.adj* = 0.001).

Nasopharyngeal resistome composition of preterm infants is linked to microbial composition

and carries hospitalization signatures

To investigate the impact of microbial composition on resistome composition, we taxonomically

profiled the infant nasopharyngeal samples using MetaPhlAn3 (manuscript under preparation,

Rajar et al. 2023). We noticed that microbial community types, as determined by DMM models,

were highly linked with the resistotypes (chi-square test: *p* < 0.0001). Resistotype R2 was very

specific to the *Serratia* community type. Resistotype R1 was associated with the community types

mainly driven by *Streptococcus* and *Gemella* and Resistotype R3 was enriched in community types

predominated by *Staphylococcus* and *Cutibacterium* (Fig. 5B). These community types were also

significantly associated to the overall composition of the resistome (PERMANOVA: R^2 = 13.01%,

p.adj = 0.001; Fig. 5C). Additionally, result from Procrustes analysis showed a strong and

significant association between the microbial taxa abundance and ARG abundance profiles

(correlation coefficient (r) = 0.85, *p* = 0.001, m^2 = 0.26), underlining that bacterial community

composition structured the resistome composition in the nasopharyngeal microbiome of preterm

infants. However, the correlation was more robust in the antibiotic-naive infants (r = 0.94, *p* =

0.001, m^2 = 0.11) as compared to antibiotic-treated infants (r = 0.85, *p* = 0.001, m^2 = 0.27; Fig.

6A, B).

Next, we performed pairwise Spearman's correlation analysis between ARG and microbial species

abundances to predict the origin or potential microbial host of ARGs in nasopharyngeal samples.

The strongest positive correlation (r ≥ 0.8, *p.adj* < 0.05) was found between *SST-1, AAC(6')-Ic,*

tet(41), *MexI ARGs* and *Serratia marcescens/nematodiphila* (Fig. 6C). It is worth noting that
during our study, a clinically detected outbreak of *S. marcescens* occurred in the NICU a few
months after infant enrollment had commenced. When present at T1, the carriage of these ARGs
persisted in 20 out of 22 preterm infants across both the naïve and treated groups, extending long
after hospital discharge as observed at ~8 to 10 months chronological age (6 months corrected age
(T6)) (Fig. 6D). Occasional time points in which the ARGs associated with *S. marcescens* showed
variations from T1 were observed in few of the samples. Whether these are true variations is
difficult to ascertain as metagenomic studies can only detect DNA sequences that are present in
the specific sample and to a depth that is possible to be captured by the present analytical
approaches.

The highly prevalent ARGs like *patA, patB and rlmA(II)* within Resistotype 1 were observed to
exhibit a strong co-occurrence with *Streptococcus mitis/oralis* and *Gemella*
haemolysans/sanguinis (Fig. 6C). *Staphylococcus aureus* and *Staphylococcus epidermidis*
correlated only with *blaZ* beta-lactam ARG. Interestingly, these associations are in accordance
with information in the literature and AMR gene databases on the known microbial host of these
ARGs, highlighting the potential of such analysis approach to predict the origin of ARG in
metagenomes. However, to reduce the bias due to false ranking of features with many zero values,
this analysis is only applicable on the features (ARGs or taxa) present in high prevalence across
samples.

Lastly, we also examined whether some of these strongly associated bacterial species with ARGs
would also directly correlate with our overall resistome outcomes. We plotted the relative
abundance of each species versus the total ARG abundance (sum of RPKM) and the number of
unique ARGs from all samples. Our analysis revealed that only the relative abundance of
Streptococcus mitis (rmcorr correlation coefficient ($r^{rm} = 0.29$, *p.adj* = 0.01) showed a moderately
linear association with the total ARG abundance while adjusting for inter-individual variability
(Supplementary Fig. 7). However, we did not observe a significant correlation between the relative
abundance of *S. mitis* and the number of unique ARGs ($r^{rm} = 0.19$, *p.adj* = 0.12). To statistically
assess whether *S. mitits* relative abundance explains the dispersion of resistomes across samples,
we conducted PERMANOVA test and found that *S. mitis* significantly explains 11.69% (*p.adj* =

0.001) of the variation in the overall resistome composition. Only *S. marcescens* (R^2 = 11.8%,

p.adj = 0.001) exhibited similar variation in the resistome.

DISCUSSION

The nasopharynx represents a highly accessible microbial community and also serves as a crucial

diagnostic window in combating respiratory infections and AMR [29]. It is also the primary

ecological niche for common respiratory pathogens [30]. To date, one of the closest attempts for

an in-depth investigation of ARGs in the respiratory tract using a culture-free approach identified

ARGs in 86% of the samples. However, this was based on the microbiomes of the anterior nares

and from an adult population (Human Microbiome Project) [31], which are different from the

groups and collection sample sites used in our study. In a recent study focusing on infants, ARGs

were detected in 64% of nasopharynx samples [32]. Samples in this study were enriched for

streptococci, thus precluding general conclusions on the extent of ARG presence in the natural

microbiome. In our study, ARGs were present in 95% of the preterm infant samples from the

nasopharynx, regardless of whether they were exposed to antibiotics or not. Although our study

differs in several aspects from the ones above, the combined findings indicate that the respiratory

tract is likely an important reservoir of ARGs and an integral component of human respiratory

microbiomes. For preterm infants, the detection of ARGs in the nasopharynx during the first few

hours of life might occur due to microbiome colonization from multiple sources such as

transgenerational transfer from mothers and the environment [33, 34]. In addition, preterm infants

usually have longer stay in the hospital after birth than term infants. This most likely contributes

to an increased risk for respiratory tract colonization with drug resistant microorganisms, as

recently reported in relation to the gut resistome [35].

We found that ARGs conferring multidrug resistance via efflux pumps encompassed a large

proportion of the nasopharyngeal resistome in the preterm infants. Similar findings have been

previously reported in the gut and oral resistome of preterm and term infants by other metagenomic

studies [7, 33, 36, 37]. In addition, we detected many of the same aminoglycoside-modifying

enzymes, tetracycline protection proteins, and beta-lactamases ARGs identified previously in the

nasopharynx of South African infants, even though the samples in the South African study were

culture-enriched for streptococci and were from a country with higher AMR challenges than the

origin of samples used in the present study. Such similarities in resistome composition may, at

least in part, be explained by the large proportion of streptococci found in the natural microbial communities of the infant nasopharynx. Together, these findings suggest the existence of a core infant nasopharyngeal resistome.

Staphylococcus aureus is one of the leading pathogens causing neonatal sepsis, and one of the priority drug resistant pathogens by the WHO [38]. We found a strong correlation between the abundance of *Staphylococcus aureus/epidermidis* and the PC1-beta-lactamase (*blaZ*) gene. This is a gene found in the chromosome and mobile plasmids of staphylococci, thus representing high risk ARGs. Another strong correlation was found between *Serratia* species and specific resistance genes such as *SST-1*, which confers resistance against cephalosporins. Other resistance genes correlated with this group included aminoglycoside (*AAC(6')-Ic)* and tetracycline (*tet(41)*) resistance genes. Of note, many clinical isolates of *Serratia spp.*, especially *S. marcescens*, have been reported to harbor these ARGs on their chromosomes or plasmids [39-41]. The beta-lactamase encoding gene *SST-1*, for instance, is found in 33% of the sequenced genomes of *S. marcescens*, while the aminoglycoside acetyltransferase *AAC(6')-Ic* and *tet(41*) are found in 96% and 67% of *S. marcescens* available genomes, respectively [16]. This holds particular relevance, given that *S. marcescens* is linked to nosocomial infections and hospital outbreaks, particularly in NICUs [42]. Of note, multidrug resistant *Serratia spp.* are listed among the most critical bacteria by the WHO for which innovative new treatments are urgently needed [43].

Our results showed a significant effect of early broad-spectrum antibiotic regimen (ampicillin + gentamicin) on the total abundance, diversity, and composition of ARGs in the nasopharynx of preterm infants. These effects on the resistome were short-lived and only observed directly after the cessation of antibiotic treatment (T2). In a recent study investigating the impact of antibiotics used in the first week of life on intestinal resistome, Reyman et al. reported that three other antibiotic combinations (penicillin + gentamicin, co-amoxiclav + gentamicin, and amoxicillin + cefotaxime) also had a transient impact on the infant resistome [44]. As the effects of antibiotics on microbiome and resistome composition are unique to each individual and dependent on the antibiotics used and the body site, caution should be taken when extending the interpretation of such general findings. Interestingly, the transient effects of ampicillin + gentamicin on the nasopharyngeal resistome were more discernible in preterm infants whose mothers were exposed to antibiotics during pregnancy before delivery (prenatal antibiotics).

Moreover, we also observed a more profound effect of prenatal antibiotic exposure on the nasopharyngeal resistome of preterm infants at T2 (after cessation of early antibiotics), which persisted until the last sampling time point before discharge from the NICU (T4-T5), but not at 6 months corrected age. Although very few studies have focused on the association between prenatal antibiotic exposure and the resistome, a recent study based on the gut indicates a possible correlation of prenatal antibiotics with increased abundance of ARGs, at least when used 5 to 6 weeks before birth [7]. Additionally, in concordance with other studies on the infant gut [33, 45], our data showed no significant impact of intrapartum antibiotic therapy on the overall nasopharyngeal resistome composition. The nasopharyngeal samples of preterm infants at 6 months corrected age were still clustered according to the birth weight group (similar to T1, Supplementary Fig. 5C), suggesting a strong but transient influence of other environmental factors, such as antibiotic exposures. Nonetheless, there is a paucity of relevant studies on nasopharynx that can corroborate our findings, as they are mainly restricted to the infant gut resistome [33-35, 44-46].

One of the strengths of this study is that it presents the most comprehensive information to date on the characteristics and dynamics of resistome development in the nasopharynx, including results for the first six months of life (corrected) – a critical period of airway development in infants. In addition, the naive group was enrolled at the same time as our treated group from the same hospital, thus removing potential confounding differences in enrollment years or sampling duration. On the other hand, the small number of samples at each time point and inter-individual response to antibiotic exposure may be insufficient to unravel significant associations between antibiotic regimen and resistome development. Moreover, the baseline samples for the antibiotic-treated group ideally would represent samples taken before treatment. However, antibiotic treatment initiation for sEONS and clinical stabilization of the preterm infants were a high priority that needed to precede nasopharynx sample collection. To partially address this issue, we included only preterm infants with samples collected no more than 24 hours after antibiotic initiation. Even so, this choice may have resulted in the selection of more clinically stable infants, which could possibly limit the generalizability of our results. Despite this approach, the differences in resistome changes between baseline and after treatment found in our study may underestimate the true impact of antibiotics on the resistome. We rigorously adjusted for such confounding factors and inter-individual variation using multivariable methods or matched analysis to ensure the reliability of

our observations as a consequence of early antibiotics. Nevertheless, we cannot isolate early
antibiotic effects from other adverse pre-life events that coincide with premature birth, such as
prenatal and antepartum maternally administered antibiotics. Furthermore, for low biomass
samples such as those used in our study, sensitive approaches such as deep sequencing are
necessary to obtain meaningful microbiome information. This increases the risk of detecting
potential DNA contaminants. Although it is not possible to eliminate the potential identification
of contaminants, we adopted various control procedures, starting from sample collection to
analysis, to ensure that the data presented in our study is a reliable representation of the ARGs
found in the nasopharynx.

In conclusion, our findings suggest that early-life treatment with broad-spectrum antibiotics for
sEONS can cause perturbations in the nasopharyngeal resistome of preterm infants. However, the
overall perturbations were transient, as we did not observe a significant alteration in diversity,
abundance, composition, or carriage persistence of high-risk ARGs in treated preterm infants
compared to controls at 6 months of corrected age. These are nevertheless potentially relevant,
given the vulnerability of preterm infants in their first days of life. The individual variability in the
distribution of high-risk ARGs and the changes in the resistome following antibiotic treatment
highlight the potential benefits of developing resistome profiling tools for point-of-care use, which
could aid in informing and tailoring antibiotic treatment decisions [47]. In addition, prolonged
carriage of ARGs associated with nosocomial *S. marcescens* was noted. Future multicenter studies
with long-term follow-up are needed to assess the ecological side-effects of hospitalization and
different antibiotic regimens, including alternatives to broad-spectrum antimicrobial therapy on
infant respiratory resistome development. Understanding the development of the respiratory
resistome and the impact of antibiotic therapies is relevant for striking a balance between the
conflicting goals of minimizing the risk of serious infectious complications to preterm birth on one
side, and minimizing the selection of organisms with ARGs on the other. In general, the
disproportionate contribution of respiratory pathogens to drug-resistant deaths at all ages
highlights the importance of future studies focusing on this yet rather unexplored reservoir of
pathogens and ARGs.

REFERENCES

1. Murray, C., et al., *& Naghavi, M.(2022). Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis.* The Lancet. **399**(10325): p. 629-655.
2. Von Wintersdorff, C.J., et al., *Dissemination of antimicrobial resistance in microbial ecosystems through horizontal gene transfer.* Frontiers in microbiology, 2016. **7**: p. 173.
3. Kilian, M., et al., *Parallel evolution of Streptococcus pneumoniae and Streptococcus mitis to pathogenic and mutualistic lifestyles.* MBio, 2014. **5**(4): p. e01490-14.
4. Pailhoriès, H., et al., *Antibiotic resistance in chronic respiratory diseases: from susceptibility testing to the resistome.* European Respiratory Review, 2022. **31**(164).
5. Teo, S.M., et al., *The infant nasopharyngeal microbiome impacts severity of lower respiratory infection and risk of asthma development.* Cell host & microbe, 2015. **17**(5): p. 704-715.
6. Toivonen, L., et al., *Antibiotic treatments during infancy, changes in nasal microbiota, and asthma development: population-based cohort study.* Clinical Infectious Diseases, 2021. **72**(9): p. 1546-1554.
7. Li, X., et al., *The infant gut resistome associates with E. coli, environmental exposures, gut microbiome maturity, and asthma-associated bacterial composition.* Cell Host & Microbe, 2021. **29**(6): p. 975-987. e4.
8. Rajar, P., et al., *Microbial DNA extraction of high-host content and low biomass samples: Optimized protocol for nasopharynx metagenomic studies.* Frontiers in Microbiology, 2022. **13**.
9. Eisenhofer, R., et al., *Contamination in low microbial biomass microbiome studies: issues and recommendations.* Trends in microbiology, 2019. **27**(2): p. 105-117.
10. Krueger, F., *Trim Galore: a wrapper tool around Cutadapt and FastQC.* Trim Galore, 2012.
11. Langmead, B. and S.L. Salzberg, *Fast gapped-read alignment with Bowtie 2.* Nature methods, 2012. **9**(4): p. 357-359.
12. Li, H., et al., *The sequence alignment/map format and SAMtools.* bioinformatics, 2009. **25**(16): p. 2078-2079.
13. Quinlan, A.R. and I.M. Hall, *BEDTools: a flexible suite of utilities for comparing genomic features.* Bioinformatics, 2010. **26**(6): p. 841-842.
14. Andrews, S., *FastQC: a quality control tool for high throughput sequence data*. 2010, Babraham Bioinformatics, Babraham Institute, Cambridge, United Kingdom.
15. Ewels, P., et al., *MultiQC: summarize analysis results for multiple tools and samples in a single report.* Bioinformatics, 2016. **32**(19): p. 3047-3048.
16. Alcock, B.P., et al., *CARD 2020: antibiotic resistome surveillance with the comprehensive antibiotic resistance database.* Nucleic acids research, 2020. **48**(D1): p. D517-D525.
17. Doster, E., et al., *MEGARes 2.0: a database for classification of antimicrobial drug, biocide and metal resistance determinants in metagenomic sequence data.* Nucleic acids research, 2020. **48**(D1): p. D561-D569.
18. Allaire, J., *RStudio: integrated development environment for R.* Boston, MA, 2012. **770**(394): p. 165-171.
19. R Core Team, R. and R.C. Team, *R: a language and environment for statistical computing. R Foundation for Statistical Computing; 2020*. 2021.

620   20.   McMurdie, P.J. and S. Holmes, *phyloseq: an R package for reproducible interactive*
621   *analysis and graphics of microbiome census data.* PloS one, 2013. **8**(4): p. e61217.
622   21.   Wickham, H., *ggplot2.* Wiley interdisciplinary reviews: computational statistics, 2011.
623   **3**(2): p. 180-185.
624   22.   Oksanen, J., et al., *Vegan: Community Ecology Package (R Package Version 2.6-2). 2022*.
625   23.   Holmes, I., K. Harris, and C. Quince, *Dirichlet multinomial mixtures: generative models for*
626   *microbial metagenomics.* PloS one, 2012. **7**(2): p. e30126.
627   24.   Lahti, L. and S. Shetty, *Introduction to the microbiome R package.* 2018.
628   25.   Bakdash, J.Z. and L.R. Marusich, *Repeated measures correlation.* Frontiers in psychology,
629   2017. **8**: p. 456.
630   26.   Kolde, R. and M.R. Kolde, *Package 'pheatmap'.* R package, 2018. **1**(10).
631   27.   Zhang, A.-N., et al., *An omics-based framework for assessing the health risk of*
632   *antimicrobial resistance genes.* Nature communications, 2021. **12**(1): p. 4765.
633   28.   Long, Y., et al., *High Carriage Rate of the Multiple Resistant Plasmids Harboring Quinolone*
634   *Resistance Genes in Enterobacter spp. Isolated from Healthy Individuals.* Antibiotics, 2022.
635   **11**(1): p. 15.
636   29.   Flynn, M. and J. Dooley, *The microbiome of the nasopharynx.* Journal of Medical
637   Microbiology, 2021. **70**(6).
638   30.   Man, W.H., W.A. de Steenhuijsen Piters, and D. Bogaert, *The microbiota of the respiratory*
639   *tract: gatekeeper to respiratory health.* Nature Reviews Microbiology, 2017. **15**(5): p. 259-
640   270.
641   31.   Consortium, H.M.P., *Structure, function and diversity of the healthy human microbiome.*
642   Nature, 2012. **486**(7402): p. 207-214.
643   32.   Manenzhe, R.I., et al., *Longitudinal changes in the nasopharyngeal resistome of South*
644   *African infants using shotgun metagenomic sequencing.* PloS one, 2020. **15**(4): p.
645   e0231887.
646   33.   Parnanen, K., et al., *Maternal gut and breast milk microbiota affect infant gut antibiotic*
647   *resistome and mobile genetic elements.* Nat Commun, 2018. **9**(1): p. 3891.
648   34.   Yassour, M., et al., *Natural history of the infant gut microbiome and impact of antibiotic*
649   *treatment on bacterial strain diversity and stability.* Science translational medicine, 2016.
650   **8**(343): p. 343ra81-343ra81.
651   35.   Gasparrini, A.J., et al., *Persistent metagenomic signatures of early-life hospitalization and*
652   *antibiotic treatment in the infant gut microbiota and resistome.* Nature microbiology,
653   2019. **4**(12): p. 2285-2297.
654   36.   Rahman, S.F., et al., *Machine learning leveraging genomes from metagenomes identifies*
655   *influential antibiotic resistance genes in the infant gut microbiome.* MSystems, 2018. **3**(1):
656   p. e00123-17.
657   37.   Sukumar, S., et al., *The development of the oral resistome during the first decade of life.*
658   2022.
659   38.   Tacconelli, E., *Global Priority List of Antibiotic-Resistant Bacteria to Guide Research,*
660   *Discovery, and Development.* 2017.
661   39.   Thompson, S.A., et al., *Novel tetracycline resistance determinant isolated from an*
662   *environmental strain of Serratia marcescens.* Applied and environmental microbiology,
663   2007. **73**(7): p. 2199-2206.

40.   Matsumura, N., S. Minami, and S. Mitsuhashi, *Sequences of homologous β-lactamases*
from clinical isolates of Serratia marcescens with different substrate specificities.
Antimicrobial agents and chemotherapy, 1998. **42**(1): p. 176-179.

41.   Shaw, K., et al., *Characterization of the chromosomal aac (6')-Ic gene from Serratia*
marcescens. Antimicrobial agents and chemotherapy, 1992. **36**(7): p. 1447-1455.

42.   Cristina, M.L., M. Sartini, and A.M. Spagnolo, *Serratia marcescens infections in neonatal*
intensive care units (NICUs). International journal of environmental research and public
health, 2019. **16**(4): p. 610.

43.   WHO, I., *Global priority list of antibiotic-resistant bacteria to guide research, discovery,*
and development of new antibiotics. Who, 2017. **2017**: p. 1-7.

44.   Reyman, M., et al., *Effects of early-life antibiotics on the developing infant gut microbiome*
and resistome: a randomized trial. Nat Commun, 2022. **13**(1): p. 893.

45.   Lebeaux, R.M., et al., *The infant gut resistome is associated with E. coli and early-life*
exposures. BMC Microbiol, 2021. **21**(1): p. 201.

46.   Gibson, M.K., et al., *Developmental dynamics of the preterm infant gut microbiota and*
antibiotic resistome. Nature Microbiology, 2016. **1**: p. 16024.

47.   Leggett, R.M., et al., *Rapid MinION profiling of preterm microbiota and antimicrobial-*
resistant pathogens. Nature Microbiology, 2020. **5**(3): p. 430-442.

DATA AVAILABILITY

The raw sequencing data (after removal of human DNA) for all samples are publicly available at

NCBI SRA under BioProject ID: PRJNA1009231.

ACKNOWLEDGEMENTS

We would like to thank the parents of participating infants, and the clinical staff at the Neonatal

Intensive Care units at Ullevål hospital for their assistance with sample collection. The sequencing

service for this work was provided by the Norwegian Sequencing Centre

(www.sequencing.uio.no) and the computations were performed on resources provided by Sigma2

- the National Infrastructure for High Performance Computing and Data Storage in Norway.

FUNDING

This work was financed by the Norwegian Research Council (NFR) (project number: 273833), the

Olav Thon Foundation (project number: 421258), and the Faculty of Dentistry at the University of

Oslo.

(A)

Fig. 1: Resistome composition in the nasopharynx microbiome of preterm infants. Stacked bar plot showing the relative abundance of (A) antibiotic resistance genes (ARGs) clustered at the

Class level for the naive and antibiotic-treated groups over time; (B) top 30 most abundant ARGs identified across samples at baseline (T1) in early antibiotics treated (left) and naive (right) groups. All other ARGs are grouped into the "Other" category. MLS = macrolide-lincosamide-streptogramin.

Fig. 2: Impact of early antibiotic treatment on nasopharynx total ARG abundance and ARG α-diversity Boxplots showing the **(A)** total ARG abundance (sum of RPKM) and **(B)** ARG Shannon diversity per sample, stratified by antibiotic treatment and time points. The lower and

upper horizontal box lines correspond to the first and third quartiles, with the line in the middle of the boxes representing the median. Whiskers extending from the boxes represent the range of data points within 1.5 times the interquartile range below the first quartile and within 1.5 times the interquartile range above the third quartile. (Statistical significance: LME mixed effect model. Adjusted p-values (p): $*p < .05$, $**p < .01$ and $***p < .001$). RPKM = reads per kilobase of reference gene per million bacterial reads.

Fig. 3: β-diversity PCA plots of overall nasopharyngeal resistome composition stratified per time point. Principal component analysis (PCA) based on Euclidean distances of CLR-transformed ARG abundances (*i.e.,* Aitchison dissimilarity) between samples, visualizing the overall nasopharyngeal resistome composition stratified for antibiotic-treated infants and naive at time point (A) T1 (baseline) and (B) T2 (immediately after cessation of early antibiotic therapy). Each data point represents the resistome composition of one sample. The ellipses represent the 95% confidence interval around the centroid of resistome composition for each group. Statistical significance (adjusted p-values) and effect size (R^2) of the differences in beta diversity were assessed using PERMANOVA. In panel (C), the results of the PERMANOVA-tests for the later time points are presented. Data points of early antibiotic-treated infants are colored red and those of naive infants are blue. PERMANOVA = permutational multivariate analysis of variance; p.adj = adjusted p-values.

Fig. 4: Factors influencing the overall nasopharyngeal resistome composition in the preterm infants. PCA visualizations of Aitchison dissimilarities between all the nasopharyngeal samples from birth until 6 months corrected age at ARG level. Each data point corresponds to a sample, colored by (A) Infant, (B) PMA, (C) Time points and (D) Prenatal and post-natal antibiotic exposure and shaped according to early antibiotic exposure or naive groups. Effect sizes (R^2) and corresponding adjusted p-values (p.adj, corrected using the Benjamini-Hochberg method) are calculated using PERMANOVA-tests (as shown in the bottom right of each plot). Percentage refers to the percentage of variance explained by the principal component. PMA = postmenstrual age; PC1 = principal component 1; PC2 = principal component 2; PERMANOVA = permutational multivariate analysis of variance; p.adj = adjusted p-values.

Fig. 5: Association between resistotypes, community types, and ARG abundance profiles. The PCA plot showing the association between overall ARG abundance profile with respect to their (A) resistotypes and (C) community types identified among all the nasopharyngeal samples using the DMM models. R^2 was obtained using the PERMANOVA test. (B) Samples proportions for each resistotype shown as a function of community type. DMM = dirichlet multinomial mixture; Strep = *Streptococcus* ; Staph = *Staphylococcus*.

Fig. 6: The resistome variation in nasopharynx is explained by microbiome variation. The ARG and microbial abundance profiles were correlated with each other for antibiotic treated (A) and naive (B) groups using Procrustes analysis. The length of line connecting two points denotes the degree of dissimilarity or distance between the microbiome (filled square) and resistome (filled circle) composition of the same sample. The line and points are colored based on time points in both groups. The correlation coefficients and significance were calculated using the *protest* function in vegan package. (C) Heatmap depicts the Spearman's pairwise correlation coefficients (*r*) between individual ARGs and microbial species from nasopharyngeal samples. Hierarchical clustering using Euclidean distance was performed on both rows and columns. Correlation coefficients (r) was shown only where adjusted p-value < 0.05. (D) Timeline of ARGs associated with *S. marcescens* in the nasopharynx samples of each preterm infant across time points. Each ARG is represented by a different color and shape.

Supplementary Fig. 1: Sample collection and antibiotic exposure for preterm infants in this study. Sampling timeline of 198 nasopharyngeal samples collected from 36 preterm infants over first six months (corrected age), together with information regarding the time of early antibiotic exposure (red and blue colored circles), exposure to subsequent antibiotic treatment (green colored circles) and included/excluded samples. The x-axis represents the days of life (i.e., age) and the y-axis represents the infants. NP = nasopharyngeal; AB = antibiotics.

Supplementary Fig. 2: Statistical analysis design used in the study. ARGs = antibiotic resistance genes; PERMANOVA = permutational multivariate analysis of variance.

Supplementary Fig. 3: Rarefaction curve for the number of unique ARGs identified at subsampled sequencing depth. The red line indicates the median sequencing depth for all samples, while the gray shading represents one standard deviation.

Supplementary Fig. 4: Mean normalized relative abundances of different ARG mechanisms found in the nasopharynx of early antibiotic treated and naive preterm infants at different time points (T1-T6).

Supplementary Fig. 5: Heatmap showing the log10-transformed RPKM abundance of potential high-risk antibiotic resistance genes (ARGs) identified in the nasopharynx of all preterm infants across samples. The x-axis represents the names of samples, which are arranged by treatment group and time points. RPKM = reads per kilobase of reference gene per million bacterial reads.

Supplementary Fig. 6: PCA visualizations of beta diversity analysis using the Aitchison distances at time points: (A) T1, (B) T2 and (C) T6 (6 month corrected age). Each point corresponds to a sample colored based on birth weight group (A, C) and prenatal and postnatal antibiotic exposure (B). Ellipses indicate 95% confidence intervals (CI). Adjusted p-values and effect size (R^2) are based on PERMANOVA test. CLR = centered-log ratio; BW= Birth Weight.

Supplementary Fig. 7: Box plots of the relative abundances of the main ARG contributors are shown for each Resistotype cluster identified across all the preterm infants' nasopharyngeal resistomes using DMM models. The X-axis displays the three identified clusters, and the Y-axis represents the relative abundance of the ARGs. DMM = dirichlet multinomial mixture.

Supplementary Fig. 8: Scatterplots showing the association between correlated species and overall resistome outcomes across all infant samples from birth to 6 months corrected age. The total ARG abundance in RPKM was positively correlated with the relative abundance of *Streptococcus mitis* (A) and non-linearly negatively associated with the relative abundance of *Serratia marcescens* (C). Plots (B) and (D) depict the relationship between the number of unique ARGs and the relative abundance of *Streptococcus mitis* and *Serratia marcescens*, respectively. Plot (A) and (C) are colored by the relative abundance of *Streptococcus mitis*, while plots (B) and (D) are colored by the relative abundance of *Serratia marcescens*. RPKM = reads per kilobase of reference gene per million bacterial reads.

NP cohort	Naive	Early-antibiotics	p
n	21	15	
Male, n (%)	12 (57%)	11 (73%)	0.484
GA, weeks (mean, SD)	31, 6/7	29 5/7, 1	**<0.001**
BW, grams (mean, SD)	1562, 188	1306, 277	**<0.05 (0.0022)**
BW, group, n (%)			**<0.05 (0.013)**
ELBW	0 (0%)	2 (13%)	
VLBW	8 (38%)	10 (67%)	
LBW	13 (62%)	3 (20%)	
Mode of delivery, n (%)			0.694
Vaginal	4 (19%)	4 (27%)	
C-section	17 (81%)	11 (73%)	
Antibiotics during pregnancy, n (%)			**<0.001**
None	21 (100%)	6 (40%)	
< 10 days	0	7 (47%)	
>= 10 days	0	2 (13%)	
Intrapartum antibiotics (IAP), n (%)			0.50
None documented	2 (10%)	0 (0%)	
Given	19 (90%)	15 (100%)	
ROM, hours (means, SD)	3, 12	432, 733	**<0.05 (0.0108)**
Apgar10 (mean, SD)	9, 1	8, 1	0.3039
Nutrition (discharge to 6 months CA)			**<0.05 (0.039)**
fully breastfed	7 (33%)	4 (29%)	
at least 50% mother milk, but also some formula	3 (14%)	6 (42%)	
mostly formula and some mothers milk	6 (29%)	0 (0%)	
formula	2 (10%)	4 (29%)	
Lost to follow up	3 (14%)	1 (7%)	0.626

Table 1: Baseline characteristics of preterm infants analyzed in this study

Time points	T1	T2	T3	T4	T5	T6	Total
Days of life (DOL)	0-3	4-9	12-19	20-31	32-76	229-288	
Extracted samples (before library prep)	35	37	36	35	23	32	**198**
Exclude samples (Low DNA yield)	2	1	1	0	0	0	**4**
Exclude samples (Low DNA yield) (%)	5.7%	2.7%	2.8%	0%	0%	0%	**2%**
Sequenced samples	33	36	35	35	22	32	**194**
Samples failed during Resistome profiling	5	1	2	0	1	1	**10**
Samples with double sampling time points	0	1	0	1	1	0	**3**
Included in the final analysis	28	34	33	34	21	31	**181**

Supplementary Table 1. Sample flowchart. Flowchart showing number of nasopharyngeal aspirate samples from preterm infants (*n=36*) available at several steps in the study including at sampling, DNA extraction, sequencing and bioinformatics analysis.

Supplementary Table 2: Sequence reads statistics

Sample	Raw reads	Reads after Trimming	Human reads	Remaining (Clean) reads	Reads aligning to CARD database	% of clean reads aligned to CARD
S_103	26080501	26067155	16513464	9553691	17435	0.1825
S_104	22764460	22752830	20748076	2004754	10262	0.5119
S_105	20970229	20964496	3419609	17544887	506	0.0029
S_106	28332496	28321717	15003675	13318042	6043	0.0454
S_119	27475272	27457851	25839804	1618047	105	0.0065
S_120	22266609	22253052	20695679	1557373	82	0.0053
S_121	26201196	26188557	22868983	3319574	11179	0.3368
S_122	33790528	33772729	28966515	4806214	6608	0.1375
S_123	28686631	28676796	1679787	26997009	22095	0.0818
S_124	36678862	36580731	32369824	4210907	330	0.0078
S_125	37839541	37794781	27178195	10616586	302677	2.851
S_127	27688873	27682506	593223	27089283	181147	0.6687
S_128	31005030	30983206	27167769	3815437	15888	0.4164
S_129	28691309	28680175	17478848	11201327	17478	0.156
S_130	33163819	33076882	19478139	13598743	401	0.0029
S_137	31372046	31329501	14342527	16986974	13164	0.0775
S_138	21667917	21662586	846609	20815977	504320	2.4228
S_140	24503529	24497786	2632209	21865577	26203	0.1198
S_141	32823359	32806877	26585210	6221667	27575	0.4432
S_142	26789831	26776365	22169024	4607341	3279	0.0712
S_154	33191448	33139128	27972056	5167072	39874	0.7717
S_156	26884193	26874230	546850	26327380	586046	2.226
S_157	37364064	37340342	32241091	5099251	30575	0.5996
S_158	21680884	21674017	651114	21022903	25474	0.1212
S_159	32047093	32031086	26619128	5411958	4165	0.077

S_160	37292025	37257965	16861312	20396653	12196	0.0598
S_161	45235114	45148815	41399286	3749529	1241	0.0331
S_162	67632600	67539141	62221411	5317730	28064	0.5277
S_163	38901430	38869031	36914900	1954131	22451	1.1489
S_164	44585351	44483261	8951337	35531924	288242	0.8112
S_165	25628648	25540450	10013320	15527130	61843	0.3983
S_166	25531087	25512936	21798951	3713985	3399	0.0915
S_167	23680492	23663333	6734120	16929213	116337	0.6872
S_168	45431318	45401429	35864464	9536965	59284	0.6216
S_169	55896867	55855030	52825510	3029520	9542	0.315
S_170	30624654	30570075	19822065	10748010	26921	0.2505
S_171	27481415	27469324	11777603	15691721	2894	0.0184
S_172	37856150	37836322	25830529	12005793	16131	0.1344
S_173	32620409	32598666	2520144	30078522	144584	0.4807
S_174	29843735	29836353	8305978	21530375	41500	0.1928
S_175	28853583	28843865	8770011	20073854	42924	0.2138
S_176	40254779	40230614	38052826	2177788	4703	0.216
S_177	35925057	35889012	19426456	16462556	18834	0.1144
S_178	20274695	20263845	4157655	16106190	27812	0.1727
S_179	28497251	28482477	685432	27797045	64980	0.2338
S_180	23519660	23515651	1775004	21740647	87325	0.4017
S_181	23879524	23867846	23099772	768074	4218	0.5492
S_182	27132318	27127749	3087081	24040668	45811	0.1906
S_183	31972863	31964880	2266406	29698474	50764	0.1709
S_184	31108627	31101186	1126947	29974239	57650	0.1923
S_185	31622571	31616517	3735586	27880931	60664	0.2176
S_186	46757985	46730502	44193952	2536550	4820	0.19
S_187	30402571	30396836	3688615	26708221	47750	0.1788
S_188	33765529	33756376	1114356	32642020	62286	0.1908
S_189	33083199	33073575	2904430	30169145	57677	0.1912

S_190	30060393	30040023	26669757	3370266	18826	0.5586
S_191	33250280	33222469	16861645	16360824	101214	0.6186
S_193	47659319	47614810	14786449	32828361	56496	0.1721
S_214	52537220	52508963	49589032	2919931	5637	0.1931
S_215	29571723	29543447	8765633	20777814	47796	0.23
S_216	35053190	35033550	6769117	28264433	76971	0.2723
S_217	39517504	39487561	38163679	1323882	625	0.0472
S_218	30675257	30665911	1270181	29395730	56564	0.1924
S_219	27547953	27529357	972788	26556569	111748	0.4208
S_220	33157900	33149181	484150	32665031	66541	0.2037
S_221	21346517	21337082	831108	20505974	31847	0.1553
S_222	26286828	26273899	25340090	933809	919	0.0984
S_223	33944591	33939850	414140	33525710	66865	0.1994
S_224	40663366	40642802	38200609	2442193	2399	0.0982
S_225	31198454	31171888	11201529	19970359	96199	0.4817
S_226	28486798	28471399	21921787	6549612	11262	0.1719
S_227	23681140	23676886	758241	22918645	45401	0.1981
S_228	22840259	22827956	12959279	9868677	18766	0.1902
S_229	32833037	32813316	29804790	3008526	4748	0.1578
S_230	32797647	32777151	27045604	5731547	8726	0.1522
S_234	23368571	23345781	15243726	8102055	2282	0.0282
S_235	29631358	29616146	9188612	20427534	924	0.0045
S_236	24316377	24307110	10727801	13579309	5898	0.0434
S_237	31936082	31906503	3105384	28801119	81934	0.2845
S_238	27183251	27160033	9761467	17398566	61606	0.3541
S_241	21273396	21265742	2856634	18409108	35437	0.1925
S_242	37922507	37895061	34772369	3122692	17337	0.5552
S_243	49645886	49619252	46731003	2888249	14157	0.4902
S_244	66190561	66123529	63015475	3108054	9000	0.2896
S_245	24627154	24616003	12421581	12194422	31173	0.2556

S_247	42327108	42314690	484772	41829918	230967	0.5522
S_248	47197213	47176488	26244459	20932029	30315	0.1448
S_249	51501346	51474520	13469371	38005149	93014	0.2447
S_250	32355912	32328838	13242506	19086332	8192	0.0429
S_251	27317957	27289175	17759907	9529268	914	0.0096
S_252	61163417	61088642	47722474	13366168	36220	0.271
S_253	45402778	45377662	39162898	6214764	9007	0.1449
S_254	32164676	32143130	6335296	25807834	88170	0.3416
S_255	39009857	38993377	25088212	13905165	25987	0.1869
S_256	44877026	44838912	37951047	6887865	9660	0.1402
S_257	34093993	34087358	3603064	30484294	64242	0.2107
S_258	21659477	21650879	1178683	20472196	111693	0.5456
S_259	33243536	33235008	1377165	31857843	84420	0.265
S_26	30270838	30256072	26479542	3776530	14701	0.3893
S_260	35892679	35854245	17290178	18564067	36661	0.1975
S_266	33878209	33871694	778911	33092783	52270	0.1579
S_267	29882106	29867961	3938949	25929012	127336	0.4911
S_268	32997828	32992621	2002198	30990423	54318	0.1753
S_269	48795323	48743359	18996479	29746880	175072	0.5885
S_27	49542796	49483919	35050025	14433894	172882	1.1978
S_270	30056925	30048511	785470	29263041	22344	0.0764
S_271	51738432	51716699	48204756	3511943	5587	0.1591
S_278	29095206	29082097	17430590	11651507	147	0.0013
S_279	29498360	29486657	2167755	27318902	214738	0.786
S_28	37996875	37978461	36894669	1083792	1639	0.1512
S_280	39383489	39340791	14014208	25326583	176136	0.6955
S_281	20907270	20894105	2682760	18211345	106040	0.5823
S_282	31922829	31901011	7462386	24438625	93232	0.3815
S_283	37846534	37832711	1040092	36792619	139836	0.3801
S_284	29863131	29846583	24399378	5447205	67	0.0012

S_285	34228242	34214505	14192796	20021709	73436	0.3668
S_286	41765816	41726214	30517072	11209142	66965	0.5974
S_287	32011138	32000174	2080267	29919907	149226	0.4988
S_288	34869195	34833718	19684842	15148876	58922	0.389
S_289	39559968	39542758	1775686	37767072	105574	0.2795
S_29	38045611	38021180	34100541	3920639	5300	0.1352
S_290	37264718	37245276	26298029	10947247	18539	0.1693
S_291	32072825	32065914	1094390	30971524	56532	0.1825
S_292	63816388	63807235	3838178	59969057	109957	0.1834
S_293	29500665	29493003	5380549	24112454	41039	0.1702
S_294	33573848	33566767	1050194	32516573	68572	0.2109
S_295	29125110	29118244	889487	28228757	46684	0.1654
S_296	34647449	34641389	2501224	32140165	55314	0.1721
S_297	27768840	27758959	1480206	26278753	145877	0.5551
S_298	33622191	33613344	2308277	31305067	72539	0.2317
S_299	22661274	22654821	287602	22367219	66621	0.2979
S_30	28267450	28256137	5690538	22565599	6201	0.0275
S_300	20301912	20296120	296296	19999824	37094	0.1855
S_301	24829638	24825480	540037	24285443	50436	0.2077
S_302	31874316	31867866	833258	31034608	76222	0.2456
S_303	39962863	39916697	18377375	21539322	92682	0.4303
S_304	26868511	26861459	297886	26563573	62060	0.2336
S_305	32243814	32235699	429181	31806518	71735	0.2255
S_306	28410746	28404637	5017057	23387580	48409	0.207
S_307	27609393	27603414	1085057	26518357	56119	0.2116
S_308	32232733	32227923	351812	31876111	63333	0.1987
S_309	26565516	26560390	766174	25794216	57990	0.2248
S_310	36884546	36844010	16364759	20479251	43776	0.2138
S_311	24334581	24314320	1204496	23109824	50555	0.2188
S_312	31069699	31062500	976160	30086340	1325	0.0044

S_313	36354561	36333646	28072594	8261052	178	0.0022
S_314	39811266	39782498	37746037	2036461	205	0.0101
S_315	54249580	54222485	51664787	2557698	7333	0.2867
S_316	44061982	44039723	40399113	3640610	7186	0.1974
S_317	44116478	44095092	42622292	1472800	633	0.043
S_331	29644028	29624401	13828912	15795489	31192	0.1975
S_332	39672217	39644362	38196480	1447882	827	0.0571
S_333	26467120	26441980	13851310	12590670	97356	0.7732
S_334	28679182	28648634	11319308	17329326	105763	0.6103
S_335	29706172	29701470	1533135	28168335	46960	0.1667
S_342	4190563	4185620	997864	3187756	873	0.0274
S_343	26076305	26063824	20907404	5156420	4327	0.0839
S_344	28905903	28893140	19415003	9478137	8856	0.0934
S_345	36832919	36821990	776032	36045958	136753	0.3794
S_346	61044890	61004312	59371452	1632860	2000	0.1225
S_347	69251411	68623791	64379680	4244111	1229	0.029
S_354	27433235	27422512	16225329	11197183	53	0.0005
S_356	27229735	27221424	4719264	22502160	86767	0.3856
S_357	27546962	27535322	18225311	9310011	27877	0.2994
S_364	64520057	64429899	62001636	2428263	979	0.0403
S_365	24055590	24045067	10406711	13638356	24613	0.1805
S_366	55484020	55321634	52415053	2906581	4562	0.157
S_367	28470266	28444380	8614772	19829608	96588	0.4871
S_368	20964782	20957580	7515333	13442247	25918	0.1928
S_369	31720586	31709062	3007642	28701420	42042	0.1465
S_53	34434233	34414250	32019563	2394687	7072	0.2953
S_55	26470453	26463326	7666968	18796358	2006	0.0107
S_86	28423053	28408167	23773990	4634177	13285	0.2867
S_87	66938909	66905387	64680249	2225138	6394	0.2874
S_88	30976709	30958246	4982552	25975694	72811	0.2803

S_89	25774477	25760044	24970857	789187	46	0.0058
S_90	46594816	46555570	45060476	1495094	395	0.0264
S_91	36674226	36641370	35795529	845841	1172	0.1386
S_93	53178273	53147304	51682818	1464486	2822	0.1927
S_95	33533344	33517467	32593464	924003	1464	0.1584
S_96	26810898	26796813	22625616	4171197	492	0.0118

Supplementary Table 3: PERMANOVA results

Univariate analysis					
Factor	**Df**	**SumOfSqs**	**R2**	**F**	**p.adj**
Early antibiotics	1	3574	0.02283	4.1813	0.001
Individual ID	35	59041	0.37705	2.5075	0.001
Time points (age groups)	5	12832	0.08195	3.1241	0.001
Mode of Delivery	1	600	0.00383	0.6884	0.756
Gender	1	1856	0.01185	2.1468	0.02
Birth weight (BW) group	2	6393	0.04083	3.7885	0.001
Days of life (DOL)	1	2466	0.01575	2.8645	0.005
Post-menstrual age (PMA)	10	15136	0.09666	1.8191	0.001
Gestational age (GA) at birth	3	6313	0.04032	2.4787	0.001
Prenatal maternal antibiotics duration (categorized)	2	8162	0.05212	4.8942	0.001
Prenatal maternal antibiotics	1	7008	0.04475	8.386	0.001
Early antibiotics duration	1	4640	0.02963	5.4662	0.001
Ruptured membranes duration	1	1692	0.0108	1.9551	0.046
Apgar score	1	1412	0.00902	1.6285	0.082
Intrapartum maternal antibiotics	1	2311	0.01476	2.6813	0.007

Multivariate analysis					
Factor	**Df**	**SumOfSqs**	**R2**	**F**	**p.adj**
Individual ID	31	47028	0.30033	2.255	0.001
Early antibiotics	1	947	0.00605	1.1577	0.31
Early antibiotics duration	1	2009	0.01283	2.4664	0.009
Time points (age groups)	5	13213	0.08438	3.4531	0.001
Post-menstrual age (PMA)	10	15261	0.09746	1.959	0.001
Prenatal maternal antibiotics	1	5700	0.0364	6.9391	0.001
Prenatal maternal antibiotics duration (categorized)	2	6961	0.04445	4.25	0.001