Kinematic signatures of fast feedback trajectory control during goal-directed finger, arm, and saccadic eye movements

Niranjan Chakrabhavi [a,c,*], Varsha Vasudevan [b], Varadhan SKM [c], Ashitava Ghosal [d], Aditya Murthy [a,*]

[a] Centre for Neuroscience, Indian Institute of Science, Bengaluru 560012, India. [b] Department of Bioengineering, Indian Institute of Science, Bengaluru, 560012, India. [c] Department of Applied Mechanics, Indian Institute of Technology Madras, Chennai 600036, India. [d] Department of Mechanical Engineering, Indian Institute of Science, Bengaluru 560012, India. [*]Corresponding Authors – Niranjan Chakrabhavi (niranjanc@iisc.ac.in) and Aditya Murthy (adi@isc.ac.in).

Abbreviated title: Fast feedback enables rapid trajectory control.

Author Contribution: NC conceived the study; VV and AM designed the saccade experiment; VV collected saccade data. NC and VSKM designed the finger experiment; NC and AM designed the arm experiment; NC collected data for finger and arm experiment. NC performed the data analysis; AM and AG supervised the analysis; NC and AM wrote the manuscript; NC, AM, AG, VV, and VSKM reviewed the manuscript.

First Author: Niranjan Chakrabhavi (ORCID ID: 0000-0003-2583-8689).

Conflict of Interest Statement: The authors declare no conflict of interest.

Keywords: motor control, trajectory planning, variability, and decorrelation.

This pdf file includes:
1. Figures 1 to 10
2. Tables 1 to 3

Ethical Statement: All the experiments discussed in this manuscript have been ethically

approved and all the participants provided their consent for the experiments. (Check methods

section for further details).

Data Availability: Data for this research article is available on request from the corresponding

author. The data is not available publicly due to privacy and ethical restrictions.

Acknowledgements

We would like to thank all the subjects who participated in this study and colleagues in

the labs of VSKM and AM for their assistance during the experiments. This research was

supported by the Department of Biotechnology – Indian Institute of Science Partnership

Program grant (DBT–IISc, *BT/PR27952/INF/22/212/2018*), Intramural funds from IISc

(Institute of Eminence grant from the Ministry of Education), and Science and Engineering

Research Board grant (SERB, *CRG/2022/000553*). NC was supported by the Prime Minister's

Research Fellowship from the Ministry of Education (PMRF, *PM/MHRD-18-16074.03*).

Abstract

Goal-directed eye, hand, and finger movements follow invariant kinematics consisting of approximate straight-line trajectories and bell-shaped velocity profiles. A fundamental unresolved issue is whether these trajectories are planned or whether they are a consequence of trajectory-free online control. We address this question using Spearman's rank correlation, zero-crossing rate, and z-scores, and analyze within-trial variability to investigate differences in the time evolution of trajectories during the presence or absence of a goal in finger and whole-arm reaching movements, along with analyzing rapid goal-directed saccadic eye movements. We found that the central nervous system (CNS) implements control to follow an average trajectory, where goal-directed movements show an enhanced degree of trajectory control. Further, we found behavioral signatures of rapid control that might operate on these planned trajectories as early as ~60 ms in finger movements and ~70 ms in whole-arm reaching movements. Such early signatures of control suggest that the system could exploit internal feedback along with fast feedback processes to implement meaningful corrections during simple voluntary unperturbed movements. The analysis also revealed that the controller gains varied along the movement and peaked distinctly at early (20 %) and late (90 %) phases of finger and arm movements, implying that trajectory control may be accomplished through implicit way-point objectives during the execution of the movement. These later corrections were missing during the late phase of eye movements, pointing to a predominant role of delayed sensory feedback mediating corrections in the context of finger and arm movements during this period.

Introduction

Normative goal-directed movements are characterized by invariant-kinematic patterns that consist of straight-line trajectories with bell-shaped velocity profiles that scale with movement amplitude and speed (Morasso, 1981). Numerous psychophysical and neurophysiological studies highlight the notion that movement attributes are represented/programmed in the central nervous system well before execution (Keele, 1968; Rosenbaum, 1985; Dick *et al.*, 1986; Morris *et al.*, 1994; Moran & Schwartz, 1999; Padoa-Schioppa *et al.*, 2002; Churchland *et al.*, 2006; Summers & Anson, 2009). Such results have motivated the use of internal models wherein a complete specification of a trajectory can be made, in principle, before its implementation (Bizzi *et al.*, 1984; Wolpert *et al.*, 1995; Wolpert & Kawato, 1998; Desmurget & Grafton, 2000; Wolpert & Ghahramani, 2000). However, other influential models of motor control suggest that only certain critical features of the movement such as the target or desired displacement need to be specified in advance (Todorov & Jordan, 2002; Desmurget & Grafton, 2003; Scott, 2004); here, the neural representation specifying movement trajectories may only be generated dynamically in real-time during movement execution as a consequence of solving a cost function based on sensory feedback that is sensitive to the demands of the task. Thus, it remains unclear whether movement attributes such as trajectories are planned before movement execution or whether they are derived from online feedback control.

One feature that, in principle, can distinguish between planned and derived movement trajectories is the timing of their respective control signals that are necessary to mitigate the effects of motor noise, which is a ubiquitous property of muscle and neural activity. In the context of planned movement trajectories, a comparison between desired and estimated current states can be used as an error signal to control movements in the absence of actual sensory information (Gordon & Ghez, 1987; Kawato *et al.*, 2003; Richardson *et al.*, 2011). In contrast,

sensory feedback control relies on relatively delayed sensory feedback to make meaningful

corrections during the movement (Todorov & Jordan, 2002; Wolpert *et al.*, 2011). While such

rapid control has been studied in the context of fast saccadic eye movements (Kawato, 1999;

Gaveau *et al.*, 2003; Richardson *et al.*, 2011) as well as rapid corrections to task-related errors

(Cluff & Scott, 2015; Scott *et al.*, 2015; LoTemplio *et al.*, 2023), distinct signatures of such

rapid control in the context of unperturbed normative movements have not been experimentally

demonstrated to the best of our knowledge. Furthermore, in both saccadic eye movements and

skeletal reaching movements, it is unclear whether these corrections are guided towards a

planned trajectory (Messier & Kalaska, 1999; Richardson *et al.*, 2011) and whether such

trajectory corrections are constantly implemented along the movement or whether the system

prefers hotspots of control during certain phases of movement (Liu & Todorov, 2007; Kuang

& Gail, 2015).

We developed two novel statistical measures to quantify the extent of

control/corrections to trajectories during normative goal-directed movements by investigating

unique patterns among within-trial variability. Firstly, we used Spearman's rank correlation to

track the relative locations of trajectories to one another, along the duration of the movement,

and associated the decrease in correlation (or the lack of) to quantify the extent of online

control. This measure of control signifies online corrections independent of a planned

trajectory. Secondly, we used z-scores and zero-crossing rate to quantify the propensity of

trajectories to approach an average trajectory, i.e., to examine the system's ability to follow an

average planned trajectory (trajectory control). Using these statistical measures we showed

presence or absence of rapid control across different movement modalities such as precise

finger movements (Chakrabhavi & Skm, 2019), whole arm simple reaching movements, and

across different movement contexts i.e., in the presence or absence of a goal/task, between

varied skill of movements using the dominant and non-dominant arms, across different

movement speeds, and during fast saccadic eye movements (Varsha *et al.*, 2021).

Materials and Methods

A total of four experiments have been conducted and analyzed for the current study,

comprising a variety of finger, whole arm, and saccadic eye movements. Different task

conditions have been tested for the presence or absence of control towards a planned trajectory

during goal and non-goal finger movements (Chakrabhavi & Skm, 2019), goal and return

movements by the arm, movements performed by dominant and non-dominant arms and at

varying movement speeds (fast and slow), and saccadic eye movements in various directions

(Varsha *et al.*, 2021).

Experiment 1: Cyclic flexion and extension movements in fingers

The experiment was conducted by Chakrabhavi and SKM (2019), and the data have

been adopted for the current study. The participants were asked to sit on a height-adjustable

chair and were made to rest their forearms and palms on a flat surface in neutral (pronation-

supination) forearm posture (Fig. 1A, left). The forearm and the thumb were constrained to the

experimental setup while the index, middle, ring, and little fingers were suspended in air and

free to move. Electro-magnetic 3D tracking sensors (Micro Sensor 1.8™, Liberty tracking

system, Polhemus Inc., Colchester, VT, USA) were attached to the dorsal aspects of phalanges

on all four fingers, and the data from the distal phalanges have been used in this study. The

experiment was designed on the LabVIEW interface (National Instruments Corp, Austin, TX,

USA) and was acquired at a 100 Hz sampling frequency. Eight healthy right-handed subjects

participated in the study (5 males and 3 females), and the experiment was approved by the

institute ethics committee at IIT Madras, and written consent was obtained from all the

participants before the start of the experiment.

138   During an attempt, subjects were asked to maintain a wrist posture (0^0 neutral,

30^0 flexion, or 30^0 extension). They were instructed to make cyclic flexion-extension

movements at the metacarpophalangeal joint with the one of the four fingers (index, middle,

ring and little) between 75 ± 10 % of its maximum flexion-extension limits, and to ignore any

involuntary movements in the other non-instructed fingers due to the phenomenon of finger

interdependence/enslavement (Häger-Ross & Schieber, 2000; Li *et al.*, 2004). The movements

performed by the instructed fingers were considered goal-directed, and the involuntary

movements in the non-instructed fingers were considered non-goal-directed movements. Each

attempt lasted for 32 seconds, with an initial 2 seconds of rest followed by 30 seconds of cyclic

flexion-extension movements (Fig. 1A, right). Real-time visual feedback of the distal phalange

of the instructed finger was provided on the screen, along with its movement landmarks, and

an audio metronome was played at 1.5 Hz to ensure the amplitude and timing of the instructed

finger movement, respectively. There were 36 attempts (3 wrist postures x 4 instructed fingers

x 3 attempts each) in total. Data from the distal phalanges of each of the four fingers at neutral

wrist posture (0^0) were used in this study. Specifically, each attempt was segmented into several

half-cycles comprising movements from flexion (local peaks of middle finger trace in Fig. 1A,

right) to subsequent extension (local troughs of middle finger trace in Fig. 1A, right) of

instructed finger and the same time stamps were used to segment the movements of non-

instructed fingers. The half cycles were then resampled to 100 equally spaced segments to

analyze control with respect to a normalized percentage time scale as in Figs. 3 and 4, whereas

the first 300 ms were used for the analysis on an actual time scale in Fig. 3C.

Experiment 2,3 – Discrete horizontal reaching movements by dominant and non-dominant arms and fast and slow movements

Participants: Twenty-five healthy right-handed subjects (15 subjects for experiment 2, and 10 subjects for experiment 3; 21 males; Age: 21-32) participated in the study. The experiment was approved by the institute ethics committee at IISc Bengaluru, and all the participants provided written informed consent before the start of the experiment. The Edinburgh Handedness Inventory (Oldfield, 1971) was used to evaluate their dominant arm.

Experimental Setup: Subjects sat on a chair with a backrest to minimize the movements of the clavicle (collarbone). Their dominant or non-dominant arm was required to rest on a smooth elevated table to facilitate horizontal reaching movements (Fig. 1B, left). They were mounted with five electromagnetic 3D motion tracking sensors (Teardrop Mini sensors, 240Hz, Liberty tracking system, Polhemus Inc., Colchester, USA) on the surface of the skin at the sternoclavicular joint (neck), right or left acromion process (right or left shoulder) depending on the task of interest, lateral epicondyle (lateral aspects of the elbow), dorsal tubercle of radius (wrist) and the dorsal aspect of the hand respectively, by a 3M micropore surgical tape. The sensor on the dorsal side of the hand was used for the current analysis. The lights were switched off in the experimental room to occlude the subject's vision of their arm posture during the movement. Subjects were required to hold an endpoint robotic manipulandum (KINARM, 240 Hz, BKIN Technologies Ltd., Kingston, Canada) and perform a center-out discrete horizontal planar reaching movement as required by the task. During the reach movement, subjects were required to look at the stimulus screen (LCD Screen, 60 Hz) in front of them, where a circular-shaped cursor on the screen was mapped onto the position of the held manipulandum. All the experimental protocols were coded and executed on TEMPO/VideoSYNC Software (Reflective Computing, St. Louis, MO, USA). The dataset was post-processed using MATLAB interface (MathWorks, Massachusetts, USA).

Task Structure: A white fixation box (hollow square, 2 cm) appeared at the center of the

workspace where the subjects were required to position the pink-colored cursor inside it by

moving the manipulandum with one of their arms (Fig. 1B, right). Once they fixated at the

center for 500 ms, a red-colored target (filled square, 1 cm) appeared on the screen at an

eccentricity of 12 cm from the center at any of the eight directions (0^0, 45^0, 90^0, 135^0, 180^0,

225^0, 270^0, or 315^0). Subjects made smooth, unperturbed goal-directed movement towards the

target location as soon as the central fixation box disappeared, which was after a hold time of

1000 ms. After initiating the movement, they were given 1000 ms to reach the target location

(within a 2 cm square window). Once they reached the target, they were required to stay there

for 500 ms to complete one trial successfully. The successful completion of the trial was

indicated to the subjects by providing an auditory tone. The inter-trial interval was set to 1500

ms, where the cursor had to be brought back to the central fixation box for the initiation of the

subsequent trial. These were called return movements and were not associated with either the

primary task or reward. Sample movement traces of goal and return movements by the

dominant arm and non-dominant arms are shown in Fig. 1B. If the subject did not fixate

enough at the center or at the target, or if they were not able to execute movements within the

required time, the trial was aborted. The experiment was conducted such that ~40 successful

trials were recorded in each of the eight directions (5 out of 15 subjects performed ~60 trials

towards 0^0 and 45^0, and ~20 trials towards other directions). The same task was now repeated

with the other arm, and the sequence of recruitment of the arms was randomized across

subjects.

Experiment 3 on fast and slow movements was designed similarly with some minor

modifications. The target color was used to cue the subjects about the speed of movement –

green for fast (reach within 400 ms) and red (reach from 600-1000 ms) for slow movements,

and that the trials were randomly interleaved. For this experiment, the subjects used their

dominant right arm and performed movements toward 0^0 and 45^0 directions only.

Data pre-processing: The motion data from the 3D motion tracking sensors on the hand

was low-pass filtered with a cut-off frequency of 20 Hz using a fourth-order zero-lag

Butterworth filter (Scholz *et al.*, 2000). The time at which the hand crossed 5 % of its maximum

speed during the acceleration and deceleration phases, respectively, was noted (Georgopoulos

et al., 1982; Scott *et al.*, 2001; Chakrabhavi & Skm, 2019). The time of onset and the end of

movement were identified by descending along the velocity curve from the 5 % mark to

whenever it reached zero for the first time. The motion data was then resampled to 100 equally

spaced segments to normalize for time (percentage time) across various trials in the analysis of

Figs. 5 – 9, while the data for the first 500 ms during the movement was used for the analysis

of the actual time scale in Figs. 5C and 7D. For Figs. In 5A, 6A, and 6D, the fixation condition

was defined as the time between movement onset (t = 0) to 1000 ms prior to the onset, in the

negative direction of time, when the cursor was inside the fixation box. We used this as a

reference to compare goal-directed movements that were performed subsequently.

Experiment 4 – Saccadic Eye Movements

The experiment was performed by Varsha et al. (2021) at IISc Bengaluru, and the data

have been used for this study as well. A total of 20 healthy subjects participated in this study.

Participants sat in front of a monitor, with their heads fixed (Fig. 1C, left), and their eye

movements were recorded using an eye tracker (240 Hz, ISCAN, Boston, USA). The

experiment was approved by the ethics committee at IISc Bengaluru, and participants provided

informed consent.

On a given trial, the subjects had to fixate at the central fixation box for 400 ± 60 ms

after which the box disappeared, and a green target appeared in the periphery. The subjects

were asked to initiate saccadic eye movements towards the target as soon as possible and

received a green tick and a beep sound on successful trial completion. The targets were shown

at 12^0 eccentricities at various orientations (0^0, 30^0, 45^0, 60^0, 90^0, 120^0, 135^0, 150^0, and 180^0)

and eccentricities of 6^0, 8.5^0, and 10.4^0 along the orientations of 0^0 and 90^0. The saccadic eye

movements of 12^0 eccentricities were considered for the analysis (sample trajectories in Fig.

1C, right). The onset and the end of movement were adopted as it is from Varsha et al., (2021).

------------------------------- *Insert Figure 1 here* -------------------------------

Data analysis

The displacement data (in the direction of flexion-extension) of the four fingers (index,

middle, ring, and little) from Experiment 1 (Chakrabhavi & Skm, 2019), the displacement in

X and Y directions (horizontal movement) of the hand from Experiments 2 and 3, and the

displacement in X and Y directions of saccadic eye movements in Experiment 4, were used for

the analysis.

The number of trials in each of the fingers was selected such that the initial postural

variability was equated between the goal and the corresponding non-goal fingers. For the goal

and return movements by the arm, the initial variability in hand postures was equalized along

both X and Y coordinates. For dominant and non-dominant arm movements, the equalization

of initial variability was performed by considering the similarity of movements with respect to

both arms. For example, the initial variability along 0^0 directed movements by the dominant

arm was equated with the 180^0 directed movements performed by the non-dominant arm. The

initial variability was similar for fast and slow movements as the trials were randomly

interleaved. A standard F-test was used to compare variability between two

conditions/distributions, and the farthest point from the mean of the distribution, with more

significant variability, was eliminated iteratively until there were no significant differences in

variability between the two distributions (Gopal *et al.*, 2017). The equalization of initial

variability was done to eliminate the influence of initial variance on the correlation analysis

and restricted to reporting differences in correlation due to differences in movement execution.

For any given task, and at every normalized time point, each trial was assigned a z-

score using Eq. 1, by subtracting its amplitude from the mean and dividing it by the inter-trial

standard deviation (Fig. 2). Subtraction from the mean was performed to track how the

trajectories across different trials moved relative to the average trajectory, normalized by the

standard deviation, through the course of movement (Figs. 7A and 8A). A sample of movement

trajectories (grey) of the arm, directed towards 0^0, is shown in Fig. 2C, and the z-scores of the

corresponding red and blue trajectories are shown in Fig. 2D. The motivation of this procedure

was to analyze a unitless measure to study variability.

$$z_{i,j} = \frac{A_{i,j} - \mu_j}{\sigma_j}$$　　　　　**(Eq. 1)**

where $z_{i,j}$ is the z-score of trial $-i$ at time $-j$ of the trajectory with amplitude $A_{i,j}$. Parameters

μ_j and σ_j are the inter-trial average and inter-trial standard deviation of amplitude (A) across n

trials at time j. $A_{n \, x \, T}$ is the matrix representing the amplitude of trajectories with n trials and T

= 101, normalized time instances.

To investigate the structure of such variability, we performed Spearman's rank

correlation (Hollander *et al.*, 2013) between the z-scores at the start of movement $z(0)$, with

the z-scores at subsequent time instances $z(j)$. In Fig. 2, we have depicted two distinct cases

of online control. On the left side (Fig. 2A), the representative trajectories crossover each other

such that their relative ranks change during the movement. For example, the rank order of

trajectories - red, green, blue, and pink (top to bottom) at time $t = 0$, changes to green, pink,

red, and blue at time $t = j$. This leads to decorrelation among their ranks and a decrease in

Spearman's correlation coefficient (ρ) because of the implementation of active online control.

Conversely, on the right side (Fig. 2B), the order of trajectories remains the same throughout

the movement, and the trajectories never converge towards each other, thereby resulting in high

ρ (close to 1) and weaker/lack of implementation of online control. We performed the

correlation analysis from movement onset to the end of movement (t = 0 to 100 %), to obtain

a percentage time varying trajectory of correlation coefficient, $\rho(t)$. We chose Spearman's

correlation coefficient as we were particularly interested in the crossovers among trajectories,

independent of changes in inter-trial standard deviation (Figs. 7A and 8A) during the movement

(Janse *et al.*, 2021). This method is independent of whether we compute the z - score of the

trajectories or do it on the actual amplitudes. We investigated the average decorrelation ($\bar{\rho}$)

over the entire movement duration to quantify the extent of online control. We then fit a logistic

function $\rho fit(t)$, over $\rho(t)$ for each of the subject and used parameter β to quantify the rate of

decrease of $\rho(t)$ using Eq. 2.

294
$$\rho fit(t) = a + \frac{1}{1+e^{\beta(\rho(t)-c)}}$$
(**Eq. 2**)

-------------------------------- *Insert Figure 2 here* --------------------------------

To substantiate our hypothesis, we computed a dispersion measure for the time-

normalized trajectories ($A_{n \times T}$) by taking the ratio of average inter-trial standard deviation over

the time course with the average range of movement during the *n* trials in Eq. 3. We divided

variability with the range to account for differences in movement amplitude in the finger

experiment. We then correlated the average decorrelation ($\bar{\rho}$) with its average dispersion

measure (or with standard deviation in Figs. 3B, 5B, 7C, 8C, 9C, and 10B), expecting that an

increased decorrelation leads to lower dispersion.

305
$$Dispersion = \frac{\frac{1}{T}\sum_{j=1}^{T}\sqrt{\frac{\sum_{i=1}^{n}\left(A_{i,j}-\mu_j\right)^2}{n-1}}}{\frac{1}{n}\sum_{i=1}^{n}|\max(A_i)-\min(A_i)|}$$ (**Eq. 3**)

where μ_j is the average of amplitude (A) across n trials at time j and A_i is the trajectory on trial

$-i$.

The timing of control in Figs. 3C, 5C, and 7D – E, were done similarly to finding the

end of the lag phase and beginning of the growth phase in a growth curve using the 'tangent

method' (Bertrand, 2019; Smug *et al.*, 2024). A tangent was constructed at the point of

maximum rate of decorrelation in the Spearman's correlation coefficient curve, $\rho(t)$, and the

initiation of control was identified as the time at which the tangent met the $\rho = 1$, or the

horizontal line.

For Figs. 3D, 5D, and 10C, the correlation analysis was performed using a similar

technique described earlier but between consecutive 10 % - -time instances. At any percentage

time point, the value of Spearman's correlation (ρ_c) was computed between the z-scores of

trajectories at that time point with respect to an earlier 10 % - time instance. For example, at t

= 20 %, correlation analysis was performed between postures at 20 % and 10 % movement

duration. This analysis was restricted to goal-directed movements.

We further investigated whether the implemented online control was intended to follow

an average planned trajectory. In Fig. 2D, the $z = 0$ represents an average trajectory (dashed

line), and the red trajectory crosses the dashed line three times, showing a strong trajectory

control. In contrast, the blue line does not cross the dashed line at all, showing weaker trajectory

control. To quantify this effect, we computed a zero-crossing rate (ZCR) from the z-scores of

movement trajectories, which was defined as per Eq. 4. A larger ZCR implies a greater

trajectory control. For arm movements, the ZCR, in Figs. 6A and 6C were averaged across X

and Y coordinates and along all the movement directions.

The above measure of *ZCR* computed on a one-dimensional dataset was extended to

two-dimensional data (ZCR2) in the context of arm movements (Figs. 6D-E, 7F, 8D, and 9D)

and saccadic eye movements (Fig. 10D), where trajectories were produced in a spatial plane.

This was done by forming a line segment between the trial-averaged 2D spatial locations at

any two timestamps t1 and t2 (the average line, the 2D equivalent of a 1D - $z = 0$ line). For any

given trial, we approximated the trajectory between the same two timestamps as another line

segment. A zero-crossing was marked if the corresponding trajectory line intersected with the

average line. Once the number of zero-crossings had been counted for the entire duration and

all the trials, *ZCR2* was computed using Eq. 4. We also computed *ZCR and ZCR2* during the

early (0 to 33 %), middle (34 to 66 %), and late (67 to 100 %) phases of movements in Figs.

4C, 6C, 6E and 10D.

$$ZCR \ (or \ ZCR2) = \frac{Total \ number \ of \ zero \ crossings}{Total \ number \ of \ trials} \qquad \textbf{(Eq. 4)}$$

To support evidence of trajectory control, the starting postures of goal and non-goal

movements in fingers (Fig. 4B) and arm (Figs. 6B and 7G) were divided into two groups, the

ones which were farther away ($|z| > 0.5$, dark and light purple in Fig. 4B, and red and blue in

Fig. 6B) and the ones closer ($|z| <= 0.5$, black and grey) to the average initial posture ($z = 0$).

The average z-scores (across trials) were tracked along the time course of movement to check

the propensity of these two groups to converge towards an average planned trajectory ($z = 0$,

between 0 – 100 % movement phases). If the central nervous system intended to follow an

average trajectory, the group that was farther away from the initial posture should show greater

convergence towards the average trajectory than the closer ones. The difference in z-scores at

0^{th} (initial) and 20^{th} percentile time points was used to quantify this effect.

 Statistics

The data were checked for normality using the Kolmogorov-Smirnov test. We

performed a pairwise t-test for paired comparisons and a two-sample *t*-test for independent

comparisons if the data were normal. The Wilcoxon signed rank test and Wilcoxon rank-sum

test for paired and independent comparisons were used otherwise. For more than two sample

paired comparisons, we performed repeated measures ANOVA. Multiple pairwise comparisons

were reported with Bonferroni corrections.

Results

Online control of goal and non-goal finger movements

The differences in the magnitude of online control among goal and non-goal fingers

were quantified by computing the average decorrelation ($\bar{\rho}$) during the movement (Fig. 3A). A

two-way repeated measures ANOVA (4 finger-tasks x 2 finger-type, goal or non-goal) showed

that the $\bar{\rho}$ of the goal finger was significantly different from the non-goal finger ($P < 0.001$).

By comparing the goal and the corresponding non-goal fingers in a pairwise method, it was

observed that the extent of decorrelation was highest in the goal fingers (Table 1).

The dynamics of online control in goal and non-goal fingers were teased apart by

computing the rate of decrease (β) in correlation co-efficient, $\rho(t)$ (Fig. 3A). A logistic function

was fit to the dataset to find β (median explained variance = 0.95). A two-way repeated

measures ANOVA (4 finger-tasks x 2 finger-types, goal or non-goal) showed that the rate of

decrease of $\rho(t)$ was significantly different across goal and non-goal fingers ($P < 0.001$).

Pairwise comparisons showed that the rate of decrease of $\rho(t)$ of the goal-directed fingers was

significantly higher than the non-goal, non-neighboring fingers but was not different from that

of neighboring fingers (Table 2).

376   -------------------------------- *Insert Figure 3 here* --------------------------------

378   Further, we checked if the average decorrelation in finger trajectories was

systematically associated with the average trajectory dispersion (Fig. 3B) and found that they

were tightly correlated (Spearman's $\rho = 0.9$, $P < 0.001$), and the average dispersion of goal

finger (median = 0.073) was significantly less than that of non-goal neighboring (median =

0.143, $P < 0.001$) and non-neighboring (median = 0.573, $P < 0.001$) fingers. Thus, a higher

degree of trajectory decorrelation was associated with lesser dispersion in trajectories and vice

versa. As goal fingers were associated with varying movement amplitudes, we compared $\rho(t)$

of all four goal fingers to check the sensitivity of the correlation analysis to the differences in

the range of movement among them. A one-way repeated measures ANOVA on the goal fingers

revealed that the $\bar{\rho}$ were not significantly different ($P = 0.72$), and the rate of decrease of $\rho(t)$

were also not significantly different across goal fingers ($P = 0.52$), but the range of movement

was significantly different between them ($P < 0.001$), with index (mean ± SEM: 10.63 ± 0.27

cm) and the little (mean ± SEM: 7.42 ± 0.27 cm) finger having the highest and lowest

movement ranges, respectively.

393   --------------------------------- *Insert Table 1 here* ---------------------------------

395   Next, we quantified the differences in the initiation of control by performing the

396   correlation analysis on the actual timing using the tangent method. We found that the decrease

397   in $\rho(t)$ of the goal finger (dark purple) was initiated at 60.31 ± 1.93 ms (mean ± SEM). In

398   comparison, decorrelation of the non-goal neighboring (medium purple) and non-neighboring

399   (light purple) fingers were initiated at 115.25 ± 15.13 ms and 163.64 ± 11.67 ms, respectively

400   (Fig. 3C).

401
402   -------------------------------- *Insert Table 2 here* ----------------------------------

403
Motivated by the previous findings, we examined whether control was constant or

occurred during distinct phases along the movement duration. For this, we performed the

correlation analysis on the goal finger movements on consecutive 10 % - time instances and

found that the controller acted discretely with two prominent troughs in $\rho_c(t)$ at 20^{th} (binominal

P < 0.001) and 90^{th} percentile time (binominal *P* < 0.001) points (Fig. 3D), suggesting the

presence of early and late phases of control.

Trajectory control of goal-directed finger movements

We tested the zero-crossing rate (*ZCR*) of goal and non-goal fingers during the four

different instructed finger tasks to probe the propensity of trajectories to follow an average

trajectory (Fig. 4A, and Fig. 2D for the computation method). A two-way repeated measures

ANOVA (4 finger-tasks x 2 finger-type, goal or non-goal) revealed that the *ZCR* of the goal

finger was significantly different from that of non-goal fingers (*P* < 0.001). By performing a

pairwise comparison between goal and non-goal fingers, it was found that the goal finger had

the highest ZCR in the four tasks (Table 3).

419
420   -------------------------------- *Insert Figure 4 here* ----------------------------------

421
To support this result further, we divided the initial postures of goal (solid line) and

non-goal (dashed line) fingers into two groups (Fig. 4B), the ones farther away ($|z| > 0.5$,

purple) and the ones closer ($|z| <= 0.5$, black) to the average initial posture ($z = 0$) and found

that the away group had a significantly greater propensity to converge towards an average

trajectory (mean ± SEM: 0.43 ± 0.02) than the closer ones (0.09 ± 0.02, $P < 0.001$). Also, the

farther away group had a significantly greater convergence in the goal finger as compared to

that in non-goal fingers (0.02 ± 0.01, $P < 0.001$). To substantiate the evidence of a temporal

structure of control in Fig. 3D, we computed the *ZCR* of goal finger movements across three

movement segments (Fig. 4C): early (0 to 33 %, light grey), middle (34 to 66 %, medium grey),

and late (67 to 100 %, dark grey) phases of movement duration and found that the *ZCR* was

significantly higher during early (mean ± SEM: 0.44 ± 0.01) and late (0.49 ± 0.01) phases as

compared to the middle phase (0.27 ± 0.02; Adj. $P < 0.001$ for both the comparisons).

--------------------------------- *Insert Table 3 here* ----------------------------------

Online control of goal-directed whole-arm movements

We investigated whether the correlation analysis performed on finger movements could

be extended to whole arm reaching movements and if there were systematic control signatures

in goal movements. We performed the correlation analysis (Fig. 5A) and found that the $\bar{\rho}$ in

goal movements (mean ± SEM: 0.38 ± 0.02) was significantly less than that during fixation

(0.98 ± 0.003, $P < 0.001$). A logistic function was fit over $\rho(t)$ (median explained variance =

0.94) to find the rate of decrease (β) in correlation, and it was found to be significantly higher

in goal movements (mean ± SEM: 0.143 ± 0.005) relative to fixation (0.063 ± 0.028, $P < 0.005$).

Further, the average decrease in correlation in goal movements was significantly associated

with trajectory dispersion (Pearson's $r = 0.57$, $P = 0.02$, Fig. 5B). We estimated the timing at

which trajectory correction was initiated in goal-directed movements using the tangent method.

We found that it occurred at 69.56 ± 3.45 ms (mean ± SEM, Fig. 5C).

Further, to investigate whether such online control was constant or varied across

different phases of movement, we performed the correlation analysis of goal-directed

movements on consecutive 10% - time instances, $\rho_c(t)$. As was observed in finger movements,

we found two distinct troughs at 20^{th} (binomial P < 0.001) and around 90^{th} (trough shared

between 80^{th} and 90^{th} percentiles, binomial P < 0.001 for both time stamps, shared across the

participants) percentile time instances, indicating early and late phase of control, respectively

(Fig. 5D).

Trajectory control of goal-directed whole-arm movements

As we did with finger movements, we computed the zero-crossing rate (*ZCR*) for goal

movements and compared them during fixation and found that the *ZCR* of goal movements

(mean ± SEM: 1.52 ± 0.03) was significantly greater than that during fixation (0.22 ± 0.02; *P*

< 0.001, Fig. 6A). Similar to the analysis in the previous section, we divided the initial postures

of goal movements into two groups (Fig. 6B), the ones farther away ($|z|$ > 0.5, red) and the

ones closer ($|z|$ <= 0.5, black) to the average initial posture (z = 0) and found that the away

group had a significantly greater propensity to converge towards an average trajectory (mean

± SEM: 0.57 ± 0.02) than the closer ones (0.12 ± 0.02, *P* < 0.001).

To substantiate the evidence for distinct phases of control, the *ZCR* of goal-directed arm

movements was computed across three movement segments (Fig. 6C): early (0 to 33 %, light

red), middle (34 to 66 %, medium red), and late (67 to 100 %, dark red) phases of movement

duration. We found that the *ZCR* was significantly higher during the early (mean ± SEM: 0.59

± 0.01) and late (0.53 ± 0.02) phases as compared to the middle phase (0.40 ± 0.01; Adj. *P* <

0.001 for both comparisons), like that of goal-directed finger movements.

The above zero-crossing analysis was extended to two-dimensional data with

trajectories in the XY plane by computing ZCR2 (see methods). The results were qualitatively

similar to the one-dimensional analysis. We found that the *ZCR2* of goal movements (red, mean

± SEM: 1.23 ± 0.04) was significantly greater than during fixation (grey, Fig. 6D, 0.05 ± 0.01,

P < 0.001). We further divided the *ZCR2* measure of goal movements into three segments (Fig.

6E): early (0 to 33 %, light red), middle (34 to 66 %, medium red), and late (67 to 100 %, dark

red) and found that *ZCR2* during middle phase (mean ± SEM: 0.36 ± 0.02) was significantly

less than the early (0.43 ± 0.02, Adj. *P* = 0.005) and late phases (0.44 ± 0.02, Adj. *P* < 0.001).

484   -------------------------------- *Insert Figure 6 here* --------------------------------

Differences in control between goal and return movements

As the differences in control measures between fixation and reaches may be related to

differences in postural versus movement control, we tested whether these signatures of active

control were different across similar movements with varying contexts. For this, we took

advantage of return movements that were made back to the fixation spot. The maximum inter-

trial standard deviation of return movements (mean ± SEM: 1.46 ± 0.05 cm) was significantly

greater than goal movements (mean ± SEM: 1.17 ± 0.05 cm, *P* < 0.001, Fig. 7A). We found

that $\bar{\rho}$ in goal movements (mean ± SEM: 0.38 ± 0.02) was significantly less than that of return

movements (0.41 ± 0.01, *P* = 0.04, Fig. 7B). This difference in $\bar{\rho}$ was prominent in the first 50

% of the movement duration (goal: 0.5 ± 0.02, return: 0.61 ± 0.01, *P* < 0.001). A logistic

function was fit over $\rho(t)$ (median explained variance = 0.97) to find that the rate of decrease

(β) was significantly higher in goal movements (mean ± SEM: 0.143 ± 0.05) than in return

movements (0.099 ± 0.006, *P* < 0.001). Also, the $\bar{\rho}$ in goal movements was significantly

associated with their respective trajectory dispersion (Pearson's *r* = 0.53, *P* = 0.002).

Interestingly, the $\bar{\rho}$ during the first 20 % - time duration was significantly less in goal

movements (mean ± SEM: 0.81 ± 0.01) than the return movements (0.91 ± 0.01, $P < 0.001$)

and it was significantly associated with the respective maximum standard deviation (Pearson's

$r = 0.464$, $P = 0.009$), which interestingly occurred much later during the movement (Fig. 7C).

This association was conserved for the duration as early as 10 % into the movement execution

(Pearson's $r = 0.426$, $P = 0.018$). These differences between goal and return movements were

not because of movement kinematics, as the maximum movement speeds were not significantly

different across the conditions ($P = 0.62$). Further, the maximum rate of change in $\rho(t)$ for goal

movements (mean ± SEM: 128.33 ± 5.62 ms) was significantly earlier than that during return

movements (201.38 ± 8.02 ms, $P < 0.001$). Using the tangent method, we found that online

control during goal movements was initiated significantly faster than the return movements by

52.58 ± 4.75 ms (Fig. 7D-E, mean ± SEM, $P < 0.001$). We also computed *ZCR2* to find the

propensity of the trajectories to follow an average trajectory. We found that the *ZCR2* of goal

movements (red, mean ± SEM: 1.24 ± 0.04) was significantly greater than the return

movements (1.16 ± 0.04, $P = 0.034$, Fig. 7F). In addition, the farther away group in goal

movements (mean ± SEM: 0.57 ± 0.02, Fig. 7G) had significantly greater propensity to

converge towards the average trajectory than during return movements (0.33 ± 0.02, $P < 0.001$).

--------------------------------- *Insert Figure 7 here* ----------------------------------

Differences in control between dominant and non-dominant arm movements

As expected, the trajectories of the dominant arm (average SD, mean ± SEM: 0.63 ±

0.02) had significantly less inter-trial variability (Fig. 8A) than the trajectories of the non-

dominant arm (0.69 ± 0.02, $P = 0.004$). Although movement kinematics were not significantly

different between dominant (maximum speed, mean ± SEM: 23.38 ± 0.98 cm/s) and non-

dominant arms (24.38 ± 0.72 cm/s, $P = 0.33$), the $\bar{\rho}$ was significantly lesser in the trajectories

of the dominant arm (mean ± SEM: 0.38 ± 0.02) as compared that of the non-dominant arm

(0.41 ± 0.02, $P = 0.029$, Fig. 8B). Differences in the average SD were also significantly

associated with the differences in average decorrelation between dominant and non-dominant

arms (Pearson's $r = 0.648$, $P = 0.009$, Fig. 8C). However, no significant differences in the rate

of decorrelation (β) between dominant (mean ± SEM: 0.145 ± 0.005) and non-dominant arms

(0.151 ± 0.006, $P = 0.19$) was observed. Further, the *ZCR2* (Fig. 8D) was also not significantly

different between the dominant arm (red, mean ± SEM: 1.31 ± 0.05) and non-dominant arm

movements (1.30 ± 0.04, $P = 0.42$). Taken together, we found greater online control in

dominant arm movements, but this did not translate to significant differences in trajectory

control.

536
--------------------------------- *Insert Figure 8 here* ---------------------------------

Invariance of control across different speeds of movement

We subjected the correlation analysis to different movement speeds, between fast and

slow conditions (Fig. 9A) to check how sensitive the correlation analysis was to differences in

signal-dependent noise that was expected to be greater for higher movement speeds. As

instructed, the maximum speed of fast movements (mean ± SEM: 37.03 ± 0.69 cm/s) was

significantly higher than the slow movements (mean ± SEM: 16.58 ± 0.39 cm/s, $P < 0.001$).

Correspondingly, the max SD was higher during fast movements (mean ± SEM: 0.93 ± 0.02

cm) when compared to the slow condition (0.82 ± 0.03 cm, $P = 0.002$). Nevertheless, we found

that $\rho(t)$ for fast and slow movements were largely overlapping and that the $\bar{\rho}$ for fast

movements (mean ± SEM: 0.46 ± 0.04) was not significantly different from that of slow

movements (0.48 ± 0.05, $P = 0.317$, Fig. 9B). Further, the average decorrelation was not

significantly associated with the max SD across two speeds of movement (Fig. 9C, r = 0.249,

P = 0.289), Despite that, the rate of decrease in correlation (β) being greater for slower

movements (mean ± SEM: 0.130 ± 0.008) when compared to fast movements (0.108 ± 0.005,

P = 0.011). We also computed *ZCR2* measure (Fig. 9D) and found no significant differences

across fast (mean ± SEM: 0.92 ± 0.08) and slow (1.06 ± 0.12, P = 0.089) movements.

555
-------------------------------- *Insert Figure 9 here* ----------------------------------

Control of goal directed saccadic eye movements

We subjected discrete goal-directed saccadic eye movements to a similar set of analyses

as that of arm movements. We found that the trajectories decorrelated significantly, indicating

active online control. The average correlation ($\bar{\rho}$) was 0.31 ± 0.01 (mean ± SEM), significantly

less than 1 (Fig. 10A, P < 0,001). The average decorrelation among trajectories was

significantly associated with the average inter-trial standard deviation (Fig. 10B, Pearson's r =

0.544, P = 0.013), suggesting that the online trajectory correction was related to a reduction in

trajectory variability. The decorrelation analysis on subsequent 10% time instances ($\rho_c(t)$)

showed that the saccadic movements did not follow a prominent w-shaped control structure

like that of arm movements but rather had control only during the early phase (10% time) of

movement (Fig. 10C, binomial P < 0.001). Although there was a second significant trough at

40% time (P < 0.001), it was not as prominent as that at 10%. Further, we computed *ZCR2* for

early, middle and late phases of movement (Fig. 10D) and found that the trajectory control was

predominant in the early phase (mean ± SEM: 0.44 ± 0.02) as compared to middle (0.26 ± 0.01,

Adj. P < 0.001) and late phases (0.15 ± 0.01, Adj. P < 0.001), in support of the previous

analysis. It also decreased from middle to late phases of movement (Adj. P < 0.001).

-------------------------------- *Insert Figure 10 here* --------------------------------

Overall, we used the time course of decorrelation between the initial and subsequent

time points as a measure of active online control, the rate of zero crossings around a mean

trajectory and the trend of z-scores as signatures of trajectory control, and compared these

measures in different contexts of finger, arm and saccadic eye movements to show that 1) active

online control (or corrections) in trajectories was initiated as rapidly as ~60 ms in goal-directed

finger movements and ~70 ms in goal-directed whole-arm reach movements, providing

behavioral kinematic evidence for fast feedback control during normative movements. 2) Such

rapid control also maintained the movements along an average planned trajectory. 3) Control

was found to be focused at two distinct phases: at early (20 %) and late (around 90 %) phases

along the time course of finger and arm movement. 4) Online control was sensitive to the level

of skill (across dominant and non-dominant) but was not sensitive to the range (across

instructed fingers) or the speed (across fast and slow) movements. 5) Rapid online control,

potentially mediated by internal feedback, was observed in saccadic eye movements, and

signatures of control were prominent in the early phase and decreased through the middle and

late phases of movement, unlike the w-shaped control structure in the arm movements.

Discussion

Since Woodworth (1899), the detailed analysis of the kinematics of limb trajectories

has provided insights into understanding the relative contributions of motor planning relative

to feedback control (Carlton, 1981; Abrams *et al.*, 1990; Abrams & Pratt, 1993; Chua & Elliott,

1993). In addition to assessing the ability of the motor system to regulate variability throughout

the trajectory (Khan *et al.*, 2006), the coefficient of determination (R^2) has been used before as

a within-trial estimate of the extent to which early kinematic markers (Carlton, 1981; Gordon

& Ghez, 1987; Messier & Kalaska, 1999; Desmurget *et al.*, 2005), such as peak acceleration

and peak velocity, to determine end-point distributions. In this context, online control has been

inferred by measuring the extent to which such markers do not predict the endpoint (i.e., lower

R^2). These results, and the validity of the approach, have been validated by manipulating factors

such as the availability of visual feedback (Khan & Franks, 2003).

The correlation analysis used in this study uses a similar logic with some important

differences that allowed us to reveal novel aspects of control during reaching movements. First,

the earliest kinematic markers typically used, peak acceleration at ~150 ms, are too late to

reveal fast feedback control. Here, in contrast to previous works (Messier & Kalaska, 1999;

Richardson *et al.*, 2011; Kuang & Gail, 2015) that have emphasized the relationship of

kinematic markers to the end of movement, we measured how quickly trajectories diverged

from an initial starting posture. This analysis ensured that the starting variability between

different task conditions was not significantly different and that any difference in control would

be revealed at the earliest possible time. Further, the comparison with non-goal movements

was particularly useful, since it did not involve manipulation of target locations (Wolpert &

Ghahramani, 2000; Gritsenko & Kalaska, 2010) or postures (Pruszynski *et al.*, 2008, 2011) that

would emphasize differences in sensory feedback control as opposed to internally driven

control. Finally, and importantly, the absence of any difference in the extent of decorrelation

for fast versus slow movements suggests that signal-dependent noise (Harris & Wolpert, 1998)

per se was not a factor contributing to the observed decorrelation signal (Fig. 9).

Trajectory control during goal-directed arm movements

In contrast to internal models of trajectory control, optimal feedforward/feedback

control can also explain straight-line trajectories and is supported by recent work showing the

absence of corrective movements to a nominal trajectory in the presence of a perturbation

(Cluff & Scott, 2015). Although this result cannot be easily reconciled with our results, we

suggest that a potential drawback of perturbation studies to assess trajectory control is the

assumption that the original normative plan is still relevant during corrections. In the context

of target shift experiments, at least, this assumption is questionable as the original plan is

expected to be modified or canceled (Mirabella *et al.*, 2008; Murthy *et al.*, 2009). By extension,

we assume a similar logic would also hold for external perturbations of the hand. Nevertheless,

and interestingly, even in the study by Cluff and Scott (2015), when trajectory specifications

such as timing and path were part of the goal, corrections were directed to a normative

trajectory. In this latter context, our experimental findings broadly agree with them (Cluff &

Scott, 2015) as our goal-directed finger and arm movements had timing constraints as

compared to the non-goal movements.

Further, the demonstration of relatively fast task-related sensory feedback control (~50

ms) (Wolpert *et al.*, 1998; Pruszynski *et al.*, 2008), which could, in principle, shape trajectories

during voluntary movement execution involving active participation of the motor cortex

(Pruszynski *et al.*, 2011), is also a key prediction of optimal feedback control models. However,

since movement in optimal feedback models (in the absence of trajectory costs) is driven by

costs associated with the current posture and the target position, they don't necessarily require

movements to crossover an average trajectory. This is in contrast with our observations of zero

crossings, especially during the early phase, and that the trajectories starting farther away had

a rapid propensity to converge towards the average $z = 0$ line (Figs. 4 and 6). This suggests the

presence of an average planned trajectory computed from the initial fixation to the target

location in the exocentric space. Interestingly, this convergence was prominent for goal

movements as compared to non-goal or return movements, implying that they were not a

consequence of a regression to the mean effect (Figs. 4B, 7F, and 7G). Furthermore, similar

zero crossings across dominant and non-dominant arms (Fig. 8D) and across different

movement speeds (Fig. 9D) suggest that the trajectory plan is rather robust across various

contexts and is particularly sensitive to goal-directedness.

The other important aspect of control we tested is whether the CNS implements a fixed

extent of control or if it varies along different phases of movement. Modeling of perturbation

studies on arm movements has suggested that the extent of feedback controller gains is low

near movement initiation, while they increase towards the end of the movement as the hand

reaches the target (Liu & Todorov, 2007). In the context of finger and arm movements, we

found that there were two distinct phases of control (Figs. 3D and 5D), which is reflected at

time points where there is a relatively large change of decorrelation in the early (trough at the

20^{th} percentile) and late (trough at 90^{th} percentile) phases. This was also supported by the

relatively greater magnitudes of zero crossing rate during early and late phases of movement

(Figs. 4C, 6C, and 6E). This pattern is reminiscent of via-point control (Bizzi *et al.*, 1984),

where, post-perturbation, the arm neither corrected towards the target nor the starting position

but ended up at an intermediate point between them. This further reinforces the notion of a

trajectory that can be efficiently corrected at two extreme postural positions when the velocities

of movements are relatively low and state estimation is more robust. Such distinct troughs were

not observed in a previous study since no accuracy constraints were enforced towards the end

of movement (Kuang & Gail, 2015). Implementing control close to the target location is

obviously beneficial and has been known since the time of Bernstein's seminal experiments

(Bongaardt & Meijer, 2000), and having a high degree of control earlier on has the obvious

advantage of clamping variability to manageable levels before the system is swamped by

signal-dependent noise. Indeed, we observed that the initial 10 % and 20 % of the response is

predictive of the subsequent trajectory variability (Fig. 7C). However, the relative absence of

decorrelation between the early and late phases of control suggests that trajectory control is

largely intermittent. Thus, we suggest that optimal control models could incorporate such

trajectory constraints in their cost function (Mistry *et al.*, 2013).

Trajectory control during goal-directed eye movements

For saccadic eye movements, we found that the trajectory control was predominantly

during the early phase (trough at the 10th percentile) and gradually decreased along the time

course of movement, suggesting that the structure of control is effectively different from that

of arm movements. The association between the average decorrelation and the average inter-

trial standard deviation suggests that such decorrelations are likely to be related to the control

of these saccadic movements. Although we could not perform the timing analysis due to the

lack of enough data points, given that the average duration of saccade was 56.5 ± 2.2 ms (mean

$\pm$ SEM) and that most of the decorrelation and zero crossings were during the early phase of

movement, we suggest that the control could be solely driven by internal feedback consistent

with previous work (Guthrie *et al.*, 1983; Wurtz, 2008; Richardson *et al.*, 2011; Varsha *et al.*,

2021).

Interestingly, the absence of the 2^{nd} later peak in saccades relative to arm and finger

movements further suggests that such a later phase of control is likely to be mediated by

sensory/visual feedback control, which is seen in arm and finger movements and is unavailable

in the case of fast saccadic movements. This is also supported by the time evolution of inter-

trial variability, which is bell-shaped in arm movements (Figs. 7A and 8A) while it fans out

with time in saccadic eye movements (Varsha *et al.*, 2021). Nevertheless, in this context, the

presence of zero crossings, particularly during the initial phases of the saccadic movement, is

consistent with some degree of trajectory control (Vasudevan *et al.*, 2023). Although this

finding contrasts with standard models of saccadic control that rely on either displacement or

end-of-movement (Harris & Wolpert, 1998), it is consistent with recent experimental evidence

of trajectory control (Katnani *et al.*, 2012; Smalianchuk *et al.*, 2018) in monkeys as well as a

recently proposed model of saccadic control that simulates kinematic variability in humans

during eye-hand coordination tasks (Vasudevan *et al.*, 2023).

Fast feedback control in normative voluntary movements

Several studies have indicated signatures of relatively fast feedback control involving

reflexive arm movements that occur between 50 – 100 ms after the beginning of movements

during tasks that require posture/ trajectory stabilization following mechanical perturbations

(Messier & Kalaska, 1999; Pruszynski *et al.*, 2008, 2011; Nashed *et al.*, 2014; Cluff & Scott,

2015). However, whether these feedback signals are also at play during unperturbed normative

voluntary goal-directed movements is still not clear. In contrast to such fast sensory feedback,

internal feedback control has been shown to implement corrections using an efference copy of

the initiated motor command to estimate the hand position via a forward model in the absence

of delays associated with sensory feedback (Desmurget & Grafton, 2000). To the best of our

knowledge, these rapid control strategies have been shown to occur when goals and decisions

are changed during movements (Pisella *et al.*, 2000; Gaveau *et al.*, 2003; Desmurget *et al.*,

2005; LoTemplio *et al.*, 2023) to help in fast error corrections. In the context of normative

movements, such rapid control has been only studied in the context of fast saccadic eye

movements using a similar analysis (Richardson *et al.*, 2011). Evidence of rapid control has

also been shown in isometric force application (Gordon & Ghez, 1987), but it has not been

demonstrated in normative voluntary reaching movements.

In our study, the initiation of control was indexed as the time point of significant

decorrelation using the tangent method and this occurred as early as ~60 ms in voluntary finger

movements (Fig. 3C), ~70 ms for voluntary whole-arm reach movements (Figs. 5C and 7D)

and predominantly during the early phase of saccadic eye movements. We interpret this rapid

control in reaching movements to be a consequence of internal feedback supplemented with

fast sensory feedback processes since the onset of goal based decorrelation occurs less than

~50 ms (movement duration) in the case of saccadic eye movements as well as in some subjects

performing the reach movement task (two participants in Fig 7E), which is too fast to allow for

fast sensory feedback to operate. Such rapid control in behavior is posited to be mediated even

more significantly earlier by the neural substrates, owing to conduction and electromechanical

delays (Cavanagh & Komi, 1979; Van Acker *et al.*, 2016). However, this rapid onset of

decorrelation was typically followed with a later stage that is consistent with fast sensory

feedback control with a maximum rate of decorrelation at ~94 ms for the goal finger and ~128

ms for goal movements of the arm (Pruszynski *et al.*, 2011; Scott *et al.*, 2015).

Distinctions and similarities across finger, arm, and saccadic eye movements

The first experiment (Chakrabhavi & Skm, 2019) required time-synchronized cyclic flexion

and extension movements with the goal/instructed finger while it induced involuntary non-goal

movements in the neighboring and non-neighboring fingers because of finger enslavement

(Häger-Ross & Schieber, 2000; Li *et al.*, 2004). In this context, there was a pragmatic

expectation of a trajectory controller for the goal-directed finger movements, and we found

evidence for the same when compared with non-goal movements in the enslaved fingers.

Nevertheless, whether this form of control was specific to cyclic finger movements or whether

it generalized to simple reaching movements with the arm (Schaal *et al.*, 2004), where it is not

clear if the specification of a trajectory is necessary for the control of such movements (Wolpert

& Ghahramani, 2000; Todorov & Jordan, 2002; Liu & Todorov, 2007). This was an open

question we attempted to address in this study. In this context, a previous study (Sendhilnathan

et al., 2021) on saccades comparing movements towards a target as a goal, while task-irrelevant

movements during the inter-trial interval were considered non-goal, found significant

differences in the kinematic variability between goal and non-goal conditions. Neural

correlates of such behavioral differences were also observed in the activity of frontal eye field

neurons that showed no decorrelation in beta band power prior to non-goal saccades, in contrast

to goal-directed saccades. Further, the Fano factor in frontal eye field neurons was much

reduced for goal-directed movements compared to non-goal saccades, which correlated with

the larger kinematic variability observed for non-goal saccades (Sendhilnathan *et al.*, 2021).

Such neural and kinematic differences observed between goal and non-goal saccadic

movements motivated the current approach to study whole arm reaching movements by

considering movements towards a target as goal-directed while return movements during the

inter-trial interval, comprising of exertions towards the center, which was not in contention of

a reward or completion of a trial, be considered a proxy for non-goal condition. Although the

return movements could potentially be guided in anticipation of a reward on the subsequent

trial, nevertheless, goal movements in both finger and arm experiments were directed towards

completion of the trial and were rewarded, while the non-goal/return movements were not.

Moving ahead, it would be interesting to disambiguate the effects of trajectory control from

processes mediating the anticipation of reward. Further, the experimental paradigm involving

saccadic eye movements (Varsha *et al.*, 2021) did not have a hold time. The extent of hold time

is seen to have distinct effects on movement planning (Kapoor & Murthy, 2008). We observed

a sharp decrease in correlation in this case, unlike a gradual sigmoidal-like decrease in

experiments 2 and 3. We speculate that hold time allows precisely timed corrections, unlike the

ballistic spontaneous corrections that we see in experiment 4. Despite these differences in the

task design, we found distinct signatures of control in finger, arm and saccadic eye movements

that could potentially utilize fast internal feedback control to mediate meaningful corrections

towards a planned trajectory during the execution of movements.

References

Abrams, R.A., Meyer, D.E., & Kornblum, S. (1990) Eye-hand coordination: Oculomotor control in rapid aimed limb movements. *Journal of Experimental Psychology: Human Perception and Performance*, **16**, 248–267.

Abrams, R.A. & Pratt, J. (1993) Rapid Aimed Limb Movements: Differential Effects of Practice on Component Submovements. *Journal of Motor Behavior*, **25**, 288–298.

Bertrand, R.L. (2019) Lag Phase Is a Dynamic, Organized, Adaptive, and Evolvable Period That Prepares Bacteria for Cell Division. *Journal of Bacteriology*, **201**, 10.1128/jb.00697-18.

Bizzi, E., Accornero, N., Chapple, W., & Hogan, N. (1984) Posture control and trajectory formation during arm movement. *J. Neurosci.*, **4**, 2738–2744.

Bongaardt, R. & Meijer, O.G. (2000) Bernstein's Theory of Movement Behavior: Historical Development and Contemporary Relevance. *Journal of Motor Behavior*, **32**, 57–71.

Carlton, L.G. (1981) Processing visual feedback information for movement control. *Journal of Experimental Psychology: Human Perception and Performance*, **7**, 1019–1030.

Cavanagh, P.R. & Komi, P.V. (1979) Electromechanical delay in human skeletal muscle under concentric and eccentric contractions. *Europ. J. Appl. Physiol.*, **42**, 159–163.

Chakrabhavi, N. & Skm, V. (2019) Wrist Posture Does Not Influence Finger Interdependence. *J Appl Biomech*, **35**, 410–417.

Chua, R. & Elliott, D. (1993) Visual regulation of manual aiming. *Human Movement Science*, **12**, 365–401.

Churchland, M.M., Santhanam, G., & Shenoy, K.V. (2006) Preparatory activity in premotor and motor cortex reflects the speed of the upcoming reach. *J Neurophysiol*, **96**, 3130–3146.

Cluff, T. & Scott, S.H. (2015) Apparent and Actual Trajectory Control Depend on the Behavioral Context in Upper Limb Motor Tasks. *J. Neurosci.*, **35**, 12465–12476.

Desmurget, M. & Grafton, S. (2000) Forward modeling allows feedback control for fast reaching movements. *Trends in Cognitive Sciences*, **4**, 423–431.

Desmurget, M. & Grafton, S. (2003) Feedback or feedforward control: End of a dichotomy. In *Taking Action: Cognitive Neuroscience Perspectives on Intentional Acts*. The MIT Press, Cambridge, MA, US, pp. 289–338.

Desmurget, M., Turner, R.S., Prablanc, C., Russo, G.S., Alexander, G.E., & Grafton, S.T. (2005) Updating Target Location at the End of an Orienting Saccade Affects the Characteristics of Simple Point-to-Point Movements. *Journal of Experimental Psychology: Human Perception and Performance*, **31**, 1510–1536.

Dick, J.P.R., Benecke, R., Rothwell, J.C., Day, B.L., & Marsden, C.D. (1986) Simple and complex movements in a patient with infarction of the right supplementary motor area. *Movement Disorders*, **1**, 255–266.

Gaveau, V., Martin, O., Prablanc, C., Pélisson, D., Urquizar, C., & Desmurget, M. (2003) On-line modification of saccadic eye movements by retinal signals. *NeuroReport*, **14**, 875.

Georgopoulos, A.P., Kalaska, J.F., Caminiti, R., & Massey, J.T. (1982) On the relations between the direction of two-dimensional arm movements and cell discharge in primate motor cortex. *J. Neurosci.*, **2**, 1527–1537.

Gopal, A., Jana, S., & Murthy, A. (2017) Contrasting speed-accuracy tradeoffs for eye and hand movements reveal the optimal nature of saccade kinematics. *Journal of Neurophysiology*, **118**, 1664–1676.

Gordon, J. & Ghez, C. (1987) Trajectory control in targeted force impulses. III. Compensatory adjustments for initial errors. *Exp Brain Res*, **67**, 253–269.

Gritsenko, V. & Kalaska, J.F. (2010) Rapid online correction is selectively suppressed during movement with a visuomotor transformation. *J Neurophysiol*, **104**, 3084–3104.

Guthrie, B.L., Porter, J.D., & Sparks, D.L. (1983) Corollary Discharge Provides Accurate Eye Position Information to the Oculomotor System. *Science*, **221**, 1193–1195.

Häger-Ross, C. & Schieber, M.H. (2000) Quantifying the Independence of Human Finger Movements: Comparisons of Digits, Hands, and Movement Frequencies. *J. Neurosci.*, **20**, 8542–8550.

Harris, C.M. & Wolpert, D.M. (1998) Signal-dependent noise determines motor planning. *Nature*, **394**, 780–784.

Hollander, M., Wolfe, D.A., & Chicken, E. (2013) *Nonparametric Statistical Methods*. John Wiley & Sons.

Janse, R.J., Hoekstra, T., Jager, K.J., Zoccali, C., Tripepi, G., Dekker, F.W., & van Diepen, M. (2021) Conducting correlation analysis: important limitations and pitfalls. *Clinical Kidney Journal*, **14**, 2332–2337.

Kapoor, V. & Murthy, A. (2008) Covert inhibition potentiates online control in a double-step task. *Journal of Vision*, **8**, 20.

Katnani, H.A., Opstal, A.J.V., & Gandhi, N.J. (2012) Blink Perturbation Effects on Saccades Evoked by Microstimulation of the Superior Colliculus. *PLOS ONE*, **7**, e51843.

Kawato, M. (1999) Internal models for motor control and trajectory planning. *Current Opinion in Neurobiology*, **9**, 718–727.

Kawato, M., Kuroda, T., Imamizu, H., Nakano, E., Miyauchi, S., & Yoshioka, T. (2003) Internal forward models in the cerebellum: fMRI study on grip force and load force coupling. In *Progress in Brain Research*, *Neural Control of Space Coding and Action Production*. Elsevier, pp. 171–188.

Keele, S.W. (1968) Movement control in skilled motor performance. *Psychological Bulletin*, **70**, 387–403.

Khan, M., Franks, I., Elliott, D., Lawrence, G., Chua, R., Bernier, P., Hansen, S., & Weeks, D. (2006) Inferring online and offline processing of visual feedback in target-directed movements from kinematic data. *Neuroscience & Biobehavioral Reviews*, **30**, 1106–1121.

Khan, M.A. & Franks, I.M. (2003) Online Versus Offline Processing of Visual Feedback in the Production of Component Submovements. *Journal of Motor Behavior*, **35**, 285–295.

Kuang, S. & Gail, A. (2015) When adaptive control fails: Slow recovery of reduced rapid online control during reaching under reversed vision. *Vision Research*, On-line Visual Control of Action, **110**, 155–165.

Li, Z.-M., Dun, S., Harkness, D.A., & Brininger, T.L. (2004) Motion enslaving among multiple fingers of the human hand. *Motor Control*, **8**, 1–15.

Liu, D. & Todorov, E. (2007) Evidence for the Flexible Sensorimotor Strategies Predicted by Optimal Feedback Control. *J. Neurosci.*, **27**, 9354–9368.

LoTemplio, S.B., Lopes, C.L., McDonnell, A.S., Scott, E.E., Payne, B.R., & Strayer, D.L. (2023) Updating the relationship of the Ne/ERN to task-related behavior: A brief review and suggestions for future research. *Frontiers in Human Neuroscience*, **17**.

Messier, J. & Kalaska, J.F. (1999) Comparison of variability of initial kinematics and endpoints of reaching movements. *Exp Brain Res*, **125**, 139–152.

Mirabella, G., Pani, P., & Ferraina, S. (2008) Context influences on the preparation and execution of reaching movements. *Cognitive Neuropsychology*, **25**, 996–1010.

Mistry, M., Theodorou, E., Schaal, S., & Kawato, M. (2013) Optimal control of reaching includes kinematic constraints. *Journal of Neurophysiology*, **110**, 1–11.

Moran, D.W. & Schwartz, A.B. (1999) Motor cortical representation of speed and direction during reaching. *J Neurophysiol*, **82**, 2676–2692.

Morasso, P. (1981) Spatial control of arm movements. *Experimental Brain Research*, **42**.

Morris, M.E., Summers, J.J., Matyas, T.A., & Iansek, R. (1994) Current Status of the Motor Program. *Physical Therapy*, **74**, 738–748.

Murthy, A., Ray, S., Shorter, S.M., Schall, J.D., & Thompson, K.G. (2009) Neural Control of Visual Search by Frontal Eye Field: Effects of Unexpected Target Displacement on Visual Selection and Saccade Preparation. *Journal of Neurophysiology*, **101**, 2485–2506.

Nashed, J.Y., Crevecoeur, F., & Scott, S.H. (2014) Rapid Online Selection between Multiple Motor Plans. *J. Neurosci.*, **34**, 1769–1780.

Oldfield, R.C. (1971) The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia*, **9**, 97–113.

Padoa-Schioppa, C., Li, C.-S.R., & Bizzi, E. (2002) Neuronal Correlates of Kinematics-to-Dynamics Transformation in the Supplementary Motor Area. *Neuron*, **36**, 751–765.

Pisella, L., Gréa, H., Tilikete, C., Vighetto, A., Desmurget, M., Rode, G., Boisson, D., & Rossetti, Y. (2000) An "automatic pilot" for the hand in human posterior parietal cortex: toward reinterpreting optic ataxia. *Nat Neurosci*, **3**, 729–736.

Pruszynski, J.A., Kurtzer, I., Nashed, J.Y., Omrani, M., Brouwer, B., & Scott, S.H. (2011) Primary motor cortex underlies multi-joint integration for fast feedback control. *Nature*, **478**, 387–390.

Pruszynski, J.A., Kurtzer, I., & Scott, S.H. (2008) Rapid Motor Responses Are Appropriately Tuned to the Metrics of a Visuospatial Task. *Journal of Neurophysiology*, **100**, 224–238.

Richardson, B.A., Ratneswaran, A., Lyons, J., & Balasubramaniam, R. (2011) The time course of online trajectory corrections in memory-guided saccades. *Exp Brain Res*, **212**, 457–469.

Rosenbaum, D.A. (1985) Motor Programming: A Review and Scheduling Theory. In Heuer, H., Kleinbeck, U., & Schmidt, K.-H. (eds), *Motor Behavior: Programming, Control, and Acquisition*. Springer, Berlin, Heidelberg, pp. 1–33.

Schaal, S., Sternad, D., Osu, R., & Kawato, M. (2004) Rhythmic arm movement is not discrete. *Nat Neurosci*, **7**, 1136–1143.

Scholz, J.P., Schöner, G., & Latash, M.L. (2000) Identifying the control structure of multijoint coordination during pistol shooting. *Experimental Brain Research*, **135**, 382–404.

Scott, S.H. (2004) Optimal feedback control and the neural basis of volitional motor control. *Nat Rev Neurosci*, **5**, 532–545.

Scott, S.H., Cluff, T., Lowrey, C.R., & Takei, T. (2015) Feedback control during voluntary motor actions. *Current Opinion in Neurobiology*, Motor circuits and action, **33**, 85–94.

Scott, S.H., Gribble, P.L., Graham, K.M., & Cabel, D.W. (2001) Dissociation between hand motion and population vectors from neural activity in motor cortex. *Nature*, **413**, 161–165.

Sendhilnathan, N., Basu, D., Goldberg, M.E., Schall, J.D., & Murthy, A. (2021) Neural correlates of goal-directed and non–goal-directed movements. *Proceedings of the National Academy of Sciences*, **118**, e2006372118.

Smalianchuk, I., Jagadisan, U.K., & Gandhi, N.J. (2018) Instantaneous Midbrain Control of Saccade Velocity. *J. Neurosci.*, **38**, 10156–10167.

Smug, B.J., Opalek, M., Necki, M., & Wloch-Salamon, D. (2024) Microbial lag calculator: A shiny-based application and an R package for calculating the duration of microbial lag phase. *Methods in Ecology and Evolution*, **15**, 301–307.

Summers, J.J. & Anson, J.G. (2009) Current status of the motor program: Revisited. *Human Movement Science*, Disruptions of Handwriting, **28**, 566–577.

Todorov, E. & Jordan, M.I. (2002) Optimal feedback control as a theory of motor coordination. *Nature Neuroscience*, **5**, 1226–1235.

Van Acker, G.M., Luchies, C.W., & Cheney, P.D. (2016) Timing of Cortico-Muscle Transmission During Active Movement. *Cereb Cortex*, **26**, 3335–3344.

Varsha, V., Murthy, A., & Padhi, R. (2021) A Stochastic Optimal Control Model with Internal Feedback and Velocity Tracking for Saccadic Eye Movements. *Biomedical Signal Processing and Control*, **68**, 102679.

Vasudevan, V., Gopal, A., Jana, S., Padhi, R., & Murthy, A. (2023) Velocity tracking control best explains the modulation of saccade kinematics during eye-hand coordination. *Eur J Neurosci*, **58**, 2232–2247.

Wolpert, D.M., Diedrichsen, J., & Flanagan, J.R. (2011) Principles of sensorimotor learning. *Nature Reviews Neuroscience*, **12**, 739–751.

Wolpert, D.M. & Ghahramani, Z. (2000) Computational principles of movement neuroscience. *Nat Neurosci*, **3**, 1212–1217.

Wolpert, D.M., Ghahramani, Z., & Jordan, M.I. (1995) An internal model for sensorimotor integration. *Science*, **269**, 1880–1882.

Wolpert, D.M. & Kawato, M. (1998) Multiple paired forward and inverse models for motor control. *Neural Netw*, **11**, 1317–1329.

Wolpert, D.M., Miall, R.C., & Kawato, M. (1998) Internal models in the cerebellum. *Trends in Cognitive Sciences*, **2**, 338–347.

Woodworth, R.S. (1899) Accuracy of voluntary movement. *The Psychological Review: Monograph Supplements*, **3**, i–114.

Wurtz, R.H. (2008) NEURONAL MECHANISMS OF VISUAL STABILITY. *Vision Res*, **48**, 2070–2089.

Figures

Figure 1: **Task design of finger, arm, and saccadic eye experiments and their corresponding raw trajectories.** *A) Experiment 1: Cyclic finger flexion-extension – Left:* The subjects rested their forearm and palms on a fixed platform in a neutral posture (pronation-supination), the thumb finger was restrained to the setup, and the four fingers (index, middle, ring, and little fingers) were suspended and free to move. On any given trial, the goal finger (middle finger, depicted with red color, in this case) was instructed to produce cyclic flexion (peaks) and extension (troughs) movements to specified landmarks shown on the stimulus screen (see methods). The phenomenon of finger enslavement led to involuntary movements in the other non-goal fingers (index, ring and little fingers). *Right:* Sample traces of goal/instructed middle finger task. The traces were from the sensors attached to the distal phalanges of the fingers. In this figure, the goal-directed middle finger movement (red color trace) engaged non-goal movements in the index, ring, and little fingers. *B) Experiment 2,3: Discrete horizontal whole arm reaching movements – Left:* The experimental setup consisted of a robotic arm that the subject held onto and performed simple reaching movements in the horizontal plane by looking at the stimulus presented on a screen placed in front. 3D motion

trackers were mounted on the neck, shoulder, elbow, and wrist joints and on the dorsal side of
the hand. The sensor on the hand was considered for the analysis in this study. *Right:* The
experimental protocol started with the subject fixating a pink circular cursor in a central
fixation box for 500 ms. A red-colored target appeared at an eccentricity of 12cm on the screen.
The target location was block randomized to appear at any of the 8 directions (0, 45, 90, 135,
180, 225, 270, and 315 degrees from the center). Subjects had to stay at the fixation for 1000
ms during hold time, and the central fixation box disappeared, marking the go cue to begin the
movement. Subjects made simple, smooth movements towards the target location within
1000ms and stayed there for an additional 500ms to complete the trial. These movements were
considered goal-directed movements. They then had to get back to the central fixation box for
the start of the subsequent trial. These movements were considered return movements and were
not rewarded. The intertrial interval was 1500 ms. *Bottom:* Sample traces of goal (DG) and
return (DR) movements by the dominant arm and the goal movements by the non-dominant
arm (NDG) of one subject (10 trials each). The goal-directed movements were from the center
to a target location, while the return movements were from the target to the center. *C)*
Experiment 4: Discrete saccadic eye movements – Left: The subjects sat on a chair, head fixed,
and performed discrete saccadic eye movements from the center to a target (randomized, 9
directions – 0^0, 30^0, 45^0, 60^0, 90^0, 120^0, 135^0, 150^0 and 180^0) at an eccentricity of 12 degrees,
while their saccadic movements were recorded using an eye tracker. The saccade had to be
initiated as soon as the target appeared on the screen. *Right:* Sample trajectories of saccades
along the 9 directions (10 trials each) by one sample subject.
996

Figure 2: Behavioural signatures of online control and average trajectory planning. The colors red, blue, green, and pink in A and B represent individual trials of a given task. A) If the CNS implements active online control/correction based on its initial posture, the z-scores at the start of the movement z(0), are posited to change along the time course (z(j) at any time t = j), resulting in crossovers among the trajectories. These crossovers could be captured by a decrease in Spearman's correlation coefficient (ρ). By performing the correlation analysis between movement onset and the subsequent time instances t = $0,...,j,j+1,...T$ (where T is the end of movement), we obtain a time-varying $\rho(t)$. A low value of ρ indicates greater control. B) A case where the CNS does not implement online control, i.e., the trajectories starting farther away from the average remain farther away. The correlation coefficient ρ remains high and closer to 1 along the time course. C) Sample trajectories of the X coordinate of hand movements, directed along 0^0, of a single subject, plotted against percentage time. The dashed black line represents the average trajectory, and red and blue are two selected trials. D) Plot of z-scores for the red and blue trials/trajectories and the average trajectory, $z = 0$ (dashed black line), as depicted in C. To investigate if the implemented online control was directed towards an average planned trajectory, we computed the number of times the trajectories crossed the z = 0 line for a given set of trials (zero-crossing rate - ZCR, see methods). For example, the red trial has three zero-crossings during the movement and is under stronger trajectory control, i.e., a greater propensity towards the average trajectory, while the blue trial had no zero crossings and is said to be under weaker trajectory control.

1020
<u>**Figure 3:**</u> **Spearman's correlation analysis of goal (bold) and non-goal finger movements.**
A) The trajectories of the correlation coefficient, $\rho(t)$, of the four fingers during the goal-
directed index finger task, the middle finger task, the ring finger task, and the little finger task
(mean ± SEM). The decrease in $\rho(t)$ was significantly higher in the goal finger in all four tasks.
Colors: green - index, red - middle, blue - ring, and orange - little fingers. B) Plot of average
decorrelation ($\bar{\rho}$) vs trajectory dispersion showed that the goal fingers (circles) had the lowest
$\bar{\rho}$ and lowest dispersion followed by that of non-goal neighboring fingers (diamonds) and non-
neighboring (triangles) fingers. C) The trajectories, $\rho(t)$ (mean ± SEM), of goal finger, and
non-goal neighboring and non-neighboring fingers plotted from movement onset to 300ms into
the movement, and the online control was initiated at 60.31 ± 1.93 ms, 115.25 ± 15.13 ms, and
163.64 ± 11.67 ms (mean ± SEM), respectively. D) Correlation analysis of goal finger between
consecutive 10 % time instances ($\rho_c(t)$, mean ± SEM). The two distinct phases of control,
early control at 20 % and late control at 90 % of movement duration, were observed. Colors in
Fig. B to D: dark, medium, and light shades of purple represent goal, non-goal neighboring and
non-goal non-neighboring fingers, respectively.
1036
1037
1038

1039
1040
Figure 4: The zero-crossing rate (*ZCR*) of goal (bold) and non-goal finger movements and
the average z-scores. A) *ZCR* computed from *z*-scores of goal and non-goal fingers (mean ±
SEM) during goal-directed index, middle, ring, and little finger tasks. *ZCR* was the highest in
the goal finger of any task (further details in the Results, letter G on the bar represents the
corresponding goal finger during any task). Colors: green - index, red - middle, blue - ring, and
orange - little fingers. B) *z* - scores of goal (solid line) and non-goal (dashed line) fingers for
trajectories (mean ± SEM) starting away ($|z| > 0.5$, dark and light purple respectively) and
closer ($|z| <= 0.5$, black and grey respectively) from the average initial posture ($z = 0$). The
trajectories that started farther away from the average trajectory had a greater propensity to
come towards the average trajectory ($P < 0.001$), which is a potential signature of trajectory
control. This convergence towards average trajectory was prominent among goal movements
compared to non-goal movements ($P < 0.001$). C) *ZCR* of goal fingers plotted for early (0 to
33 %, light grey), middle (34 to 66 %, medium grey), and late (67 to 100 %, dark grey)
movement phases (mean ± SEM). We found that the *ZCR* was significantly greater in the early
and late phases as compared to the middle phase (Adj. $P < 0.001$, both cases).
1056

1057
Figure 5: **Spearman's correlation analysis of goal-directed arm movements.** A) The
correlation co-efficient, $\rho(t)$, plotted (mean ± SEM) for goal movements (red) was found to
decrease along the movement duration while it stayed close to 1 during fixation condition
(black). Goal movements were directed towards the target while the fixation was the time from
movement onset to 1000 ms prior to the onset (in the negative direction of time) while the
cursor was inside the fixation box. B) The average decorrelation ($\bar{\rho}$) was significantly
associated with trajectory dispersion (Pearson's r = 0.57, P = 0.02). A lower correlation value
corresponded with a lower trajectory dispersion. C) The correlation co-efficient, $\rho(t)$, plotted
on actual time scale (mean ± SEM) for goal (red) movements. The time at which a tangent line
(see methods) from $\rho(t)$ of goal movements intersected with the ρ = 1, dashed black line was
at 69.56 ± 3.45 ms (mean ± SEM). D) Correlation analysis of goal movements between
consecutive 10 % - time instances ($\rho_c(t)$, mean ± SEM). The two distinct phases of control
i.e., early control at 20[th] percentile and late control at 80[th] and 90[th] percentile along the
movement duration were observed.
1072

1073
Figure 6: The zero-crossing rate (*ZCR and ZCR2*) and the average z-scores of goal-
directed arm movements. A) The 1-dimensional *ZCR* (mean ± SEM) of goal-directed
movements (red) was significantly higher than during fixation (black, $P < 0.001$). B) The goal
movements were segregated into ones starting farther away ($|z| > 0.5$, red) and closer ($|z| <=$
0.5, black) to the average trajectory ($z = 0$) and we found that the trajectories starting farther
away had greater propensity towards the center than the closer ones (mean ± SEM). C) *ZCR*
(mean ± SEM) of goal movements computed at the early (0 to 33 %, light red), middle (34 to
66 %, medium red), and late (67 to 100 %, dark red) movement phases. The *ZCR* was
significantly greater during early and late phases as compared to the middle phase of movement
(Adj. $P < 0.001$, both cases). D) The 2-dimensional *ZCR2* (mean ± SEM) of goal-directed
movements (red) was significantly higher than during fixation (black, $P < 0.001$). E) The *ZCR2*
(mean ± SEM) was also significantly greater during the early and late phases, compared to the
middle phase (Adj. $P = 0.005$ and Adj. $P < 0.001$, respectively).

1087
1088

<u>Figure 7</u>: Differences in control between goal (red) vs return (blue) reach movements. A)
The trajectory of inter-trial standard deviation (mean ± SEM) for goal and return movements.
Return movements had greater SD than the goal movements. B) The trajectories of the
correlation coefficient, $\rho(t)$, of goal and return movements (mean ± SEM). The average
decrease in correlation ($\bar{\rho}$) was significantly higher in goal movements (P = 0.04). C)
Correlation between $\bar{\rho}$, computed from 0 to 20 % time, with the maximum inter-trial standard
deviation of goal and return trajectories. The initial decorrelation among trajectories was
significantly associated with the maximum inter-trial variability (r = 0.464, P = 0.009) and that
the goal movements had lower $\bar{\rho}$ and lesser variability than return movements. D) The
variations in the correlation coefficient $\rho(t)$ on the actual time scale from movement onset to
500ms into the movement (mean ± SEM), showed a steeper and earlier decrease for goal as
compared to return movements. E) The initiation of control, marked by the tangent method,
showed a significantly earlier onset of control during goal movements as compared to return
movements by 52.58 ± 4.75 ms (mean ± SEM, P < 0.001). F) *ZCR2* (mean ± SEM) of goal
movements was significantly higher than that of return movements (P = 0.034). G) The
convergence of z-scores (mean ± SEM) towards the average trajectory in the farther away
group was significantly greater among goal movements (solid line, red) as compared to return
movements (P < 0.001, dashed line, blue). The solid black line and dashed grey line belong to
the closer group of goal and return movements, respectively.

1108

1109
Figure 8: **Differences in control between dominant (red) and non-dominant arms (black).**
A) Plot of inter-trial standard deviation (mean ± SEM) of trajectories of dominant and non-
dominant arms. The non-dominant arm had a higher average SD than the dominant arm ($P =$
0.004). B) Correlation coefficient, $\rho(t)$ (mean ± SEM), of dominant and non-dominant arms.
The average decorrelation was greater in the dominant arm as compared to the non-dominant
arm ($P = 0.029$). C) The differences in inter-trial SDs were significantly correlated with the
differences in the average decorrelation between dominant and non-dominant arms (r = 0.648,
$P = 0.009$). ND – Non-Dominant, D – Dominant. D) The *ZCR2* (mean ± SEM) measure was
not significantly different between the two conditions ($P = 0.42$).
1119

1120
<u>Figure 9:</u> Control analysis on fast and slow reaching movements. A) Plot of correlation
coefficient trajectories, $\rho(t)$ (mean ±SEM), of fast (green) and slow (red) movements by the
dominant arm. The $\rho(t)$ of fast and slow movements overlapped with each other. B) The
average decorrelation ($\bar{\rho}$) was not significantly different between fast and slow movements (P
= 0.317). C) The average decorrelation was not significantly associated with the maximum SD
($r = 0.249$, $P = 0.289$), even though the maximum SD was higher during fast movements as
compared to slow movements ($P = 0.002$). D) The *ZCR2* was not significantly different
between fast and slow movements ($P = 0.089$).
1129

1130
Figure 10: Distinct signatures of control in saccadic eye movements. A) The trajectory of
the correlation coefficient ($\rho(t)$) of saccadic eye movements (red, mean ±SEM, averaged
across 9 directions). We find signatures of rapid online control with a steep decrease in the
correlation coefficient. B) The average decrease in correlation ($\bar{\rho}$) was significantly associated
with the average inter-trial standard deviation (r = 0.544, P = 0.013). C) The decorrelation
between consequent 10% - time instances ($\rho_c(t)$, blue, mean ± SEM) showed it was highest
during the early phase of movement and gradually decreased with the time course of
movement. This signature of control is unlike that found in finger and arm movements. D) The
2D zero-crossing rate measure (*ZCR2*) of saccades during early (orange), middle (green), and
late (sky blue) phases of movement. Trajectory control was the highest during the early phase
and decreased along the middle (P <0.001) and late phases (P< 0.001) as seen in Figure C.
1142

Tables

Goal Finger	$\bar{\rho}$	Non-Goal Finger	$\bar{\rho}$	Adj. *P*
Index	0.40 ± 0.05	Middle	0.76 ± 0.09	0.017
		Ring	0.89 ± 0.04	<0.001
		Little	0.93 ± 0.02	<0.001
Middle	0.36 ± 0.06	Index	0.56 ± 0.03	0.057
		Ring	0.72 ± 0.03	0.003
		Little	0.87 ± 0.03	<0.001
Ring	0.33 ± 0.05	Index	0.84 ± 0.04	<0.001
		Middle	0.51 ± 0.05	0.060
		Little	0.71 ± 0.04	<0.001
Little	0.35 ± 0.04	Index	0.92 ± 0.02	<0.001
		Middle	0.78 ± 0.04	<0.001
		Ring	0.59 ± 0.07	0.016

Table 1: Pairwise comparison of average decorrelation ($\bar{\rho}$) for goal and non-goal fingers (mean ± SEM). The *P* value in the last column is adjusted for multiple comparisons using Bonferroni correction. The $\bar{\rho}$ The index finger was found to be significantly less than the non-goal fingers.

Goal Finger	Avg. Rate (β)	Non-Goal Finger	Avg. Rate (β)	Adj. P
Index	0.077 ± 0.012	Middle	0.043 ± 0.024	0.330
		Ring	0.030 ± 0.009	0.028
		Little	0.027 ± 0.006	0.012
Middle	0.115 ± 0.032	Index	0.060 ± 0.013	0.276
		Ring	0.057 ± 0.010	0.255
		Little	0.034 ± 0.006	0.060
Ring	0.118 ± 0.032	Index	0.029 ± 0.004	0.046
		Middle	0.078 ± 0.022	0.230
		Little	0.035 ± 0.005	0.045
Little	0.131 ± 0.031	Index	0.024 ± 0.006	0.005
		Middle	0.033 ± 0.006	0.025
		Ring	0.053 ± 0.016	0.020

1150

1151 **Table 2: Pairwise comparison of rate of decrease in correlation (β) for goal and non-goal
1152 fingers (mean ± SEM).** The *P* value in the last column is adjusted for multiple comparisons
1153 using Bonferroni correction. The average rate of decrease in $\rho(t)$ in goal finger was found to
1154 be significantly higher than the non-goal non-neighboring fingers but was not different from
1155 the neighboring fingers.

1156

Goal Finger	Zero Crossing Rate	Non-Goal Finger	Zero Crossing Rate	Adj. P
Index	1.18 ± 0.04	Middle	0.52 ± 0.13	0.004
		Ring	0.28 ± 0.11	<0.001
		Little	0.20 ± 0.07	<0.001
Middle	1.16 ± 0.05	Index	0.84 ± 0.09	0.045
		Ring	0.74 ± 0.05	0.002
		Little	0.37 ± 0.07	<0.001
Ring	1.24 ± 0.04	Index	0.43 ± 0.06	<0.001
		Middle	0.99 ± 0.07	0.033
		Little	0.71 ± 0.08	<0.001
Little	1.19 ± 0.05	Index	0.30 ± 0.04	<0.001
		Middle	0.50 ± 0.06	<0.001
		Ring	0.80 ± 0.11	0.010

Table 3: Pairwise comparison of zero crossing rate (*ZCR*) of goal and non-goal fingers (mean ± SEM). The *P* value in the last column is adjusted for multiple comparisons using Bonferroni correction. The *ZCR* in goal finger was found to be significantly higher than the non-goal fingers.