Internal feedback enables rapid trajectory control during goal-directed finger and whole-arm reaching movements

Niranjan Chakrabhavi [a,b,*], Varadhan SKM [b], Ashitava Ghosal [c], Aditya Murthy [a,*]

[a]Centre for Neuroscience, Indian Institute of Science, Bengaluru 560012, India. [b]Department of Applied Mechanics, Indian Institute of Technology Madras, Chennai 600036, India. [c]Department of Mechanical Engineering, Indian Institute of Science, Bengaluru 560012, India.

[*]Corresponding Authors – Niranjan Chakrabhavi (niranjanc@iisc.ac.in) and Aditya Murthy (adi@isc.ac.in).

Author Contribution: NC conceived the study; NC and VSKM designed the finger experiment; NC and AM designed the arm experiment; NC collected data and performed the data analysis; AM and AG supervised the analysis; NC and AM wrote the manuscript; AG and VSKM reviewed the manuscript.

Competing Interest Statement: The authors declare no competing interest.

Classification: Major – Biological Sciences, Minor – Neuroscience.

Keywords: motor control, trajectory planning, variability, and decorrelation.

This pdf file includes:
1. Main text
2. Figures 1 to 4

Supporting Information: This article contains supporting information.

Abstract

Goal-directed behaviors involve the execution of a variety of eye, hand, and finger movements that follow invariant kinematics consisting of straight-line trajectories and bell-shaped velocity profiles, despite the presence of neuromotor noise and distinct biomechanics. A fundamental unresolved issue in the field is to determine and delineate the extent to which these trajectories are planned or whether trajectories are a consequence of online control. In this study, we addressed this question using a unitless geometric measure along with the rate of zero-crossings to analyze within-trial variability to investigate differences in the time course of trajectories during the presence or absence of a goal in finger and whole-arm reaching movements. We found that the CNS implements control to follow an average trajectory, where the goal-directed movements showed an enhanced degree of trajectory control than the non-goal movements. Further, by performing the analysis on the actual timing, we found behavioral signatures of internal feedback control that might operate on these planned trajectories as early as 30 ms in finger movements and 16.6 ms in whole-arm reach movements. The analysis also revealed that the controller gains varied along the movement and peaked distinctly at early (20%) and late (90%) phases of movement, suggesting that the CNS breaks down the movement into accomplishing virtual way-point objectives during the execution of movement.

Significance Statement

The extent to which movements reflect the unfolding of a prespecified trajectory plan or whether trajectories arise from the characteristics of online control has remained a vexing issue for motor theorists. Using novel metrics of control (decorrelation and zero-crossing rate) and a novel comparison between goal and non-goal conditions in finger and whole-arm reaching movements, we demonstrate that the CNS implements rapid internal feedback control to follow

44 a planned trajectory. Interestingly, the extent of trajectory control was not continuous but was

45 implemented in distinct stages at the early and late phases of movement. Taken together these

46 results provide novel constraints for computational theories of motor control.

Introduction

Normative goal-directed movements are characterized by invariant-kinematic patterns

that consist of straight-line trajectories with bell-shaped velocity profiles that scale with

movement amplitude and speed (1). Numerous psychophysical and neurophysiological studies

highlight the notion that movement attributes are represented/programmed in the central

nervous system well before execution (2–10). Such results have motivated the use of internal

models wherein a complete specification of a trajectory can be made, in principle, before its

implementation (11–15). However, other influential models of motor control suggest that only

certain critical features of the movement such as the endpoint or desired displacement need to

be specified in advance (16–18); here, the neural representation specifying movement

trajectories may only be generated dynamically in real-time during movement execution as a

consequence of sensory feedback that is sensitive to the demands of the task. Thus, it remains

unclear whether movement attributes such as trajectories are planned prior to movement

execution or whether they derive from online feedback control.

One feature that in principle can distinguish between planned and derived movement

trajectories is the timing of their respective control signals that are necessary to mitigate the

effects of motor noise that is a ubiquitous property of muscle and neural activity. Whereas in

the context of planned movement trajectories, fast internal feedback control is posited to use

an efference copy of the planned motor command to make predictions on the sensory

consequences (internal feedback) to control movements in the absence of actual sensory

information (19–21). In contrast, sensory feedback control relies on relatively delayed sensory

(> 50 ms) feedback to make meaningful corrections during the movement (22, 23). While

internal feedback back control has been extensively studied in fast saccadic eye movements

(21, 24, 25) distinct signatures of such control in the context of normative arm movements in

the absence of sensory perturbations have not been experimentally demonstrated.

Using the distinction between the temporal properties of these two forms of control, we

used a novel unitless-geometric approach and compared the profiles in movements which were

either goal/task driven versus a similar movement in the absence of a goal/task to determine

whether trajectory control is based exclusively on sensory feedback (26, 27) or can it also

involve internal feedback control as predicted by some computational theories of motor control

(14, 28). By computing the z-scores of kinematic trajectories and by tracking how they changed

over the time course of movement, using Spearman's decorrelation and the zero-crossing rate,

we show that the motor system can plan a kinematic trajectory which can be guided by internal

feedback control. Further, by examining the temporal changes in decorrelation over the

trajectory we showed that trajectory control is most prominent during the initial and final

phases of the movement.

Materials and Methods

Experiment 1 – Individual cyclic flexion-extension movement of fingers

A detailed description of the experiment can be found in our previously published

article (29). In brief, participants sat on a height-adjustable chair and positioned their right

dominant arm at ~30^0 shoulder flexion and ~120^0 elbow extension as they rested their forearm

and palm on a flat surface in neutral (pronation-supination) forearm posture (Fig. S1A). The

forearm and the thumb were constrained by a strap while the other four fingers of interest were

suspended in air and free to move. The movements of phalanges of all four fingers were

recorded using electromagnetic tracking sensors (Micro Sensor 1.8^{TM}, Liberty tracking system,

Polhemus Inc., Colchester, VT, USA). The task was designed on LabVIEW (National

Instruments Corp, Austin, TX, USA) and the data was sampled at a frequency of 100 Hz. A

total of eight healthy right-handed subjects (5 males) participated in this study. The experiment

was approved by the institute ethics committee at IIT Madras, and all the participants provided

a written informed consent before the start of the experiment.

On any given attempt, the subjects were required to maintain a predefined wrist posture

(0^0 neutral, 30^0 flexion or 30^0 extension) and asked to make cyclic flexion-extension

movements (0.75 Hz movement frequency) at the metacarpophalangeal joint with one of the

instructed fingers (among index, middle, ring, and little) between 75±10% of its maximum

flexion-extension limits and to ignore any movements in the non-instructed fingers. For this

analysis, we considered movements performed by instructed finger as goal-directed and the

movements in non-instructed finger as non-goal condition. Each attempt lasted for 32s with an

initial 2s of rest followed by 30s of cyclic flexion-extension movements (Fig. S1B). Real-time

visual feedback of the distal phalange of the instructed finger was provided on the screen, along

with its movement landmarks, and an audio metronome was played at 1.5 Hz to ensure the

amplitude and timing of the instructed finger movement, respectively. There were 36 attempts

(3 wrist postures x 4 instructed fingers x 3 attempts each) in total. Data from the distal

phalanges of each of the four fingers at neutral wrist posture (0^0) were used in this study.

Specifically, each attempt was segmented into several half-cycles comprising of movements

from flexion (local peaks of middle finger trace in Fig. S1B) to subsequent extension (local

troughs of middle finger trace in Fig. S1B) of instructed finger and the same time stamps were

used to segment the movements of non-instructed fingers. The half cycles were then resampled

to 100 equal spaced segments for analysis in Fig. 2 and the first 300 ms were used for the

analysis on actual timing.

Experiment 2,3 – Horizontal reaching movements by dominant and non-dominant arms

and fast and slow movements

Participants: Twenty-five healthy right-handed subjects (15 subjects for experiment 2,

and 10 subjects for experiment 3; 21 males; Age [min-max]: 21-32) participated in the study.

The experiment was approved by the institute ethics committee at IISc Bengaluru, and all the

participants provided a written informed consent before the start of the experiment. The

Edinburgh Handedness Inventory (30) was used to evaluate their dominant arm.

Experimental Setup: Subjects were made to sit on a chair with a backrest to minimize

the movements of the clavicle (collar bone) and their dominant or the non-dominant arm was

required to rest on a smooth elevated table to facilitate horizontal reaching movements (Fig.

S2A). They were mounted with five electromagnetic 3D motion tracking sensors (Teardrop

Mini sensors, 240Hz, Liberty tracking system, Polhemus Inc., Colchester, USA) on the surface

of the skin at the sternoclavicular joint (neck), right or left acromion process (right or left

shoulder) depending on the task of interest, lateral epicondyle (lateral aspects of the elbow),

dorsal tubercle of radius (wrist) and the dorsal aspect of the hand respectively, by a 3M

micropore surgical tape. The sensor on the dorsal side hand was used for the current analysis.

The lights were switched off in the experimental room to occlude the subject's vision of their

arm posture during the movement. Subjects were required to hold an endpoint robotic

manipulandum (KINARM, 240 Hz, BKIN Technologies Ltd., Kingston, Canada) and perform

a center out discreet horizontal planar reaching movement as required by the task. During the

reaching movement, the subjects were required to look at the stimulus screen (LCD Screen, 60

Hz) in front of them where a circular shaped cursor on the screen was mapped onto the position

of the held manipulandum. All the experimental protocols were coded and executed on

TEMPO/VideoSYNC Software (Reflective Computing, St. Louis, MO, USA). The dataset was

post-processed using MATLAB interface (MathWorks, Massachusetts, USA).

Task Structure: A white fixation box (hollow square, 2cm) appeared at the center of the

workspace where the subjects were required to position the pink-colored cursor inside it by

moving the manipulandum with one of their arms (Fig. S2B). Once they fixated at the center

for 500 ms, a red-colored target (filled square, 1cm) appeared on the screen at an eccentricity

of 12 cm from the center at any of the 8 directions (0^0, 45^0, 90^0, 135^0, 180^0, 225^0, 270^0, or 315^0).

Subjects made smooth unperturbed goal directed movement towards the target location as soon

as the central fixation box disappeared, which was after a delay period of 1000ms. After

initiating the movement, they were given 1000ms to reach the target location. Once they

reached the target, they were required to stay there for 500ms to successfully complete one

trial. The successful completion of the trial was indicated to the subjects by providing a beep

sound. The intertrial interval was set to 1500ms where the cursor had to be brought back to the

center of the screen in the absence of a stimulus and wait for the next trial. These were

considered as non-goal movements as there was no target location and they were not rewarded

for the movement (31). Sample movement traces of goal and non-goal conditions by dominant

arm and the goal directed movements by non-dominant arm are shown in Fig. S2C. If the

subject did not fixate enough at the center or at the target, or if they were not able to execute

movements within the required time, the trial was aborted. The experiment was conducted such

that 40 successful trials were recorded in each of the 8 directions. The same task was now

repeated with the other arm and that the sequence of recruitment of the arms were randomized

across subjects to study differences between dominant and non-dominant arms. Experiment-3

on fast and slow movements was designed similarly with some minor modifications. The target

color was used to cue the subjects about the speed of movement – green for fast (within 400ms)

and red (from 600-1000ms) for slow movements and that the trials were randomly interleaved.

For this experiment, the subjects used their dominant right arm and performed movements

towards 0^0 and 45^0 directions only.

Data pre-processing: The motion data from the Polhemus sensor on the hand was low-

pass filtered with a cut-off frequency of 20 Hz using a fourth order zero lag Butterworth filter

(32). The time at which the hand crossed 5% of its maximum speed during the acceleration and

deceleration phases respectively were noted (29, 33, 34). The time of onset and the end of

movement were identified by descending along the velocity curve from the 5% mark to

whenever it reached zero for the first time. The motion data was then resampled to 100 equally

spaced segments to normalize for time across various trials in analysis of Fig. 3, and 4, and the

data for the first 500ms during the movement was used for the analysis of actual timing in Fig.

3D.

Data analysis

The displacement data (in the direction of flexion- extension) of the four fingers (index,

middle, ring, and little) from experiment 1 (29), the displacement in X and Y directions

(horizontal movement) of the hand from experiment 2 and experiment 3 were used for the

analysis. The number of trials in each of the fingers were selected such that the initial postural

variability was equated between the goal and the corresponding non-goal fingers. For the goal

and non-goal whole-arm movements, the initial variability in hand postures were equated along

both X and Y co-ordinates of each of the directions of movement (Fig. S2D). For dominant

and non-dominant arm movements, the equalization of initial variability was performed by

considering similarity of movements with respect to both arms. For example, the initial

variability along 0^0 directed movements by dominant arm were equated with the 180^0 directed

movements performed by the non-dominant arm (note that 13 out 15 subjects were analyzed

for differences across arms as 2 of them lacked enough number of trials). The initial variability

was similar for fast and slow movements as the trials were randomly interleaved. The

equalization of initial variability was done to eliminate the influence of initial variance on the

decorrelation analysis and restricted to report differences in decorrelation caused due to

differences in movement execution.

For any given task, and at every normalized time point, each trial was assigned a z-

score by subtracting its amplitude from the mean and divided it by the inter-trial standard

deviation (Fig. S3). Subtraction from the mean was performed to track how the trajectories

across different trials moved relative to the average trajectory, and that they were normalized

by the standard deviation, as it changed through the course of movement (Fig. S6A). A sample

of movement trajectories (grey) of the arm, directed towards 0^0, is shown in Fig. S3A and the

z-scores of the corresponding red and blue trajectories are shown in Fig. S3B. The motivation

of this procedure was to analyze the within trial variability by observing changes in the z-scores

of each trial along the duration of movement by providing a unitless geometric approach to

study such variability.

To investigate the structure of within trial variability we performed Spearman's rank

correlation between the z-scores at the start of movement $z(0)$, with the z-scores at subsequent

time instances $z(i)$. In Fig. 1, we have depicted two distinct cases of trajectory control. On the

left side (Fig. 1A), the representative trajectories crossover each other such that their relative

ranks change during the movement. For example, the rank order of trajectories - red, green,

blue, and pink (top to bottom) at time $t = 0$, changes to green, pink, red, and blue at time $t = i$.

This leads to decorrelation among their ranks and a decrease in Spearman's correlation

coefficient (ρ), suggesting for the implementation of active trajectory control, which is intended

to follow an average trajectory. Conversely, on the right side (Fig. 1B), the order of trajectories

remains the same throughout the movement and that the trajectories never converge towards

their average trajectory, thereby resulting in high ρ (close to 1) and weaker/lack of

implementation of such trajectory control. We perform the decorrelation analysis from

movement onset to the end of movement (t = 0 to 100%), to obtain a time varying trajectory

of correlation coefficient, $\rho(t)$. We chose the Spearman's correlation coefficient as we were

particularly interested in the crossovers among trajectories, independent of changes in inter-

trial standard deviation (Fig. S6A) during the time course of movement. To indicate as to why

this analysis provides novel insights as compared to that of previous studies (21, 35, 36), we

performed the decorrelation analysis between any of the time points with the end of movement

for goal and non-goal conditions (Fig. S7). We investigated the average decorrelation ($\bar{\rho}$) over

the entire movement duration to quantify the extent of control. We then fit a logistic function

$\rho fit(t)$, over the trajectories of $\rho(t)$ for each of the subjects and used parameter β to quantify

the rate of decrease of $\rho(t)$ using Eq. 1.

$$\rho fit(t) = a + \frac{1}{1+e^{\beta(\rho(t)-c)}} \quad (1)$$

To substantiate our hypothesis, we computed a dispersion measure for the time

normalized trajectories ($A_{n \, x \, T}$, where A represents the amplitude of trajectories with n trials

and T = 101, number of normalized time instances) by taking the ratio of average inter-trial

standard deviation over the time course, with the average range of movement during the n trials,

in Eq. 2. We divided variability with the range, to account for differences in movement

amplitude in the finger experiment. We then correlated the average decorrelation ($\bar{\rho}$) with its

average dispersion measure (or with maximum standard deviation in Fig. 3B) expecting that a

reduction in $\bar{\rho}$ leads to lower trajectory dispersion.

$$237 \quad Dispersion = \frac{\frac{1}{T}\sum_{j=1}^{T}\sqrt{\frac{\sum_{i=1}^{n}\left(A_{i,j}-\mu_j\right)^2}{n-1}}}{\frac{1}{n}\sum_{i=1}^{n}|\max(A_i)-\min(A_i)|} \quad (2)$$

Where μ_j is the average of amplitude (A) across n trials at time j.

In Fig. S3B, the $z = 0$, represents an average trajectory (dashed line), and the red

trajectory crosses the dashed line three times showing a strong trajectory control while the blue

does not cross the dashed line at all showing weaker trajectory control. To quantify this effect,

we computed a zero-crossing rate (ZCR) from the z-scores of movement trajectories, which

was defined as per Eq. 3. A zero-crossover was considered valid only if the amplitude of

transition (Δz) at the time of cross-over was greater than 0.01. The threshold was chosen as the

ZCR under this value did not yield significant differences between goal and non-goal conditions

and can be attributed to noise. For arm movements, the ZCR plotted in Fig. 3 was further

averaged across X and Y co-ordinates and along all the movement directions.

$$248 \quad ZCR = \frac{Total\ number\ of\ valid\ zero\ crossings}{Total\ number\ of\ trials} \quad (3)$$

To further support the evidence of trajectory control, the starting postures of goal

movements in fingers (Fig. S5A) and arm (Fig. S5B), were divided into two groups, the ones

which were farther away ($|z| > 0.5$, blue) and the ones closer ($|z| <= 0.5$, black) to the average

initial posture ($z = 0$). The average z-scores (across trials) were tracked along the time course

of movement to check the propensity of these two groups to converge towards an average

planned trajectory ($z = 0$, from time $0 - 100$ %). If the central nervous system intended to

follow an average trajectory, the group which was farther away from the initial posture should

show greater convergence towards the average trajectory than the closer ones. The difference

in z-scores at 0^{th} (initial) and 20^{th} percentile time points were used to quantify this effect.

For Fig. 2E and 3F, the decorrelation analysis was performed using a similar technique

described earlier but between subsequent 10 % - time instances. At any percentage time point,

the value of ρ was computed between the z-scores of trajectories at that time point with respect

to an earlier 10%-time instance. At $t = 0$, ρ was equal to 1, as comparison was made between

the z-scores of initial postures with itself, as a reference.

Statistics

The data were checked for normality using the Kolmogorov-Smirnov test. We

performed pairwise t-test for paired comparisons and two-sample t-test for independent

comparisons if the data were normal, and Wilcoxon signed rank test and Wilcoxon rank-sum

test for paired and independent comparisons otherwise. For more than two sample paired

comparisons, we performed a repeated measures ANOVA. Multiple pairwise comparisons

were reported with Bonferroni corrections.

Results

Trajectory control of goal and non-goal finger movements

The differences in the magnitude of control among goal and non-goal fingers were

quantified by computing the average decorrelation ($\bar{\rho}$) during the movement (Fig. 2A). A two-

way repeated measures ANOVA (4 finger-tasks x 2 finger-type, goal or non-goal) showed that

the $\bar{\rho}$ of goal finger was significantly different from the non-goal finger ($P < 0.001$). By

comparing goal and the corresponding non-goal fingers in a pairwise method it was observed

that the extent of decorrelation was highest in the goal fingers (Table S1).

The dynamics of control in goal and non-goal fingers were teased apart by computing

the rate of decrease (β) in correlation co-efficient, $\rho(t)$ (Fig. 2A). A logistic function was fitted

to the dataset to find β (median explained variance = 0.95). A two-way repeated measures

ANOVA (4 finger-tasks x 2 finger-types, goal or non-goal) showed that the rate of decrease of

$\rho(t)$ was significantly different across goal and non-goal fingers ($P < 0.001$). Broadly, by

performing pairwise comparisons, it was found that the rate of decrease of $\rho(t)$ in the goal

directed finger was significantly higher than the non-goal non-neighboring fingers but was

from different from that of neighboring fingers (Table S2).

Further, we checked if the average decorrelation in finger trajectories was

systematically associated with the average trajectory dispersion (Fig. 2B) and found that they

were tightly correlated (Spearman's $\rho = 0.9$, $P < 0.001$), and the average dispersion of goal

finger (median = 0.073) was significantly less than that of non-goal neighboring (median =

0.143, $P < 0.001$) and non-neighboring (median = 0.573, $P < 0.001$) fingers (Fig. 2B). Thus, a

higher degree of trajectory decorrelation was associated with lesser dispersion in trajectories

and vice versa. As goal fingers were associated with varying movement amplitudes, we

compared $\rho(t)$ of all four goal fingers to check the sensitivity of the decorrelation analysis to

the differences in the range of movement among them (Fig. S4). A one-way repeated measures

ANOVA revealed that the $\bar{\rho}$ (Fig. S4B) were not significantly different ($P = 0.72$) and the rate

of decrease of $\rho(t)$ (Fig. S4C) were also not significantly different across the goal fingers (P

= 0.52) but the range of movement (Fig. S4D) were significantly different between them ($P <$

0.001), with index (mean ± SEM: 10.63 ± 0.27 cm) and little (mean ± SEM: 7.42 ± 0.27 cm)

fingers having the highest and lowest movement ranges, respectively.

We then tested the zero-crossing rate (ZCR) of goal and non-goal fingers during the

four different instructed finger tasks to probe the propensity of trajectories to follow an average

trajectory (Fig. 2A and Fig. S3 for the method of computation). A two-way repeated measures

ANOVA (4 finger-tasks x 2 finger-type, goal or non-goal) revealed that the ZCR of goal finger

was significantly different from that of non-goal fingers ($P < 0.001$). By performing pairwise

comparison between goal and non-goal fingers it was found that the goal finger had the highest

ZCR in all the four tasks (Table S3). To support this further, we divided initial postures of goal

fingers into two groups (Fig. S5A), the ones farther away ($|z| > 0.5$, blue) and the ones closer

($|z| <= 0.5$, black) to the average initial posture ($z = 0$) and found that the away group had a

significantly greater propensity to converge towards an average trajectory than the closer ones

($P < 0.001$).

Next, we quantified the differences in the time of control by performing the

decorrelation analysis on the actual timing, from movement onset to 300 ms into the movement.

We found that $\rho(t)$ of the goal finger (dark purple) diverged significantly from that of non-goal

neighboring (medium purple) and non-neighboring (light purple) fingers at 50ms and 30ms,

respectively (Fig. 2D).

Motivated by the previous findings, we examined whether trajectory control was

continuous or occurred in distinct phases along the movement duration. For this, we performed

the decorrelation analysis on the goal finger movements on consecutive 10% - time instances

and found that the controller acted discreetly with two prominent troughs in $\rho(t)$ at 20^{th}

(binominal $P < 0.001$) and 90^{th} percentile time (binominal $P < 0.001$) points (Fig. 2E)

suggesting the presence of early and late phases of control. To substantiate the above finding,

we computed the ZCR of goal finger movements across three movement segments; early (0 to

33 %), middle (34 to 66 %), and late (67 to 100 %) phases of movement duration and found

that the ZCR was significantly higher during early (mean $\pm$ SEM: 0.42 ± 0.01) and late ($0.47 \pm$

0.01) phases as compared to the middle phase (0.25 ± 0.02; $P < 0.001$ for both the

comparisons). Also, it was found that the early phase of control (trough at 20 % - 0.78 ± 0.02)

was not significantly different from that of late phase (trough at 90 % - 0.77 ± 0.03, $P = 0.69$).

Trajectory control of goal and non-goal whole-arm movements

We investigated whether the above analysis could be extended to whole-arm reach

movements and if there were systematic changes in trajectory control for goal and non-goal

movements (Fig. 3; see methods and Fig. S2 for a description of goal and non-goal

movements). We performed the decorrelation analysis (Fig. 3A) and found that $\bar{\rho}$ in goal

movements (mean ± SEM: 0.38 ± 0.02) was significantly less than that of non-goal movements

(0.41 ± 0.01, $P = 0.04$; Fig. S6C). This difference in $\bar{\rho}$ was prominent in the first 50 % of the

movement duration (goal: 0.5 ± 0.02, non-goal: 0.61 ± 0.01, $P < 0.001$). Like the results from

finger experiment, the $\bar{\rho}$ in goal movements was significantly associated with their respective

average trajectory dispersion (Pearson's $r = 0.53$, $P = 0.002$, Fig. S6B) and that the trajectory

dispersion of goal directed movements (mean ± SEM: 0.168 ± 0.003) was significantly less

than that of non-goal movements (0.177 ± 0.003, $P = 0.002$, Fig. S6D). A logistic function

was fit over $\rho(t)$ (median explained variance = 0.97) to find the rate of decrease (β) and it was

found to be significantly higher in goal movements (mean ± SEM: 0.143 ± 0.05) than the non-

goal movements (0.099 ± 0.006, p < 0.001).

As expected, the maximum inter-trial standard deviation of goal movements (mean ±

SEM: 1.17 ± 0.05 cm) was significantly less than the non-goal movements (1.46 ± 0.05 cm, P

< 0.001, Fig. 3B and S6A). We wanted to check if this difference in variability could be

explained by differences in decorrelation between goal and non-goal movements during the

early phases of movement. The $\bar{\rho}$ during the first 20 % - time duration was significantly less in

goal movements (mean ± SEM: 0.81 ± 0.01) than the non-goal movements (0.91 ± 0.01, $P <$

0.001) and found that it was significantly associated with the maximum standard deviation

(Pearson's $r = 0.464$, $P = 0.009$) which occurred much later during the movement (Fig. S6A).

Interestingly, this association was conserved for the duration as early as 10 % into the

movement execution (Pearson's $r = 0.426$, $P = 0.018$).

We also computed the zero-crossing rate (ZCR) for goal and non-goal arm movements

and found that the ZCR of goal movements (mean ± SEM: 1.48 ± 0.03) was significantly greater

than the non-goal movements (1.32 ± 0.02; $P < 0.001$, Fig. 3C). Similar to analysis in the

previous section, we divided initial postures of goal movements into two groups (Fig. S5B),

the ones farther away ($|z| > 0.5$, blue) and the ones closer ($|z| <= 0.5$, black) to the average

initial posture ($z = 0$) and found that the away group had a significantly greater propensity to

converge towards an average trajectory than the closer ones ($P < 0.001$).

To estimate the time of divergence between $\rho(t)$ of goal from non-goal movements, the

decorrelation analysis was done on actual time scale, from start to 500 ms into the movement

(Fig. 3D) and we observed that they diverged at a median time of 16.67 ms across subjects

(Fig. 3E). Further, the maximum change in $\rho(t)$ for goal movements (mean ± SEM: 128.33 ±

5.62 ms) and was significantly earlier than for non-goal movements (201.38 ± 8.02 ms, $P <$

0.001). These differences between goal and non-goal movements were not because of

movement kinematics as the maximum movement speeds were similar across the conditions

($P = 0.62$, Fig. S2E).

Further, to investigate whether the controller was continuous or discrete, we performed

the decorrelation analysis of goal directed movements on consecutive 10 % - time instances.

As was observed for finger movements, we found two distinct troughs at 20[th] (binomial $P <$

0.001) and around 90[th] (trough shared between 80[th] and 90[th] percentiles, binomial $P < 0.001$ for

both time stamps, shared across the participants) percentile time instances, indicating early and

late phase of control respectively (Fig. 3F). Subsequently, the ZCR of goal directed arm

movements were computed across three movement segments; early (0 to 33 %), middle (34 to

66 %), and late (67 to 100 %) phases of movement duration and found that the *ZCR* was

significantly higher during early (mean ± SEM: 0.57 ± 0.01) and late (0.52 ± 0.01) phases as

compared the middle phase (0.38 ± 0.05; *P* < 0.001 for both the comparisons), similar to that

of finger movements. Also, the trough during early phase of control (mean ± SEM: 0.78 ± 0.01)

was significantly less than that of the late phase (0.82 ± 0.01, *P* < 0.001).

Invariance in the extent of decorrelation across arms and for different speeds

We tested the decorrelation analysis across dominant and non-dominant arms to see if

there were systematic changes in trajectory control with acquisition of skill (Fig. 4A and Fig.

S2 for methods). We found that the trajectories overlapped to a large extent and that the $\bar{\rho}$ of

dominant arm (mean ± SEM: 0.36 ± 0.02) was not significantly different from that of the non-

dominant arm (0.39 ± 0.02, *P* = 0.11, Fig. 4B) and the average effect size was small (Cohen's

d - 0.34).

Next, we subjected the decorrelation analysis to different movement speeds, between

fast and slow conditions (Fig. 4C and Fig. S2 for methods) to check how sensitive decorrelation

analysis was to differences in signal dependent noise that was expected to be greater for high

movement speeds. We found that $\rho(t)$ for fast and slow movements were largely overlapping

and that the $\bar{\rho}$ for fast movements (mean ± SEM: 0.44 ± 0.03) was not significantly different

from that of slow movements (0.46 ± 0.05, *P* = 0.377, Fig. 4D) and the average effect size was

small (Cohen's *d* - 0.49).

Taken together, the decorrelation analysis was sensitive to presence or absence of a

goal in finger and whole-arm reaching movements, but it was invariant to changes in skill or

speed of movement.

Discussion

In this study, we used the rate of zero crossings around a mean trajectory as a proxy of trajectory control, and the time course of decorrelation between the initial and subsequent time points as a measure of control and compared these measures in the context of goal and non-goal movements to show that 1) the CNS plans an average trajectory during goal-directed finger and whole-arm reach movements. 2) Trajectory control was found to act at two distinct phases: at early (20 %) and late (around 90 %) phases along the time course of movement. 3) The control (or corrections) in trajectories was initiated as rapidly as 30 ms in goal-directed finger movements and 16.6 ms in goal-directed whole-arm reach movements, providing behavioral evidence for internal feedback control during normative movements.

Relation to previous studies

Since Woodsworth (37), the detailed analysis of the kinematics of limb trajectories has provided insights to understand the relative contributions of motor planning from feedback control (38–41). In addition to assessing the ability of the motor system to regulate variability throughout the trajectory (42), the coefficient of determination (R^2) has been used before as a within-trial estimate of the extent to which early kinematic markers (20, 40, 43, 44) such as peak acceleration and peak velocity determine end-point distributions. In this context, online control has been inferred by measuring the extent to which such markers do not predict the endpoint (i.e., lower R^2). Although inferential, such claims have been validated by manipulating factors such as the availability of visual feedback (45). The decorrelation analysis used in this study uses a similar logic with some important differences that allowed us to reveal novel aspects of trajectory control. First, the earliest kinematic markers typically used are peak acceleration which being (~150 ms) is too late to reveal fast feedback control. Here, in contrast

to previous works (35, 44, 46) that have emphasized the relationship of kinematic markers to

the end of movement, we measured how quickly trajectories diverged from an initial starting

posture. This analysis was facilitated by ensuring that the starting variability between goal and

non-goal movements were the same and that any difference in control would be revealed at the

earliest possible time. Performing decorrelation analysis with respect to the initial posture was

critical to demonstrate effects of fast feedback control as the decorrelation analysis with respect

to the end of movement was not sensitive to differences in the early half of the movement that

we wanted to address in this paper (Fig. S7). Further, the comparison with non-goal

movements that have distinct neural correlates at least in the context of saccade movements

(31), as well distinct variability profiles (Fig. S6A) was particularly useful, since it did not

involve manipulation of target locations (11, 47) or postures(48, 49) that would emphasize

differences in sensory feedback control as opposed to internal feedback control. In this context,

although the lack of any difference in the decorrelation profiles between dominant and non-

dominant arms was surprising to us given the previous literature (50–53), both tasks were

primarily goal driven. Thus, the use of goal and non-goal movements was an important factor

to reveal differences in trajectory control using the decorrelation analysis. Also, systematic

differences between goal and non-goal condition suggest that the crossovers were not merely

because of the phenomenon of regression towards the mean, as both movements started with

the same initial variability. Second, in previous studies it is unclear whether to attribute the

lack of explained variability to trajectory compensation or to signal dependent noise as well as

to nonlinearities that would affect R^2 estimates. Here, the latter potential confound was

mitigated using a Spearman's correlation, while the former potential confound was addressed

by showing correlations between the early decorrelation signal and trajectory dispersions. Also,

if the decorrelation was susceptible to noise, it should have been greater for non-goal

movements, while we found that it was the other way round. Finally, the absence of any

difference in the extent of decorrelation for fast versus slow movements suggests that signal-

dependent noise per se was not a factor.

Trajectory control during goal-directed movements

Although the notion of a straight-line trajectory plan derives from behavioral studies

(3, 4, 54), this notion has been questioned in the context of optimal feedback control models

that are able to predict realistic trajectories even in the absence of cost functions that don't

involve trajectory specifications (27, 55, 56). Further, the demonstration of relatively fast task-

related sensory feedback control (~50 ms) (48, 57) which could in principle shape trajectories

during voluntary movement execution involving active participation of motor cortex (26) also

is key prediction of optimal feedback control models. However, since movement in optimal

feedback models is driven by costs associated with the current posture and the target position,

they don't necessarily require movements to crossover an average trajectory. This is in contrast

with our observations of zero crossings, which suggest the presence of a mean planned

trajectory computed from the initial fixation to the target location. The presence of zero

crossings is unlikely to be a consequence of noise since they are more robustly expressed in

goal directed finger and whole-arm movements, despite normalizing for the initial postural

variability. Thus, we believe the relative paucity of zero crossings and lower decorrelation

primarily reflects the absence of robust trajectory control due to the absence of a visual target

in non-goal movements.

The other important aspect we tested is whether the CNS implements a fixed extent

control along the movement duration or if it is time varying. Modelling of perturbation studies

on arm movements has suggested that the extent of feedback controller gains are low near

movement initiation while they increase towards the end of movement as the hand reaches the

target (27). In our study, however, we found that there were two distinct phases of control

which is reflected at time points where there is a relatively large change of decorrelation, which

was in the (Fig 2E and 3F), the early (trough at 20^{th} percentile) and late (trough at 90^{th}

percentile) phases. This pattern is reminiscent of virtual via-point control (22) and further

reinforces the notion of a mean trajectory that can be efficiently corrected at two extreme

postural positions when the velocities of movements are relatively low and state estimation is

more robust. Such distinct troughs were not observed in a previous study since the initial

variability was clamped and no accuracy constraints were enforced towards the end of

movement (35). Further, while implementing control close to the target location is obviously

beneficial and has been known since the time of Bernstein's seminal experiments (58), having

a high degree of control earlier on has the obvious advantage of clamping variability to

manageable levels before the system is swamped by signal-dependent noise. Indeed, we

observed that the initial 10 % and 20 % of the response is predictive of the subsequent trajectory

variability (Fig. 3B). However, the relative absence of any decorrelation between the early and

late phases of control suggests that trajectory control is intermittent. In this context, the absence

of trajectory control during the middle phase of the movement is consistent with the predictions

of optimal control that don't require explicit trajectory control.

Internal feedback control vs fast sensory feedback control

One of the critical aspects that we addressed in this work is the timing of feedback

control that have bearing on the nature of control signals that enable normative goal directed

movements. Several studies have indicated signatures of relatively fast feedback control

involving reflexive arm movements that occur between 50 – 100 ms after the beginning of

movements (26, 44, 48) during tasks that require posture stabilization following mechanical

perturbations. However, whether these feedback signals are also at play during voluntary goal-directed movements is still not clear. On the other hand, internal feedback control implements corrections to trajectories using an efference copy of the initiated motor command (19) to estimate the hand position via a forward model in the absence of delays associated with sensory feedback (59). However, to the best of our knowledge internal feedback has been shown to occur when goals and decisions are changed during movements (24, 43, 60, 61) and help in fast error corrections. In the context of normative movements such internal feedback control has been only studied in the context of fast saccadic eye movements using a similar decorrelation analysis (21). Evidence of internal feedback control has been shown in isometric force application (20) but it has not been demonstrated in normative voluntary reaching movements. In our study, the initiation of control was indexed as the time point when significant differential decorrelations were observed between goal and non-goal movements, and this occurred as early as 30 ms in voluntary finger movements and at 16.6ms for voluntary whole-arm reach movements. We interpret this rapid control in reach movements to be a signature of internal feedback control. However, this early phase of decorrelation was followed with a later stage that is consistent with fast feedback control with peaks at ~94 ms for goal finger and ~128 ms for goal movements of arm (49).

References

1. P. Morasso, Spatial control of arm movements. *Experimental Brain Research* **42** (1981).

2. S. W. Keele, Movement control in skilled motor performance. *Psychological Bulletin* **70**, 387–403 (1968).

3. D. A. Rosenbaum, "Motor Programming: A Review and Scheduling Theory" in *Motor Behavior: Programming, Control, and Acquisition*, H. Heuer, U. Kleinbeck, K.-H. Schmidt, Eds. (Springer, 1985), pp. 1–33.

4. J. P. R. Dick, R. Benecke, J. C. Rothwell, B. L. Day, C. D. Marsden, Simple and complex
movements in a patient with infarction of the right supplementary motor area. *Movement*
Disorders **1**, 255–266 (1986).

5. M. E. Morris, J. J. Summers, T. A. Matyas, R. Iansek, Current Status of the Motor Program.
Physical Therapy **74**, 738–748 (1994).

6. J. J. Summers, J. G. Anson, Current status of the motor program: Revisited. *Human Movement*
Science **28**, 566–577 (2009).

7. D. W. Moran, A. B. Schwartz, Motor cortical representation of speed and direction during
reaching. *J Neurophysiol* **82**, 2676–2692 (1999).

8. C. Padoa-Schioppa, C.-S. R. Li, E. Bizzi, Neuronal Correlates of Kinematics-to-Dynamics
Transformation in the Supplementary Motor Area. *Neuron* **36**, 751–765 (2002).

9. M. M. Churchland, G. Santhanam, K. V. Shenoy, Preparatory activity in premotor and motor
cortex reflects the speed of the upcoming reach. *J Neurophysiol* **96**, 3130–3146 (2006).

10. P. Cisek, Preparing for Speed. Focus on "Preparatory Activity in Premotor and Motor Cortex
Reflects the Speed of the Upcoming Reach." *Journal of Neurophysiology* **96**, 2842–2843 (2006).

11. D. M. Wolpert, Z. Ghahramani, Computational principles of movement neuroscience. *Nature*
Neuroscience **3**, 1212–1217 (2000).

12. D. M. Wolpert, M. Kawato, Multiple paired forward and inverse models for motor control.
Neural Networks **11**, 1317–1329 (1998).

13. D. M. Wolpert, Z. Ghahramani, M. I. Jordan, An internal model for sensorimotor integration.
Science (New York, N.Y.) **269**, 1880–2 (1995).

14. Desmurget, Grafton, Forward modeling allows feedback control for fast reaching movements.
Trends in cognitive sciences **4**, 423–431 (2000).

15. E. Bizzi, N. Accornero, W. Chapple, N. Hogan, Posture control and trajectory formation during
arm movement. *J. Neurosci.* **4**, 2738–2744 (1984).

16. S. H. Scott, Optimal feedback control and the neural basis of volitional motor control. *Nature*
Reviews Neuroscience **5**, 532–546 (2004).

17. E. Todorov, M. I. Jordan, Optimal feedback control as a theory of motor coordination. *Nature*
neuroscience **5**, 1226–35 (2002).

18. M. Desmurget, S. Grafton, Feedback or feedforward control: End of a dichotomy. *Taking*
action: Cognitive neuroscience perspective on intentional acts, 289–338 (2003).

19. M. Kawato, *et al.*, "Internal forward models in the cerebellum: fMRI study on grip force and
load force coupling" in *Progress in Brain Research*, Neural Control of Space Coding and Action
Production., (Elsevier, 2003), pp. 171–188.

20. J. Gordon, C. Ghez, Trajectory control in targeted force impulses. III. Compensatory
adjustments for initial errors. *Exp Brain Res* **67**, 253–269 (1987).

21.   B. A. Richardson, A. Ratneswaran, J. Lyons, R. Balasubramaniam, The time course of online
trajectory corrections in memory-guided saccades. *Exp Brain Res* **212**, 457–469 (2011).

22.   E. Todorov, M. I. Jordan, Optimal feedback control as a theory of motor coordination. *Nature
Neuroscience* **5**, 1226–1235 (2002).

23.   D. M. Wolpert, J. Diedrichsen, J. R. Flanagan, Principles of sensorimotor learning. *Nature
Reviews Neuroscience* **12**, 739–751 (2011).

24.   V. Gaveau, *et al.*, On-line modification of saccadic eye movements by retinal signals.
NeuroReport **14**, 875 (2003).

25.   M. Kawato, Internal models for motor control and trajectory planning. *Current Opinion in
Neurobiology* **9**, 718–727 (1999).

26.   J. A. Pruszynski, *et al.*, Primary motor cortex underlies multi-joint integration for fast feedback
control. *Nature* **478**, 387–390 (2011).

27.   D. Liu, E. Todorov, Evidence for the Flexible Sensorimotor Strategies Predicted by Optimal
Feedback Control. *J. Neurosci.* **27**, 9354–9368 (2007).

28.   C. M. Harris, D. M. Wolpert, Signal-dependent noise determines motor planning. *Nature* **394**,
780–784 (1998).

29.   N. Chakrabhavi, V. Skm, Wrist Posture Does Not Influence Finger Interdependence. *J Appl
Biomech* **35**, 410–417 (2019).

30.   R. C. Oldfield, The assessment and analysis of handedness: The Edinburgh inventory.
Neuropsychologia **9**, 97–113 (1971).

31.   N. Sendhilnathan, D. Basu, M. E. Goldberg, J. D. Schall, A. Murthy, Neural correlates of goal-
directed and non–goal-directed movements. *Proceedings of the National Academy of Sciences*
118, e2006372118 (2021).

32.   J. P. Scholz, G. Schöner, M. L. Latash, Identifying the control structure of multijoint
coordination during pistol shooting. *Experimental Brain Research* **135**, 382–404 (2000).

33.   S. H. Scott, P. L. Gribble, K. M. Graham, D. W. Cabel, Dissociation between hand motion and
population vectors from neural activity in motor cortex. *Nature* **413**, 161–165 (2001).

34.   A. P. Georgopoulos, J. F. Kalaska, R. Caminiti, J. T. Massey, On the relations between the
direction of two-dimensional arm movements and cell discharge in primate motor cortex. *J.
Neurosci.* **2**, 1527–1537 (1982).

35.   S. Kuang, A. Gail, When adaptive control fails: Slow recovery of reduced rapid online control
during reaching under reversed vision. *Vision Research* **110**, 155–165 (2015).

36.   M. Krüger, A. Straube, T. Eggert, The Propagation of Movement Variability in Time: A
Methodological Approach for Discrete Movements with Multiple Degrees of Freedom.
Frontiers in Computational Neuroscience **11** (2017).

37.   R. S. Woodworth, Accuracy of voluntary movement. *The Psychological Review: Monograph
Supplements* **3**, i–114 (1899).

38.   R. A. Abrams, D. E. Meyer, S. Kornblum, Eye-hand coordination: Oculomotor control in rapid
598   aimed limb movements. *Journal of Experimental Psychology: Human Perception and*
599   *Performance* **16**, 248–267 (1990).

600   39.   R. A. Abrams, J. Pratt, Rapid Aimed Limb Movements: Differential Effects of Practice on
601   Component Submovements. *Journal of Motor Behavior* **25**, 288–298 (1993).

602   40.   L. G. Carlton, Processing visual feedback information for movement control. *Journal of*
603   *Experimental Psychology: Human Perception and Performance* **7**, 1019–1030 (1981).

604   41.   R. Chua, D. Elliott, Visual regulation of manual aiming. *Human Movement Science* **12**, 365–401
605   (1993).

606   42.   M. A. Khan, *et al.*, Inferring online and offline processing of visual feedback in target-directed
607   movements from kinematic data. *Neuroscience & Biobehavioral Reviews* **30**, 1106–1121
608   (2006).

609   43.   M. Desmurget, *et al.*, Updating Target Location at the End of an Orienting Saccade Affects the
610   Characteristics of Simple Point-to-Point Movements. *Journal of Experimental Psychology:*
611   *Human Perception and Performance* **31**, 1510–1536 (2005).

612   44.   J. Messier, J. F. Kalaska, Comparison of variability of initial kinematics and endpoints of
613   reaching movements. *Exp Brain Res* **125**, 139–152 (1999).

614   45.   M. A. Khan, I. M. Franks, Online Versus Offline Processing of Visual Feedback in the Production
615   of Component Submovements. *Journal of Motor Behavior* **35**, 285–295 (2003).

616   46.   B. A. Richardson, A. Ratneswaran, J. Lyons, R. Balasubramaniam, The time course of online
617   trajectory corrections in memory-guided saccades. *Exp Brain Res* **212**, 457–469 (2011).

618   47.   V. Gritsenko, J. F. Kalaska, Rapid online correction is selectively suppressed during movement
619   with a visuomotor transformation. *J Neurophysiol* **104**, 3084–3104 (2010).

620   48.   J. A. Pruszynski, I. Kurtzer, S. H. Scott, Rapid Motor Responses Are Appropriately Tuned to the
621   Metrics of a Visuospatial Task. *Journal of Neurophysiology* **100**, 224–238 (2008).

622   49.   J. A. Pruszynski, *et al.*, Primary motor cortex underlies multi-joint integration for fast feedback
623   control. *Nature* **478**, 387–390 (2011).

624   50.   R. L. Sainburg, D. Kalakanis, Differences in control of limb dynamics during dominant and
625   nondominant arm reaching. *J Neurophysiol* **83**, 2661–2675 (2000).

626   51.   R. L. Sainburg, S. Y. Schaefer, Interlimb Differences in Control of Movement Extent. *Journal of*
627   *Neurophysiology* **92**, 1374–1383 (2004).

628   52.   R. L. Sainburg, Evidence for a dynamic-dominance hypothesis of handedness. *Exp Brain Res*
629   **142**, 241–258 (2002).

630   53.   X. Xiao, H. J. Hu, L. F. Li, L. Li, Comparison of dominant hand to non-dominant hand in
631   conduction of reaching task from 3D kinematic data: Trade-off between successful rate and
632   movement efficiency. *Math Biosci Eng* **16**, 1611–1624 (2019).

54.   P. Morasso, Spatial control of arm movements. *Experimental Brain Research* **42** (1981).

55.   V. Vasudevan, A. Gopal, S. Jana, R. Padhi, A. Murthy, Velocity tracking control best explains the
modulation of saccade kinematics during eye–hand coordination. *European Journal of*
Neuroscience **58**, 2232–2247 (2023).

56.   B. Berret, A. Conessa, N. Schweighofer, E. Burdet, Stochastic optimal feedforward-feedback
control determines timing and variability of arm movements with or without vision. *PLOS*
Computational Biology **17**, e1009047 (2021).

57.   D. M. Wolpert, R. C. Miall, M. Kawato, Internal models in the cerebellum. *Trends in Cognitive*
Sciences **2**, 338–347 (1998).

58.   R. Bongaardt, O. G. Meijer, Bernstein's Theory of Movement Behavior: Historical Development
and Contemporary Relevance. *Journal of Motor Behavior* **32**, 57–71 (2000).

59.   M. Desmurget, S. Grafton, Forward modeling allows feedback control for fast reaching
movements. *Trends in Cognitive Sciences* **4**, 423–431 (2000).

60.   L. Pisella, *et al.*, An "automatic pilot" for the hand in human posterior parietal cortex: toward
reinterpreting optic ataxia. *Nat Neurosci* **3**, 729–736 (2000).

61.   S. B. LoTemplio, *et al.*, Updating the relationship of the Ne/ERN to task-related behavior: A
brief review and suggestions for future research. *Frontiers in Human Neuroscience* **17** (2023).

Figures

Trajectories cross over; Active Trajectory Control	Parallel Trajectories; Lack of Trajectory Control

Figure 1: A behavioural signature of active trajectory control. The colors red, blue, green, and pink represent individual trials of a given task. A) If the CNS has a trajectory plan (an average trajectory) that it intends to follow during the task, the z-scores at the start of the movement $z(0)$, is posited to change along the time course ($z(i)$ at any time t $= i$), due to the implementation of trajectory control. The crossover among trajectories is captured by a decrease in the Spearman's correlation coefficient (ρ). By performing the decorrelation analysis between movement onset and the subsequent time instances t $= 0,...,i,i+1,...T$ (where T is the end of movement), we obtain a time varying $\rho(t)$. A low value of ρ indicates greater trajectory control and such control also entails trajectories to have lower trajectory dispersion. The extent of trajectory control can also be captured by counting the number of zero-crossings in the z-scores, and that it would be more when the system implements a greater degree of control. B) A case where the CNS does not follow an average trajectory. The correlation coefficient ρ remains high and closer to 1 along the time course, accompained with a greater trajectory dispersion and lesser zero crossings.

Figure 2: Decorrelation analysis of goal (bold) and non-goal finger movements. A) The trajectories of correlation co-efficient $\rho(t)$ of the goal directed index, middle ring, and little finger tasks (mean ± SEM). The decrease in $\rho(t)$ is significantly higher in the goal finger of any task. B) Plot of average decorrelation ($\bar{\rho}$) vs average trajectory dispersion shows that the goal fingers (circles) had the lowest $\bar{\rho}$ and lowest trajectory dispersion followed by that of non-goal neighboring fingers (diamonds) and non- neighboring (triangles) fingers. C) The zero-crossing rate (*ZCR*) computed from *z*-scores of goal and non-goal fingers (mean ± SEM) during goal directed index, middle, ring, and little finger tasks. *ZCR* was the highest in the goal finger of any task (further details in the Results, letter G on the bar represents the corresponding goal finger during any task). D) The trajectories of $\rho(t)$ (mean ± SEM) of goal finger, and non-goal neighboring and non-neighboring fingers plotted from movement onset to 300ms into the movement. The $\rho(t)$ in goal fingers significantly deviated from neighboring and non-neighboring fingers at 50ms and 30ms, respectively. E) Decorrelation analysis of goal finger between consecutive 10%-time instances. The two distinct phases of control, early control at 20% and late control at 90% of movement duration were observed. Color in Fig. A and C: green-index, red-middle, blue-ring, and orange-little fingers. Colors in Fig. B, D and E: dark, medium, and light shades of purple represent goal, non-goal neighboring and non-goal non-neighboring fingers respectively.

<u>Figure 3:</u> Decorrelation analysis of goal (red) vs non-goal reach movements (blue). A) The decorrelation trajectories $\rho(t)$ of goal and non-goal movements (mean ± SEM). B) Correlation between $\bar{\rho}$, computed from 0 to 20% time, with the maximum inter-trial standard deviation of goal (red) and non-goal (blue) trajectories. The initial decorrelation among trajectories was significantly associated with the maximum inter-trial variability ($r = 0.464$, $P = 0.009$) and that the goal movements had lower $\bar{\rho}$ and lesser trajectory variability than non-goal movements. C) The zero-crossing rate (*ZCR,* mean ± SEM) was significantly greater during goal movements than that of non-goal movements ($P<0.001$). D) The variations in decorrelation trajectories $\rho(t)$ on actual time scale from movement onset to 500ms into the movement (mean ± SEM), showed a steeper decrease for goal as compared to non-goal movements. E) Histogram of the time at which the $\rho(t)$ of goal movements started diverting from the non-goal movements. The diversion happened very early during the movement (median time =16.67ms). F) Decorrelation analysis of goal movements between consecutive 10%-time instances (mean ± SEM). Two distinct phases of control, early control at 20% and late control at 90% of movement duration were observed.

Figure 4: Decorrelation analysis on reaching movements by dominant and non-dominant arms and different movement speeds. A) The decorrelation trajectories $\rho(t)$ (mean ±SEM) of dominant (red) and non-dominant (black) arms. The trajectories were found to overlap along the time course of movement. B) The average decorrelation ($\bar{\rho}$) was not significantly different across dominant and non-dominant arms ($P = 0.11$). C) Plot of $\rho(t)$ trajectories (mean ±SEM) of fast (green) and slow (red) movements by the dominant arm. It was observed that the trajectories overlapped with each other. D) The average decorrelation ($\bar{\rho}$) was not significantly different between fast and slow movements ($P = 0.38$).

717