1 **Transcriptomic sexual conflict at two evolutionary timescales revealed by experimental**
2 **evolution in *Caenorhabditis elegans***
3
4
5 Short Title: Sexual conflict at two evolutionary timescales
6
7

8 Katja R. Kasimatis[1,2]*, John H. Willis[1], Christine A. Sedore[1], and Patrick C. Phillips[1]*
9
10 [1]Institute of Ecology and Evolution, University of Oregon, Eugene, Oregon, United States of
11 America
12
13 [2]Current address: Department of Biology, University of Virginia, Charlottesville, Virginia,
14 United States of America
15
16 *Corresponding authors:
17 Email: kasimatis@virginia.edu (KRK)
18 Email: pphil@uoregon.edu (PCP)
19

ABSTRACT

Sex-specific regulation of gene expression is the most plausible way for generating sexually differentiated phenotypes from an essentially shared genome. However, since genetic material is shared, sex-specific selection in one sex can have an indirect response in the other sex. From a gene expression perspective, this tethered response can move one sex away from their wildtype expression state and impact potentially many gene regulatory networks. Here, using experimental evolution in the model nematode *Caenorhabditis elegans,* we explore the coupling of direct sexual selection on males with the transcriptomic response in males and females over microevolutionary timescales to uncover the extent to which post-insemination reproductive traits share a genetic basis between the sexes. We find that differential gene expression evolved in a sex-specific manner in males, while in females indirect selection causes an evolved response. Almost all differentially expressed genes were downregulated in both evolved males and females. Moreover, 97% of significantly differentially expressed genes in males and 69% of significantly differentially expressed genes in females have wildtype female-biased expression profile. Changes in gene expression profiles were likely driven through *trans*-acting pathways that are shared between the sexes. We found no evidence that the core dosage compensation machinery was impacted by experimental evolution. Together these data suggest a de-feminization of the male transcriptome and masculinization of the female transcriptome driven by direct selection on male sperm competitive ability. Our results indicate that on short evolutionary timescales sexual selection can generate putative sexual conflict in expression space.

INTRODUCTION

Sexual dimorphism, ubiquitous across multicellular organisms, is rooted in the seeming

conundrum that the sexes can be dramatically different yet share largely the same genome [1,2].

On a functional level, this means that gene expression must be tightly regulated in a sex-specific

manner [3,4]. On an evolutionary scale, it also means that selection acting differentially on one

sex can have an immediate correlated response to selection on attributes of the other sex [5,6].

This correlation underpins the classic Fisherian model of runaway sexual selection, as direct

selection acting on a male display trait generates self-reinforcing indirect selection on female

preference if they share some form of joint inheritance, either through pleiotropy or linkage [7–

9]. Indirect selection can in principle generate a correlated evolutionary response on any number

of female traits, however, particularly on polygenic traits for which alleles are more likely to be

pleiotropic [10]. For example, males of *Pteridophora alberti*, the King of Saxony bird-of-

paradise, have highly elongated feathers equal to several body lengths on either side of their

heads. Curiously, females display similar, but much smaller, feathers on their heads, possibly as

residual responses to sexual selection in males [11,12]. Much like these birds, opposite-sex

perturbation would be expected to be fairly short-lived on an evolutionary timescale, as natural

selection against misexpression in the wrong sex leads to heightened sexual dimorphism [13].

The degree to which mating traits have additive genetic covariance, then, determines the

opportunity for sexual conflict [14,15]. While the evolution of sexual dimorphism has been

extensively studied for macro-level secondary sexual characteristics, such as plumage, territorial

behavior, and body size, much less is known at the molecular level, particularly about the

coupling between the primary domains of dimorphism: the response to sexual selection at the

evolutionary level and the evolution sex-specific gene regulation at the functional level. We

address that coupling here with an experimental evolution framework using the nematode

Caenorhabditis elegans.

In internally fertilizing species, there is an additional layer to the reproductive process to

those classically studied under sexual selection, namely post-insemination reproductive

interactions. Here the male "trait" of interest can be any cell or protein that affects male

fertilization success, while female "preference" is realized in any tissue, cell, or protein that

interacts with the male ejaculate to affect female fertilization success. Sexual selection on male

reproductive proteins has been inferred through molecular evolution analyses [16–19]. These

studies indicate that many seminal fluid proteins evolve rapidly due to sexually antagonistic

coevolution, which in turn implies that direct selection is acting on both female and male post-

insemination traits. However, identifying the specific female traits with which these male

reproductive proteins interact has proved challenging. In this case, transcriptomic analysis can be

useful for identifying wide-spread changes in gene expression after mating to uncover polygenic

responses to selection within and across generations.

Few studies have measured direct selection on male post-insemination traits and the

correlated response in females in a similar manner to male display and female preference traits.

We previously used experimental evolution to select for increased sperm competitive ability in

C. elegans males [20]. Experimental evolution provides an opportunity to reproducibly isolate

directional selection on male characteristics from the potentially correlated responses in females

using timescales when the opportunity for sexual conflict is likely to be highest. We found that

after 30 generations of evolution, males showed a strong, rapid response to post-insemination

selection at both the phenotypic and genomic levels [20]. Specifically, post-insemination

reproductive success increased by greater than four-fold in the evolved populations. This

reproductive fitness increase was underlain by a polygenic response of 82 significance peaks

represented by 57 genes, 10 intergenic regions, and 15 putative pseudogenes. Together we found

that when the androdioecious *C. elegans* mating system is genetically engineered to reflect the

ancestral diecious mating state, there is overwhelming selection to improve maleness.

Here we test if this strong, direct selection acting on male sperm competitive ability had

an indirect impact on females. Specifically, we quantify the impact of direct post-insemination

sexual selection on the transcriptomes of mated ancestral and evolved males and compare how

selection impacts the transcriptome relative to the genome. Additionally, we examine the

transcriptomes of mated females using a fully factorial crossing design of ancestral and evolved

individuals to determine: i) whether strong, direct selection on sperm competition generates a

response in females, and if so, ii) is the female response evolved or plastic, and iii) is the female

response indicative of sexual conflict.

RESULTS

A strong-inference framework for expression evolution

We selected four experimental evolution replicates that evolved under direct selection on sperm

competitive ability for 30 generations to test for changes in the transcriptional profile of mated

males and mated females. Ancestral and evolved males were crossed with the corresponding

generation of ancestral or evolved females to assess how direct selection on males altered the

male transcriptome. For females, we used a full factorial crossing design of ancestral and

evolved females mated with ancestral and evolved males to determine if the female

transcriptome changes and whether those changes were a result of selection (direct or indirect) or

female plasticity. Populations of individuals for each cross were mated for 48 hours after which

individuals of the focal sex were isolated for transcriptomic analysis. The ancestor$_F$-ancestor$_M$

cross represents the baseline transcriptomic profile of both males and females after mating before

experimental evolution occurred. The evolved$_F$-evolved$_M$ cross represents the transcriptomic

profile of males and females after 30 generations of evolution. If direct sexual selection on males

had no impact on a gene's expression level, then there would not be differential expression

between the ancestor$_F$-ancestor$_M$ and evolved$_F$-evolved$_M$ cross for that gene. In males, such genes

would not contribute to the phenotype under selection, namely sperm competitive ability. In

females, such genes would have no genetic covariance between females and males.

If instead, in females, a gene is differentially expressed between the ancestor$_F$-ancestor$_M$

and evolved$_F$-evolved$_M$ crosses, then females are responding to the sexual selection exerted on

males. If the female transcriptional changes are a plastic response to increased sperm

competitiveness, then male generation should be a major predictor, that is females of the

evolved$_F$-ancestor$_M$ cross should have expression profiles similar to the ancestor$_F$-ancestor$_M$

cross and females of the ancestor$_F$-evolved$_M$ should have expression profiles similar to the

evolved$_F$-evolved$_M$ cross. Alternatively, if female transcriptional changes are a result of either

direct selection in females induce by sperm competitiveness or indirect selection generated by

genetic covariation between the sexes, then the expectation is that ancestral females will have

similar expression profiles and evolved females will have similar expression profiles, regardless

of male generation.

Direct selection on males repeatably downregulates the evolved male transcriptome

We analyzed changes in male gene expression for each experimental evolution replicate

separately (Fig S1, Fig S2). Principal component analysis (PCA) was performed to compare

samples basted on male generation (*i.e.*, G0 ancestor or G31 evolved). In each replicate, the first

principal component partitioned samples by male generation and explained 68% to 82% of the

total variance in gene expression (Fig S1). Thus, there was a strong effect of direct selection on

the male transcriptome. Differential gene expression analysis showed an overall signature of

downregulation with 50% (Replicate B2) to 91% (Replicate A3) of genes being downregulated

in evolved males. However, the degree of overlap between significantly differentially expressed

genes (DEs) varied greatly among replicates (Fig S3A). Ninety-eight percent of DEs in Replicate

A3 were significant in at least one other replicate, while only 60% of DEs in Replicate A2 were

shared with at least one other replicate. Replicate A2 also had the most DEs (n = 2,181) and

showed the strongest increase in post-mating reproductive success [20], indicating replicate-

specific variation in the transcriptional response.

Differential evolution between experimental evolution replicates is not surprising and

could be cause by multiple processes, including selection acting on different segregating genetic

variants and genetic drift. To focus on the consistent and repeatable gene expression changes that

are most likely linked with the phenotype under selection, we combined all experimental

evolution replicates. The final combined dataset in males consisted of three independent

ancestor$_F$-ancestor$_M$ crosses, and 12 independent evolved$_F$-evolved$_M$ crosses.

PCA again clustered samples by male generation, such that all ancestral males clustered

and all evolved males clustered (Fig 1A). The first principal component explained 42% of the

variation in total gene expression and separated males by generation. The combined analysis,

therefore, supports the individual replicate analysis and shows a strong effect of direct selection

on the male transcriptome. We analyzed differential gene expression using a generalized linear

model framework using male generation and correcting for average differences among

evolutionary replicates. Differential expression analysis showed 589 genes to be significantly

differentially expressed based on male generation of which 125 had at least a two-fold change in

expression (Fig 1B; File S1). All but 18 DEs were represented in individual replicate analyses,

which indicates that the transcriptional changes identified are likely due to selection (Fig S3B).

Replicate did not contribute to significant differential gene expression (File S1).

Only five genes were significantly upregulated (mean fold increase = 1.0) in evolved

males relative to ancestral males. They are located on autosomes and do not have ontology terms

that clearly connect them to reproduction. The other 584 genes (99%) were downregulated (mean

fold decrease = 1.5) in evolved males relative to ancestral males. The widespread signature of

downregulation is not an artifact of biased count distribution (Fig S4A). Rather, the genes which

are highly upregulated in evolved males tended to occur in specific experimental evolution

replicates, highlighting the importance of multiple levels of replication when identifying

consistent patterns of selection. To determine if SNPs in the coding region or the *cis*-regulatory

region of these genes were responsible for the changes in expression, we examined allele

frequencies in our previously published genomic mapping data for these populations [20]. None

of the DEs had significant allele frequency changes in the coding region. Two genes (*apn-1* and

apa-2) had SNPs in the *cis*-coding region, however, none of these SNPs showed a significant

allele frequency change over the course of experimental evolution (File S2). Together, these data

indicate that direct selection on males is downregulating the male transcriptome through *trans*-

acting pathways.

To assess the functional pathways in which these downregulated genes were involved, we

examined their associated gene ontology (GO) terms and KEGG Pathways. GO enrichment

analysis identified 20 ontology terms that were significantly enriched, many of which were

related to some form of binding: DNA binding (n = 4), RNA binding (n = 2), and protein binding

(n = 1) (File S3). Eight KEGG pathways were significantly enriched, including four in the

replication and repair class, three in the metabolism class, and one in the endocrine system class

(File 3).

Female gene expression changes correlate with males based on female generation

First, to assess variation in the evolutionary response, we analyzed changes in female expression

for each experimental evolution replicate separately. PCA clustered females by generation in

each replicate (Fig S5). The first principal component explained 30% to 46% of the total

variance in gene expressed based on female generation alone. Male generation did not contribute

to variance partitioning in the first two principal components. Replicated B3 had two outlier

transcriptome replicates, which were censored in the downstream analyses (Fig S5). Differential

expression analysis supported that female generation alone determined expression changes as the

male term did not identify DEs (Fig S6). Female DEs recapitulated the male signature of

downregulation with 92% to 96% of DEs being downregulated in evolved females. The majority

of DEs overlapped with at least one other experimental evolution replicate, suggesting less

replicate-specific evolution in females than in males (Fig S3C).

Although the separate analysis of each replicate helps to highlight variance in evolution

response, to more formally address evolutionary changes in the transcriptome that are repeatable

across replicates, we used a combined analysis after correcting for among-replicate variance. The

final censored, combined dataset in females therefore consisted of three independent ancestor$_F$-

ancestor$_M$ crosses, 12 independent ancestor$_F$-evolved$_M$ crosses, 11 independent evolved$_F$-

ancestor$_M$ crosses, and 11 independent evolved$_F$-evolved$_M$ crosses. PCA of the full dataset again

clustered samples by female generation along the first principal component, which explained

28% of the total variance in gene expression levels (Fig 1B). Male generation did not contribute

to variance partitioning in the first two principal components, which captured 46% of the total

variance. Thus, the combined PCA supports the individual replicate analyses and indicates that

the female response is evolved rather than a plastic response based on male generation.

To formally examine potential differences of ancestral and evolved males on the female

transcriptional response, as well as the influence of evolutionary response to heightened sperm

competition of the females themselves, differential expression for individual genes was analyzed

using a generalized linear model using female and male generation, correcting for average

differences among evolutionary replicates. Neither male generation nor replicate contributed to

significant changes in gene expression (Fig 1D; File S1). Rather, female generation determined

differential gene expression, again demonstrating an evolved expression response (Fig 1D). One

hundred genes showed significant differential gene expression of which eight genes had at least a

two-fold change in expression (Fig 1D). Only six genes were significantly upregulated in

evolved (mean fold increase = 0.7) females relative to ancestral females. Five of these genes are

uncharacterized in function. The other 94 DEs were downregulated (mean fold decrease = 1.3) in

evolved females relative to ancestral females (Fig 1D). Again, widespread downregulation is not

an artifact of biased count distribution (Fig S4B). None of the significantly differentially

expressed genes had significant allele frequency changes in the coding region or the *cis*-

regulatory region (File S1). Thus, changes in gene expression were again likely regulated

through *trans*-acting pathways. No GO terms or KEGG pathways were significantly enriched in

female DEs.

Indirect selection on females drives transcriptome evolution

Since the female transcriptome response is evolved, we next examined whether the changes in

female expression were due to direct selection in response to increased sperm competitiveness or

indirect selection as a correlated response to the direct selection acting on males. To this end, we

examined at the correlation in differential gene expression changes between the sexes. Genes that

were only significantly differentially expressed in males (*i.e.*, unique male DEs) were not

correlated with female changes in gene expressed (n = 565, p = 0.27, adjusted R^2 = 0.0004; Fig

2). Rather, these genes changed expression in males alone. This widespread male-specific

transcriptomic response indicates that direct selection on males predominantly changed the male

transcriptome in a sex-specific manner.

232   Genes that were only significantly differentially expressed in females (*i.e.*, unique female

DEs) were, however, significantly correlated with corresponding changes in male expression (n=

76, $F_{4,74}$ = 21.3, p < 0.001, adjusted R^2 = 0.21). Therefore, these genes are largely being

downregulated in females and males, though not significantly so in males (Fig 2). This

correlation holds when all female DEs – unique and shared between the sexes (n = 24) – are

included ($F_{1,98}$ = 42.16, p < 0.001, adjusted R^2 = 0.29). Together, these data support that indirect

selection on females was driving changes in the female transcriptome as female expression

changes were not independent from those in male, while male expression changes occurred in a

sex-specific manner.

Correlated selection de-feminizes the transcriptome of males and females

We hypothesized that selection was driving this widespread downregulation via either female-

biased or non-sex biased genes. In either case, the transcriptomic effect would be a less female-

like and a more male-like transcriptome to support the increase in male reproductive fitness. To

test this hypothesis, we examined the relationship between changes in gene expression after

experimental evolution and sex-biased expression in wildtype females and males [21]. The

majority (n = 571) of male DEs showed some degree of female-biased expression in wildtype

individuals (Fig 3A). A strong negative correlation exists between wildtype sex-bias and gene

expression changes in males, such that more female-biased genes in wildtype were more strongly

downregulated after experimental evolution regardless of whether the gene was located on an

autosome or the X chromosome ($F_{1,568}$ = 1562, p < 0.0001, adjusted R^2 = 0.73). The reduction in

female-bias indicates a de-feminization of the male transcriptome.

253   Similarly, the majority (n = 69) of female DEs also showed some degree of female-biased

expression in wildtype individuals (Fig 3B). Interestingly, 67 of the downregulated genes (81%)

were located on the X chromosome, of which 57 were female-biased (Fig 3B). There is a

negative relationship between sex-biased expression and differential gene expression on the X

chromosome, such that more female-biased genes had a greater fold decrease in expression ($F_{1,65}$

= 12.4, p < 0.001, adjusted R^2 = 0.15). This relationship suggests a pleiotropic effect in a *trans-*

regulatory pathway impacting regulation of X-linked genes. No relationship exists between

autosomal genes and differential gene expression (p = 0.14), though the low number of

autosomal genes likely makes this comparison under-powered. Together, these data support a

correlated expression response in females and indicate a skew towards a less female-like

transcriptome in evolved females.

Significantly more DEs are located on the X chromosome in females than in males ($\chi^2 =$

319.5, d.f. = 1, p < 0.0001) (Fig S7). One major *trans*-regulatory pathway that could explain the

correlation between downregulated genes and the X chromosome is dosage compensation. In

Caenorhabditis nematodes, hermaphrodites/females have two copies of the X chromosome, each

of which are partially downregulated to match the dosage of the single X chromosome in males

[22]. To determine if whole-scale changes in dosage compensation are responsible for the

downregulated pattern in evolved females, we correlated differential gene expression with gene

expression levels for dosage compensated and non-dosage compensated X-linked genes [23].

Only 31 (n_{male} = 2, n_{female} = 20, n_{shared} = 9) of the significant genes located on the X chromosome

were present in both data sets (Fig. 4). Of these, seven were dosage compensated in wildtype

hermaphrodites/females (XX individuals) and three were dosage compensated in both sexes.

These genes were further downregulated in evolved individuals, particularly in females,

potentially suggesting an expression level lower than males. Nineteen genes in total were not

dosage compensated in wildtype hermaphrodites/females, but were downregulated in evolved

females by on average 1.4-fold. Similarly, the mean fold decrease in wildtype expression in

males (XO individuals) for these same genes is 1.9-fold. Eight genes in total were not dosage

compensated in wildtype males and were downregulated in evolved males by an average 2.4-

fold. While not definitive, these results point to a potential breakdown in proper dosage

compensation accompanying widespread transcriptome downregulation.

The core dosage compensation genes *sdc-2*, *sdc-3*, *dpy-27*, and *dpy-30* did not have

significant changes in gene expression after experimental evolution (File S1) nor were there

significant allele frequency changes at or near these genes [20]. Additionally, there were no

significant allele frequency changes at the *rex* or *dox* binding sites, indicating that the dosage

compensation machinery itself is likely not the cause of widespread downregulation in evolved

females. Therefore, alternative pathways must be involved in this de-feminization of the male

and female transcriptomes resulting from direct sexual selection acting on males.

DISCUSSION

Sexual selection acts in a sex-specific manner yet can have an indirect evolutionary response in the other sex due to genetic associations created by their sharing the same genome (apart from heterogametic sex chromosomes, when present). When sexual selection acts on one sex, the other sex can be pulled away from their sex-specific optimum as a correlated response to selection, generating conflict between the sexes. Here we explored how the transcriptome of newly evolved males responded to direct sexual selection on sperm competitive ability [20] and whether this male-specific selection generated indirect selection on females by quantifying the ancestral and evolved transcriptomes of males and females mated with different experimental evolution generations.

Genes with significant expression changes between ancestral and evolved males were almost exclusively downregulated. Moreover, most of these genes are female-biased in wildtype individuals. These results suggest that the evolved male transcriptome moved away from a female-like state. This transcriptomic response supports our previous fitness data [20] and indicates that over the course of experimental evolution these newly derived males improved their reproductive ability by becoming more like males under obligate male-female (dioecious) mating. This highlights the first, and fairly ancient, time horizon for sexual conflict revealed by our data. The transition from the ancestral dioecious mating system within *C. elegans* has led to patterns of gene expression that favor hermaphroditic function—which is essentially very female-like—over male-specific function. Selection for increased male function generated in the populations analyzed here [20], as well as that observed in other studies [24–26], leads to a rapid improvement in male reproductive traits such as sperm size and competitive ability. Our results illustrate that these macro-phenotypic effects are reflected directly in the transcriptome via a shift in the balance of expression-level conflict away from female-biased function.

Further, as a byproduct of selection for improved male fitness, females displayed widespread concomitant signature of downregulation in the evolved female transcriptome. This represents sexual conflict on the second time horizon. Selection on regulatory elements that allow enhanced male performance leads to a short-term correlated response to selection on the female transcriptome as well, making it more male-like. Whether we might expect sex-specific transcriptional regulation to rebalance itself over time depends on the specific fitness impacts of

these transcriptional changes on females, which are difficult to assess here. Overall, the results

indicate a direct de-feminization of the male transcriptome and an indirect masculinization of the

female transcriptome generated by direct selection on male fertilization success. Since gene

expression levels impact trait function and, in turn, fitness, short-term sexual conflict was likely

generated as a byproduct of direct sexual selection in males as female expression was pushed off

its global optimum. This give and take is likely to be an important feature structuring

transcriptional regulation in many male-female species.

The factorial female mating design used here allows for changes in gene expression to be

categorized as a plastic response to increased sperm competitiveness in males or an evolved

response in the females themselves. If females were predisposed to increased sperm interactions,

then male generation should have been a major contributor to gene expression changes. Such a

scenario is analogous to a pre-existing bias model of sexual selection [27] and can lead to direct

selection on female preference/resistance [14,15]. However, we found no evidence that male

evolutionary history impacted differential gene expression. Rather, all expression changes were

determined based on whether a female was from the ancestral or evolved populations, indicating

that differential gene expression was a result of indirect selection on females during experimental

evolution. Under this scenario optimal gene expression differs between the sexes, such that direct

selection optimizing the male transcriptome generates an indirect response in females due to a

between-sex genetic covariance in expression [6,13,28]. For our post-insemination model, selection

was directed on sperm competitive ability. Since selection on males generated a polygenic

response and the expression changes observed were likely due to *trans*-acting pathways, it is

possible that multiple reproductive trait may be impacted.

Interestingly, there was a nearly complete lack of overlap between the genic changes

identified in Kasimatis et al. (2022) [20] and the expression changes characterized here. One

potential reason for this discrepancy is that the genic changes were altering expression of

upstream transcription factors that are expressed earlier in development. Kasimatis et al. (2022)

[20] identified genomic changes in the coding sequence of or nearby intergenic region of four

transcription factors. Of particular note is the transcription factor *sea-1*, which is involved in sex

determination and dosage compensation through hypoactivation of the X chromosome [29].

Genomic analysis identified seven SNPs with a significant allele frequency change located

within a 30 base pair region of intron 2 [20]. It is possible that this region changes regulation of

sea-1 itself or alters splicing in a way that increase the hypoactivation function of *sea-1* and thus
is responsible for the widespread downregulation of X-linked genes.

The majority of downregulated genes in females were located on the X chromosome.
Because males are the heterogametic sex in *C. elegans*, alleles affecting gene expression on the
X chromosome are directly exposed to selection without any potential shielding of an alternative
allele due to dominance, as would occur in heterozygous females. Thus, selection can be more
effective on X-bearing genes, especially when there is sex-specific selection [30–32]. The
response we observe here is thus consistent with the dominance theory of the rapid evolution of
sex chromosomes and their role in speciation [30,33].

Despite the preponderance of signal on the X chromosome, we found no evidence that
mutations in the core dosage compensation pathway were the cause of this widespread
downregulation. Downregulation of both dosage compensated and non-dosage compensated
genes suggests a potential dysregulation of dosage compensation. These changes further support
a less wildtype hermaphrodite/female-like state. Wildtype dosage compensation levels may not
represent an expression level that is optimal for both sexes [34–36]. Sexual conflict over dosage
compensation creates an obvious additional, but perhaps underappreciated, layer to
understanding the evolution of sexual dimorphism across the genotype-phenotype-fitness
landscape.

Previous experimental evolution studies suggest that male-limited evolution over tens of
generations can decrease female fitness [37–39]. Conversely, female-limited or reduced sexual
selection evolution tends to increase female fitness with little to no effect on male fitness [40–
42]. These previous studies therefore suggest that sex-specific selection acting through males
may have a higher potential to generate sexual conflict. However, this previous work has been
blind to the traits under selection and therefore the genetic architecture underlying the sexual
conflict response. Our work to bridge these layers of the genomic, transcriptomic, and fitness
responses suggests a Red Queen scenario [43,44] relating genome evolution and transcriptional
dimorphism during post-insemination interactions. Specifically, on microevolutionary timescales
strong sexual selection on males can rapidly improve "maleness" at the genomic and phenotypic
levels [20] as well as the transcriptomic level, as shown here. In turn, this selection on males
pushes females away from their fitness optimum through indirect selection [45,46]. Females may
then respond through direct selection, which creates a macroevolutionary picture of a stable

female-like state. Thus, examining the functional layers through which sexual dimorphism arises

is critical for understanding the context in which sexual conflict can occur.

Sexual selection drives the evolution of some of the most elaborate phenotypic variation

and complex behaviors. We show that sexual selection is just as powerful an evolutionary force

on post-insemination traits. In particular, on microevolutionary timescales, females may not have

the opportunity to counter the effects of indirect selection leading to females being pulled away

from their optimal state. Given the seemingly high degree of pleiotropy between female and

male reproductive networks, sexual conflict may evolve very rapidly in populations experiencing

strong sexual selection.

393   **MATERIALS AND METHODS**

Worm culture and mating design

We previously evolved six replicated populations of *C. elegans* pseudo-females (*fog-2*) and

males under enhanced post-insemination competition for 30 generations [20]. The complete

experimental evolution details for the between-strain post-insemination selection only (BS-PO)

regime can be found in Kasimatis et al. (2022) [20]. Briefly, evolving males mated with females

for 24 hours after which male sterility was induced to prevent further sperm transfer. Fully fertile

competitor males were then added to the population to generation sperm competition. After a 24-

hour sperm competition period, progeny coming from the evolving males only were collected

and propagated to the next generation. Hence, selection was acting on sperm defensive capability

and sperm longevity. Here we chose four evolved experimental evolution replicates (A2, A3, B2,

B3), which spanned the range of reproductive increase, for transcriptomic analysis.

To generate mated male transcriptomes, we crossed ancestral (strain PX632) females to

ancestral (strain PX632) males and evolved (G31 replicates A2, A3, B2, B3) females to evolved

(G31 replicates A2, A3, B2, B3) males. The evolved crosses were performed for each

experimental evolution replicate. To start a generation, age synchronized larval stage 1 (L1)

worms were plated onto a 10 cm NGM-agar plates seeded with OP50 *Escherichia coli* at 20ºC

with a density of 1,000 worms per plate [47,48]. Late larval stage 4 (L4) females (n = 40) and

males (n = 40) were isolated onto medium NGM-agar plates (60 mm diameter) seeded with

100μL OP50 *E. coli*. Three mating plates were set up for each cross. Mating plates were kept at

20ºC for 48 hours. After the 48-hour mating period, males were collected and pooled (n = 120)

from all three mating plates per cross for bulk RNA isolation. Three biological replicate mating

assays were conducted for each cross, each coming from an independent age synchronization

event (n = 15 total male samples).

To generate mated female transcriptomes, we used a factorial design of ancestral (strain

PX632) and evolved (G31 replicates A2, A3, B2, B3) females mated to ancestral (strain PX632)

and evolved (G31 replicates A2, A3, B2, B3) males. Crosses were performed for each

experimental evolution replicate following the same experimental procedure as males. After the

48-hour mating period, females were collected and pooled (n = 120) from all three mating plates

per cross for bulk RNA isolation. Three biological replicate mating assays were conducted for

each cross, each coming from an independent age synchronization event (n = 39 total female

samples).

Transcriptome sequencing, mapping, and gene calling

We performed bulk mRNA sequencing of mated males and females, separately, on day 3 of

adulthood. Worms were isolated into 150uL of S-Basal buffer and then pelleted to ~20uL. They

were preserved with 250uL Tri-reagent and flash frozen in liquid nitrogen. The samples went

through 10 freeze-thaw cycle before RNA was isolated. RNA was isolated using the KAPA

mRNA HyperPrep Kit (KK8580) with Illumina compatible adapters. Libraries were prepared

using the TrueSeq RNA Library Prep kit (Illumina) starting from 100 ng of RNA. 100bp single

reads for males and 100bp paired-end reads were sequenced on an Illumina HiSeq 4000 at the

University of Oregon Genomics and Cell Characterization Core Facility (Eugene, OR).

Reads were trimmed using skewer v0.2.2 [49] to remove low quality bases and TrueSeq

adapters (parameters: -x AGATCGGAAGAGCACACGTCTGAACTCCAGTCA -y

AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT -t 12 -l 30 -r 0.01 -d 0.01 -q 10). The

trimmed reads were mapped to the *C. elegans* N2 reference genome (PRJNA13758-WS274) [29]

through two passes using STAR v2.5.3a [50]. A table of gene counts was made using HTSeq

v0.9.1 [51] (parameters: -s no -r pos -i ID -t gene -f) based on *C. elegans* N2 reference genome

"longest isoform" annotations.

Differential gene expression analysis

To compare-and-contrast changes in gene expression over time based on the generation (*i.e.,* G0

or G31) of individuals at the time of mating, we tested for differential gene expression using the

package DESeq2 v1.34.0 [52] in R [53]. DESeq2 uses the generalized linear model framework to

estimate the degree to which genes are differentially expressed. We first analyzed each

experimental evolution replicate independently. For males, we fit the model: gene counts ~

$\text{male}_{\text{Anc,Evo}}$. Since we had a full factorial design for mated females, we fit the model: gene counts

~ $\text{female}_{\text{Anc,Evo}}$ + $\text{male}_{\text{Anc,Evo}}$. In females, experimental evolution replicate B3 had two outlier

mating replicates (20_S36_L005 and 21_S37_L005), which were subsequently censored from

the dataset (Fig S5). We then combined all experimental evolution replicates to capture only the

expression changes that were repeatable across experimental evolution and reanalyzed the

differential gene expression model. Normalized count data and gene expression fold change

values were extracted from the DESeq2 linear models. However, significance values were

determined by fitting a generalized linear model in R with generation and replicate as fixed

effects. This approach is more robust to variation between experimental evolution replicates and

therefore provides a more conservative estimate significantly differentially expressed genes.

Significance was based on a Bonferroni cut-off. These censored, combined data were used for

subsequent analyses (Files S1).

We crossed referenced the genes with significant differential expression with the

genomic significance peaks identified in Kasimatis et al. (2022) [20]. Additionally, we examined

whether the *cis*-regulatory regions of significant differentially expressed genes – defined as 2kb

upstream of the start codon – had any SNPs with a significant change in allele frequency, but

none of these met the significance peak threshold defined in Kasimatis et al. (2022) [20] (File

S2).

We tested for enrichment of gene ontology terms and KEGG (Kyoto Encyclopedia of

Genes and Genomes) pathways in significantly differentially expressed genes using the package

goseq [54] in R [53]. goseq performs gene ontology analysis taking transcript length biases into

account. Median transcript lengths were calculated using from

TxDb.Celegans.UCSC.ce11.ensGene [55] based on the UCSC Genome Browser *C. elegans*

genome version "ce11". Gene ontology enrichment was calculated using the "Wallenius" method

with a 0.05 false-discovery rate cutoff [56] (File S3).

Sex-biased gene expression and dosage compensation

We compared the change in expression of significant genes with their wildtype expression

profiles in *C. elegans fog-2* females and males [21] to examine any potential changes in sex-

biased expression profiles. The relationship between sex-biased expression and differentiation

gene expression in each sex was analyzed by fitting a linear model in R.

The dosage compensation profiles for *C. elegans* hermaphrodites (XX individuals) and

males (XO individuals) were taken from Jans et al. (2009), as were the positions of *rex* and *dox*

binding sites. We then correlated dosage compensation profiles with all genes in our

transcriptomic dataset. The positions of *rex* and *dox* sites were compared with genomic SNP data

from Kasimatis et al. (2022) [20] to determine if any significant allele frequency changes

occurred in these regions.

Data Accessibility: Raw transcriptomic reads are available at NCBI SRA under accession

number PRJNA1013082. Differential expression analysis results for the combined data are in

File S1. Cross-referencing of *cis*-regulatory SNPs are available in File S2. Gene ontology and

KEGG pathway enrichment results are available in File S3. All scripts are publicly available on

the GitHub repository: https://github.com/katjakasimatis/expevol_femaletranscriptomics. The

experimental evolution ancestral strain (PX632) and evolved replicates are available from the

Phillips lab upon request.

Acknowledgements: We thank Ruben Lancaster and Alex Smith for experimental assistance

and Anastasia Teterina for transcriptome analysis advice. We thank Locke Rowe and Stephen

Wright for their insightful discussion.

Funding: This research was supported by NIH grant R35GM131838 to PCP.

Author Contributions: KRK and PCP devised the project. KRK and CS performed the crosses.

JHW prepared the transcriptomic libraries. KRK analyzed the data. KRK wrote the manuscript

with the support of the other authors.

REFERENCES

1. Wyman MJ, Stinchcombe JR, Rowe L. A multivariate view of the evolution of sexual

dimorphism. J Evol Biology. 2013;26: 2070–2080.

2. Kasimatis KR, Nelson TC, Phillips PC. Genomic signatures of sexual conflict. J Hered.

2017;108: 780–790.

3. Mank JE. The transcriptional architecture of phenotypic dimorphism. Nat Rev Genet. 2017;1:

1–7.

4. Grath S, Parsch J. Sex-biased gene expression. Annu Rev Genet. 2016;50: 29–44.

5. Lande R, Arnold SJ. The measurement of selection on correlated characters. Evolution.

1983;37: 1210–1236.

6. Lande R, Arnold SJ. Evolution of mating preference and sexual dimorphism. Journal of

Theoretical Biology. 1985;117: 651–664.

7. Fisher RA. The genetical theory of natural selection. Oxford University Press; 1930.

8. Lande R. Models of speciation by sexual selection on polygenic traits. Proc Nat AcadSci.

1981;78: 3721–3725.

9. Kirkpatrick M. Sexual selection and the evolution of female choice. Evolution. 1982;36: 1–12.

10. Turelli M. Effects of pleiotropy on predictions concerning mutation-selection balance for

polygenic traits. Genetics. 1985;111: 165–195.

11. Cooper W, Forshaw J. The birds of paradise and bower birds. Sydney: Collins; 1977.

12. Darwin C. The descent of man and selection in relation to sex. Murray; 1871.

13. Lande R. Sexual dimorphism, sexual selection, and adaptation in polygenic characters. Evol.

1980;34: 292–305.

14. Arnqvist G, Rowe L. Sexual conflict. Princeton University Press; 2005.

15. Gavrilets S, Arnqvist G, Friberg U. The evolution of female mate choice by sexual conflict.

Proc Royal Soc B. 2001;268: 531–539.

16. Chapman T. Seminal fluid-mediated fitness traits in *Drosophila*. Heredity. 2011;87: 511–

521.

17. Begun DJ, Whitley P, Todd BL, Waldrip-Dail HM, Clark AG. Molecular population genetics

of male accessory gland proteins in *Drosophila*. Genetics. 2000;156: 1879–1888.

18. Clark NL, Aagaard JE, Swanson WJ. Evolution of reproductive proteins from animals and

plants. Reproduction. 2006;131: 11–22.

19. Swanson WJ, Vacquier VD. Reproductive protein evolution. Annu Rev Ecol Syst. 2002;33:
161–179.

20. Kasimatis KR, Moerdyk-Schauwecker MJ, Lancaster R, Smith A, Willis JH, Phillips PC.
Post-insemination selection dominates pre-insemination selection in driving rapid evolution of
male competitive ability. PLoS Genetics. 2022;18: e1010063.

21. Albritton SE, Kranz A-L, Rao P, Kramer M, Dieterich C, Ercan S. Sex-biased gene
expression and evolution of the x chromosome in nematodes. Genetics. 2014;197: 865–883.

22. Meyer BJ. The X chromosome in *C. elegans* sex determination and dosage compensation.
Current Opinion in Genetics & Development. 2022;74: 101912.

23. Jans J, Gladden JM, Ralston EJ, Pickle CS, Michel AH, Pferdehirt RR, et al. A condensin-
like dosage compensation complex acts at a distance to control expression throughout the
genome. Gene Dev. 2009;23: 602–618.

24. LaMunyon CW, Ward S. Larger sperm outcompete smaller sperm in the nematode
Caenorhabditis elegans. Proc R Soc Lond B. 1998;265: 1997–2002.

25. Palopoli MF, Peden C, Woo C, Akiha K, Ary M, Cruze L, et al. Natural and experimental
evolution of sexual conflict within *Caenorhabditis* nematodes. BMC Evol Biol. 2015;15: 1–13.

26. Carvalho S, Phillips PC, Teotónio H. Hermaphrodite life history and the maintenance of
partial selfing in experimental populations of *Caenorhabditis elegans*. BMC Evol Biol. 2014;14:
117.

27. Ryan MJ, Fox JH, Wilczynski W, Rand AS. Sexual selection for sensory exploitation in the
frog *Physalaemus pustulosus*. Nature. 1990;343: 66–67.

28. Barker BS, Phillips PC, Arnold SJ. A test of the conjecture that g-matrices are more stable
than b-matrices. Evolution. 2010;64: 2601–2613.

29. Harris TW, Arnaboldi V, Cain S, Chan J, Chen WJ, Cho J, et al. WormBase: a modern model
organism information resource. Nucleic Acids Research. 2019;29: 82.

30. Charlesworth B, Coyne JA, Barton NH. The relative rates of evolution of sex chromosomes
and autosomes. Am Nat. 1987;130: 113–146.

31. Mank JE. Sex chromosomes and the evolution of sexual dimorphism: Lessons from the
Genome. Am Nat. 2009;173: 141–150.

32. Vicoso B, Charlesworth B. Effective population size and the faster-X effect: an extended
model. Evolution. 2009;63: 2413–2426.

33. Charlesworth B. The evolution of sex chromosomes. Science. 1991;251: 1030–1033.

34. Wright AE, Mank JE. Battle of the sexes: Conflict over dosage-sensitive genes and the origin

of X chromosome inactivation. Proc Natl Acad Sci. 2012;109: 5144–5145.

35. Mank JE, Hosken DJ, Wedell N. Some inconvenient truths about sex chromosome dosage

compensation and the potential role of sexual conflict. Evolution. 2011;65: 2133–2144.

36. Gu L, Walters JR. Evolution of sex chromosome dosage compensation in animals: A

beautiful theory, undermined by facts and bedeviled by details. Genome Biol Evol. 2017;9:

2461–2476.

37. Rice WR. Sexually antagonistic male adaptation triggered by experimental arrest of female

evolution. Nature. 1996;381: 232–234.

38. Prasad NG, Bedhomme S, Day T, Chippindale AK. An evolutionary cost of separate genders

revealed by male-limited evolution. The Am Nat. 2007;169: 29–37.

39. Pischedda A, Chippindale AK. Direct benefits of choosing a high fitness mate can offset the

indirect costs associated with intralocus sexual conflict. Evolution. 2017; 1–27.

40. Pitnick S, Brown WD, Miller GT. Evolution of female remating behaviour following

experimental removal of sexual selection. Proc Royal Soc B. 2001;268: 557–563.

41. Pitnick S, Miller GT, Reagan J, Holland B. Males' evolutionary responses to experimental

removal of sexual selection. Proc Royal Soc Lond Ser B Biological Sci. 2001;268: 1071–1080.

42. Holland B, Rice WR. Experimental removal of sexual selection reverses intersexual

antagonistic coevolution and removes a reproductive load. Proc Nat AcadSci. 1999;96: 5083–

5088.

43. Valen LV. A new evolutionary law. Evolutionary theory. 1973;1: 1–30.

44. Brockhurst MA, Chapman T, King KC, Mank JE, Paterson S, Hurst GDD. Running with the

Red Queen: the role of biotic conflicts in evolution. Proc Royal Soc B. 2014;281: 20141382–

20141382.

45. Rice WR, Holland B. The enemies within: intergenomic conflict, interlocus contest evolution

(ICE), and the intraspecific Red Queen. Behav Ecol Sociobiol. 1997;41: 1–10.

46. Rice WR, Chippindale AK. Intersexual ontogenetic conflict. J Evol Biol. 2001;14: 685–693.

47. Brenner S. The genetics of *Caenorhabditis elegans*. Genetics. 1974;77: 71–94.

48. Kenyon C, Chang J, Gensch E, Rudner A, Tabtiang R. A *C. elegans* mutant that lives twice

as long as wild type. Nature. 1993;366: 461–464.

49. Jiang H, Lei R, Ding S-W, Zhu S. Skewer: a fast and accurate adapter trimmer for next-

generation sequencing paired-end reads. BMC Bioinformatics. 2014;15: 182–12.

50. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal RNA-seq aligner. Bioinformatics. 2013;29: 15–21.

51. Putri GH, Anders S, Pyl PT, Pimanda JE, Zanini F. Analysing high-throughput sequencing

data in Python with HTSeq 2.0. Bioinformatics. 2022;38: 2943–2945.

52. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-

seq data with DESeq2. Genome Biol. 2014;15: 31–21.

53. Team RC. R: A language and environment for statistical computing. Foundation for

Statistical Computing; 2020. Available: https://www.R-project.org/

54. Young MD, Wakefield MJ, Smyth GK, Oshlack A. Gene ontology analysis for RNA-seq:

accounting for selection bias. Genome Biol. 2010;11: R14.

55. Team B, Maintainer B. TxDb.Celegans.UCSC.ce11.ensGene: Annotation package for TxDb

object(s). R package version 346. 2019. doi:10.18129/b9.bioc.txdb.celegans.ucsc.ce11.ensgene

56. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. J R Stat Soc: Ser B (Methodol). 1995;57: 289–300.

Figure 1. Differential gene expression is partitioned by ancestral (purple) and evolved (green) individuals. **A)** Principal component analysis of the variance stabilized dataset partitions male generation along PC1. **B)** Differential gene expression was analyzed based on male generation and experimental evolution replicate. Only the male term contributed to gene expression changes (File S1). Genes with a significant change in expression from ancestral to evolved males (shown in green) were determined using a Bonferroni cut-off. Positive expression changes correspond to upregulation in evolved males, while negative expression changes correspond to downregulation

in evolved males. **C)** Principal component analysis of the variance stabilized dataset partitions

female generation along PC1. **D)** Differential gene expression was analyzed based on female

generation, male generation, and experimental evolution replicate. Only the female term

contributed to gene expression changes. Genes with a significant change in expression from

ancestral to evolved females (shown in green) were determined using a Bonferroni cut-off.

Positive expression changes correspond to upregulation in evolved females, while negative

expression changes correspond to downregulation in evolved females.

Figure 2. A comparison of changes in gene expression levels between females and males. Positive expression changes correspond to upregulation in evolved females/males, while negative expression changes correspond to downregulation in evolved females/males. Genes that are not significantly differentially expressed in either sex are shown in gray. Genes that are significantly differentially expressed in males but not in females (*i.e.*, unique male) are shown in blue. Genes that are significantly differentially expressed in females but not in males (*i.e.*, unique female) are shown in pink. Genes that are significantly differentially expressed in both sexes (*i.e.*, shared) are shown in purple. There is a significant relationship between the change in gene expression in females and males for all female DEs ($F_{1,98} = 42.16$, $p < 0.001$, adjusted $R^2 = 0.29$).

Figure 3. Downregulated genes are female-biased. **A)** There is a significant negative correlation

between wildtype sex-bias in expression and changes in expression after experimental evolution

in males. This relationship holds regardless of genomic location. Positive wildtype gene

expression values indicate female-biased expression and negative values indicate male-biased

expression. Both sex-biased expression (female to male) and experimental evolution fold change

are plotted on a $\log_2$ scale. **B)** There is a negative correlation between significant differentially

expressed genes located on the X chromosome (shown in coral) and sex-biased expression in

females. Only 22 genes were located on autosomes (shown in blue).

Figure 4. Downregulation of X-linked genes is impacting dosage compensation. **A)** The

relationship between normalized expression of dosage compensated (circles) and not dosage

compensated (triangles) X-linked genes in wildtype hermaphrodites/females (XX individuals)

versus differential gene expression in males. Genes with significant differential gene expression

in males only are shown in blue and shared genes are shown in purple. Both axes are plotted on a

$\log_2$ scale. **B)** The relationship between normalized expression of dosage compensated (circles)

and not dosage compensated (triangles) X-linked genes in wildtype hermaphrodites/females (XX

individuals) versus differential gene expression in females. Genes with significant differential

gene expression in females only are shown in pink and shared genes are shown in purple.

Figure S1. Principal component analysis of the variance stabilized data for each experimental

evolution replicate partitions male generation along PC1.

Figure S2. Differential gene expression in males for each experimental evolution replicate was analyzed based on male generation. Genes with a significant change in expression from ancestral to evolved males (shown in green) were determined using a Bonferroni cut-off. Positive expression changes correspond to upregulation in evolved males, while negative expression changes correspond to downregulation in evolved males.

Figure S3. The overlap of significant genes across individual experimental evolution replicates

in: **A)** males and **C)** females. The overlap between the individual experimental evolution

replicate analyses and the combined analysis in: **B)** males and **D)** females.

Figure S4. Distribution of normalized gene expression counts against the change in gene

expression for the combined analysis of: **A)** males and **B)** females. Triangles represent points

that fall outside of the y-axis range.

Figure S5. Principal component analysis of the variance stabilized data for each experimental evolution replicate partitions female generation along PC1. Experimental evolution replicate B3 had two outlier transcriptome replicates (circled) and were censored from the final, combined dataset.

Figure S6. Differential gene expression in females for each experimental evolution replicate was analyzed based on female generation and male generation. Genes with a significant change in expression from ancestral to evolved females (shown in green) were determined using a Bonferroni cut-off. Positive expression changes correspond to upregulation in evolved females, while negative expression changes correspond to downregulation in evolved females.

Figure S7. Chromosomal location of significantly differentially expressed genes. **A)** In males,

the majority of DEs are located on the autosomes. Unique male DEs are shown in blue and

shared DEs are shown in purple. **B)** In females, the majority of DEs are located on the X

chromosome. Unique female DEs are shown in pink and shared DEs are shown in purple.

File S1. Differential expression analysis for the combined male and combined female datasets.

File S2. Genomic SNPs located in the cis-regulatory region of DEs.

File S3. Gene ontology analysis and KEGG Pathway analysis for DEs.