1 **Post-insemination sexual selection in males indirectly masculinizes the female**
2 **transcriptome**
3
4

5 Katja R. Kasimatis[1,2]*, John H. Willis[1], and Patrick C. Phillips[1]*
6
7 [1]Institute of Ecology and Evolution, University of Oregon, Eugene, Oregon, United States of
8 American
9
10 [2]Current address: Department of Ecology and Evolutionary Biology, University of Toronto,
11 Toronto, Ontario, Canada
12
13 *Corresponding authors:
14 Email: k.kasimatis@utoronto.ca (KRK)
15 Email: pphil@uoregon.edu (PCP)

16
17
18 Short Title: Sexual Selection in Males Masculinizes the Female Transcriptome
19
20 Keywords: sexual selection, sexual conflict, sex chromosome evolution, transcriptomics, *C.*
21 *elegans*
22
23
24 Abstract Word Count: 231 words
25 Total Word Count: 5,648 words
26

ABSTRACT

Sex-specific regulation of gene expression is the most plausible way for generating sexually

differentiated phenotypes from an essentially shared genome. However, since genetic material is

shared, sex-specific selection in one sex can have an indirect response in the other sex. From a

gene expression perspective, this tethered response can move one sex away from their wildtype

expression state and impact potentially many gene regulatory networks. Here, using

experimental evolution in the model nematode *Caenorhabditis elegans,* we explore the coupling

of direct sexual selection on males with the transcriptomic response in females over

microevolutionary timescales to uncover the extent to which post-insemination reproductive

traits share a genetic basis between the sexes. We find that differential gene expression is driven

by female ancestral or evolved generation alone and that male generation has no impact on

changes in gene expression. Almost all differentially expressed genes were downregulated in

evolved females. Moreover, 80% of these gene were located on the X chromosome and have

wildtype female-biased expression profiles. Changes in gene expression profiles were likely

driven through *trans*-acting pathways that are shared between the sexes. We found no evidence

that the core dosage compensation machinery was impacted by experimental evolution. Together

these data suggest masculinization of the female transcriptome driven by direct selection on male

sperm competitive ability. Our results indicate that on short evolutionary timescales sexual

selection can generate sexual conflict in expression space.

LAY SUMMARY

47
Sexual selection drives the evolution of some of the most dramatic phenotypic differences
between the sexes. Such sexual dimorphism is so common across multicellular organisms that
we often overlook how remarkable it is for shared genetic material to create numerous and
complex sex differences. At an evolutionary level, sexual dimorphism furthers the opportunity
for sex-specific selection to optimize the fitness of a given sex. As a consequence, sex-specific
selection, such as sexual selection, can have an indirect evolutionary response in the other sex
due to genetic associations created by the sexes sharing the same genome. This correlated
evolutionary response can create sexual conflict by shifting a sex away from their fitness
optimum. At the functional level, sexual dimorphism is generated is through sex-specific
regulation of gene expression. Bridging the evolutionary response to sexual selection with the
evolution of sex-specific gene regulation during post-mating interactions has proved challenging.
We previously used experimental evolution to increase male fertility by directly selecting for
increased sperm competitive ability. In this study, we examined the effect of this direct selection
on males on gene expression patterns in females. Differential gene expression was determined by
whether a female was ancestral or evolved generation, indicating that gene expression changes
were an evolved response due to indirect selection on females. Significantly differentially
expressed genes were downregulated in evolved females. These genes tended to be female-
biased in wildtype individuals and located on the X chromosome. The downregulation of X-
linked genes suggests expression levels in females equal to or lower than that in males. Together
these results indicate a less female-like transcriptome after experimental evolution. This supports
a sexual conflict scenario by which direct sexual selection on males indirectly masculinizes the
female transcriptome over short evolutionary timescales.

                                        INTRODUCTION

Sexual dimorphism, ubiquitous across multicellular organisms, is rooted in the seeming

conundrum that the sexes can be dramatically different yet share largely the same genome

(Wyman *et al.*, 2013; Kasimatis *et al.*, 2017). On a functional level, this means that gene

expression must be tightly regulated in a sex-specific manner (Grath & Parsch, 2016; Mank,

2017). On an evolutionary scale, it also means that selection acting differentially on one sex can

have an immediate correlated response to selection on attributes of the other sex (Lande &

Arnold, 1983, 1985). This correlation underpins the classic Fisherian model of runaway sexual

selection, as direct selection acting on a male display trait generates self-reinforcing indirect

selection on female preference if they share some form of joint inheritance, either through

pleiotropy or linkage (Fisher, 1930; Lande, 1981; Kirkpatrick, 1982). Indirect selection can in

principle generate a correlated evolutionary response on any number of female traits, however,

particularly on polygenic traits for which alleles are more likely to be pleiotropic (Turelli, 1985).

For example, males of *Pteridophora alberti*, the King of Saxony bird-of-paradise, have highly

elongated feathers equal to several body lengths on either side of their heads. Curiously, females

display similar, but much smaller, feathers on their heads, possibly as residual responses to

sexual selection in males (Darwin, 1871; Cooper & Forshaw, 1977). Much like these birds,

opposite-sex perturbation would be expected to be fairly short-lived on an evolutionary

timescale, as natural selection against misexpression in the wrong sex leads to heightened sexual

dimorphism (Lande, 1980). The degree to which mating traits have additive genetic covariance,

then, determines the opportunity for sexual conflict (Gavrilets *et al.*, 2001; Arnqvist & Rowe,

2005). While the evolution of sexual dimorphism has been extensively studied for macro-level

secondary sexual characteristics, such as plumage, territorial behavior, and body size, much less

is known at the molecular level, particularly about the coupling between the primary domains of

dimorphism: the response to sexual selection at the evolutionary level and the evolution sex-

specific gene regulation at the functional level. We address that coupling here with an

experimental evolution framework using the nematode *Caenorhabditis elegans*.

In internally fertilizing species, there is an additional layer to the reproductive process to

those classically studied under sexual selection, namely post-insemination reproductive

interactions. Here the male "trait" of interest can be any cell or protein that affects male

fertilization success, while female "preference" is realized in any tissue, cell, or protein that

interacts with the male ejaculate to affect female fertilization success. Sexual selection on male reproductive proteins has been inferred through molecular evolution analyses (Begun *et al.*, 2000; Swanson & Vacquier, 2002; Clark *et al.*, 2006; Chapman, 2011). These studies indicate that many seminal fluid proteins evolve rapidly due to sexually antagonistic coevolution, which in turn implies that direct selection is acting on both female and male post-insemination traits. However, identifying the specific female traits with which these male reproductive proteins interact has proved challenging. In this case, transcriptomic analysis can be useful for identifying wide-spread changes in gene expression after mating to uncover polygenic responses to selection within and across generations.

Few studies have measured direct selection on male post-insemination traits and the correlated response in females in a similar manner to male display and female preference traits. We previously used experimental evolution to select for increased sperm competitive ability in *C. elegans* males (Kasimatis *et al.*, 2022). Experimental evolution provides an opportunity to reproducibly isolate directional selection on male characteristics from the potentially correlated responses in females using timescales when the opportunity for sexual conflict is likely to be highest. We found that after 30 generations of evolution, males showed a strong, rapid response to post-insemination selection at both the phenotypic and genomic levels (Kasimatis *et al.*, 2022). Specifically, post-insemination reproductive success increased by greater than four-fold in the evolved populations. This reproductive fitness increase was underlain by a polygenic response of 82 significance peaks represented by 57 genes, 10 intergenic regions, and 15 putative pseudogenes. Here we test if this strong, direct selection acting on male sperm competitive ability had an indirect impact on females. Specifically, we examine the transcriptomes of mated females using a fully factorial crossing design of ancestral and evolved individuals to determine: i) whether strong, direct selection on sperm competition generates a response in females, and if so, ii) is the female response evolved or plastic, and iii) is the female response indicative of sexual conflict.

MATERIALS AND METHODS

Worm culture and mating design

We previously evolved six replicated populations of *C. elegans* pseudo-females (*fog-2*) and males under enhanced post-insemination competition for 30 generations (Kasimatis *et al.*, 2022).

The complete experimental evolution details for the between-strain post-insemination selection

only (BS-PO) regime can be found in Kasimatis *et al.* (2022). Briefly, evolving males mated

with females for 24 hours after which male sterility was induced to prevent further sperm

transfer. Fully fertile competitor males were then added to the population to generate sperm

competition. After a 24 hour sperm competition period, progeny coming from the evolving males

only were collected and propagated to the next generation. Hence, selection was acting on sperm

defensive capability and sperm longevity. Here we chose four evolved experimental evolution

replicates (A2, A3, B2, B3), which spanned the range of reproductive increase, for female

transcriptomic analysis.

For each experimental evolution replicate, we used a factorial design of ancestral (strain

PX632) and evolved (G31 replicates A2, A3, B2, B3) females mated to ancestral (strain PX632)

and evolved (G31 replicates A2, A3, B2, B3) males. To start a generation, age synchronized

larval stage 1 (L1) worms were plated onto a 10 cm NGM-agar plates seeded with OP50

Escherichia coli at 20ºC with a density of 1,000 worms per plate (Brenner, 1974; Kenyon *et al.*,

1993). Late larval stage 4 (L4) females (n = 40) and males (n = 40) were isolated onto medium

NGM-agar plates (60 mm diameter) seeded with 100μL OP50 *E. coli*. Three mating plates were

set up for each cross. Mating plates were kept at 20ºC for 48 hours. After the 48 hour mating

period, females were collected and pooled (n = 120) from all three mating plates per cross for

bulk RNA isolation. Three biological replicate mating assays were conducted for each cross,

each coming from an independent age synchronization event (n = 39 total samples).

Transcriptome sequencing, mapping, and gene calling

We performed bulk RNA sequencing of mated females on day 3 of adulthood. Worms were

isolated into 150uL of S-Basal buffer and then pelleted to ~20uL. They were preserved with

250uL Tri-reagent and flash frozen in liquid nitrogen. RNA was isolated using the Direct-zol

RNA Miniprep kit (Zymo). Libraries were prepared using the TrueSeq Stranded mRNA Library

Prep kit (Illumina) starting from 100 ng of RNA. 100bp paired-end reads were sequenced on an

Illumina HiSeq 4000 at the University of Oregon Genomics and Cell Characterization Core

Facility (Eugene, OR).

Reads were trimmed using skewer v0.2.2 (Jiang *et al.*, 2014) to remove low quality bases

and TrueSeq adapters (parameters: -x AGATCGGAAGAGCACACGTCTGAACTCCAGTCA -

y AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT -t 12 -l 30 -r 0.01 -d 0.01 -q 10). The

trimmed reads were mapped to the *C. elegans* N2 reference genome (PRJNA13758-WS274)

(Harris *et al.*, 2019) through two passes using STAR v2.5.3a (Dobin *et al.*, 2013). A table of

gene counts was made using HTSeq v0.9.1 (Putri *et al.*, 2022) (parameters: -s no -r pos -i ID -t

gene -f) based on *C. elegans* N2 reference genome "longest isoform" annotations.

Differential gene expression analysis

To compare-and-contrast changes in gene expression over time based on the generation (*i.e.,* G0

or G31) of individuals at the time of mating, we tested for differential gene expression using the

package DESeq2 v1.34.0 (Love *et al.*, 2014) in R (R Core Team, 2020). DESeq2 uses the

generalized linear model framework to estimate the degree to which genes are differentially

expressed. We fit the model: gene counts ~ female$_{Anc,Evo}$ + male$_{Anc,Evo}$ + female$_{Anc,Evo}$:male$_{Anc,Evo}$.

We first analyzed each experimental evolution replicate independently. The male (ancestor

versus evolved) and interaction (ancestor$_{F,M}$ versus evolved$_{F,M}$) contrasts for each experimental

evolution replicate both showed no gene expression changes based on a Bonferroni cut-off

applied to adjusted p-values. The female (ancestor versus evolved) contrast had 51, 42, 23, and

32 significantly differentially expressed genes in experimental evolution replicates A2, A3, B2,

and B3, respectively (File S1). Principal component analysis of the variance stabilized data

supported variance partitioning by female, but not male, generation (Fig. S1). Experimental

evolution replicate B3 had two outlier mating replicates (20_S36_L005 and 21_S37_L005),

which were subsequently censored from the dataset. Given the consistency in results, we pooled

all experimental evolution replicates and reanalyzed the differential gene expression model. To

account for multiple testing, p-values were adjusted using a Benjamin and Hochberg correction

and significance was based on a Bonferroni cut-off. These censored, pooled data are represented

in the results (Files S1 and S2).

The molecular and biological functions of the genes with significant differential

expression were determined using gene ontology analysis in UniProt (Consortium, 2021) and

QuickGO (Huntley *et al.*, 2015). General reproduction keywords were taken from the WormBase

gene descriptions (Harris *et al.*, 2019). We tested for enrichment of gene ontology terms in

significantly differentially expressed genes using the package goseq (Young *et al.*, 2010) in R (R

Core Team, 2020). goseq performs gene ontology analysis taking transcript length biases into

account. Median transcript lengths were calculated using from

TxDb.Celegans.UCSC.ce11.ensGene (Team & Maintainer, 2019) based on the UCSC Genome

Browser *C. elegans* genome version "ce11". Gene ontology enrichment was calculated using the

"Wallenius" method with a 0.05 false-discovery rate cutoff (Benjamini & Hochberg, 1995) (File

S3).

We crossed referenced the genes with significant differential expression with the

genomic significance peaks identified in Kasimatis *et al.* (2022). Additionally, we examined

whether the *cis*-regulatory regions of significant differentially expressed genes – defined as 2kb

upstream of the start codon – had any SNPs with a significant change in allele frequency, but

none of these met the significance peak threshold defined in Kasimatis *et al.* (2022) (File S4).

Sex-biased gene expression and dosage compensation

We compared the change in expression of significant genes with their wildtype expression

profiles in *C. elegans fog-2* females and males (Albritton *et al.*, 2014) to examine any potential

changes in sex-biased expression profiles. To determine if significantly differentially expressed

genes were more female- or male-biased, we used a χ^2 test on the 1x3 contingency table of

significant genes with wildtype female-biased, non-biased, or male-biased expression. The

relationship between sex-biased expression and differentiation gene expression was analyzed by

fitting a linear model in R.

The dosage compensation profiles for *C. elegans* hermaphrodites (XX individuals) and

males (XO individuals) were taken from Jans et al. (2009), as were the positions of *rex* and *dox*

binding sites. We then correlated dosage compensation profiles with all genes in our

transcriptomic dataset. To determine if difference in dosage compensation profiles varied

between significant and non-significant differentially expressed genes, we used a χ^2 test on the

2x2 contingency table. The positions of *rex* and *dox* sites were compared with genomic SNP data

from Kasimatis *et al.* (2022) to determine if any significant allele frequency changes occurred in

these regions.

<div align="center">**RESULTS**</div>

A strong-inference framework for expression evolution

We used a full factorial crossing design of ancestral and evolved females mated with ancestral

and evolved males from four experimental evolution replicates that were evolved under direct

selection on sperm competitive ability. Populations of individuals for each cross were mated for

48 hours after which females were isolated for transcriptomic analysis. The ancestor$_F$-ancestor$_M$

cross represents the baseline transcriptomic profile of females after mating before experimental

evolution occurred. The evolved$_F$-evolved$_M$ cross represents the transcriptomic profile of females

after 30 generations of evolution. If direct sexual selection on males had no impact on a gene's

expression level in females, then there would not be differential expression between the

ancestor$_F$-ancestor$_M$ and evolved$_F$-evolved$_M$ cross for that gene. Such genes would have no

genetic covariance between females and males.

If instead a gene is differentially expressed between the ancestor$_F$-ancestor$_M$ and

evolved$_F$-evolved$_M$ crosses, then females are responding to the sexual selection exerted on males.

If the female transcriptional changes are a plastic response to increased sperm competitiveness,

then male generation should be a major predictor, that is females of the evolved$_F$-ancestor$_M$ cross

should have expression profiles similar to the ancestor$_F$-ancestor$_M$ cross and females of the

ancestor$_F$-evolved$_M$ should have expression profiles similar to the evolved$_F$-evolved$_M$ cross.

Alternatively, if female transcriptional changes are a result of either direct selection in females

induce by sperm competitiveness or indirect selection generated by genetic covariation between

the sexes, then the expectation is that ancestral females will have similar expression profiles and

evolved females will have similar expression profiles, regardless of male generation.

We analyzed changes in gene expression for each experimental evolution replicate

separately (Fig S1). Since the same pattern was observed, we pooled all four experimental

evolution replicates, excluding the two outlier transcriptome replicates in experimental evolution

replicate B3, for greater power. The final censored, pooled dataset therefore consisted of three

independent ancestor$_F$-ancestor$_M$ crosses, 12 independent ancestor$_F$-evolved$_M$ crosses, 11

independent evolved$_F$-ancestor$_M$ crosses, and 11 independent evolved$_F$-evolved$_M$ crosses.

Female generation determines gene expression changes

Principal component analysis (PCA) was performed to compare samples based on female and

male generation (*i.e.,* G0 or G31) during mating. The first principal component explained 28% of

the total variance in gene expression levels based on female generation alone, such that all

ancestral females clustered together and all evolved females clustered together, regardless of

male generation (Fig 1A). Male identity did not contribute to variance partitioning in the first

two principal components, which captured 46% of the total variance.

Differential gene expression was analyzed using a model that fit female generation, male

generation, and the interaction between female and male generations. Neither male generation

nor the interaction term contributed to changes in gene expression (Fig 1B). Rather, female

generation determined differential gene expression, indicating an evolved expression response

(Fig 1B). Sixty-four genes showed significant differential gene expression of which 11 genes had

at least a two-fold change in expression (Fig 1B; Fig 2). Only three genes were significantly

upregulated in evolved females (mean fold increase = 1.2). All three genes are uncharacterized in

function and have no ontology terms relating them to reproduction (File S2). The other 61

significant genes were downregulated (mean fold decrease = 1.5) in evolved females relative to

ancestral females (Fig 1B; Fig 2). To determine if SNPs in the coding region or the *cis*-

regulatory region of these genes were responsible for the changes in expression, we examined

allele frequencies in our previously published genomic mapping data for these populations

(Kasimatis *et al.*, 2022). None of the significantly differentially expressed genes had significant

allele frequency changes in the coding region or the *cis*-regulatory region (File S2 & S4). Thus,

changes in gene expression were likely regulated through *trans* acting pathways.

To assess the functional pathways and biological processes in which these downregulated

genes were involved, we examined their associated gene ontology (GO) terms (File S1). The

most common molecular functions were DNA binding (n = 4), hydrolase activity (n = 4), and

transferase activity (n = 4). The most common biological process was vesicle-mediated transport

(n = 9) followed by membrane organization (n = 3) and lipid metabolic processes (n = 3). Six

genes were pseudogenes. Seventeen genes (28%) were uncharacterized in both molecular

function and biological process. Nineteen genes (31%) had a keyword relating them to

reproduction in their WormBase description (Harris *et al.*, 2019). GO enrichment analysis

identified 10 ontology terms that were significantly enriched in differentially expressed genes,

including intracellular membrane-bounded organelle (n_{DE} = 30, n_{total} = 4792, p < 0.00001),

membrane-bounded organelle (n_{DE} = 30, n_{total} = 4952, p < 0.00001), and vesicle-mediated

transport (n_{DE} = 10, n_{total} = 470, p < 0.00001) (File S3).

X Chromosome Masculinization

If widespread downregulation in evolved females was a result of direct selection in females, then

we hypothesize that selection would be countering the effects of increased sperm

competitiveness by either upregulating female-biased genes or downregulating male-biased

genes. We examined the relationship between changes in gene expression and sex-biased

expression in wildtype females and males (Albritton *et al.*, 2014). The majority (n = 46) of genes

with a significant expression change (excluding the four pseudogenes) showed some degree of
female-biased expression in wildtype individuals ($\chi^2 = 53.5$, d.f. = 2, p < 0.0001) (Fig 3A). Four
genes had male-biased expression, and only three were male-biased by at least an order of
magnitude difference. Interestingly, 48 of the downregulated genes (80%) were located on the X
chromosome, of which 40 were female-biased (Fig 3A). There is a negative relationship between
sex-biased expression and differential gene expression on the X chromosome, such that more
female-biased genes had a greater fold decrease in expression (t = -2.71, d.f. = 46, p < 0.01). This
relationship suggests a pleiotropic effect in a *trans*-regulatory pathway impacting regulation of
X-linked genes. No relationship exists between autosomal genes and differential gene expression
(p = 0.26), though the low number of autosomal genes likely makes this comparison under-
powered. Together, these data indicate a skew towards a less female-like transcriptome in
evolved females.
One major *trans*-regulatory pathway that could explain the correlation between
downregulated genes and the X chromosome is dosage compensation. In *Caenorhabditis*
nematodes, hermaphrodites/females have two copies of the X chromosome, each of which are
partially downregulated to match the dosage of the single X chromosome in males (Meyer,
2022). To determine if whole-scale changes in dosage compensation are responsible for the
downregulated pattern in evolved females, we correlated differential gene expression with gene
expression levels for dosage compensated and non-dosage compensated X-linked genes ($\chi^2 =$
4.97, d.f. = 1, p = 0.03) (Jans *et al.*, 2009). Only 23 (45%) of the significant genes located on the
X chromosome were present in both data sets (Fig. 3B). Of these, seven were dosage
compensated in wildtype hermaphrodites/females (XX individuals). These genes were further
downregulated in evolved females, potentially suggesting an expression level lower than males.
Sixteen genes were not dosage compensated in wildtype hermaphrodites/females, but were
downregulated in evolved females by on average 1.5-fold. Similarly, the mean fold decrease in
wildtype expression in males (XO individuals) for these same genes is 1.8-fold. The core dosage
compensation genes *sdc-2* (log$_2$Fold Change = 0.02; p$_{adj}$ = 0.9), *sdc-3* (log$_2$Fold Change = -0.15;
p$_{adj}$ = 0.9), *dpy-27* (log$_2$Fold Change = -0.07; p$_{adj}$ = 1.0), and *dpy-30* (log$_2$Fold Change = 0.04;
p$_{adj}$ = 1.0) did not have significant changes in gene expression after experimental evolution nor
were there significant allele frequency changes at or near these genes (Kasimatis *et al.*, 2022).
Additionally, there were no significant allele frequency changes at the *rex* or *dox* binding sites,

indicating that the dosage compensation machinery itself is likely not the cause of widespread

downregulation in evolved females. Therefore, alternative pathways must be involved in this

masculinization of the female transcriptome resulting from direct sexual selection acting on

males.

323   **DISCUSSION**

Sexual selection acts in a sex-specific manner yet can have an indirect evolutionary response in

the other sex due to genetic associations created by their sharing the same genome (apart from

heterogametic sex chromosomes when present). When sexual selection acts on one sex, the other

sex can be pulled away from their sex-specific optimum as a correlated response to selection,

generating conflict between the sexes. Here we explored whether direct sexual selection on male

sperm competitive ability (Kasimatis *et al.*, 2022) generated indirect selection in females by

quantifying the ancestral and evolved transcriptomes of females mated with different male

generations.

The factorial mating design used here allowed for changes in gene expression to be

categorized as a plastic response to increased sperm competitiveness in males or an evolved

response in the females themselves. If females were predisposed to increased sperm interactions,

then male generation should have been a major contributor to gene expression changes. Such a

scenario is analogous to a pre-existing bias model of sexual selection (Ryan *et al.*, 1990) and can

lead to direct selection on female preference/resistance (Gavrilets *et al.*, 2001; Arnqvist & Rowe,

2005). However, we found no evidence that male evolutionary history impacted differential gene

expression. Rather, all expression changes were determined based on whether a female was from

the ancestral or evolved populations, indicating that differential gene expression was a result of

indirect selection on females during experimental evolution. This scenario is analogous to

Fisherian sexual selection where direct selection on male traits can generate an indirect response

on female preference for those traits if an additive genetic covariance exists (Lande, 1981). For

our post-insemination model, selection was directed on sperm competitive ability. The correlated

female trait is not immediately clear based on GO analysis. However, membrane-bounded

organelle, vesicle-mediated transport, and lipid metabolism may be related to oogenesis, while

DNA binding may be related to transcriptional activity. Since selection on males generated a

polygenic response and the expression changes observed were likely due to *trans*-acting

pathways, it is possible that more than one female reproductive trait may be involved.

Interestingly, there was a complete lack of overlap between the genic changes identified

in Kasimatis *et al.* (2022) and the expression changes characterized here. One potential reason

for this discrepancy is that the genic changes were altering expression of upstream transcription

factors that are expressed earlier in development. Kasimatis *et al.* (2022) identified genomic

changes in the coding sequence of or nearby intergenic region of four transcription factors. Of

particular note is the transcription factor *sea-1*, which is involved in sex determination and

dosage compensation through hypoactivation of the X chromosome (Harris *et al.*, 2019).

Genomic analysis identified seven SNPs with a significant allele frequency change located

within a 30 base pair region of intron 2 (Kasimatis *et al.*, 2022). It is possible that this region

changes regulation of *sea-1* itself or alters splicing in a way that increase the hypoactivation

function of *sea-1* and thus is responsible for the widespread downregulation of X-linked genes.

Genes with significant expression changes between ancestral and evolved females were

almost exclusively downregulated. Moreover, most of these genes are female-biased in wildtype

individuals. These results suggest that the evolved female transcriptome moved away from a

female-like state, supporting indirect rather than direct selection on females. The majority of

downregulated genes were located on the X chromosome. Because males are the heterogametic

sex in *C. elegans*, alleles affecting gene expression on the X chromosome are directly exposed to

selection without any potential shielding of an alternative allele due to dominance, as would

occur in heterozygous females. Thus, selection can be more effective on X-bearing genes,

especially when there is sex-specific selection (Charlesworth *et al.*, 1987; Mank, 2009; Vicoso &

Charlesworth, 2009). The response we observe here is thus consistent with the dominance theory

of the rapid evolution of sex chromosomes and their role in speciation (Charlesworth *et al.*,

1987; Charlesworth, 1991).

Despite the preponderance of signal on the X chromosome, we found no evidence that

mutations in the core dosage compensation pathway were the cause of this widespread

downregulation. Downregulation of dosage compensated genes suggests potentially lower

expression in evolved females relative to evolved males. In contrast, downregulation of non-

dosage compensated genes moved the evolved female expression level closer to the wildtype

male expression level. These changes further support a less female-like and more male-like

379 transcriptome. Since expression in males was not quantified, we cannot definitively say that

380 these genes were not equally downregulated in both sexes to maintain the sex bias and dosage

381 ratios. However, under such a scenario male generation should have been a significant predictor

382 of gene expression changes, which was not observed. Overall, the results indicate an indirect

383 masculinization of the female transcriptome generated by direct selection on male fertilization

384 success. Since gene expression levels impact trait function and, in turn, fitness, short-term sexual

385 conflict was likely generated as a byproduct of direct sexual selection in males as female

386 expression was pushed off its global optimum.

387   Previous experimental evolution studies suggest that male-limited evolution over tens of

388 generations can decrease female fitness (Rice, 1996; Prasad *et al.*, 2007; Pischedda &

389 Chippindale, 2017). Conversely, female-limited or reduced sexual selection evolution tends to

390 increase female fitness with little to no effect on male fitness (Holland & Rice, 1999; Pitnick *et*

391 *al.*, 2001a, 2001b). These previous studies therefore suggest that sex-specific selection acting

392 through males may have a higher potential to generate sexual conflict. However, this previous

393 work has been blind to the traits under selection and therefore the genetic architecture underlying

394 the sexual conflict response. Our work to bridge these layers of the genomic, transcriptomic, and

395 fitness responses suggests a Red Queen scenario (Valen, 1973; Brockhurst *et al.*, 2014) relating

396 genome evolution and transcriptional dimorphism during post-insemination interactions.

397 Specifically, on microevolutionary timescales strong sexual selection in males pushes females

398 away from their fitness optimum through indirect selection (Rice & Holland, 1997; Rice &

399 Chippindale, 2001). Females may then respond through direct selection, which creates a

400 macroevolutionary picture of a stable female-like state. Thus, examining the functional layers

401 through which sexual dimorphism arises is critical for understanding the context in which sexual

402 conflict can occur.

403   Sexual selection drives the evolution of some of the most elaborate phenotypic variation

404 and complex behaviors. We show that sexual selection is just as powerful an evolutionary force

405 on post-insemination traits. In particular, on microevolution timescales, females may not have

406 the opportunity to counter the effects of indirect selection leading to females being pulled away

407 from their optimal state. Given the seemingly high degree of pleiotropy between female and

408 male reproductive networks, sexual conflict may evolve very rapidly in populations experiencing

409 strong sexual selection.

Acknowledgements: We thank Ruben Lancaster and Alex Smith for experimental assistance and Anastasia Teterina for transcriptome analysis advice. We thank Locke Rowe and Stephen Wright for their insightful discussion.

Funding: This research was supported by NIH grant R35GM131838 to PCP and a Natural Sciences and Engineering Research Council of Canada (NSERC) Banting Postdoctoral Fellowship to KRK.

Author Contributions: KRK and PCP devised the project. KRK performed the crosses. JHW prepared the transcriptomic libraries. KRK analyzed the data. KRK wrote the manuscript with the support of the other authors.

Data Accessibility: Raw transcriptomic reads will be made available at NCBI prior to publication. Differential expression analysis results for all data are in File S1. Analysis of significant genes are in File S2. Gene ontology enrichment results are available in File S3. Cross-referencing of *cis*-regulatory SNPs are available in File S4. All scripts are publicly available on the GitHub repository: https://github.com/katjakasimatis/expevol_femaletranscriptomics. The experimental evolution ancestral strain (PX632) and evolved replicates are available from the Phillips lab upon request.

References

Albritton, S.E., Kranz, A.-L., Rao, P., Kramer, M., Dieterich, C. & Ercan, S. (2014) Sex-biased

gene expression and evolution of the x chromosome in nematodes. *Genetics*, **197**, 865–883.

Arnqvist, G. & Rowe, L. (2005) *Sexual conflict*. Princeton University Press. Princeton, NY.

Begun, D.J., Whitley, P., Todd, B.L., Waldrip-Dail, H.M. & Clark, A.G. (2000) Molecular

population genetics of male accessory gland proteins in *Drosophila*. *Genetics*, **156**, 1879–

1888.

Benjamini, Y. & Hochberg, Y. (1995) Controlling the false discovery rate: a practical and

powerful approach to multiple Testing. *Journal of the Royal Statistical Society: Series B*

(Methodological), **57**, 289–300.

Brenner, S. (1974) The genetics of *Caenorhabditis elegans*. *Genetics*, **77**, 71–94.

Brockhurst, M.A., Chapman, T., King, K.C., Mank, J.E., Paterson, S. & Hurst, G.D.D. (2014)

Running with the Red Queen: the role of biotic conflicts in evolution. *Proceedings of the*

Royal Society B: Biological Sciences, **281**, 20141382.

Chapman, T. (2011) Seminal fluid-mediated fitness traits in *Drosophila*. *Heredity*, **87**, 511–521.

Charlesworth, B. (1991) The evolution of sex chromosomes. *Science*, **251**, 1030–1033.

Charlesworth, B., Coyne, J.A. & Barton, N.H. (1987) The relative rates of evolution of sex

chromosomes and autosomes. *American Naturalist*, **130**, 113–146.

Clark, N.L., Aagaard, J.E. & Swanson, W.J. (2006) Evolution of reproductive proteins from

animals and plants. *Reproduction*, **131**, 11–22.

Consortium, U. (2021) UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids*

Research, **49**, D480–D489.

Cooper, W. & Forshaw, J. (1977) *The Birds of Paradise and Bower Birds*. Collins, Sydney.

Darwin, C. (1871) *The Descent of Man and Selection in Relation to Sex*. Murray. Murray.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., *et al.* (2013) STAR:

ultrafast universal RNA-seq aligner. *Bioinformatics*, **29**, 15–21.

Fisher, R.A. (1930) *The Genetical Theory of Natural Selection*. Oxford University Press. Oxford,

UK.

Gavrilets, S., Arnqvist, G. & Friberg, U. (2001) The evolution of female mate choice by sexual

conflict. *Proceedings of the Royal Society B: Biological Sciences*, **268**, 531–539.

Grath, S. & Parsch, J. (2016) Sex-biased gene expression. *Annual review of genetics*, **50**, 29–44.

Harris, T.W., Arnaboldi, V., Cain, S., Chan, J., Chen, W.J., Cho, J., *et al.* (2019) WormBase: a
modern model organism information resource. *Nucleic Acids Research*, **29**, 82.

Holland, B. & Rice, W.R. (1999) Experimental removal of sexual selection reverses intersexual
antagonistic coevolution and removes a reproductive load. *Proceedings of the National*
Academy of Sciences, **96**, 5083–5088.

Huntley, R.P., Sawford, T., Mutowo-Meullenet, P., Shypitsyna, A., Bonilla, C., Martin, M.J., *et*
al. (2015) The GOA database: gene ontology annotation updates for 2015. *Nucleic Acids*
Research, **43**, D1057-63.

Jans, J., Gladden, J.M., Ralston, E.J., Pickle, C.S., Michel, A.H., Pferdehirt, R.R., *et al.* (2009) A
condensin-like dosage compensation complex acts at a distance to control expression
throughout the genome. *Genes & Development*, **23**, 602–618.

Jiang, H., Lei, R., Ding, S.-W. & Zhu, S. (2014) Skewer: a fast and accurate adapter trimmer for
next-generation sequencing paired-end reads. *BMC bioinformatics*, **15**, 182.

Kasimatis, K.R., Moerdyk-Schauwecker, M.J., Lancaster, R., Smith, A., Willis, J.H. & Phillips,
P.C. (2022) Post-insemination selection dominates pre-insemination selection in driving
rapid evolution of male competitive ability. *PLoS Genetics*, **18**, e1010063.

Kasimatis, K.R., Nelson, T.C. & Phillips, P.C. (2017) Genomic signatures of sexual conflict. *The*
Journal of heredity, **108**, 780–790.

Kenyon, C., Chang, J., Gensch, E., Rudner, A. & Tabtiang, R. (1993) A *C. elegans* mutant that
lives twice as long as wild type. *Nature*, **366**, 461–464.

Kirkpatrick, M. (1982) Sexual selection and the evolution of female choice. *Evolution*, **36**, 1.

Lande, R. (1980) Sexual dimorphism, sexual selection, and adaptation in polygenic characters.
Evolution, **34**, 292–305.

Lande, R. (1981) Models of speciation by sexual selection on polygenic traits. *Proceedings of*
the National Academy of Sciences, **78**, 3721–3725.

Lande, R. & Arnold, S.J. (1983) The measurement of selection on correlated characters.
Evolution, **37**, 1210–1226.

Lande, R. & Arnold, S.J. (1985) Evolution of mating preference and sexual dimorphism. *Journal*
of Theoretical Biology, **117**, 651–664.

Love, M.I., Huber, W. & Anders, S. (2014) Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biology*, **15**, 31–21.

Mank, J.E. (2009) Sex chromosomes and the evolution of sexual dimorphism: lessons from the
genome. *The American Naturalist*, **173**, 141–150.

Mank, J.E. (2017) The transcriptional architecture of phenotypic dimorphism. *Nature Reviews*
Genetics, **1**, 1–7.

Meyer, B.J. (2022) The X chromosome in C. elegans sex determination and dosage
compensation. *Current Opinion in Genetics & Development*, **74**, 101912.

Pischedda, A. & Chippindale, A.K. (2017) Direct benefits of choosing a high fitness mate can
offset the indirect costs associated with intralocus sexual conflict. *Evolution*, 1–27.

Pitnick, S., Brown, W.D. & Miller, G.T. (2001a) Evolution of female remating behaviour
following experimental removal of sexual selection. *Proceedings of the Royal Society B:*
Biological Sciences, **268**, 557–563.

Pitnick, S., Miller, G.T., Reagan, J. & Holland, B. (2001b) Males' evolutionary responses to
experimental removal of sexual selection. *Proceedings of the Royal Society of London. Series*
B: Biological Sciences, **268**, 1071–1080.

Prasad, N.G., Bedhomme, S., Day, T. & Chippindale, A.K. (2007) An evolutionary cost of
separate genders revealed by male-limited evolution. *The American Naturalist*, **169**, 29–37.

Putri, G.H., Anders, S., Pyl, P.T., Pimanda, J.E. & Zanini, F. (2022) Analysing high-throughput
sequencing data in Python with HTSeq 2.0. *Bioinformatics*, **38**, 2943–2945.

R Core Team. (2020) *R: A language and environment for statistical computing*. Foundation for
Statistical Computing. Foundation for Statistical Computing.

Rice, W.R. (1996) Sexually antagonistic male adaptation triggered by experimental arrest of
female evolution. *Nature*, **381**, 232–234.

Rice, W.R. & Chippindale, A.K. (2001) Intersexual ontogenetic conflict. *Journal of Evolutionary*
Biology, **14**, 685–693.

Rice, W.R. & Holland, B. (1997) The enemies within: intergenomic conflict, interlocus contest
evolution (ICE), and the intraspecific Red Queen. *Behavioral Ecology and Sociobiology*, **41**,
1–10.

Ryan, M.J., Fox, J.H., Wilczynski, W. & Rand, A.S. (1990) Sexual selection for sensory
exploitation in the frog Physalaemus pustulosus. *Nature*, **343**, 66–67.

Swanson, W.J. & Vacquier, V.D. (2002) Reproductive protein evolution. *Annual Review of*
Ecology and Systematics, **33**, 161–179.

Team, B. & Maintainer, B. (2019) TxDb.Celegans.UCSC.ce11.ensGene: Annotation package for

TxDb object(s). *R package version 3.4.6.*

Turelli, M. (1985) Effects of pleiotropy on predictions concerning mutation-selection balance for

polygenic traits. *Genetics*, **111**, 165–195.

Valen, L.V. (1973) A new evolutionary law. *Evolutionary theory*, **1**, 1–30.

Vicoso, B. & Charlesworth, B. (2009) Effective population size and the faster-X effect: an

extended model. *Evolution*, **63**, 2413–2426.

Wyman, M.J., Stinchcombe, J.R. & Rowe, L. (2013) A multivariate view of the evolution of

sexual dimorphism. *Journal of Evolutionary Biology*, **26**, 2070–2080.

Young, M.D., Wakefield, M.J., Smyth, G.K. & Oshlack, A. (2010) Gene ontology analysis for

RNA-seq: accounting for selection bias. *Genome Biology*, **11**, R14.

Figure 1. Differential gene expression is explained by female generation. **A)** Principal

component analysis of the variance stabilized dataset partitions female generation along PC1. **B)**

Differential gene expression was analyzed based on female generation, male generation, and the

interaction between female and male generations. Only the female contrast contributed to gene

expression changes. Genes with a significant change in expression from ancestral to evolved

females (shown in pink) were determined using a Bonferroni cut-off. Positive expression

changes correspond to upregulation in evolved females, while negative expression changes

correspond to downregulation in evolved females.

Figure 2. A heatmap of gene expression levels in significant differentially expressed genes.

Ancestral females (light purple) and evolved females (dark purple) group together, while

ancestral (light green) and evolved (dark green) males are interspersed. Sixty-one of the 64

significant genes are downregulated in evolved females. Pseudogenes are labeled in gray.

Figure 3. Downregulated genes are female-biased. **A)** There is a negative correlation between

significant differentially expressed genes located on the X chromosome (shown in coral) and

sex-biased expression. Only 13 genes were located on autosomes (shown in blue). Both sex-

biased expression (female to male) and experimental evolution fold change are plotted on a $\log_2$

scale. **B)** The relationship between normalized expression of dosage compensated (circles) and

not dosage compensated (triangles) X-linked genes in wildtype hermaphrodites/females (XX

individuals) versus differential gene expression. Genes with significant differential gene

expression are shown in pink. The dashed line represents the mean fold decrease of non-dosage

compensated genes in wildtype males (XO individuals). Both axes are plotted on a $\log_2$ scale.

Figure S1. Principal component analysis of the variance stabilized data for each experimental

evolution replicate partitions female generation along PC1. Experimental evolution replicate B3

had two outlier transcriptome replicates (circled) and were censored from the final, pooled

dataset.