1

Cell-type-specific plasticity shapes neocortical dynamics for motor learning

Shouvik Majumder[1], Koichi Hirokawa[1], Zidan Yang[1], Charles R. Gerfen[2], Lorenzo Fontolan[3,4],

Sandro Romani[4], Anant Jain[1], Ryohei Yasuda[1], and Hidehiko K. Inagaki[1]

9   [1] Max Planck Florida Institute for Neuroscience, Jupiter, FL 33458, USA.

10   [2] National Institute of Mental Health, Bethesda, MD 20814, USA.

11   [3]Turing Centre for Living Systems, Aix- Marseille University, INSERM, INMED U1249, Marseille,

12   France

13   [4] Janelia Research Campus, HHMI, Ashburn VA 20147, USA.

Abstract:	197 words	
Main text:	3739 words	
Figures:		4
Extended Data Figures:		10
Extended Data Table:		4

Correspondence:

Hidehiko K. Inagaki

Hidehiko.inagaki@mpfi.org

Abstract

Neocortical spiking dynamics control aspects of behavior, yet how these dynamics emerge during motor learning remains elusive. Activity-dependent synaptic plasticity is likely a key mechanism, as it reconfigures network architectures that govern neural dynamics. Here, we examined how the mouse premotor cortex acquires its well-characterized neural dynamics that control movement timing, specifically lick timing. To probe the role of synaptic plasticity, we have genetically manipulated proteins essential for major forms of synaptic plasticity, Ca^{2+}/calmodulin-dependent protein kinase II (CaMKII) and Cofilin, in a region and cell-type-specific manner. Transient inactivation of CaMKII in the premotor cortex blocked learning of new lick timing without affecting the execution of learned action or ongoing spiking activity. Furthermore, among the major glutamatergic neurons in the premotor cortex, CaMKII and Cofilin activity in pyramidal tract (PT) neurons, but not intratelencephalic (IT) neurons, is necessary for learning. High-density electrophysiology in the premotor cortex uncovered that neural dynamics anticipating licks are progressively shaped during learning, which explains the change in lick timing. Such reconfiguration in behaviorally relevant dynamics is impeded by CaMKII manipulation in PT neurons. Altogether, the activity of plasticity-related proteins in PT neurons plays a central role in sculpting neocortical dynamics to learn new behavior.

Introduction

Neural computations are mediated by time-varying and coordinated spiking activity across a population of neurons, referred to as neural dynamics. For example, during the planning of volitional movement, the premotor cortex exhibits slowly varying neural activity that determines the type and timing of upcoming movement, referred to as preparatory activity[1,2]. Previous research has shown flexible reconfiguration of dynamics, including preparatory activity, during motor learning[3-16]. But the neural mechanisms that reshape the neocortical dynamics to enable acquisitions of new behavior remain elusive.

Network architectures, i.e., synaptic connections, constrain neural dynamics[17]. Therefore, altering specific connections in the network architecture through synaptic plasticity is the primary theory for learning[12,14,18]. First, experience- and learning-dependent synaptic plasticity has been observed across many brain areas[7,19-27]. Second, essentially all neurons are equipped with molecular pathways that mediate plasticity[28-32]. Third, manipulations of synaptic plasticity influence learning, although this has primarily been investigated in the hippocampus and cerebellum and far less in the neocortex[11,33-40].

The neocortex contains diverse neuronal cell types across layers, characterized by unique gene expression profiles and anatomical features[41-43]. These cell types often carry distinct information and contribute to different aspects of neural dynamics and behavior in expert animals performing tasks[44-47]. The function of synaptic plasticity during learning may also vary across neocortical cell types to shape these cell-type-dependent dynamics and behavior.

To probe the role of synaptic plasticity across cell types during motor learning, we performed a series of acute genetic manipulations of proteins required for synaptic plasticity. We studied learning of motor timing in mice, focusing on the premotor cortex, as it provides several advantages. First, neocortical dynamics in the premotor cortex and their causal roles on behaviors, including timing behavior, have been well-

established in expert animals[1,2,48–52]. In addition, the premotor cortex has been implicated in learning across

tasks and species[10,53–55], making it an ideal site to examine the function of plasticity in reconfiguring

dynamics during learning. Finally, animals are adept at learning motor timing, making it a quick and robust

system to study learning. Leveraging this model system with molecular manipulations and high-density

electrophysiology, we identified the key cell type required to shape behaviorally relevant neocortical

dynamics and timing of action.

Results

CaMKII activity in ALM is necessary to learn new lick timing

We developed an operant motor timing task, in which water-restricted mice learn to delay their lick time in

order to acquire water rewards (Fig.1a and Methods). Trial onset was signaled by an auditory cue (3 kHz

tone, 0.6 s), and a lick after a following unsignaled delay epoch was rewarded (rewarded trials). An early

lick during the delay epoch aborted the trial without a reward. Training proceeded in two stages. First, mice

were trained to lick after the cue onset with a minimal delay (0.1 s, 'cue association'). Second, the delay

duration was gradually increased ('delay training'; reaching criterion performance, 30% rewarded trials in

the last 100 trials with a given delay duration, resulted in a delay increase of 0.1 s; Methods). Following

this protocol, mice learned to delay lick timing within and across sessions (Fig. 1b; lick time reached 1.20

± 0.31 s, mean ± SEM, in 6 days of delay training, n = 6 mice). High-speed videography (300 Hz) revealed

that animals moved their jaw and tongue before the full tongue protrusion (Extended Data Fig. 1a-f). Yet,

the delayed lick time after training was primarily due to the withholding of orofacial movement (Extended

Data Fig. 1a-f).

Next, we examined whether synaptic plasticity in the anterior-lateral motor cortex (ALM; AP 2.5mm ML

1.5 mm from Bregma), a premotor cortical area responsible for orofacial movement[5,56], is required for this

Figure 1. CaMKII activity in ALM is required for delay learning.

a. Task design.

b. Example learning curve. Distributions of lick time per trial bin are shown (50 trials; Methods). Vertical dotted lines separate sessions.

c. paAIP2 manipulation during learning. The blue light was on during the delay training but not cue association. Day 0, the last day of cue association. The median lick times of the last 100 trials are shown. Thick lines, mean ± SEM. Thin lines, individual mice (n > 5 for 'ALM' and 'No' per condition, and n = 4 for 'M1'). *: $p < 0.05$ (bootstrap followed by *Bonferroni* correction). See Extended Data Table 3 for the exact n and *p*-values. Right, example learning curves of mice with paAIP2 expression in ALM (top, light off; bottom, light on). The format is the same as in b.

d. Distribution of first lick time on the second day of cue association. 'Cue' and 'No cue', trials with and without a cue, respectively (Methods; cue starts at the trial onset in cue trials). Different distributions between cue and no cue trials indicate successful cue association. Shade, SEM. $p_{cue\ with\ vs.\ w/o\ light} = 0.571$ (ranksum test, n = 3 and 5 mice, light on and off respectively).

e. ALM paAIP2 manipulation in expert animals. Comparing the first lick time within a session (100 trials before and after the onset of blue light). Thick lines, mean ± SEM. Thin lines, individual sessions (n = 4 mice, 20 sessions). $p = 0.117$ (bootstrap). See Methods for the shuffle procedure.

f. Knocking out CaMKIIα expression in ALM. Thick lines, mean ± SEM. Thin lines, individual mice (n = 6 mice per condition). *: $p < 0.05$ (bootstrap followed by *Bonferroni* correction).

learning. To this end, we blocked the activity of CaMKII, a Ca^{2+}-dependent kinase required to induce major

types of synaptic plasticity and learning[27,29,30,32,33]. A transient manipulation of CaMKII activity using a

genetically-encoded photoactivatable competitive inhibitor of CaMKII, paAIP2, blocks the induction of

synaptic plasticity[39] in the presence of blue light without influencing the excitability of neurons (Extended

Data Fig. 2a-f).

Control mice without paAIP2 expression learned new lick timing regardless of blue light illumination of

ALM (470nm, 0.5mW, 0.2 Hz; Fig. 1c, first column; Methods). In contrast, AAV-mediated bilateral

paAIP2 expression in ALM blocked learning when blue light was illuminated during the delay training, but

not in the absence of blue light (Fig. 1c, second column; Extended Data Table 3; Extended Data Table 1

and 2). Notably, during light illumination, mice continued licking after the cue without a change in the

variability of lick time (Fig.1c and Extended Data Fig 1g-j). This implies that paAIP2 manipulation does

not block animals' ability to explore different lick times across trials (which is a prerequisite for

reinforcement learning[57]). Instead, the manipulation interferes with the ability to directionally shift the

distribution of lick times, presumably guided by rewards. Similar manipulation in M1 (AP 0.0mm ML 1.5

mm from Bregma) did not block learning (Fig. 1c, third column), implying that CaMKII activity in ALM

is required for delay learning.

The ALM paAIP2 manipulation did not interfere with the cue association (Fig. 1d). In expert mice trained

for two weeks with a fixed delay at 1.5 s (Methods), paAIP2 manipulation did not alter the distribution of

lick timing, implying that CaMKII activity in ALM is not required for mice to execute learned delayed licks

(Fig. 1e). Electrophysiological recording of ALM in expert mice confirmed that paAIP2 manipulation does

not directly perturb ongoing spiking dynamics during behavior (Extended Data Fig. 2g-o). Altogether,

CaMKII activity in ALM is specifically required for learning new lick timing but not for the initial cue

association, the execution of learned action, or ongoing spiking activity.

CaMKIIα isoform is required for synaptic plasticity and learning, while other CaMKII isoforms are

implicated in different cellular functions[30,58]. Since paAIP2 binds to the kinase domain homologous across

the CaMKII family, paAIP2 likely blocks all of them. To test whether CaMKIIα in ALM is required for

learning, we acutely knocked it out in adult ALM by injecting AAV-hsyn-cre in CaMKIIα conditional

knockout mice[59] (Fig. 1f and Extended Data Fig. 3c). These mice learned cue association, yet could not

learn to delay lick timing (Fig. 1f), consistent with the paAIP2 manipulation. Altogether, learning new lick

timing requires CaMKIIα activity in ALM.

Synaptic plasticity proteins in PT but not IT cells are required for motor learning

The major excitatory cell types in the premotor cortex include IT and PT neurons, projecting within and

outside the telencephalon, respectively (Fig. 2a). Both of these cell types highly express CaMKIIα in ALM[42]

(Extended Data Fig. 3). The key question is whether synaptic plasticity in these diverse populations has

redundant or specialized roles in learning. We developed an AAV expressing paAIP2 in a cre-dependent

manner to manipulate CaMKII activity in distinct neocortical cell types (Fig. 2b). Strikingly, paAIP2

manipulation of PT neurons in ALM completely blocked delay learning without affecting the distribution

of cue-triggered lick, similar to the bulk manipulation across cell types (Fig. 2b and c; Extended Data Fig.1).

We observed consistent behavioral effects whether we labeled PT neurons using Sim1-cre KJ18 transgenic

line[60] or manipulated individual PT subtypes using retrograde AAV (PT_{upper} and PT_{lower} neurons with

distinct subcortical projection[46]; 2700 ± 208 manipulated cells/animal, mean $\pm$ SEM; Fig.2c and Extended

Data Fig. 4). In contrast, paAIP2 manipulations in IT neurons (layer 2/3 and 5 IT neurons) did not block

learning (Fig. 2c).

To test the necessity of the CaMKIIα isoform across cell types, we acutely knocked out CaMKIIα using

CRISPR/Cas9. We injected AAV expressing guide RNA against CaMKIIα[30] in ALM of adult mice

Figure 2. CaMKII activity in PT neurons is required for delay learning.

a. Network architecture and cell types in ALM.

b. Expression of paAIP2 in ALM PT_{upper} neurons.

c. The effect of paAIP2 manipulation in distinct cell types in ALM. 'No' and 'ALM' are duplicated from Figure 1c for comparison purposes. Thick lines, mean ± SEM. Thin lines, individual mice (n > 3 mice per condition). *: $p < 0.05$ (bootstrap followed by *Bonferroni* correction; Extended Data Table 3).

d. Top, cell-type-specific KO of CaMKIIα using CRISPR/Cas9. Bottom, immunohistochemical validation of loss of CaMKIIα protein expression in PT_{upper} neurons.

e. The effect of CaMKIIα KO in PT_{upper} and IT neurons. Thick lines, mean ± SEM. Thin lines, individual mice (n > 4 mice per condition). *: $p < 0.05$ (bootstrap followed by *Bonferroni* correction; Extended Data Table 3)

f. The effect of inactivating Cofilin in PT_{upper} and IT neurons. Thick lines, mean ± SEM. Thin lines, individual mice (n > 4 per condition). *: $p < 0.05$ for both Cofilin-SN light on vs. off comparison and Cofilin-SN light on vs. unconjugated light on conditions (bootstrap followed by *Bonferroni* correction; Extended Data Table 3).

expressing Cas9[61] in PT_{upper} or IT neurons (Methods). After a month of AAV injection, a significant

proportion of cells that co-express Cas9 and guide RNA lost CaMKIIα protein expression (Fig. 2d and

Extended Data Fig. 3). PT_{upper}-specific knockout of CaMKIIα blocked delay learning without any effect on

cue association, whereas IT-specific knockout did not affect learning, consistent with the paAIP2

manipulations (Fig. 2e).

As an independent approach to manipulate synaptic plasticity, we inactivated Cofilin, a protein required for

actin remodeling during major forms of synaptic plasticity[40]. To this end, we leveraged Cofilin conjugated

with SuperNova (Cofilin-SuperNova), a monomer photosensitizing fluorescent protein for Chromophore

Assisted Light Inactivation (CALI)[40]. PT_{upper}-specific expression of Cofilin-SuperNova and orange light

illumination (595 nm, 0.75 mW, 1 min every 10 min) during the delay training blocked learning without

affecting cue-triggered licks (Fig. 2f). In contrast, no illumination or unconjugated SuperNova with

illumination did not affect learning, implying that CALI-mediated inactivation of Cofilin blocked the

learning. Consistent with the CaMKII manipulation, manipulating Cofilin in IT neurons did not affect

learning. Altogether, three cell-type specific acute genetic manipulations imply that synaptic plasticity in

PT but not IT neurons is necessary to learn new lick timing.

Evolution of ALM preparatory dynamics during delay learning

Since preparatory activity determines the upcoming actions[2], ALM preparatory dynamics are likely tailored

during learning new lick timing. To characterize preparatory activity in our behavioral task, we performed

acute high-density extracellular electrophysiological recordings in ALM during learning and in expert mice

(Fig. 3 and Extended Data Fig. 5; with rigorous quality control in spike sorting, Extended Data Fig. 6).

Among the 3203 putative pyramidal neurons we recorded from 30 mice, we focused on 1613 neurons with

preparatory activity (defined as neurons with significant activity between cue and lick, signed-rank test, p

< 0.05; Methods; Extended Data Fig. 7 and Table. 3).

Figure 3. Evolution of ALM dynamics during delay learning.

a. Spiking activity of example neurons in expert mice. Lines with different colors indicate the mean spike rate of trials with different first lick times (vertical dotted lines in the same color indicate the corresponding lick time). Top, activity in absolute time. Bottom, activity in relative time, following temporal warping (all trials were warped to have identical lick time; Methods).

b. Grand average peri-stimulus time histogram (PSTH) of positively-modulated ALM preparatory neurons (cells with the mean spike rate between cue and lick significantly higher than the baseline). The same format as in **a**. Shade, SEM. Trial types with more than 50 neurons are shown. See Extended Data Table 3 for the number of neurons analyzed.

c. Z-scored spiking activity of preparatory ALM neurons in trials with first lick time within 0.95-1.1 s (first column) and 1.25-1.4 s (second column). All preparatory neurons with these two trial types are shown (n = 344 cells). The difference between the first and second columns after temporal warping (third column).

d. Pearson's correlation of population activity between two trial types shown in **c**.

e-h. Same as in **a-d** for during learning. **g-h**: N = 269 cells. Note that many cells show a decrease in spiking activity as mice lick later (third column), unlike in the expert mice (**c**), consistent with the population average (**f**).

i. Comparison of cue response between trials with different lick times (calculated based on the grand average PSTH shown in **b** and **f**). *P*-value; hierarchical bootstrap testing the significance of correlation coefficient between lick time vs. cue-response (with a null hypothesis that there is no correlation). Error bar, SEM (hierarchical bootstrap). Left, schema showing time windows used to calculate cue response and pre-lick activity. SR, spike rate. T_{cue}, time from the cue. T_{lick}, time from lick.

j. Same as in **i** for pre-lick activity.

k. The time between 10% to 90% activity is calculated for the grand average PSTH shown in **b** and **f** (Methods). *P*-value; hierarchical bootstrap testing the significance of correlation coefficient between lick time vs. the time between 10-90% activity. Error bar, SEM (hierarchical bootstrap).

First, we leveraged trial-to-trial variability of lick time (CV = 0.41 ± 0.04; mean ± SEM; Extended Data

Fig. 1k) to characterize how ALM preparatory dynamics change as a function of lick time in expert mice.

Many ALM neurons showed ramping spiking dynamics starting at the cue onset and reaching a peak around

the lick onset[2,51] (Fig. 3a, cell 175, and Fig. 3b). When expert mice licked at different timing, the slope of

ramping activity was altered without a change in the peak activity level (Fig 3a and b; trials with different

lick times are shown in different colors). Consistently, temporal warping of spiking dynamics to normalize

the lick time resulted in near-identical activity patterns across trials at both single-cell and population levels

(even for non-ramping up cells, e.g., cell 232 in Fig. 3a; Fig 3b and c and Extended Data Fig. 5; Methods).

Thus, the ALM preparatory dynamics are temporally stretched or compressed to trigger licking at different

times in expert animals, consistent with previous observations in other tasks[50–52,62,63].

Next, we examined whether similar reconfiguration in ALM dynamics underlies learning. At the beginning

of delay training, mice licked early, and ALM neurons showed a transient response to the cue (Fig. 3e and

f and Extended Data Fig.5; blue traces). Since strong ALM activity drives a lick[5,44], this high amplitude

cue response may explain immediate licks following the cue.

As mice licked later during learning, ALM dynamics progressively changed (Fig. 3e-f; from blue to red

traces). First, the spiking activity following the cue decreased ('cue response'; Fig. 3i), which may explain

the loss of immediate licks. In contrast, the activity around the lick onset ('pre-lick activity') stayed high

across trials (Fig. 3j), which may function as a 'threshold' activity level to trigger lick[50]. Second, slow

dynamics (persistent or ramping activity) emerged, and filled the extended temporal gap between the cue

and the delayed lick (Fig. 3k and Extended Data Fig. 7; the time between 10% to 90% activity level

increased). These two modes of reconfigurations continued throughout delay training, converting the

transient cue response into a ramping activity observed in the expert mice (Fig. 3f). Unlike in experts, ALM

dynamics during learning were distinct across lick timings after the temporal warping (Fig. 3e-g).

During learning, Pearson's correlation of ALM population activity is high across time points after the cue between trials with different lick times (Fig. 3h). This implies similar population activity patterns constitute preparatory activity across lick times (while the amplitude of activity changes). Consistently, a large proportion of the ALM preparatory activity is explained along a single dimension (65-80%; Extended Data Fig. 8), implying that learning reconfigures dynamics in a low-dimensional space[64]. In contrast, in the expert mice, population activity patterns changed between cue to lick, and thus the dimensionality is higher (Fig. 3d and Extended Data Fig. 8). Altogether, the reconfiguration of ALM preparatory dynamics during learning is qualitatively different from that in expert animals.

Blocking CaMKII activity in ALM PT neurons impedes the evolution of dynamics

We next asked how the CaMKII manipulation in PT neurons influences the reconfigurations of ALM preparatory dynamics. To this end, we performed extracellular electrophysiological recordings in ALM in conjunction with PT neuron-specific paAIP2 manipulation during delay training (Fig. 4a).

To quantify how ALM dynamics change over training, we analyzed activity chronologically (Extended Data Fig. 9a). In control mice, the dynamics slowed down over three days of delay training, consistent with the delayed lick time (Fig.4b and c, black). In addition, the cue response significantly decreased within a session on days 2 and 3 of delay training (Fig. 4d; note that the cue response reaches the 'floor' as mice lick later, Fig. 3i, which may explain a stronger reduction on day 2). These reconfigurations of preparatory dynamics were significantly attenuated in the animals with PT neuron-specific paAIP2 manipulations (Fig. 4c and d and Extended Data Fig. 9a). Since preparatory activity precedes movement, the impedance in its reconfiguration likely explains the lack of learning during the paAIP2 manipulation.

Figure 4. CaMKII manipulation in ALM PT neurons impedes the evolution of preparatory activity.

a. Schema. Recording of extracellular activity in ALM during paAIP2 manipulation of PT$_{upper}$ (or PT$_{lower}$) neurons.

b. Time to first lick during delay training with recording (different cohorts of mice from those tested without recording in Fig.2c). Lines, mean ± SEM. *: p = 0.476, 0.005, <0.001 for day 1, 2, and 3 of delay training, respectively, comparing control vs. PT-specific paAIP2 manipulation (both PT cell types were pooled for statistics because we did not detect a quantitative difference between these two groups; hierarchical bootstrap with a null hypothesis that the increase in lick time within a session is not larger in control). See Extended Data Table 3 for comparisons of each PT cell type and sample size. Early and late trials, first 75 and last 75 trials in the session.

c. The time between 10% to 90% activity of positively-modulated ALM neurons is compared across manipulation types. Lines, mean ± SEM. *: p = 0.400, 0.095, 0.009 for day 1, 2, and 3, respectively, comparing control vs. PT-specific paAIP2 manipulation (both PT cell types were pooled; hierarchical bootstrap with a null hypothesis that the increase within a session is not larger in control). See Extended Data Table 3 for comparisons of each PT cell type and sample size.

d. The change in cue amplitude of positively-modulated ALM neurons within sessions (late trials – early trials; Δcue response) of. Lines, mean ± SEM. * in red: p = 0.273, 0.006, 0.362 for day 1, 2, and 3, respectively, comparing control vs. PT-specific paAIP2 manipulation (both PT cell types were pooled; hierarchical bootstrap with a null hypothesis that the decrease within a session is not larger in control). * in black: p = 0.039, <0.001, 0.002 on day 1, 2, and 3, respectively, for hierarchical bootstrap with a null hypothesis that Δcue response is non-negative in control mice.

e. Schema of the model (see Methods for details). All synaptic connections are excitatory, except for the ones from GABAergic neurons. "t" in the learning rule, trial.

f. Top, dynamics of ALM neurons across lick times in the model (the mean of IT and PT neurons activity). Different color indicates activity in trials with different lick times. Dotted lines, corresponding lick times. Bottom, schemas of the energy landscape in early and late lick trials (along a long-time constant dimension that captures preparatory activity). Note that the full dynamics (top) is not monotonic due to activity along other directions.

g. The amplitude of cue input (g$_1$), synaptic weight of cue to PT synapse (g$_2$), and lick timing (g$_3$) during learning in the model.

Lick time and ALM cue response varies across trials, even during paAIP2 manipulations (Extended Data

Fig. 7 and 9). In rare trials in which mice happened to lick late during PT neuron-specific paAIP2

manipulation, ALM dynamics were similar to that in the control (Extended Data Fig. 7- 9). This implies

that paAIP2 manipulation in PT neurons does not perturb spontaneous fluctuations in ALM dynamics.

Instead, it is required to directionally reconfigure dynamics for learning.

Potential mechanisms for synaptic plasticity to change lick time

How can synaptic plasticity in PT neurons reconfigure preparatory dynamics and drive learning? We

generated a network model of ALM (Fig. 4e and Extended Data Table 4) constrained by previous findings:

IT neurons form strong connections within the neocortex[65]. PT neurons, in contrast, do not project back to

IT neurons but project to the thalamus and brainstem[65] (for simplicity, we combined PT subtypes in the

network model). The thalamic nuclei receiving PT input project back to ALM, and this thalamocortical

loop maintains the preparatory activity[66,67]. The projection of PT neurons to the brainstem drives a lick

when activity is high[44,46,47].

This model implemented preparatory dynamics as ALM activity transitioning from the baseline activity to

the threshold activity level that triggers a lick[2] (Fig. 4f). At the beginning of a trial, the transient cue input

rapidly pushes the activity (ball) out of the 'baseline' stable point, corresponding to the transient cue

response (thick arrows, Fig. 4f bottom). After this transient excursion, the ball slowly rolls down the energy

landscape until it reaches the threshold activity level to trigger a lick, which explains the slow ramping

dynamics (thin arrows, Fig. 4f bottom). Thus, the amplitude of the cue response influences the following

ramping dynamics (the energy landscape illustrates activity along a single dimension and does not explain

the full dynamics, e.g., non-monotonic transient changes are affected by other dimensions). We varied the

amplitude of the cue input across trials (Fig. 4e and g_1), which changed the preparatory activity, and thus,

induced across-trial variability in lick times as in the data. This allowed exploration of lick times required

for learning[57].

By imposing 1) a simple reward-dependent plasticity rule[68] in the excitatory synapse between cue input and

PT neurons and 2) the delay training protocol we applied to mice, the model reproduced the experimental

observations (Fig. 4e-g). A reward was provided when the network happened to 'lick' later than the delay

due to a fluctuation in cue input. The reward-dependent plasticity reduced the cue response and delayed the

lick time in the following trials. The reduction in cue response was caused by the potentiation of the synapse,

which paradoxically reduced the response due to strong excitatory inputs from thalamus/ALM neurons to

the GABAergic neuron (alternatively, a model with synaptic depression without paradoxical effect

reproduced similar dynamics and learning; Extended Data Fig. 10c-e). Thus, the plasticity allowed the

network to exploit lick time guided by reward, and iterations of this cycle resulted in a gradual and

directional change in preparatory dynamics and lick time (Fig. 4f and g). The model displayed no learning

without the synaptic plasticity in PT neurons, reproducing experimental observations (Fig. 4g, blue). The

same network could reproduce expert dynamics by varying tonic input[69] (Extended Data Fig. 10).

Altogether, the model replicates the key experimental observations in this paper and proposes a potential

mechanistic link among them: how synaptic plasticity in PT neurons shapes neocortical dynamics and

behavior, i.e., decreased ALM cue response (Fig. 4d), generated slow dynamics (Fig. 4c), and delayed lick

time (Fig. 4b).

Discussion

We provided lines of evidence supporting synaptic plasticity in PT neurons as a learning mechanism in the

motor timing task. First, we performed three types of acute cell-type-specific genetic manipulations

targeting proteins required for synaptic plasticity, all of which blocked delay learning when PT but not IT

neurons were manipulated. Acute manipulations are unlikely to recruit developmental effects or

compensatory mechanisms[70]. Importantly, these manipulations did not affect the execution of learned

behavior, implying that the effect is specific to learning[36,71]. Second, we performed a series of

electrophysiological recordings *in* and *ex vivo* (Extended Data Fig. 2), confirming that transient paAIP2

manipulation does not affect the excitability of neurons or ongoing spiking activity. Instead, the paAIP2

manipulation impedes the directional reconfiguration of ALM dynamics, leading to rewarded lick, without

perturbing trial-to-trial fluctuations in dynamics and behavior (Fig. 4 and Extended Data Fig.7-9). This is

consistent with a view that CaMKII activity in PT neurons consolidates dynamics followed by a reward,

presumably via synaptic plasticity of inputs that led to successful actions (Fig 4, model).

Synaptic plasticity has been observed across nearly all brain areas and cell types, including putative IT

neurons[12,19,21], seemingly suggesting global and redundant learning mechanisms. Thus, it was unexpected

for a subset of cell types in one cortical area to be necessary for learning.

PT neurons occupy a unique position in the neocortex: they integrate the cortical input and control the

information going through the thalamocortical loop that mediates preparatory activity[46,66]. Plasticity in PT

neurons is, thus, well suited to reconfigure preparatory activity. Interestingly, putative PT neurons

(complex-tuft cells) show active structural synaptic plasticity across areas and conditions[19,20,72]. In addition,

localized synaptic plasticity, especially at the output nodes of a network like PT neurons, is theoretically

beneficial[73,74]. Thus, PT neurons could be a universal learning site across neocortical areas and behavioral

tasks. PT neurons encode behavior more reliably than other cortical cell types[45], which could result from

behaviorally relevant synaptic plasticity in these neurons.

Nonetheless, the necessity of one cell type does not exclude the necessity of other cell types and brain areas.

Indeed, we discovered that at least two PT subtypes are required for learning. Neural dynamics are

orchestrated across brain areas, including the striatum and cerebellum. Diverse cell types across these areas

may implement different aspects of motor learning[23,71,75–77]. Molecular pathways underlying synaptic

plasticity could also be diverse, and we do not rule out plasticity without CaMKII and Cofilin. Leveraging

genetic manipulations targeting various molecules underlying synaptic plasticity, it is tempting to test how

multiregional cell types coordinate diverse types of learning.

A common approach in neuroscience is to manipulate spiking activity and test its effect on behavior. Yet,

spiking activity inevitably propagates to connected brain areas and often interferes with movement. For

instance, silencing spikes in ALM alters spiking activity across motor-related brain areas and blocks

licks[47,78], which complicates the interpretation of behavioral effects even if it blocks learning[79]. The paAIP2

manipulations, instead, do not directly affect the ongoing spiking activity or execution of learned behavior.

Molecularly-based plasticity manipulations have been used to test the causality of brain areas for

learning[11,38,39,76,80]. We further advanced such methods with cell-type-specific manipulation and *in vivo*

electrophysiology. This approach could effectively map learning mechanisms across behavioral tasks and

brain areas.

Animals learn appropriate types and timing of movement. CaMKII manipulations in ALM blocked learning

of lick timing, without affecting the acquisition of tongue movement to the lick port or cue association

(Fig.1). Thus, different mechanisms may underlie learning movement types. Interestingly, learning

movement types, such as sensory-motor association and learning new kinematics, generates new M1

activity patterns during movement across days[3–8,13]. Although we observed low-dimensional

reconfiguration of ALM preparatory dynamics within sessions (Fig. 3), longitudinal recordings are required

to test the emergence of activity patterns over days. In addition, our manipulations do not distinguish the

type of synaptic plasticity responsible for learning. Furthermore, the pathways providing reward

information and learning rule in PT neurons are unknown. The long-term goal is to explore these questions

by directly and longitudinally monitoring synaptic plasticity and spiking activity of PT neurons during

learning.

Impairments in learning and memory have devastating consequences in life. PT neurons are the homologous

population of neurons vulnerable to amyotrophic lateral sclerosis and frontotemporal disorder in humans[81].

Identifying neocortical cell types required for learning, our findings and approach could be a foundation

for future translational research of various diseases and injuries that affect learning.

Acknowledgment

We thank N. Li, L. Colgan, T. Wang, D. Fitzpatrick, and Inagaki lab members for comments on the

manuscript, M. Inagaki and L. Walendy for animal training, V. Grimaldi and R. DiCicco for histology, P.

Scarpinato for DeepLabCut analysis, J. Yu for molecular biology, and H. Shearin and other MPFI ARC

members for animal care. This work was funded by ZIA MH002497-34 (C.R.G.), Howard Hughes Medical

Institute (S.R.), Max Planck Florida Institute for Neuroscience (H.K.I. and R.Y.), Max Planck Free Floater

Program (H.K.I.), NIH New Innovator Award (NINDS and OD; 1DP2NS132108; H.K.I.), Searle Scholars

Program (H.K.I.), Klingenstein-Simons Fellowship (H.K.I.), and McKnight Scholar Award (H.K.I.).

Author Contributions

S.M., K.H., and H.K.I. planned and performed experiments and analyzed data. Z.Y., S.M., and H.K.I.

developed the motor timing task. C.R.G. and K.H. performed histology. F.L. and S.R. performed network

modeling. A.J. and R.Y. performed ex vivo recordings. S.M. and H.K.I. wrote the paper with input from

all the authors.

Competing interests

The authors declare no competing interests.

Author Information

The authors declare no competing interests. Correspondence and requests for materials should be addressed

to Hidehiko.inagaki@mpfi.org.

METHOD DETAILS

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mice

This study is based on both male and female mice (age > P60, except for acute slice recordings that were P28). We used seven mouse lines: C57BL/6J (JAX# 000664), Tlx-Cre PL56 (RRID: MMRRC_041158-UCD)[60], GRP-Cre KH288 (RRID: MMRRC_031183-UCD)[60], Sim1-cre KJ18 (MGI: 4367070)[60], LSL-Cas9 (B6J.129(B6N)-Gt(ROSA)26Sortm1(CAG-cas9-EGFP)Fezh/J, JAX# 026175)[61], CaMKIIα-cKO (JAX# 006575)[59] and, Vgat-ChR2-EYFP (JAX# 014548)[82]. Transgenic mice with non-C57BL/6J backgrounds were backcrossed to C57BL/6J for at least three generations. See Supplementary Table 1 for mice used in each experiment.

All procedures were in accordance with protocols approved by the MPFI IACUC committee. We followed the published water restriction protocol[83] (with a modification of a minimum of 0.6 ml water per day to compensate for the high humidity in Florida). Mice were housed in a 12:12 reverse light: dark cycle and behaviorally tested during the dark phase. A typical behavioral session lasted between 1 and 2 hours. Mice obtained all of their water in the behavior apparatus. Mice were implanted with a titanium headpost for head fixation and single-housed.

Virus injection

We followed published protocols (dx.doi.org/10.17504/protocols.io.bctxiwpn) for virus injection. See Supplementary Table 1 for detailed descriptions of viruses and injection coordinates. See Supplementary Table 2 for a list of viruses used in this research[39,40,84].

Behavior

A day before the training of the motor-timing task, we habituated mice with the experimental setup. A water-restricted mouse was head-fixed and placed in a training rig. 100 water drops (approximately 2 µL/drop) were delivered through a lickport at random timing (inter-trial intervals (ITI) sampled from an exponential distribution with a mean of 7.5s) without any sensory cue.

Next, we performed two-step training of the motor-timing task: cue association (2-3 days) and delay training (~6 days). Both training phases shared the following general task structure: At the beginning of each trial, an auditory cue was presented, which consisted of three repeats of pure tones (3 kHz, 150 ms duration with 100 ms inter-tone intervals). A delay epoch started from the onset of the cue presentation. Licking during the delay epoch aborted the trial without water reward, followed by a 1.5 s timeout epoch. Licking during the 3 s answer epoch following the delay was considered a 'correct lick', and a water reward (approximately 2 µL/drop) was delivered, followed by a 1.5s consumption epoch. Trials without lick during the delay and answer epochs were considered 'no response' trials. Trials were separated by ITI randomly sampled from an exponential distribution with a mean of 2.2 s with 0.3 s offset (with a maximum ITI of 5 s). This prevented mice from predicting the trial onset without cue. Animals had to withhold licks during the full ITI epoch for the next trial to begin (if they licked, ITI was repeated). In a fraction of randomly interleaved trials, the auditory cue and water reward were omitted to assess spontaneous lick rate ('no cue' trials).

The cue association phase lasted for 2 or 3 sessions with 250-300 trials each. 15% of trials were no cue trials. The delay epoch duration was set to a minimum (0.1 s). After the second session, an animal was considered to have learned the cue association if the median lick time (the first lick time from the cue onset) in the last 100 trials was lower than 0.5 s. The few mice that failed to learn cue association within 3 sessions were excluded from the subsequent experiments (6/198 mice).

The cue association was followed by delay training which lasted for at least 6 days or until the animal
reached a 1.8 s delay duration (beyond this delay, lick timing became too variable and unstable with our
training condition). Sessions were terminated if animals stopped licking for 50 consecutive cue trials or
reached 1000 trials. 5% of trials were no cue trials. The delay duration started at 0.1 s and automatically
increased based on the performance: if the probability of correct trials in the last 100 cued trials with the
same delay duration exceeded 30%, the delay duration was increased by 0.1s. The delay duration at the
beginning of each session was set to 0.1s less than the final delay duration in the previous session (with a
minimum delay duration of 0.1 s), as animals did not reach the criteria performance for the final delay
duration on the last day.
411
412 For the electrophysiological recording of expert mice (Fig. 3 and Extended Data Fig.2g-o), animals were
413 first trained up to 1.8 s delay following the protocol above. Then, the animals were trained with a 1.5 s
414 delay duration for at least two weeks until their performance was stable with a masking flash (see
415 *Optogenetics*). The delay duration was fixed at 1.5 s during the recording.
416
417 To avoid bias, 1) behavior was automatically controlled by Bpod (Sanworks) and custom MATLAB codes,
418 2) experimenters were blinded to the genotype of mice, and 3) control and experimental animals were run
419 in parallel.
420
Optogenetics
For paAIP2 experiments (Fig. 1 and 2c), animals were implanted with a clear skull cap[83]. Stainless steel
tubes (0.062" OD, 0.052" ID, 8988K36, McMaster CARR) were cut into 3mm long pieces and glued
directly on top of the clear skull cap above the target region (ALM or M1) to serve as sleeves for optic
fibers. 470 nm LED light (M470F3, Thorlabs) at 0.5 mW power (power measured at the fiber tip) was
delivered bilaterally through fiber optics (NA 0.39, 400 μm core diameter, M98L01, Thorlabs). The light
was on for 1 second every 5 seconds (0.2 Hz) throughout the training session (LED illumination timing was
independent of behavior). For control light-off sessions, the LED light was directed away from the skull
with the same illumination protocol (i.e., mice could see flickering light as in the light-on condition).
430
431 For SN-CALI experiments (Fig. 2f), the same clear skull protocol was used. 595 nm LED light (M595F2,
432 Thorlabs) at 0.75 mW power was delivered bilaterally for 1 minute every 10 minutes through optic fibers.
433
434 For paAIP2 manipulation during electrophysiological recordings (Fig. 4), 470nm LED light was delivered
435 through an optic probe holder (OFPH_100_500-0.63_FC, Doric lenses) and probe tip (0.63NA, 500 μm
436 core diameter, OPT_500-0.63_FLT, Doric lenses). The location of the optic probe tips was adjusted to
437 cover the whole craniotomy (~ 1.5 mm). 470 nm light at 3 mW power (at the probe tip) was delivered
438 bilaterally for 1 second every 5 seconds (0.2 Hz).
439
440 For paAIP2 manipulation in expert mice (Fig.1e and Extended Data Fig. 2g-o), the light was turned on at
441 least 30 minutes after the session onset. To prevent mice from distinguishing photostimulation and control
442 trials, a 'masking flash' (1 ms pulses at 10 Hz) was delivered near the eyes using 470 nm LEDs (Luxeon
443 Star) throughout the session.
444
Extracellular electrophysiology
A small craniotomy (diameter, 1-1.5 mm) was made over the recording sites a day before the first recording
session. Extracellular spikes were recorded using 64 ch two-shank silicon probes (H-2, Cambridge
Neurotech). Voltage signals were multiplexed, recorded on a PCI6133 board (National instrument), and
digitized at 400 kHz (14-bit). The signals were demultiplexed into 64 voltage traces sampled at 25 kHz
and stored for offline analysis. All recordings were made with the open-source software SpikeGLX
(http://billkarsh.github.io/SpikeGLX/). During recordings, the craniotomy was immersed in a cortex buffer

(125 mM NaCl, 5 mM KCl, 10 mM glucose, 10 mM HEPES, 2 mM $MgSO_4$, 2 mM $CaCl_2$; adjust pH to
7.4). Brain tissue was allowed to settle for at least five minutes before recordings.

454
455   ***Histology***
456   Mice were perfused transcardially with PBS, followed by 4% PFA / 0.1 M PBS. Brains were post-fixed
457   overnight and transferred to 20 or 30 % sucrose PB before sectioning on a freezing microtome. Coronal 50
458   μm free-floating sections were processed using standard fluorescent immunohistochemical techniques.

459
For images in Extended Data Fig. 4, all sections were stained with NeuroTrace® 435/455 Blue Fluorescent
Nissl Stain (Thermo Fisher Scientific, N21479). The fluorescent label was amplified with chicken anti-GFP
(Thermo Fisher Scientific, A10262) and goat anti-chicken 488 (Thermo Fisher Scientific, A11039). Slide-
mounted sections were imaged on a Zeiss microscope with a Ludl motorized stage controlled with
Neurolucida software (MBF Bioscience, Williston VT). Imaging was done with a 10× objective and a
Hamamatsu Orca Flash 4 camera. Every section from the frontal pole through the brainstem were imaged.

466
For the CaMKIIα staining (Extended Data Fig. 3), GFP signal was amplified with rabbit anti-GFP (Thermo
Fisher Scientific, A11122, RRID: AB_221569, 1:500) and goat anti-rabbit 488 secondary antibodies
(Thermo Fisher Scientific, A11008, RRID: AB_143165, 1:500). The CaMKIIα was labeled with anti-
CaMKII-α (6G9) (Cell Signaling Technology, #50049, RRID: AB_2721906, 1:250) and goat anti-rabbit
647 secondary antibodies (Thermo Fisher Scientific, A21236, RRID: AB_2535805, 1:500). Sections were
imaged with a confocal laser scanning microscope (ZEISS, LSM 980) using a 20× objective.

473
Acute slice recording
AAV was injected into P28 mice to label PT_{upper} neurons in ALM bilaterally (n = 5; see Extended Data
Table 1 for injection coordinate). Two to three weeks later, blue light (0.5mW 0.2Hz for one hour)
illumination was given to one hemisphere under isoflurane sedation. Immediately after this, a slice was
prepared and recorded (we also performed ex vivo illumination of brain slices, which yielded similar
results; data not shown). Mice were perfused intracardially with a chilled choline chloride solution (124
mM Choline Chloride, 2.5 mM KCl, 26 mM $NaHCO_3$, 3.3 mM $MgCl_2$, 1.2 mM NaH_2PO_4, 10 mM Glucose,
and 0.5 mM $CaCl_2$, pH 7.4 equilibrated with 95 % O_2 / 5 % CO_2). The brain was removed and placed in the
choline chloride solution. Transverse slices (300 μm) from both hemispheres containing the ALM were cut
using a vibratome (Leica) and maintained in a submerged chamber at 32 °C for 1 h and then at room
temperature in oxygenated artificial cerebrospinal fluid (ACSF: 127 mM NaCl, 2.5 mM KCl, 2 mM $CaCl_2$,
1mM $MgCl_2$, 25 mM $NaHCO_3$, 1.25 mM NaH_2PO_4 and 25 mM glucose).

486
paAIP2-GFP labeled ALM PT_{upper} neurons were visualized using epifluorescent illumination. Whole-cell
current-clamp recordings were obtained in labeled neurons using a Multiclamp 700B amplifier. Patch
pipettes (3-6 ΩM) were filled with a potassium gluconate solution (130 mM K gluconate, 10 mM Na
phosphocreatine, 4 mM $MgCl_2$, 4 mM NaATP, 0.3 mM MgGTP, 3 mM L- Ascorbic acid, 10 mM HEPES.
pH 7.2, 320 mOsm). These experiments were performed at room temperature (~25°C) in oxygenated ACSF.
Recordings were digitized at 10 kHz and filtered at 2 kHz. Current injections were given in 100 pA
increments from -100 to 400 pA. Threshold, AP half-width, and time of AP half-width were analyzed in
the current step where the first AP was observed. Recordings were performed in both hemispheres (one
with illumination before recording and the other without), and the experimenter was blinded to the identity
of the illuminated hemisphere during recording. All data were acquired and analyzed with custom-written
C# and MATLAB codes.

498
Molecular Biology
For pAAV-CaMKII -DIO-mEGFP-P2A-paAIP2, we subcloned the mEGFP-P2A-paAIP2 sequence from
the pAAV-CaMKII-mGFP-P2A-paAIP2 (addgene: 91718) into the pAAV-CaMKII-FLEX-MCS plasmid
(MPFI molecular core) using AscI and BamHI. For pAAV-U6CaMKIIgRNA-hsyn-mScarlet, we

synthesized U6-CaMKIIα gRNA sequence[30] and inserted it into pAAV-hSyn-mScarlet (addgene: 131001)
using XbaI and ApaI. AAV2/9 based on these plasmids were packaged by UNC vector core.
505
For validation of efficacy of gRNA (Extended Data Fig. 3a and b), we transfected Neuro-2A cells in 6 well
plates with pCAG-cre, pLenti-Cas9-GFP (addgene: 86145), and pAAV-U6CaMKIIgRNA-hsyn-mScarlet
using Lipofectamine[TM] 2000 (Invitrogen). Two days after lipofection, GFP+ mScarlet+ cells were sorted
(BD FACSAria™ Fusion), and genome DNA was extracted using DNeasy Blood and Tissue Kit
(QIAGEN). Index was added using PCR with Nextera XT Index Kit (Illumina). We used MiSeq 300 cycle
ver. 2 (Illumina) for the sequencing.
512
513
514
515 **QUANTIFICATION AND STATISTICAL ANALYSIS**
516
517 *Behavioral analysis*
518 Mice often ignored several cue trials at the beginning of each session. In addition, sated mice stopped
519 licking at the end of sessions. To analyze behavior while mice are engaged in the task, we analyzed all trials
520 between the first occurrence of 5 consecutive cue trials with licks and 20 trials before the last occurrence
521 of 3 consecutive no-response trials.
522
523 We analyzed the time of the first lick after the cue (referred to as 'lick time'). Electrical lick ports measured
524 lick time, detecting the tongue's contact with the lick port. To plot learning curves in Fig.1b and c (inset),
525 all cue trials within each session were binned every 50 trials. Then, histograms of lick time were computed
526 for each bin and were normalized to the respective peaks.
527
528 To compare learning across conditions (Figs.1, 2 and Extended Data Fig. 1 and 7), we used the last 100 cue
529 trials in each session to compute the median first lick time, coefficient of variation (standard deviation
530 divided by the mean) of the first lick time, and no response rate. To calculate lick time, 'no-response' trials
531 were excluded. If mice reached a 1.8 s delay before 6 days of the delay training, the training was stopped,
532 and the lick timing on the last behavior session was duplicated for analysis.
533
534 To compare the effect of paAIP2 manipulation in expert mice, we analyzed 3 blocks of trials for within-
535 animal comparisons (Fig.1e, and Extended Data Fig.2k and l). In Fig. 1e, we had two control conditions:
536 'Light off', data of expert mice without paAIP2 manipulation (light onset was artificially and randomly
537 assigned from those in Light on sessions); 'Light on, shuffle', data of light on sessions, but the light onset
538 was randomly reassigned across sessions.
539
Videography analysis
High-speed (300 Hz) videography of orofacial movement (side view) was acquired using a CMOS camera
(Chameleon3 CM3-U3-13Y3M-CS, FLIR) with IR illumination (940nm LED). We used DeepLabCut[85] to
track the movement of the tongue and jaw. Movements along the dorsoventral direction were analyzed and
plotted in Extended Data Fig 1. Trajectories were normalized: the mean position before the cue was
subtracted from trajectories and divided by the minimum value (thus, the downward movement of the
tongue and jaw is upward in the plot). The onset of jaw movement in each trial is the first time point after
the cue when the normalized movement trajectory exceeds 0.15. The onset of tongue movement is when
DeepLabCut first detects the tongue after the cue.
549
Extracellular recording analysis
JRClust[86] (https://github.com/JaneliaSciComp/JRCLUST) with manual curations was used for spike
sorting. We used a combination of quality metrics to rigorously select single units for analysis (Extended

Data Fig. 6). First, a false positive rate was estimated according to the inter-spike interval violation[87], and
units with a false positive rate < 0.15 were selected. Second, we extracted spike features (2 principal
components of spike shape recorded in 3 neighboring recording sites) of all spikes recorded at the same or
adjacent recording sites. Then, we calculated the Mahalanobis distance of spike features of each spike from
the center of the spike cluster (unit) of interest. We performed the receiver operating characteristic (ROC)
analysis of Mahalanobis distance to distinguish spikes within vs. outside the cluster of interest. A recording
session was divided into time bins containing consecutive 1000 spikes of the unit of interest. AUC was
calculated for every time bin. Units with a mean AUC (across time bins) > 0.9 and without any time bin
with AUC<0.75 were selected. Third, units with low spike rates (less than 0.1 spikes per s) were excluded
from the analysis. Units passing all these criteria were deemed single units.
563
We recorded 3689 neurons passing the quality metrics in ALM from 30 mice during learning. Putative
pyramidal neurons (units with spike width > 0.5 ms [88]; 3203 among 3689 neurons) were analyzed. To define
preparatory cells, we compared the spike rates in the baseline (0 to 0.2 s before the cue) vs. during the task
(from cue to first lick) in trial 21-95 of the session ('Early trials' in Fig. 4a). Cells with a p-value lower than
0.05 (signed-rank test) were considered task-modulated cells and analyzed in Figs. 3 and 4 (See Extended
Data Table 3 for details). Cells with positively and negatively task-modulated cells (cells with significant
increase or decrease in spike rate during the task) were analyzed separately (except for Fig.3 c, d, g, and h)
so that they do not cancel out. Since there were more positively modulated cells (Extended Data Fig. 7e),
the main figures show the analysis of positively modulated cells.
573
For the peri-stimulus time histograms (PSTHs) in Fig. 3 and Extended Data Fig. 5, trials were pooled based
on the first lick time. Analyzed time ranges were T = [0.05~0.20, 0.20~0.35, 0.35~0.50, 0.50~0.65,
0.65~0.80, 0.80~0.95, 0.95~1.1] second for recording during learning, and T = [0.95~1.1, 1.1~1.25,
1.25~1.4, 1.4~1.55, 1.55~1.7, 1.7~1.85, 1.85~2.0] second for recording in expert mice. Ten trials with the
first lick within the time range were randomly selected, and the spike rate was averaged. If the number of
trials within the range was less than 10, the range was not included in the analysis. PSTHs were smoothed
with a 50 ms causal boxcar filter. For Extended Data Fig. 2h, PSTHs were smoothed with a 20 ms causal
boxcar filter to detect fast changes in activity. SEM was based on hierarchical bootstrap: first, we randomly
selected animals with replacement; second, we randomly selected sessions of each animal with
replacement; and third, we randomly selected cells within each session with replacement (1000 iterations).
584
To temporally warp the PSTH (Fig. 3), we linearly scaled the spike timing in the time range T_{cue} > 0.1 by
$(LT_{target}$ - 0.1)/$(LT_{trial\ to\ be\ warped}$ - 0.1), where T_{cue} denotes time after the cue, and LT denotes the first lick time
in each trial type (LT_{target} was 1 for Fig.3a,b, and f). After the warping, we calculated PSTH and smoothed
it with a 50 ms causal boxcar filter. We did not warp $0 < T_{cue} < 0.1$ as the onset of cue response showed the
same temporal profile regardless of the lick time (most likely determined by the latency of cue input to
ALM[47]). Because of this, we did not warp the first trial type with a lick latency between 0.05~0.20 s during
learning.
592
Cue response (Fig. 3 and 4, Extended Data Fig.2n) was normalized for the baseline spike rate:
$\overline{SR_{0.1 < T\ cue < 0.15}}$ - $\overline{SR_{-0.2 < T\ cue < 0}}$, where SR_T denotes mean spike rate in the time range indicated in T.
Similarly, pre-lick activity ($\overline{SR_{-0.05 < T\ lick < 0}}$) was normalized by the baseline spike rate. To calculate
'Time between 10-90% activity' (Fig. 3 and 4), PSTH was normalized by the minimum and maximum
spike rate between cue and lick. Then, the time from the first time point crossing 10% to the last time point
crossing 90% was measured (in the case of negative modulated cells, Extended Data Fig. 7, activity was
flipped). Mean speed (Extended Data Fig. 7) is the mean of absolute change in spike rate between cue to
lick (here, spike rate was calculated in 50 ms time bin). We performed hierarchical bootstrap for statistics
and SEM (1000 iterations).
602

603   For the correlation analysis (Fig. 3d and h, and Extended Data Fig. 7g), we looked for an $n \times 1$ unit vector
604     $r_{T, \text{trial type}}$, representing the spike rate of all n neurons at time point T of a trial type of interest. Then, we
605   calculated Pearson's correlation of r between the two trial types indicated in the plot across time points (in
606   the case of Extended Data Fig. 7g, autocorrelation was calculated).
607
608   For recording in the expert mice (Extended Data Fig. 2), we recorded 716 neurons that passed the quality
609   metrics in ALM in 20 sessions from 4 expert mice. 647 putative pyramidal neurons were analyzed.
610

611   ***Ramping direction analysis (Extended Data Fig. 8)***
612   To calculate ramping mode (**RM**) for a population of n recorded neurons, we looked for an $n \times 1$ unit vector
613   that maximally distinguished the mean activity before the trial onset (0 - 0.2 s before cue; $r_{\text{before cue}}$) and the
614   mean activity before the first lick (0 – 0.15 s before the first lick; $r_{\text{before lick}}$) in the *n-dimensional* activity
615   space. We defined a population ramping vector: $w = r_{\text{before lick}} - r_{\text{before cue}}$. **RM** is w normalized by its norm.
616

617   In each recording session, we randomly selected 24 cue trials with the first lick time above 0.25 s. We
618   define RM based on the activity of these randomly selected trials, and projected activity in different sets of
619   trials for the cross-validation. To calculate the "activity explained", we calculated the squared sum of the
620   spike rate after subtracting the baseline (mean spike rate 0 – 0.2 s before cue onset) across neurons. We
621   calculated the squared sum of the activity along RM after subtracting the baseline.
622

623   ***Histology Analysis***
624   For Extended Data Fig. 3e, cell counting and signal analysis were conducted using MATLAB. Cells were
625   binary classified into GFP and/or RFP positive/negative based on the fluorescence signal intensity. The
626   CaMKIIα signal intensity within the cell was normalized by the signal in the surrounding areas
627   (median of areas within 12-60 pixels (0.6 μm/pixel) from the cell).
628

For Extended Data Fig. 4, imaged sections were processed with NeuroInfo software (MBF Bioscience,
Williston, VT) to align the serial sections into a whole brain volume. Neurons throughout the ALM cortex
were detected and marked using the automatic cell detection function in NeuroInfo software, which uses a
modified Laplacian of Gaussian (LOG) algorithm that detects labeled neuron perikarya based on size and
fluorescence intensity with a neural network that eliminates signal artifacts. The accuracy of cell counts
compared to visual detection is approximately 95%.
635

Network Model
Using a dynamical systems approach[89,90], we consider four variables representing the average membrane
currents (h) and spike rates ($r = f(h)$, where $f(h)$ is the neural activation function) of neuronal
populations in different regions of the brain. Specifically, we modeled two excitatory populations
(pyramidal tract, h_{PT}, and intratelencephalic, h_{IT}) and one inhibitory population (h_{In}) in the premotor
cortex, one excitatory population in the thalamic nuclei receiving PT input (h_{th}). The average membrane
current dynamics of population k are described by the following nonlinear differential equation:

643
$$\tau_k \dot{h}_k(t) = -h_k(t) + \sum_j W_{kj} f\big(h_j(t)\big) + I_k + b_k I^{cue}(t) + c_k I^{tonic}$$

Where τ_k is the membrane time constant of population k, W_{kj} are elements of the connectivity matrix
between presynaptic population j and postsynaptic population k, I_k^{DC} is the baseline input current. $I^{cue}(t)$
is the external input current provided to ALM neurons elicited by the auditory cue via the synaptic weights
b_k, and I^{tonic} is the non-contextual tonic input provided to the thalamus via the synaptic weights c_k. We
hypothesized the pathway providing cue input is different from thalamic nuclei maintaining preparatory
activity according to previous work[47]. For all populations, we used a threshold-linear activation function:

650
$$f\big(h_j(t)\big) = [h(t)]_+ = \begin{cases} 0 & if\ h(t) \le 0 \\ h(t) & if\ h(t) > 0 \end{cases}$$

651
The baseline input currents were chosen to be negative for all four populations so that the network dynamics
displays a baseline stable fixed point (lower attractor) for $h_j(t) < 0\ \forall\ j$. The connectivity matrix $\boldsymbol{W}$
respected the biological constraints found in the literature (see Extended Data Table 4), and was chosen so
that a stable fixed point (higher attractor) existed for $h_j(t) > 0\ \forall\ j$. The transition from the lower to the
higher attractor was triggered by $I^{cue}(t)$ which took the form of a 100ms long boxcar function. Ramping
activity emerged as the dynamics evolved towards the higher attractor following the cue. The lick onset
was defined upon reaching a spike rate threshold ($r_{PT}^* = 15$ spikes per second) motivated by the activity of
ALM (Fig.3).

660
We imposed synaptic plasticity between the input relaying the auditory cue and PT neurons (b_{PT}) to
implement learning. To update b_{PT}, we used a heavily simplified learning algorithm. We hypothesized that
the external inputs to the network $I^{cue}(t)$ and I^{tonic} were subject to fluctuations across trials. In particular,
the peak value of the cue signal I^{cue} and I^{tonic} were each drawn from normal distributions with means
μ^{cue}, μ^{tonic} and standard deviations $\sigma^{cue}, \sigma^{tonic}$, respectively(see Extended Data Table 4). Because of
these fluctuations, the spike rate of PT neurons will reach the threshold at a different speed in each trial.
The synaptic weight b_{PT} was either exclusively potentiated (model shown in Fig.4), or depressed (model
shown in Extended Data Fig. 10c-e), at the end of rewarded trials, i.e., trials in which the PT spike rate
exceeded the hard threshold after the delay duration according to the update rule:
670
$$b_{PT}^{tr+1} = b_{PT}^{tr} + \Delta b_{PT}\ (1 - c)$$
671 where $\Delta b_{PT} = \eta\ [b_{PT}^{max} - b_{PT}^{tr}]_+\ I^{cue}$ for potentiation ($\Delta b_{PT} = \eta\ [b_{PT}^{min} - b_{PT}^{tr}]_+\ I^{cue}$ for depression), η
672 being the learning rate. c is the recent reward rate (mean of the last five trials; we added this term to prevent
673 the network from learning when the reward rate was 100%). As in the behavioral experiments with mice,
674 when the reward rate in the last 100 trials reached 30%, the delay duration was increased by 100 ms.

675
To model the behavior and the neural dynamics in expert mice (Extended Data Fig. 10a, b, and e), we used
the same network parameters that were found at the end of the learning process. To vary the lick time, we
changed the non-contextual tonic input $I^{tonic}(t)$, which is compatible with a previous paper modeling
dynamics in expert animals performing a timing task[69]. In the synaptic potentiation model (Extended Data
Fig. 10a-b), stronger tonic inputs lead to slower dynamics, allowing the model to capture the behavior of
expert mice with increasing delay durations. In the synaptic depression model (Extended Data Fig. 10e),
weaker tonic inputs lead to slower dynamics.

683
684 ***Statistics***
685 The sample sizes are similar to the sample sizes previously published in the field. No statistical methods
686 were used to determine the sample size. During spike sorting, experimenters could not tell the trial type and
687 therefore were blind to conditions. All *signed rank* and *ranksum* tests were two-sided. All bootstrapping
688 was done at least over 1,000 iterations.

689
690 ***Reagent and data availability***
691 The new plasmids reported in this paper will be posted to Addgene before publication. The recording data
692 in NWB format and example codes will be shared on DANDI around the time of publication.

693
694

Extended Data Figure 1

Extended Data Figure 1. Characterization of behavior during delay training

a. Tracking of jaw and tongue, using high-speed videography and DeepLabCut[85]. Six example animals (2 per experimental condition) are shown. Left, video clip overlayed with example trajectories of jaw (green) and tongue (magenta) (15 trials). Middle, trajectories of jaw and tongue at the beginning of delay training (from trial 50 to 65 on day 1 of delay training). Trajectories are aligned to lick onset, where lick onset is defined as the moment tongue touches the lick port for the first time after the cue. Black cross, cue (trial) onset. Green cross, the onset of jaw movement (Methods; analyzed in panels **c** and **e**). Right, the same as in the middle but for later in delay training (trial 50 to 65 from the end of day 3 of delay training).

b-f. Summary of the first lick time (**b**), jaw movement onset (**c**), tongue protrusion onset (**d**), time from jaw movement onset to the first lick (**e**), and time from tongue protrusion onset to the first lick (**f**) (Methods). Comparing the first and last 100 trials within a session. Thick lines, mean ± SEM. Thin lines, individual mice (n = 7, 5, and 7 for control, PT_{upper} paAIP2 manipulation, and PT_{lower} paAIP2 manipulation, respectively). The delayed tongue and jaw movement (i.e., withholding of orofacial movement after the cue onset) mainly explains the delayed lick time.

g-i. Characterization of behavior during paAIP2 manipulation. The coefficient of variance (C.V.) of first lick time (**g**), no-response rate (**h**), and number of trials per session (**i**) are shown for animals analyzed in Fig. 2c. Lines, mean ± SEM. No change in C.V. between control vs. paAIP2 manipulation indicates that the manipulation did not affect the variability of lick timing. No significant increase in the no-response rate and no significant decrease in the number of trials in paAIP2 manipulation implies that the manipulation did not block cue-triggered lick and motivation (note that all significant changes are in the opposite direction). *: $p < 0.05$ (bootstrap followed by *Bonferroni* correction; Extended Data Table 3); n.s.: $p > 0.05$ for all comparisons.

j. Relationship between mean vs. standard deviation of first lick time. The standard deviation of lick time scales with its mean, adheres to Weber's law[49,51]. This is why we analyzed CV (standard deviation divided by mean) to evaluate across-trial variability in lick time (e.g., **g**). Circle, the first 100 or last 100 trials of each session. Lines, fit with linear regression. The slope of the regression line: 0.49 (0.43 – 0.56), 0.53 (0.45-0.64), and 1.00 (0.77-1.35), mean (5 – 95 % confidence interval, bootstrap) for during learning (control), during learning (paAIP2 in ALM with light on), and expert, respectively. Thus, ALM paAIP2 manipulation does not affect the variability of lick time.

Extended Data Figure 2

Extended Data Figure 2. The paAIP2 manipulation does not affect excitability or ongoing spiking activity

    a. Retrograde labeling of PT_{upper} neurons in ALM with GFP (with paAIP2) in an acute cortical slice. Right top, example cell. Right bottom, example dendritic shaft.

    b. Representative traces of whole-cell current-clamp recordings from paAIP2 labeled PT_{upper} neurons at three different current steps (−100, 0, and 300 pA) before (black) and after (blue) 60 minutes of *in vivo* blue light stimulation (1 sec on, 4 sec off; Methods).

    c. The mean number of action potentials (AP) evoked by depolarizing current steps. n = 13 cells after stimulation and 12 cells before stimulation. Error bar, SEM.

    d. AP threshold showed no difference between control and stimulation ($p = 0.1225$, ranksum test).

    e. AP half-width showed no difference between control and stimulation ($p = 0.3981$, ranksum test).

    f. Time to the first AP half-width from AP threshold. There was no difference between control and stimulation ($p = 0.7795$, ranksum test).

    g. Schematic for the extracellular electrophysiological recording of ALM with paAIP2 manipulation in expert mice (delay duration, 1.5 s). Blue light (470 nm, 1 s on, 4 s off, 3 mW; Methods) was turned on >30 mins after the session onset.

    h. Blue light-triggered average of ALM activity. Thick line, average PSTH of all putative pyramidal ALM neurons (n = 647 neurons, 4 animals, 20 sessions). Blue bar, blue light pulse (1 s). Shade, SEM. We observed a non-significant increase in spike rate (~0.2 spikes per s, on average) around 100ms after the light onset. This could be a visual response, as we observed a similar response in mice without paAIP2 expression (data not shown).

    i. Average spike rate of individual neurons before and during blue light pulses, calculated over 1s time windows before and during the blue light pulse. Circle, neuron. Dashed line, unit line. No difference in spike rate was observed before and during blue light ($p = 0.69$, signed rank test), indicating that paAIP2 manipulation does not directly affect spiking activity.

    j. Median lick time (5 trial sliding windows; aligned to the first trial with blue light illumination). Thick line, average across sessions (20 sessions from 4 animals). Shaded area, SEM. Vertical dashed lines separate 3 phases in the session: pre-light (0 - 100 trials before blue light onset), post-light phase I (0 - 100 trials after blue light onset), and post-light phase II (100 - 200 trials after blue light onset). The last trial corresponds to ~40 mins of blue light illumination.

    k. Median lick times in the 3 phases in **j**. Thick line, mean ± SEM. Thin lines, individual sessions. No significant difference was observed (*p*-value, signed rank test), indicating that paAIP2 manipulation did not directly affect lick time in expert mice.

    l. Coefficient of variance of the first lick time in the 3 phases in **j**. The same format as in **k**. No significant difference was observed (*p*-value, signed rank test), indicating that paAIP2 manipulation does not directly affect the variance of lick time. The no-response rate did not change significantly, either (data not shown).

    m. Grand average PSTH aligned to the cue onset (for trials with lick times between 0.55 - 2.63 s, corresponding to 10% to 90% quantiles of the lick time distribution in these sessions). n = 357 significantly modulated neurons in ALM. Lines, mean. Shade, SEM.

    n. Comparisons of task-related activity (cue response: mean spike rate over 100-150 ms after cue onset minus baseline spike rate) before and after blue light illumination. Circle, neuron. Dashed line, unit line. We found no significant differences (signed rank test), suggesting that paAIP2 manipulation does not affect task-related activity in expert mice.

    o. The same as **n** but for pre-lick activity (mean spike rate over 50 ms preceding the first lick minus baseline spike rate). We found no significant differences (signed rank test), suggesting that paAIP2 manipulation does not affect task-related activity in expert mice.

Extended Data Figure 3

Extended Data Figure 3. Histological validation of CRISPR/Cas9 KO of CaMKIIα

    a. Sequencing of genomic DNA in Neuro2A cells transfected with guide RNA against CaMKIIα and Cas9 (Methods).

    b. The proportion of amplicon with Indel mutation.

    c. CaMKIIα immunohistochemistry of ALM in CaMKIIα conditional knock out (cKO) mouse injected with AAV-hsyn-cre. Loss of signal was observed around the injection site (ALM; the center of the image), validating the CaMKIIα antibody.

    d. CaMKIIα immunohistochemistry of ALM with cell-type-specific CRISPR/Cas9 KO of CaMKIIα. Loss of CaMKIIα was observed only in cells expressing both Cas9 (green) and guide RNA (magenta, in right panels). Black areas smaller than PT/IT neurons are presumably glia, inhibitory neurons, or neuropile of manipulated cells.

    e. Quantification of CaMKIIα immunostaining signal (Methods).

Extended Data Figure 4

Extended Data Figure 4. Histological validation of cell-type-specific manipulation
a. paAIP2-GFP expression in ALM PT neurons and some layer 2/3 neurons in Sim1-cre KJ18 mice.
b. paAIP2-GFP expression in ALM PT_{upper} neurons (due to the tropism of AAV_{retro}[46], layer 6 neurons
are only sparsely labeled).
c. paAIP2-GFP expression in ALM PT_{lower} neurons.
d. paAIP2-GFP expression in ALM layer 5 IT neurons in Tlx-cre PL56 mice.
e. paAIP2-GFP expression in layer 2/3 IT and some layer 5 IT neurons in Grp-cre KH288 mice.
791
Top, paAIP2-GFP expression in ALM. Bottom panels, paAIP2-GFP expression in 4 coronal sections
from anterior to posterior showing expression in the frontal cortex, striatum and corpus callosum,
thalamus, and medulla. See Extended Data Table 1 for injection coordinates. Signal in the corpus
callosum (white arrows) in **d** and **e** confirms the labeling of IT neurons. Whereas the lack of signal in
the corpus callosum in **b** and **c** confirms a lack of co-labeling of IT neurons. Signal in the thalamus and
internal capsule (white arrowheads) in **a**, **b**, and **c** confirms the labeling of PT neurons. In contrast, the
lack of signal in the thalamus and internal capsule in **d** and **e** confirms a lack of co-labeling of PT
neurons. The labeling of PT neurons in Sim1-cre KJ18 mice (**a**) was sparser than the labeling of PT_{upper}
or PT_{lower} neurons by AAV_{retro} (i.e., **b** and **c**). In addition, we note that Sim1-cre KJ18 labels some layer
2/3 neurons in ALM. Scale bars are identical across panels in each row.

Extended Data Figure 5

802   **Extended Data Figure 5. Spiking activity patterns of single cells**
803   Example neurons recorded during learning and in experts. The data format is the same as in **Fig. 3a**. Note
804   that cells recorded in different animals often show similar reconfiguration of dynamics (e.g., Cell 43, 859,
805   and 480 during learning without paAIP2 manipulation), indicating the reconfiguration is general across
806   animals.

Extended Data Figure 6

Extended Data Figure 6. Quality metrics of spike-sorted units

Drifts in the recording affect spike sorting quality. We implemented rigorous quality metrics evaluating cluster isolation across time points within a recording session to minimize the effect of recording drift in our analysis.

- **a.** An example unit that passed the quality standard. a_1, projection of spike feature along the first PC. Time, the time within the recording session. Black dots, spikes belonging to the unit. Gray dots, all the other spikes sharing the peak channel (i.e., recorded at the same or adjacent recording sites). a_2, spike shape of the unit belonging to this unit. Randomly selected 250 spikes are overlaid. a_3, the inter-spike interval of spikes belonging to the unit (see Methods for false positive rate). a_4, Mahalanobis distance (calculated based on spike features) of each spike from the center of the unit cluster. CDF is shown for spikes belonging to the unit (blue) and all other spikes sharing the peak channel (black). Blue dotted line, 95% point of spikes belonging to the unit. Black dotted line, the distance where the false positive rate reaches 5%. a_5, ROC analysis of the Mahalanobis distance (distinguishing spikes belonging or not belonging to the unit; Methods). AUC is calculated for each time window containing every consecutive 1000 spikes belonging to the cluster.
- **b.** The same as in **a** for different unit.
- **c-d.** The same as in **a** for different units not passing the quality standard due to high FA in inter-spike interval analysis (c) or mean AUC lower than the threshold (d).
- **e** The histogram of mean AUC (AUC averaged across all time windows) across all manually curated units. Quality criterion: mean AUC > 0.9 (dashed line).
- **f** The histogram of the false-positive rate calculated according to the inter-spike interval across all manually curated units.
- **g** No single time window should have AUC < 0.75 to pass the quality criteria. The proportion of AUC below the threshold (across all manually curated units) is shown for all manually curated units.

Extended Data Figure 7

Extended Data Figure 7. ALM dynamics during learning

a-d. Behavioral performance of the animals during ALM recording (different cohorts of mice from those analyzed in Fig2c; the same cohorts of mice analyzed in Fig.4). Time to first lick (**a**), no-response rate (**b**), coefficient of variation (**c**), and number of trials per session (**d**) are shown. Thick lines, mean $\pm$ SEM. Thin lines, animals. N = 10, 5, 7 mice for control, PT_{upper}, and PT_{lower} manipulation, respectively. *: p< 0.05 control vs. PT_{upper} manipulation or control vs. PT_{lower} manipulation conditions (indicated by the color of *; bootstrap followed by *Bonferroni* correction; Extended Data Table 3), n.s., p >0.05 for all comparisons. The result is consistent with behavior without recording (Fig. 2 and Extended Data Fig. 1).

e. The proportion of cells with positive or negative-modulated preparatory activity (spike rate between cue to the first lick significantly higher or lower from that in the baseline, repsectively; two-sided signed-rank test, $p < 0.05$). Error bar, SEM (hierarchical bootstrap). $p > 0.05$ for all comparisons of proportion (on days 1, 2, and 3) between control vs. manipulations (hierarchical bootstrap). Thus, the paAIP2 manipulation does not affect the proportion of preparatory neurons.

f. Sike rate significantly decreased during learning. $(\overline{SR_{trial\ type\ 1}} - \overline{SR_{trial\ type\ 2}})/\overline{SR_{both\ trial\ types}}$ was calculated for each neuron and the distributions of this value are shown as CDF, where $\overline{SR}$ denotes the mean spike rate between cue to first lick. During the training, a significant proportion of neurons decreased spiking activity as mice licked later (blue; $p = 1.18 \times 10^{-6}$, signed-rank test, n = 269 cells; median -0.103; based on neurons shown in Fig. 3g). In contrast, in the expert, there was no significant change in spike rate (black; $p = 0.27$, signed-rank test, n = 344 cells; median: -0.001; based on neurons shown in Fig. 3c) even when the fold-difference in lick time between trial types is roughly matched to that in Day1&2 (purple; $p = 0.69$, signed-rank test, n = 39 cells; median: 0.006). Altogether decrease in spike rate is unique to during learning.

g. Autocorrelation of ALM population activity (top; trials with lick time between 0.50 and 0.65 s) to estimate time-constant of population activity (bottom; different trials are shown in different colors). The correlation of population activity between time points: $(T_{cue} + T_{lick})/2$ vs. following time points, is shown, where T_{cue} and T_{lick} denote the time of cue and lick, respectively. Vertical dotes lines, lick times. Consistent with **Fig. 3**, the time constant of population dynamics increased as mice learned to lick later.

h. Grand average peri-stimulus time histogram (PSTH) of negatively-modulated ALM preparatory neurons. The same format as in **Fig. 3b**. Shade, SEM. Trial types with more than 25 neurons are shown.

i. Characterization of PSTH of positively and negatively modulated ALM preparatory neurons. The format is the same as in **Fig. 3i-k**, and the data of paAIP2 manipulation is overlaid. Mean speed, the average spike rate (SR; spikes per sec) change between cue to lick. Dotted line, expected mean speed (mean pre-lick activity divided by lick time). The mean speed (first column) decreases, and time between 10-90% activity (second column) increases as animals lick later, consistent with a view that dynamics are temporally stretched. The absolute values of the cue response (third column) decrease during learning, whereas pre-lick activity (fourth column) is stable in both positively and negatively modulated cells. Trial types with more than 50 trials were analyzed.

a

b Day 1&2 of training

c Expert

d PT$_{upper}$ paAIP2 manipulation
Day 1&2 of training

e PT$_{lower}$ paAIP2 manipulation
Day 1&2 of training

Extended Data Figure 8

Extended Data Figure 8. Low dimensional population activity during learning
a. Projection of ALM spiking activity to the ramping mode, which maximally distinguishes the
population activity between the baseline and pre-lick (Methods). Top, grand average PSTH aligned
to cue (left) or lick (right). Bottom, projection to the ramping mode. n = 153 preparatory neurons
in ALM (neurons with >= 10 trials for all lick times were analyzed).
b. The square sum of total spiking activity (black) and the square of projection along ramping mode
(blue) at each time point, indicating the large proportion of spiking activity can be explained by
activity along this mode. Shade, SEM (hierarchical bootstrap). Var explained: mean ± SEM.
c. Same as **b** for Expert. N = 123 preparatory neurons in ALM (neurons with >=10 trials for all lick
times were analyzed).
d. Same as **b** for PT_{upper} paAIP2 manipulation, indicating the large proportion of spiking activity can
be explained by activity along this mode even during the paAIP2 manipulation. N = 117 preparatory
neurons in ALM (neurons with >=10 trials for all lick times were analyzed).
e. Same as **b** for PT_{lower} paAIP2 manipulation. N = 140 preparatory neurons in ALM (neurons with
>=10 trials for all lick times were analyzed).

Extended Data Figure 9

Extended Data Figure 9. Similar ALM activity across experimental conditions when mice licked around the same time

    a. Comparisons of the grand average PSTHs of positively-modulated ALM preparatory on day 1 early trials vs. day 3 late trials (regardless of lick time; the same dataset is quantified in **Fig. 4c-d**). PSTHs are aligned to cue (top) or lick (bottom). Control shows a clear change in cue response and ramping activity (red arrows). Early and late trials, first 75 and last trials in the session. See Extended Table 3 for number of neurons analyzed.

    b. When mice lick around the same time (regardless of the day of training), the grand average PSTHs of ALM are similar regardless of paAIP2 manipulation. The grand average PSTHs of positively-modulated ALM preparatory neurons across manipulation types are shown. Activity in trial types with different lick timings is shown in different colors. The paAIP2 manipulation sessions had fewer trials with later licks. The same format as in **Fig.3f**. Shade, SEM. The control data is duplicated from **Fig. 3f** for comparison purposes.

    c. The standard deviation of cue response (related to **Fig. 4d**; calculated based on early trials; late trials yield similar results). The standard deviation and CV (not shown) of cue response are not significantly different between control and paAIP2 manipulation. $p = 0.284, 0.322, 0.274$ (day 1, 2, 3 comparing control vs. PT_{upper} paAIP2 manipulation) and $0.830, 0.800, 0.592$ (day 1, 2, 3 comparing control vs. PT_{lower} paAIP2 manipulation; hierarchical bootstrap). Although the across-trial fluctuation in cue response is the same, the mean cue response changed between early and lick trials only in control (**Fig. 4d**). Thus, the paAIP2 manipulation blocks the directional change in cue response without affecting the across-trial fluctuation.

    d. Clustering analysis to test whether we can distinguish ALM activity patterns between control and paAIP2 manipulation conditions. Z-scored spiking activity of ALM neurons recorded in control animals (WT) and animals with PT-specific paAIP2 manipulations is shown. Neurons from each experimental condition were subsampled and pooled (94 neurons per condition were randomly sampled without replacement; two different groups of neurons were randomly subsampled in the WT condition for cross-validation and denoted WT and WT2). Then we performed hierarchical clustering of the mean activity pattern of these neurons (here, the number of clusters was set to 10). Mean spiking activity in trials with a first lick time between 0.2 and 0.35 s was analyzed. The cluster identity of each neuron is shown in different colors in the middle column, and the type of experimental condition is shown in the right column (see legend in the figure for color scheme). Each cluster contains neurons recorded in all experimental conditions, implying that ALM neurons with similar activity patterns were recorded across experimental conditions.

    e. Fraction of neurons in each cluster in panel **d**. The cluster ID is sorted based on the proportion of neurons in WT data. The correlation coefficients of the proportion of neurons between experimental conditions, indicating similarity in the composition of activity patterns, are shown at the top.

    f. We have repeated this correlation analysis shown in **d-e** 1000 times (sampling random subsets of neurons without replacement; we have tested different numbers of clusters: 5, 10, or 20, all of which yielded similar results) to plot the mean and SEM. C, control: the correlation between WT vs. WT2, showing the upper bound of the correlation coefficient with this procedure and sample size (as neurons are sampled from the same dataset). Both PT_{upper} and PT_{lower} paAIP2 manipulation conditions have similar correlation coefficients to control ($p > 0.05$), indicating that activity patterns in ALM are indistinguishable across conditions when mice lick around the same time, i.e., paAIP2 manipulations in PT neurons do not change task-related activity patterns in ALM.

Network model of expert mice (re. Fig4e-g; model with synaptic potentiation)

a

Input

Cue

Network & dynamics

ALM

Outcome

Lick

Non-contextual tonic input

→ σ_{tonic}

Thalamus

IT cell   GABAergic cell
PT cell   Thalamic cell

b

Average PSTH of IT & PT cells in the model

Cue   Lick

Spikes per s

Time from cue (s)

0   2

Energy landscape

Early lick trials

Cue   Lick

Rolling **steep** landscape

Late lick trials

Cue   Lick

Rolling **shallow** landscape

Network model with synaptic depression between synapse$_{cue \to PT}$

c

Average PSTH of IT & PT cells during learning

Cue   Lick

Spikes per s

Time from cue (s)

0   1

d₁

Cue input

1000   2000   3000

d₂

Synaptic weight (cue → PT)

0.7

0.5

1000   2000   3000

d₃

Time from cue (s)

Control
No plasticity

Distribution of the first lick time

Delay duration

1000   2000   3000

Number of trials

e

Average PSTH of IT & PT cells in expert

Cue   Lick

Spikes per s

Time from cue (s)

0   2

Extended Data Figure 10

Extended Data Figure 10. A model of expert dynamics and a model with synaptic depression

    a. Schema of the expert model (see Methods for details). The network structure is identical to that for learning (**Fig. 4e**). Following literature[51], we provided a non-contextual tonic input to vary lick times across trials (no plasticity is imposed as this is a model of expert).

    b. Dynamics of ALM neurons in the model (top) and corresponding energy landscape (bottom). Different color indicates activity in trials with different lick times. The amplitude of tonic input changed the slope of the landscape, which changed the speed of dynamics. This reproduced the 'temporal scaling' of ramping dynamics consistent with experimental data (**Fig.3**) and previous report[51].

    c. The same format as in **Fig.4f**, but for a network model with synaptic depression between synapses between cue and PT neurons. The network architecture is identical to that in **Fig.4e**, but the reward-dependent plasticity results in a synaptic depression instead of potentiation (Methods).

    d. The same format as in **Fig.4g**, but for the network model with LTD between synapse between cue and PT neurons.

    e. The same format as in **b**, but for the network model with LTD between synapse between cue and PT neurons. Altogether, similar to the potentiation model, the depression model can reproduce the experimental observations.

References

1. Vyas, S., Golub, M. D., Sussillo, D. & Shenoy, K. V. Computation Through Neural Population Dynamics. *Annual Review of Neuroscience* **43**, 249–275 (2020).
2. Inagaki, H. K. *et al.* Neural Algorithms and Circuits for Motor Planning. *Annual Review of Neuroscience* **45**, 249–271 (2022).
3. Laubach, M., Wessberg, J. & Nicolelis, M. A. Cortical ensemble activity increasingly predicts behaviour outcomes during learning of a motor task. *Nature* **405**, 567–571 (2000).
4. Costa, R. M., Cohen, D. & Nicolelis, M. A. L. Differential corticostriatal plasticity during fast and slow motor skill learning in mice. *Curr Biol* **14**, 1124–1134 (2004).
5. Komiyama, T. *et al.* Learning-related fine-scale specificity imaged in motor cortex circuits of behaving mice. *Nature* **464**, 1182–1186 (2010).
6. Huber, D. *et al.* Multiple dynamic representations in the motor cortex during sensorimotor learning. *Nature* **484**, 473–478 (2012).
7. Peters, A. J., Chen, S. X. & Komiyama, T. Emergence of reproducible spatiotemporal activity during motor learning. *Nature* **510**, 263–267 (2014).
8. Masamizu, Y. *et al.* Two distinct layer-specific dynamics of cortical ensembles during learning of a motor task. *Nat Neurosci* **17**, 987–994 (2014).
9. Halverson, H. E., Khilkevich, A. & Mauk, M. D. Relating Cerebellar Purkinje Cell Activity to the Timing and Amplitude of Conditioned Eyelid Responses. *J Neurosci* **35**, 7813–7832 (2015).
10. Makino, H. *et al.* Transformation of Cortex-wide Emergent Properties during Motor Learning. *Neuron* **94**, 880-890.e8 (2017).
11. Adler, A., Zhao, R., Shin, M. E., Yasuda, R. & Gan, W.-B. Somatostatin-Expressing Interneurons Enable and Maintain Learning-Dependent Sequential Activation of Pyramidal Neurons. *Neuron* **102**, 202-216.e7 (2019).
12. Peters, A. J., Liu, H. & Komiyama, T. Learning in the Rodent Motor Cortex. *Annu Rev Neurosci* **40**, 77–97 (2017).
13. Peters, A. J., Lee, J., Hedrick, N. G., O'Neil, K. & Komiyama, T. Reorganization of corticospinal output during motor learning. *Nat Neurosci* **20**, 1133–1141 (2017).
14. Papale, A. E. & Hooks, B. M. Circuit Changes in Motor Cortex During Motor Skill Learning. *Neuroscience* **368**, 283–297 (2018).
15. Wagner, M. J. *et al.* Shared Cortex-Cerebellum Dynamics in the Execution and Learning of a Motor Task. *Cell* **177**, 669-682.e24 (2019).
16. Sun, X. *et al.* Cortical preparatory activity indexes learned motor memories. *Nature* **602**, 274–279 (2022).
17. Abbott, L. F. & Dayan, P. *Theoretical Neuroscience: Computational and Mathematical Modeling of Neural Systems*. (MIT Press, 2005).
18. Takeuchi, T., Duszkiewicz, A. J. & Morris, R. G. M. The synaptic plasticity and memory hypothesis: encoding, storage and persistence. *Philos Trans R Soc Lond B Biol Sci* **369**, 20130288 (2014).
19. Holtmaat, A., Wilbrecht, L., Knott, G. W., Welker, E. & Svoboda, K. Experience-dependent and cell-type-specific spine growth in the neocortex. *Nature* **441**, 979–983 (2006).
20. Hofer, S. B., Mrsic-Flogel, T. D., Bonhoeffer, T. & Hübener, M. Experience leaves a lasting structural trace in cortical circuits. *Nature* **457**, 313–317 (2009).
21. Xu, T. *et al.* Rapid formation and selective stabilization of synapses for enduring motor memories. *Nature* **462**, 915–919 (2009).

996   22. Yang, G., Pan, F. & Gan, W.-B. Stably maintained dendritic spines are associated with
997   lifelong memories. *Nature* **462**, 920–924 (2009).
998   23. Yin, H. H. *et al.* Dynamic reorganization of striatal circuits during the acquisition and
999   consolidation of a skill. *Nat Neurosci* **12**, 333–341 (2009).
24. Xiong, Q., Znamenskiy, P. & Zador, A. M. Selective corticostriatal plasticity during
acquisition of an auditory discrimination task. *Nature* **521**, 348–351 (2015).
25. Biane, J. S., Takashima, Y., Conner, J. M., Scanziani, M. & Tuszynski, M. H.
Reorganization of Recurrent Layer 5 Corticospinal Networks Following Adult Motor
Training. *J. Neurosci.* (2019) doi:10.1523/JNEUROSCI.3442-17.2019.
26. Sanes, J. N. & Donoghue, J. P. Plasticity and primary motor cortex. *Annu Rev Neurosci* **23**,
393–415 (2000).
27. Wilbrecht, L., Holtmaat, A., Wright, N., Fox, K. & Svoboda, K. Structural Plasticity
Underlies Experience-Dependent Functional Plasticity of Cortical Circuits. *J. Neurosci.* **30**,
4927–4932 (2010).
28. Malenka, R. C. & Bear, M. F. LTP and LTD: an embarrassment of riches. *Neuron* **44**, 5–21
(2004).
29. Lisman, J., Yasuda, R. & Raghavachari, S. Mechanisms of CaMKII action in long-term
potentiation. *Nat. Rev. Neurosci.* **13**, 169–182 (2012).
30. Incontro, S. *et al.* The CaMKII/NMDA receptor complex controls hippocampal synaptic
transmission by kinase-dependent and independent mechanisms. *Nat Commun* **9**, 2069
(2018).
31. Matsuzaki, M., Honkura, N., Ellis-Davies, G. C. R. & Kasai, H. Structural basis of long-term
potentiation in single dendritic spines. *Nature* **429**, 761–766 (2004).
32. Yasuda, R., Hayashi, Y. & Hell, J. W. CaMKII: a central molecular organizer of synaptic
plasticity, learning and memory. *Nat Rev Neurosci* **23**, 666–682 (2022).
33. Silva, A. J., Paylor, R., Wehner, J. M. & Tonegawa, S. Impaired spatial learning in alpha-
calcium-calmodulin kinase II mutant mice. *Science* **257**, 206–211 (1992).
34. Silva, A. J., Stevens, C. F., Tonegawa, S. & Wang, Y. Deficient hippocampal long-term
potentiation in alpha-calcium-calmodulin kinase II mutant mice. *Science* **257**, 201–206
(1992).
35. Zeeuw, C. I. D. *et al.* Expression of a Protein Kinase C Inhibitor in Purkinje Cells Blocks
Cerebellar LTD and Adaptation of the Vestibulo-Ocular Reflex. *Neuron* **20**, 495–508 (1998).
36. Luft, A. R., Buitrago, M. M., Ringer, T., Dichgans, J. & Schulz, J. B. Motor skill learning
depends on protein synthesis in motor cortex after training. *J Neurosci* **24**, 6515–6520
(2004).
37. Nabavi, S. *et al.* Engineering a memory with LTD and LTP. *Nature* **511**, 348–352 (2014).
38. Hayashi-Takagi, A. *et al.* Labelling and optical erasure of synaptic memory traces in the
motor cortex. *Nature* **525**, 333–338 (2015).
39. Murakoshi, H. *et al.* Kinetics of endogenous CaMKII required for synaptic plasticity
revealed by optogenetic kinase inhibitor. *Neuron* **94**, 37-47.e5 (2017).
40. Goto, A. *et al.* Stepwise synaptic plasticity events drive the early phase of memory
consolidation. *Science* **374**, 857–863 (2021).
41. Shepherd, G. M. G. Corticostriatal connectivity and its role in disease. *Nat Rev Neurosci* **14**,
278–291 (2013).
42. Tasic, B. *et al.* Shared and distinct transcriptomic cell types across neocortical areas. *Nature*
563, 72–78 (2018).

43. Matho, K. S. *et al.* Genetic dissection of the glutamatergic neuron system in cerebral cortex.
Nature **598**, 182–187 (2021).
44. Li, N., Chen, T.-W., Guo, Z. V., Gerfen, C. R. & Svoboda, K. A motor cortex circuit for
motor planning and movement. *Nature* **519**, 51–56 (2015).
45. Baker, A. *et al.* Specialized subpopulations of deep-layer pyramidal neurons in the
neocortex: bridging cellular properties to functional consequences. *Journal of Neuroscience*
38, 5441–5455 (2018).
46. Economo, M. N. *et al.* Distinct descending motor cortex pathways and their roles in
movement. *Nature* **563**, 79–84 (2018).
47. Inagaki, H. K. *et al.* A midbrain-thalamus-cortex circuit reorganizes cortical dynamics to
initiate movement. *Cell* S0092-8674(22)00146–5 (2022) doi:10.1016/j.cell.2022.02.006.
48. Mauk, M. D. & Buonomano, D. V. The neural basis of temporal processing. *Annu. Rev.*
Neurosci. **27**, 307–340 (2004).
49. Buhusi, C. V. & Meck, W. H. What makes us tick? Functional and neural mechanisms of
interval timing. *Nat. Rev. Neurosci.* **6**, 755–765 (2005).
50. Murakami, M., Vicente, M. I., Costa, G. M. & Mainen, Z. F. Neural antecedents of self-
initiated actions in secondary motor cortex. *Nat. Neurosci.* **17**, 1574–1582 (2014).
51. Wang, J., Narain, D., Hosseini, E. A. & Jazayeri, M. Flexible timing by temporal scaling of
cortical responses. *Nat Neurosci* **21**, 102–110 (2018).
52. Banerjee, A., Chen, F., Druckmann, S. & Long, M. A. Neural dynamics in the rodent motor
cortex enables flexible control of vocal timing. 2023.01.23.525252 Preprint at
https://doi.org/10.1101/2023.01.23.525252 (2023).
53. Cao, V. Y. *et al.* Motor Learning Consolidates Arc-Expressing Neuronal Ensembles in
Secondary Motor Cortex. *Neuron* **86**, 1385–1392 (2015).
54. Kantak, S. S., Stinear, J. W., Buch, E. R. & Cohen, L. G. Rewiring the brain: potential role
of the premotor cortex in motor control, learning, and recovery of function following brain
injury. *Neurorehabil Neural Repair* **26**, 282–292 (2012).
55. Berlot, E., Popp, N. J. & Diedrichsen, J. A critical re-evaluation of fMRI signatures of motor
sequence learning. *eLife* **9**, e55241 (2020).
56. Guo, Z. V. *et al.* Flow of cortical activity underlying a tactile decision in mice. *Neuron* **81**,
179–194 (2014).
57. Sutton, R. S. & Barto, A. G. *Reinforcement Learning: An Introduction*. (A Bradford Book,
2018).
58. Wayman, G. A. *et al.* Calmodulin-kinases: modulators of neuronal development and
plasticity. *Neuron* **59**, 914–931 (2008).
59. Hinds, H. L., Goussakov, I., Nakazawa, K., Tonegawa, S. & Bolshakov, V. Y. Essential
function of alpha-calcium/calmodulin-dependent protein kinase II in neurotransmitter release
at a glutamatergic central synapse. *Proc Natl Acad Sci U S A* **100**, 4275–4280 (2003).
60. Gerfen, C. R., Paletzki, R. & Heintz, N. GENSAT BAC cre-recombinase driver lines to
study the functional organization of cerebral cortical and basal ganglia circuits. *Neuron* **80**,
1368–1383 (2013).
61. Platt, R. J. *et al.* CRISPR-Cas9 knockin mice for genome editing and cancer modeling. *Cell*
159, 440–455 (2014).
62. Maimon, G. & Assad, J. A. A cognitive signal for the proactive timing of action in macaque
LIP. *Nat. Neurosci.* **9**, 948–955 (2006).

63. Inagaki, H. K., Fontolan, L., Romani, S. & Svoboda, K. Discrete attractor dynamics
underlies persistent activity in the frontal cortex. *Nature* **566**, 212–217 (2019).
64. Hennig, J. A. *et al.* How learning unfolds in the brain: toward an optimization view. *Neuron*
109, 3720–3735 (2021).
65. Kiritani, T., Wickersham, I. R., Seung, H. S. & Shepherd, G. M. G. Hierarchical connectivity
and connection-specific dynamics in the corticospinal-corticostriatal microcircuit in mouse
motor cortex. *J Neurosci* **32**, 4992–5001 (2012).
66. Guo, Z. V. *et al.* Maintenance of persistent activity in a frontal thalamocortical loop. *Nature*
545, 181–186 (2017).
67. Collins, D. P., Anastasiades, P. G., Marlin, J. J. & Carter, A. G. Reciprocal Circuits Linking
the Prefrontal Cortex with Dorsal and Ventral Thalamic Nuclei. *Neuron* **98**, 366-379.e4
(2018).
68. Loewenstein, Y. & Seung, H. S. Operant matching is a generic outcome of synaptic plasticity
based on the covariance between reward and neural activity. *Proc Natl Acad Sci U S A* **103**,
15224–15229 (2006).
69. Wang, J., Narain, D., Hosseini, E. A. & Jazayeri, M. Flexible timing by temporal scaling of
cortical responses. *Nat. Neurosci.* **21**, 102–110 (2018).
70. Lisberger, S. G. Cerebellar LTD: A Molecular Mechanism of Behavioral Learning? *Cell* **92**,
701–704 (1998).
71. Kawai, R. *et al.* Motor cortex is required for learning but not for executing a motor skill.
Neuron **86**, 800–812 (2015).
72. Wang, L., Conner, J. M., Rickert, J. & Tuszynski, M. H. Structural plasticity within highly
specific neuronal populations identifies a unique parcellation of motor learning in the adult
brain. *Proc Natl Acad Sci U S A* **108**, 2545–2550 (2011).
73. Jaeger, H. & Haas, H. Harnessing Nonlinearity: Predicting Chaotic Systems and Saving
Energy in Wireless Communication. *Science* **304**, 78–80 (2004).
74. Masse, N. Y., Rosen, M. C., Tsao, D. Y. & Freedman, D. J. Rapid learning with highly
localized synaptic plasticity. 2022.05.09.491102 Preprint at
https://doi.org/10.1101/2022.05.09.491102 (2022).
75. Boyden, E. S., Katoh, A. & Raymond, J. L. Cerebellum-dependent learning: the role of
multiple plasticity mechanisms. *Annu Rev Neurosci* **27**, 581–609 (2004).
76. Wolff, S. B. E., Ko, R. & Ölveczky, B. P. Distinct roles for motor cortical and thalamic
inputs to striatum during motor skill learning and execution. *Science Advances* **8**, eabk0231
(2022).
77. Shmuelof, L. & Krakauer, J. W. Are We Ready for a Natural History of Motor Learning?
Neuron **72**, 469–476 (2011).
78. Bollu, T. *et al.* Cortex-dependent corrections as the tongue reaches for and misses targets.
Nature **594**, 82–87 (2021).
79. Otchy, T. M. *et al.* Acute off-target effects of neural circuit manipulations. *Nature* **528**, 358–
363 (2015).
80. Kakegawa, W. *et al.* Optogenetic Control of Synaptic AMPA Receptor Endocytosis Reveals
Roles of LTD in Motor Learning. *Neuron* **99**, 985-998.e6 (2018).
81. Suzuki, N. *et al.* The mouse C9ORF72 ortholog is enriched in neurons known to degenerate
in ALS and FTD. *Nat Neurosci* **16**, 1725–1727 (2013).
82. Zhao, S. *et al.* Cell type–specific channelrhodopsin-2 transgenic mice for optogenetic
dissection of neural circuitry function. *Nat Methods* **8**, 745–752 (2011).

83. Guo, Z. V. *et al.* Procedures for behavioral experiments in head-fixed mice. *PLoS ONE* **9**,
e88678 (2014).
84. Marshel, J. H. *et al.* Cortical layer-specific critical dynamics triggering perception. *Science*
365, eaaw5202 (2019).
85. Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with
deep learning. *Nat Neurosci* **21**, 1281–1289 (2018).
86. Jun, J. J. *et al.* Fully integrated silicon probes for high-density recording of neural activity.
Nature **551**, 232–236 (2017).
87. Hill, D. N., Mehta, S. B. & Kleinfeld, D. Quality Metrics to Accompany Spike Sorting of
Extracellular Signals. *J. Neurosci.* **31**, 8699–8705 (2011).
88. Guo, Z. V. *et al.* Flow of cortical activity underlying a tactile decision in mice. *Neuron* **81**,
179–194 (2014).
89. Abarbanel, H. D. & Rabinovich, M. I. Neurodynamics: nonlinear dynamics and
neurobiology. *Curr Opin Neurobiol* **11**, 423–430 (2001).
90. Sussillo, D. Neural circuits as computational dynamical systems. *Curr Opin Neurobiol* **25**,
156–163 (2014).
1149