**The heme oxygenase-1 metalloporphyrin inhibitor stannsoporfin enhances the bactericidal**

**activity of a novel regimen for multidrug-resistant tuberculosis in a murine model**

<u>**Jennie Ruelas Castillo**</u>[1], Pranita Neupane[1], Styliani Karanika[1], Stefanie Krug[1,⊥], Darla Quijada[1],

Andrew Garcia[1], Samuel Ayeh[1], Addis Yilma[1], Diego L. Costa[3, 4], Alan Sher[5], Nader Fotouhi[6],

Natalya Serbina[6], Petros C. Karakousis[1,2,7*]

[1] Center for Tuberculosis Research, Department of Medicine, Johns Hopkins University School

of Medicine, Baltimore, MD, USA

[2] Department of International Health, Johns Hopkins Bloomberg School of Public Health,

Baltimore, MD, USA

[3] Departmento de Bioquímica e Imunologia, Faculdade de Medicina de Ribeirão Preto,

Universidade de São Paulo, Ribeirão Preto, Brazil

[4] Departamento de Imunologia, Instituto de Ciências Biomédicas, Universidade de São Paulo,

São Paulo, Brazil

[5] Immunobiology Section, Laboratory of Parasitic Diseases, National Institute of Allergy and

Infectious Diseases, National Institutes of Health, Bethesda, MD, USA

[6] TB Alliance, New York, NY, USA

[7] Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School of

Public Health, Baltimore, MD, USA

[⊥] Department of Microbiology, University of Washington School of Medicine, Seattle, WA

98195, USA

[*] Correspondence:

Center for Tuberculosis Research

CRB-2

1550 Orleans St., Room 110

Baltimore, MD 21287

petros@jhmi.edu

**Abstract:**

Multidrug-resistant (MDR) *Mycobacterium tuberculosis* (Mtb) poses significant challenges to

global tuberculosis (TB) control efforts. Host-directed therapies (HDT) offer a novel approach

for TB treatment by enhancing immune-mediated clearance of Mtb. Prior preclinical studies

found that inhibition of heme oxygenase-1 (HO-1), an enzyme involved in heme metabolism,

with tin-protoporphyrin IX (SnPP) significantly reduced mouse lung bacillary burden when co-

administered with the first-line antitubercular regimen. Here we evaluated the adjunctive HDT

activity of a novel HO-1 inhibitor, stannsoporfin (SnMP), in combination with a novel MDR-TB

regimen comprising a next-generation diarylquinoline, TBAJ-876 (S), pretomanid (Pa), and a

new oxazolidinone, TBI-223 (O) (collectively, SPaO) in Mtb-infected BALB/c mice. After 4

weeks of treatment, SPaO + SnMP 5 mg/kg reduced mean lung bacillary burden by an additional

0.69 $\log_{10}$ (P=0.01) relative to SPaO alone. As early as 2 weeks post-treatment initiation, SnMP

adjunctive therapy differentially altered the expression of pro-inflammatory cytokine genes, and

CD38, a marker of M1 macrophages. Next, we evaluated the sterilizing potential of SnMP

adjunctive therapy in a mouse model of microbiological relapse. After 6 weeks of treatment,

SPaO + SnMP 10 mg/kg reduced lung bacterial burdens to $0.71 \pm 0.23 \log_{10}$ CFU, a 0.78 log-

fold greater decrease in lung CFU compared to SpaO alone (P=0.005). However, adjunctive

SnMP did not reduce microbiological relapse rates after 5 or 6 weeks of treatment. SnMP was

well tolerated and did not significantly alter gross or histological lung pathology. SnMP is a

promising HDT candidate requiring further study in combination with regimens for drug-

resistant TB.

Keywords: *Mycobacterium tuberculosis*, Host-directed therapies, Heme-oxygenase 1,

chemotherapy, drug resistance

**Introduction:**

*Mycobacterium tuberculosis* (Mtb) is the causative agent of tuberculosis (TB), a deadly disease

that claims millions of lives yearly. The emergence of multidrug-resistant TB (MDR TB), which

is resistant to at least two of the most potent first-line drugs, rifampin, and isoniazid, is a

significant challenge to global health [1], [2]. The World Health Organization estimated that in

2021, there were 465,000 new cases of MDR TB worldwide, and only 60% of these cases were

treated successfully [1]. MDR-TB treatment is lengthy, complex, expensive, and requires the use

of second-line drugs that are less effective and more toxic, highlighting the urgent need for novel

therapeutics [3], [4]. Given the global burden of MDR TB, host-directed therapies (HDT)

targeted at boosting the immune system have the potential to contribute to tuberculosis control to

prevent further cross-resistance with current antibiotics [5], [6].

Heme oxygenase-1 (HO-1) is an enzyme involved in the degradation of heme into biliverdin,

carbon monoxide (CO), and iron [7]–[10]. HO-1 is a known biomarker for active TB in humans

[11]. HO-1 also plays a crucial role in the survival of Mtb inside the host by suppressing immune

responses and promoting bacterial growth [12]. Several studies have shown that Mtb infection

induces HO-1 expression in macrophages [8], [13], leading to the generation of CO, which has

immunomodulatory properties and can suppress the production of pro-inflammatory cytokines,

such as TNF-α, IL-1β, and IL-6 [14], [15]. Additionally, the production of CO can inhibit the

activity of T cells and natural killer cells, which are essential components of the host immune

response against Mtb [16]. HO-1 induction can drive macrophage polarization towards an anti-

inflammatory, M2 phenotype, which is associated with reduced microbicidal activity [17], [18]

and increased persistence of Mtb in the host [19].

Several studies have highlighted the potential utility of HO-1 inhibitors as HDT for TB. Costa *et*

*al.* showed previously that pharmacological inhibition of HO-1 with the metalloporphyrin tin

protoporphyrin (SnPP) during the chronic phase of infection in the C57BL/6 murine model of

TB reduced the lung bacillary burden alone and also augmented the activity of the standard

regimen against drug-susceptible TB [20]. HO-1 inhibition promoted the differentiation of CD4$^+$

T cells into interferon gamma (IFN$\gamma$)-producing Th1 cells. Additionally, SnPP enhanced IFN-$\gamma$-

dependent, nitric oxide synthase 2 (NOS2)-induced nitric oxide production by macrophages,

resulting in enhanced control of bacterial growth [21]. Another compound in this class, the HO-1

inhibitor stannsoporfin (SnMP), has been in clinical development as a therapy for

hyperbilirubinemia in neonates [22]. SnMP has been shown to induce the activation,

proliferation, and maturation of naïve CD4$^+$ and CD8$^+$ T cells via interactions with CD14$^+$

monocytes in *vitro* [23].

In the current study, we compared the adjunctive bactericidal activity of SnPP and SnMP when

co-administered with a novel MDR-TB regimen containing TBAJ-876 (S), pretomanid (Pa), and

TBI-223 (O); (collectively, SPaO) or the first-line drug regimen rifampin (R), isoniazid (H), and

pyrazinamide (Z); (collectively, RHZ) against chronic lung infection with drug-susceptible Mtb

in BALB/c mice. In addition, we tested the ability of adjunctive SnMP to shorten the duration of

curative treatment in a murine model of microbiological relapse.

**Results:**

**In chronically infected BALB/c mice, adjunctive therapy with SnMP increases the bactericidal activity of SPaO**

We investigated SnPP and SnMP as adjunctive HDT agents in BALB/c mice infected with Mtb H37Rv (Fig 1A). Each compound was given alone or co-administered with human-equivalent doses of the first-line regimen RHZ for a total of 6 weeks or the enhanced-potency MDR regimen SPaO for a total of 4 weeks. In a previous study [20], SnPP was dosed at 5 mg/kg, but, in the current study, the dose was increased to 10 mg/kg to more closely approximate the drug exposures observed with SnMP 5 mg/kg. Briefly, plasma pharmacokinetics profiles were determined for SnPP and SnMP up to 24 hours following drug administration. The last time point with detectable blood concentrations was after 8 hours. SnPP 10 mg/kg and SnMP 5 mg/kg achieved similar 24-hour drug exposures in plasma (Fig S1), supporting the selection of these doses for the primary study. Over the course of the study, vehicle-treated animals maintained a steady lung colony-forming units (CFU) plateau, while each of the other regimens reduced the mean lung CFU burden steadily over time (Fig 1B). After 2 weeks of treatment (Fig 1C, Table1), monotherapy with SnPP 10 mg/kg or SnMP 5 mg/kg reduced the mean lung bacillary burden by 0.88 $\log_{10}$ (P = 0.0001) and 0.67 $\log_{10}$ CFU (P = 0.0004), respectively, relative to the vehicle control. Similarly, after 4 and 6 weeks of treatment, SnPP 10 mg/kg reduced the mean lung bacterial burden by 1.2 $\log_{10}$ CFU (P = 0.0001) and 1.1 $\log_{10}$ CFU (P = 0.0001), respectively, relative to the vehicle control (Figs 1D, 1E). Although SnMP 5 mg/kg significantly reduced the mean lung bacterial burden by 1.1 $\log_{10}$ CFU (P = 0.001; Fig 1D) relative to the vehicle control after 4 weeks of treatment, this bactericidal effect was not sustained at 6 weeks (0.4 $\log_{10}$ CFU, P = 0.09 ; Fig 1E). After 2 weeks of treatment, neither SnPP nor SnMP adjunctive therapy showed an additive effect when co-administered with SPaO (Fig 1C). However, after 4 weeks of

treatment (Fig 1D), SPaO + SnMP 5 mg/kg reduced the mean lung bacillary burden to 2.49 ±

0.12 $\log_{10}$ CFU, representing an additional 0.69 $\log_{10}$ CFU reduction compared to SPaO alone (P

= 0.01). Adjunctive treatment with SnMP 5 mg/kg or SnPP 10 mg/kg did not significantly

enhance the bactericidal activity of RHZ at 4 weeks (P = 0.52 and P = 0.89, respectively; Fig 1D)

or 6 weeks (P = 0.97 and P = 0.86, respectively; Fig 1E).

**HO-1 inhibition does not exacerbate lung inflammation and regulates pro-inflammatory**

**immune responses.**

The chronic stages of TB disease are characterized by severe lung inflammation, which results in

long-term lung dysfunction in one quarter of cases despite microbiological cure [24], [25]. After

4-6 weeks of infection, the gross pulmonary pathology of Mtb-infected BALB/c mice is

characterized by pronounced tubercular lesions. At the histopathological level, macrophage and

lymphocyte inflammatory aggregates fill the alveolar spaces, representing cellular

granulomatous lesions [26]. HO-1$^{-/-}$ knock-out mice develop a progressive inflammatory state

[13], [27], and splenocytes isolated from these mice respond to lipopolysaccharide stimulation by

producing inflammatory cytokines [4], [13]. Chronic treatment with SnPP in wild-type mice can

induce tubulointerstitial inflammation and fibrosis [28], while SnMP induces the expression of

pro-inflammatory cytokines in human peripheral blood mononuclear cells [29]. Robust

expression of pro-inflammatory cytokines can exacerbate lung inflammation, however, when

controlled, it can offer antimycobacterial protection to the host [30].

To determine whether HO-1 inhibition modulates TB-associated lung pathology, we first

analyzed the lung weight to body weight ratio (lung/body weight) of Mtb-infected mice treated

with SnPP or SnMP alone or as adjunctive HDT with RHZ or SPaO as a gross indicator of lung

inflammation. An increase in this ratio reflects an increase in lung inflammation caused by

cellular infiltration and the presence of tissue lesions consistent with disease exacerbation [26],

[31]. Antitubercular treatment with RHZ for 4 or 6 weeks or with SPaO for 2 or 4 weeks

significantly reduced ($P < 0.0001$) the mean lung/body weight ratio relative to vehicle (Fig 2A-

C). SnMP or SnPP monotherapy for 2 or 6 weeks also significantly reduced the lung/body

weight ratio relative to vehicle (Fig 2A, C), although the decrease in this ratio was not

statistically significant after 4 weeks of treatment (Fig 2B). The lung/body weight ratio was

significantly ($P < 0.0001$) correlated with the lung CFU burden in each of the treatment groups,

as assessed by the Pearson correlation coefficient ($r = 0.9074$ after 2 weeks of treatment; $r =$

0.8251 after 4 weeks of treatment; and $r = 0.9282$ after 6 weeks of treatment) (Fig S2). Grossly,

after 4 weeks of treatment, the lungs of mice treated with vehicle, SnPP or SnMP contained more

pronounced tubercle lesions compared to those of mice receiving the antitubercular drug

regimens (Fig 2D).

Given the adjunctive activity of SnMP with SPaO after 4 weeks of treatment, we next quantified

the percentage of lung surface area ($\mu m^2$) affected by inflammation at this time point using

hematoxylin and eosin staining to determine if adjunctive SnMP had any impact on macrophage

and lymphocytic infiltration in TB lung lesions (Fig 2E). The predominant inflammatory lesions

observed were cellular granulomas containing myeloid cores and lymphocytic cuffs. As

expected, the percentage of lung surface area involved by inflammation was significantly

reduced by the antitubercular regimens RHZ (10.71% $\pm$ 4.9), SPaO (5.65% $\pm$ 1.9) relative to the

vehicle control (29.26% $\pm$ 6.4; $P = 0.011$, and $P = 0.003$ respectively). However, there was no

significant difference in percentage of lung surface area involved by inflammation between

vehicle (29.26% $\pm$ 6.4) and SnMP monotherapy (29.26% $\pm$ 2.3) ($P > 0.99$) . Similarly, SnMP

adjunctive therapy did not significantly reduce the percentage of lung surface area inflammation

relative to RHZ alone (10.71% ± 4.9 vs 5.74 + 1.2, P = 0.41) or SPaO alone (5.65% ± 1.9 vs 5.74

+ 1.2, P > 0.99).

In order to further investigate the adjunctive contribution of SnMP to the bactericidal activity of

SPaO observed at 4 weeks of treatment (Table 1), we used RT-qPCR to evaluate the expression

of genes encoding pro-inflammatory cytokines and macrophage markers in lung homogenates at

week 2, at which time a significant difference in lung bacillary burden was observed between

treatment groups. Interferon-gamma (IFN-γ) and tumor necrosis factor-alpha (TNF-α) are

important macrophage-activating cytokines [32], [33]. NOS2 expression in macrophages is

upregulated in response to Mtb infection and IFN-γ activation is required for successful bacterial

load reduction following pharmacological inhibition of HO-1 [21]. SnMP monotherapy was

associated with reduced expression of *Tnf-α* (P = 0.005, Fig 3C) relative to vehicle. SPaO alone

was associated with a further reduction in expression of *Ifn-γ* (P = 0.002, Fig 3B), and *Tnf-α* (P =

0.001, Fig 3C); while SPaO + SnMP showed reduced expression of *Nos2* (P = 0.03, Fig 3A), *Ifn-*

*γ* (P = 0.006, Fig 3B), and *Tnf-α* (P = 0.03, Fig 3C) relative to vehicle. These results are

consistent with decreased cytokine production in response to reduced mean lung bacterial

burdens in these groups. *Nos2, Ifn-γ,* and *Tnf-α* expression was comparable in the lungs of mice

treated with SPaO or SPaO + SnMP (Fig 3A-C). Next, we looked for expression changes in the

mouse macrophage markers CD38 and early growth response protein 2 (EGR2), a transcriptional

regulator/M1 marker and a negative T-cell regulator/M2 marker, respectively [34], [35]. SnMP

monotherapy, but not SnMP adjunctive therapy in combination with SPaO, was associated with

significantly higher expression of *Cd38* (P < 0.0001) compared to any other treatment group (Fig

3D). In contrast, *Egr2* expression (Fig 3E) did not differ significantly among groups. Overall,

these data are consistent with reduced expression of pro-inflammatory cytokine genes as a result

of reduced mean lung CFU in the different treatment groups as compared to the vehicle group

(Table 1). However, SnMP treatment also appeared to induce the expression of *Cd38*, potentially

polarizing lung macrophages to an M1 phenotype, which is more effective in killing intracellular

Mtb.

**SnMP adjunctive therapy enhances the bactericidal efficacy of SPaO after 6 weeks of**

**treatment without altering relapse rates**

Next, we evaluated the adjunctive sterilizing activity of SnMP at different dosages (5 mg/kg and

10 mg/kg) in combination with SPaO. We selected treatment durations expected to yield > 50%

relapse rates for the backbone regimen based on the prior literature [36]. The mice were treated

for a total of 5 or 6 weeks, at which time the treatment was discontinued, and microbiological

relapse was assessed 3 months later (Fig 4A). Adjunctive therapy with SnMP 5 mg/kg (SnMP5)

did not significantly alter lung bacillary burdens relative to the SPaO regimen at the time points

assessed. However, after 6 weeks of treatment, SPaO + SnMP 10 mg/kg (SnMP10) reduced

mean lung bacterial burdens to $0.71 \pm 0.23$ $\log_{10}$ CFU, representing an additional 0.78 $\log_{10}$ CFU

reduction (P = 0.005) when compared to SPaO alone (Fig 4B, Table 1). As in the bactericidal

activity study, the lung/body weight ratios of mice treated with adjunctive SnMP5 or SnMP10

did not differ statistically from that of mice treated with the SPaO control regimen after 6 weeks

of treatment (Fig S3A). Additionally, the lung/body weight ratios were maintained in each of the

treatment groups at each relapse time point (Fig S3B). After 5 weeks of treatment, all mice

experienced microbiological relapse with SPaO or SPaO + SnMP10 (Fig 4C). Similarly, the

proportion of relapsing animals was equivalent in the groups receiving SPaO and SPaO +

SnMP10 (66%) (Fig 4D).

**Discussion:**

Inhibition of HO-1 has been proposed as a potential HDT strategy associated with reduced Mtb

growth in vitro [37] and in vivo [20], [21]. Our study investigated the adjunctive bactericidal

activity of a clinical-stage HO-1 inhibitor, SnMP, in a mouse model of chronic TB, as well as its

adjunctive sterilizing activity when combined with an MDR-TB regimen in a murine model of

microbiological relapse. We found that adjunctive therapy of SnMP enhanced the bactericidal

activity of the SPaO regimen, but not that of the RHZ regimen, without exacerbating lung

inflammation. Although adjunctive therapy with SnMP10 continued to enhance the bactericidal

activity of SPaO after 6 weeks, it did not alter relapse rates compared to SPaO alone.

In contrast with prior work [20], we did not detect additive bactericidal activity of SnPP with

RHZ in Mtb-infected mice. These discrepant findings may be explained by methodological

differences in the two studies. In the current study, the administration of R was separated from

that of HZ to limit drug interactions [38], [39], whereas the three antibiotics were administered

together in the study by Costa *et al*. Furthermore, in the study by Costa *et al.*, C57BL/6 mice

were used, while our study used BALB/c mice. Relative to BALB/c mice, C57BL/6 mice are

able to more effectively limit the replication of Mtb through the development of T-cell-mediated

immunity [40], [41]. Interestingly, this discrepancy was observed only when SnPP was given as

adjunctive therapy, as we found that SnPP monotherapy reduced lung bacillary burden, as was

reported previously [20]. These findings highlight the challenges associated with preclinical

studies of HDT for TB, as the host-directed activity of these agents may be model-specific.

Our studies showed that SnMP administered alone regulated the expression of the gene encoding

TNF-α in the lungs. Scharn *et al*. have shown previously that SnPP reduced the secretion of

proinflammatory cytokines, including TNF-α, in Mtb-infected U937 cells [37], [24]. Notably,

CD38 expression was upregulated in Mtb-infected mouse lungs following treatment with SnMP

alone. Although CD38 is necessary for immune cell activation and is expressed by many

immune cell types [42], it appears to be a specific marker for M1 macrophage phenotypes in

murine models [35]. In humans and in mice, M1 macrophages play an essential role in

propagating a host-protective response against Mtb (reviewed in Ahmad *et al.*, 2022) [43], [44].

Collectively, these results highlight an immunoregulatory role for the HO-1 inhibitor SnMP

during Mtb infection. Additional studies are needed to further characterize the molecular

mechanisms by which SnMP promotes Mtb clearance by the host.

Based on published data suggesting the SPaO regimen, at the doses of, S 12.5 mg/kg; Pa 100

mg/kg; and O 100 mg/kg, is able to eradicate lung infection in Mtb-infected mice by week 6

[36], the relapse time points of 5 and 6 weeks were selected to evaluate the adjunctive sterilizing

activity of SnMP. In the current study, we found that, although SnMP increased the bactericidal

activity of SPaO, adjunctive therapy did not reduce relapse rates compared to the background

SPaO regimen. Future studies will focus on testing higher doses of SnMP and extending the

duration of treatment to determine the potential of SnMP adjunctive therapy in shortening the

duration of curative MDR treatment.

Consistent with prior preclinical studies [20], [21], [37], we have found that HO-1 inhibition is a

promising HDT strategy for TB. We show for the first time that the novel HO-1 inhibitor, SnMP,

enhances the bactericidal activity of the novel MDR-TB regimen SPaO in mice and modulates

the expression of several pro-inflammatory cytokine and macrophage marker genes, which have

been implicated in the control of Mtb replication in vivo, although adjunctive therapy with SnMP

10 mg/kg did not reduce relapse rates relative to SPaO. SnMP was well tolerated and did not

alter gross lung pathology or histological inflammation in the lungs. Overall, this study advances

our understanding of HO-1 inhibitors as potential adjunctive HDT agents for TB, but further

research is needed to determine the potential utility of targeting this pathway against drug-

susceptible and drug-resistant TB. In addition, to enhance the translational relevance of this HDT

approach, future medicinal chemistry studies should focus on the development of formulations

with favorable oral bioavailability and toxicity profiles.

**Materials and Methods:**

<u>Pharmacokinetic analyses:</u>

Single-dose PK studies were conducted by BioDuro Inc. (Beijing, China). SnMP and SnPP were

formulated in sodium phosphate buffer (pH 7.4 – 7.8) and administered by the intraperitoneal

route to 9–11-week-old female BALB/c mice. Three mice were used per each time point with 8

time points for each dose group. Blood was collected on 0.25, 0.5, 1, 2-, 4-, 8-, and 24-hours

post-administration and levels of compounds in plasma were quantified by liquid

chromatography-tandem mass spectrometry using AB Sciex API 4000 LC/MS/MS

instrumentation. The following PK parameters were calculated from plasma drug concentrations:

$t1/2$, $tmax$, $Cmax$, $AUClast$, $AUCInf$. All parameters were determined by noncompartmental

analysis using WinNonLin software 8.0.

<u>Bacteria and growth conditions:</u>

Wild-type Mtb H37Rv was grown in Middlebrook 7H9 broth (Difco, Sparks, MD) supplemented

with 10% oleic acid-albumin-dextrose- catalase (OADC) (Difco, Sparks, MD), 0.2% glycerol,

and 0.05% Tween-80 at 37°C in a roller bottle.

_M. tuberculosis_ infection of mice:

Wild-type Mtb H37Rv cultures were grown to an $OD_{600}$ of 0.8-1, followed by 1:200 dilution in

7H9 broth (as stated above). 10 mL of the diluted bacterial culture was used as the inoculum for

aerosol infection. Female BALB/c (6-8-week-old, Jackson Laboratory) were aerosol-infected

with ~100 bacilli of Mtb H37Rv using the Glas-Col Inhalation Exposure System (Terre Haute,

IN). Animals were equally distributed in the five chambers of the Glas-Col machine to ensure

equal infection amongst all animals. The inoculating culture was plated in serial tenfold dilutions

on the day of infection on Middlebrook 7H11 agar containing 10% OADC. On the day post-

infection, 5 mice were sacrificed, the lungs were homogenized using glass homogenizers and

plated on 7H11 agar containing 10% OADC to enumerate the implanted CFU (Fig 1).

Antibiotic and HDT treatment preparation:

Protoporphyrin IX (SnPP) (Evotec, US) and stannsoporfin (SnMP) (Evotec, US) were prepared

with sodium phosphate buffer (pH to 7.4-7.8 with HCl). Rifampin (R), isoniazid (H) and

pyrazinamide (Z) (Sigma-Aldrich) were dissolved in deionized water. Pretomanid (Pa) (RTI)

was dissolved in deionized water containing cyclodextrin micelle (CM-2) formulation,10%

hydroxypropyl-β-cyclodextrin (HPCD) (Sigma) and 10% lecithin (ICN Pharmaceuticals Inc.,

Aurora, OH); TBAJ-876 (S) (Evotec, US) was prepared in 20% HPCD solution acidified with

1.5% 1N HCl, pH 3; and TBI-223 (O) (Evotec, US) was prepared in 0.5% Methylcellulose. The

vehicle administered was either CM-2 formulation containing 10% HPCD (Sigma) and 10%

lecithin (ICN Pharmaceuticals Inc., Aurora, OH) to mock Pa, 0.5% methylcellulose to mock O or

20% HPCD solution acidified with 1.5% 1N HCl to mock S. Vehicle of drugs were made in big

batches and the drugs were dissolved into their corresponding vehicles weekly.

<u>Antibiotic and HDT doses and administration:</u>

In the bactericidal activity studies, the mice received one of the following regimens beginning 28

days after aerosol infection: 1) Vehicle (negative control); 2) SnPP 10 mg/kg; 3) SnMP 5 mg/kg;

4) R 10 mg/kg, H 25 mg/kg,  Z 150 mg/kg (collectively, RHZ); 5) S 6.25 mg/kg, Pa 30 mg/kg

and O 45 mg/kg (collectively, SPaO); 6) R 10 mg/kg, H 25 mg/kg, P 150 mg/kg, SnPP 10

mg/kg; 7) R 10 mg/kg, H 25 mg/kg, P 150 mg/kg, SnMP 5 mg/kg; 8) S 6.25 mg/kg, Pa 30

mg/kg, O 45 mg/kg, SnPP 10 mg/kg; 9) S 6.25 mg/kg, Pa 30 mg/kg, O 45 mg/kg, SnMP 5 mg/kg

. The HO-1 inhibitors were given once daily by intraperitoneal injection (IP). PK analysis of the

HO-1 inhibitors was conducted at 5 mg/kg and 10 mg/kg to test which dosages achieved similar

plasma concentrations. There results also confirmed that the dosages achieved human-equivalent

doses [45]. The other drugs were given orally by esophageal cannulation once daily, except for

Pa and O, which were given twice daily separated by at least 4 hours. For the RHZ regimen, R

was given first, and two hours later, H and Z were administered. The vehicle was administered

via gavage to control mice concurrently with treatments containing active compounds to

experimental mice.

In the relapse model studies, the mice received one of the following regimens beginning 28 days

after aerosol infection: 1) Vehicle (negative control); 2) S 6.25 mg/kg, Pa 30 mg/kg, O 45 mg/kg;

3) S 6.25 mg/kg, Pa 30 mg/kg, O 45 mg/kg, SnMP 5 mg/kg; or 4) S 6.25 mg/kg, Pa 30 mg/kg, O

45 mg/kg, SnMP 10mg/kg. The discrepancies in numbers of animals between relapse groups is

due to unexpected mortality from excessive gavage volume when the drugs were prepared

separately at the start of therapy. Beginning in week 2, 2X concentrations of Pa and O were

made and, at the time of administration, Pa and O were mixed at equal volumes to make a single

gavage volume of 0.2 mL, which was tolerated well by the mice.

For both studies (except for the relapse time points, for which both lobes of the lung were

homogenized and plated), the right lung and 3/5 of the left lung of each mouse were

homogenized using glass homogenizers, and serial tenfold dilutions of lung homogenates in PBS

were plated on Middlebrook 7H11 agar containing 0.4% activated charcoal to reduce the effects

of drug carryover at the indicated time points; entire lung homogenates from relapse animals

were plated on charcoal-containing 7H11 agar for consistency. Plates were incubated at 37°C

and CFU were counted 4 weeks later. The CFU per lung data were calculated using lung

weights, which were measured at the time of organ harvest. For the animals in the relapse

groups, the entire lung was used for CFU determination. Microbiological relapse was defined as

the presence of 1 or more colonies in plated lung homogenates. All procedures were performed

according to protocols approved by the Johns Hopkins University Institutional Animal Care and

Use Committee.

Measurement of lung/body weight ratio:

Lung and body weight measurements in milligrams (mg) and grams (g), respectively, were taken

at the time of harvest using weight scales. The lung/body weight ratio was calculated by dividing

lung(g)/body(g) weight and then multiplying by 100.

<u>Quantitative analysis of lung histopathology</u>

At the time of harvest, the left lung medial apex to base was sectioned and fixed with

4% paraformaldehyde (PFA). After a 24-hour incubation in 4% PFA, the lung section was

transferred to PBS and sent in for processing. The paraffin blocks of lung samples were

processed to obtain three sections per lung spaced 12 μm apart for hematoxylin and eosin

staining. Each lung section was sectioned to 4 μm. Each slide was scanned by a brightfield

microscope at the JHU Oncology Tissue Core Center to obtain digital 40X images. Quantitative

analysis was performed by JRC and AY blinded to treatment allocation and mouse strain. For

each treatment group, a total of 15 scanned slides (3 slides per mice) were reviewed. Regions of

interest corresponding to the absolute areas of total lung surface area and inflammation were

manually drawn and their area quantified using the open-source software QuPath

(https://qupath.github.io/). Inflammation was defined as: 1. "Lymphocytic cuffs", areas of high

cell density due to the predominance of lymphocytes, which are smaller cells with scant

cytoplasm; 2. Regions dense with macrophages, which have relatively large amounts of

cytoplasm. The lung inflammation to whole surface area was quantified and multiplied by 100 to

obtain the percentage.

<u>Cytokine or cell marker gene expression:</u>

A section of the left lower lobe (~ 1/5 of the left lung) was flash-frozen in liquid nitrogen at the

time of harvest and later lysed in TRIzol using a hard tissue homogenizing kit (CK28-R,

Percellys) with the manufacturer-recommended bead-beating protocol (Percellys Evolution;

8,500 rpm, three 15-s cycles with 30-s breaks on ice after each cycle). Total RNA was extracted

using the miRNeasy minikit (217004, Qiagen). RT-qPCR sequences for each primer pair were

ordered from IDT and are listed in the supplementary material Table 2. Two-step RT-qPCR was

performed on a StepOnePlus real-time PCR system (ThermoFisher Scientific) using the high-

capacity RNA-to-cDNA kit (4387406, ThermoFisher Scientific), followed by Power SYBR

green PCR master mix (436877, Applied Biosystems). All samples were tested with three

technical replicates. Target gene expression (cycle threshold (Ct) values) were normalized

relative to the housekeeping gene *β-actin*. The Ct value for each gene in infected test samples

was then normalized to that of uninfected samples and mRNA expression fold change was

determined by the ΔΔCt method.

<u>Statistics:</u>

The differences between groups were assessed using an ordinary one-way ANOVA followed by

different multiple comparison tests or Welch's ANOVA test (depending on the spread of the

data) as stated on the figure panel description. The Prism software (GraphPad, San Diego, CA,

USA) version 9 was utilized for this analysis. Results were considered statistically significant

when the p-value was less than 0.05. The Pearson's correlation coefficient was calculated using

the Prism software.

**Supplemental material:**

**Figure S1:** Plasma pharmacokinetic parameters for SnPP and SnMP in mice.

**Figure S2:** Lung/body weight ratio correlates with lung bacterial burden after 2, 4 and 6 weeks

of treatment.

**Figure S3:** Mouse lung/body weight ratio after 6 weeks of treatment and at the relapse time

point.

**Table 1:** Lung CFU count for time course and relapse study.

**Table 2:** RT-qPCR sequences for each primer pair.

**Author Contributions:**

**Acknowledgments:** These studies were supported by NIAID/NIH grant K24AI143447 and a grant from the TB Alliance to PCK. AS is supported by the intramural research program of the NIAID. We thank the Oncology Tissue Services (SKCCC) at JHU supported by the P30 CA006973 grant for their services and for imaging the H&E-stained slides. We gratefully acknowledge the assistance of Michael Urbanowski, MD, PhD, for his training and guidance in the histopathological quantification of the lung tissues.

**Ethics statement:** The animal study was reviewed and approved by Johns Hopkins University Institutional Animal Care and Use Committee. The animal welfare assurance number is D16-00173 (A3272-01). JHU is registered with the USDA to conduct animal research and has maintained active AAALAC accreditation since 10/4/1974.

**References**:

[1]    World Health Organization, "Global Tuberculosis Report 2022," 2022. [Online]. Available: http://apps.who.int/bookorders.

[2]    K. J. Seung, S. Keshavjee, and M. L. Rich, "Multidrug-resistant tuberculosis and extensively drug-resistant tuberculosis," *Cold Spring Harb Perspect Med*, vol. 5, no. 9, Sep. 2015, doi: 10.1101/cshperspect.a017863.

[3]    E. Pontali *et al.*, "Regimens to treat multidrug-resistant tuberculosis: Past, present and future perspectives," *European Respiratory Review*, vol. 28, no. 152. European Respiratory Society, 2019. doi: 10.1183/16000617.0035-2019.

[4]    C. Lange, D. Chesov, J. Heyckendorf, C. C. Leung, Z. Udwadia, and K. Dheda, "Drug-

resistant tuberculosis: An update on disease burden, diagnosis and treatment,"

*Respirology*, vol. 23, no. 7, pp. 656–673, Jul. 2018, doi: 10.1111/resp.13304.

[5]    D. J. Frank *et al.*, "Remembering the host in tuberculosis drug development," in *Journal*

*of Infectious Diseases*, Oxford University Press, May 2019, pp. 1518–1524. doi:

10.1093/infdis/jiy712.

[6]    G. Kilinç, A. Saris, T. H. M. Ottenhoff, and M. C. Haks, "Host-directed therapy to combat

mycobacterial infections*," *Immunological Reviews*, vol. 301, no. 1. John Wiley and Sons

Inc, pp. 62–83, May 01, 2021. doi: 10.1111/imr.12951.

[7]    R. Tenhunen, H. S. Marver, and R. Schmid, "The enzymatic conversion of heme to

bilirubin by microsomal heme oxygenase.," *Proc Natl Acad Sci U S A*, vol. 61, no. 2, pp.

748–55, Oct. 1968, doi: 10.1073/pnas.61.2.748.

[8]    S. W. Ryter and A. M. K. Choi, "Heme oxygenase-1/carbon monoxide: From metabolism

to molecular therapy," *American Journal of Respiratory Cell and Molecular Biology*, vol.

41, no. 3. pp. 251–260, Sep. 01, 2009. doi: 10.1165/rcmb.2009-0170TR.

[9]    M. D. Maines, "THE HEME OXYGENASE SYSTEM: A Regulator of Second

Messenger Gases," 1997. [Online]. Available: www.annualreviews.org

[10]   S. W. Ryter and R. M. Tyrrell, "The heme synthesis and degradation pathways: role in

oxidant sensitivity. Heme oxygenase has both pro- and antioxidant properties.," *Free*

*Radic Biol Med*, vol. 28, no. 2, pp. 289–309, Jan. 2000, doi: 10.1016/s0891-

5849(99)00223-3.

[11]    N. Rockwood *et al.*, "Mycobacterium tuberculosis induction of heme oxygenase-1

expression is dependent on oxidative stress and reflects treatment outcomes," *Front*

*Immunol*, vol. 8, no. MAY, May 2017, doi: 10.3389/fimmu.2017.00542.

[12]    A. Kumar, A. Farhana, L. Guidry, V. Saini, M. Hondalus, and A. J. C. Steyn, "Redox

homeostasis in mycobacteria: the key to tuberculosis control?," *Expert reviews in*

*molecular medicine*, vol. 13. 2011. doi: 10.1017/s1462399411002079.

[13]    M. H. Kapturczak *et al.*, "Animal Model Heme Oxygenase-1 Modulates Early

Inflammatory Responses Evidence from the Heme Oxygenase-1-Deficient Mouse," 2004.

[14]    B. Wegiel *et al.*, "Carbon monoxide expedites metabolic exhaustion to inhibit tumor

growth," *Cancer Res*, vol. 73, no. 23, pp. 7009–7021, Dec. 2013, doi: 10.1158/0008-

5472.CAN-13-1075.

[15]    L. E. Otterbein *et al.*, "Carbon monoxide has anti-inflammatory effects involving the

mitogen-activated protein kinase pathway," *Nat Med*, vol. 6, no. 4, pp. 422–428, Apr.

2000, doi: 10.1038/74680.

[16]    M. P. Soares and F. H. Bach, "Heme oxygenase-1: from biology to therapeutic potential,"

*Trends Mol Med*, vol. 15, no. 2, pp. 50–58, Feb. 2009, doi:

10.1016/j.molmed.2008.12.004.

[17]    Y. Naito, T. Takagi, and Y. Higashimura, "Heme oxygenase-1 and anti-inflammatory M2

macrophages," *Archives of Biochemistry and Biophysics*, vol. 564. Academic Press Inc.,

pp. 83–88, Dec. 15, 2014. doi: 10.1016/j.abb.2014.09.005.

[18]    S. C. Funes, M. Rios, J. Escobar-Vera, and A. M. Kalergis, "Implications of macrophage

polarization in autoimmunity," *Immunology*, vol. 154, no. 2. Blackwell Publishing Ltd,

pp. 186–195, Jun. 01, 2018. doi: 10.1111/imm.12910.

[19] S. O'Leary, M. P. O'Sullivan, and J. Keane, "IL-10 blocks phagosome maturation in

Mycobacterium tuberculosis-infected human macrophages," *Am J Respir Cell Mol Biol*,

vol. 45, no. 1, pp. 172–180, Jul. 2011, doi: 10.1165/rcmb.2010-0319OC.

[20] D. L. Costa *et al.*, "Pharmacological inhibition of host heme oxygenase-1 suppresses

mycobacterium tuberculosis infection in vivo by a mechanism dependent on T

lymphocytes," *mBio*, vol. 7, no. 5, Sep. 2016, doi: 10.1128/mBio.01675-16.

[21] D. L. Costa *et al.*, "Heme oxygenase-1 inhibition promotes IFNγ- and NOS2-mediated

control of Mycobacterium tuberculosis infection," *Mucosal Immunol*, vol. 14, no. 1, pp.

253–266, Jan. 2021, doi: 10.1038/s41385-020-00342-x.

[22] American Academy of Pediatrics Subcommittee on Hyperbilirubinemia, "Management of

hyperbilirubinemia in the newborn infant 35 or more weeks of gestation.," *Pediatrics*, vol.

114, no. 1, pp. 297–316, Jul. 2004, doi: 10.1542/peds.114.1.297.

[23] T. D. Burt, L. Seu, J. E. Mold, A. Kappas, and J. M. McCune, "Naive Human T Cells Are

Activated and Proliferate in Response to the Heme Oxygenase-1 Inhibitor Tin

Mesoporphyrin," *The Journal of Immunology*, vol. 185, no. 9, pp. 5279–5288, Nov. 2010,

doi: 10.4049/jimmunol.0903127.

[24] B. W. Allwood *et al.*, "Post-tuberculosis lung health: Perspectives from the First

International Symposium," in *International Journal of Tuberculosis and Lung Disease*,

International Union Against Tuberculosis and Lung Disease (The Union), Aug. 2020, pp.

820–828. doi: 10.5588/ijtld.20.0067.

[25] J. G. Pasipanodya *et al.*, "Pulmonary impairment after tuberculosis and its contribution to

TB burden," *BMC Public Health*, vol. 10, 2010, doi: 10.1186/1471-2458-10-259.

[26]    F. J. Leong, V. Dartois, and T. Dick, Eds., *A Color Atlas of Comparative Pathology of*

504            *Pulmonary Tuberculosis*, 1st Edition. CRC Press, 2010. doi: 10.1201/EBK1439835272.

505    [27]    K. D. Poss and S. Tonegawa, "Heme oxygenase 1 is required for mammalian iron

506            reutilization," 1997. [Online]. Available: www.pnas.org.

507    [28]    J. P. Juncos *et al.*, "Anomalous renal effects of tin protoporphyrin in a murine model of

508            sickle cell disease," *American Journal of Pathology*, vol. 169, no. 1, pp. 21–31, 2006, doi:

509            10.2353/ajpath.2006.051195.

510    [29]    C. E. Bunse *et al.*, "Modulation of heme oxygenase-1 by metalloporphyrins increases anti-

511            viral T cell responses," *Clin Exp Immunol*, vol. 179, no. 2, pp. 265–276, Feb. 2015, doi:

512            10.1111/cei.12451.

513    [30]    J. M. Cicchese *et al.*, "Dynamic balance of pro- and anti-inflammatory signals controls

514            disease and limits pathology," *Immunol Rev*, vol. 285, no. 1, pp. 147–167, Sep. 2018, doi:

515            10.1111/imr.12671.

516    [31]    M. K. K. Niazi, G. Beamer, and M. N. Gurcan, "Detecting and characterizing cellular

517            responses to Mycobacterium tuberculosis from histology slides.," *Cytometry A*, vol. 85,

518            no. 2, pp. 151–61, Feb. 2014, doi: 10.1002/cyto.a.22424.

519    [32]    N. V Serbina and J. L. Flynn, "Early Emergence of CD8 T Cells Primed for Production of

520            Type 1 Cytokines in the Lungs of Mycobacterium tuberculosis-Infected Mice," 1999.

521    [33]    I. E. A. Flesch and S. H. E. Kaufmann, "Activation of Tuberculostatic Macrophage

522            Functions by Gamma Interferon, Interleukin-4, and Tumor Necrosis Factor," 1990.

523    [34]    M. Safford *et al.*, "Egr-2 and Egr-3 are negative regulators of T cell activation," *Nat*

524            *Immunol*, vol. 6, no. 5, pp. 472–480, 2005, doi: 10.1038/ni1193.

525    [35]   K. A. Jablonski *et al.*, "Novel markers to delineate murine M1 and M2 macrophages,"

526            *PLoS One*, vol. 10, no. 12, Dec. 2015, doi: 10.1371/journal.pone.0145342.

527    [36]   S. Y. Li *et al.*, "Next-Generation Diarylquinolines Improve Sterilizing Activity of

528            Regimens with Pretomanid and the Novel Oxazolidinone TBI-223 in a Mouse

529            Tuberculosis Model," *Antimicrob Agents Chemother*, vol. 67, no. 4, Apr. 2023, doi:

530            10.1128/aac.00035-23.

531    [37]   C. R. Scharn *et al.*, " Heme Oxygenase-1 Regulates Inflammation and Mycobacterial

532            Survival in Human Macrophages during Mycobacterium tuberculosis Infection ," *The*

533            *Journal of Immunology*, vol. 196, no. 11, pp. 4641–4649, Jun. 2016, doi:

534            10.4049/jimmunol.1500434.

535    [38]   J. Grosset, C. Truffot-Pernot, C. Lacroix, and B. Ji, "Antagonism between isoniazid and

536            the combination pyrazinamide-rifampin against tuberculosis infection in mice.,"

537            *Antimicrob Agents Chemother*, vol. 36, no. 3, pp. 548–51, Mar. 1992, doi:

538            10.1128/AAC.36.3.548.

539    [39]   Z. Ahmad *et al.*, "Comparison of the 'Denver regimen' against acute tuberculosis in the

540            mouse and guinea pig.," *J Antimicrob Chemother*, vol. 65, no. 4, pp. 729–34, Apr. 2010,

541            doi: 10.1093/jac/dkq007.

542    [40]   A. M. Cooper, "Cell-mediated immune responses in tuberculosis," *Annual Review of*

543            *Immunology*, vol. 27. pp. 393–422, 2009. doi: 10.1146/annurev.immunol.021908.132703.

544    [41]   J. L. Flynn and J. Chan, "Immunology of tuberculosis.," *Annu Rev Immunol*, vol. 19, pp.

545            93–129, 2001, doi: 10.1146/annurev.immunol.19.1.93.

546    [42]   W. Li *et al.*, "CD38: A Significant Regulator of Macrophage Function," *Frontiers in*

547            *Oncology*, vol. 12. Frontiers Media S.A., Feb. 17, 2022. doi: 10.3389/fonc.2022.775649.

548    [43]    F. Ahmad *et al.*, "Macrophage: A Cell With Many Faces and Functions in Tuberculosis,"

549            *Frontiers in Immunology*, vol. 13. Frontiers Media S.A., May 06, 2022. doi:

550            10.3389/fimmu.2022.747799.

551    [44]    A. Khan *et al.*, "Human M1 macrophages express unique innate immune response genes

552            after mycobacterial infection to defend against tuberculosis," *Commun Biol*, vol. 5, no. 1,

553            Dec. 2022, doi: 10.1038/s42003-022-03387-9.

554    [45]    Food and Drug Administration and GASTROINTESTINAL DRUGS ADVISORY

555            COMMITTEE and the PEDIATRIC ADVISORY COMMITTEE, "Stannsoporfin -

556            Proposed for the treatment of neonates greater than or equal to 35 weeks of gestational age

557            with indicators of hemolysis who are at risk of developing severe hyperbilirubinemia,"

558            May 2018. Accessed: Aug. 01, 2023. [Online]. Available:

559            https://fda.report/media/112936/FDA-Briefing-Information-for-the-May-3--2018-Joint-

560            Meeting-of-the-Gastrointestinal-Drugs-Advisory-Committee-and-the-Pediatric-Advisory-

561            Committee.pdf

563    **Figures:**
564

**Figure 1: HO-1 inhibitors increase the antitubercular activity of a novel MDR regimen in a**

**mouse model of chronic TB.**

A) Experimental design of mouse experiments. Each line represents a different treatment group

and length of treatment. BALB/c mice were aerosol-infected with ~100 bacilli of Mtb H37Rv.

Treatment was initiated 4 weeks after aerosol infection. B) CFU means ± standard deviation (SD)

of lung bacillary burden after treatment initiation (week 0). Scatterplot of lung mycobacterial

burden after 2 weeks C), 4 weeks D) or 6 weeks E) of treatment. Each point represents the CFU

per mouse lung from individual, the error bars represent the mean ± SD. RHZ:

rifampin/isoniazid/pyrazinamide; SPaO: TBAJ-876/pretomanid/TBI-223; SnPP: tin-

protoporphyrin IX; SnMP: stannsoporfin; CFU: colony-forming units. For panels B-E, each

experimental group consisted of four to five mice. For panels C-E, statistical analysis was

performed by an ordinary one-way ANOVA followed by Dunnett's correction to control for

multiple comparisons and Bonferroni correction for treatment groups with the same antibiotic

backbone. * = $P < 0.05$; ** = $P < 0.01$; *** = $P < 0.001$; **** = $P < 0.0001$; ns: not significant.

**Figure 2: SnMP adjunctive therapy does not exacerbate gross lung inflammation.**

A, B, C) Scatterplot of lung/body weight ratio. D) Image of the right lung at time of harvest after

4 weeks of treatment. Scale bar = 1 cm. E) Hematoxylin and eosin staining of left lung medial

apex to base sections demonstrating the histopathology of different treatment groups after 4

weeks of treatment. Top row are representative histopathology images of the groups. Bottom row

are areas of inflammation within the images above (magnification of the black boxes above). F)

Quantitative analysis of histopathology after 4 weeks of treatment for selected groups. g: grams;

RHZ: rifampin/isoniazid/pyrazinamide; SPaO: TBAJ-876 (S)/pretomanid (Pa)/TBI-223 (O);

SnPP: tin-protoporphyrin IX; SnMP: stannsoporfin. For panels A-C and F, each point represents

data from individual mice, each experimental group consisted of four to five mice, and the bar

represents the means ± SD. Statistical analysis was performed by an ordinary one-way ANOVA

followed by Tukeys multiple comparison test for panels A-C and for panel F, Welch's ANOVA

test was conducted. ns: not significant; ** = $P < 0.001$; *** = $P < 0.001$; **** = $P < 0.0001$.

**Figure 3: Genes encoding M1 macrophage markers were upregulated in the lungs after two**

**weeks of treatment with SnMP.**

RT-cPCR was performed for the expression of the following genes in snap-frozen lung

homogenates: A) *Nos2*; B) *Ifn-γ*; C) *Tnf-α*; D) *Cd38*; E) *Egr2*. The dotted line represents the

uninfected normalization of mRNA expression, SPaO: TBAJ-876 (S)/Pretomanid (Pa)/TBI-223

(O); SnMP: stannsoporfin. Each bar represents a treatment group consisting of three to five mice.

Data represent individual data points with means ± SD of the results. Panel A was analysed by

the Welch's ANOVA test while panels B-E were tested by an ordinary one-way ANOVA

followed by Tukeys multiple comparison test. ns: not significant. * = $P < 0.05$; ** = $P < 0.001$;

*** = $P < 0.001$, **** = $P < 0.0001$.

**Figure 4: Adjunctive treatment with SnMP did not help reduce relapse rates after 6 weeks**

**of treatment.**

A) Experimental design of mouse experiment. Each line represents a different treatment group

and length of treatment. B) Lung bacillary burden after 6 weeks of treatment. Relapse study

statistics containing the % relapsed animals, and animals above a 3 $\log_{10}$ CFU after completing 5

C) or 6 D) weeks of treatment and a three month wait period (+ 3M). W0= Week 0, W5= Week

5, W6= Week 6, SPaO: TBAJ-876(S)/Pretomanid(Pa)/TBI-223(O), Stannsoporfin: SnMP,

SnMP5: SnMP 5 mg/kg, SnMP10: SnMP 10 mg/kg. Data in panel B represent four to five mice

per treatment group. Statistical analysis was performed by an ordinary one-way ANOVA

followed by Tukeys multiple comparison test. Data in panel C and D represent nine to twenty-

two mice per treatment group. * = $P < 0.01$, ** = $P < 0.001$; **** = $P < 0.0001$, ns: not

significant.

**A**

BALB/C Female
6-8 Weeks

Infected with
aerosolized *Mtb*
H37Rv

▲ SPaO    ▲ SPaO + SnMP10
■ W0 UNT    ▲ SPaO + SnMP5

Groups kept for relapse endpoint

Start of
treatment:
W0

W5
Harvest

W6
Harvest

W5 + 3M
Harvest

W6 + 3M
Harvest

**B**

Lung Bacterial Burden
($log_{10}$ CFU)

****

**

ns

W0 UNT   SPaO   SPaO + SnMP5   SPaO + SnMP10

**6 weeks of treatment**

**C**

| 5 weeks + 3M group | | |
|---|---|---|
| Treatment | Animals Relapsed (%) | % Animals above 3 $log_{10}$ CFU |
| SPaO | 9/9 (100) | 100 |
| SPaO + SnMP5 | 20/22 (90) | 80 |
| SPaO + SnMP10 | 18/18 (100) | 61 |

**D**

| 6 weeks + 3M group | | |
|---|---|---|
| Treatment | Animals Relapsed (%) | % Animals above 3 $log_{10}$ CFU |
| SPaO | 6/9 (66) | 66 |
| SPaO + SnMP5 | 13/17 (76) | 46 |
| SPaO + SnMP10 | 12/18 (66) | 33 |

| Regimen | Mean lung $\log_{10}$ CFU count (±SD) at: | | | | |
|---|---|---|---|---|---|
| | dpi | W0 | W2 | W4 | W6 |
| Untreated time course | $3.19 \pm 0.04$ | $6.59 \pm 0.10$ | | | |
| Vehicle | | | $6.40 \pm 0.25$ | $6.55 \pm 0.22$ | $6.42 \pm 0.54$ |
| SnPP | | | $5.52 \pm 0.29$ | $5.30 \pm 0.30$ | $5.30 \pm 0.19$ |
| SnMP | | | $5.73 \pm 0.22$ | $5.36 \pm 0.31$ | $5.98 \pm 0.06$ |
| RHZ | | | | $4.17 \pm 0.30$ | $3.02 \pm 0.30$ |
| RHZ + SnPP | | | | $4.11 \pm 0.23$ | $3.09 \pm 0.16$ |
| RHZ + SnMP | | | | $4.01 \pm 0.04$ | $3.00 \pm 0.12$ |
| SPaO | | | $4.75 \pm 0.17$ | $3.18 \pm 0.45$ | |
| SPaO + SnPP | | | $4.64 \pm 0.13$ | $2.67 \pm 0.20$ | |
| SPaO + SnMP5 | | | $4.68 \pm 0.07$ | $2.49 \pm 0.25$ | |
| Untreated relapse | $2.52 \pm 0.10$ | $6.18 \pm 0.11$ | | | |
| SPaO | | | | | $1.49 \pm 0.27$ |
| SPaO + SnMP5 | | | | | $1.47 \pm 0.30$ |
| SPaO + SnMP10 | | | | | $0.71 \pm 0.23$ |

**Table 1: Lung CFU count for time course and relapse study.**
dpi = day post infection; W0 = week 0 (start of treatment); W2 = Week 2; W4 = Week 4; W5 = Week 5; W6 = Week 6

| Gene name | Forward primer 5' to 3' | Reverse primer 5' to 3' | Reference |
|---|---|---|---|
| β-actin | AGCTGCGTTTTACACCCTTT | AAGCCATGCCAATGTTGTCT | PMID: 32862202 |
| Nos2 | CGAAACGCTTCACTTCCAA | TGAGCCTATATTGCTGTGGCT | PMID: 32862202 |
| Ifn-γ | CGATCTTGGCTTTGCAGCT | CCTTTTTCGCCTTGCTGTTG | PMID: 32862202 |
| Tnf-α | GCCTCTTCTCATTCCTGCTTG | CTGATGAGAGGGAGGCCATT | PMID: 21810933 |
| Cd38 | TCTCTAGGAAAGCCCAGATC | AGAAAAGTGCTTCGTGGTAG | PMID: 36351893 |
| Egr2 | CGCCACACCAAGATCCAC | AGCCCCAGGACCAGAGG | PMID: 22049075 |

**Table 2: RT-qPCR sequences for each primer pair.**