Combining molecular dynamics simulations and X-ray scattering techniques for the accurate treatment of protonation degree and packing of ionizable lipids in monolayers

Miriam Grava,[†] Mohd Ibrahim,[‡] Akhil Sudarsan,[‡] Julio Pusterla,[†] Julian Philipp,[¶] Joachim O. Rädler,[¶] Nadine Schwierz,[*,‡] and Emanuel Schneck[*,†]

†*Institute for Condensed Matter Physics, TU Darmstadt, Hochschulstraße 8, 64289 Darmstadt, Germany*

‡*Institute of Physics, University of Augsburg, Augsburg, Germany*

¶*Fakultät für Physik, Ludwig-Maximilians-Universität München (LMU), München, Germany*

E-mail: nadine.schwierz@physik.uni-augsburg.de; emanuel.schneck@pkm.tu-darmstadt.de

Abstract

The pH-dependent change in protonation of ionizable lipids is crucial for the success of lipid-based nanoparticles as mRNA delivery systems. Despite their widespread application in vaccines, the structural changes upon acidification are not well understood. Molecular dynamics simulations support structure prediction but require an a-priori knowledge of the lipid packing and protonation degree. The presetting of the protonation degree is a challenging task in the case of ionizable lipids since it depends on pH and on the local lipid environment and often lacks experimental validation.

Here, we introduce a methodology of combining all-atom molecular dynamics simulations with experimental total-reflection X-ray fluorescence and scattering measurements for the ionizable lipid Dlin-MC3-DMA (MC3) in POPC monolayers. This joint approach allows us to simultaneously determine the lipid packing and the protonation degree of MC3. The consistent parameterization is expected to be useful for further predictive modeling of the action of MC3-based lipid nanoparticles.

Introduction

The pH-dependent change in protonation of ionizable lipids is fundamental for endosomal release and hence the success of lipid nano-particles (LNPs) as mRNA delivery systems.[1] Ionizable lipids are already successfully used in FDA-approved drugs for the treatment of amyloidosis and in Moderna's vaccines against SARS-CoV-2.[2,3] Hence, the design of ionizable lipids with high transfection efficiencies is expected to play a key role in modern RNA therapeutics. Despite their importance, the microscopic structural changes upon acidification, preceding the endosomal escape of RNA, have not been resolved so far. Molecular dynamics (MD) simulations are particularly suited to provide atomistic insights, however, they require an a-priori knowledge of the lipid packing and the protonation degree. The presetting of the protonation degree is not always a simple task since it depends sensitively on the environment. Three competing interactions have to be considered: (i) conformational rearrangements of the lipids, (ii) partial dehydration of the ionizable group, and (iii) interactions with polar and charged lipids or ions. Due to these contributions, it is not surprising that ionizable lipids can have significant pK_a shifts. For example, the pK_a value of Dlin-MC3-DMA (MC3) was found to be shifted from about 9.4 at infinite dilution[4] to 6.4 inside a lipid nano-particle.[5]

Several theoretical models are available for pK_a predictions. Popular approaches are based on solving the linearized Poisson-Boltzmann equation[6] or generalized Born models.[7] Such simple models work well for simple systems but fail in more complicated cases where confor-

mational rearrangements become important. MD simulations include these conformational rearrangements and, when combined with free energy methods such as free energy perturbation[8] or thermodynamic integration,[9] yield the free energy of deprotonation and hence the pK_a. Using implicit or explicit solvent, free energy simulations have been successfully applied to calculate pK_a shifts in proteins.[10,11] In addition, constant-pH MD simulations have been developed to overcome limitations of fixed protonation states in conventional MD simulations.[12,13] However, these methods are computationally expensive and their accuracy depends on the underlying force fields and the convergence of the sampling of all degrees of freedom. In particular in lipid systems, which show pH dependent structural phase transitions,[14] such convergence is difficult to achieve. In addition, the correct packing of the lipids is crucial. To correctly reproduce the lateral structure, the force fields of all components including ionizable lipids, uncharged lipids, and the water model have to be accurate. Shortcomings of the force fields, for instance in reproducing the experimental surface tension of water, will lead to deviations in the packing such that simulations can fail to reproduce experimental surface pressure-area isotherms in simple monolayer systems.[15–18] Consequently, consistent methods are required to correctly assign the protonation degree and the packing of lipids to yield robust and meaningful results. In this respect the combination of all-atom MD simulations and X-ray scattering techniques offer promising perspectives, as the latter are uniquely suited to determine the packing of lipid layers[19–21] and the protonation degree[22] by quantifying the interfacial counterion excess.[23,24]

The aim of this current work is to introduce a methodology of combining total reflection X-ray scattering and X-ray fluorescence experiments to yield a reliable prediction as benchmark for consistent modeling. We focus on a lipid monolayer composed of the phospholipid POPC and the transfection lipid MC3, one of the most widely used ionizable lipids due to its high transfection efficiency.[25] These simple monolayers, bound to the air/water interface at controlled packing density, are an ideal starting point since they allow us to quantitatively compare the results from all-atom MD simulations and scattering experiments. The joint

approach allows us to optimize the protonation degree and the lipid packing. In turn, the simulations yield a detailed insight into the layer structure at the molecular level.

Materials and methods

Chemicals

The transfection lipid DLin-MC3-DMA (MC3, see Fig. 1) was purchased from MedChem-Express (Monmouth Junction, NJ, USA). The phospholipid 1-palmitoyl-2-oleoyl-glycero-3-phosphocholine (POPC, see Fig. 1) was purchased from Sigma-Aldrich (Merck KGaA, Germany). Potassium bromide (KBr), Hydrobromic acid (HBr) 48 wt.% in H_2O and Tris(hydroxymethyl)-aminomethan (Tris-Base) were of analytical grade, purchased also from Sigma-Aldrich (Merck KGaA, Germany) and used without further purification. Chloroform (purity $\geq$ 99%, anhydrous) and methanol (purity $\geq$ 99.9%) were purchased from Sigma-Aldrich (Merck KGaA, Germany) and used as received.

Sample preparation

The mixture of MC3-POPC (20:80 mol%) was prepared by mixing the two dry components and subsequently dissolving them at a concentration of (1 mg/ml) in chloroform-methanol (7:3 v/v). This solution was then spread onto an air-aqueous interface. The resulting monolayer film was laterally compressed until reaching a final lateral pressure of $\pi = 30$ mN/m while in some cases recording isotherms (see Fig. S1 in the Supporting Information). Two solutions were used as subphases for different experiments: KBr 2 mM Tris 5 mM pH 7.5 and KBr 2mM Tris 5mM pH 5. The corresponding pHs were reached by adding few drops of a 1M HBr solution to the appropriate amounts of Tris-Base. The resulting Br^- concentration after titration was 5.8 mM for pH 7.5 and 6.8 mM for pH 5 (see Supporting Information for further details).

Grazing-incidence X-ray scattering techniques

Grazing-incidence X-ray scattering experiments (GIXOS and TRXF, see below) were carried out at the beamline P08 at storage ring PETRA III of Deutsches Elektronen-Synchrotron (DESY, Hamburg, Germany), with the settings described in references,[26,27] from which the following text is partially reproduced. The Langmuir trough (Riegler & Kirstein, Potsdam, Germany) was located in a hermetically sealed container with Kapton windows, and the temperature was kept at 20 °C by a thermostat. The container was constantly flushed with a stream of humidified helium (He) to prevent air scattering and the generation of reactive oxygen species. The synchrotron X-ray beam was monochromatized to a photon energy of 15 keV, corresponding to a wavelength of $\lambda = 0.827$ Å. The incident angle was adjusted to $\theta_i = 0.07$ °, slightly below the critical angle of total reflection, $\theta_c = 0.082$ °. A ground glass plate was placed less than 1 mm beneath the illuminated area of the monolayer in order to reduce mechanically excited surface waves. The beam footprint on the water surface was 1 mm x 60 mm as imposed by the incident beam optics.

Under total-reflection conditions, an X-ray standing wave (SW) is formed at the air/water interface. The penetration depth Λ of its evanescent tail into the aqueous hemispace is a function of the angle of incidence θ_i[28] and for the incident angle used is $\simeq 8$ nm. The exact shape $\phi(z)$ of the SW intensity along the vertical position z (for a given incident angle) follows from the interfacial electron density profile $\rho_e(z)$, and can be calculated as described previously.[21,29]

Grazing-incidence X-ray off-specular scattering (GIXOS)

Analogous to conventional X-ray reflectometry, GIXOS allows reconstructing the interfacial electron density profile (i.e., the laterally-averaged structure of the lipid layer in the direction perpendicular to the surface) from the q_z-dependent scattering intensity, however at fixed incident angle. The details of this technique are described in reference.[20,30,31] As explained more briefly in Kanduc et al.,[32] the q_z-dependence of the diffuse scattering intensity $I(q_{xy} \neq$

$0, q_z)$ recorded at low-enough yet finite q_{xy} ("out of the specular plane) with the help of a narrow slit contains information equivalent to that of the conventional reflectivity $R(q_z)$ and can be transformed as:

$$I(q_{xy} \neq 0, q_z) = I(q_z) = R(q_z)\frac{V(q_z)}{R_{\mathrm{F}}(q_z)} \tag{1}$$

to good approximation, where $R_{\mathrm{F}}(q_z)$ the reflectivity of an ideal interface between the two bulk media and $V(q_z)$ is the Vineyard function.[33] The approximation is based on the assumption of conformal topographic roughness of all surfaces, which is justified for a lipid monolayer subject to capillary wave roughness. In the present work, the GIXOS signal was measured at $q_{xy} = 0.04$ Å^{-1}.

Like conventional reflectivity curves, GIXOS curves can be computed on the basis of an assumed interfacial electron density profile $\rho_{\mathrm{e}}(z)$, via the phase-correct summation of all reflected and transmitted partial waves occurring at the density gradients.[21,29] Here, we either modeled $\rho_{\mathrm{e}}(z)$ as a set of rough slabs representing the tail and headgroup regions of the monolayer with adjustable thickness, electron density, and roughness parameters[27,31,32] or extracted $\rho_{\mathrm{e}}(z)$ directly from the MD simulations. In all cases, $\rho_{\mathrm{e}}(z)$ was then discretized and the corresponding q_z-dependent reflectivities $R(q_z)$ were calculated from the Fresnel reflection laws at each slab-slab interface using the iterative recipe of Parratt[34] or the equivalent Abebe's matrix method[35] and subsequently multiplied with $V(q_z)/R_{\mathrm{F}}(q_z)$ to obtain the theoretical GIXOS signal. For fits to the experimental data, a constant scale factor and a constant intensity background was applied to the modeled intensities, and the fitting range was limited to the consensus range of validity ($q_z > 3 \cdot q_z^{\mathrm{C}}$, where q_z^{C} is the q_z-value at the critical angle of total reflection).[31]

Total-reflection X-ray fluorescence (TRXF)

The fluorescence signal induced via photoelectric ionization by the X-ray beam under total reflection conditions was recorded with an Amptek X-123SDD detector (Amptek, Bedford, USA). The detector was placed almost parallel to the water surface and perpendicular to

the X-ray beam axis, in order to keep elastic and Compton scattering into the detector as low as possible. The center of the detector view angle was set to coincide with the beam footprint position on the water surface.

The fluorescence intensity I_F emitted by the Br^- counterions present at the interface is determined by their interfacial depth profile $c(z)$.[36] On a quantitative level, I_F is proportional to the spatial integral over the product of $c(z)$ and the known SW intensity profile $\phi(z)$ introduced above,

$$I_F = A \int_{-\infty}^{\infty} c(z)\phi(z)dz \tag{2}$$

where the prefactor A can be calibrated with a suitable reference. Here, we used the surface of a bare KBr solution as reference, for which $c(z)$ is known and can be considered constant.[21,36] Experimentally, I_F was obtained by fitting the intensity peak associated with the respective K_α or L_β emission lines of Br (at ≈ 12.0 keV and ≈ 1.5 keV) in the recorded fluorescence spectra with a Gaussian function. Constraints on the peak positions were imposed based on the tabulated line energies.[37] The standing wave $\phi(z)$ was calculated with a slab-model representation of the interfacial electron density profile,[21] whose parameters were previously obtained in fits to the GIXOS curves. Note that roughness can be neglected for the computation of $\phi(z)$ when q_z is low, as is the case under total reflection.[36]

Theoretical predictions of the protonation degree

Infinite dilution limit

The ionizable MC3 lipids are protonated according to the reaction

$$MC3^0 + H_3O^+ \leftrightarrow MC3^+ + H_2O \tag{3}$$

At infinite dilution of the lipids, the protonation degree is given by the HendersonHasselbalch equation

$$pH = \log \frac{1-\alpha}{\alpha} + pK_a. \tag{4}$$

for the protonation degree α, where we assume $pK_a = 9.47$ at infinite dilution.[4]

Self-consistent equation for the protonation degree

The pK_a is related to the equilibrium constant K_a by

$$pK_a = -\log K_a \tag{5}$$

At the monolayer/water interface, the protonation of MC3 lipids are influenced by the distribution of H_3O^+ ions and the equilibrium constant K_a is given by

$$K_a = \frac{1-\alpha}{\alpha} \cdot [H_3O^+]_{surface} \tag{6}$$

where $[H_3O^+]_{surface}$ is the concentration at the position of the ionizable group in the monolayer. The surface concentration is related to the bulk concentration via a Boltzmann distribution[38]

$$[H_3O^+]_{surface} = [H_3O^+]_{bulk} \cdot e^{-e\Phi_0/(k_B T)} \tag{7}$$

where e is the elementary charge, $k_B T$ the product of Boltzmann constant and temperature, and Φ_0 the electrostatic potential at the position of the ionizable group. Combining Eqs. 6-7 and using the definitions for pH and pK_a yields a self-consistent equation for the pH as a function of the protonation degree[39]

$$pH = \log \frac{1-\alpha}{\alpha} + pK_a - \frac{e\Phi_0}{k_B T \ln 10}. \tag{8}$$

The last term is the electrostatic contribution that takes the H_3O^+ distribution dependent on the surface potential into account. Without this term the ordinary equation for the acid deprotonation equilibrium in bulk solution is obtained (Eq. 3).

The electrostatic potential $\Phi(z)$ is calculated from the Poisson Boltzmann equation at a planar interface

$$\epsilon_0 \epsilon_r \frac{d^2\Phi(z)}{dz^2} = -\sum_{i=\pm} q_i c_0 e^{-q_i \Phi(z)/k_B T} \tag{9}$$

Here, z is the distance perpendicular to the surface, $q_\pm = \pm e$ is the ion charge in a 1:1 electrolyte, c_0 is the bulk salt concentration, ϵ_0 is the dielectric constant of vacuum, ϵ_r is the relative dielectric constant of water, and $\Phi(z)$ is the electrostatic potential. For the constant charge boundary condition, we obtain the analytical solution

$$\Phi(z) = \frac{2k_B T}{e} \ln \frac{1 + \Gamma_0 e^{-\kappa z}}{1 - \Gamma_0 e^{-\kappa z}} \tag{10}$$

where Γ_0 is related to the surface potential via

$$\Gamma_0 = \tanh\left(\frac{e\Phi_0}{4k_B T}\right) \tag{11}$$

and κ^{-1} is the Debye screening length

$$\kappa^{-1} = \sqrt{\frac{\epsilon_0 \epsilon_r k_B T}{e^2 c_0}} \tag{12}$$

The surface potential $\Phi_0 = \Phi(0)$ is obtained from

$$\Phi_0 = \frac{2k_B T}{e} \text{arcsinh}\left(\frac{\sigma}{\sqrt{8k_B T c_0 \epsilon_0 \epsilon_r}}\right) \tag{13}$$

The surface charge density σ is related to the area per lipid via $\sigma = \alpha e f_{MC3}/A_{lip}$, where f_{MC3} is the fraction of MC3 among all lipids and e is the elementary charge. The experimentally obtained ion concentrations of $c_0 = 6.8$ mM for $\alpha = 67.5$ % and $c_0 = 5.8$ mM for $\alpha = 15$ %

were used for the theoretical predictions. In the self-consistent approach, the protonation degree depends on the surface potential Φ_0. Clearly, Φ_0 depends on σ and hence on the protonation degree α and the area per lipid A_{lip}. In our approach, Φ_0 and σ were obtained from the results of the final simulations with optimized α and A_{lip} based on the experiments. σ was obtained by dividing the total number of protonated MC3 lipids in one leaflet of the monolayer by the area of the leaflet. The values of σ calculated at $\alpha = 67.5$ % and $\alpha = 15$ % were 0.029 Cm^{-2} and 0.006 Cm^{-2}, respectively. With these values, we obtain surface potentials of $\Phi_0 = 92.5$ mV and $\Phi_0 = 33.5$ mV at $\alpha = 67.5$ % and $\alpha = 15$ %, respectively.

MD Simulations

We simulated the MC3-POPC monolayers (20:80 mol %) using the Gromacs simulation package (v-2021.5).[40] The monolayers were created by translating the leaflets of an equilibrated bilayer which in turn was created using the Mem-Gen webserver.[41] The AMBER Lipid 17 force field[42] was used to describe the POPC lipids. For the cationic and neutral MC3 molecules we used our recently developed force fields.[43] These parameters have the advantage that they closely reproduce the structure of MC3 in lipid layers as judged by neutron reflectometry experiments. In addition, the parameters are compatible with the AMBER force field family. Ions were described using the Mamatkulov-Schwierz force field parameters[44] and the TIP3P water model was used.[45] The systems contained 160 POPC and 40 MC3 lipids per monolayer, and 60 water molecules per lipid. Chloride ions were added to neutralize the positive charge of the protonated MC3 lipid molecules. The systems were energy minimized using a gradient descent algorithm. The temperature of the systems was maintained at 293.15 K using the velocity rescaling thermostat with stochastic term[46] and a time constant of 1.0 ps. For the NPT simulations, semi-isotropic Parrinello-Rahman[47] was used. The Van der Waals interaction potential was cut off and shifted to zero at 1.2 nm. Electrostatic interactions were cut-off at 1.2 nm and the particle mesh ewald (PME) method[48]

with a fourier grid spacing of 0.12 nm was used to evaluate the long-range electrostatics part. All bonds involving hydrogens were constrained using the LINCS algorithm.[49] A timestep of 2.0 fs was used to integrate the equations of motion. Analysis was performed using Gromacs inbuilt routines and with MDAnalysis[50] and trajectories were visualized with VMD.[51]

To obtain the desired area per lipid, the in-plane components P_{xx} and P_{yy} of the pressure tensor were varied in 11 steps from -50 bar to -34 bar. These simulations had durations of 100 ns. A semi-isotropic barostat was employed, such that $P_{xx} = P_{yy}$. The box size in z-direction, $L_z = 20$ nm, was kept constant. The surface tension γ can be deduced from the pressure tensor components[52] as

$$\gamma = \frac{L_z}{n} \left[P_{zz} - \frac{P_{xx} + P_{yy}}{2} \right] \tag{14}$$

where $n = 2$ is the number of interfaces in the simulation box.[53] The value of γ was directly obtained using the Gromacs `gmx energy` routine. The covered range of in-plane pressure tensor components corresponds to a surface tension range of 34 mN/m $\leq \gamma \leq$ 50 mN/m. The surface pressure is calculated from $\pi = \gamma_0 - \gamma$, where $\gamma_0 = 58$ mN/m is the surface tension of TIP3P water at 293.15 K as obtained through extrapolation of the data by Vega et al.[52] The last 50 ns of each of these simulation at given value of surface pressure was used to obtain the electron density, from which the GIXOS curves were evaluated using Eq. 1. The Refnx python package[54] was used to obtain the reflectivity profiles ($R(q_z)$ and $R_F(q_z)$ in Eq. 1). The simulations which best reproduced the experimental GIXOS curves at pH 5.0 and pH 7.5 for $\alpha = 1.0$ had the parameters summarized in Table 1. The simulations with experimentally validated values of A_{lip} and α were performed for 500 ns in the NVT ensemble and had the parameters summarized in Table 2.

Table 1: Simulation parameters that best reproduce the experimental GIXOS curves for $\alpha = 1.0$.

pH	Fixed Parameters			Calculated Parameters				
	α	L_z [nm]	P_{xx}/P_{yy} [bar]	P_{zz} [bar]	L_x/L_y [nm]	A_{lip} [nm^2]	γ [mN/m]	π [mN/m]
5.0	1.0	20.0	46.0	-0.36 ± 0.72	12.10 ± 0.02	0.73 ± 0.002	45.9 ± 0.9	12.1 ± 0.9
7.5	1.0	20.0	48.0	1.26 ± 1.0	12.25 ± 0.02	0.75 ± 0.003	49.8 ± 1.2	8.2 ± 1.2

Table 2: Parameters used for the simulations with experimentally validated area per lipid and protonation degree.

pH	Fixed Parameters				Calculated Parameters				
	α	L_x/L_y [nm]	L_z [nm]	A_{lip} [nm^2]	P_{xx} [bar]	P_{yy} [bar]	P_{zz} [bar]	γ [mN/m]	π [mN/m]
5.0	0.675	12.10	20.0	0.73	-47.8 ± 0.35	-48.2 ± 0.4	-0.36 ± 0.46	47.6 ± 0.6	10.4 ± 0.6
7.5	0.150	12.25	20.0	0.75	-48.8 ± 0.6	-49.2 ± 0.8	-0.42 ± 0.56	49.1 ± 0.8	8.9 ± 0.8

Results and Discussion

The lipid monolayers contain 80 mol% of the charge-neutral zwitterionic phospholipid POPC and 20 mol% of the ionizable lipid MC3 (Fig. 1). Dependent on the pH, the ionizable MC3 can be neutral (MC3^0) or positively charged (MC3$^+$) such that the monolayer contains up to three lipid species.

Figure 1: Molecular structures of the phospholipid POPC and the ionizable lipid MC3.

Our aim is to elucidate the influence of pH on the properties of the monolayers at two different pH values, pH 7.5 and pH 5.0, representing the extracellular and the endosomal

milieu, respectively. However, to perform simulations two important parameters have to be chosen. The first one is the lateral monolayer packing in terms of the average area per lipid, A_{lip}. It should match the experimental one at a lateral pressure of $\pi = 30$ mN/m, a value representative of the packing in lipid bilayers.[55] The second one is the protonation degree α of MC3, which should be consistent with the experimental value at given pH conditions. To correctly assign A_{lip} and α, we obtain complementary information from two synchrotron-based grazing-incidence X-ray scattering techniques, namely grazing-incidence X-ray off-specular scattering (GIXOS) and total-reflection X-ray fluorescence (TRXF).[36,56,57] While GIXOS, like conventional reflectometry, precisely reveals the monolayer thickness[27] (albeit with limited insights into the conformations and lateral distributions of the different molecular species), TRXF is highly sensitive to the interfacial accumulation of counterions at charged monolayers[23,24,36] and can therefore serve to determine protonation degrees.[22] As will be shown later, the accuracy in the experimental determination of α is further enhanced by including the counterion distribution from the simulations.

Figure 2: Methodology of combining all-atom molecular dynamics simulations with experimental total reflection X-ray scattering and X-ray fluorescence for consistent modeling of the area per lipid and the protonation degree.

Fig. 2 outlines the methodology to obtain consistent values for A_{lip} and α. Initially, α is estimated based on the Henderson-Hasselbach equation which predicts $\alpha = 1$ for both pH values (Fig. 3). Note that the self-consistent equation yields an improved estimate but requires the surface charge density, which is not known a-priori. In a second step, we performed simulations with the assumed α value while systematically varying A_{lip} to identify the values that match the layer thickness obtained in the GIXOS measurements at the two pH conditions. In the next step, the pH-dependent protonation degrees α are deduced from

the TRXF measurements while imposing the known values of A_{lip} as well as the shape of the counterion distribution found in the simulations. Finally, new simulations with the experimentally validated values of A_{lip} and α are performed for both pH conditions. In the following, we discuss the results from each step in detail.

Figure 3: Dependence of the protonation degree α on pH obtained from the Henderson-Hasselbach equation and the self-consistent equation using $\mathrm{pK_a} = 9.4.$[4] Dashed straight lines indicate the predictions for the two experimental pH values, 5.0 and 7.5.

Determination of the area per lipid

Fig. 4 A shows the GIXOS curves obtained experimentally with monolayers at $\pi = 30$ mN/m on aqueous subphases with pH 7.5 (top) and with pH 5.0 (bottom). The subphases additionally contained defined concentrations of $\mathrm{Br^-}$ and $\mathrm{K^+}$ ions, as required for the simultaneous TRXF measurements discussed further below. The curves contain information on the electron density profiles of the monolayers, where the overall layer thicknesses are mainly encoded in the q_z-positions of the intensity minima. The lines superimposed to the experimental data points are the simulated GIXOS curves computed from the electron density profiles obtained in simulations. Based on the Henderson-Hasselbalch equation (Eq. 3) the protonation degree is $\alpha \approx 1$ (Fig. 3). However, significant deviations from the experimental curve (dashed line in Fig. 4 A) are obtained with simulations at the experimental surface pressure corresponding to a surface tension of $\gamma = \gamma_0 - \pi = 28$ mN/m, with $\gamma_0 = 58$ mN/m for TIP3P water

(see Methods section). Clearly, the packing of the lipids and hence A_{lip} is not reproduced. This deviation is likely caused by the TIP3P water model which, like most water models, does not yield the correct surface tension of water.[18,52] In order to provide improvement, we systematically vary A_{lip} in the simulations to obtain the best-matching value for each pH, characterized by the minimum χ^2 deviation (Fig. S5) from the experimental GIXOS data. The simulated GIXOS curves obtained after optimization (solid lines) reproduce the experimental data well, demonstrating that the simulations properly capture the overall structure of the monolayers.

For pH 7.5 and pH 5.0, the best-matching values are $A_{\text{lip}} = 0.75$ nm^2 and $A_{\text{lip}} = 0.73$ nm^2, respectively, which is in the typical range for lipid bilayers the fluid phase.[58] The associated surface pressures in the simulations are as low as $\pi \approx 12$ mN/m (pH 5.0, see Table 1) and $\pi \approx 8$ mN/m (pH 7.5, see Table 1) and deviate by a factor of ≈ 3 from the experimental value. Interestingly, the agreement is much better on the level of the surface tension, where the simulations yield $\gamma \approx 46$ mN/m (pH 5.0, see Table 1) and $\gamma \approx 50$ mN/m (pH 7.5, see Table 1), compared to the experimental value of $\gamma = \gamma_0 - \pi = 42$ mN/m, where $\gamma_0 = 72$ mN/m. As seen in Tables 1 and 2, the protonation degree of MC3 only has a minor influence on γ and π for a fixed area per lipid. In conclusion, the lateral density in terms of A_{lip} appears to be the most robust parameter for the experimental validation of MD simulations of lipid monolayers.

The electron density profile $\rho_{\text{e}}(z)$ and the counterion density profile $c(z)$ obtained in the simulations for $A_{\text{lip}} = 0.73$ nm^2 are shown in Fig. 4 C. The profiles are consistent with earlier reports on fluid lipid monolayers at comparable lateral pressures,[19] however the electron density maximum representing the lipid headgroups is slightly less pronounced than for commonly studied phospholipid layers, because MC3 has a more compact headgroup without the electron-rich P atom.

Figure 4: (A) Comparison of experimental GIXOS curves with the simulated ones for the optimized values of A_{lip}. The system consists of 20 % MC3$^+$ and 80 % POPC. The curves at different pH values are vertically shifted for clarity. The dashed black line is obtained by using the same surface pressure as in the experiments ($\pi = 30$ mN/m) and corresponds to $A_{\text{lip}} = 0.63$ nm^2. The solid blue lines correspond to the simulations with optimized area per lipid ($A_{\text{lip}} = 0.75$ nm^2 for pH = 7.5 and $A_{\text{lip}} = 0.73$ nm^2 for pH = 5.0. (B) Snapshot from a simulation with $A_{\text{lip}} = 0.63$ nm^2 and $\alpha = 1$ after 100 ns of equilibration. All other run parameters are summarized in Table 1. MC3$^+$ is represented in yellow with the head group nitrogen highlighted by yellow spheres. POPC is shown in white and chloride ions in cyan color. (C) Corresponding electron density profile $\rho_{\text{e}}(z)$. The plot also shows the counterion density profile $c(z)$ obtained in the simulations (data points). The solid line superimposed is an analytical fit with Eq. 15.

Determination of the protonation degree

The pH-dependent protonation degree of MC3 was determined experimentally through measurements of the interfacial excess of Br$^-$ counterions, Γ, which coincides with the density of positive net charges in the monolayer.[36] Br$^-$ rather than the more common Cl$^-$ was used as counterion in these experiments because the used TRXF setup was slightly more sensitive to heavier elements. Fig. 5 A shows the X-ray fluorescence spectra in a narrow energy range around the peak associated with the K_α emission line of Br. The spectra were obtained under total reflection conditions with the bare aqueous subphases and with lipid monolayers at $\pi = 30$ mN/m both at pH 5.0 and pH 7.5. The complete spectra are shown in the Supporting Information (Fig. S2). The intensities of the emission lines are much higher in the presence of the monolayer at both pH values, which reflects the accumulation of Br$^-$ ions at the interface required to compensate the positive charges of MC3$^+$. The L_β emission lines

behave consistently (see Supporting Information). The intensity is much higher at pH 5.0 than at pH 7.5, evidencing that the positive charge density is much higher at the lower pH and, thus, the protonation degree α. This observation immediately implies that α at pH 7.5 is much lower than 100 %, in contradiction to the estimation from the Henderson-Hasselbalch equation. The slight difference in the intensities observed with the bare subphases at the two pH values can be attributed to the slightly different concentration of Br^- as a result of pH titration with HBr (see Methods section).

Figure 5: (A) X-ray fluorescence spectra around the K_α emission peak of the Br^- counterions obtained with the bare aqueous subphases and with lipid monolayers at two pH values. (B) Illustration of the model used for data analysis (here pH 5.0): the two-slab representation of the experimentally obtained electron density $\rho_e(z)$, the standing wave intensity $\phi(z)$, and the counterion profile $c(z)$. For the sake of clarity, all curves are normalized to have maximal values of 1.

The following quantitative analysis takes the counterion profiles $c(z)$ from the MD simulations into account. This approach can be viewed as a refinement with respect to commonly used treatments of the counterion distribution in terms of thin adsorption layers[27,36] or related approximations.[59] Here, the simulation-based distributions were modeled as a (truncated) Gaussian function on top of the constant experimental bulk concentration c_0,

$$c(z) = c_0 + (c_{max} - c_0)e^{-(z-z_{max})^2/(2\sigma^2)} \qquad z \geq z_0 \qquad (15)$$

, where z_0 defines the interface between the tail and headgroup slabs. This functional form

was fitted to the numerical counterion profiles extracted from the simulations (see solid line in Fig. 4 C), yielding $z_{max} = z_0 + 0.56$ nm; $\sigma = 0.48$ nm for pH 5.0 and $z_{max} = z_0 + 0.53$ nm; $\sigma = 0.47$ nm for pH 7.5. With that, c_{max} remains as the only adjustable parameter, which is varied so as to match the measured fluorescence intensities I_F in the presence of the lipid monolayer and at the given pH, according to Eq. 2. For illustration, Fig. 5 B displays all ingredients required for this calculation: the two-slab model of the electron density $\rho_e(z)$, obtained from the experimental GIXOS curves as described in the Supporting Information (Fig. S3), the corresponding standing wave intensity $\phi(z)$, and the parameterized counterion profile $c(z)$. The counterion excess then simply follows as

$$\Gamma = \int_{z_0}^{\infty} [c(z) - c_0] \, dz. \tag{16}$$

We obtain $\Gamma = 0.185$ nm^{-2} for pH 5.0 and $\Gamma = 0.039$ nm^{-2} for pH 7.5. For a known counterion excess, the protonation degree α of MC3 follows as:

$$\alpha = \frac{\Gamma A_{lip}}{f_{MC3}} \tag{17}$$

where $f_{MC3} = 0.2$ is the MC3 mole fraction. With that, we obtain $\alpha = 67.5$ % for pH 5.0 and $\alpha = 14.5$ % for pH 7.5.

The obtained protonation degrees are significantly lower compared to the prediction from the Henderson-Hasselbalch equation. This is not surprising since the equation is valid at infinite dilution. In the monolayer system, interactions between polar lipids, charged lipids, ions and protons contribute to the protonation degree. The effect of the proton distribution is taken into account in the self-consistent equation 3. The protonation degrees predicted in this way are $\alpha = 99$ % for pH 5.0 and $\alpha = 68$ % for pH 7.5. Although these values provide a more meaningful estimate, they are still significantly higher than the ones experimentally observed and require an a-priori estimate of the surface charge density. To provide improvement, the interactions between the lipids in the monolayer as well as conformational rearrangement

and partial dehydration have to be taken into account. In particular the effect of partial dehydration is expected to have a significant effect since the uncharged headgroups are buried while the charged headgroups are solvent exposed (Fig. 6 A and B). In any case, reliable experimental measurements of the protonation degree are imperative for validation and improvement of theoretical predictions.

Finally, it is important to confirm that the simulations with the experimentally validated protonation degree still reproduce the experimental GIXOS curves and counterion profiles that were used in the validation procedure (Fig. 2). Otherwise additional iterations would be required. The results show that after only one iteration the results converged and the simulated GIXOS curves do not significantly respond to the updated α values (see Fig. S4 in the Supporting Information), and the peak in the counterion distribution still has the same position and width (see Fig. S7 in the Supporting Information).

Simulations with experimentally validated area per lipid and protonation degree

As the last step, MD simulations with the experimentally validated values of A_{lip} and α for the two pH values were performed, with the parameters summarized in Table 2. The correct protonation degrees were imposed by representing the MC3 molecules with suitable numbers of $MC3^0$ and $MC3^+$ variants. In practice, 6 and 27 out of the 40 MC3 molecules in each monolayer were represented with $MC3^+$ for pH 7.5 and pH 5.0, respectively, yielding $\alpha = 15.0$ % and $\alpha = 67.5$ %. These protonation degrees come as close as possible to the experimental values when considering that the numbers of lipids can only be varied in integer steps.

Fig. 6 A shows a snapshot from an equilibrated simulation with $\alpha = 67.5$ %, corresponding to pH 5.0. POPC matrix lipids are represented in white, $MC3^0$ in yellow, and $MC3^+$ in green. Panel B shows the normalized number density profiles ρ_{n} of the tails and headgroups of all lipid species and of water in the z-direction, i.e, perpendicular to the membrane plane.

The distribution of the positively charged headgroups of MC3$^+$ largely overlaps with that of POPC's zwitterionic headgroups, while the distribution of the uncharged and only weakly polar headgroups of MC3^0 is strongly shifted towards the hydrophobic region accommodating the chains. As shown in the Supporting information (Figure S8), consistent observations can be made at $\alpha = 15.0$ %, corresponding to pH 7.5. This behavior has been noticed earlier[43] and may provide the explanation for the small yet significant increase in A_{lip} when increasing the pH from 5.0 to 7.5 (see above). Namely, in contrast to the naive expectation that a reduction in the density of charged lipids would decrease lateral repulsion and thus the lipids' effective area requirement, the deep insertion of uncharged MC3^0 into the monolayer effectively increases the area requirement at high pH, where MC3^0 is more abundant.

Figure 6: Simulations with experimentally validated A_{lip} and α. (A) Side view snapshot of one of the monolayers with $\alpha = 67.5$ %. Colors: green (MC3^0), yellow (MC3$^+$) and gray (POPC). (B) Number density profiles $\rho_n(z)$ of water and of the head and tail groups of MC3^0, MC3$^+$, and POPC. Solid lines represent head groups and dashed lines represent tail groups. Dotted line: water. The top scale applies to MC3, the bottom scale to POPC and water.

We now turn to the in-plane (xy) distribution of the three lipid species. Fig. 7 A shows a top-view simulation snapshot of a monolayer after equilibration. Qualitatively, it is seen that both MC3^0 and MC3$^+$, depicted in green and yellow, respectively, distribute rather evenly over the membrane plane, which maximizes their average distance. This behavior is quantified with the help of the normalized in-plane radial distribution functions (RDFs)

of the headgroup N atoms, which are shown in panel B. Indeed, while the RDF between two POPC headgroups on approach oscillates around unity and even exhibits a pronounced next-neighbor peak at $r \approx 0.9$ nm due to dipole interactions,[60] the RDF between two $MC3^+$ headgroups systematically drops below unity on approach, albeit with a few weak oscillations. This drop in the RDF can be attributed to the electrostatic repulsion between the positively charged headgroups. The RDF between two $MC3^0$ drops below unity on approach, too, but like-charge repulsion obviously cannot be the reason because $MC3^0$ carry no net charge. One possible reason might be that the penetration of an $MC3^0$ headgroup into the hydrophobic region locally leads to molecular rearrangements that oppose to some extent the penetration of another $MC3^0$ nearby.

Another qualitative difference between the RDF of two $MC3^0$ and the other two RDFs shown in Fig. 7 B is its finite value at $r = 0$. Apparently, two $MC3^0$ headgroups can share the same in-plane (xy) coordinates because they are somewhat mobile in the third dimension (z). This is in stark contrast to POPC and $MC3^+$, whose headgroups are firmly pinned in z direction to the hydrophilic/hydrophobic interface and thus sterically repel each other laterally in the limit of small r. This interpretation is further corroborated by the mixed RDFs shown in Fig. 7 C. Strikingly, the mixed RDFs of $MC3^0$ with POPC and of $MC3^0$ with $MC3^+$ both merely exhibit a weak dip near $r = 0$, demonstrating that there is little steric exclusion because the headgroups of $MC3^0$ roam at a different "altitude" than those of POPC and $MC3^+$. In contrast, as seen from the mixed RDF of POPC with $MC3^+$, their headgroups mutually exclude each other sterically for small r and exhibit a pronounced next-neighbor peak at $r \approx 0.9$ nm due to attractive interactions between the positively charged $MC3^+$ headgroups and the negatively charged phosphate groups in the PC headgroups.[61]

Figure 7: Simulations with experimentally validated A_{lip} and α. (A) Top view snapshot of one of the monolayers with $\alpha = 67.5$ %. Colors: green (MC3^0), yellow (MC3$^+$) and gray (POPC). (B) In-plane radial distribution functions (RDF) between the head group nitrogens of a same lipid species. (C) RDF between nitrogens of different lipid species.

Conclusions

Molecular dynamics simulations of ionizable lipids provide detailed insight into their structure but require an a-priori knowledge of the lipid packing and protonation degree. We have introduced a methodology of combining all-atom molecular dynamics simulations with experimental total reflection X-ray scattering and X-ray fluorescence to make a robust and reliable predictions of MC3 lipid area and degree of protonation at pH conditions that mimic the extracellular and endosomal milieus. The combined approach yields a protonation degree $\alpha = 14.5\%$ and an average area per molecule $A_{\text{lip}} = 0.73$ nm^2 at pH 7.5. At pH 5.0, it yields $\alpha = 67.5\%$ and $A_{\text{lip}} = 0.75$ nm^2. In both cases the value for the protonation degree differs significantly from the value expected from the Henderson-Hasselbalch equation. Hence, the presetting of the protonation degree in the simulations based on simple theories can fail due to the interactions between polar lipids, charged lipids, ions and protons in complex lipid interfaces.

The simulations with optimized A_{lip} and α, obtained from the matching methodology, reveal that the headgroups of the uncharged MC3 enter deeply into the hydrophobic monolayer region while the charged MC3 headgroups are solvent-exposed, as seen from the density profiles

in Fig. 6. Lipid dehydration upon deprotonation is therefore expected to have a significant effect on the protonation degree and is likely the reason why theoretical predictions based on the Henderson-Hasselbach equation overestimate the protonation degree for these simple monolayer systems. In addition, the buried uncharged MC3 lipids are laterally almost uncorrelated from the more hydrophilic POPC headgroups and the charged MC3 lipids. This behavior is associated with a non-typical structural response of A_{lip} to pH variations, which may play a significant role for the endosomal escape and subsequent biological action of mRNA.

The joint experimental and simulation approach presented here provides a benchmark for consistent modeling. The results can be used to test theoretical models or advanced simulation methods such as free energy simulations or constant-pH simulations. The methodology is also suited to investigate changes in protonation and lipid packing in response to interactions with nucleic acids. Another challenge is to extend this methodology for predictions in three-dimensional lipid mesophases and lipid nanoparticles.

Acknowledgement

We acknowledge DESY (Hamburg, Germany), a member of the Helmholtz Association HGF, for the provision of experimental facilities. Parts of this research were carried out at PETRA III and we would like to thank Chen Shen for assistance in using P08. Beamtime was allocated for proposal I-11009842. We thank Gerald Brezesinski and Tetiana Mukhina for help with the synchrotron experiments. This research was supported by the Bundesministerium für Bildung und Forschung (BMBF) within the Röntgen-Ångström cluster project Medisoft, number 05K18EZA and 05K18WMA, and by the Emmy Noether program of the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), number 315221747. We acknowledge the GOETHE HLR for super-computing access. The authors gratefully acknowledge the scientific support and HPC resources provided by the Erlangen National

High Performance Computing Center (NHR@FAU) of the Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU) under the NHR project b119ee. NHR funding is provided by federal and Bavarian state authorities. NHR@FAU hardware is partially funded by the German Research Foundation (DFG), number 440719683.

CRediT author statement

If required

References

(1) Li, Z.; Carter, J.; Santos, L.; Webster, C.; van der Walle, C. F.; Li, P.; Rogers, S. E.; Lu, J. R. Acidification-Induced Structure Evolution of Lipid Nanoparticles Correlates with Their In Vitro Gene Transfections. ACS nano **2023**,

(2) Hou, X.; Zaks, T.; Langer, R.; Dong, Y. Lipid nanoparticles for mRNA delivery. Nature Reviews Materials **2021**, 6, 1078–1094.

(3) Miao, L.; Zhang, Y.; Huang, L. mRNA vaccine for cancer immunotherapy. Molecular Cancer **2021**, 20, 1–23.

(4) Carrasco, M. J.; Alishetty, S.; Alameh, M.-G.; Said, H.; Wright, L.; Paige, M.; Soliman, O.; Weissman, D.; Cleveland IV, T. E.; Grishaev, A.; others Ionization and structural properties of mRNA lipid nanoparticles influence expression in intramuscular and intravascular administration. Communications biology **2021**, 4, 956.

(5) Jayaraman, M.; Ansell, S. M.; Mui, B. L.; Tam, Y. K.; Chen, J.; Du, X.; Butler, D.; Eltepu, L.; Matsuda, S.; Narayanannair, J. K.; others Maximizing the potency of siRNA lipid nanoparticles for hepatic gene silencing in vivo. Angewandte Chemie **2012**, 124, 8657–8661.

(6) Tanford, C.; Kirkwood, J. G. Theory of protein titration curves. I. General equations for impenetrable spheres. Journal of the American Chemical Society **1957**, 79, 5333–5339.

(7) Roux, B.; Simonson, T. Implicit solvent models. Biophysical chemistry **1999**, 78, 1–20.

(8) Zwanzig, R. W. High-temperature equation of state by a perturbation method. I. Non-polar gases. The Journal of Chemical Physics **1954**, 22, 1420–1426.

(9) Kirkwood, J. G. Statistical mechanics of fluid mixtures. The Journal of chemical physics **1935**, 3, 300–313.

(10) Warshel, A.; Sussman, F.; King, G. Free energy of charges in solvated proteins: micro-scopic calculations using a reversible charging process. Biochemistry **1986**, 25, 8368–8372.

(11) Simonson, T.; Carlsson, J.; Case, D. A. Proton binding to proteins: p K a calculations with explicit and implicit solvent models. Journal of the American Chemical Society **2004**, 126, 4167–4180.

(12) Baptista, A. M. Comment on Explicit-solvent molecular dynamics simulation at con-stant pH: methodology and application to small amines[J. Chem. Phys. 114, 9706 (2001)]. The Journal of chemical physics **2002**, 116, 7766–7768.

(13) Donnini, S.; Ullmann, R. T.; Groenhof, G.; Grubmüller, H. Charge-neutral constant pH molecular dynamics simulations using a parsimonious proton buffer. Journal of chemical theory and computation **2016**, 12, 1040–1051.

(14) Philipp et al., pH-dependent structural transitions in cationic ionizable lipid mesophases are critical for lipid nanoparticle function. Submitted **2023**,

(15) Duncan, S. L.; Larson, R. G. Comparing experimental and simulated pressure-area isotherms for DPPC. Biophysical journal **2008**, 94, 2965–2986.

(16) Klauda, J. B.; Venable, R. M.; Freites, J. A.; OConnor, J. W.; Tobias, D. J.; Mondragon-Ramirez, C.; Vorobyov, I.; MacKerell Jr, A. D.; Pastor, R. W. Update of the CHARMM all-atom additive force field for lipids: validation on six lipid types. The journal of physical chemistry B **2010**, 114, 7830–7843.

(17) Lamberg, A.; Ollila, O. S. Comment on Structural properties of POPC monolayers under lateral compression: computer simulations analysis. Langmuir **2015**, 31, 886–887.

(18) Tempra, C.; Ollila, O. S.; Javanainen, M. Accurate simulations of lipid monolayers require a water model with correct surface tension. Journal of Chemical Theory and Computation **2022**, 18, 1862–1869.

(19) Helm, C. A.; Möhwald, H.; Kjaer, K.; Als-Nielsen, J. Phospholipid monolayer density distribution perpendicular to the water surface. A synchrotron X-ray reflectivity study. Europhysics Letters **1987**, 4, 697.

(20) Wiegart, L.; Struth, B.; Tolan, M.; Terech, P. Thermodynamic and structural properties of phospholipid Langmuir monolayers on hydrosol surfaces. Langmuir **2005**, 21, 7349–7357.

(21) Schneck, E.; Schubert, T.; Konovalov, O. V.; Quinn, B. E.; Gutsmann, T.; Brandenburg, K.; Oliveira, R. G.; Pink, D. A.; Tanaka, M. Quantitative determination of ion distributions in bacterial lipopolysaccharide membranes by grazing-incidence X-ray fluorescence. Proceedings of the National Academy of Sciences **2010**, 107, 9147–9151.

(22) Sturm, M.; Gutowski, O.; Brezesinski, G. The Effect of pH on the Structure and Lateral Organization of Cardiolipin in Langmuir Monolayers. ChemPhysChem **2022**, 23, e202200218.

(23) Shapovalov, V. L.; Brezesinski, G. Breakdown of the Gouy–Chapman Model for

Highly Charged Langmuir Monolayers: Counterion Size Effect. The Journal of Physical Chemistry B **2006**, 110, 10032–10040.

(24) Shapovalov, V. L.; Ryskin, M. E.; Konovalov, O. V.; Hermelink, A.; Brezesinski, G. Elemental Analysis within the Electrical Double Layer Using Total Reflection X-ray Fluorescence Technique. J. Phys. Chem. B **2007**, 111, 3927–3934.

(25) Cullis, P. R.; Hope, M. J. Lipid nanoparticle systems for enabling gene therapies. Molecular Therapy **2017**, 25, 1467–1475.

(26) Kanduč, M.; Schlaich, A.; de Vries, A. H.; Jouhet, J.; Maréchal, E.; Demé, B.; Netz, R. R.; Schneck, E. Tight cohesion between glycolipid membranes results from balanced water–headgroup interactions. Nature communications **2017**, 8, 1–9.

(27) Pusterla, J.; Scoppola, E.; Appel, C.; Mukhina, T.; Shen, C.; Brezesinski, G.; Schneck, E. Characterization of lipid bilayers adsorbed to functionalized air/water interfaces. Nanoscale **2022**, 14, 15048–15059.

(28) Als-Nielsen, J.; McMorrow, D. Elements of modern X-ray physics; John Wiley & Sons, 2011.

(29) Yun, W.; Bloch, J. X-ray near total external fluorescence method: Experiment and analysis. Journal of applied physics **1990**, 68, 1421–1428.

(30) Mora, S.; Daillant, J.; Luzet, D.; Struth, B. X-ray surface scattering investigation of Langmuir films: Phase transitions and elastic properties. EPL (Europhysics Letters) **2004**, 66, 694.

(31) Oliveira, R. G.; Schneck, E.; Quinn, B. E.; Konovalov, O. V.; Brandenburg, K.; Gutsmann, T.; Gill, T.; Hanna, C. B.; Pink, D. A.; Tanaka, M. Crucial roles of charged saccharide moieties in survival of gram negative bacteria against protamine revealed by

combination of grazing incidence x-ray structural characterizations and Monte Carlo simulations. Physical Review E **2010**, 81, 041901.

(32) Kanduč, M.; Schneck, E.; Stubenrauch, C. Intersurfactant H-bonds between head groups of n-dodecyl-β-d-maltoside at the air-water interface. Journal of colloid and interface science **2021**, 586, 588–595.

(33) Vineyard, G. H. Grazing-incidence diffraction and the distorted-wave approximation for the study of surfaces. Physical Review B **1982**, 26, 4146.

(34) Parratt, L. G. Surface studies of solids by total reflection of X-rays. Physical review **1954**, 95, 359.

(35) Abelès, F.; La, F. A. théorie générale des couches minces. J. Phys. Radium **1950**, 11, 307–309.

(36) Brezesinski, G.; Schneck, E. Investigating ions at amphiphilic monolayers with X-ray fluorescence. Langmuir **2019**, 35, 8531–8542.

(37) Thompson, A.; Vaughan, D.; others X-ray Data Booklet, Tables 1-3. Lawrence Berkeley National Laboratory, University of California, CA, USA (https://xdb. lbl. gov/xdb. pdf) **2001**,

(38) Caspers, J.; Goormaghtigh, E.; Ferreira, J.; Brasseur, R.; Vandenbranden, M.; Ruysschaert, J.-M. Acido-basic properties of lipophilic substances: A surface potential approach. Journal of Colloid and Interface Science **1983**, 91, 546–551.

(39) Schwierz, N.; Horinek, D.; Netz, R. R. Specific Ion Binding to Carboxylic Surface Groups and the pH Dependence of the Hofmeister Series. Langmuir **2015**, 31, 215–225, PMID: 25494656.

(40) Abraham, M. J.; Murtola, T.; Schulz, R.; Pll, S.; Smith, J. C.; Hess, B.; Lindahl, E.

GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. SoftwareX **2015**, 1-2, 19–25.

(41) Knight, C. J.; Hub, J. S. MemGen: a general web server for the setup of lipid membrane simulation systems. Bioinformatics **2015**, 31, 2897–2899.

(42) Gould, I.; A.A., S.; Dickson, C.; Madej, B.; Walker, R. Lipid17: A Comprehensive AMBER Force Field for the Simulation of Zwitterionic and Anionic Lipids. Unpublished Work **2018**,

(43) Ibrahim, M.; Gilbert, J.; Heinz, M.; Nylander, T.; Schwierz, N. Structural insights on ionizable Dlin-MC3-DMA lipids in DOPC layers by combining accurate atomistic force fields, molecular dynamics simulations and neutron reflectivity. Nanoscale **2023**, 15, 11647–11656.

(44) Mamatkulov, S.; Schwierz, N. Force fields for monovalent and divalent metal cations in TIP3P water based on thermodynamic and kinetic properties. J. Chem. Phys. **2018**, 148, 074504.

(45) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. Comparison of simple potential functions for simulating liquid water. J. Chem. Phys. **1983**, 79, 926–935.

(46) Bussi, G.; Donadio, D.; Parrinello, M. Canonical sampling through velocity rescaling. J. Chem. Phys. **2007**, 126.

(47) Parrinello, M.; Rahman, A. Polymorphic transitions in single crystals: A new molecular dynamics method. J. Appl. Phys. **1981**, 52, 7182–7190.

(48) Essmann, U.; Perera, L.; Berkowitz, M. L.; Darden, T.; Lee, H.; Pedersen, L. G. A smooth particle mesh Ewald method. J. Chem. Phys. **1995**, 103, 8577–8593.

(49) Hess, B.; Bekker, H.; Berendsen, H. J. C.; Fraaije, J. G. E. M. LINCS: A linear constraint solver for molecular simulations. J. Comput. Chem. **1997**, _18_, 1463–1472.

(50) Michaud-Agrawal, N.; Denning, E. J.; Woolf, T. B.; Beckstein, O. MDAnalysis: A toolkit for the analysis of molecular dynamics simulations. J. Comput. Chem. **2011**, _32_, 2319–2327.

(51) Humphrey, W.; Dalke, A.; Schulten, K. VMD: Visual molecular dynamics. J. Mol. Graph. **1996**, _14_, 33–38.

(52) Vega, C.; de Miguel, E. Surface tension of the most popular models of water by using the test-area simulation method. The Journal of chemical physics **2007**, _126_, 154707.

(53) Müller, P.; Bonthuis, D. J.; Miller, R.; Schneck, E. Ionic surfactants at air/water and oil/water interfaces: a comparison based on molecular dynamics simulations. The Journal of Physical Chemistry B **2021**, _125_, 406–415.

(54) Nelson, A. R. J.; Prescott, S. W. refnx: neutron and X-ray reflectometry analysis in Python. J. Appl. Crystallogr. **2019**, _52_, 193–200.

(55) Marsh, D. Lateral pressure in membranes. Biochimica et Biophysica Acta (BBA)-Reviews on Biomembranes **1996**, _1286_, 183–223.

(56) Bloch, J.; Yun, W.; Yang, X.; Ramanathan, M.; Montano, P.; Capasso, C. Adsorption of counterions to a stearate monolayer spread at the water-air interface: A synchrotron x-ray study. Physical review letters **1988**, _61_, 2941.

(57) Daillant, J.; Bosio, L.; Benattar, J.; Blot, C. Interaction of cations with a fatty acid monolayer. A grazing incidence x-ray fluorescence and reflectivity study. Langmuir **1991**, _7_, 611–614.

(58) Nagle, J. F.; Tristram-Nagle, S. Structure of lipid bilayers. Biochimica et Biophysica Acta (BBA)-Reviews on Biomembranes **2000**, _1469_, 159–195.

(59) Stefaniu, C.; Latza, V. M.; Gutowski, O.; Fontaine, P.; Brezesinski, G.; Schneck, E. Headgroup-Ordered Monolayers of Uncharged Glycolipids Exhibit Selective Interactions with Ions. The Journal of Physical Chemistry Letters **2019**, 10, 1684–1690.

(60) Kaganer, V. M.; Möhwald, H.; Dutta, P. Structure and phase transitions in Langmuir monolayers. Reviews of Modern Physics **1999**, 71, 779.

(61) Zantl, R.; Baicu, L.; Artzner, F.; Sprenger, I.; Rapp, G.; Rädler, J. O. Thermotropic phase behavior of cationic lipid- DNA complexes compared to binary lipid mixtures. The Journal of Physical Chemistry B **1999**, 103, 10300–10310.