1

Sub-nucleolar trafficking of Hendra virus matrix protein is regulated by ubiquitination and oligomerisation

Stephen M. Rawlinson[1][*][#], Tianyue Zhao[1][*], Florian A. Gomez[1], Cassandra T. David[1], Christina L. Rootes[2], Patrick F. Veugelers[1,3], Ashley M. Rozario[4], Cameron R. Stewart[2], Toby D.M. Bell[3] and Gregory W. Moseley[1][#]

[1] Department of Microbiology, Biomedicine Discovery Institute, Monash University, 19 Innovation Walk (Bldg 76), Clayton Campus, Victoria 3800, Australia.

[2] CSIRO Health & Biosecurity, Australian Centre for Disease Preparedness, Geelong, Victoria, Australia.

[3] School of Chemistry, Monash University, Clayton, Victoria 3800, Australia.

[4] Department of Rural Clinical Sciences, La Trobe Institute for Molecular Science, La Trobe University, Bendigo, Victoria 3552, Australia.

* SMR and TZ contributed equally to this work.

Correspondence should be addressed to: SMR (stephen.rawlinson@monash.edu) and GWM (greg.moseley@monash.edu).

ABSTRACT

Hendra virus (HeV) is a highly pathogenic member of the Henipavirus genus (order *Mononegavirales*), the replication cycle of which occurs primarily in the cytoplasm. The HeV matrix protein (HeV M) plays critical roles in viral assembly and budding at the plasma membrane, but also undergoes nuclear/nucleolar trafficking, to accumulate in nucleoli early in infection and, later, localise predominantly at the plasma membrane. Previously we found that HeV M protein targets specific sub-nucleolar compartments (corresponding to the FC-DFC (fibrillar centre (FC)/dense fibrillar component (DFC)) where it interacts with the nucleolar protein Treacle and modulates rRNA biogenesis by subverting the host nucleolar DNA damage response, indicating the importance of specific sub-nucleolar trafficking to infection. However, the mechanisms underlying targeting and movement between sub-nucleolar compartments by viral or cellular proteins remain poorly defined. Here, we assessed the molecular regulation of HeV M protein nucleolar/sub-nucleolar trafficking, finding that in infected cells and in cells expressing HeV M protein alone, M protein localizes into Treacle-enriched FC-DFC at early time points, and that FC-DFC localization is subsequently lost due to relocalization into the surrounding granular component (GC) of the nucleolus. Analysis using mutated M proteins and pharmacological modulation of ubiquitination indicate that this dynamic localization is regulated by ubiquitination and oligomerisation, with ubiquitination required for retention of HeV M in Treacle-enriched sub-nucleolar compartments, and oligomerisation required for egress. To our knowledge, this study provides the first direct insights into the dynamics and mechanisms of viral protein trafficking between sub-nucleolar compartments, important to the interplay between HeV M protein and host cell factors during infection.

AUTHOR SUMMARY

Henipaviruses, including Hendra (HeV) and Nipah viruses, cause deadly diseases in humans and livestock and are considered priority diseases by the World Health Organization due to their epidemic potential and lack of effective treatments. Understanding how these viruses interact with host cells is essential for developing new therapeutics. Our study examines the matrix (M) protein of henipaviruses and its interaction with the nucleolus, a cell structure that mediates ribosome production, and is a common target for various viruses, although their functions are largely unresolved. Previously, we showed that the HeV M protein targets a sub-nucleolar structure, called the FC-DFC, to modulate ribosome biogenesis. Here, we report that the M protein's movement between sub-nucleolar compartments is controlled by two processes: ubiquitination, which causes accumulation of the protein in the FC-DFC, and oligomerization, which is associated with exit. Similar mechanisms are also observed in other henipaviruses. Our findings reveal mechanisms regulating the hijacking of host cell functions by henipaviruses and suggest new potential targets for antiviral therapies. This study is the first to investigate how viral proteins move within the nucleolus, offering new insights into interactions that may be significant to multiple viruses.

INTRODUCTION

The nucleolus comprises a highly multifunctional structure with long established roles in ribosome biogenesis, as well as roles in cell cycle regulation, the DNA damage response (DDR), cellular stress responses, and signal recognition particle assembly [1, 2]. The nucleolus was recently shown to be a membrane-less organelle (MLO) comprising at least three immiscible liquid condensates that are formed by liquid-liquid phase separation (LLPS) [3]. The three components are the fibrillar

centre (FC), dense fibrillar component (DFC) and granular component (GC). The FC is surrounded

by the DFC to form functional units (FC-DFC), which are embedded within the GC (Fig. 1A) [4].

These compartments play distinct roles, including assembling a pipeline for the key steps of

ribosome biogenesis.

Consistent with its multifunctionality, the nucleolus is a common target of diverse viruses [5-7].

This targeting is proposed to enable viral exploitation of diverse processes to usurp host cell

biology and/or facilitate virus replication [5-7]. Despite the prevalence of viral protein nucleolar

targeting, functional outcomes generally remain poorly understood. The potential nucleolar

functions of viral proteins are of particular interest with respect to RNA viruses that typically have

limited coding capacity and replicate their genomes in the cytoplasm, but nevertheless target

specific proteins to nucleoli. These include the highly pathogenic non-segmented negative sense

RNA viruses (nsNSVs) Hendra (HeV) and Nipah (NiV) viruses (genus *Henipavirus*, family

Paramyxoviridae), the matrix (M) protein of which localizes to the nucleus and nucleolus during

infection [8-11].

Henipavirus M proteins form stable homodimers, and play critical roles in virus assembly in the

cytoplasm and in budding at the plasma membrane, which involves the formation of M protein

oligomers [12, 13]. The subcellular localization of M protein is dynamic, being nucleolar early in

infection before exiting the nucleolus/nucleus and accumulating at the plasma membrane for

assembly/budding [8-11]. Interestingly, transit through the nucleolus is reported to be a

prerequisite for M protein to fulfill assembly and budding functions, suggestive of a regulatory

role of nucleoli in viral release [10, 11]. Genetic screens have indicated the importance of nucleolar

proteins in infection, and proteomic datasets suggest that M protein interacts with multiple

nucleolar proteins [8, 10, 14, 15]. However, potential intranucleolar roles of HeV M protein

remained unresolved until the identification of a novel nucleolar function whereby HeV M

localises to a sub-nucleolar compartment corresponding to the FC-DFC, where it interacts with

Treacle protein and impairs ribosomal RNA (rRNA) biogenesis [14]. This process appears to be

mediated by mimicry of a cellular process that normally occurs during a DDR. Thus subcellular

trafficking underpins key functions of HeV M. However, how this trafficking is regulated,

particularly between sub-nucleolar compartments and other regions of the cell remains unresolved.

Indeed, the mechanisms regulating trafficking of proteins in general between sub-nucleolar liquid

condensates is poorly understood, with no prior studies to our knowledge, for any viral protein.

Previously we showed that substitution of residue K258 in HeV M for alanine (HeV M K258A)

impairs FC-DFC localisation/Treacle-binding and DDR modulation/budding activity, without

preventing localization to the GC, where HeV M K258A accumulates [14]. K258 forms part of a

bipartite nuclear localization sequence (NLS; often referred to as 'NLS2'; M contains at least two

NLSs; NLS1 is located at residues 82-87 [16, 17]) and is reported to be required for M

ubiquitination (Fig. 2A) [10, 11]. It has been proposed that the M protein enters the nucleus via

the NLS and accumulates within nucleoli before exiting the nucleolus and nucleus (mediated by a

nuclear export sequence (NES)). Exit of the nucleolus/nucleus is reported to be triggered by

ubiquitination requiring K258 by an unidentified ubiquitin ligase [18]. However, this model was

proposed prior to the description of the functionally important localization of HeV M to sub-

nucleolar compartments. As a result, the coordination and regulation of various trafficking steps,

including trafficking within the nucleolus, remain undefined.

The potential role of ubiquitination, and definition of other molecular mechanisms in sub-nucleolar

localization are of particular interest as mechanisms regulating nucleolar/sub-nucleolar trafficking

(which involves movement between LLPS structures) are poorly understood compared with those

for nuclear trafficking, which involves conventional protein interactions with trafficking receptors

and the nuclear pore complex. In this study, we examined the regulation of HeV M protein

trafficking between the FC-DFC and GC finding that ubiquitination plays a crucial role.

Interestingly, our data indicate that ubiquitination exerts opposing effects on sub-nucleolar and

nucleocytoplasmic localization, suppressing exit from the FC-DFC to the GC while being required

for egress from the nucleolus/nucleus. Furthermore, mutations affecting HeV M oligomerization

indicated that oligomerization is necessary for FC-DFC egress. These findings provide new

insights into the interplay of ubiquitination and oligomerization in HeV M protein

nuclear/nucleolar transit and roles in modulation of host cell biology, viral assembly and budding.

RESULTS

Ubiquitination affects sub-nucleolar trafficking of HeV M protein.

Previously HeV M protein was reported to be ubiquitinated at several sites, potentially including residue K258 (and equivalent residues in other henipaviruses), and that mutation of K258 to A or R (the latter preventing ubiquitination but retaining the positive charge) inhibits ubiquitination of several sites that were shown to be mono-ubiquitinated [10, 11]. Ubiquitination was implicated in M protein trafficking using the proteasome/ubiquitination inhibitor, MG132, which caused nuclear and nucleolar retention of HeV and NiV M proteins, and inhibited nuclear export and viral-like particle (VLP) production by NiV M proteins [10, 11].

To explore the possibility that ubiquitination regulates trafficking between sub-nucleolar condensates, we examined the effect of MG132 treatment on localization/accumulation of HeV M protein to sub-nucleolar punctate compartments (which correspond to FC-DFC; Fig. S1 and data previously reported data [14]) using confocal laser scanning microscopy (CLSM) analysis of living HeLa cells expressing GFP-fused wild-type (WT) HeV M (GFP-HeV M). GFP-HeV M accumulated with FC-DFC (Fig. 1B). Notably, FLAG-tagged HeV M protein also localised within FC-DFC (Fig. S2A) and interacted with Treacle (Fig. S2B), but not with the Treacle-binding mutant (K258A), as expected [14], suggesting that the GFP-tag is not causing artefacts.

As expected [10, 11], MG132 treatment resulted in an apparent increase in nuclear accumulation of GFP-HeV M protein (Fig. 1B); quantitative image analysis confirmed a significant increase in the nuclear to cytoplasmic fluorescence ratio (Fn/c) (Fig. 1C). This is consistent with previously reported impairment of nuclear export [10, 11]. In contrast, accumulation of M protein in the FC-

DFC appeared to be reduced (Fig. 1B), and this effect was confirmed by a reduced ratio of

fluorescence intensity of the FC-DFC compared with the GC ($F_{FC\text{-}DFC/GC}$) (Fig. 1D), an increase in

GC fluorescence indicative of movement of M protein from the FC-DFC into the surrounding GC

(Fig. 1E), and a decrease in the number of cells with FC-DFC accumulation of M protein (Fig.

1F).

To confirm that the effects of MG132 on HeV M sub-nucleolar trafficking are due to

ubiquitination, we co-transfected cells with a plasmid expressing HA-ubiquitin (HA-Ubi) to

replenish ubiquitin depletion by MG132. Expression of HA-Ubi reversed the effect of MG132 in

reducing FC-DFC accumulation in a dose-dependent fashion (Fig. 1G, H). These findings indicate

that ubiquitination promotes the accumulation of HeV M protein within the FC-DFC, while

reduced ubiquitination leads to its egress from the FC-DFC and accumulation in the GC. The

observed increase in GC fluorescence is consistent with previous reports of an apparent

enhancement of nuclear and nucleolar localization of M protein following MG132 treatment,

proposed to reflect decreased export from the nucleus and corresponding decrease in egress from

the nucleolus[10]. Our data suggest that the accumulation of diffuse (GC) fluorescence in the

nucleolus following MG132 treatment is not solely due to reduced nuclear/nucleolar egress, but

also increased egress from the FC-DFC to the GC. Notably, the opposing effects of ubiquitination

on FC-DFC and nuclear/nucleolar GC localization indicate different mechanisms affecting

trafficking between the compartments, such that FC-DFC localization is not simply the result of

altered protein concentration in the nucleus/nucleolar compartment, but is specifically and

distinctly regulated by ubiquitination.

Previously, proteosome inhibitors were shown to reduce NiV titers during live virus infections,

indicating that ubiquitination plays a critical role in NiV infection [11]. To test if similar

mechanisms occur during HeV infection, HeLa cells were infected with HeV at MOIs of 0.5 or 5,

followed by treatment with proteosome inhibitors MG132 (Fig. 1I) and Bortezomib (Fig. 1J). Both

inhibitors reduced virus titers in a dose-dependent manner at both MOIs, with statistically

significant effects at both MOIs for Bortezomib and at MOI 0.5 MG132; the reduction observed

for MG132 at MOI 5 was dose-dependent but not significant. These findings confirm the

importance of ubiquitination in HeV infection, similar to what is observed with NiV [11].

Conservative substitution of K258 to R reduces FC-DFC targeting by HeV M.

We previously showed that K258A mutation in HeV M protein (HeV M K258A) abolishes its

targeting to the FC-DFC, resulting in accumulation within the GC and loss of binding to the FC-

DFC-enriched protein, Treacle [14]. This suggested that K258 forms part of a targeting signal due

to its positive charge and/or affects sub-nucleolar localization due to its ubiquitination [10, 11].

The above data (Fig. 1) indicate that ubiquitination is required for the retention of HeV M within

the FC-DFC. Mutation at K258 is reported to affect mono-ubiquitination at several sites, which

has been suggested to indicate ubiquitination likely occurs at K258 but also impacts on other

mono-ubiquitination sites in M protein [11]. To examine whether the effects we observed on FC-

DFC localization following MG132 relate to ubiquitination at K258 or associated sites, we

compared the effects on subcellular localization of HeV M by the substitutions K258A (which

removes the positive charge and the potential ubiquitination site) and K258R (which retains a

positive charge but lacks the lysine of the potential ubiquitination site) (Fig. 2A). Previous studies

on equivalent mutations in NiV M protein indicated that the positive charge is important to

function of the nuclear localization sequence (NLS) and nucleolar accumulation, while ubiquitination regulates nuclear export [11]. However, no effects on sub-nucleolar localization was reported, although our data indicate that ubiquitination has opposing effects on nuclear/nucleolar accumulation and FC-DFC accumulation (above).

CLSM analysis of cells expressing GFP-fused HeV M WT, K258A or K258R variants (Fig. 2B), indicated sub-nucleolar accumulation of WT M protein in c. 90% of cells, consistent with localization to FC-DFC (Fig. 2C). As expected [14], HeV M K258A protein did not localize/accumulate within FC-DFC, but accumulated within the GC in 100% of cells (Fig. 2B). In contrast, HeV M K258R displayed an intermediate phenotype, with a substantial proportion of cells (c. 60%) showing FC-DFC accumulation similar to WT and the remainder lacking FC-DFC accumulation similar to K258A (Fig. 2B, C). Consistent with this, the $F_{FC\text{-}DFC/GC}$ ratio for HeV M K258A (c. 1.0) was significantly lower than that for HeV M WT (c. 1.8), while HeV M K258R showed an intermediate phenotype (c. 1.4) (Fig. 2D). The reduced $F_{FC\text{-}DFC/GC}$ for HeV M K258R resulted from the presence of a K258A-like sub-population (for which the $F_{FC\text{-}DFC/GC}$ was equivalent to that for K258A) and the fact that the $F_{FC\text{-}DFC/GC}$ for the population with apparent FC-DFC accumulation was significantly lower than the $F_{FC\text{-}DFC/GC}$ for HeV WT protein (p < 0.01) (Fig. 2D); thus, even in cells where HeV M K258R localized to the FC-DFC, this localization was impaired compared with the HeV M WT protein. Overexpression of HA-Ubi, treatment with MG132, or a combination of these conditions did not result in any significant FC-DFC accumulation of HeV M K258A (Fig. 2E, F), consistent with the positive charge at residues 258 being essential for FC-DFC localization [14]. Interestingly, MG132-treatment of cells expressing GFP-HeV-M-K258R resulted in a significant reduction in FC-DFC accumulation to reach levels

similar to HeV M K258A, and HeV M WT with MG132/HA (Fig. 1H). Expression of HA-Ubi

reversed this effect (Fig. 2G, H). Together, these data imply that ubiquitination dependent on K258

is required for efficient FC-DFC localization, but that ubiquitination at other K258-independent

sites (either in M protein or other cellular proteins), is also required.

Dynamic localization of HeV M in sub-nucleolar compartments is regulated by K258.

The effects of the K258 mutation on HeV M protein accumulation in FC-DFC suggest potential

impacts on a targeting sequence and/or affinity for specific components within the FC-DFC. For

NiV M protein, K258 is proposed to be part of a NLS, which typically consists of short stretches

of basic residues. Several basic residues (R244, R245, R256, R257, K258 in NiV M) are highly

conserved among henipavirus M proteins [10, 11]. Thus, K258 may play a crucial role in nuclear

import through the NLS and contribute to an overlapping targeting sequence for nucleoli/sub-

nucleolar FC-DFC [11, 17]. Previous data indicated that ubiquitination dynamically regulates the

nuclear localization of HeV M [10]. Moreover, NiV M protein undergoes dynamic and temporal

regulation of localization during infection, being nuclear/nucleolar early in infection before

nucleolar exit/nuclear export, and eventual accumulation and budding at the plasma membrane [9,

11]. Similarly, we found that in HeV-infected cells, HeV M localizes to sub-nucleolar

compartments (FC-DFC) early during infection (7 hours post-infection (p.i.)), but becomes more

diffuse in the nucleolus (i.e. accumulated into the GC), and with greater nuclear accumulation at

24 hours p.i. (Fig. 3A), consistent with observations for GFP HeV M WT protein and observations

of dynamic nuclear/nucleolar localization of M protein in NiV infected cells [11]. Thus, we

speculated that the observed differences in FC-DFC localization between HeV M WT and the

K258 mutants (assessed at 24 hours post-transfection in Fig. 1 and Fig. 2) might be attributable to

the dynamic regulation of various M protein trafficking signals related to the changes in HeV M

localization during infection.

To investigate this, we assessed the sub-nucleolar localization of HeV M WT and mutant proteins

at time points from 8 h to 72 h p.t. (Fig. 3B-D). The WT HeV M protein exhibited clear

accumulation in FC-DFC (typically multiple structures in each nucleolus) at 8 h p.t. (c. 85% of

cells), which progressively diminished over the course of the experiment, accompanied by a more

diffuse GC distribution with only around 10% of cells exhibiting accumulation of M protein in

multiple FC-DFC at 72 h p.t. (Fig. 3E). This is consistent with a dynamic interaction whereby M

protein initially enters the FC-DFC and then undergoes gradual egress to the GC. Measurement of

the $F_{FC\text{-}DFC/GC}$ confirmed a progressive loss of FC-DFC localization (Fig. 3F).

Consistent with roles of K258 in NLS activity of NiV M protein [10], the Fn/c for HeV M K258A

was reduced compared with WT at 8 and 16 h p.t., supporting its involvement in nuclear import

(Fig. 3G; Fn/c c. 2 for WT, compared with Fn/c c. 1 for K258A at both timepoints). Further

analysis revealed that the reduction in nuclear localization of the K258A mutant was due to a

significant proportion of cells with higher fluorescence intensity in the cytoplasm (Fc) than in the

nucleus (Fn) at early time points, in contrast to cells expressing WT and K258R M protein (Fig.

3H; c. 50-60% of cells expressing K258A M protein showed Fn > Fc between 8-24 h p.t., whereas

in cells expressing WT M protein, > 85% of cells showed Fn > Fc at all timepoints). However,

over time, the K258A mutant gradually exhibited a proportion of cells with Fn > Fc similar to WT

and K258R (nearly 100% of cells at 48 and 72 h p.t.), suggesting a delay in nuclear import of

K258A compared to the other variants.

Despite reduced nuclear accumulation at early time points, HeV M K258A was strongly nucleolar

at all time points, consistent with reduced nucleolar egress. However, no accumulation in FC-DFC

was observed at any time point, and some images indicated absence of fluorescence from these

structures (e.g. white arrow, 48 h p.t., Fig. 3C). Thus, HeV M protein can specifically partition

between sub-nucleolar phase-separated compartments, dependent on K258, and this is independent

of the accumulation in the nucleus, consistent with distinct mechanisms of trafficking/localization.

Notably, similar dynamics were observed for WT NiV M, with the percentage of cells showing

FC-DFC accumulation reducing over time (c. 25% of cells showing FC-DFC accumulation at 72

h p.t.), while NiV M-K258A remained excluded from FC-DFC at all timepoints (Fig. S3).

HeV M K258R accumulated to higher levels in the nucleus than the cytoplasm at early time points

compared with K258A (Fig. 3D, upper panels and 3G), similar to WT M. This is consistent with

a requirement for the positive charge in the NLS for efficient nuclear import, as reported for NiV

M protein [11, 17]. Additionally, HeV M K258R accumulated to very high levels in the nucleus

at later time points (48 h), consistent with an impaired nuclear export mechanism [10, 11]. HeV

M K258R also showed clear FC-DFC localization in Treacle-enriched compartments at 8 h p.t.

(Fig. 3D), similar to (but moderately reduced compared with) WT HeV M protein, followed by

loss of FC-DFC localization over time. Thus, the presence of a basic residue at position 258 is

necessary for initial entry and accumulation within the FC-DFC.

While FC-DFC localization of WT and K258R HeV M protein diminished following the initial

accumulation, the apparent rate of loss was greater for HeV M K258R, such that by 24 h p.t. (Fig.

3E) c. 25% of HeV M K258R-expressing cells displayed FC-DFC localization compared with c.

60% for WT HeV M. By 48 and 72 h p.t. < 5% and 0%, respectively, of HeV M K258R-expressing

cells displayed FC-DFC localization, and nucleoli with apparent exclusion from FC-DFC

structures were apparent (e.g. 48 h p.t., Fig. 3D, white arrow, similar to observations for HeV M

K258A. Calculation of the $F_{FC-DFC/GC}$ ratio confirmed a significant decrease in FC-DFC

localization by both HeV M WT and K258R over the course of the experiment, with a more rapid

decrease for the latter (Fig. 3F). Thus, it appears that HeV M localises initially to the FC-DFC,

dependent primarily on the presence of a positive charge at position 258. HeV M then relocalizes

to the GC, and this process is accelerated in HeV M containing the K258R substitution that is

impaired for ubiquitination, consistent with ubiquitination supporting retention into the FC-DFC.

FC-DFC accumulation is enhanced by inhibition of HeV M protein oligomerization.

Protein oligomerization has been associated with regulation of subcellular localization, including

nuclear import and export of proteins such as p53 [19] and signal transducer and activator of

transcription (STAT) proteins [20]. Moreover, oligomerization is implicated in the formation of

MLOs through LLPS [21, 22]. Based on the crystal structure of HeV M protein, residue R57 of

HeV M has been implicated in oligomerization *via* packing of HeV M dimers [13]. Mutation of

R57 to D or E impairs the formation of VLPs by M protein, suggesting roles for oligomerization

in the assembly and budding processes [13]. Assembly and budding is also dependent on K258

[10, 11], suggesting that nuclear and nucleolar localization/trafficking, and budding are

intrinsically linked and/or involve coordinated regulation. Additionally, it has been hypothesized

that ubiquitin modification at K258 may be necessary for M protein oligomerization [18].

However, roles of oligomerization in HeV M nuclear/nucleolar trafficking have not been

investigated.

To examine this we analysed the localisation of HeV M WT or R57D as above, which indicated

pronounced accumulation of HeV M R57D into the FC-DFC compared to the WT protein (Fig.

4A). The Fn/c was not significantly different between HeV M WT and R57D indicating similar

function of the nuclear trafficking signals (Fig. 4B). However, the ratio of nucleolar to nuclear

fluorescence (Fnu/n) (Fig. 4C) and $F_{FC-DFC/GC}$ (Fig. 4D) was significantly increased for HeV M

R57D compared to WT HeV M protein. Thus, R57D enhances the accumulation of HeV M in the

FC-DFC, consistent with a role for oligomerization in regulating the localization of HeV M to

nucleolar condensates. These data further indicate that the mechanisms underlying M protein FC-

DFC localization differ from those regulating nuclear import/export. Interestingly, in contrast to

results for WT M protein, MG132 treatment did not reduce FC-DFC accumulation (Fig. 4E, F).

This suggests that the impaired ability of HeV M R57D protein to oligomerize both enhances FC-

DFC localization, and causes it to become insensitive to the effects of reduced ubiquitination.

These data are consistent with egress of M protein from the FC-DFC being dynamically regulated

by ubiquitination, but also being dependent on the ability of M protein to undergo oligomerization.

Accumulation of HeV M protein in the FC-DFC correlates with inhibitory function toward

rRNA biogenesis and with binding to Treacle.

HeV M protein accumulation in FC-DFC enables subversion of the nucleolar DDR pathway

through targeting of Treacle. This function is strongly impaired by the K258A mutation [14].

R57D and K258R enhance (Fig. 4) and suppress (to a level intermediate between WT and K258A

protein; Fig. 2), respectively, accumulation within the FC-DFC, suggesting that

oligomerization/ubiquitination regulate functional FC-DFC interactions. To assess this directly,

we used the Click-iT RNA Imaging Kit to measure the effects of M protein and mutants on rRNA

biogenesis (Fig. 5A, B), and co-immunoprecipitation (co-IP) assays to assess interactions with

Treacle (Fig. 5C), as previously described [14, 23].

Consistent with previous findings, WT HeV M protein, but not HeV M K258A significantly

impaired rRNA biogenesis at 24 h p.t.; the level of inhibition was comparable with that observed

for rRNA silencing by Treacle knockdown (c. 20-30% inhibition) [14, 24] (Fig. 5A, B). In contrast,

R57D, significantly enhanced inhibition of rRNA biogenesis compared with WT M protein (c.

30% inhibition), consistent with strong FC-DFC localization and apparent loss of ubiquitin-

regulated localization. The K258R mutation resulted in a phenotype for rRNA silencing that was

intermediate between WT and K258A M proteins, correlating with their differing capacities to

localize to the FC-DFC. Notably, the ability of WT M protein to inhibit rRNA biogenesis was lost

by 72 h p.t. (Fig. S4), matching with dynamic loss of FC-DFC localization (Fig. 3). As expected,

rRNA biogenesis in cells expressing HeV M K258A was unchanged between 24 and 72 h p.t.,

correlating with consistent lack of localization to the FC-DFC, while R57D was found to inhibit

rRNA biogenesis at both 24 and 72 h p.t. (consistent with a sustained FC-DFC localization over

time, see below and Fig. 6). Thus, the extent of FC-DFC accumulation appears to correspond with

inhibition of rRNA biogenesis.

Consistent with the differing localization and functional impact of the mutated proteins, co-IP and

immunoblotting (IB) analysis indicated increased interaction of HeV M R57D protein with

Treacle, compared with WT HeV M protein (Fig. 5C; c. 1.6-fold more Treacle precipitated with

R57D). Furthermore, Treacle interaction of HeV M K258R was reduced compared with WT

protein (c. 15% of WT levels), but was greater than HeV M K258A (c. 5 %), indicating an

intermediate phenotype. Notably, Treacle is reported to form multiple isoforms [25], and only the

upper band of Treacle appeared to IP with the M proteins (Fig. 5C), perhaps indicating interaction

with specific Treacle isoforms. Taken together, these data indicate that FC-DFC accumulation,

Treacle binding, and the inhibition of rRNA biogenesis by the HeV M protein are correlated,

consistent with regulation of M protein function in rRNA production *via* ubiquitin and

oligomerization dependent FC-DFC localization/Treacle interaction.

Loss of HeV M FC-DFC accumulation does not relate to disruption or loss of FC-DFC.

Loss of HeV M FC-DFC localization (Fig. 3) could be attributed to two possible mechanisms: (1)

egress of the protein from intact FC-DFC structures, or (2) depletion of FC-DFC structures through

events such as fusion or disassembly/disruption of the liquid bodies. To investigate these

possibilities, we analysed cellular FC-DFCs directly by fixation and immunostaining of cells for

Treacle at time points from 8-72 h p.t. to express HeV M WT, K258A, K258R, or R57D (Fig. 6A-

D).

In cells expressing GFP-HeV M WT protein the appearance of FC-DFCs was similar throughout

the experiment (Fig. 6A). At early time points, HeV M WT protein strongly colocalized with

Treacle in FC-DFC, but this diminished over time (indicated by a reduced percentage of cells with

detectable colocalization of HeV M and Treacle FC-DFCs), although multiple Treacle-enriched

FC-DFCs lacking HeV M association remained detectable in nucleoli (Fig. 6A, E). Thus, HeV M

protein appears to transit through FC-DFC, where it interacts with Treacle, before egress, with no

significant disruption of FC-DFC structures. As expected, HeV M K258A showed no

colocalization/accumulation in Treacle FC-DFC at any time point, despite the presence of multiple

FC-DFC. HeV M K258R protein showed similar results to WT, but with more rapid egress, as

expected (Fig. 3C), and no evident loss of FC-DFC (Fig. 6C, E). By 24-48 h p.t., co-localization

was barely detectable, similar to K258A (Fig. 6E). In contrast, HeV M R57D exhibited clear

colocalization with Treacle in all detectable FC-DFCs from 8 to 48 h p.t., with only a minor

downward trend (colocalization with FC-DFC in c. 95% of cells measured) at 72 h p.t. (Fig. 6D,

E). These findings are consistent with a requirement for oligomerization for HeV M protein to

egress the FC-DFC, and with the capability of HeV M R57D to maintain efficient rRNA silencing

function at 72 h p.t., by which time WT M protein has largely lost such function. Thus, it appears

that HeV M protein transits through intact FC-DFC, with the loss of colocalization due to

trafficking rather than disruption or major structural change to FC-DFC. These data further support

that dynamic localization of M protein to FC-DFC, and interaction with Treacle, underlie specific

silencing of rRNA biogenesis.

HeV M proteins do not affect the structure or distribution of FC-DFC.

The lack of major disruption of FC-DFC during HeV M protein accumulation or egress aligns with

our previous findings [14]. Specifically, using single molecule localization microscopy (SMLM)

with direct stochastic optical reconstruction microscopy (*d*STORM) of cells immunostained for

Treacle, we found that HeV M WT protein accumulation in FC-DFC (24 h p.t.) does not cause

'gross' changes to FC-DFC numbers or to the structure of FC-DFC [14]. Based on the enhanced

FC-DFC localization and function in rRNA silencing observed for HeV M R57D, we reasoned

that, if M protein does have any structural effects on FC-DFC, they would be exaggerated by

R57D. We thus analysed cells expressing GFP-HeV M WT, R57D or K258A protein, and cells

expressing GFP alone or non-transfected cells, by immunostaining for Treacle followed by

*d*STORM (Fig. 7A). Expression of HeV M WT, K258A (as previously [14]), K258R and R57D

proteins had no significant impact on the size of Treacle-labelled FC-DFC (Fig. 7B). These data

are consistent with specific intermolecular interactions of M protein with Treacle within FC-DFC

regulating rRNA biogenesis, and the duration/extent of functional modulation being controlled by

ubiquitination (and potentially multimerization)-dependent egress of M protein from FC-DFC.

Ubiquitination regulates sub-nucleolar trafficking of M proteins of multiple henipaviruses

Previously, we showed that the FC-DFC accumulation, Treacle binding, and inhibition of rRNA

biogenesis are conserved among M proteins of multiple henipaviruses (including NiV, Cedar

(CedV), and Mojiang (MojV) viruses), albeit with some differences in the extent of FC-DFC

accumulation [23]. To determine if ubiquitin-dependence of FC-DFC accumulation is conserved

in different henipaviruses, we assessed the effects of MG132 as above (e.g. Fig. 1). Similar to HeV

M, MG132 treatment significantly impaired NiV M FC-DFC accumulation (Fig. 8A, B) and

reduced the percentage of cells with FC-DFC accumulation (Fig. 8C) consistent with the homology

of HeV and NiV M proteins (~90% amino acid identity). Consistent with our previous report [23],

CedV M showed the highest accumulation in FC-DFC and lowest GC accumulation of the M

proteins assessed; $F_{FC-DFC/GC}$ accumulation of CedV M protein was significantly reduced (but

remained higher than that of HeV or NiV M proteins) following MG132 treatment, and the

percentage of cells with FC-DFC accumulation of CedV M protein remained c. 100% (Fig. 8C).

MojV M showed the lowest accumulation in FC-DFC (consistent with previous data [23]),

resulting only a minor and non-significant reduction in $F_{FC-DFC/GC}$ (Fig. 8B); however, there was a

significant reduction in the percentage of cells with clear FC-DFC accumulation of MojV M

protein following MG132 treatment (Fig. 8D).Taken together, these data indicate conserved roles

of ubiquitination in regulating henipavirus M protein localization to the FC-DFC, although the

extent of accumulation differs between M proteins, correlating with evolutionary divergence (c.

61% and 60% similarity of CedV and MojV M proteins, respectively, compared with HeV M

protein).

<u>DISCUSSION</u>

Here we have found that HeV M protein dynamically transits through the FC-DFC, indicating that

the previously identified translocation through the nucleus/nucleolus involves additional sub-

nucleolar trafficking between LLPS structures, with the different stages of trafficking regulated by

post-translational modification. This transit enables regulated interactions with Treacle and other

host factors, enabling functional regulation of rRNA synthesis by modulating the nucleolar DDR,

as well as virus assembly and budding. To our knowledge, this study presents the first data on the

mechanisms governing the trafficking of a viral protein within sub-nucleolar LLPS structures, and

expands and refines the model for M protein trafficking. Specifically, we found that M trafficking

to FC-DFC requires a basic residue at residue 258, and that its egress from FC-DFC requires

oligomerization that is regulated, at least in part, by the ubiquitination status of the M protein.

Importantly, while ubiquitination has previously been reported to regulate nuclear and overall

nucleolar localization[10], indicating that ubiquitination is required for efficient nuclear export and

nucleolar egress, our data, in contrast indicates that ubiquitination is required for nucleolar

retention/accumulation into the FC-DFC. These findings highlight that localization of proteins to

specific sub-nucleolar compartments involves highly specific mechanisms.

Taken together, our data support a model for M protein sub-nucleolar trafficking in which (1) entry

of M protein dimers into the FC-DFC and functional interaction with Treacle to modulate rRNA

biogenesis requires a basic charge at residue 258, and (2) exit from the FC-DFC requires

oligomerization of the M protein within the FC-DFC. However, the retention/egress of M from the

FC-DFC is dynamically regulated, at least in part, by the ubiquitination status of M protein. Our

data support a likely role for ubiquitination of K258, but also indicate that other ubiquitination

sites, either within M protein or on host proteins, also contribute, as FC-DFC localization by HeV

M K258R protein is impaired but remains sensitive to inhibition of ubiquitination. These

observations are consistent with previous data indicating that M protein can be mono-ubiquitinated

at at least four sites, and K258R mutation inhibits ubiquitination at several of these sites (proposed

to include K258 itself) which is likely to contribute to the impaired FC-DFC localization of this

mutant. However, at least one site remains functional which may account for the residual

accumulation of HeV M K258R into FC-DFC that is lost following MG132 treatment [10, 11].

Our data are consistent with a model whereby M protein exits the FC-DFC in the form of oligomers

but egress is negatively regulated by ubiquitination of M protein; however, it is also possible that

ubiquitination of host proteins also regulates interactions with M protein required for exit.

Movement of proteins between LLPS MLO structures such as nucleolar sub-compartments does

not use conventional translocation processes associated with membrane-enclosed organelles (e.g.

movement *via* pores/channels), but rather depends on partitioning through physicochemical

properties and interactions with MLO-resident molecules [4, 26]. This likely accounts for the poor

definition of nucleolar 'targeting sequences' compared with NLS and NES motifs that form

specific interactions with trafficking receptor proteins (importins and exportins) [27].

Ubiquitination and oligomerization play important roles in the formation and regulation of LLPS

[28, 29], and, notably, our data indicate a potential role for oligomerization, possibly regulated by

ubiquitination, in coordinating M protein's interactions/localization into sub-nucleolar liquid

bodies. Thus, ubiquitination/oligomerisation may alter the physicochemical properties of M

protein or interactions with constituents of different nucleolar condensates, as well as affecting

importin/exportin interactions and interactions at budding sites. The differing nature of the

mechanisms of sub-nucleolar trafficking and nucleocytoplasmic trafficking are consistent with our

observations that ubiquitination has differing effects on exit from the FC-DFC to the GC, nucleolus

to the nucleus, and nucleus to the cytoplasm. Thus, specific orchestration of ubiquitination,

deubiquitination, oligomerization, and LLPS interactions may underlie appropriate temporal

regulation of transport between these compartments, enabling specific control of rRNA silencing,

virus replication, and assembly/budding, aligning with different stages of the viral life cycle [13].

As viruses typically mimic or hijack cellular processes, our findings likely have implications

beyond viral infection. The intricate regulation of M protein in multiple intranuclear compartments

is unlikely to have evolved solely to control the concentration in the cytoplasm for viral processes

such as assembly and budding. Rather, it suggests a specific coordination of sub-nuclear functions,

including DDR subversion, where M protein appears to mimic cellular NBS1 [14]. Our findings

on the regulation of sub-nucleolar partitioning of M protein, including roles of positively charged

residues (typical of nuclear/nucleolar targeting signals), ubiquitination and oligomerisation

identifies mechanisms that may be relevant to cellular proteins that transit sub-nucleolar

compartments, including those of the DDR. In the light of current advances toward therapeutic

modulation of cellular and viral LLPS structures [30-33], our data also has the potential to

contribute to novel antiviral approaches for currently incurable Henipavirus infection, and possibly

other nucleolus-related pathologies [34, 35].

499
500
MATERIALS AND METHODS

Cell culture, transfection and treatment

HEK-293T (ATCC: CRL-3216) and HeLa (ATCC: CCL-2) cells were cultured in Dulbecco's

Modified Eagle Medium (DMEM) supplemented with 10% Fetal Calf Serum (FCS), 2 mM

Glutamax, 50 U/mL Penicillin, and 50 μg/mL Streptomycin. The cells were maintained at 37 °C

with 5% CO_2. HEK-293T and HeLa cells were grown to 80-90% confluency before transfection

using Lipofectamine 2000 and Lipofectamine 3000, respectively, according to the manufacturer's

instructions (ThermoFisher Scientific). For free ubiquitin depletion, transfected HeLa cells were

treated with 50 μM MG132 or 0.5% DMSO for control at 18 h p.t. for 6 h before CLSM imaging

analysis. MG132 was purchased from Sigma (M7449-200UL) as a 10 mM ready-made solution in

DMSO. Experiments for co-expression of HA-Ubi, HeLa cells were transfected with the same

total amount of DNA of 2500 ng: 1000 ng of GFP-HeV M co-transfected with 1500 ng total of

HA-alone plasmid and/or HA-Ubi, with the HA/HA-Ubi ratio varied.

Virus infections

Wild-type HeV (Hendra virus/horse/1994/Hendra) was used for all virus work and performed at

the CSIRO Australian Centre for Disease Preparedness (CSIRO-ACDP) in Biosafety Level (BSL)-

4 laboratories. For analysis of IF, HeLa cells were seeded onto coverslips and mock- or HeV-

infected (MOI 5) prior to fixation at 7 h and 24 h p.i. using 4% paraformaldehyde (1h, RT) and

permeabilization with 0.1% TritonX-100 for 10 min. IF labelling was performed using a mouse

primary antibody to HeV M (1:500; developed internally (Ref#: 1805-21-1527) and an anti-mouse

AlexaFluor 488 secondary antibody. DNA was visualised using a DAPI.

For tissue culture infective dose ($TCID_{50}$) analysis HeLa cells were seeded into 96-well plates prior

to HeV-infection the next day at MOI 0.5 or 5. At 18 h p.i. cells were treated with DMSO, MG132

(1 nM, 10 nM or 100 nM) or Bortezomib (5 nM, 50 nM or 500 nM). At 25 h p.i., additional

Bortezomib was added to Bortezomib samples, as previously done for NiV[11]. At 42 h p.i.

supernatants were collected and $TCID_{50}$/ml was determined as previously[36].

Constructs

Mammalian cell expression of N-terminal GFP tagged HeV-M (Accession Number AEB21196.1),

and mutants were generated by directional cloning of the M gene cDNA into the multiple cloning

site of the pEGFP-C1 vector, as previously described[17]. Site-directed mutagenesis of R57D

residue was performed by applying QuickChange method, using *Pfu* DNA Polymerase from

Promega. The following oligonucleotides were used to introduce the single R57D mutation: 5'-

caagatctataccccaggtgcaaatgagGACaaattcaacaactacatgtacatg-3'   and   3'-

catgtacatgtagttgttgaatttGTCctcatttgcacctggggtatagatcttg-5'. DNA sequencing was performed to

confirm the correct mutation. The plasmid for expression of HA-ubiquitin (HA-Ubi) has been

published previously [37].

Confocal laser scanning microscopy (CLSM) and image analysis

For CLSM imaging analysis, HeLa cells were seeded on 1.5 (0.17 mm) thickness glass coverslips

and transfected with the indicated constructs at 80-90% confluency. Imaging was performed at the

indicated time p.t. or 24 h p.t., if not specified. CLSM was conducted using a Nikon C1 inverted

confocal microscope with a 60× oil immersion objective (NA 1.4) at Monash Micro Imaging

Facility. Live-cell CLSM imaging was performed within a heated chamber at 37 °C.

CLSM images were analyzed using ImageJ freeware software. The mean fluorescence of the

nucleus (Fn), cytoplasm (Fc), nucleolus (Fnu; whole nucleolus), FC-DFC ($F_{FC\text{-}DFC}$), GC (F_{GC}), and

background fluorescence (Fb) were determined. After subtracting the background fluorescence

(Fb) from all values, the nuclear to cytoplasmic (Fn/c), nucleolar to nuclear (Fnu/n), and FC-DFC

to GC ($F_{FC\text{-}DFC/GC}$) fluorescence ratios were calculated. In cells where accumulation into sub-

nucleolar compartments was not evident (e.g., cells expressing K258A- or K258R-mutated M

protein), two distinct areas in the diffuse region of the nucleolus were selected to represent the

"FC-DFC" and the "GC" for image analysis. The F_{GC} analysis was based on images captured under

the same microscopy and software settings.

The percentage of cells with FC-DFC accumulation was determined by dividing the number of

cells showing any nucleolus with FC-DFC ≥ 1 by the total number of cells expressing the indicated

proteins in each sample. Data are presented as mean $\pm$ S.E.M. (standard error of the mean) or mean

± SD (standard deviation), as indicated in the figure legend. Statistical analysis (Student's t-test)

was performed using GraphPad Prism software.

Immunofluorescence (IF)

For IF staining, cells grown on glass coverslips were washed twice gently with PBS at 24 h p.t.,

fixed with 4% (w/v) paraformaldehyde at room temperature (RT) for 15 min, permeabilized using

0.25% Triton X-100 (v/v in PBS) at RT for 5 min, and blocked with 1% bovine serum albumin

(BSA) in PBS at RT for 1 h. For samples expressing FLAG-M proteins, an additional 5 min

incubation with 5 µg/ml proteinase K was performed after fixation. The cells were then incubated

with primary antibody specific to either Treacle (1:100; Cat # 11003-1-AP, Proteintech), UBF1

(1:500; Cat# Ab244287; Abcam), Nucleolin (1:200; Cat#14574, CST), anti-FLAG (1:250;

Cat#F1804, Sigma) or NPM1 (1:200; Cat# 32-5200, ThermoFisher Scientific) at RT for 1.5 h.

Subsequently, cells were incubated with goat anti-rabbit or anti-mouse 568 AlexaFluor conjugate

secondary antibody (Cat # A-11011/A-11004, ThermoFisher Scientific) at a 1:1000 dilution in the

dark at RT for 1.5 h. DNA staining was performed using Hoechst 33342 at a 1:2000 dilution of a

20 mM stock solution (Cat # 62249, ThermoFisher Scientific). The cells were mounted onto

microscope glass slides (Lomb Scientific) using Mowiol reagent.

5-ethnyl uridine (EU) incorporation assays.

Levels of rRNA synthesis were determined using an image-based technique (Click iT RNA Alexa

Fluor 594 Imaging kit, Thermo-Fisher, Cat# C10330) as previously described [14, 23, 24, 38].

Cells were incubated for 1 h in the presence of EU before fixation in 4% paraformaldehyde at RT

for 12 min and permeabilization in 0.25% Triton X-100 for 5 min at RT. Samples were processed

according to the manufacturer's recommendations to label incorporated EU with Alexa Fluor 594.

DNA was labeled using Hoechst 33342. Cells were imaged by CLSM to detect labeling of nascent

rRNA by measuring the fluorescence intensity of Alexa Fluor 594 within nucleoli. Quantitative

analysis was performed using ImageJ software to determine the mean EU fluorescence of nucleoli,

which were identified using a combination of DNA, GFP, and transmitted light channels. Relative

rRNA synthesis levels were determined by measuring nucleolar EU levels of both GFP-expressing

and non-expressing cells from the same sample (as an internal control) and expressed as nucleolar

EU of GFP-expressing cells relative to non-expressing cells.

Immunoprecipitation (IP) and immunoblotting (IB)

HEK-293T cells were seeded in 6 cm dishes and transfected to express the indicated GFP-fused

proteins. At 24 h p.t., cells were harvested and lysed in 10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5

mM EDTA, 0.5% NP-40, and 1x Protease Inhibitor Cocktail (PIC; Sigma-Aldrich

Cat#11697498001) for 20 min at 4°C. The lysate was cleared by centrifugation at 20,000 xg for 6

min at 4°C, and 10% of the supernatant was collected as the 'input' sample. The remaining lysate

was incubated with GFP-Trap magnetic beads (ChromoTek) as previously described for 45 min at

4°C on a rotary mixer [14]. The beads were then washed three times with dilution buffer (10 mM

Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 1x PIC) and resuspended in 2x SDS-PAGE sample

buffer to solubilize the proteins. The entire process from the addition of lysis buffer to the addition

of 2x SDS-PAGE sample buffer was completed in less than 1.5 h to minimize degradation of

Treacle, which is commonly observed (e.g. [24, 38, 39]).

Immunoprecipitates and cell lysates (input) were analyzed by SDS-PAGE and immunoblotting

(IB). Proteins in IP and input samples were separated on 10% SDS-PAGE gels and transferred

onto nitrocellulose membranes using the Bio-Rad Trans-Blot Turbo Transfer System according to

the manufacturer's instructions. After protein transfer, the membranes were blocked in blocking

buffer (5% non-fat milk in PBS with 0.1% Tween-20) at RT for 1 h, incubated with the indicated

primary antibodies at RT for 2 h or at 4°C overnight on a rocker, followed by incubation with

HRP-conjugated secondary antibodies (goat anti-rabbit or anti-mouse) at RT for 1 h. The

membranes were then imaged using a Gel Doc XR+ Gel Documentation System (Bio-Rad). The

primary antibodies used for IB were anti-Treacle (Cat # 11003-1-AP, Proteintech; 1:2000), anti-

GFP (Cat # 11814460001, Roche; 1:2000), and anti-β-tubulin (Cat # T8328, Sigma; 1:2000). The

secondary antibodies used for IB were goat anti-rabbit (Cat # AP307P) and goat anti-mouse (Cat

AP308P) IgG HRP-conjugated antibodies (Merck Millipore). The Treacle bands in the IP were

quantitated using ImageLab (Version 6.0) software.

_d_STORM imaging and analysis

HeLa cells were fixed 24 h p.t. with 4% paraformaldehyde for 10 min then permeabilized with

0.1% Triton X-100 for 10 min. Fixed cells were blocked in 2% BSA/PBS for 30 min, then stained

with anti-Treacle primary antibodies (3 μg/ml, 1 h) and Alexa Fluor 647 conjugated secondary

antibodies (5 μg/ml, 45 min). Samples were imaged with a switching buffer of 100 mM

mercaptoethylamine (MEA) in PBS made to pH 8.5 [40]. _d_STORM imaging was performed on a

home-built super-resolution set-up as previously described [40, 41]. Briefly, the setup comprised

an inverted fluorescence microscope (Olympus IX81, 100X 1.49 NA TIRF objective) with an

EMCCD camera (Andor iXon). Blue laser excitation (Toptica iBeam 488 nm) was used to identify

GFP-positive cells. High power red laser excitation (Oxxius LaserBoxx 638 nm) was used to

induce reversible photoswitching of Alexa Fluor 647 fluorophores in the fixed samples. Single

molecule 'blinking' events were captured at 100 Hz for between 10,000 – 20,000 frames and

processed using rapi*d*STORM [42] to render a 2D coordinate map as the super-resolved *d*STORM

images of subnucleolar Treacle. For analysis, images were first smoothed in ImageJ using the

Gaussian Blur function (Sigma (Radius) = 0.5) to account for single molecule localization

precision error. Images were converted to 8-bit (greyscale values 2-255 threshold) then converted

into binary images. Using the "Distance Map" function in ImageJ, individual subnucleolar

compartments were discretized based on pixel density by applying a threshold for greyscale values

2-255. The "Analyse Particles" function was then used to identify and measure subnucleolar

compartments.

FIGURE LEGENDS

Figure 1   **Ubiquitination regulates FC-DFC accumulation of HeV M and impacts on**

virus production. (A) Schematic of a nucleolus showing the three primary sub-compartments:

fibrillar centre (FC), dense fibrillar component (DFC) and granular component (GC). The region

composed of the FC and DFC compartments is referred to as the FC-DFC. (B) HeLa cells

transfected to express GFP-HeV M protein were treated 18 h post transfection (p.t.) with MG132

or without (DMSO) for 6 h before CLSM analysis. Representative images are shown for each

condition; yellow boxes are magnified in the zoom panel. Yellow arrowheads indicate nucleoli;

white arrowheads indicate localization of M protein to sub-nucleolar compartments consistent with

FC-DFC. Images such as those in B were analysed to determine: (C) the nuclear to cytoplasmic

(Fn/c) fluorescence ratio; (D) the ratio of fluorescence of the FC-DFC to that of the GC ($F_{FC\text{-}DFC/GC}$); (E) the fluorescence intensity of the GC (F_{GC}) (arbitrary units (a.u.)), and (F) the % of HeV

M-expressing cells with apparent accumulation in FC-DFC (histogram shows the percentage of M

protein-expressing cells containing at least one nucleolus with evident accumulation of M protein

into one or more FC-DFC). Histograms for C, D and E show mean ± S.E.M., n ≥ 24 cells for each

condition (data from a single assay, consistent with two independent experiments); histogram in F

shows mean percentage ± SD from two independent assays; ≥ 73 cells for each condition. (G)

HeLa cells co-transfected with plasmid to express GFP-HeV M and with differing amounts of HA

or HA-ubiquitin (HA-Ubi) expression plasmid (1500 ng total HA/HA-Ubi plasmid transfected,

comprising HA-Ubi and/or HA, as indicated) and treated without (DMSO) or with MG132. (H)

Images such as those in G were used to calculate the $F_{FC\text{-}DFC/GC}$. Data from a single assay (n = 24

cells per sample), representative of two independent assays. (I, J) HeLa cells infected with HeV at

MOI 0.5 or MOI 5 and treated with increasing concentrations of proteosome inhibitors, (I) MG132

or (J) Bortezomib (Bort), prior to collection at 42 h p.i. and determination of HeV titres

(TCID50/ml ± S.E.M., n = 6). Statistical analysis used Student's t-test; * p < 0.05; ** p < 0.01;***

p < 0.001; **** p < 0.0001; ns, non-significant.

Figure 2 **K258R mutation impacts sub-nucleolar trafficking of HeV M protein**. (A)

Schematic of the HeV M protein indicating the bipartite NLS (residues 244-258; critical basic

residues are in bold and underlined) and residue K258, mutation of which regulates the ubiquitin

status (Ubi) of M protein by removing the proposed ubiquitin site at 258, and affecting other sites.

Mutations used in this study are indicated (grey boxes). (B) CLSM images of living HeLa cells

expressing the indicated GFP-fused proteins (24 h p.t.); for HeV M WT and K258A, images are

representative of 90-100% of cells in > 29 fields of view; for HeV M K258R two major populations

(each representing c. 40-60% of the population) were observed, corresponding to either a "FC-

DFC accumulation" (upper panel) or "K258A-like" (lower panel) phenotype. Nucleoli are

highlighted by the yellow box, which is magnified in the zoom panel. White arrowheads indicate

accumulation within FC-DFC. (C) Images such as those in B were analysed to determine the

percentage of cells with clear FC-DFC accumulation of M protein (mean percentage $\pm$ SD, n = 3

separate assays, each sampling $\geq$ 119 cells). Student's t-test with Welch's correction was used to

determine significance; ** $p < 0.01$.) (D) Images were analysed to determine $F_{FC\text{-}DFC/GC}$ (mean $\pm$

S.E.M., n $\geq$ 90 cells for each condition, from three independent assays; green line in K258R

indicates mean of all samples). The two distinct populations in the K258R sample are indicated.

Comparison of K258R "WT-like" population with WT samples using Student's t-test showed a

significant difference ($p < 0.01$). (E-H) Images of HeLa cells co-transfected to express GFP-HeV

M K258A (E) or K258R (G) with different amounts of HA or HA-ubiquitin-expressing plasmid

imaged by CLSM (as in Fig. 1G, H). Images such as these were analyzed to determine the $F_{FC\text{-}}$

$_{DFC/GC}$ of K258A (F) and K258R (H) expressing cells (n $\geq$ 17 cells for F, and n $\geq$ 21 for H; data

from one assay, representative of two independent assays). Statistical analysis used Student's t-

test; * $p < 0.05$; ** $p < 0.01$; **** $p < 0.0001$; ns, non-significant.

Figure 3   **HeV M protein undergoes dynamic localization to the FC-DFC which is**

impacted by K258R mutation. (A) HeLa cells were mock-infected or infected with HeV (MOI

5) prior to fixation and immunostaining for HeV M protein at 7 h and 24 h post-infection (p.i.).

Nuclei were detected using DAPI. Microscope settings and image correction are identical between

equivalent mock and HeV infected images. (B-C) HeLa cells transfected to express the indicated

proteins were analysed live at 8, 16, 24, 48 and 72 h p.t. by CLSM. Images representative of major

phenotypes are shown for each condition; yellow boxes are magnified in the zoom panel. Yellow

arrowheads indicate nucleoli; white arrowheads indicate accumulation of M protein in FC-DFC.

Images such as those in B-C were analysed to determine: (E) the percentage of M protein-

expressing cells with FC-DFC accumulation in any nucleolus in the cell (mean ± SD from three

independent assays, n ≥ 59 cells for each condition); (F) $F_{FC-DFC/GC}$ (mean ± S.E.M., n ≥ 55 cells

for each condition from three independent assays, except for 8 h p.t. (WT, K258A, K258R), 16 h

p.t. K258A, and 48 h p.t. WT samples, where data is from two assays; (G) Fn/c (mean ± S.E.M.;

n ≥ 24 cells) and (H) percentage of cells with Fn > Fc (mean ± SD from three independent assays,

except for 8 h p.t. (WT, K258A, K258R), 16 h p.t. K258A, and 48 h p.t. WT samples, where data

is from two assays; n ≥ 23 cells for each condition). Student's t-test was used to compare values

for WT and K258R at each timepoint in D and E; * p< 0.05; *** p < 0.001; **** p < 0.0001; ns,

non-significant.

Figure 4   **R57D mutation enhances accumulation of HeV M in FC-DFC and is**

insensitive to ubiquitination depletion. (A) CLSM images of live HeLa cells expressing the

indicated proteins (24 h p.t.). Images are representative of 90-100% of cells in > 9 fields of view

from a single assay, and typical of ≥ 4 independent assays. Hoechst 33342 (blue) was used to

identify DNA/nuclei. Nucleoli are highlighted by the yellow box, which is magnified in the zoom

panel. Yellow arrowheads indicate nucleoli; white arrowheads indicate accumulation in FC-DFC.

(B-D). Images such as those shown in A were used to determine (B) Fn/c; (C) nucleolar to nuclear

fluorescence ratio (Fnu/n; Fnu is the mean fluorescence intensity for the entire area of the

nucleolus); (D) $F_{FC-DFC/GC}$. B, C and D show mean ± S.E.M., n ≥ 24 cells for each condition. (E)

CLSM images of HeLa cells expressing the indicated protein and treated without (DMSO) or with MG132. (F) Images such as those in E were analysed to determine $F_{FC-DFC/GC}$ (n ≥ 14 cells from one assay; data representative of three independent assays). Statistical analysis used Student's t-test; **** $p < 0.0001$; ns, non-significant.

Figure 5   **Silencing of rRNA synthesis and Treacle-binding by HeV M protein/mutants correlate with FC-DFC accumulation.** (A) HeLa cells were transfected to express the indicated GFP-fused proteins before addition of EU reagent (23 h p.t.) for 1h, fixation (24 h p.t.), labeling to detect nascent RNA (EU fluorescence) and DNA (Hoechst 33342), and imaging by CLSM. (B) Images such as those in A were used to determine EU fluorescence in the nucleoli of GFP-positive cells, relative to that in non-GFP expressing cells in the same sample (mean relative EU fluorescence ± S.E.M., n ≥ 51 cells from two independent experiments). * $p < 0.05$; *** $p < 0.001$; **** $p < 0.0001$; ns, non-significant. (C) HEK-293T cells were transfected to express GFP or the indicated GFP-fused proteins prior to lysis (24 h p.t.) and IP for GFP. Cell lysate (input) and IP samples were subjected to IB using the indicated antibodies. Quantitation of Treacle band in IPs, relative to that detected in IP for GFP-HeV M WT are indicated.

Figure 6   **M protein FC-DFC accumulation decreases over time without loss of FC-DFC compartments.** (A-D) HeLa cells were transfected to express the indicated proteins before fixation at 8, 16, 24, 48 and 72 h p.t. and immunostaining using anti-Treacle antibody (red) and imaging via CLSM. Hoechst 33342 (blue in Merge panels) was used to localize DNA/nuclei. Representative images are shown for each condition; yellow arrowheads indicate nucleoli; unfilled white arrowheads indicate Treacle in FC-DFC; white arrowheads indicate accumulation of M

protein into FC-DFC; yellow arrows indicate colocalization of Treacle and HeV M protein in FC-

DFC. (E) Images such as those in A-D were analysed to determine the percentage of cells

expressing HeV M protein with evident colocalization of HeV M protein and Treacle in FC-DFC

(n ≥ 23 cells for each condition).

Figure 7   **HeV M R57D protein does not affect the structure or distribution of Treacle-**

enriched FC-DFC. (A) HeLa cells expressing the indicated proteins or non-transfected (NT) were

fixed and immunolabeled for Treacle before imaging using SMLM with *d*STORM. Fluorescence

microscopy was used to identify cells expressing GFP/GFP-fusion proteins (green). Yellow boxes

in fluorescence images show zoomed region imaged using SMLM (Zoom 1); yellow boxes in

Zoom 1 images show additional zoomed region (Zoom 2). Scale bars correspond to 5000 nm

(Zoom 1) and 500 nm (Zoom 2). SMLM images such as those shown in A were used to determine

(B) Tukey boxplots showing median area and the 95% confidence interval of Treacle-enriched

FC-DFC (n ≥ 505 FC-DFC for each sample). Statistical analysis used Kolmogorov-Smirnov tests

as normality testing indicated a non-parametric distribution.

Figure 8 FC-DFC accumulation of M proteins of multiple henipaviruses is regulated by

ubiquitination. (A) CLSM images of live HeLa cells transfected to express the indicated proteins.

Representative images are shown for each condition, with yellow boxes magnified in the zoom

panel. Hoechst 33342 was used to stain nuclei/DNA. (B) Images such as those in A were used to

determine $F_{FC\text{-}DFC/GC}$ (mean $F_{FC\text{-}DFC/GC}$ ± S.E.M., from one assay (n ≥ 13), representative of three

independent   assays). (C) The   percentage   of   M   protein-expressing   cells   with   apparent

accumulation of M protein in FC-DFC (mean percentage ± S.E.M. from three independent assays;

each sample was determined from n ≥ 13 cells).

<u>ACKNOWLEDGEMENTS</u>

We acknowledge the facilities and technical assistance of Monash Micro Imaging (Monash

University) with confocal microscopy. This study was supported by National Health and Medical

(NHMRC) Research Council grants 1125704, 1160838 and 1079211 to GWM; Australian

Research Council (ARC) grants DP210100998, DP150102569 to GWM, and DP170104477 to

TDMB; and the Miegunyah Trust Grimwade Fellowship to GWM.

1.   Boisvert FM, van Koningsbruggen S, Navascues J, Lamond AI. The multifunctional

nucleolus. Nat Rev Mol Cell Biol. 2007;8(7):574-85. doi: 10.1038/nrm2184. PubMed PMID:

17519961.

2.   Iarovaia OV, Minina EP, Sheval EV, Onichtchouk D, Dokudovskaya S, Razin SV, et al.

Nucleolus: A Central Hub for Nuclear Functions. Trends Cell Biol. 2019;29(8):647-59. Epub

2019/06/10. doi: 10.1016/j.tcb.2019.04.003. PubMed PMID: 31176528.

3.   Feric M, Vaidya N, Harmon TS, Mitrea DM, Zhu L, Richardson TM, et al. Coexisting

Liquid   Phases   Underlie   Nucleolar   Subcompartments.   Cell.   2016;165(7):1686-97.   Epub

2016/05/24.   doi: 10.1016/j.cell.2016.04.047.   PubMed   PMID: 27212236;   PubMed   Central

PMCID: PMCPMC5127388.

4.   Lafontaine DLJ, Riback JA, Bascetin R, Brangwynne CP. The nucleolus as a multiphase

liquid   condensate.   Nat   Rev   Mol   Cell   Biol.   2021;22(3):165-82.   Epub   2020/09/03.   doi:

10.1038/s41580-020-0272-6. PubMed PMID: 32873929.

5.   Hiscox JA. RNA viruses: hijacking the dynamic nucleolus. Nat Rev Microbiol.

2007;5(2):119-27. Epub 2007/01/17. doi: 10.1038/nrmicro1597. PubMed PMID: 17224921;

PubMed Central PMCID: PMCPMC7097444.

6.   Iarovaia OV, Ioudinkova ES, Velichko AK, Razin SV. Manipulation of Cellular Processes

via Nucleolus Hijaking in the Course of Viral Infection in Mammals. Cells. 2021;10(7). Epub

2021/07/03. doi: 10.3390/cells10071597. PubMed PMID: 34202380; PubMed Central PMCID:

PMCPMC8303250.

7.   Rawlinson SM, Moseley GW. The nucleolar interface of RNA viruses. Cell Microbiol.

2015;17(8):1108-20. doi: 10.1111/cmi.12465. PubMed PMID: 26041433.

8.   Deffrasnes C, Marsh GA, Foo CH, Rootes CL, Gould CM, Grusovin J, et al. Genome-wide

siRNA Screening at Biosafety Level 4 Reveals a Crucial Role for Fibrillarin in Henipavirus

Infection. PLoS Pathog. 2016;12(3):e1005478. doi: 10.1371/journal.ppat.1005478. PubMed

PMID: 27010548; PubMed Central PMCID: PMCPMC4806981.

9.   Monaghan P, Green D, Pallister J, Klein R, White J, Williams C, et al. Detailed

morphological characterisation of Hendra virus infection of different cell types using super-

resolution and conventional imaging. Virol J. 2014;11:200. doi: 10.1186/s12985-014-0200-5.

PubMed PMID: 25428656; PubMed Central PMCID: PMCPMC4254186.

10.   Pentecost M, Vashisht AA, Lester T, Voros T, Beaty SM, Park A, et al. Evidence for

ubiquitin-regulated nuclear and subnuclear trafficking among Paramyxovirinae matrix proteins.

PLoS Pathog. 2015;11(3):e1004739. doi: 10.1371/journal.ppat.1004739. PubMed PMID:

25782006; PubMed Central PMCID: PMCPMC4363627.

11.   Wang YE, Park A, Lake M, Pentecost M, Torres B, Yun TE, et al. Ubiquitin-regulated

nuclear-cytoplasmic trafficking of the Nipah virus matrix protein is important for viral budding.

PLoS Pathog. 2010;6(11):e1001186. doi: 10.1371/journal.ppat.1001186. PubMed PMID:

21085610; PubMed Central PMCID: PMCPMC2978725.

12.   Battisti AJ, Meng G, Winkler DC, McGinnes LW, Plevka P, Steven AC, et al. Structure

and assembly of a paramyxovirus matrix protein. Proc Natl Acad Sci U S A. 2012;109(35):13996-

4000. Epub 2012/08/15. doi: 10.1073/pnas.1210275109. PubMed PMID: 22891297; PubMed

Central PMCID: PMCPMC3435216.

13.   Liu YC, Grusovin J, Adams TE. Electrostatic Interactions between Hendra Virus Matrix

Proteins Are Required for Efficient Virus-Like-Particle Assembly. J Virol. 2018;92(13). Epub

2018/04/27. doi: 10.1128/JVI.00143-18. PubMed PMID: 29695428; PubMed Central PMCID:

PMCPMC6002731.

14.   Rawlinson SM, Zhao T, Rozario AM, Rootes CL, McMillan PJ, Purcell AW, et al. Viral

regulation of host cell biology by hijacking of the nucleolar DNA-damage response. Nat Commun.

2018;9(1):3057. Epub 2018/08/05. doi: 10.1038/s41467-018-05354-7. PubMed PMID: 30076298;

PubMed Central PMCID: PMCPMC6076271.

15.   Sun W, McCrory TS, Khaw WY, Petzing S, Myers T, Schmitt AP. Matrix proteins of

Nipah and Hendra viruses interact with beta subunits of AP-3 complexes. J Virol.

2014;88(22):13099-110. doi: 10.1128/JVI.02103-14. PubMed PMID: 25210190; PubMed Central

PMCID: PMCPMC4249102.

16.   Donnelly CM, Vogel OA, Edwards MR, Taylor PE, Roby JA, Forwood JK, et al.

Henipavirus Matrix Protein Employs a Non-Classical Nuclear Localization Signal Binding

Mechanism. Viruses. 2023;15(6). Epub 2023/06/28. doi: 10.3390/v15061302. PubMed PMID:

37376602; PubMed Central PMCID: PMCPMC10303457.

17.   McLinton EC, Wagstaff KM, Lee A, Moseley GW, Marsh GA, Wang LF, et al. Nuclear

localization and secretion competence is conserved amongst henipavirus matrix proteins. J Gen

Virol. 2017. doi: 10.1099/jgv.0.000703. PubMed PMID: 28056216.

18.   Watkinson RE, Lee B. Nipah virus matrix protein: expert hacker of cellular machines.

FEBS Lett. 2016;590(15):2494-511. doi: 10.1002/1873-3468.12272. PubMed PMID: 27350027.

19.   Liang SH, Clarke MF. Regulation of p53 localization. Eur J Biochem. 2001;268(10):2779-

83. Epub 2001/05/19. doi: 10.1046/j.1432-1327.2001.02227.x. PubMed PMID: 11358492.

20.   Gorby C, Martinez-Fabregas J, Wilmes S, Moraga I. Mapping Determinants of Cytokine

Signaling via Protein Engineering. Front Immunol. 2018;9:2143. Epub 2018/10/16. doi:

10.3389/fimmu.2018.02143.   PubMed   PMID: 30319612;   PubMed   Central   PMCID:

PMCPMC6170656.

21.   Banani SF, Lee HO, Hyman AA, Rosen MK. Biomolecular condensates: organizers of

cellular biochemistry. Nat Rev Mol Cell Biol. 2017;18(5):285-98. Epub 2017/02/23. doi:

10.1038/nrm.2017.7. PubMed PMID: 28225081; PubMed Central PMCID: PMCPMC7434221.

22.   Boeynaems S, Alberti S, Fawzi NL, Mittag T, Polymenidou M, Rousseau F, et al. Protein

Phase Separation: A New Phase in Cell Biology. Trends Cell Biol. 2018;28(6):420-35. Epub

2018/04/01. doi: 10.1016/j.tcb.2018.02.004. PubMed PMID: 29602697; PubMed Central PMCID:

PMCPMC6034118.

23.   Rawlinson SM, Zhao T, Ardipradja K, Zhang Y, Veugelers PF, Harper JA, et al.

Henipaviruses and lyssaviruses target nucleolar treacle protein and regulate ribosomal RNA

synthesis. Traffic. 2023;24(3):146-57. Epub 2022/12/09. doi: 10.1111/tra.12877. PubMed PMID:

36479968.

24.   Larsen DH, Hari F, Clapperton JA, Gwerder M, Gutsche K, Altmeyer M, et al. The NBS1-

Treacle complex controls ribosomal RNA transcription in response to DNA damage. Nat Cell Biol.

2014;16(8):792-803. doi: 10.1038/ncb3007. PubMed PMID: 25064736; PubMed Central PMCID:

PMCPMC4962910.

25.   So RB, Gonzales B, Henning D, Dixon J, Dixon MJ, Valdez BC. Another face of the

Treacher Collins syndrome (TCOF1) gene: identification of additional exons. Gene. 2004;328:49-

57. Epub 2004/03/17. doi: 10.1016/j.gene.2003.11.027. PubMed PMID: 15019983.

26.   Ditlev JA, Case LB, Rosen MK. Who's In and Who's Out-Compositional Control of

Biomolecular Condensates. J Mol Biol. 2018;430(23):4666-84. Epub 2018/08/14. doi:

10.1016/j.jmb.2018.08.003. PubMed PMID: 30099028; PubMed Central PMCID:

PMCPMC6204295.

27.   Emmott E, Hiscox JA. Nucleolar targeting: the hub of the matter. EMBO Rep.

2009;10(3):231-8. doi: 10.1038/embor.2009.14. PubMed PMID: 19229283; PubMed Central

PMCID: PMCPMC2658561.

28.   Luo YY, Wu JJ, Li YM. Regulation of liquid-liquid phase separation with focus on post-

translational modifications. Chem Commun (Camb). 2021;57(98):13275-87. Epub 2021/11/25.

doi: 10.1039/d1cc05266g. PubMed PMID: 34816836.

29.   Mohanty P, Kapoor U, Sundaravadivelu Devarajan D, Phan TM, Rizuan A, Mittal J.

Principles Governing the Phase Separation of Multidomain Proteins. Biochemistry. 2022. Epub

2022/07/09. doi: 10.1021/acs.biochem.2c00210. PubMed PMID: 35802394.

30.   Conti BA, Oppikofer M. Biomolecular condensates: new opportunities for drug discovery

and RNA therapeutics. Trends Pharmacol Sci. 2022;43(10):820-37. Epub 2022/08/27. doi:

10.1016/j.tips.2022.07.001. PubMed PMID: 36028355.

31.   Mitrea DM, Mittasch M, Gomes BF, Klein IA, Murcko MA. Modulating biomolecular

condensates: a novel approach to drug discovery. Nat Rev Drug Discov. 2022;21(11):841-62.

Epub 2022/08/17. doi: 10.1038/s41573-022-00505-4. PubMed PMID: 35974095; PubMed Central

PMCID: PMCPMC9380678 discovery company that studies condensates, and have a financial

stake in the company.

32.   Risso-Ballester J, Galloux M, Cao J, Le Goffic R, Hontonnou F, Jobart-Malfait A, et al. A

condensate-hardening drug blocks RSV replication in vivo. Nature. 2021;595(7868):596-9. Epub

2021/07/09. doi: 10.1038/s41586-021-03703-z. PubMed PMID: 34234347.

33.   Wang S, Dai T, Qin Z, Pan T, Chu F, Lou L, et al. Targeting liquid-liquid phase separation

of SARS-CoV-2 nucleocapsid protein promotes innate antiviral immunity by elevating MAVS

activity. Nat Cell Biol. 2021;23(7):718-32. Epub 2021/07/10. doi: 10.1038/s41556-021-00710-0.

PubMed PMID: 34239064.

34.   Maehama T, Nishio M, Otani J, Mak TW, Suzuki A. Nucleolar stress: Molecular

mechanisms and related human diseases. Cancer Sci. 2023;114(5):2078-86. Epub 2023/02/11. doi:

10.1111/cas.15755. PubMed PMID: 36762786; PubMed Central PMCID: PMCPMC10154868.

35.   Stochaj U, Weber SC. Nucleolar Organization and Functions in Health and Disease. Cells.

2020;9(3). Epub 2020/02/29. doi: 10.3390/cells9030526. PubMed PMID: 32106410; PubMed

Central PMCID: PMCPMC7140423.

36.   Stewart CR, Marsh GA, Jenkins KA, Gantier MP, Tizard ML, Middleton D, et al.

Promotion of Hendra virus replication by microRNA 146a. J Virol. 2013;87(7):3782-91. Epub

2013/01/25. doi: 10.1128/JVI.01342-12. PubMed PMID: 23345523; PubMed Central PMCID:

PMCPMC3624204.

37.   Frank D, Garnish SE, Sandow JJ, Weir A, Liu L, Clayer E, et al. Ubiquitylation of RIPK3

beyond-the-RHIM can limit RIPK3 activity and cell death. iScience. 2022;25(7):104632. Epub

2022/07/09. doi: 10.1016/j.isci.2022.104632. PubMed PMID: 35800780; PubMed Central

PMCID: PMCPMC9254354.

38.   Mooser C, Symeonidou IE, Leimbacher PA, Ribeiro A, Shorrocks AK, Jungmichel S, et

al. Treacle controls the nucleolar response to rDNA breaks via TOPBP1 recruitment and ATR

activation. Nat Commun. 2020;11(1):123. Epub 2020/01/09. doi: 10.1038/s41467-019-13981-x.

PubMed PMID: 31913317; PubMed Central PMCID: PMCPMC6949271.

39.   Ciccia A, Huang JW, Izhar L, Sowa ME, Harper JW, Elledge SJ. Treacher Collins

syndrome TCOF1 protein cooperates with NBS1 in the DNA damage response. Proc Natl Acad

Sci U S A. 2014;111(52):18631-6. doi: 10.1073/pnas.1422488112. PubMed PMID: 25512513;

PubMed Central PMCID: PMCPMC4284564.

40.   Whelan DR, Bell TD. Image artifacts in single molecule localization microscopy: why

optimization of sample preparation protocols matters. Sci Rep. 2015;5:7924. doi:

10.1038/srep07924. PubMed PMID: 25603780; PubMed Central PMCID: PMCPMC4300460.

41.   Rozario AM, Zwettler F, Duwé S, Hargeaves RB, Brice A, Dedecker P, et al. 'Live and

large': super-resolution optical fluctuation imaging (SOFI) and expansion microscopy (ExM) of

microtubule remodelling by rabies virus P protein. Aust J Chem. 2020;73(8):686-92.

42.   Wolter S, Loschberger A, Holm T, Aufmkolk S, Dabauvalle MC, van de Linde S, et al.

rapidSTORM: accurate, fast open-source software for localization microscopy. Nat Methods.

2012;9(11):1040-1. doi: 10.1038/nmeth.2224. PubMed PMID: 23132113.

 SUPPLEMENTARY FIGURE LEGENDS

Figure S1. GFP-HeV M protein accumulation colocalizes with FC-DFC marker UBF1 but

not GC markers.

HeLa cells were transfected to express GFP-HeV M and fixed at 24 h p.t. with 4%

paraformaldehyde before immunostaining for the nucleolar markers UBF1 (FC-DFC localization)

and NPM1 and nucleolin (GC localization). Yellow boxes are magnified in the zoom panel.

Hoechst 33342 (blue) was used to identify DNA/nuclei.

Figure S2. FLAG-HeV M binds Treacle and localizes to sub-nucleolar compartments. (A)

HeLa cells transfected to express FLAG-HeV M were fixed with 4% paraformaldehyde and treated

with or without proteinase K, before immunostaining for FLAG (green) and Treacle (red). Yellow

boxes are shown magnified in the zoom panel. Hoechst 33342 (blue) was used to identify

DNA/nuclei. (B) 293 FLP-In™ cells stably transfected to enable inducible expression of 3xFLAG-

HeV M or 3xFLAG-HeV M-K258A were treated with tetracycline (+Tet) to induce expressions,

or not treated (-Tet), for 24 h before lysis and IP for FLAG; IPs were analysed by IB using the

indicated antibodies.

Figure S3. FC-DFC accumulation of NiV M protein decreases over time.

(A) HeLa cells transfected to express GFP-NiV M WT or K258A proteins were analyzed live at

8, 16, 24, 48, and 72 h p.t. by CLSM. Images representative of major phenotypes are shown for

each condition, with yellow boxes magnified in the zoom panel.

(B) Images such as those in (A) were used to determine the percentage of cells with apparent FC-

DFC accumulation. The number of cells analyzed to determine the percentage for each sample is

indicated on the graph. n ≥ 18 cells were analysed for each time point.

<u>Figure S4.</u> Inhibition of rRNA biogenesis by HeV M protein correlates with dynamic

localization to FC-DFC. HeLa cells were transfected to express the indicated proteins before

addition of EU reagent. Sample were labeled to detect nascent RNA (EU fluorescence) and imaged

by CLSM. EU fluorescence in the nucleoli of GFP-positive cells relative to that in non-GFP

expressing cells in the same sample (mean relative EU fluorescence ± S.E.M., n ≥ 22 cells) ** $p<$

0.01; ns, non-significant.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8