GreenMemTraX: A User-Friendly Software for Super-Resolution Analysis of Membrane Protein Dynamics and Nanoscale Organization in Plants

Leander Rohr[1], Alexandra Ehinger[1], Luiselotte Rausch[1], Nina Glöckner Burmeister[2], Alfred J. Meixner[3], Klaus Harter[1], Birgit Kemmerling[1], and Sven zur Oven-Krockhaus[1,3,*]

1 Center for Plant Molecular Biology (ZMBP), University of Tübingen, 72076 Tübingen, Germany

2 Department of Biomedicine, Aarhus University, 8000 Aarhus, Denmark

3 Institute for Physical and Theoretical Chemistry, University of Tübingen, 72076 Tübingen, Germany

*Correspondence: Sven zur Oven-Krockhaus (sven.zur-oven-krockhaus@uni-tuebingen.de)

ABSTRACT

Super-resolution microscopy provides deep insights into the dynamics of single membrane proteins and the nanoscale organization of plasma membrane proteins in living cells. However, in addition to plant-specific challenges in imaging applications, the elaborate analysis of single-molecule data complicates its widespread use in plant cell biological research. With GreenMemTraX, we provide a user-friendly, all-in-one, open-source software package that guides users through the steps of localization, tracking, mobility, and cluster analyses of plant plasma membrane proteins based on single particle tracking Photoactivated Localization Microscopy (sptPALM) data. GreenMemTraX dramatically simplifies data management following the FAIR (Findable, Accessible, Interoperable, Reusable) principles and increases the ease of use and reproducibility of the evaluated results. In addition, batch processing with detailed parameter output is available for all analysis paths, maintaining the consistency of raw data with the calculated results. The applicability of GreenMemTraX is demonstrated by evaluating sptPALM data for two representative plant plasma membrane proteins, namely the brassinosteroid receptor Brassinosteroid Insensitive 1 (BRI1) and the Receptor-Like Protein 44 (RLP44). Both were tested with the photoconvertible or photoactivatable fluorophores mEos3.2, PA-GFP, and PATagRFP and expressed in the *Nicotiana benthamiana* transient expression system and in stably transformed *Arabidopsis thaliana* plants. To facilitate plant applications,

34 **GreenMemTraX provides a number of preset parameter settings to address**
35 **plant-specific obstacles.**

36 **Keywords:** GreenMemTraX, software, super-resolution microscopy, localization,
37 single particle tracking, sptPALM, photoconvertible fluorophores, photoactivatable
38 fluorophores, cluster analysis, plasma membrane proteins, BRI1, RLP44, plant

INTRODUCTION

The plasma membrane (PM) is one of the most important compartments, as it controls
both homeostasis and mass transport and is the primary mediator of signals into or
out of the cell. Most of these processes are controlled by membrane proteins.
Analyzing their dynamics, organization and function is a fascinating area of research
in plants and life sciences in general. Only a few techniques allow the in vivo analysis
of membrane proteins with high spatiotemporal resolution. Suitable microscopy
techniques include fluorescence recovery after bleaching (FRAP) (Axelrod et al., 1976;
Peters et al., 1974), fluorescence correlation spectroscopy (FCS) (Magde et al., 1972),
and single particle tracking (spt) (Manzo and Garcia-Parajo, 2015) of labelled
membrane proteins, the latter being mainly driven by the advent of super-resolution
technologies such as photoactivated localization microscopy (PALM) (Betzig et al.,
2006; Hess et al., 2006). Application of sptPALM (Manley et al., 2008) is comparably
underrepresented in plant cell biological research because of the structural
peculiarities of plant cells having a cell wall.

Since sptPALM is a single-molecule microscopy technique, it depends on the detection
of the fluorescence signal from individual fluorophores with sufficient contrast. This
entails high demands on the imaging hardware and the brightness of the fluorophores.
Commonly used very bright and stable fluorophores in animal cell biology, such as
Alexa Fluor or Janelia Fluor dyes, pass through the PM into the cell and can be bound
to proteins of interest using a variety of methods (Lelek et al., 2022). However, these
organic dyes cannot pass the plant cell wall, so plant scientists predominantly rely on
genetically encoded fluorophores fused to the proteins of interest. Consequently, the
number of fluorophores usable for sptPALM in plant approaches is limited (Hosy et al.,
2015; McKenna et al., 2019).

The best contrast for imaging membrane proteins is usually achieved in animal cells
under total internal reflection (TIRF) as their PM can be placed directly above the
coverslip. However, due to the additional, about 1 to 10 μm thick cell wall of plant cells,
the plant PM is not in the optimal range for TIRF (below 150 nm). Therefore, alternative
illumination methods such as highly inclined thin illumination (HiLo) (Tokunaga et al.,
2008) or variable angle epifluorescence microscopy (VAEM) (Konopka and Bednarek,
2008) must be used, which in turn have detrimental effects on the signal-to-noise ratio.

These technical difficulties have contributed to the fact that dynamic analysis of
membrane proteins in plant cells at very high resolution has only recently taken off
(Bayle et al., 2021; Gronnier et al., 2019; Hosy *et al.*, 2015; Jaillais and Ott, 2020;
Martiniere and Zelazny, 2021; McKenna *et al.*, 2019). With the improvement of
technical possibilities in microscopy and the increasing acceptance of single particle
tracking, other analytical methods such as cluster analysis of proteins in the PM of
plant cells are now coming into focus, which, in addition to dynamic parameters such
as diffusion coefficients, can also provide information on the spatial distribution of
proteins in cell membranes (Gronnier et al., 2017; Perraki et al., 2018; Smokvarska et
al., 2023).

Despite the progress made in recent years, the analysis of sptPALM data is a
complicated, multistep process. First, the positions of the individual labelled
membrane proteins in each image must be determined with high precision
(localization), followed by the assignment of localizations across multiple images to
trajectories (tracking). The mobilities or proportions of mobile/immobile proteins are
then calculated from the analysis of these trajectories. Subsequently, the number,
shape, and size of crowded regions in the membrane are determined based on the
localizations or the tracks (cluster analysis). Each of these steps actually represents a
separate area of research. Localization depends on physical camera parameters and
localization algorithms that must incorporate noise statistics for the low photon counts
typical of single-molecule microscopy. Tracking can be performed with a variety of
algorithms, and the chosen parameters have a large impact on the data evaluation,
representation and thus their interpretation. Mobility analysis is initially performed
using mean square displacement (MSD) plots (Qian et al., 1991). Still, this method is
not always applicable and is highly dependent on the parameters used. Combining
sptPALM data with cluster analysis is a relatively recent development; several

methods having advantages and disadvantages, leading to different data evaluation,
representation and interpretation. This procedural complexity and challenges prevent
broader application of super-resolution techniques such as sptPALM in plant cell
biology.

Although practical guides have been published recently (Bayle *et al.*, 2021), there is
still a lack of all-in-one and comprehensive software packages that guide the user
through all analyses without excessive specialized knowledge. The available software
is usually limited to or specializes in only one of the above sub-steps, with compatibility
problems between the different solutions. Furthermore, the individual software
modules are typically designed for broad use but are predominantly adapted to animal
research. In addition, the large number of adjustable parameters require advanced
bioinformatics and microscopy expert knowledge. Focusing on plant PM dynamics and
organization allows concentrating on key parameters, for ease of use and
transparency of the analyses.

In this work we present an all-in-one comprehensive open-source software package,
named GreenMemTraX, guiding the scientist user-friendly and intuitively through the
steps of sptPALM-based localization, tracking, mobility- and cluster analyses of
membrane proteins in plant cells. Moreover, the storage of the sptPALM data and
metadata follow FAIR (Findable, Accessible, Interoperable and Reusable) guiding
principles (Wilkinson et al., 2016) for scientific data management and stewardship. We
explored three genetically encoded fluorophores (mEos3.2, PA-GFP and PATagRFP)
of different photophysical characteristics, highly suitable for sptPALM studies in living
plant cells. The PA-GFP and PATagRFP pair also enable dual-color sptPALM
applications in the future. The applicability of GreenMemTraX is exemplified with the
temporal and spatial dynamics of two representative plant membrane proteins, the
brassinosteroid receptor Brassinosteroid Insensitive 1 (BRI1) and the Receptor-Like
Protein 44 (RLP44).

124 RESULTS

125 Design and properties of GreenMemTraX
Main principles

GreenMemTraX is run as an executable program or as an application in MATLAB
(Mathworks). The entire code is available on GitHub for maximum transparency and
the accompanying manual answers any analysis-specific questions (both will be
published after peer-review). For each implemented main analysis step (location,
tracking, cluster analysis) an extensive literature research was performed to identify
and test the most suitable method or algorithm for GreenMemTraX. As already
mentioned in the introduction, existing software applications usually focus on a specific
part of the analysis, which is processed there in detail. For the integration into
GreenMemTraX, we choose software modules that deliver robust results quickly and
that have achieved good results in peer-reviewed evaluations (Chenouard et al., 2014;
Nieves et al., 2023; Sage et al., 2019). The published codes were partially adapted to
the requirements in GreenMemTraX; alternatively, new software code was written
following the steps described in the respective publications (see below for details).

Localization
The sptPALM raw data usually consists of time series images that are saved as TIFF
image stacks. Typical formats are 100x100 to 400x400 pixels (corresponding to 10x10
to 40x40 µm) with 2 000 to 10 000 images and a frame rate between 20 and 50 Hz.
These parameters can vary depending on the specific experiment but are mainly
governed by the optimal magnification of the imaging system for single molecule
detection, the choice of the frame rate that achieves sufficient contrast, the size of the
imaged PM region of interest, and the photophysical characteristics of the chosen
fluorescent proteins (FPs). To better represent the biological variance, many short
movies of a tracked membrane protein from a multitude of different cells are better
than a few long movies of only a few cells.

In the first step, the fluorophore-tagged proteins are localized in each image with high
precision using algorithms developed for super-resolution microscopy. For this
purpose, we adapted the core of the SMAP software (Ries, 2020) for 2D localization
only. Its algorithm for determining single molecule positions uses a robust maximum
likelihood estimation (MLE) of Gaussian point spread function (PSF) models, which
was shown to be very accurate (Sage *et al.*, 2019) and also uses the processing unit
of a graphics card for outstanding processing speed. In addition, the fitter can also
account for the pixel-specific noise of commonly used complementary metal-oxide
semiconductor (CMOS) cameras. Starting in the first tab of the software, after

supplying the raw data files, the user can start the localization process for a list of files

using default parameters (Figure 1A). An auxiliary window can be opened to check the

performance of the algorithm on selected single images and adjust the parameters if

necessary. At the end of the localization process, three histograms show (i) the PSF

sizes, (ii) photons and (iii) localization accuracy for all detected and fitted fluorescence

spots. These histograms can be used to check the quality of the raw data and exclude

outliers from further analysis, for example, localizations with unusually large PSFs due

to poor focusing or aggregation artefacts. After review and possible adjustment or

removal of invalid data, the molecule positions can be passed on to the next analysis

step.

Figure 1. Graphical user interface of GreenMemTraX, with tabs for localization, tracking and
mobility analysis, analysis of nanoscale organization and batch processing.

(A) Localization tab. After loading in the sptPALM files (1), the localization procedure is executed based
on user-defined parameters (2), with its results visualized as three histograms (PSF sizes (3), photon
numbers (4) and localization accuracy (5)) for all detected and fitted fluorescence spots. This serves as
quality control and allows the user to filter out, e.g., badly localized single molecule positions. The
selected files and their remaining localizations can then be transferred to the next tab, the **(B)** Tracking

tab. Here, the track building algorithm uses the specified parameters (6) to connect the detected
localization to trajectories, which are displayed in the overview image (7) and a user-defined magnified
view (8). Different layers and visualization options can be chosen (9), and the resulting images stored
as vector graphics. A binary mask can be defined for each file (10) to exclude tracks (for example, those
outside the cell area) from further analysis. The mobility analysis via MSD plots is also performed on
this tab, based on user-specified parameters (11), visualized as the mean MSD plot (12), averaging the
MSD data of the selected file(s), and the distribution of logarithmized, individual diffusion coefficients
(13), as a collection of derived diffusion coefficients from fitting the MSD curves of each individual track.
The fitting results representing the average diffusion coefficient are displayed below and might differ
considerably when multiple diffusive populations are present. Selected files can then be transferred to
the next tab, the **(C)** Cluster Analysis tab. The user can choose from several clustering algorithms (14)
and set their specific parameters (15). The visualization of the results (16) depends on the chosen
method and can be adjusted with several options (17), (18). For some algorithms, parameter estimates
can be calculated (19) based on the chosen data. **(D)** Batch Analysis tab. All user-specified parameters
from all previous tabs are stored in a systematic overview window (20) that can be opened from any
tab. Input files for batch processing are also added here and are shown as tables (21) for inspection
and the assignment of noise files and masks. The entire file list can then be automatically processed,
beginning with the localization procedure and may include tracking, mobility and cluster analysis as
chosen by the user, summarized in (22). All parameters and results are stored in a comprehensive
spreadsheet file, a section of which is shown as the inset (23). Additional options for all tabs are detailed
in the GreenMemTraX manual.

Tracking

In the second tab of GreenMemTraX **(Figure 1B)**, the individual protein localizations
are linked through time to obtain trajectories or tracks. The generally low localization
density in sptPALM data allows for a simple but very fast tracking algorithm that can
also bridge detection gaps typical of single-molecule microscopy. The corresponding
code was adapted from a MATLAB program published by Jean-Yves Tinevez (Tinevez,
2011)**.** The principle of this simple Linear Assignment Problem (Simple LAP) tracker is
also part of his software TrackMate (Tinevez et al., 2017). After performing the
tracking, the resulting trajectories can be visualized (Figure 1B, upper plots), e.g.,
coloring them according to their duration, displacement, mobility and other
characteristics. In case of obvious connection errors, the tracking can be repeated with
adjusted track building parameters.

Mobility analysis

The tracking tab of GreenMemTraX also provides mean square displacement (MSD)
analysis, which is the most widely used method for extracting diffusion coefficients and
motion patterns for this type of data (Manzo and Garcia-Parajo, 2015). A more detailed
description of this method was described by (Saxton and Jacobson, 1997). Only
selected tracks with a certain minimum length are forwarded for the analysis, removing
short track artefacts that originate predominantly from background fluorophores. The
mean MSD over all tracks is calculated and the diffusion coefficient estimated via a
linear fit (Figure 1B, lower left plot) that commonly only includes the first few points of
an MSD curve. While this type of ensemble analysis gives a first impression of protein

mobility, a more detailed analysis is additionally performed by calculating MSD curves
for each track, thus estimating individual diffusion coefficients for each detected
membrane protein. Their distribution can then be plotted as a histogram (Figure 1B,
lower right plot), revealing, among other things, the existence of subpopulations that
were hidden in the mean MSD plot. Due to the small number of data points for
individual tracks, a goodness-of-fit threshold value (the adjusted R^2) can be specified
for the linear fit to reject inconclusive results. In addition to storing the peak lg(D) value
for each file examined, the relative proportion of multiple populations can also be
estimated. Performing the MSD analysis adds new options to the tracking images,
such as coloring tracks based on their mobility or splitting them visually into mobile
and immobile tracks.

Cluster analysis
While mobility analysis provides information about the dynamics of membrane
proteins, their spatial organization can also be obtained by cluster analysis based on
the single molecule data. This is performed in a separate tab of GreenMemTraX,
where various computational methods can be applied (Figure 1C). Voronoi tessellation
is performed as in the SR-Tesseler software (Levet et al., 2015) or in ClusterViSu
(Andronov et al., 2016). Other methods include density-based spatial clustering of
applications with noise (DBSCAN) (Ester et al., 1996) and the recently introduced
nanoscale spatiotemporal indexing clustering (NASTIC) (Wallis et al., 2023). So far,
however, no method has become firmly established in cell biological research, partly
because the size, shape, and density of membrane clusters could not be satisfactorily
validated by an independent technique. In addition, the above algorithms define cluster
sizes differently, making it difficult to provide absolute values for the cluster
dimensions. Nevertheless, comparative analyses of membrane protein organization
are possible when they are performed with the same method. When applying the
different clustering methods, the user can base the analysis on two different data sets:
(i) all localized fluorophore positions in the sample or (ii) the centroids of the assembled
tracks. Option (i) disregards the assignment of localizations to individual membrane
proteins, which can result in varying numbers of cluster points for each protein, while
option (ii) assigns each detected membrane protein to one cluster point, at the
expense of a smaller set of points. This choice largely depends on the available data
quantity, but option (ii) should be generally preferred for its more consistent
assignment. Only the NASTIC method is based solely on the protein tracks

themselves, since their spatial overlap is used for cluster analysis. GreenMemTraX
implements all the above algorithms, so the users can follow their preferences or
compare different algorithms. Again, the results of the applied analysis are visualized,
can be color-coded with different settings and exported for publication.

Batch processing
The last tab contains the batch analysis (Figure 1D), which automatically processes
the raw sptPALM data selected by the user and then stores all analysis results in a
consistent format. A summary of all parameters used for the individual steps in the
analysis of the membrane protein of interest are listed in an intuitive selection
structure. If necessary, parameters can still be changed here, and the subsequent
batch analysis is performed based on these parameters collectively for all analysis
steps (localization, tracking, mobility and cluster analysis). The parameter list is also
attached to the collected results, so that all steps and settings can be traced back
according to the FAIR principle.

Fluorophores for in planta sptPALM analysis

The choice of suitable fluorophores is a crucial step in all fluorescence imaging
techniques. The requirements are even higher for single-molecule techniques
including sptPALM, as the signal of individual fluorescent molecules must be detected
with sufficient contrast. In plant cells, the rather impenetrable cell wall largely
precludes the use of organic dyes and quantum dots, so that genetically encoded FPs
must be used, which are fused translationally to the membrane protein of interest.
Conversely, FPs exhibit comparatively low photostability and brightness. Moreover,
sptPALM analyses of membrane proteins require photoactivatable or -convertible FP
variants, where the density of visible fluorophores per image is controlled by an
additional activation laser, further limiting the pool of applicable FPs. Genetic
constructs encoding photoconvertible mEos2 fusions have been useful for sptPALM-
based analyses of membrane proteins in plant cells (Gronnier *et al.*, 2017; Hosy *et al.*,
2015; Smokvarska *et al.*, 2023). However, (Zhang et al., 2012) showed that mEos2
has the disadvantage that it easily forms oligomers and aggregates when fused to
membrane proteins. We therefore implemented the improved version mEos3.2 that is
truly monomeric, is brighter, matures faster and displays higher photon budget and
label density (Zhang *et al.*, 2012). The native green form of mEos3.2 can be converted
to a red form using light of ~400 nm wavelength, allowing to adjust the density of the

visible fluorophores in the red imaging channel. However, as mEos3.2 occupies both
the green and red part of the spectrum, it is not compatible to combine it with other
FPs for potential dual-color applications. Based on their photophysical characteristics
(Table 1), we propose the use of photoactivatable (PA-) GFP (Patterson and Lippincott-
Schwartz, 2002) and PATagRFP (Subach et al., 2010) as additional fluorophores for
sptPALM applications in plant cells. Both PA FPs are non-fluorescent in their
respective native forms and can be converted to their spectrally distinct fluorescent
forms with light of ~400 nm, enabling simultaneous imaging of differently labelled
proteins in two color channels. The three fluorophore genes were codon-optimized for
their use in plant cells (Supplemental Table 1).

Table 1. Basic spectral characteristics of the fluorophores mEos2, mEos3.2, PA-GFP and
PATagRFP.

For the mEos fluorophores, both the native green (G) and the photoconverted red (R) form are listed,
while for PA-GFP and PATagRFP, only the activated forms are presented. A more detailed compilation
of their photophysical characteristics can be found in Supplemental Table 2.

	Excitation maximum (nm)	Emission maximum (nm)	reference
mEos2 (G)	506	519	(McKinney et al., 2009)
mEos2 (R)	573	584	(McKinney *et al.*, 2009)
mEos3.2 (G)	507	516	(Zhang *et al.*, 2012)
mEos3.2 (R)	572	580	(Zhang *et al.*, 2012)
PA-GFP	504	517	(Patterson and Lippincott-Schwartz, 2002)
PATagRFP	562	595	(Subach *et al.*, 2010)

Biological applications using GreenMemTraX

Introduction of the intrinsic PM proteins used for proof-of-principle sptPALM analyses

In order to demonstrate the applicability of GreenMemTraX for sptPALM analyses, two
well-described intrinsic membrane proteins were used, namely the brassinosteroid
(BR) receptor BRASSINOSTEROID INSENSITIVE 1 (BRI1) and the RECEPTOR-
LIKE PROTEIN 44 (RLP44).

BRI1 initiates well-understood signaling pathways in plant cells. Upon binding of BR
to BRI1's extracellular domain, its interaction with the co-receptor BRI1-ASSOCIATED
KINASE 1 (BAK1) is enhanced. This leads to a re-arrangement of proteins within the
complex, eventually resulting in auto- and transphosphorylation of their kinase
domains and its full signaling activity (Gou and Li, 2020; Wolf, 2020). The outcomes
of BR activation of the BRI1/BAK1 complex are on the one hand the differential
regulation of BR-responsive genes via a nucleo-cytoplasmic signaling cascade (Mora-

Garcia et al., 2004; Vert and Chory, 2006; Yin et al., 2005; Zhu et al., 2017) and on the
other hand a rapid acidification of the apoplast via the activation of PM-resident P-type
proton pumps (Caesar et al., 2011; Grosseholz et al., 2022; Witthöft et al., 2011)

RLP44 is proposed to be a cell wall integrity sensor that controls cell wall homeostasis
by interplay with BRI1 and BAK1 (Wolf et al., 2012; Wolf et al., 2014). In fact, we were
recently able to demonstrate the existence of a ternary RLP44/BRI1/BAK1 complex in
the PM of living plant cells using a spectral Förster resonance energy transfer (FRET)
and FRET-Fluorescence-lifetime imaging microscopy (FLIM) approach (Glöckner et
al., 2022). Additionally, RLP44 has been linked to phytosulfokine signaling. The
corresponding receptor Phytosulfokine Receptor 1 (PSKR1) is also proposed to form
a complex with RLP44 and BAK1 (Gomez et al., 2021; Holzwart et al., 2018).

GreenMemTraX analysis of BRI1 receptor dynamics and nano-structured organization
in the Nicotiana benthamiana transient expression system
For initial assessment, fluorescent fusion proteins were transiently expressed in
Nicotiana benthamiana (*N. benthamiana*) epidermal leaf cells. Transient expression
assays also provide robust results comparable with stably transformed Arabidopsis
lines as shown for the photoswitchable fluorophore mEos2 (Gronnier *et al.*, 2017;
Perraki *et al.*, 2018).

Figure 2. All BRI1-fluorophore fusions accumulate in the PM of epidermal cells of transiently transformed *Nicotiana benthamiana* leaves and their spatiotemporal properties can be analysed by GreenMemTraX.

(A) Representative images of sptPALM movies, showing, from left to right, single fusion proteins of BRI1-mEos3.2, BRI1-PA-GFP and BRI1-PATagRFP. The BRI1 fusion proteins were expressed via the native BRI1 promoter (*pBRI1*). Scale bar = 1 μM. **(B)** Representative magnifications of areas containing tracks generated by GreenMemTraX (random coloring per track) after localization of the BRI1 fusion proteins, with panels ordered as above. Scale bar = 100 nm. **(C)** Diffusion coefficients for the indicated BRI1 fusion proteins in the PM obtained with GreenMemTraX. For statistical evaluation, the data were checked for normal distribution and then analyzed applying the Kruskal-Wallis test followed by the Steel-Dwass post-hoc test, with n ≥ 17 for each fusion protein derived from at least six plants, measured on at least four different days. For detailed statistics see Supplemental Table 3. **(D)** The track data was used for cluster analysis in GreenMemTraX, exemplified by BRI1-mEos3.2 clusters that were assigned using the NASTIC algorithm. The top image represents one cell with random coloring per track and cluster boundaries highlighted in white. Scale bar = 2 μm. The bottom is a magnified view of the respective marked region in the top image. Scale bar = 500 nm.

Endogenous promotor driven constructs encoding the BRI1 and RLP44 fusions with the fluorophores mEos3.2, PA-GFP, and PATagRFP (see Materials) were transiently expressed in *N. benthamiana* epidermal leaf cells with the even distribution of activated fluorophores (Figure 2A) necessary for a comprehensive evaluation of sptPALM data (Bayle *et al.*, 2021).

Using GreenMemTraX (for detailed parameters see Supplemental Table 4), the data were processed, and trajectories generated (Figure 2B). After MSD analysis, the diffusion coefficients were calculated (Figure 2C).

Although the three fluorophores are sptPALM-optimized versions of different
precursors from various marine organisms, all three BRI1 fusion proteins showed
comparable diffusion coefficients in *N. benthamiana* epidermal leaf cells (Figure 2C).
This demonstrates that all three fluorophores can be reliably used for sptPALM studies
in plant cells. In live cell microscopy, the best signal-to-noise ratio was obtained with
mEos3.2, due to its excellent photophysical properties (Supplemental Table 2). On the
other hand, PA-GFP and PATagRFP can be combined - due to their non-overlapping
spectra - for dual-color sptPALM experiments.

As shown in Figure 2D, GreenMemTraX allows the use of sptPALM single-molecule
data to group BRI1 protein tracks into clusters.

In conclusion, we demonstrate that the transient expression system in *N. benthamiana*
is a suitable choice for testing mEos3.2, PA-GFP and PATagRFP fusions of PM
proteins in plant cells.

GreenMemTraX analysis of RLP44 dynamics and nano-structured organization in PM
of transgenic in Arabidopsis thaliana
To the best of our knowledge, neither mEos3.2 nor PATagRFP were used before for
sptPALM experiments in transgenic plants. In contrast, PA-GFP has been applied
previously (Martiniere et al., 2012; McKenna *et al.*, 2019).

For the generation of stable Arabidopsis lines via *Agrobacteria*-mediated
transformation, endogenous promoter (here *pRLP44*) constructs with the *RLP44*
cDNA and the codon-optimized versions of our fluorophores (see Methods) were used.

As shown in Figure 3A, all three RLP44 fusions successfully accumulated in
transgenic *Arabidopsis thaliana* (*A. thaliana)* hypocotyl cells in densities optimal for
sptPALM data evaluation. Using GreenMemTraX, trajectories were generated (Figure
3B) and the diffusion coefficients were calculated via MSD analysis. All three RLP44
protein fusions resulted in comparable diffusion coefficients (Figure 3C). We noticed
minor fluorophore-dependent differences in the RLP44 diffusion coefficients, likely
reflecting that the plant cells for the data sets may have had different cellular and
physiological stages.

All tagged RLP44 proteins allowed data processing suitable for cluster analysis using
GreenMemTraX (Figure 3D).

In summary, we demonstrate that mEos3.2-, PA-GFP- and PATagRFP-tagged
membrane proteins are suitable for sptPALM experiments in transgenic *A. thaliana*
and allow generation of data that can easily be analyzed by GreenMemTraX.

409 *GreenMemTraX application examples to detect significant changes in protein mobility*
410 Most sptPALM analyses rely on comparing dynamic parameters of proteins, for
411 example, in response to environmental conditions such as hyper-osmotic stress
412 (Martiniere et al., 2019). We conducted sptPALM measurements in different light

conditions. GreenMemTraX-based data evaluation using transgenic *A. thaliana* seedlings accumulating RLP44-mEos3.2 under the control of the *RLP44* promoter (Figure 4) were performed in seedlings either grown in white light or darkness to determine how the developmental state (photomorphogenic versus skotomorphogenic) influences the dynamic properties of RLP44-mEos3.2 in the PM. As shown in Figure 4A, RLP44-mEos3.2 moved significantly faster in in the PM of cells of the upper part of dark-grown hypocotyls compared to the lower part. The diffusion coefficient of RLP44-mEos3.2 in the PM of hypocotyl cells of light-grown seedlings, where a differentiation of different cell types along the axis is hardly possible, was in between the diffusion coefficients gained from the etiolated seedlings. These in planta generated data show that tissue-specific differences of the RLP44-mEos3.2 dynamics can be detected and analyzed by GreenMemTraX.

The effect of other environmental factors on protein mobility can also be evaluated by GreenMemTraX. Here we chose the hyper-osmotic stress response of plant cells after addition of sorbitol, a non-toxic, widely used, non-metabolic osmolyte already applied for sptPALM (Hosy *et al.*, 2015). Sorbitol induces a clear cell volume loss (plasmolysis) when used at a concentration of 300 mM. Under milder hyper-osmotic conditions (100 mM), the cells experience a reduction in turgor without major changes in their volume (Martiniere *et al.*, 2019). Thus, the increasing concentration of sorbitol from 100 to 300 mM induces a progressive separation of the PM from the cell wall which changes the dynamics of almost all intrinsic PM proteins (Hosy *et al.*, 2015). Based on this knowledge, treatments with 100 mM and 300 mM sorbitol or mock treatment were performed with light-grown and dark-grown seedlings for 20 minutes. Then, the sptPALM data for RLP44-mEos3.2 were measured in hypocotyl cells of light-grown seedlings and cells of the upper hypocotyl of dark-grown seedlings and evaluated using GreenMemTraX. Whereas the concentration of 100 mM sorbitol did not significantly affect the diffusion coefficient of RLP44-mEos3.2 compared to the mock treatment (0 mM), 300 mM sorbitol significantly increased its dynamic behavior in all measured cells (Figure 4B).

These examples represent common analyses of sptPALM data in plant research, and demonstrate the applicability of GreenMemTraX, which was used to perform all analysis steps, including localization, tracking and protein mobility analysis.

Figure 4. Evaluation of sptPALM data by GreenMemTraX reveals tissue-specific and environmental differences in the dynamics of RLP44-mEos3.2 in epidermal cells of the *Arabidopsis* hypocotyl.

(A) Diffusion coefficients of RLP44-mEos3.2 in epidermal hypocotyl cells of light-grown and dark-grown *A. thaliana* seedlings. The locations of the recorded measurements are indicated by arrows in the corresponding seedling sketches. For dark-grown seedlings, the center of the distance from the apical hook to the seed was used as the reference point. Based on this, the centers to the apical hook and to the seed were used as selected regions, named 'upper' and 'lower', respectively. For statistical evaluation, the data were checked for normal distribution and equal variances and the statistical analysis was performed using the Kruskal-Wallis test, with n ≥ 30, from nine plants, measured on three different days. For detailed statistics see Supplemental Table 3. **(B)** Diffusion coefficients of RLP44-mEos3.2 recorded from epidermal hypocotyl cells of light-grown or dark-grown *A. thaliana* seedlings after 20 min. treatment with sorbitol solution of 100 mM or 300 mM or mock-treatment (0 mM). For the recording of the data in the dark-grown seedlings the upper part of the hypocotyl cells was chosen (see (A)). For statistical evaluation, the data were checked for normal distribution. The statistical analysis was performed using the Kruskal-Wallis test followed by the Steel-Dwass post-hoc test, with n ≥ 14, from three different plants. For detailed statistics see Supplemental Table 3.

DISCUSSION

GreenMemTraX is an all-in-one software package for sptPALM data evaluation, optimized for plant cell biologists

With GreenMemTraX, we provide a user-friendly open-source software package that guides users through the steps of localization, tracking, analysis of mobility and nanoscale organization of membrane proteins on the basis of sptPALM data. GreenMemTraX is tailored for use by cell biologists who want to perform sptPALM analyses in plant cells. Due to the cell wall, only areas close to the microscope slide can be analyzed in plants such as the plasma membrane and plasma membrane associated proteins. This narrows down the large number of adjustable parameters that are usually applied in sptPALM research in animal or human cell cultures and tissues and will significantly lower the hurdle for sptPALM application in plant cell biology.

GreenMemTraX has a default set of analysis parameters that is well suited for plant PM proteins. However, all settings can be adjusted by the users to their specific experimental requirements. Moreover, batch processing with detailed parameter and data output is available for all analysis paths, maintaining the consistency of the sptPALM data with the calculated results. The detailed but clearly structured parameter organization also allows to investigate the influence of individual parameters in the overall analysis process on the results, providing a method to evaluate the reliability of the analyses. Importantly, the data management procedure of GreenMemTraX follows the FAIR guiding principles for scientific data management and stewardship. Due to its open-source characteristics, GreenMemTraX also allows users to customize or extend the program according to their needs.

GreenMemTraX was programmed in MATLAB, a software language that is easy to work with and widely used in academia. Care was taken to make the code straightforward to understand and all programming steps are commented to make it easier to insert changes or add extensions, which could also make GreenMemTraX interesting for scientists working with animal or human cell cultures.

PATagRFP, PA-GFP and mEos3.2 are suitable FPs for sptPALM analyses in living plant cells

The rather impenetrable cell wall almost completely excludes the use of organic dyes and quantum dots for sptPALM applications in plant cells. Plant scientists therefore rely on genetically encoded photoactivatable or photoconvertible FPs fused to the protein of interest. With mEos3.2, PA-GFP and PATagRFP, three different FPs are now available that do not aggregate, deliver satisfactory signal-to-noise ratios during imaging and are therefore applicable for sptPALM applications both in a transient transformation system (*N. benthamiana*) and in stably transformed *A. thaliana* plants. We expect that these fluorophores will also work in other plant species. Of particular interest is the combination of PA-GFP and PATagRFP, which will enable dual-color sptPALM in the future for simultaneous recording and spatiotemporal analysis of two different fusion proteins in the PM. Because of its superior photophysical properties, we recommend the use of mEos3.2 for single FP fusion protein analyses. Especially when using mEos3.2, but also the other FPs, the sptPALM technology shows its superiority in terms of spatiotemporal resolution of membrane proteins in living plant cells compared to other methods such as FRAP and FCS (Bayle *et al.*, 2021).

GreenMemTraX allows the rapid in vivo analysis of membrane proteins in transiently and stably transformed plant cells

We used two functionally well-studied membrane proteins, namely BRI1 and RLP44, for our proof-of-principle experiments regarding the usability of the three FPs and applicability of GreenMemTraX. Both proteins were evaluated in either transiently transformed *N. benthamiana* epidermal leaf cells or the epidermis of the hypocotyl of stable transgenic *A. thaliana* seedlings. The accumulation of the BRI1 and RLP44 fusion proteins were driven by the respective endogenous promoters. Strong over-accumulation of fusion proteins driven by constitutively active promoters leads to overcrowded sptPALM movies. Analysis of these movies results in misconnected tracks that affects the entire GreenMemTraX analysis causing inadequate results in subsequent steps. Therefore, we recommend using weaker promoters whenever possible.

GreenMemTraX offers different algorithms (Voronoi tessellation, DBSCAN, NASTIC) for the analysis of the nanoscale organization of a given membrane protein. We

recommend the NASTIC algorithm as it is specifically designed to work with track data.
Because of the sensitivity to parameter changes, care must be taken when regarding
the results of the nanoscale evaluation output by all of these algorithms as absolute
values. However, relative comparisons of the nanoscale organization for a given
membrane protein are possible if there are no changes in the parameter settings
between the experiments.

GreenMemTraX captures developmental and physiological effects on the in vivo dynamics of membrane proteins

During the GreenMemTraX analysis of the RLP44-mEos3.2 dynamics in epidermal
cells of light-grown *Arabidopsis* hypocotyls, we observed high data variability. This led
to the hypothesis that cell-specific effects depending on the cell's position in the organ,
i.e., on its developmental or physiological state, have an influence on the membrane
dynamics of the fusion protein to be investigated. This hypothesis was substantiated
by the technically easier access to the potential positional effects in the dynamics of
RLP44-mEos3.2 in epidermal cells along the hypocotyl axis of dark-grown seedlings:
The diffusion coefficient of RLP44-mEos3.2 is significantly higher in the epidermal cells
of the upper part than in those of the lower part of the dark-grown hypocotyl.
Furthermore, the diffusion coefficient in the lower part is similar to that found in the
epidermal cells of light-grown hypocotyls. GreenMemTraX can easily capture
differences in mobilities and allows sptPALM assays in different tissues that may vary
in their membrane properties.

As previously shown by (Hosy *et al.*, 2015; Martiniere *et al.*, 2019), the treatment of
Arabidopsis root cells with increasing concentration of the osmolyte sorbitol increases
the diffusion coefficient of membrane proteins such as aquaporins. Using
GreenMemTraX, we could confirm these findings and add potential effects of the
differentiation states of the analyzed tissue. This shows that reproducible and robust
data analysis is provided by our software tool.

In summary, sptPALM becomes more easily accessible and analyzable with
GreenMemTraX. The fluorophores mEos3.2, PA-GFP and PATagRFP have proven to
be suitable for sptPALM in planta and allow analysis of membrane proteins in transient
expression systems as well as in stable transformed plants. They are suitable for
treatment experiments and applicable in different tissues allowing also developmental

assays. The fast and easy evaluation of sptPALM data by GreenMemTraX will greatly

facilitate the comprehensive investigation of the dynamics and nanoscale organization

of membrane proteins in the future.

METHODS

Plasmid construction

All expression clones were constructed using GoldenGate assembly with BB10 as the

vector (Binder et al., 2014). Promoter sequences were obtained with the help of the

Integrated Genome Browser (Freese et al., 2016). Level I modules were generated by

PCR amplification of the desired sequences and then blunt-end cloned into pJET1.2

(Thermo Fisher Scientific). Fluorophores were designed as C-terminal fusions

(D-E module) using either a glycine/serine or a glycine/alanine rich linker. The coding

sequences of BRI1 and RLP44 were constructed as B-D modules, eliminating the

need for a B-C dummy module. A full list of used constructs can be found in

Supplemental Table 5. The correctness of Level I constructs was confirmed by Sanger

sequencing. Cut-ligations for the Level II generation were performed with 40 cycles,

without using bovine serum albumin as described by (Binder *et al.*, 2014). Reactions

were transformed into TOP10 cells (Thermo Fisher Scientific), and colony correctness

was verified via restriction enzyme analysis and partial Sanger sequencing.

Plant material and growth conditions

The transgenic *A. thaliana* lines used in this study were all in the Columbia (Col-0)

background. Stable lines were generated using the Floral dip method according to

(Zhang et al., 2006). Transgenic seeds were propagated either based on the presence

of the pFAST marker by binocular visual inspection or by selection of survivors on

½ Murashige and Skoog (MS) plates containing 1 % (w/v) sucrose and 0.8 % (w/v)

phytoagar supplemented with 25 µM hygromycin. For sptPALM measurements, seeds

were sterilized with a solution of 70 % ethanol (v/v) and 0.05 % Triton X-100 for

30 minutes followed by a 10-minute treatment with absolute ethanol. Seeds were sown

on ½ MS plates (+1 % sucrose and 0.8 % phytoagar) and stratified at 4 °C for at least

24 hours. For measurements of dark-grown seedlings, seeds were exposed to light

from the growth chamber for two hours before being wrapped in aluminum foil until the

day of measurement. Light-grown seedlings were cultivated in growth chambers at

20 °C under long-day conditions (16 hours light / 8 hours dark). The duration of growth

is indicated in the respective figures. The *N. benthamiana* plants used in this study

were also cultivated under controlled conditions. Proteins were transiently expressed

using the AGL *Agrobacterium tumefaciens* strain (Lifeasible), as previously described

(Hecker et al., 2015; Ladwig et al., 2015), without the washing step with sterile water.

The plants were infiltrated with the respective construct at an OD600 of 0.1, in a ratio

of 1:1 with the silencing inhibitor p19. After watering, the plants were kept in ambient

conditions and were imaged three days after infiltration.

Sample preparation and movie acquisition

All sptPALM measurements with transiently transformed *N. benthamiana* were

performed three days after infiltration. A small leaf area was cut out, excluding veins,

and placed between two coverslips (Epredia 24x50 mm #1 or equivalent) with a drop

of water. This "coverslip sandwich" was then placed on the specimen stage, lightly

weighted down by a brass ring to help flatten the uneven cell layers, especially in

N. benthamiana. Seven-day-old *A. thaliana* seedlings were used to acquire data from

stable *Arabidopsis* lines. Depending on the experiment, either light-grown or dark-

grown plants were used. Similar to the handling of *N. benthamiana* leaf discs, the

seedlings were placed between coverslips and brass rings.

The custom-built microscope platform for sptPALM acquisition is described in detail in

the Supplemental Information. Briefly, lasers of different wavelengths and their

intensities are controlled by an acousto-optic transmission filter. A laterally translatable

lens in the excitation beam path allows to adjust the VAEM illumination of the sample

utilizing a high NA objective. The emitted light from the sample is separated from the

excitation light by a multi-band beam splitter and is detected by an sCMOS camera.

Depending on the fluorophore fusion, the following filters were inserted into the

emission beam path: (i) mEos3.2: 568 LP Edge Basic Longpass Filter, 584/40 ET

Bandpass; (ii) PA-GFP: 488 LP Edge Basic Longpass Filter, 525/50 BrightLine HC;

(iii) PATagRFP: 568 LP Edge Basic Longpass Filter, 600/52 BrightLine HC (all AHF

analysentechnik AG). The excitation power arriving at the sample was measured

(PM100D with S120C, Thorlabs) in epifluorescence mode after the objective to keep

it constant for the respective experiment sets. If necessary, photoconversion or

photoactivation was performed using 405 nm excitation at varying low intensities (for
detailed acquisition parameters see Supplemental Table 4). The magnification of the
optical system was adjusted so that the length of one camera pixel corresponds to
100 nm in the sample plane. Viable regions of interest were screened in a larger area
of 51.2 x 51.2 µm by adjusting the focal plane and the VAEM angle with a frame rate
of 10 Hz, while recording was performed with 12.8 x 12.8 µm and frame rates between
20 and 50 Hz, recording between 2 500 and 5 000 frames per movie (see
Supplemental Table 4). For each measurement day, noise files (a series of dark
images) were recorded with the corresponding frame rates.

Raw data processing and analysis with GreenMemTraX

Subsets of each experimental data set were loaded into GreenMemTraX to inspect
the quality of the data and to find appropriate parameters for each analysis step as
described above. On the Batch Analysis tab, all sptPALM raw data files that share the
same parameter set were processed together (see Supplemental Table 4 for a detailed
list of applied analysis parameters). Samples that showed significantly low numbers
of localizations or tracks compared to others in the same batch were discarded.

SUPPLEMENTAL INFORMATION
635 Document S1. Supplemental information.

FUNDING
637 Our research was supported by the German Research Foundation (DFG) via the CRC
638 1101 to A.J.M., B.K., K.H. and S.z.O.-K., and by individual DFG grants to K.H. (HA
639 2146/22, HA 2146/23). We also thank the DFG for grants for scientific equipment
640 (FUGG: INST 37/991-1, INST 37/992-1, INST 37/819-1, INST 37/965-1).

AUTHOR CONTRIBUTIONS
Conceptualization, A.J.M., B.K., K.H., and S.z.O.-K.; Software, S.z.O.-K.; Formal
Analysis, L.R. and S.z.O.-K.; Data Curation, L.R., LI.R., and S.z.O.-K.; Methodology,
S.z.O.-K.; Investigation, L.R., LI.R., and S.z.O.-K.; Writing – Original Draft, K.H., L.R.,
and S.z.O.-K.; Writing – Review & Editing, A.J.M., B.K., K.H., L.R., LI.R., and S.z.O.-K.;

Visualization, L.R. and S.z.O.-K.; Funding Acquisition, A.J.M., B.K., and K.H.;
Resources, A.E., A.J.M., and N.G.B.; Supervision, A.J.M., B.K., and K.H.

ACKNOWLEDGMENTS

649   No conflict of interest is declared.

REFERENCES

Andronov, L., Orlov, I., Lutz, Y., Vonesch, J.L., and Klaholz, B.P. (2016).
ClusterViSu, a method for clustering of protein complexes by Voronoi tessellation in
super-resolution microscopy. Sci. Rep. **6**:24084. 10.1038/srep24084.

Axelrod, D., Koppel, D.E., Schlessinger, J., Elson, E., and Webb, W.W. (1976).
Mobility Measurement by Analysis of Fluorescence Photobleaching Recovery
Kinetics. Biophys. J. **16**:1055-1069. 10.1016/S0006-3495(76)85755-4.

**Bayle, V., Fiche, J.B., Burny, C., Platre, M.P., Nollmann, M., Martiniere, A., and
Jaillais, Y.** (2021). Single-particle tracking photoactivated localization microscopy of
membrane proteins in living plant tissues. Nat. Protoc. **16**:1600-+. 10.1038/s41596-
020-00471-4.

**Betzig, E., Patterson, G.H., Sougrat, R., Lindwasser, O.W., Olenych, S.,
Bonifacino, J.S., Davidson, M.W., Lippincott-Schwartz, J., and Hess, H.F.** (2006).
Imaging intracellular fluorescent proteins at nanometer resolution. Science **313**:1642-
1645. 10.1126/science.1127344.

**Binder, A., Lambert, J., Morbitzer, R., Popp, C., Ott, T., Lahaye, T., and Parniske,
M.** (2014). A modular plasmid assembly kit for multigene expression, gene silencing
and silencing rescue in plants. PLoS One **9**:e88218. 10.1371/journal.pone.0088218.

**Caesar, K., Elgass, K., Chen, Z., Huppenberger, P., Witthoft, J., Schleifenbaum,
F., Blatt, M.R., Oecking, C., and Harter, K.** (2011). A fast brassinolide-regulated
response pathway in the plasma membrane of Arabidopsis thaliana. Plant J. **66**:528-
540. 10.1111/j.1365-313X.2011.04510.x.

**Chenouard, N., Smal, I., de Chaumont, F., Maska, M., Sbalzarini, I.F., Gong, Y.H.,
Cardinale, J., Carthel, C., Coraluppi, S., Winter, M., et al.** (2014). Objective
comparison of particle tracking methods. Nat. Methods **11**:281-U247.
10.1038/nmeth.2808.

Ester, M., Kriegel, H.-P., Sander, J., and Xu, X. (1996). A density-based algorithm for
discovering clusters in large spatial databases with noise. Proceedings of the Second
International Conference on Knowledge Discovery and Data Mining:226–231.

Freese, N.H., Norris, D.C., and Loraine, A.E. (2016). Integrated genome browser:
visual analytics platform for genomics. Bioinformatics **32**:2089-2095.
10.1093/bioinformatics/btw069.

Glöckner, N., zur Oven-Krockhaus, S., Rohr, L., Wackenhut, F., Burmeister, M.,
Wanke, F., Holzwart, E., Meixner, A.J., Wolf, S., and Harter, K. (2022). Three-
Fluorophore FRET Enables the Analysis of Ternary Protein Association in Living Plant
Cells. Plants **11**10.3390/plants11192630.

Gomez, B.G., Holzwart, E., Shi, C., Lozano-Duran, R., and Wolf, S. (2021).
Phosphorylation-dependent routing of RLP44 towards brassinosteroid or
phytosulfokine signalling. J. Cell Sci. **134**10.1242/jcs.259134.

Gou, X., and Li, J. (2020). Paired Receptor and Coreceptor Kinases Perceive
Extracellular Signals to Control Plant Development. Plant Physiol. **182**:1667-1681.
10.1104/pp.19.01343.

Gronnier, J., Legrand, A., Loquet, A., Habenstein, B., Germain, V., and Mongrand,
S. (2019). Mechanisms governing subcompartmentalization of biological membranes.
Curr. Opin. Plant Biol. **52**:114-123. 10.1016/j.pbi.2019.08.003.

Gronnier, J., Crowet, J.M., Habenstein, B., Nasir, M.N., Bayle, V., Hosy, E., Platre,
M.P., Gouguet, P., Raffaele, S., Martinez, D., et al. (2017). Structural basis for plant
plasma membrane protein dynamics and organization into functional nanodomains.
eLife **6**10.7554/eLife.26404.

Grosseholz, R., Wanke, F., Rohr, L., Glöckner, N., Rausch, L., Scholl, S., Scacchi,
E., Spazierer, A.J., Shabala, L., Shabala, S., et al. (2022). Computational modeling
and quantitative physiology reveal central parameters for brassinosteroid-regulated
early cell physiological processes linked to elongation growth of the Arabidopsis root.
eLife **11**10.7554/eLife.73031.

Hecker, A., Brand, L.H., Peter, S., Simoncello, N., Kilian, J., Harter, K., Gaudin, V.,
and Wanke, D. (2015). The Arabidopsis GAGA-Binding Factor BASIC
PENTACYSTEINE6 Recruits the POLYCOMB-REPRESSIVE COMPLEX1
Component LIKE HETEROCHROMATIN PROTEIN1 to GAGA DNA Motifs. Plant
Physiol. **168**:1013-1024. 10.1104/pp.15.00409.

Hess, S.T., Girirajan, T.P.K., and Mason, M.D. (2006). Ultra-high resolution imaging
by fluorescence photoactivation localization microscopy. Biophys. J. **91**:4258-4272.
10.1529/biophysj.106.091116.

Holzwart, E., Huerta, A.I., Glockner, N., Garnelo Gomez, B., Wanke, F., Augustin,
S., Askani, J.C., Schurholz, A.K., Harter, K., and Wolf, S. (2018). BRI1 controls
vascular cell fate in the Arabidopsis root through RLP44 and phytosulfokine signaling.
Proc. Natl. Acad. Sci. U.S.A. **115**:11838-11843. 10.1073/pnas.1814434115.

Hosy, E., Martiniere, A., Choquet, D., Maurel, C., and Luu, D.T. (2015). Super-
resolved and dynamic imaging of membrane proteins in plant cells reveal contrasting
kinetic profiles and multiple confinement mechanisms. Mol. Plant **8**:339-342.
10.1016/j.molp.2014.10.006.

Jaillais, Y., and Ott, T. (2020). The Nanoscale Organization of the Plasma Membrane
and Its Importance in Signaling: A Proteolipid Perspective(1)([OPEN]). Plant Physiol.
182:1682-1696. 10.1104/pp.19.01349.

Konopka, C.A., and Bednarek, S.Y. (2008). Variable-angle epifluorescence
microscopy: a new way to look at protein dynamics in the plant cell cortex. Plant J.
53:186-196. 10.1111/j.1365-313X.2007.03306.x.

Ladwig, F., Dahlke, R.I., Stuhrwohldt, N., Hartmann, J., Harter, K., and Sauter, M.
(2015). Phytosulfokine Regulates Growth in Arabidopsis through a Response Module
at the Plasma Membrane That Includes CYCLIC NUCLEOTIDE-GATED
CHANNEL17, H+-ATPase, and BAK1. Plant Cell **27**:1718-1729.
10.1105/tpc.15.00306.

Lelek, M., Gyparaki, M.T., Beliu, G., Schueder, F., Griffie, J., Manley, S.,
Jungmann, R., Sauer, M., Lakadamyali, M., and Zimmer, C. (2022). Single-
molecule localization microscopy (vol 1, 39, 2021). Nat. Rev. Methods Primers
210.1038/s43586-022-00161-3.

Levet, F., Hosy, E., Kechkar, A., Butler, C., Beghin, A., Choquet, D., and Sibarita,
J.B. (2015). SR-Tesseler: a method to segment and quantify localization-based super-
resolution microscopy data. Nat. Methods **12**:1065-1071. 10.1038/nmeth.3579.

Magde, D., Webb, W.W., and Elson, E. (1972). Thermodynamic Fluctuations in a
Reacting System - Measurement by Fluorescence Correlation Spectroscopy. Phys.
Rev. Lett. **29**:705-&. DOI 10.1103/PhysRevLett.29.705.

Manley, S., Gillette, J.M., Patterson, G.H., Shroff, H., Hess, H.F., Betzig, E., and
Lippincott-Schwartz, J. (2008). High-density mapping of single-molecule trajectories

with photoactivated localization microscopy. Nat. Methods **5**:155-157.
10.1038/Nmeth.1176.

Manzo, C., and Garcia-Parajo, M.F. (2015). A review of progress in single particle
tracking: from methods to biophysical insights. Rep. Prog. Phys. **78**10.1088/0034-
4885/78/12/124601.

Martiniere, A., and Zelazny, E. (2021). Membrane nanodomains and transport
functions in plant. Plant Physiol. **187**:1839-1855. 10.1093/plphys/kiab312.

Martiniere, A., Fiche, J.B., Smokvarska, M., Mari, S., Alcon, C., Dumont, X.,
Hematy, K., Jaillais, Y., Nollmann, M., and Maurel, C. (2019). Osmotic Stress
Activates Two Reactive Oxygen Species Pathways with Distinct Effects on Protein
Nanodomains and Diffusion. Plant Physiol. **179**:1581-1593. 10.1104/pp.18.01065.

Martiniere, A., Lavagi, I., Nageswaran, G., Rolfe, D.J., Maneta-Peyret, L., Luu,
D.T., Botchway, S.W., Webb, S.E., Mongrand, S., Maurel, C., et al. (2012). Cell wall
constrains lateral diffusion of plant plasma-membrane proteins. Proc. Natl. Acad. Sci.
U.S.A. **109**:12805-12810. 10.1073/pnas.1202040109.

McKenna, J.F., Rolfe, D.J., Webb, S.E.D., Tolmie, A.F., Botchway, S.W., Martin-
Fernandez, M.L., Hawes, C., and Runions, J. (2019). The cell wall regulates
dynamics and size of plasma-membrane nanodomains in Arabidopsis. Proc. Natl.
Acad. Sci. U.S.A. **116**:12857-12862. 10.1073/pnas.1819077116.

McKinney, S.A., Murphy, C.S., Hazelwood, K.L., Davidson, M.W., and Looger, L.L.
(2009). A bright and photostable photoconvertible fluorescent protein. Nat. Methods
6:131-133. 10.1038/Nmeth.1296.

Mora-Garcia, S., Vert, G., Yin, Y., Cano-Delgado, A., Cheong, H., and Chory, J.
(2004). Nuclear protein phosphatases with Kelch-repeat domains modulate the
response to brassinosteroids in Arabidopsis. Genes Dev. **18**:448-460.
10.1101/gad.1174204.

Nieves, D.J., Pike, J.A., Levet, F., Williamson, D.J., Baragilly, M., Oloketuyi, S., de
Marco, A., Griffie, J., Sage, D., Cohen, E.A.K., et al. (2023). A framework for
evaluating the performance of SMLM cluster analysis algorithms. Nat. Methods
20:259-+. 10.1038/s41592-022-01750-6.

Patterson, G.H., and Lippincott-Schwartz, J. (2002). A photoactivatable GFP for
selective photolabeling of proteins and cells. Science **297**:1873-1877. DOI
10.1126/science.1074952.

Perraki, A., Gronnier, J., Gouguet, P., Boudsocq, M., Deroubaix, A.F., Simon, V.,
German-Retana, S., Legrand, A., Habenstein, B., Zipfel, C., et al. (2018). REM1.3's
phospho-status defines its plasma membrane nanodomain organization and activity
in restricting PVX cell-to-cell movement. PLoS Pathog. **14**:e1007378.
10.1371/journal.ppat.1007378.

Peters, R., Peters, J., and Tews, K.H. (1974). Microfluorimetric Studies on
Translational Diffusion of Proteins in Erythrocyte-Membrane. Pflugers Arch. - Eur. J.
Physiol. **347**:R36-R36.

Qian, H., Sheetz, M.P., and Elson, E.L. (1991). Single-Particle Tracking - Analysis of
Diffusion and Flow in 2-Dimensional Systems. Biophys. J. **60**:910-921. Doi
10.1016/S0006-3495(91)82125-7.

Ries, J. (2020). SMAP: a modular super-resolution microscopy analysis platform for
SMLM data. Nat. Methods **17**:870-872. 10.1038/s41592-020-0938-1.

Sage, D., Pham, T.A., Babcock, H., Lukes, T., Pengo, T., Chao, J., Velmurugan, R.,
Herbert, A., Agrawal, A., Colabrese, S., et al. (2019). Super-resolution fight club:
assessment of 2D and 3D single-molecule localization microscopy software (vol 16,
pg 387, 2019). Nat. Methods **16**:561-561. 10.1038/s41592-019-0453-4.

Saxton, M.J., and Jacobson, K. (1997). Single-particle tracking: Applications to
membrane dynamics. Annu. Rev. Biophys. Biomol. Struct. **26**:373-399. DOI
10.1146/annurev.biophys.26.1.373.

Smokvarska, M., Bayle, V., Maneta-Peyret, L., Fouillen, L., Poitout, A., Dongois,
A., Fiche, J.B., Gronnier, J., Garcia, J., Hofte, H., et al. (2023). The receptor kinase
FERONIA regulates phosphatidylserine localization at the cell surface to modulate
ROP signaling. Sci. Adv. **9**10.1126/sciadv.add4791.

Subach, F.V., Patterson, G.H., Renz, M., Lippincott-Schwartz, J., and Verkhusha,
V.V. (2010). Bright Monomeric Photoactivatable Red Fluorescent Protein for Two-
Color Super-Resolution sptPALM of Live Cells. J. Am. Chem. Soc. **132**:6481-6491.
10.1021/ja100906g.

Tinevez, J.-Y. (2011). simpletracker. https://github.com/tinevez/simpletracker.

Tinevez, J.Y., Perry, N., Schindelin, J., Hoopes, G.M., Reynolds, G.D., Laplantine,
E., Bednarek, S.Y., Shorte, S.L., and Eliceiri, K.W. (2017). TrackMate: An open and
extensible platform for single-particle tracking. Methods **115**:80-90.
10.1016/j.ymeth.2016.09.016.

Tokunaga, M., Imamoto, N., and Sakata-Sogawa, K. (2008). Highly inclined thin
illumination enables clear single-molecule imaging in cells (vol 5, pg 159, 2008). Nat.
Methods **5**:455-455. DOI 10.1038/nmeth0508-455.

Vert, G., and Chory, J. (2006). Downstream nuclear events in brassinosteroid
signalling. Nature **441**:96-100. 10.1038/nature04681.

Wallis, T.P., Jiang, A., Young, K., Hou, H., Kudo, K., McCann, A.J., Durisic, N.,
Joensuu, M., Oelz, D., Nguyen, H., et al. (2023). Super-resolved trajectory-derived
nanoclustering analysis using spatiotemporal indexing. Nat. Commun. **14**:3353.
10.1038/s41467-023-38866-y.

Wilkinson, M.D., Dumontier, M., Aalbersberg, I.J., Appleton, G., Axton, M., Baak,
A., Blomberg, N., Boiten, J.W., Santos, L.B.D., Bourne, P.E., et al. (2016).
Comment: The FAIR Guiding Principles for scientific data management and
stewardship. Sci. Data **3**10.1038/sdata.2016.18.

Witthöft, J., Caesar, K., Elgass, K., Huppenberger, P., Kilian, J., Schleifenbaum,
F., Oecking, C., and Harter, K. (2011). The activation of the Arabidopsis P-ATPase 1
by the brassinosteroid receptor BRI1 is independent of threonine 948 phosphorylation.
Plant Signaling Behav. **6**:1063-1066. 10.4161/psb.6.7.15650.

Wolf, S. (2020). Deviating from the Beaten Track: New Twists in Brassinosteroid
Receptor Function. Int. J. Mol. Sci. **21**10.3390/ijms21051561.

Wolf, S., Mravec, J., Greiner, S., Mouille, G., and Hofte, H. (2012). Plant cell wall
homeostasis is mediated by brassinosteroid feedback signaling. Curr. Biol. **22**:1732-
1737. 10.1016/j.cub.2012.07.036.

Wolf, S., van der Does, D., Ladwig, F., Sticht, C., Kolbeck, A., Schurholz, A.K.,
Augustin, S., Keinath, N., Rausch, T., Greiner, S., et al. (2014). A receptor-like
protein mediates the response to pectin modification by activating brassinosteroid
signaling. Proc. Natl. Acad. Sci. U.S.A. **111**:15261-15266. 10.1073/pnas.1322979111.

Yin, Y., Vafeados, D., Tao, Y., Yoshida, S., Asami, T., and Chory, J. (2005). A new
class of transcription factors mediates brassinosteroid-regulated gene expression in
Arabidopsis. Cell **120**:249-259. 10.1016/j.cell.2004.11.044.

Zhang, M.S., Chang, H., Zhang, Y.D., Yu, J.W., Wu, L.J., Ji, W., Chen, J.J., Liu, B.,
Lu, J.Z., Liu, Y.F., et al. (2012). Rational design of true monomeric and bright
photoactivatable fluorescent proteins. Nat. Methods **9**:727-U297.
10.1038/Nmeth.2021.

Zhang, X., Henriques, R., Lin, S.S., Niu, Q.W., and Chua, N.H. (2006).
Agrobacterium-mediated transformation of Arabidopsis thaliana using the floral dip
method. Nat. Protoc. **1**:641-646. 10.1038/nprot.2006.97.

Zhu, J.Y., Li, Y., Cao, D.M., Yang, H., Oh, E., Bi, Y., Zhu, S., and Wang, Z.Y. (2017).
The F-box Protein KIB1 Mediates Brassinosteroid-Induced Inactivation and
Degradation of GSK3-like Kinases in Arabidopsis. Mol. Cell **66**:648-657 e644.
10.1016/j.molcel.2017.05.012.

849