**Translation as a Biosignature**

**Authors:** Jordan M. McKaig[1], MinGyu Kim[2], Christopher E. Carr[1,2*]

**Affiliations:**

[1]School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA,

30332, USA. [2]Daniel Guggenheim School of Aerospace Engineering, Georgia Institute of

Technology, Atlanta, GA, 30332, USA.

*Correspondence to: cecarr@gatech.edu. Address: ESM Building, Room G10, 620 Cherry St

NW, Atlanta, GA 30332, USA

**One Sentence Summary:**

Life, defined as a chemical system capable of Darwinian evolution, likely requires an apparatus

to translate heritable instructions into cellular machinery, and we propose to detect this as a

biosignature of extant life beyond Earth.

**Word Counts:** Abstract (200), Main Text (6952)

**Abstract**

Life on Earth relies on mechanisms to store heritable information and translate this information

into cellular machinery required for biological activity. In all known life, storage, regulation, and

translation are provided by DNA, RNA, and ribosomes. Life beyond Earth, even if ancestrally or

chemically distinct from life *as we know it* may utilize similar structures: it has been proposed

that charged linear polymers analogous to nucleic acids may be responsible for storage and

regulation of genetic information in non-terran biochemical systems. We further propose that a

ribosome-like structure may also exist in such a system, due to the evolutionary advantages of

separating heritability from cellular machinery. Here, we use a solid-state nanopore to detect

DNA, RNA, and ribosomes, and demonstrate that machine learning can distinguish between

biomolecule samples and accurately classify new data. This work is intended to serve as a proof

of principal that such biosignatures (i.e., informational polymers or translation apparatuses)

could be detected, for example, as part of future missions targeting extant life on Ocean Worlds.

A negative detection does not imply the absence of life; however, detection of ribosome-like

structures could provide a robust and sensitive method to seek extant life in combination with

other methods.

**Key words:** RNA world, Darwinian evolution, nucleic acids, agnostic life detection

**Main Text:**

**1. Introduction**

Life is commonly defined as a "self-sustaining chemical system capable of Darwinian evolution"

(Benner, 2010; Joyce, 1994; Luisi, 1998; Neveu et al., 2018). In all known forms of life,

Darwinian evolution occurs via heritable changes encoded into molecular features (i.e., nucleic

acids), which are then translated into cellular machinery and other proteins. These products of

translation are responsible for a variety of biological functions within and outside the cell

including maintenance, growth, reproduction, regulation, structure, storage, and transport.

In all known life, the ribosome is integral to the "central dogma of biology," which traces

the flow of genetic information into organismal structure and function: genetic information is

stored in DNA, which is transcribed into RNA, which is then translated into protein. Specifically,

the ribosome is responsible for the process of translation, using mRNA-encoded information to

synthesize proteins. All known life is descended from a common ancestor (Pace, 2001), and the

structure and genetic code of the ribosome is highly conserved across evolutionary time and the

tree of life. Indeed, the most highly-conserved sequences across biological systems correspond to

16S and 23S ribosomal subunits and transfer RNAs, and these sequences have experienced little

change over the past 4 billion years (Bray et al., 2018; Isenbarger et al., 2008). Ribosomes are

found within all cells regardless of place in the tree of life or level of activity, and are

instrumental for a cell's ability to carry out life-sustaining processes such as growth,

reproduction, and metabolism (Bernstein and Baserga, 2004; Bowman et al., 2020).

This conservation of ribosome-associated genomes has been posited as evidence for the

existence of an RNA-protein world before DNA became the dominant form of nucleic acid-

based information storage (Fournier et al., 2010; Goldman et al., 2010; Harish and Caetano-

Anollés, 2012; Hsiao et al., 2009; Jerome et al., 2022). This "RNA world" model for the origin

of life posits that RNAs may have acted as both heritable material as well as cellular machinery

prior to the evolution of the ribosome or other more sophisticated biological structures. This

feature persists today, with RNA molecules being capable of both storing genetic information

and catalyzing biochemical reactions. However, there is good reason to assume that life beyond

Earth or an alternate origin of life could utilize a heritable storage system that is physically

distinct from translation-performing cellular machinery, as discussed in section 2.1.

Translation-performing molecules similar to ribosomes could function as an agnostic

biosignature, providing evidence for "life as we don't know it" that may not share a common

origin or physiochemical basis with life on Earth. Given the uncertain range of possibilities for

alien biochemistry, morphology, and other characteristics, various methodologies to search for

non-terran lifeforms have been proposed that are not specific to known biochemistry. Such

strategies can identify potential agnostic biosignatures via direct detection, such as using DNA

probes (apatamers) to bind to repeating patterns within compounds (Johnson et al., 2018). They

can also be elucidated by analyzing observed molecules: for example, the probability of a

molecule of a certain complexity forming abiotically can be quantified (Marshall et al., 2021),

and identification of such molecules can be accomplished via mass spectrometry (Chou et al.,

2021). Such methods generally rely on the assumption that observation of abundant molecules

with sufficiently high complexity can indicate a biotic origin.

Molecules with specific structural or functional features have been proposed as potential

agnostic biosignatures. Such features are generally analogous to what is known about terran life

without presupposing any origin or biochemical basis. For example, information-encoding

charged linear polymers could serve as a mechanism to store, regulate, and transmit genomic

information, analogous to nucleic acids on Earth (Benner, 2017). We extend this point, and argue

that the "Darwinian evolution" definition of life not only implies a mechanism to store heritable

information, but further necessitates a mechanism to translate this information into material that

is useful for the biological system. In terran life, this need is satisfied by the ribosome, a

molecule that is composed of RNA and proteins and is responsible for using genetic information

to synthesize polypeptides and proteins through a process known as translation. The ribosome

has a very specific structure and size (although it is important to note that eukaryotic ribosomes

are 25-50% larger in diameter), and we similarly argue that non-terran life may use a molecule of

similar structure and size. Here, we demonstrate that DNA and RNA can be detected with solid-

state nanopore instrumentation. We also describe the case for targeting the translational

apparatus as a biosignature, and demonstrate the ability to target our only known example, the

ribosome, using solid-state nanopores. Finally, we show that machine learning can discriminate

between different classes of biomolecules (DNA, RNA, and ribosomes) based on the features

associated with their solid-state nanopore detections.

**2. The Case for the Translational Apparatus as a Biosignature**

*2.1 What would life look like without translation?*

Any life with separate heritable material and cellular machinery must have a translation

system to convert instructions into machinery. If these functions are not separate, then heritable

material must also act as machinery, including both self-copying and carrying out all other life

functions (metabolism, maintenance, growth, evolution). While ribozymes have been discovered

and engineered that are capable of copying short RNA sequences (Been and Cech, 1988; Ekland

and Bartel, 1996; Horning and Joyce, 2016; Sczepanski and Joyce, 2014), and RNA sequences

can be inherited via viral replication (Rampersad and Tennant, 2018), we are not aware of any

truly self-replicating informational polymer system of non-trivial complexity (i.e., capable of

replicating nucleic acid strands longer than tens of bases). Furthermore, such systems would be

unlikely to competitively support the chemical richness required to subsume all the roles of

cellular machinery (e.g., an RNA-only RNA world in the absence of proteins). In other words, if

such a system evolved, it would likely be displaced by a system with separate coding and

machinery due to selective pressures, just as may have happened with life as we know it (Gilbert,

1986; Orgel, 1968).

Separation of coding and machinery would decouple conflicting functions. To perform a

wide range of complex functions, cellular machinery requires chemical diversity, whereas

enabling copying requires chemical simplicity to facilitate copying robustness. Even while RNA

can serve as both as an informational storage molecule and as cellular machinery (as evidenced

by ribosomes, tRNAs, and other ribozymes), a physically separate translation apparatus might be

expected to arise because decoupling these two functions offers significant benefits, as observed

by these functions being decoupled in all known life.

Furthermore, separation among classes of functional biomolecules allows for their

maintenance and enables structural differences that align with function. For example, linear

charged polymers are considered as a potential universal heritable information storage format for

aqueous life (Benner, 2017). In terran life, such polymers exist as nucleic acids, which store

genetic information in linear base sequences. Their topology contributes to copying fidelity:

branched or otherwise more complicated structures would pose difficulties for effective copying.

Additionally, the charge of nucleic acids separates the properties of the molecule from its

information content, so that the molecule can be effectively copied regardless of information

content. Conversely, the ribosome has a complicated and specific structure that aids in its

preservation and enables its highly specialized role in translation. The assembly of two types of

biopolymers (polynucleotides and polypeptides) yields a functional and stable ribosome capable

of conducting its important catalytic role, leveraging the qualities of both nucleic acid and

protein to enable translation (Runnels et al., 2018). The differing charges across the ribosome

similarly allows it to function by enabling compatibility with the biomolecules that interact with

various parts of the ribosome (Leininger et al., 2021). Furthermore, self-complementarity (the

propensity for a biomolecule to bind with itself) among biopolymers enables self-preservation

and functional separation of different types of biomolecules. Meanwhile, heterogenous

assemblages of different types of biopolymers can beget more complex functionality and

"partner-preservation", where mutualistic assemblies of different biomolecule types can confer

protective effects on each other (Runnels et al., 2018). All known life requires a translation

apparatus. However, let us consider life that does not have this layer of abstraction, such that

hereditary molecules also serve as cellular machinery. As alluded to above, this introduces

complexity into copying: changes in information content would result in changes in machinery,

which could overly constrain such life's exploration of evolutionary space. In short, life without

translation would be limited in complexity, capabilities, and evolvability (Cuevas-Zuviría et al.,

2023), the latter of which is central to our definition of life.

*2.2 Detection Approaches*

A classic approach for detection and identification of bacteria and archaea involves using

polymerase chain reaction (PCR) to amplify and sequence the gene encoding for 16S ribosomal

RNA (rRNA), which is highly conserved across prokaryotes (Isenbarger et al., 2008). This

genomic approach is integral for the study of microbial communities on Earth (Kim and Chun,

2014), since it targets a genomic sequence that exists in nearly all bacteria and archaea, or for

some choices of primers, certain large clades of organisms. Similarly, the eukaryotic 18S rRNA

gene can be used to identify study eukaryotic community members. Ribosome profiling

techniques can be used to sequence mRNA fragments that are being actively translated, enabling

study of translation with single-cell resolution (Heiman et al., 2008; VanInsberghe et al., 2021).

However, targeting specific ribosome-associated RNA sequences is only suitable for seeking

lifeforms that are ancestrally related to known life. One instance where this may be relevant

would be for searching for Martian microbial communities that share an origin with life on Earth,

e.g., transferred between the planets via lithopanspermia (Carr, 2022; Mileikowsky et al., 2000).

Analysis of rRNA can be used to identify active taxa and assess activity in environmental

microbial communities, although caveats exist for these use cases. The relationship between

rRNA concentrations and microbial activity is complex and therefore simple correlations do not

often exist between the two, and any existing relationships are not consistent across taxa

(Blazewicz et al., 2013). Furthermore, in certain environmental conditions (e.g., low

temperature, low water activity, high osmolarity), the rRNA itself may persist after

environmental conditions have changed, influencing conclusions about which specific microbes

are actually currently active in an environment (Schostag et al., 2020). Although drawing explicit

conclusions about which microbial taxa are currently active is difficult from rRNA analysis

alone, detection of ribosomal material can serve as an indicator that microbial activity has

occurred recently or is actively underway in a habitat.

Content-agnostic approaches such as can be used to detect ribosomes without making

assumptions about their genetic associations. Microscopy techniques, including fluorescence

microscopy and cryoelectron microscopy, can provide information about ribosomal structure and

morphology, location and organization within cells, and study of translation (Bai et al., 2013;

Bakshi et al., 2012; Fu et al., 2011; Murphy et al., 2020). However, such technologies are

destructive, require extensive sample preparation, or are susceptible to image artifacts.

Additionally, the size, sensitivity, and complexity of these instruments present difficulties for

adapting these types of microscopes for *in situ* space exploration applications.

Dynamic light scattering (DLS) can be used to characterize molecules in an aqueous

sample by measuring their diffusion coefficient, which in turn can be used to infer molecule size

and shape (Stetefeld et al., 2016). For example, DLS can be used to interrogate ribosomal shape

(Bruining and Fijnaut, 1979), track structural changes (Patel and Cunningham, 2002), and make

comparisons between ribosomal types (Müller et al., 1986). DLS is a good strategy for making

accurate size and shape particle measurements in minutes with a small amount of sample ($<3\mu L$).

Additionally, sample preparation is minimal, and this technique is compatible with high-

concentration and turbid samples. However, measurements are sensitive to temperature and

solvent viscosity, and DLS is not capable of achieving sufficiently high resolution to distinguish

between closely-related molecules, such as monomers vs. dimers. Finally, DLS is sensitive to

changes in size and concentration (Jose et al., 2019). Overall, DLS is a good technique for

interrogating pure ribosomal samples at high concentration, but is not viable for identifying or

characterizing ribosomes within a heterogenous sample or when present at low abundance.

Solid-state nanopores (SSN) can provide a useful method for ribosomal detection with

single-molecule or single-particle resolution. This instrumentation detects individual molecules

or particles (hereafter referred to in the generic sense as particles) in an aqueous sample of

known conductivity by applying an ionic current to a nanometer-scale pore in a synthetic

membrane, using a voltage or pressure differential to pass particles through the pore, then

measuring how the particles disrupt the ionic current. The magnitude of the change in

conductance and duration of the blockage can then be used to determine the relative size of each

particle (**Figure 1**). SSN instrumentation is distinct from biological nanopore sequencing (e.g.,

the MinION from Oxford Nanopore Technologies), which uses controlled translocation of

nucleic acids regulated by biological components with specific base sequences inferred from

changes in ionic current flow. Instead, due to its larger, non-specific pore, SSN tools can analyze

a wide size range of particles and are not constrained to a specific class of molecules. This

benefit is also a challenge to achieving molecular specificity and to achieving single base

resolution, as discussed below.

In recent years, major advancements have been made in SSN technologies to improve

their sensitivity, selectivity, and throughput capacity, enabling development of applications in

biology, chemistry, and medicine (Xue et al., 2020). Particularly of interest to this work and

other astrobiological applications, SSN-based methodologies have been used to detect

biomolecules such as nucleic acids and ribosomes in laboratory conditions. Raveendran et al.

(2020) used nanopipettes fitted with ~60 nm nanopores on the tips to detect eukaryotic 80S

ribosomes (derived from human and *Drosophila melanogaster* cell cultures) to detect ribosomes

and distinguish ribosomes from polysomes (groups of ribosomes bound together by mRNA

strands). Rudenko et al. (2011) used a nanopore-microfluidic chip to detect individual

prokaryotic 50S *Escherichia coli* ribosomes and control their rate of nanopore translocation.

Similarly, Rahman et al. (2019) used an optofluidic chip to control the flow of ribosomes, DNA,

and proteins through a nanopore and measure these biomolecules as they translocated the

nanopore. Xia et al. (2022) used a nanopore drilled into a silicon nitride membrane on a glass

chip to detect and characterize DNA that was extracted from Martian analog soils, directly

demonstrating SSN applications for astrobiological life detection.

Machine learning (ML) can be applied to solid-state pore measurements to classify and

identify biological particles, given appropriate training data. Xia et al. (2021) applied multiple

ML models to their SSN data sets to differentiate signals of monosaccharides and disaccharides,

which enabled them to determine the composition of polysaccharide compounds. Arima et al.

(2020) achieved over 99% accurate classification of viruses using ML. Using a molybdenum

disulfide membrane, instead of the commonly-used silicon nitride membrane, Farimani et al.

used ML to identify individual amino acids within polypeptide chains (Farimani et al., 2018). On

the micropore scale, ML can be similarly used to identify and classify cells. Using solid-state

micropore data, Hattori et al. applied ML to discern between five clinically-relevant bacterial

species based on physical features including cell shape, surface charge density, mass, surface

proteins, and motility (Hattori et al., 2020). Tsutsui et al. tagged bacterial walls with synthetic

peptides and analyzed them with a solid-state micropore, then used machine learning to identify

bacterial features and discriminate between strains (Tsutsui et al., 2018).

Solid-state nanopores are non-specific and can be used to detect not just biomolecules

such as nucleic acids and ribosomes, but also non-biological particles. Due to this non-

specificity, it has been proposed that solid-state nanopores may be useful for detecting agnostic

biosignatures (Bywaters et al., 2017). It has also been proposed that alien lifeforms with a

separate origin and alternative biochemistry from terran life may store genetic information in

charged linear polymers that are structurally and functionally analogous, but not necessarily

chemically similar, to nucleic acids (Benner, 2017). We argue here that translational machinery

analogous to ribosomes may be present and could additionally serve as an agnostic biosignature.

Solid-state nanopore instrumentation can detect and characterize both nucleic acids and

ribosomes, and hold potential for detecting similar structures produced through alternate

biochemistries.

Here we explore the utility of ribosomes (and agnostic translational machinery) as a

biosignature, and demonstrate nucleic acid and ribosomal detection using solid-state nanopore

instrumentation. Additionally, we propose that detecting one or more peaks indicative of

translational machinery, in a sample that has been agitated to sufficiently lyse any existing cells,

could indicate the presence of life.

*2.3 Expected Size of Translational Apparatus for Life as We Don't Know It*

Life beyond Earth, even if ancestrally or chemically distinct from life as we know it may

utilize a structure similar in size to the ribosome, balancing the energetic costs of assembling

translation machinery with the minimum complexity required for such a machine. In short, the

translation machinery must have some minimum size related to the complexity of its translation

function. In addition, there is likely a selective advantage to the translation machinery not being

far larger than the minimum size, due to the energetic and time cost of synthesis and assembly

(Li et al., 2014; Shore and Albert, 2022). Additionally, different life forms within the

descendants of a particular origin of life may have translation apparatus that differ in size, similar

to how prokaryotic and eukaryotic ribosomes have different sizes on Earth. The ancient

ribosomal core, which was capable of translation and has been evolutionarily conserved since

before the last universal common ancestor (LUCA), is about 2 MDa in size. Contemporary

prokaryotic ribosomes are about 2.5 MDa, with a slightly larger size due to additional proteins

and a slightly longer rRNA sequence (about 100 nucleotides larger than the core) (Bowman et

al., 2020). Smaller bacterial ribosomes also exist, with certain ribosomal proteins being absent in

bacteria with exceptionally small genomes (Nikolaeva et al., 2021). Eukaryotic ribosomes range

in size from about 3.5 to 4 MDa, similarly attributable to longer rRNA sequences (hundreds to

thousands of nucleotides larger than the core) and additional proteins (Bowman et al., 2020). It

has been proposed that the size difference between prokaryotes and eukaryotes reflects additional

complexity in eukaryotic regulation of translation (Lafontaine and Tollervey, 2001).

Comparisons of eukaryotic and prokaryotic ribosomes can allow for identification of the

common ancestral core, and molecular models can be used to infer how ribosomes may have

evolved (Petrov et al., 2014). Modeling suggests that various ribosomal capabilities for

interpreting genetic encoding and creating polypeptides in response were sequentially added,

resulting in a molecule composed of functional proteins and nucleic acids (Petrov et al., 2015).

The tight size distribution and number of ribosomal proteins (~50 in prokaryotes, ~80 in

eukaryotes) may reflect selection for optimal duplication efficiency (Shore and Albert, 2022).

We do not presume any chemical specificity for ribosome-like machinery; rather, we argue that

function and approximate size may be conserved across multiple origins of life with molecules

that may differ in structure or composition.

**3. Methods**

*3.1 Reagents and Reagent Preparation*

Aliquots of thawed biomolecule samples (single-stranded RNA [ssRNA] ladder [NEB

#N0362], 1 kilobase [kb] double-stranded RNA [dsRNA] ladder [NEB #N0363], 1 kilobase

double-stranded DNA [dsDNA] ladder [Thermo Scientific #SM0311], high range dsDNA ladder

[Thermo Scientific #  SM1351], DNA plasmid pUC19 [NEB #N3041], *E. coli* ribosome [NEB

P0763S]) were added to Ontera Start-Up Buffer (2M LiCl, conductivity 11.96 S/m).

*3.2 Solid-State Nanopore Operation*

The Ontera NanoCounter was operated using the associated software (NanoCounter

v0.19.1). Nanopores were conditioned following manufacturer instructions, using the Ontera

Start-Up Buffer (2M LiCl, conductivity 11.96 S/m) as the conditioning buffer. Biomolecule

samples and negative controls were loaded into nanopore and analyzed following manufacturer

instructions.

*3.3 Detection of dsDNA, ssRNA, dsRNA, Ribosomes*

For a given experiment, the same pore was loaded with a sequence of samples after

conditioning, using an alternating pattern of negative control (0.2µm-filtered buffers without

added biomolecules) (run time 3-5 minutes), followed by a sample (run time 6-10 minutes),

followed by the next negative control. This was done to assess buffer cleanliness, flush out and

assess any carryover from prior samples, and quantify the risk of false positives. The

NanoCounter software used an automatic event extraction program to identify events with a

signal-to-noise ratio high enough to consider the event to be a particle translocating the

nanopore.

*3.4 Application of Machine Learning to Differentiate Biomolecule Types*

To prepare the SSN data for machine learning classifier training, event metadata was

compiled for all biomolecule events with a signal-to-noise ratio (SNR) greater than 5. The

MATLAB Classification Learner app (with MATLAB_R2023a) was used to train several

machine learning algorithms on the data. Data was partitioned by the toolbox, with 25% used for

holdout validation, 10% used for testing, and the remining 65% used for training. The response

variable was the sample identifier ("sample_name"), and the predictor variables were dwell time

("dwell_sec"), signal-to-noise ratio ("SNR"), mean current amplitude during event

("mean_amp_pA"), maximum current amplitude during event ("max_amp_pA"), median current

amplitude during event ("med_amp_pA"), standard deviation of amplitude ("std_amp_pA"), and

area of the event disruption ("area_pA_sec"). The Ontera Nanocounter software recorded current

values in picoamps and associated conductivity values in nanosiemens; since the applied voltage

was constant across experiments and thus conductivity always scaled with current, only current

values were used for the machine learning training to avoid redundancy. The datasets and code

used to run this analysis can be downloaded at https://github.gatech.edu/PXL/TAB**.**

**4. Results**

*4.1 Solid-State Nanopore Detection Achieved a Low False Positive Rate*

Negative controls were run before every test sample to assess background and carryover

from previous runs. When testing sterilized and filtered brines on new nanopores, event

abundance was low (<10 events per minute). Events detected in this case are likely particle

contaminants introduced during buffer preparation, sample loading, or present within the flow

channels of the nanopore. Event abundance in the negative controls when ran between DNA

samples was similarly low, as shown in **Figure 2.** A higher abundance of events than usual was

sometimes detected in pores when running negative controls between biomolecule samples

(**Table S1**). This was especially prominent in cases where clogging (for example, high-

concentration ribosome samples) and biomolecule precipitation (for example, RNA in lithium

chloride) were suspected to occur.

*4.2 Different Biomolecule Classes and Sizes Have Distinct Signatures*

DNA and RNA samples were suspended in 2M LiCl and detected using the Ontera

NanoCounter. The DNA samples were plasmid pUC19 (a circular fragment with a length of

2,676 base pairs), 1 kb dsDNA ladder (linear DNA fragments of 14 different sizes ranging from

250 to 10,000 base pairs), and high range dsDNA ladder (linear DNA fragments of 8 different

sizes ranging from 10,171 to 48,502 base pairs). The RNA samples were single-stranded RNA

(ssRNA) ladder (linear ssRNA fragments of 7 different sizes ranging from 500 to 9,000 bases)

and double-stranded RNA (dsRNA) ladder (linear dsRNA fragments of 7 different sizes ranging

from 21 to 500 base pairs). Events were automatically extracted from these runs using the

NanoCounter software, totaling 2,284 events over 6.62 minutes for plasmid pUC19, 919 events

over 5.93 minutes for 1 kb dsDNA ladder, 3,115 events over 5.99 minutes for high range dsDNA

ladder, 1,874 events over 10.03 minutes for ssRNA, and 1,398 events over 11.01 minutes for

dsRNA (**Table S1**). Key features were also identified, most importantly dwell time (the amount

of time that the molecule spent translocating the nanopore) and change in conductance (resulting

from the current disruption of the molecule translocating the nanopore). Given the low

background noise when running biomolecule-free buffer solutions (**Figure 2, Table S1**), we

assume that all events from these runs are indeed the target biomolecule translocating the

nanopore.

DNA events displayed specific features that were indicative of biomolecule size and

structure **(Figure 3)**. The change in conductance for pUC19 was generally approximately double

that of the HR and 1kb ladders. This is thought to be attributed to 4 strands (2 dsDNA strands)

passing through the nanopore at once for a circular DNA plasmid, as compared to the 2 strands

(1 dsDNA strand) that would pass through at once with a linear ladder. The lower-range ladder

(1kb) had a shorter range of dwell times compared to the upper-range ladder (HR), which is

indicative of the respective ladder fragment lengths.

The ssRNA and dsRNA events also differed in their change in conductance and dwell

time, with ssRNA samples generally having a slightly larger change in conductance and slightly

smaller dwell time than the dsRNA samples (**Figure 4**). However, these differences are less

discernable for the RNA samples than the DNA samples. This may be attributable to the

differences in molecule size and structure between the two types of DNA and RNA tested: the

two DNA samples differed in conformation (circular plasmid of a specific length vs. linear

fragments of varying lengths), while both RNA samples consisted of linear fragments spanning

similar ranges of sizes but in double or single stranded form. Strand number or topology can be

expected to influence persistence length and thus secondary structure (e.g., ssRNAs are highly

flexible). Additionally, high numbers of events in the buffers immediately after running RNA

suggests that precipitation and resolubilization may have occurred for these samples, which

could similarly result in low-quality events due to interactions with the nanopore.

Similarly, *E. coli* ribosomes were suspended in 2M LiCl and measured with the Ontera

NanoCounter. 17,077 events were extracted from 10.03 minutes of run time (**Table S1**). **Figure**

**5** contains a heat map of events from this run (Panel A), individual examples of events

interpreted to be an intact ribosome and a ribosomal fragment (Panel B), and overlaid events

from the regions of the heat map interpreted to be intact ribosomes (Panel C) and ribosomal

fragments (Panel D). Of the 17,077 events, 6,885 are thought to represent intact ribosomes

translocating the nanopore (defined as events with a change in conductance greater than 8 nS).

The large number of events with changes in conductance smaller than 5 nS are thought to be

ribosomal fragments. Within this results space, there is a high concentration of events with a

change in conductance between 1-5 nS and a dwell time between 0.00001-0.0001 seconds. These

are interpreted as protein fragments. Some events have a larger dwell time (0.0001-0.1 seconds),

which closely aligns with the reign occupied by RNA events (**Figure 4**). Thus, it is thought that

these fragments may be RNA products of ribosome fragmentation.

Overall, based on dwell time and change in conductance, signals from each biomolecule

class cluster together, although there is some overlap (**Figure 6**). Machine learning was

implemented in attempts to better differentiate between sample types.

*4.3 Machine Learning Algorithms Can Differentiate Biomolecule Types and Structures*

The Classification Learner app identified the validation and testing accuracy of a variety

of machine learning classifier methods, including neural networks, support vector machines, k-

nearest neighbors, kernel approximation, naïve Bayes, discriminant, logistic regression, and tree-

based models (**Table S2**). To provide an illustrative example, detailed results from the medium

tree model are shown. This specific model was selected for its high interpretability, as other top-

scoring algorithms (e.g. neural networks, support vector machines, k-nearest neighbors) can be

more difficult to interpret. The medium tree was specifically selected from the applied decision

tree models (fine tree, medium tree, coarse tree) as it balances ease of interpretation with

accuracy – the fine tree model had higher accuracy levels (94.41% validation accuracy and

95.02% testing accuracy, compared to the medium tree's 92.89% validation accuracy and

93.34% testing accuracy), but the fine tree involved a much more complex array of decisions.

Additionally, with limited training data, it is difficult to preclude overfitting, which is more

likely to occur for finer trees. The medium tree (**Figure S1**) contained 19 splits to classify

biomolecules based on event features (**Table S3**). It is possible to reach the same biomolecule

classification by following different paths throughout the tree, which may be attributable to the

presence of different molecule morphologies within a single sample (e.g. intact vs. fragmented

ribosome, supercoiled vs. unsupercoiled plasmid, bent vs. straight DNA/RNA fragment). The

medium tree's confusion matrix (**Figure 7**) demonstrates validation performance, showing how

biomolecules were generally classified correctly by this model. This specific model is shown

here due to its interpretability, but other models achieved higher performance and warrant further

exploration.

**5. Discussion**

Here, we argue that ribosomes can serve as a biosignature indicative of terran-like life,

and another instance of cellular translational machinery of similar size and structure could serve

as an agnostic biosignature. In a case of "life as we don't know it," a higher abundance of

particles within a certain size range compared to the rest of the sample could serve as an agnostic

biosignature. Using solid-state nanopore instrumentation, we demonstrate that such cellular

machinery can be detected and distinguished from other biomolecules in aqueous samples.

Such measurement is sensitive (capable of correctly identifying target molecules in the

sample) as evidenced by successful detection of various biomolecules, and specific (able to

discern whether a sample does not contain target molecules) as evidenced by the low abundance

of events within the negative controls. When measuring negative controls, no special precautions

were taken (e.g. filtration of buffers), suggesting the potential for lowering the false-positive

event rate further. Target biomolecules were evident within heat map distributions as specific

peaks corresponding to different molecule sizes and conformations. This is especially true for

ribosomes, which exist in very specific quantized sizes for prokaryotes and eukaryotes and are

often highly abundant, e.g., 25,000 to 31,000 per $\mu m^3$ for active *E. coli* cells, as reported by

Bakshi et al. 2012. High abundance of ribosomes is not limited to active cells. Slow-growing

ultra-small archaea of volume 0.03 $\mu m^6$ (effective spherical diameter of 0.4 µm) have an average

of 92 ribosomes/cell (Comolli et al., 2009), or 3000 per $\mu m^3$. Bacterial endospores contain

ribosomes at a similar abundance to active cells, even when mRNA has been degraded

(Chambon et al., 1968). This high abundance of ribosomes across different types and activity

levels of cells may allow for the distinguishing of ribosomes within cellular lysate or

environmental samples treated to lyse cells. Conversely, abiotic material that was similarly

treated would be expected to have a more even particle size distribution.

Certain features could confound a search for translational machinery as a biosignature,

including other cellular components and abiotic features. The severity and implications of such

mistaken identities vary. If another cellular component (e.g., genetic information storage

molecules, functional proteins, other cellular organelles) is mistaken as a translational apparatus,

then life has still been detected. However, if a particle of abiotic origin is mistaken for a

ribosome-like molecule, this could cause a false-positive life detection result. For example, in

Earth's atmosphere, mechanically generated particles tend to be in the coarse-mode (>1 µm) size

range, with particle sizes in the nanometer to tens of nanometer range forming from nucleation

and thus likely to exhibit a continuous size range rather than high abundance at a specific size.

Accumulation of nucleated particles can result in larger sizes (typically hundreds of nanometers).

However, in astrobiological contexts, additional caution is warranted. While 99% of dust in the

Enceladus plume is expected to be below 10 nm with a peak at 2 nm (Meier et al., 2014), the

presence of hydrothermal activities at Enceladus was subsequently inferred from Cassini-

detected silica nanoparticles in the 4-16 nm size range (Hsu et al., 2015). One mitigation in this

case would be to filter a sample to retain cells and reject free silica nanoparticles prior to lysis.

An additional approach would be to heat-treat a sample and observe if the particles are heat

tolerant or exhibit denaturation or degradation. Further and varied analysis of any detected

molecules would be necessary to discern their composition, function, and potential biogenicity.

While outside the focus of this paper, future work should evaluate the potential abiotic

production of particles in different settings to assess the risk of false positives.

The results presented here are intended to serve as a proof-of-concept study that lays the

foundation for more extensive and robust work, with multiple avenues identified for further

experimentation. One factor that warrants further study is buffer-biomolecule compatibility. In

this study, it was particularly difficult to achieve low event abundance in the negative control

buffer samples ran between RNA and ribosome samples. This is likely due to effects of the

lithium chloride buffer on these biomolecules, which is thought to have caused molecular

precipitation that hindered flow cell flushing and subsequent molecule detection. Similarly, it

was difficult to discern ssRNA and dsRNA events by looking at current disruptions, possibly to

due to precipitation of the RNA fragments and/or interaction with the nanopore. It seems that

most of the detected events were attributable to true translocations of RNA strands through the

nanopore, since the event rate was much higher for the RNA samples than the blank runs

immediately before and after (**Table S1**). However, many of these events had a relatively low

signal-to-noise ratio compared to what was observed for other samples. Refining buffer and pH

compatibility with RNA (and other target biomolecules) is a future target for optimization.

Pore size constrained the size of biomolecules that can be analyzed with this system, and

may have played a role in the ribosomal fragmentation observed. This study used nanopores

embedded within pre-fabricated flow cells that arrived in a desiccated state, which we then

hydrated and subjected to an electrochemical conditioning process that widened and stabilized

the pore. This conditioning process would result in a pore with a diameter between 20-45

nanometers, which could not be significantly altered. This was sufficient to allow the passage of

single strands of nucleic acids (~2 nm diameter) and prokaryotic ribosomes (21 nm diameter). It

is important to match pore and biomolecule size as closely as possible: if the nanopore is too

small, it inhibits free flow of molecules through the nanopore. Conversely, if the nanopore is too

large, signals from translocating molecules may be obscured by noise – the closer the size of the

biomolecule is to the size of the nanopore (without being large enough to clog the nanopore), the

higher the signal-to-noise ratio (Xia et al., 2022). With the nanopores used in this study, we were

able to detect nucleic acids, but may be able to achieve stronger signals with a smaller nanopore.

Additionally, we were able to detect ribosomes, but also observed the presence of smaller

particles thought to be ribosomal fragments. Such fragmentation is hypothesized to have been

caused by buffer chemistry and exacerbated by RNA precipitation and clogging of the nanopore.

Using a larger nanopore in future studies may help alleviate this issue, especially if coupled with

a buffer that is more compatible with ribosomes. Overall, having access to a wide variety of

nanopore sizes will be useful for future work with molecules of various sizes, structures, and

biochemistry. Future analysis of biomolecules with SSN should also include a variety of saline

solutions. Nucleic acids and ribosomes are both structurally sensitive to ionic concentration and

composition (Shukla and Mikkola, 2020; Spitnik-Elson and Atsmon, 1969; Tian et al., 2023),

and biomolecule translocation itself can also be drastically influenced by ionic conditions (Bell

et al., 2016; Kowalczyk et al., 2011). More extensive study of SSN performance under various

saline conditions will be necessary to achieve better understanding of how solution composition

influences results.

Furthermore, we demonstrate that different biomolecule classes and structures can be

differentiated and predicted by decision tree machine learning algorithms with over 90%

accuracy on both validation and test data. Both a fine and a medium tree were generated, with

the medium tree giving a more comprehensible assessment of how biomolecules could be

discerned at the cost of less decisions and lower accuracy, and the fine tree yielding higher

accuracy classification by making more decisions, at the cost of a very lengthy list of nodes and

the potential for overfitting. Other less-interpretable machine learning models (for example,

neural networks) yielded even higher accuracy, further indicating the potential that machine

learning classification holds for SSN data. This is useful for summarizing and transmitting data

during future missions with highly constrained data budgets, such as at or within Ocean Worlds.

These machine learning results are intended to justify more rigorous future application of

machine learning-based classification of solid-state nanopore data, including exploring the utility

of various algorithms, hyperparameter tuning, and feature engineering. In addition to this

computational work, it will be critical to test these algorithms on datasets from different SSN

runs under various conditions and instrumentation to ensure that accurate results are consistently

obtained. Since the dataset for each biomolecule class was generated by a single experimental

run of that specific class, the experimental conditions (e.g., pore size, buffer chemistry, etc.) may

influence event data independently of the influence of the specific biomolecule type. Generation

of more data by analyzing different biomolecule types, structures, and sizes under various buffer

conditions and with different nanopores will be instrumental in refining such a machine learning-

based classification.

There are a few major caveats for applying this laboratory-based detection of terran

biomolecules towards astrobiological life detection. First, the utility of ribosomes or ribosome-

like translational machinery as a biosignature relies on the assumption that such "life as we don't

know it" synthesizes biomolecules for cellular functioning via translation or an analogous

mechanism. Although it is not known how life may exist without ribosomes or a ribosome-like

particles, we acknowledge the possibility that alternate biochemistries, cellular structures, and

life strategies could be associated with life forms that would not be detectable by searching for

translational machinery. Additionally, this work assumes that such a translational apparatus will

remain stable during cellular lysis, extraction, and analysis conditions. If translational machinery

from an alien lifeform were to be destroyed or otherwise disrupted by such processes, this would

increase the chance of false negative results. Finally, unknown abiotic conditions could produce

confounding structures with similar signatures, resulting in false positive readings. This

possibility highlights the need for stringently using negative controls.

We argue that ribosomal detection could serve as one line of evidence for life, and such

measurement could be conducted in conjunction with other life detection analysis. Such

detection could be particularly useful as a pre-screening tool for samples, subjecting it to this

biologically-agnostic particle measurement technique to determine size distribution and potential

biogenicity of samples. This could aid in sample selection or prioritization before passing it

forward to more targeted analyses (e.g., DNA sequencing, microscopy, mass spectroscopy, or

other compositional analyses). Furthermore, we demonstrate how machine learning can be

utilized to classify biomolecules based on various features of translocation events measured by a

solid-state nanopore device. Overall, SSN presents one strategy to interrogate various

components of a sample and search for patterns or features that may indicate biogenicity.

Negative results from an attempt to detect ribosomes would not necessarily preclude the

possibility for life forms to be present within the sample. Opportunities for *in situ* sample

preparation may be limited by lack of knowledge of the biochemistry of any putative

biomolecules, so it is possible that chemical incompatibilities could disrupt the structure of any

molecules of interest. Additionally, "life as we don't know it" may sustain cellular processes

through another mechanism or structure. However, detection of ribosomes or ribosome-like

structures with a solid-state nanopore instrument could support a hypothesis that samples are

from a biologically active ecosystem, especially if both ribosomes and abundant molecules with

RNA-like properties are identified.

Perhaps the most compelling astrobiological application of solid-state nanopore

instrumentation is the search for signs of life in potentially habitable saline environments on

Mars, Europa, or Enceladus. Salinity exerts a major effect on the ability of an organism or a

biomolecule to persist in a system: microbial activity can be detected throughout wide ranges of

salinity levels and salt compositions, and biological structures such as intact cells and

biomolecules can be preserved even beyond the saline limits for active life (Klempay et al.,

2021; Oren, 2008). Thus, many have postulated that saline environments should be a major

target for astrobiological exploration, given their biological relevance and presence throughout

the solar system (Davila and Schulze-Makuch, 2016; Neveu et al., 2020; Pappalardo et al., 2013;

Phillips et al., 2023). Although saline environments are biologically relevant, they can also pose

challenges for conducting science in such environments. Indeed, many analytical techniques are

impeded by salt and require desalination treatments (Lawrence et al., 2023). Given SSN's

intrinsic requirement that samples be suspended in a conductive solution, it may be possible to

use this technique to analyze environmental biomolecule samples within their original saline

solutions or diluted with a high salt buffer of choice, without complex processing such as salt

removal. Furthermore, Ocean Worlds are expected to be low in available free energy, and thus

any existing biosphere would likely be low in biomass (Jones et al., 2018). SSN's low limit of

detection and high sensitivity makes it a promising technology for achieving the single-molecule

resolution that would be necessary to effectively search for intact biomolecules in such

environments.

Solid-state nanopores are of interest for future astrobiological space exploration due to

their ability to detect biomolecules with high sensitivity and low false positive rates, as well as

their potential utility in searching for agnostic biosignatures. Beyond its applications for

astrobiological life detection, SSN has other qualities that overall make it well-suited for space

payloads in general, including high portability, minimal resource requirements, low mass and

power requirements, potential for analysis on minimally-prepared samples, and capability for

rapid analysis. Interplanetary space missions with the explicit goal of searching for

extraterrestrial life are expected in the coming decades (National Academies of Sciences,

Engineering, and Medicine, 2022). With further research and development, a solid-state

nanopore instrument may be a compelling addition to a future mission seeking to detect signs of

extant life in the solar system.

**Acknowledgments:** The authors thank Chen Chen for valuable contributions and comments on

the machine learning components of this paper. **Author contributions:** J.M.M. and C.E.C.

developed the concept; J.M.M. and C.E.C. performed the experiments; J.M.M. and C.E.C.

analyzed the NanoCounter data; J.M.M., M.K., and C.E.C. performed the machine learning

analysis. J.M.M. and C.E.C. wrote the paper. All authors edited and approved the paper.

Biomolecule graphics (appearing in Figures 1, 3, 4) were created with BioRender.com. **Author**

**Disclosure Statement:** We declare no competing interests. **Funding:** J.M.M. was supported by

NASA award 80NSSC21K0234 to C.E.C and NASA FINESST award 80NSSC22K1320 to

J.M.M. and C.E.C. This work was also supported by NASA award 80NSSC19K1028 to C.E.C.

**References:**

Arima, A., Tsutsui, M., Yoshida, T., Tatematsu, K., Yamazaki, T., Yokota, K., Kuroda, S., Washio, T., Baba, Y., Kawai, T., 2020. Digital Pathology Platform for Respiratory Tract Infection Diagnosis via Multiplex Single-Particle Detections. ACS Sens. 5, 3398–3403. https://doi.org/10.1021/acssensors.0c01564

Bai, X., Fernandez, I.S., McMullan, G., Scheres, S.H., 2013. Ribosome structures to near-atomic resolution from thirty thousand cryo-EM particles. eLife 2, e00461. https://doi.org/10.7554/eLife.00461

Bakshi, S., Siryaporn, A., Goulian, M., Weisshaar, J.C., 2012. Superresolution Imaging of Ribosomes and RNA Polymerase in Live Escherichia coli Cells. Molecular Microbiology 85, 21. https://doi.org/10.1111/j.1365-2958.2012.08081.x

Been, M.D., Cech, T.R., 1988. RNA as an RNA Polymerase: Net Elongation of an RNA Primer Catalyzed by the *Tetrahymena* Ribozyme. Science 239, 1412–1416. https://doi.org/10.1126/science.2450400

Bell, N.A.W., Muthukumar, M., Keyser, U.F., 2016. Translocation frequency of double-stranded DNA through a solid-state nanopore. Phys. Rev. E 93, 022401. https://doi.org/10.1103/PhysRevE.93.022401

Benner, S.A., 2017. Detecting Darwinism from Molecules in the Enceladus Plumes, Jupiter's Moons, and Other Planetary Water Lagoons. Astrobiology 17, 840–851. https://doi.org/10.1089/ast.2016.1611

Benner, S.A., 2010. Defining Life. Astrobiology 10, 1021–1030. https://doi.org/10.1089/ast.2010.0524

Bernstein, K.A., Baserga, S.J., 2004. The Small Subunit Processome Is Required for Cell Cycle Progression at G1. Mol Biol Cell 15, 5038–5046. https://doi.org/10.1091/mbc.E04-06-0515

Blazewicz, S.J., Barnard, R.L., Daly, R.A., Firestone, M.K., 2013. Evaluating rRNA as an indicator of microbial activity in environmental communities: limitations and uses. ISME J 7, 2061–2068. https://doi.org/10.1038/ismej.2013.102

Bowman, J.C., Petrov, A.S., Frenkel-Pinter, M., Penev, P.I., Williams, L.D., 2020. Root of the Tree: The Significance, Evolution, and Origins of the Ribosome. Chem. Rev. 120, 4848–4878. https://doi.org/10.1021/acs.chemrev.9b00742

Bray, M.S., Lenz, T.K., Haynes, J.W., Bowman, J.C., Petrov, A.S., Reddi, A.R., Hud, N.V., Williams, L.D., Glass, J.B., 2018. Multiple prebiotic metals mediate translation. Proceedings of the National Academy of Sciences 115, 12164–12169. https://doi.org/10.1073/pnas.1803636115

Bruining, J., Fijnaut, H.M., 1979. A rotational diffusion coefficient of the 70s ribosome determined by depolarized laser light scattering. Biophysical Chemistry 9, 345–353. https://doi.org/10.1016/0301-4622(75)80049-4

Bywaters, K.B., Schmidt, H., Vercoutere, W., Deamer, D., Hawkins, A.R., Quinn, R.C., Burton, A.S., Mckay, C.P., 2017. Development of Solid-State Nanopore Technology for Life Detection. Presented at the Annual Astrobiology Science Conference 2017 (AbSciCon) 2017, Mesa, AZ.

Carr, C.E., 2022. Resolving the History of Life on Earth by Seeking Life As We Know It on Mars. Astrobiology. https://doi.org/10.1089/ast.2021.0043

Chambon, P., Deutscher, M.P., Kornberg, A., 1968. Biochemical Studies of Bacterial
Sporulation and Germination. Journal of Biological Chemistry 243, 5110–5116.
https://doi.org/10.1016/S0021-9258(18)91998-8
Chou, L., Mahaffy, P., Trainer, M., Eigenbrode, J., Arevalo, R., Brinckerhoff, W., Getty, S.,
Grefenstette, N., Da Poian, V., Fricke, G.M., Kempes, C.P., Marlow, J., Sherwood Lollar,
B., Graham, H., Johnson, S.S., 2021. Planetary Mass Spectrometry for Agnostic Life
Detection in the Solar System. Frontiers in Astronomy and Space Sciences 8.
Comolli, L.R., Baker, B.J., Downing, K.H., Siegerist, C.E., Banfield, J.F., 2009. Three-
dimensional analysis of the structure and ecology of a novel, ultra-small archaeon. ISME
J 3, 159–167. https://doi.org/10.1038/ismej.2008.99
Cuevas-Zuviría, B., Adam, Z.R., Goldman, A.D., Kaçar, B., 2023. Informatic Capabilities of
Translation and Its Implications for the Origins of Life. J Mol Evol.
https://doi.org/10.1007/s00239-023-10125-0
Davila, A.F., Schulze-Makuch, D., 2016. The Last Possible Outposts for Life on Mars.
Astrobiology 16, 159–168. https://doi.org/10.1089/ast.2015.1380
Ekland, E.H., Bartel, D.P., 1996. RNA-catalysed RNA polymerization using nucleoside
triphosphates. Nature 382, 373–376. https://doi.org/10.1038/382373a0
Farimani, A.B., Heiranian, M., Aluru, N.R., 2018. Identification of amino acids with sensitive
nanoporous MoS2: towards machine learning-based prediction. npj 2D Mater Appl 2, 1–
9. https://doi.org/10.1038/s41699-018-0060-8
Fournier, G.P., Neumann, J.E., Gogarten, J.P., 2010. Inferring the Ancient History of the
Translation Machinery and Genetic Code via Recapitulation of Ribosomal Subunit
Assembly Orders. PLOS ONE 5, e9437. https://doi.org/10.1371/journal.pone.0009437
Fu, J., Munro, J.B., Blanchard, S.C., Frank, J., 2011. Cryoelectron microscopy structures of the
ribosome complex in intermediate states during tRNA translocation. Proceedings of the
National Academy of Sciences 108, 4817–4821.
https://doi.org/10.1073/pnas.1101503108
Gilbert, W., 1986. Origin of life: The RNA world. Nature 319, 618–618.
https://doi.org/10.1038/319618a0
Goldman, A.D., Samudrala, R., Baross, J.A., 2010. The evolution and functional repertoire of
translation proteins following the origin of life. Biology Direct 5, 15.
https://doi.org/10.1186/1745-6150-5-15
Harish, A., Caetano-Anollés, G., 2012. Ribosomal History Reveals Origins of Modern Protein
Synthesis. PLOS ONE 7, e32776. https://doi.org/10.1371/journal.pone.0032776
Hattori, S., Sekido, R., Leong, I.W., Tsutsui, M., Arima, A., Tanaka, M., Yokota, K., Washio, T.,
Kawai, T., Okochi, M., 2020. Machine learning-driven electronic identifications of single
pathogenic bacteria. Sci Rep 10, 15525. https://doi.org/10.1038/s41598-020-72508-3
Heiman, M., Schaefer, A., Gong, S., Peterson, J., Day, M., Ramsey, K.E., Suárez-Fariñas, M.,
Schwarz, C., Stephan, D.A., Surmeier, D.J., Greengard, P., Heintz, N., 2008.
Development of a BACarray translational profiling approach for the molecular
characterization of CNS cell types. Cell 135, 738–748.
https://doi.org/10.1016/j.cell.2008.10.028
Horning, D.P., Joyce, G.F., 2016. Amplification of RNA by an RNA polymerase ribozyme.
Proceedings of the National Academy of Sciences 113, 9786–9791.
https://doi.org/10.1073/pnas.1610103113

Hsiao, C., Mohan, S., Kalahar, B.K., Williams, L.D., 2009. Peeling the Onion: Ribosomes Are
Ancient Molecular Fossils. Molecular Biology and Evolution 26, 2415–2425.
https://doi.org/10.1093/molbev/msp163
Hsu, H.-W., Postberg, F., Sekine, Y., Shibuya, T., Kempf, S., Horányi, M., Juhász, A., Altobelli,
N., Suzuki, K., Masaki, Y., Kuwatani, T., Tachibana, S., Sirono, S., Moragas-
Klostermeyer, G., Srama, R., 2015. Ongoing hydrothermal activities within Enceladus.
Nature 519, 207–210. https://doi.org/10.1038/nature14262
Isenbarger, T.A., Carr, C.E., Johnson, S.S., Finney, M., Church, G.M., Gilbert, W., Zuber, M.T.,
Ruvkun, G., 2008. The Most Conserved Genome Segments for Life Detection on Earth
and Other Planets. Orig Life Evol Biosph 38, 517–533. https://doi.org/10.1007/s11084-
008-9148-z
Jerome, C.A., Kim, H.-J., Mojzsis, S.J., Benner, S.A., Biondi, E., 2022. Catalytic Synthesis of
Polyribonucleic Acid on Prebiotic Rock Glasses. Astrobiology.
https://doi.org/10.1089/ast.2022.0027
Johnson, S.S., Anslyn, E.V., Graham, H.V., Mahaffy, P.R., Ellington, A.D., 2018. Fingerprinting
Non-Terran Biosignatures. Astrobiology 18, 915–922.
https://doi.org/10.1089/ast.2017.1712
Jones, R.M., Goordial, J.M., Orcutt, B.N., 2018. Low Energy Subsurface Environments as
Extraterrestrial Analogs. Frontiers in Microbiology 9.
Jose, N., Deshmukh, G.P., Ravindra, M.R., 2019. Dynamic Light Scattering: Advantages and
Applications. Dynamic Light Scattering 3.
Joyce, G., 1994. Foreword. In Origins of Life: The Central Concepts, edited by Deamer D.W.
and Fleischaker G.R. Jones and Bartlett Publishers, Boston.
Kim, M., Chun, J., 2014. Chapter 4 - 16S rRNA Gene-Based Identification of Bacteria and
Archaea using the EzTaxon Server, in: Goodfellow, M., Sutcliffe, I., Chun, J. (Eds.),
Methods in Microbiology, New Approaches to Prokaryotic Systematics. Academic Press,
pp. 61–74. https://doi.org/10.1016/bs.mim.2014.08.001
Klempay, B., Arandia☐Gorostidi, N., Dekas, A.E., Bartlett, D.H., Carr, C.E., Doran, P.T., Dutta,
A., Erazo, N., Fisher, L.A., Glass, J.B., Pontefract, A., Som, S.M., Wilson, J.M., Schmidt,
B.E., Bowman, J.S., 2021. Microbial diversity and activity in Southern California salterns
and bitterns: analogues for remnant ocean worlds. Environ Microbiol 23, 3825–3839.
https://doi.org/10.1111/1462-2920.15440
Kowalczyk, S.W., Grosberg, A.Y., Rabin, Y., Dekker, C., 2011. Modeling the conductance and
DNA blockade of solid-state nanopores. Nanotechnology 22, 315101.
https://doi.org/10.1088/0957-4484/22/31/315101
Lafontaine, D.L.J., Tollervey, D., 2001. The function and synthesis of ribosomes. Nat Rev Mol
Cell Biol 2, 514–520. https://doi.org/10.1038/35080045
Lawrence, J.D., Mullen, A.D., Bryson, F.E., Chivers, C.J., Hanna, A.M., Plattner, T., Spiers,
E.M., Bowman, J.S., Buffo, J.J., Burnett, J.L., Carr, C.E., Dichek, D.J., Hughson, K.H.G.,
King, W., Lightsey, E.G., Ingall, E., McKaig, J., Meister, M.R., Pierson, S., Tomar, Y.,
Schmidt, B.E., 2023. Subsurface Science and Search for Life in Ocean Worlds. Planet.
Sci. J. 4, 22. https://doi.org/10.3847/PSJ/aca6ed
Leininger, S.E., Rodriguez, J., Vu, Q.V., Jiang, Y., Li, M.S., Deutsch, C., O'Brien, E.P., 2021.
Ribosome Elongation Kinetics of Consecutively Charged Residues Are Coupled to
Electrostatic Force. Biochemistry 60, 3223–3235.
https://doi.org/10.1021/acs.biochem.1c00507

Li, G.-W., Burkhardt, D., Gross, C., Weissman, J.S., 2014. Quantifying absolute protein
synthesis rates reveals principles underlying allocation of cellular resources. Cell 157,
624–635. https://doi.org/10.1016/j.cell.2014.02.033
Luisi, P.L., 1998. About Various Definitions of Life. Origins of Life and Evolution of
Biospheres 28, 613–622.
Marshall, S.M., Mathis, C., Carrick, E., Keenan, G., Cooper, G.J.T., Graham, H., Craven, M.,
Gromski, P.S., Moore, D.G., Walker, Sara.I., Cronin, L., 2021. Identifying molecules as
biosignatures with assembly theory and mass spectrometry. Nat Commun 12, 3033.
https://doi.org/10.1038/s41467-021-23258-x
Meier, P., Kriegel, H., Motschmann, U., Schmidt, J., Spahn, F., Hill, T.W., Dong, Y., Jones,
G.H., 2014. A model of the spatial and size distribution of Enceladus′ dust plume.
Planetary and Space Science 104, 216–233. https://doi.org/10.1016/j.pss.2014.09.016
Mileikowsky, C., Cucinotta, F.A., Wilson, J.W., Gladman, B., Horneck, G., Lindegren, L.,
Melosh, J., Rickman, H., Valtonen, M., Zheng, J.Q., 2000. Natural Transfer of Viable
Microbes in Space: 1. From Mars to Earth and Earth to Mars. Icarus 145, 391–427.
https://doi.org/10.1006/icar.1999.6317
Müller, J.J., Zalkova, T.N., Ziwer, D., Misselwitz, R., Gast, K., Serdyuk, I.N., Welfle, H.,
Damaschun, G., 1986. Comparison of the structure of ribosomal 5S RNA from E. coli
and from rat liver using X-ray scattering and dynamic light scattering. Eur Biophys J 13,
301–307. https://doi.org/10.1007/BF00254212
Murphy, E.L., Singh, K.V., Avila, B., Kleffmann, T., Gregory, S.T., Murray, B.E., Krause, K.L.,
Khayat, R., Jogl, G., 2020. Cryo-electron microscopy structure of the 70S ribosome from
Enterococcus faecalis. Sci Rep 10, 1–11. https://doi.org/10.1038/s41598-020-73199-6
National Academies of Sciences, Engineering, and Medicine, 2022. Origins, Worlds, and Life: A
Decadal Strategy for Planetary Science and Astrobiology 2023-2032. The National
Academies Press, Washington, D.C. https://doi.org/10.17226/26522
Neveu, M., Anbar, A.D., Davila, A.F., Glavin, D.P., MacKenzie, S.M., Phillips-Lander, C.M.,
Sherwood, B., Takano, Y., Williams, P., Yano, H., 2020. Returning Samples From
Enceladus for Life Detection. Frontiers in Astronomy and Space Sciences 7.
Neveu, M., Hays, L.E., Voytek, M.A., New, M.H., Schulte, M.D., 2018. The Ladder of Life
Detection. Astrobiology 18, 1375–1402. https://doi.org/10.1089/ast.2017.1773
Nikolaeva, D.D., Gelfand, M.S., Garushyants, S.K., 2021. Simplification of Ribosomes in
Bacteria with Tiny Genomes. Molecular Biology and Evolution 38, 58–66.
https://doi.org/10.1093/molbev/msaa184
Oren, A., 2008. Microbial life at high salt concentrations: phylogenetic and metabolic diversity.
Aquat. Biosyst. 4, 2. https://doi.org/10.1186/1746-1448-4-2
Orgel, L.E., 1968. Evolution of the genetic apparatus. J Mol Biol 38, 381–393.
https://doi.org/10.1016/0022-2836(68)90393-8
Pace, N.R., 2001. The universal nature of biochemistry. Proceedings of the National Academy of
Sciences 98, 805–808. https://doi.org/10.1073/pnas.98.3.805
Pappalardo, R.T., Vance, S., Bagenal, F., Bills, B.G., Blaney, D.L., Blankenship, D.D.,
Brinckerhoff, W.B., Connerney, J.E.P., Hand, K.P., Hoehler, T.M., Leisner, J.S., Kurth,
W.S., McGrath, M.A., Mellon, M.T., Moore, J.M., Patterson, G.W., Prockter, L.M.,
Senske, D.A., Schmidt, B.E., Shock, E.L., Smith, D.E., Soderlund, K.M., 2013. Science
Potential from a Europa Lander. Astrobiology 13, 740–773.
https://doi.org/10.1089/ast.2013.1003

Patel, V.B., Cunningham, C.C., 2002. Altered Hepatic Mitochondrial Ribosome Structure
Following Chronic Ethanol Consumption. Archives of Biochemistry and Biophysics 398,
41–50. https://doi.org/10.1006/abbi.2001.2701
Petrov, A.S., Bernier, C.R., Hsiao, C., Norris, A.M., Kovacs, N.A., Waterbury, C.C., Stepanov,
V.G., Harvey, S.C., Fox, G.E., Wartell, R.M., Hud, N.V., Williams, L.D., 2014.
Evolution of the ribosome at atomic resolution. Proceedings of the National Academy of
Sciences 111, 10251–10256. https://doi.org/10.1073/pnas.1407205111
Petrov, A.S., Gulen, B., Norris, A.M., Kovacs, N.A., Bernier, C.R., Lanier, K.A., Fox, G.E.,
Harvey, S.C., Wartell, R.M., Hud, N.V., Williams, L.D., 2015. History of the ribosome
and the origin of translation. Proceedings of the National Academy of Sciences 112,
15396–15401. https://doi.org/10.1073/pnas.1509761112
Phillips, M.S., McInenly, M., Hofmann, M.H., Hinman, N.W., Warren-Rhodes, K., Rivera-
Valentín, E.G., Cabrol, N.A., 2023. Salt Constructs in Paleo-Lake Basins as High-Priority
Astrobiology Targets. Remote Sensing 15, 314. https://doi.org/10.3390/rs15020314
Rahman, M., Stott, M.A., Harrington, M., Li, Y., Sampad, M.J.N., Lancaster, L., Yuzvinsky,
T.D., Noller, H.F., Hawkins, A.R., Schmidt, H., 2019. On demand delivery and analysis
of single molecules on a programmable nanopore-optofluidic device. Nat Commun 10,
3712. https://doi.org/10.1038/s41467-019-11723-7
Rampersad, S., Tennant, P., 2018. Replication and Expression Strategies of Viruses. Viruses 55–
82. https://doi.org/10.1016/B978-0-12-811257-1.00003-6
Raveendran, M., Leach, A.R., Hopes, T., Aspden, J.L., Actis, P., 2020. Ribosome Fingerprinting
with a Solid-State Nanopore. ACS Sens 5, 3533–3539.
https://doi.org/10.1021/acssensors.0c01642
Rudenko, M.I., Holmes, M.R., Ermolenko, D.N., Lunt, E.J., Gerhardt, S., Noller, H.F., Deamer,
D.W., Hawkins, A., Schmidt, H., 2011. Controlled gating and electrical detection of
single 50S ribosomal subunits through a solid-state nanopore in a microfluidic chip.
Biosensors and Bioelectronics 29, 34–39. https://doi.org/10.1016/j.bios.2011.07.047
Runnels, C.M., Lanier, K.A., Williams, J.K., Bowman, J.C., Petrov, A.S., Hud, N.V., Williams,
L.D., 2018. Folding, Assembly, and Persistence: The Essential Nature and Origins of
Biopolymers. J Mol Evol 86, 598–610. https://doi.org/10.1007/s00239-018-9876-2
Schostag, M.D., Albers, C.N., Jacobsen, C.S., Priemé, A., 2020. Low Turnover of Soil Bacterial
rRNA at Low Temperatures. Frontiers in Microbiology 11.
Sczepanski, J.T., Joyce, G.F., 2014. A cross-chiral RNA polymerase ribozyme. Nature 515, 440–
442. https://doi.org/10.1038/nature13900
Shore, D., Albert, B., 2022. Ribosome biogenesis and the cellular energy economy. Current
Biology 32, R611–R617. https://doi.org/10.1016/j.cub.2022.04.083
Shukla, S.K., Mikkola, J.-P., 2020. Use of Ionic Liquids in Protein and DNA Chemistry. Front
Chem 8, 598662. https://doi.org/10.3389/fchem.2020.598662
Spitnik-Elson, P., Atsmon, A., 1969. Detachment of ribosomal proteins by salt: I. Effect of
conditions on the amount of protein detached. Journal of Molecular Biology 45, 113–124.
https://doi.org/10.1016/0022-2836(69)90214-9
Stetefeld, J., McKenna, S.A., Patel, T.R., 2016. Dynamic light scattering: a practical guide and
applications in biomedical sciences. Biophys Rev 8, 409–427.
https://doi.org/10.1007/s12551-016-0218-6
Tian, F.-J., Zhang, C., Zhou, E., Dong, H.-L., Tan, Z.-J., Zhang, X.-H., Dai, L., 2023.
Universality in RNA and DNA deformations induced by salt, temperature change,

stretching force, and protein binding. Proceedings of the National Academy of Sciences
120, e2218425120. https://doi.org/10.1073/pnas.2218425120
Tsutsui, M., Tanaka, M., Marui, T., Yokota, K., Yoshida, T., Arima, A., Tonomura, W.,
Taniguchi, M., Washio, T., Okochi, M., Kawai, T., 2018. Identification of Individual
Bacterial Cells through the Intermolecular Interactions with Peptide-Functionalized
Solid-State Pores. Anal. Chem. 90, 1511–1515.
https://doi.org/10.1021/acs.analchem.7b04950
VanInsberghe, M., van den Berg, J., Andersson-Rolf, A., Clevers, H., van Oudenaarden, A.,
2021. Single-cell Ribo-seq reveals cell cycle-dependent translational pausing. Nature
597, 561–565. https://doi.org/10.1038/s41586-021-03887-4
Xia, K., Hagan, J.T., Fu, L., Sheetz, B.S., Bhattacharya, S., Zhang, F., Dwyer, J.R., Linhardt,
R.J., 2021. Synthetic heparan sulfate standards and machine learning facilitate the
development of solid-state nanopore analysis. Proceedings of the National Academy of
Sciences 118, e2022806118. https://doi.org/10.1073/pnas.2022806118
Xia, Z., Patchin, M., McKay, C.P., Drndić, M., 2022. Deoxyribonucleic Acid Extraction from
Mars Analog Soils and Their Characterization with Solid-State Nanopores. Astrobiology
22, 992–1008. https://doi.org/10.1089/ast.2021.0051
Xue, L., Yamazaki, H., Ren, R., Wanunu, M., Ivanov, A.P., Edel, J.B., 2020. Solid-state
nanopore sensors. Nat Rev Mater 5, 931–951. https://doi.org/10.1038/s41578-020-0229-6

**Figure Legends**

A. Biomolecules translocating nanopore

B. Sample output from biomolecules translocating nanopore

842 **Figure 1. Examples of DNA and ribosome detection using a solid-state nanopore (SSN) in**

843 **2M LiCl.** Molecules passing through a SSN (A) cause momentary blockages in the membrane's

844 ionic current (B). In later analysis, the change in amplitude is reported as the change in

845 conductance (since the voltage applied is constant). Note the order of magnitude difference in the

846 vertical axis.

Figure 2. Overlaid event data (conductance vs. dwell time) for biomolecule (A) and buffer

(B) samples in 2M LiCl. Buffer and biomolecule samples were run in the following order: buf1,

pUC19, buf2, DNA1kb, buf3, DNAHR, buf4 (Table S1). Buffer (negative control) event counts

demonstrate low carry-over between different samples as evidenced by low (<10) event

abundance for run times of approximately 3 minutes for each buffer, and 6-10 minutes for each

biomolecule sample.

**Figure 3. Event data for DNA plasmid pUC19 (A-C), 1 kilobase ladder (D-F), high-range ladder (G-I), and all DNA types (J-K) in 2M LiCl.** Left panels (A, D, G, J) show selected event examples. Middle panels (B, E, H, K) show heat maps of change in conductance vs. dwell

time (time molecule spends disrupting nanopore's ionic current). Heat map color scales

correspond to probability of an event being in that region of the heat map. Red boxes indicate

regions of interest for right panels. Right panels (C, F, I) show all overlaid events within red

boxes with a signal-to-noise ratio greater than 5, with time scaled to 1. Panel L shows schematic

representations of the three measured types of DNA. In Panel K, each probability distribution

(pUC19, 1kb ladder, high range ladder) is evenly weighted to facilitate comparison between the

different classes.

**Figure 4. Event data for RNA samples: double-stranded RNA (A-C), single-stranded RNA**

**(D-F), and both RNA types (G-H).** Left panels (A, D, G) show representative events. Middle

panels (B, E, H) show heat maps of change in conductance vs. dwell time (time molecule spends

disrupting nanopore's ionic current). Heat map color scales correspond to probability of an event

being in that region of the heat map. Red boxes indicate regions of interest for right panels. Right

panels (C, F) show all overlaid events within red boxes, with time scaled to 1. Panel I shows

schematic representations of the two measured types of RNA. In Panel H, each probability

distribution (dsRNA ladder, ssRNA ladder) is evenly weighted to facilitate comparison between

the different classes.

**Figure 5. Event data for *E. coli* ribosome in 2M LiCl.** Panel A shows heat map of change in

conductance vs. dwell time (time molecule spends disrupting nanopore's ionic current). Heat

map color scale corresponds to a probability of an event being in that region of the heat map.

Panel B shows example events interpreted to be an intact ribosome (corresponding to Panel C)

and a ribosomal fragment (corresponding to Panel D). Panels C and D show overlaid events

within the red boxes C and D on Panel A, plotting all events with a signal-to-noise ratio greater

than 10, with time scaled to 1. Event data around a current baseline of 0 pA can be observed in C

and D throughout the middle of the scaled events (e.g. events with a current signal close to the

baseline along time axis points of 0.4 - 0.6). This is due to instances of the automatic event

extraction selecting an event with 2 translocations (due to the translocations happening close to

each other temporally).

**Figure 6. Scatter plots of conductance vs. dwell time for various biomolecule types, sizes,**

**and conformations.** Panel A includes all ribosome, DNA, and RNA samples. Panel B excludes

ribosomes so that distribution of DNA and RNA events can be seen more clearly.

|  | DNA1kb | DNAhr | dsRNA | pUC19 | ribosome | ssRNA | | TPR | FNR |
|---|---|---|---|---|---|---|---|---|---|
| **PPV** | 63.0% | 89.0% | 56.2% | 92.2% | 98.5% | 88.5% | | | |
| **FDR** | 37.0% | 11.0% | 43.8% | 7.8% | 1.5% | 11.5% | | | |
| **DNA1kb** | 153 | 34 | 24 | 11 | 6 | | | 67.1% | 32.9% |
| **DNAhr** | 61 | 624 | 37 | 7 | 8 | 1 | | 84.6% | 15.4% |
| **dsRNA** | 1 | 17 | 194 | 6 | 5 | 30 | | 76.7% | 23.3% |
| **pUC19** | 7 | 10 | 8 | 507 | 43 | | | 88.2% | 11.8% |
| **ribosome** | 21 | 15 | 32 | 10 | 4211 | | | 98.2% | 1.8% |
| **ssRNA** | | 1 | 50 | 9 | | 238 | | 79.9% | 20.1% |

**Figure 7. Confusion matrix for Medium Tree model.** This matrix was generated by the Medium Tree model by the MATLAB Classification Learner app (MATLAB_R2023a). True Class vs. Predicted Class matrix shows the raw numbers of events for each true/predicted class combination. The upper matrix shows the likelihood that an event is correctly classified given the true class by showing the Positive Predictive Value (PPV), the percentage of correctly classified events within all events of that class, and the False Discovery Rate (FDR), the percentage of incorrectly classified events within all events of that class. The right matrix shows how many of the events were correctly predicted within a predicted class by showing the True Positive Rate (TPR), the percentage of correctly classified events within all events assigned to that class, and the False Negative Rate (FNR), the percentage of incorrectly classified events within all events assigned to that class.