Decoding host-microbiome interactions through co-expression network analysis within the

non-human primate intestine

Mika Uehara[1], Takashi Inoue[2], Sumitaka Hase[1], Erika Sasaki[2, 4], Atsushi Toyoda[3], Yasubumi

Sakakibara[1*]

1) Department of Biosciences and Informatics, Keio University, Yokohama, Kanagawa 223-

8522, Japan

2) Department of Marmoset Biology and Medicine, Central Institute for Experimental Animals,

Kawasaki, Kanagawa 210-0821, Japan

3) Department of Genomics and Evolutionary Biology, National Institute of Genetics, Mishima,

Shizuoka 411-8540, Japan

4) Laboratory for Marmoset Neural Architecture, RIKEN Center for Brain Science, 2-1

Hirosawa, Wako-shi, Saitama 351-0198, Japan.

*Corresponding Author

Yasubumi Sakakibara

3-14-1 Hiyoshi, Kohoku-ku, Yokohama, 223-8522, Japan

Phone/Fax: +81-45-566-1791

E-mail: yasu@bio.keio.ac.jp

ABSTRACT

The gut microbiome affects the health status of the host through complex interactions with the

host's intestinal wall. These host-microbiome interactions may spatially vary along the physical

and chemical environment of the intestine, but these changes remain unknown. This study

investigated these intricate relationships through a gene co-expression network analysis based on

dual transcriptome profiling of different intestinal sites—cecum, transverse colon, and rectum—

of the primate common marmoset. We proposed a gene module extraction algorithm based on

graph theory to find tightly interacting gene modules of the host and the microbiome from a vast

co-expression network. The 27 gene modules identified by this method, which include both host

and microbiome genes, not only produced results consistent with previous studies regarding the

host-microbiome relationships, but also provided new insights into microbiome genes acting as

potential mediators in host-microbiome interplays. Specifically, we discovered associations

between the host gene *FBP1*, a cancer marker, and polysaccharide degradation-related genes

(*pfkA* and *fucI*) coded by *Bacteroides vulgatus*, as well as relationships between host B-cell

specific genes (*CD19*, *CD22*, *CD79B*, *PTPN6*) and a tryptophan synthesis gene (*trpB*) coded by

Parabacteroides distasonis. Furthermore, our proposed module extraction algorithm surpassed

existing approaches by successfully defining more functionally related gene modules, providing

insights for understanding the complex relationship between the host and the microbiome.

IMPORTANCE We unveiled the intricate dynamics of the host-microbiome interactions along

the colon by identifying closely interacting gene modules from a vast gene co-expression

network, constructed based on simultaneous profiling of both host and microbiome

transcriptomes. Our proposed gene module extraction algorithm, designed to interpret inter-

species interactions, enabled the identification of functionally related gene modules

encompassing both host and microbiome genes, which was challenging with conventional

modularity maximization algorithms. Through these identified gene modules, we discerned

previously unrecognized bacterial genes that potentially mediate in known relationships between

host genes and specific bacterial species. Our findings underscore the spatial variations in host-

microbiome interactions along the colon, rather than displaying a uniform pattern throughout the

colon.

INTRODUCTION

The mutual interaction between the intestinal microbiome and the host constitutes a complex

biological process, with intertwined dependencies that are essential for the survival and health of

both. The interactions between the intestinal microbiome and the host are formed by various

factors such as the effects of metabolites produced by bacteria on the host, the binding of

bacterial-derived proteins and host receptors, and the binding of bacterial cell surface structures

and host receptors. Studies on host-microbiome interaction through bacterial-derived proteins

have discovered that a novel protein P9 secreted by *Akkermansia muciniphila* binds to host

ICAM-2, thereby increasing the secretion of glucagon-like peptide 1 (GLP-1), suggesting that

the interaction between P9 and ICAM-2 could be a potential therapeutic target in type 2 diabetes

patients (1). Excluding certain gram-negative bacteria, it has also been reported that bacterial

lipopolysaccharides (LPS), which are components of the gram-negative bacterial cell wall, bind

to the Toll-like receptor 4 (TLR4) complex and can induce innate immune responses (2). Among

these interactions, the host-microbiome interaction mediated by metabolites is particularly well-

studied. For instance, primary bile acids, produced in the host's liver cells, are converted into

secondary bile acids by the microbiome in the colon. This microbiome-derived secondary bile

acid is known to regulate the homeostasis of regulatory T cells, which are immune-suppressive

cells in the host (3). Through these processes, the intestinal microbiome and host mutually

interact and establish a symbiotic relationship while forming a metabolic pathway or

physiological response.

To understand such complex interactions between the host and the microbiome, it is

beneficial to analyze combined profiles of both the microbiome and the host. Recent studies have

reported on dual profiling of host and microbiome gene expression levels to understand the

interactions between the host and microbiome (4, 5, 6). Previously, the focus was primarily on

identifying bacterial species related to host phenotypes (such as diseases) based on the relative

abundance of microbiome (7). However, profiling both host and microbiome gene expression

levels has provided further insights into which genes of which bacterial species are related to

host physiological functions and phenotypes. Gene co-expression network analysis is one of the

useful approaches for comprehensively deciphering complex interactions from the vast number

of gene expression profiles of the host and microbiome. A gene co-expression network is

constructed by representing genes as nodes and connecting genes with correlated expression

levels via edges. Representing gene relationships as a network enables the extraction of closely

interconnected gene modules (subgraphs) and the identification of hub genes connected to

multiple genes, thereby enhancing interpretability. For instance, a study (5) characterizing the

host-microbiome interaction during influenza virus infection identified modules from the host's

gene co-expression network. It discovered correlations between host gene modules involved in

interferon signal transduction and the sulfonamide antibiotic resistance genes of the microbiome.

In another study (6), network analysis was applied to the gene expression profiles of the host and

microbiome in the nasopharyngeal airway during bronchiolitis. Networks were constructed for

both the host and microbiome and modules were identified from these networks. The study

revealed that host modules were involved in regulating host-T-cell activity. Additionally,

microbiome modules involved in microbial-BCAA metabolism and oxidative stress response

were associated with disease severity.

Despite such progress, several limitations remain in understanding the interactions

between the host and the intestinal microbiome. First, there's a lack of understanding of the

spatial variation in host-microbiome interactions. The colon, rich in microbiota, can be

subdivided into the cecum, ascending colon, transverse colon, descending colon, sigmoid colon,

and rectum. Each section possesses distinct functions and differs in its physicochemical

environment, including variations in nutrient content, pH, and luminal flow (8, 9, 10) and the

microbiome could vary as well. Indeed, studies using animal models have reported that the

composition and functional fluctuation of the microbiome varies due to environmental changes

depending on intestinal location (biogeographical location) (11, 12, 13). However, these studies

focused solely on the microbiome, and the coordinated spatial variation of both the microbiome

and host gut cells along the intestinal tract has yet to be clarified.

Second, there is a lack of computational methods for extracting significant modules

(subgraphs) from the host-microbiome gene co-expression network. Analyzing a massive gene

co-expression network integrating the gene expression profiles of both the host and microbiome

helps interpret the interactions between the genes of the host and the microbiome. Especially in a

huge gene co-expression network, it is crucial to extract densely connected subgraphs (modules)

to divide the network into understandable units; however, existing algorithms for extracting

modules from the network are insufficient for accurately capturing the interaction between the

host and microbiome because they were not designed with the purpose of understanding

interactions between organisms (14, 15, 16, 17).

Lastly, there are concerns regarding the applicability of the results of studies using animal

models to humans. While mouse models have been the basis of biological research for a long

time due to ethical constraints, experiments in controlled environments, and considerations of

reproducibility, there are several issues and limitations to their use. The physiology, anatomy, and

pharmacology of mice differ from those of humans, and there are differences in the composition

and function of the intestinal microbiome between mice and humans (13, 18). Particularly

considering the complexity of host-microbiome interactions, it is challenging to precisely

replicate the relationship between humans and their intestinal microbiome. In contrast, the

common marmoset, a small New World primate, is considered a useful model in preclinical trials

because it is genetically close to humans and has common physiological and anatomical

characteristics (19). Moreover, the common marmoset is the only non-human primate in which a

germ-free state has been successfully achieved (20), and it has a fecal microbiota similar to that

of humans compared to other animal models, including mice (13), potentially broadening the

scope of intestinal microbiome research.

To address these challenges, this study aimed to reveal the interactions between the host

and the microbiome along multiple sites of the common marmoset colon—cecum, transverse

colon, and rectum—by developing a module extraction method applicable to the gene co-

expression network that represents the complex interactions between the host and the

microbiome. Through simultaneous profiling of host and microbiome transcripts in the intestinal

sites of the common marmoset, we constructed an integrated gene co-expression network,

incorporating both host and microbiome genes. From this network, we further identified densely

covariant gene modules (subgraphs) to interpret the interactions between genes. As conventional

methods for extracting gene modules from the network were insufficient for interpreting the

interactions between organisms examined in this study, we proposed a module extraction

algorithm based on graph theory. Our proposed method successfully extracted more modules

containing functionally related genes of both the host and microbiome compared to existing

algorithms, underlining its suitability for predicting host-microbiome interactions. This method

extracted 27 gene modules containing both host and microbiome, indicating that the gut

microbiome is functionally closely involved in the variation of host gene expression along the

colon.

RESULTS

Overview of the analysis pipeline integrating host and microbiome transcriptomes:

Identification of closely interacting gene modules from gene co-expression networks

To investigate the host and microbiome interactions along the colon, we proposed an analysis

method integrating the host transcriptome and the microbiome metatranscriptome (Fig. 1). First,

we computed gene expression profiles from the host transcriptome and microbiome

metatranscriptome obtained via dual RNA-seq and created three types of gene co-expression

networks: host-host, host-microbiome, and microbiome-microbiome. By unifying these three

networks, we constructed a host and microbiome gene co-expression network, with the genes of

the host and microbiome serving as nodes, and covariant genes interconnected. Second, from the

gene co-expression network, we identified gene modules containing closely interacting genes

from both the host and microbiome. We aimed to interpret the interactions between host genes

and microbiome genes within modules extracted from a gene co-expression network, based on

the assumption that functionally similar genes are co-expressed (21). Conventional module

extraction methods are not designed to understand interspecies interactions (14, 15, 16, 17) and

do not distinguish between host gene and microbiome gene nodes in the network. Consequently,

modules composed solely of host genes or exclusively of microbiome genes can be extracted,

making it challenging to understand the interactions between the host and the microbiome when

such single species modules are present. Therefore, we proposed a clique-finding-based module

extraction method that takes into account the structure of three networks: host-host, host-

microbiome, and microbiome-microbiome. This method begins by exploring cliques from the

gene co-expression network. In graph theory, a "clique" refers to a subgraph within the network

where all nodes are fully connected. For overlapping cliques, the method calculates the

cliqueness in each of the original three networks and merges them if the cliqueness of all

networks exceeds a certain threshold, thereby obtaining the final gene modules (refer to the

MATERIALS AND METHODS section for details).

Computing gene expression profiles of the host and the microbiome from dual RNA-seq data

First, we extracted high-quality RNA (RNA integrity number (RIN) scores ≥8.6; Table S1) from

the intestinal sites (cecum, transverse colon, and rectum) of five common marmosets

(Supplementary Note Section 1) and carried out dual RNA-seq with NovaSeq 6000 v1.5 (100 bp

PE). The obtained RNA read data amounted to a total of 4,724 M reads (mean 157,450,033

paired-end reads/sample; Table S2; Supplementary Note Section 2).

Next, as the RNA read data obtained by dual RNA-seq contains a mixture of reads

derived from both the host and the microbiome, it is necessary to separate these read data in

silico (Table S3; Supplementary Note Section 3). The pipeline established in this study separates

the host-derived read data and the microbiome-derived read data by aligning the RNA reads to

the host reference genome (Fig. 1B). To avoid transcriptomic cross-contamination of read data

between host and microbiome read data, we adjusted the pipeline parameters using simulated

RNA read data from the host and the microbiome (Supplementary Note Section 4). As a result,

we achieved a precision of 1.0 in separating the read data of the host and the microbiome, and

our established pipeline completely separated the reads derived from the host and those from the

microbiome (Table S9 posted at https://doi.org/10.5281/zenodo.10425960).

After the above preprocessing, we calculated the gene expression levels (transcripts per

million (TPM)) for both the host and the microbiome. As a result, we detected 16,578 host

expressed genes and 8,533 microbiome expressed genes in total (Supplementary Note Section 5

and 6). To examine the expression patterns across different sites and across different individuals

for both host genes and microbiome genes, we applied Principal Component Analysis (PCA) to

the gene expression profiles for visualization. There is a clear separation between the host gene

expression profiles for each intestinal site, and the host samples were not grouped based on

individuals (Figures 2A and B). The microbiome profiles at the cecum and rectum, which are the

beginning and end of the colon, respectively, were separated by PC1, with samples from the

transverse colon, situated between the cecum and rectum, spanning across both sites (Figure 2C).

For the microbiome profiles as well as for the host ones, samples were not separated based on

individuals (Figure 2D). These results suggest that both host and microbiome gene expression

profiles are more variable across different sites compared to those across individuals.

Spatial variations in host gene expression among intestinal sites

Of the 16,578 host genes detected in three regions of the common marmoset's intestinal tract—

cecum, transverse colon, and rectum—a total of 2,615 genes showed significant expression

variations among these sites (FDR < 0.01) (Table S4; Supplementary Note Section 5).

Furthermore, upon conducting functional pathway analysis, 40 pathways were identified, all of

which were significantly upregulated in the cecum compared to the transverse colon (normalized

enrichment score > 1.5, FDR < 0.1; Table S10 posted at

https://doi.org/10.5281/zenodo.10425960). In particular, pathways with high normalized

enrichment scores, such as Fc epsilon RI signaling pathway, T cell receptor signaling pathway,

leukocyte transendothelial migration, Yersinia infection, and B cell receptor signaling pathway

were detected, indicating that pathways related to immune functions were upregulated in the

cecum relative to the transverse colon (Fig. 3A). Additionally, a substantial number of genes

involved in these functional pathways were found to be significantly upregulated in the cecum

(FDR < 0.01). For example, among the genes specifically upregulated in the cecum, we found

genes related to B cell receptor signaling. These include *CD19*, which acts as a coreceptor for the

B cell antigen receptor complex and augments B cell activation (22); *CD79A* and *CD79B*,

components of the B cell antigen receptor that generate signals upon antigen recognition (23);

and *CD22* and *CD72*, which inhibit the signal transduction of the B cell antigen receptor (24,

25). In addition, genes associated with T cell receptor signaling were also found as being highly

expressed specifically in the cecum. These include *CD28*, a co-stimulatory receptor for T cell

receptors that activates T cells (26); *CD3D*, a component of CD3 that plays a role in signal

transduction during T cell activation (27); *CD40LG*, which is induced on the T cell surface

following T cell receptor (TCR) activation (28); and *CTLA4*, which attenuates T cell activation

and fine-tunes immune responses (29) (Fig. 3B).

Interpreting host-microbiome interactions through spatial gene co-expression network

analysis within the intestine

We constructed the unified gene co-expression network composed of 1,359 gene nodes, which

embodies co-expression information on host-host, host-microbiome, and microbiome-

microbiome interactions (Table S11, S12, and S13 posted at

https://doi.org/10.5281/zenodo.10425960; Supplementary Note Section 7). Subsequently, to

identify densely co-expressed gene sets within this network, we implemented the clique-based

module extraction algorithm proposed in this study. As a result, we then identified 27 gene

modules containing both host and microbiome genes (Table S14 and S15 posted at

https://doi.org/10.5281/zenodo.10425960). Furthermore, upon conducting a functional pathway

enrichment analysis on these gene modules, we found that 70.4% (19 out of 27) of gene modules

could be functionally characterized (Supplementary Note Section 8).

The modules we identified were characterized by functional pathways including the B

cell receptor signaling pathway, fructose and mannose metabolism, oxidative phosphorylation,

and biosynthesis of amino acids (Fig. 4). In addition, we provided visualizations of the bacterial

species that contribute to microbiome gene expression within these identified modules, together

with host and microbiome genes (Fig.5, 6, 7 and 8).

Module_07, as shown in Figure 5, was identified as a module with a positive correlation

between the host gene Fructose-1,6-bisphosphatase 1 (*FBP1*), which is highly expressed in the

transverse colon, and 40 microbiome genes. These 40 gut microbiome genes included those

associated with fructose-mannose metabolism, such as K01840 (*manB*), K00850 (*pfkA*), and

K01818 (*fucI*), as well as those related to oxidative phosphorylation, such as K00241 (*sdhC*,

frdC), K00426 (*cydB*), K02112 (*atpD*), and K02121 (*ntpE*, *atpE*). The most substantial

contributor to the expression of these microbiome genes was *Bacteroides vulgatus*.

FBP1 is known to have reduced expression in various tumor tissues, including the esophagus

(30), liver (31), lung (32), and ovarian cancer (33). It has been confirmed that its expression

increases with fructose treatment in intestinal organoids (34). Previous studies have identified

FBP1 as a differentially expressed gene in the small intestine of germ-free mice, suggesting that

the intestinal microbiome influences the expression of the host's *FBP1* gene (35), but the

bacterial species and genes affecting the expression of the *FBP1* gene have not been identified.

Bacteroides vulgatus, which contributed to the expression of fructose-mannose metabolism-

related microbiome genes observed in Module_07, has been reported to release fructose by

decomposing polysaccharides (36, 37). In this module, the gene K00850 (*pfkA*) involved in the

fructo-oligosaccharide degradation and the gene K01818 (*fucI*) involved in the 2'-fucosyllactose

degradation were detected, revealing a novel association between polysaccharide degradation

genes encoded by *Bacteroides vulgatus* and the host gene *FBP1*.

In Module_01, the B cell receptor signaling pathway was enriched (Fig. 6). This module featured

covariation between host B-cell specific genes, including *CD19*, *CD22*, *CD79B*, and *PTPN6*, which were

highly expressed in the cecum, and the tryptophan synthesis gene K01696 (*trpB*) from the microbiome.

The upregulation of these host B-cell specific genes indicates an increase in B cells within the cecum.

This tryptophan synthesis gene K01696 (*trpB*) expression was contributed to by *Parabacteroides*

distasonis. Tryptophan is an essential amino acid that, upon uptake by intestinal cells and

subsequent metabolism, can activate the Aryl hydrocarbon receptor and influence B cell immune

responses (38). Tryptophan is also known to ameliorate autoimmune diseases, such as

inflammatory bowel disease (39). *Parabacteroides distasonis* has been suggested to be involved

in various autoimmune conditions in several studies (40, 41). For instance, *Parabacteroides*

distasonis levels have been shown to be lower in multiple sclerosis (MS) patients compared to

healthy controls, and transplanting the intestinal microbiota from MS patients to germ-free mice

resulted in increased severity of autoimmune encephalomyelitis symptoms (40). While previous

studies have reported associations between tryptophan and autoimmune diseases via B-cell

immune responses, as well as between autoimmune diseases and *Parabacteroides distasonis*, this

study identifies a correlation between the tryptophan synthesis gene K01696 (*trpB*) from

Parabacteroides distasonis and B-cell specific genes (*CD19*, *CD22*, *CD79B*, and *PTPN6*).

In Module_21, a group consisting of three host genes and 40 microbiome genes was

identified. The three host genes included *MMP10*, two uncharacterized genes LOC103795407

and *CCDC162P*, all of which showed high expression in the cecum (Fig. 7). The microbiome

genes that showed a positive correlation with these genes included K01491 (*folD*), K00605

(*gcvT*), K01803 (*tpiA*), K00831 (*serC*), K15633 (*gpmI*), K00024 (*mdh*), all of which are involved

in carbon metabolism pathways. *Bacteroides vulgatus* was identified as the bacteria contributing

to the expression of these microbiome genes. The *MMP10* gene plays a crucial role in

maintaining epithelial barrier function and is necessary for resolving colonic epithelial damage

(42). A study examining the host transcriptome and microbiota composition of colorectal cancer

tumors found high expression of *MMP10* in tumor tissues and a significant increase in the genus

Bacteroides, indicating a link between the host gene *MMP10* and the genus *Bacteroides* (43).

These findings are consistent with our observations. Furthermore, in the identified Module_21,

not only the relationship between the host gene *MMP10* and *Bacteroides vulgatus*, but also a

group of microbiome genes related to carbon metabolism pathways showing a positive

correlation with the host gene *MMP10* was identified. One hypothesis explaining the observed

positive correlation between carbon metabolism-related genes encoded by *Bacteroides vulgatus*

and the host gene *MMP10* is that the microbiome may form colonies within the host's mucosal

layer. Mucin, the principal component of mucus secreted by the gastrointestinal mucosal

epithelium, contains galactose and promotes bacterial settlement and adhesion (44), serving as a

nutrient source for certain bacteria. *Bacteroides vulgatus* possesses mucosal adherence factors,

allowing access to host-derived carbon sources (45). The upregulation of *MMP10* in the cecum

might represent a protective response against colonization by *Bacteroides vulgatus*. The

detection of carbon metabolism-related genes of *Bacteroides vulgatus* in this module_21

suggests they might be utilizing carbon sources derived from mucin or dietary glucose.

Specifically, while *Bacteroides vulgatus* cannot degrade human mucin independently, it can

acquire carbon sources from mucin in co-culture with *Akkermansia muciniphila* (46), which

possesses genes detected in this module. When *Akkermansia muciniphila* converts galactose to

glucose, *Bacteroides vulgatus* might then activate these carbon metabolism-related genes to

utilize this glucose (Fig. S1). Our findings indicate a relationship between the carbon

metabolism-related genes of *Bacteroides vulgatus* and the host gene *MMP10*; however, further

experiments are required to ascertain whether the upregulation of *MMP10* serves a protective

role against colonization by *Bacteroides vulgatus* and *Akkermansia muciniphila* or if *Bacteroides*

vulgatus utilizes mucin as a carbon source.

In Module_18, three host genes and 54 microbiome genes were identified, and the

expression of these microbiome genes was further influenced by five bacterial species. The

relationships between these host genes and microbiome genes, as well as between the host genes

and bacterial species, were consistent with existing knowledge about bile acid synthesis and

degradation (Fig. 8). This module included the host gene *CYP27A1*, which is involved in bile

acid production (47), and *NR1H3*, a host gene that codes for a bile acid receptor (48). These

genes were upregulated in the cecum and transverse colon compared to the rectum. The

microbiome genes interacting with these host genes included K03664 (*smpB*), a gene essential

for bacterial survival when upregulated in the presence of secondary bile acid deoxycholic acid,

K02835 (*prfA*), which positively regulates the expression of a gene coding for bile acid

hydrolase, and K03527 (*ispH*, *lytB*), a gene involved in bile acid sensitivity. Additionally,

Bacteroides vulgatus contributed to the expression of all these microbiome genes, which aligns

with previous studies showing an increase in *Bacteroides* spp. correlating with the expression of

the host gene *NR1H3* (49).

DISCUSSION

In this study, we extracted gene modules from a host and microbiome gene co-expression

network based on dual transcriptome profiling. The module extraction method proposed in this

study is based on clique finding and takes into account the structure of three types of networks:

host-host, host-microbiome, and microbiome-microbiome. This approach enables the extraction

of modules containing both host genes and microbiome genes. Conventional module extraction

methods do not differentiate between host gene nodes and microbiome gene nodes within the

network, potentially resulting in modules composed exclusively of either host genes or

microbiome genes. Indeed, when conventional module extraction methods such as the Newman

algorithm (15), the Louvain algorithm (16), the Leiden algorithm (17), and Weighted Correlation

Network Analysis (WGCNA) (18) were applied to our dataset (Supplementary Note Section 9),

we observed that these modules included those composed solely of host genes or microbiome

genes (Table S5). Notably, about half of the genes in WGCNA modules were found in modules

comprising only one species. While the purpose of module extraction is to interpret the

interactions among genes within a module, these single-species modules do not allow for an

interpretation of the relationship between host genes and microbiome genes. Furthermore, to

verify whether functionally similar genes were being extracted, we conducted enrichment

analysis on the extracted modules (Supplementary Note Section 8). The results showed that our

method had the highest proportion of functionally characterized modules (enriched modules)

(Table S6).

The gene modules extracted, which encompassed both host and microbiome genes,

frequently exhibited relationships where a small number of genes regulated a larger number of

genes between the host and microbiome genes. These observations are thought to reflect the

characteristics of the interactions between the host and the microbiome. Several studies have

reported the impact of host gene mutations on the intestinal microbiome. For instance, it has

been reported that the host gene *VDR*, which codes for the Vitamin D receptor, influences the

colonization of *Parabacteroides* (50), and that the host gene *ABO* is associated with

Faecalicatena lactaris (51). In another study investigating the relationship between the intestinal

microbiome and host gene loci through quantitative trait locus analysis, it was suggested that

Akkermansia muciniphila regulates the expression of *ATF3*, which codes for a transcription

factor that plays a crucial role in metabolic and immune regulation, and this bacterium is related

to only four genes (52). These studies highlight the fact that a single or a small number of host

genes are controlling, or are controlled by, the intestinal microbiome. Conversely, a small

number of microbiome genes can affect the host's metabolic (53), fitness (54), and longevity (55)

phenotypes. For example, structural mutations in the gene related to butyrate transport in

Anaerostipes hadrus are strongly associated with a reduction in the host's metabolic risk (53),

and the intestinal microbiome's thiamine biosynthesis genes, transferred by horizontal

transmission, influence the host's fitness (54). Considering these previous studies and our results,

it is suggested that the interaction between the host and the microbiome involves widespread

influence of one's genes on the activity of the other.

The common marmoset used in this study bears anatomical and pharmacological

similarities to humans (19), and the fecal microbiome of the common marmoset is more similar

to that of humans compared to other animal models (13). While the common marmoset's

cecum—one of the study's targeted intestinal sites—does not possess an appendage as

pronounced as the human appendix, it is known to share significant similarities with the human

appendix in terms of microanatomical structure and immune reactivity (56, 57). The results of

this study using common marmosets could therefore hold higher applicability to humans

compared to other animal models.

In conclusion, we generated host transcriptome and microbiome metatranscriptome

profiles of the common marmoset and performed gene co-expression network analysis. The

algorithm we proposed for extracting gene modules in the gene co-expression network enabled

the identification of dense, functionally related host-microbiome co-expression modules, which

was challenging using the conventional module extraction algorithms. The identified gene

modules provided new insights into relationships between host genes and intestinal microbiota

and microbial genes bridging interactions between the host and bacteria. Our findings underscore

that interactions between the host and microbiome are not uniform throughout the colon but are

localized, indicating spatial variation in host-microbiome interactions along the colon. These

insights could be crucial in understanding mechanisms behind localized gut diseases, such as

colorectal cancer and inflammatory bowel disease, and in developing localized therapies (58, 59,

60, 61).

MATERIALS AND METHODS

Animal experiments

Common marmosets were housed at the Central Institute for Experimental Animals (Kawasaki,

Japan). Five marmosets were euthanized with intravenous administration of pentobarbital

overdose, and the digestive tract was isolated (Table S7, Supplementary Note Section 1). Three

gastrointestinal site samples (cecum, transverse colon and rectum), including the mucosal layer,

were used for dual RNA-seq. These three sites were targeted at the beginning, middle and end of

the colon, which has an abundant microbiome. The animal experiment protocol was approved by

the CIEA Institutional Animal Care and Use Committee (approval no. 17031, 18032, and 21084).

The study was conducted in accordance with the guidelines of CIEA that comply with the

Guidelines for Proper Conduct of Animal Experiments published by the Science Council of

Japan. Animal care was conducted in accordance with the Guide for the Care and Use of

Laboratory Animals (Institute for Laboratory Animal Resources, 2011).

RNA extraction and dual RNA sequencing of host and microbiome

RNA libraries were prepared after removing rRNA from the extracted high-quality Total RNA

(RNA integrity number (RIN) scores ≥ 8.6; Table S1). Illumina NovaSeq 6000 v1.5 sequencing

yielded a total of 472 Gnt of paired-end reads (100 nt $\times$ 2). This dataset included an average of

314.9 M reads per sample (Table S2 and S8; Supplementary Note Section 2).

Construction of host-microbiome co-expression network and identification of gene modules

412 In this study, we utilized gene co-expression network analysis to identify interacting gene

413 modules of host and microbiome (Supplementary Note Section 7). We established three types of

414 networks: host-host, host-microbiome, and microbiome-microbiome, wherein genes were

415 considered as nodes and co-expressing genes were interconnected (Fig. 1D). Networks were

416 defined by an adjacency matrix, where the adjacency a_{ij} between gene i and gene j was defined

417 as follows based on co-expression similarity s_{ij}:

418
$$a_{ij} = \begin{cases} 1\,, \text{if } s_{ij} > \alpha \\ 0\,, \text{otherwise} \end{cases}$$

419 The co-expression similarity s_{ij} was computed using the Spearman's correlation coefficient

420 calculated for genes i and j represented by vectors of TPM values for host-host and microbiome-

421 microbiome networks, and absolute values of Spearman's correlation coefficients for host-

422 microbiome networks. The adjacency threshold α was set at 0.80. These three types of networks

423 were then integrated into one.

424  We developed an algorithm to identify densely interacting gene modules from the host-

425 microbiome co-expression network. This algorithm, based on clique finding, progressively

426 merges cliques (Fig. 1E). In graph theory, a "clique" refers to a subgraph within the network

427 where all nodes are fully connected. In the context of a gene co-expression network, all genes

428 within a clique indicate a strong co-expression (correlation) amongst them. In this study, we

429 focus only on networks and cliques that include gene nodes from both the host and microbiome,

430 encompassing three types of networks: host-host, host-microbiome, and microbiome-

431 microbiome. Our algorithm identified modules of interacting genes in the gene co-expression

432 network as aggregates of overlapping cliques. It was implemented in three steps: (1) Cliques

433 within the network were found; (2) Beginning with the largest cliques, we examined their

434 overlap with any other cliques. If overlap occurred, we computed the cliqueness scores

independently for each of the three types of networks (host-host, host-microbiome, and

microbiome-microbiome networks) upon merging the overlapping cliques. The cliqueness scores

for the host-host, host-microbiome, and microbiome-microbiome networks were denoted as C_{hh},

C_{hm}, and C_{mm}, respectively; (3) Cliques were merged if they satisfied the condition: $C_{hh} > \tau$,

$C_{mm} > \tau$, and $C_{mm} > \tau$. The threshold τ for cliqueness was set at 0.60 based on parameter

determination (Supplementary Note Section 7). If there were multiple cliques with minimum

cliqueness scores exceeding the threshold, the clique with the highest cliqueness score was

selected. Steps (2) and (3) were repeated until no cliques could be merged.

For the host-host networks the cliqueness score C_{hh} between cliques A and B was defined as:

444
$$C_{hh} = \frac{E}{\frac{1}{2}N(N-1)}$$

Here, N is defined as the number of host gene nodes in the union of cliques A and B, represented

as $N = |N_A \cup N_B|$. N_A and N_B denote the number of host gene nodes within clique A and clique

B, respectively. E corresponds to the total number of edges between the host nodes included in

cliques A and/or B, formulated as $E = E_A + E_B + E_{AB}$. Here, E_A and E_B represent the number of

edges between host gene nodes within clique A and clique B, respectively, while E_{AB} denotes the

number of edges connecting across host gene nodes within clique A and clique B. The cliqueness

score for microbiome-microbiome networks C_{mm} was calculated in the same way.

For the host-microbiome network, the cliqueness score C_{hm}, was defined as:

453
$$C_{hm} = \frac{E}{N_{host}N_{microbiome}}$$

454   where N_{host} and $N_{microbiome}$ represent the number of host and microbiome gene nodes

455   respectively in the union of cliques A and B. E corresponds to the total number of edges between

456   the host or microbiome gene nodes included in cliques A and/or B.

Data and source code availability

All raw sequence data have been submitted to the DDBJ under project PRJDB16352 from the

Ministry of Education, Culture, Sports, Science and Technology of Japan. The gene expression

profiles of the host and microbiome are available at https://doi.org/10.5281/zenodo.8264866. The

algorithm for module extraction is available at https://github.com/MikaUhr/MEIGN.git.

ACKNOWLEDGEMENTS

We thank the scientists of the Central Institute for Experimental Animals for providing support for

animal experiments and the members of the Sakakibara Lab at Keio University for helpful

discussions.

This work was supported by grants from the Japan Agency for Medical Research and Development

(AMED PRIME) Grant Number JP19gm6010006 and JST, CREST Grant Number JPMJCR20S3,

Japan. M. Uehara has received funding from JSPS KAKENHI Grant Numbers JP20J21477.

M.U. performed the experiments, conducted the bioinformatics analysis and cowrote the paper;

T.I. and E.S. provided common marmoset samples; M.U. and S.H. performed RNA extraction; A.T.

performed deep sequencing with high-throughput sequencer; Y.S. designed and supervised the

research, analysed the data, and cowrote the paper. All authors have read and approved the

manuscript.

The authors declare no competing interests.

REFERENCES

1. Yoon HS, Cho CH, Yun MS, Jang SJ, You HJ, Kim JH, et al. 2021. Akkermansia muciniphila secretes a glucagon-like peptide-1-inducing protein that improves glucose homeostasis and ameliorates metabolic disease in mice. Nat Microbiol 6:563–73.

2. Vatanen T, Kostic AD, D'Hennezel E, Siljander H, Franzosa EA, Yassour M, et al. 2016. Variation in Microbiome LPS Immunogenicity Contributes to Autoimmunity in Humans. Cell 165:842–53.

3. Song X, Sun X, Oh SF, Wu M, Zhang Y, Zheng W, et al. 2020. Microbial bile acid metabolites modulate gut RORγ+ regulatory T cell homeostasis. Nature 577:410–5.

4. Zhang L, Forst CV, Gordon A, Gussin G, Geber AB, Fernandez PJ, et al. 2020. Characterization of antibiotic resistance and host-microbiome interactions in the human upper respiratory tract during influenza infection. Microbiome 8:1–12.

5. Fujiogi M, Raita Y, Pérez-Losada M, Freishtat RJ, Celedón JC, Mansbach JM, et al. 2022. Integrated relationship of nasopharyngeal airway host response and microbiome associates with bronchiolitis severity. Nat Commun 13:4970.

6. Lloyd-Price J, Arze C, Ananthakrishnan AN, Schirmer M, Avila-Pacheco J, Poon TW, et al. 2019. Multi-omics of the gut microbial ecosystem in inflammatory bowel diseases. Nature 569:655–62.

7. Priya S, Burns MB, Ward T, Mars RAT, Adamowicz B, Lock EF, et al. 2022. Identification of shared and disease-specific host gene–microbiome associations across human diseases using multi-omic integration. Nat Microbiol 7:780–95.

8. Cremer J, Arnoldini M, Hwa T. 2017. Effect of water flow and chemical environment on microbiota growth and composition in the human colon. Proc Natl Acad Sci 114:6438-43.

9. Dupont HL, Jiang ZD, Dupont AW, Utay NS. 2020. The intestinal microbiome in human
health and disease. Trans Am Clin Climatol Assoc 131:178.

10. Hu S, Wang Y, Lichtenstein L, Tao Y, Musch MW, et al. 2010. Regional differences in
colonic mucosa-associated microbiota determine the physiological expression of host
heat shock proteins. Am J Physiol Gastrointest Liver Physiol 299:G1266-75.

11. Yasuda K, Oh K, Ren B, Tickle TL, Franzosa EA, Wachtman LM, et al. 2015.
Biogeography of the intestinal mucosal and lumenal microbiome in the rhesus macaque.
Cell Host Microbe 17:385–91.

12. Kozik AJ, Nakatsu CH, Chun H, Jones-Hall YL. 2019. Comparison of the fecal, cecal,
and mucus microbiome in male and female mice after TNBS-induced colitis. PLoS One
14:1–14.

13. Uehara M, Inoue T, Kominato M, Hase S, Sasaki E, Toyoda A, et al. 2022. Intraintestinal
Analysis of the Functional Activity of Microbiomes and Its Application to the Common
Marmoset Intestine. mSystems 7:e00520-22.

14. Newman ME. 2004. Detecting community structure in networks. Eur Phys J B 38:321–
30.

15. Blondel VD, Guillaume JL, Lambiotte R, Lefebvre E. 2008. Fast unfolding of
communities in large networks. J Stat Mech 10008.

16. Traag VA, Waltman L, Van Eck NJ. 2019. From Louvain to Leiden: guaranteeing well-
connected communities. Sci Rep 9:5233.

17. Langfelder P, Horvath S. 2008. WGCNA: an R package for weighted correlation network
analysis. BMC bioinformatics 9:1-3.

18. Hugenholtz F, de Vos WM. 2018. Mouse models for human intestinal microbiota

research: a critical evaluation. Cell Mol Life Sci 75:149–60.

19. Kishi N, Sato K, Sasaki E, Okano H. 2014. Common marmoset as a new model animal

for neuroscience research and genome editing technology. Dev Growth Differ 56:53–62.

20. Inoue T, Okahara N, Okahashi N, Sato K, Ueda M, Atarashi K, et al. 2021. Generation of

germ-free common marmosets. Research Square. Preprint.

https://assets.researchsquare.com/files/rs-428622/v1_covered.pdf?c=1631871455

21. Stuart JM, Segal E, Koller D, Kim SK. 2003. A gene-coexpression network for global

discovery of conserved genetic modules. Science 302:249–255.

22. Wang K, Wei G, Liu D. 2012. CD19: a biomarker for B cell development, lymphoma

diagnosis and therapy. Exp Hematol Oncol 1:1–7.

23. Huse K, Bai B, Hilden VI, Bollum LK, Våtsveen TK, Munthe LA, et al. 2022.

Mechanism of CD79A and CD79B support for IgM+ B cell fitness through B cell

receptor surface expression. J Immunol 209:2042–53.

24. Clark EA, Giltiay N V. 2018. CD22: a regulator of innate and adaptive B cell responses

and autoimmunity. Front Immunol 9:2235.

25. Pan C, Baumgarth N, Parnes JR. 1999. CD72-deficient mice reveal nonredundant roles of

CD72 in B cell development and activation. Immunity 11:495–506.

26. Alegre ML, Frauwirth KA, Thompson CB. 2001. T-cell regulation by CD28 and CTLA-

4. Nat Rev Immunol 1:220–8.

27. Yang Y, Zang Y, Zheng C, Li Z, Gu X, Zhou M, et al. 2020. CD3D is associated with

immune checkpoints and predicts favorable clinical outcome in colon cancer.

Immunotherapy 12:25–35.

28. Hubbard N, Hagin D, Sommer K, Song Y, Khan I, Clough C, et al. 2016. Targeted gene

editing restores regulated CD40L function in X-linked hyper-IgM syndrome. Blood

127:2513–22.

29. Sobhani N, Tardiel-Cyril DR, Davtyan A, Generali D, Roudi R, Li Y. 2021. CTLA-4 in

regulatory T cells for cancer immunotherapy. Cancers 13:1–18.

30. He Y, Hua R, Li B, Gu H, Sun Y, Li Z. 2021. Loss of FBP1 promotes proliferation,

migration, and invasion by regulating fatty acid metabolism in esophageal squamous cell

carcinoma. Aging 13:4986–8.

31. Hirata H, Sugimachi K, Komatsu H, Ueda M, Masuda T, Uchi R, et al. 2016. Decreased

expression of fructose-1,6-bisphosphatase associates with glucose metabolism and tumor

progression in hepatocellular carcinoma. Cancer Res 76:3265–76.

32. Dong Y, Huaying S, Danying W, Chihong Z, Ruibin J, Xiaojiang S, et al. 2018.

Significance of methylation of FBP1 gene in non-small cell lung cancer. Biomed Res Int

2018:13–9.

33. Li H, Qi Z, Niu Y, Yang Y, Li M, Pang Y, et al. 2021. FBP1 regulates proliferation,

metastasis, and chemoresistance by participating in C-MYC/STAT3 signaling axis in

ovarian cancer. Oncogene 40:5938–49.

34. Hou Y, Wei W, Guan X, Liu Y, Bian G, He D, et al. 2021. A diet-microbial metabolism

feedforward loop modulates intestinal stem cell renewal in the stressed gut. Nat Commun

12:271.

35. Mardinoglu A, Shoaie S, Bergentall M, Ghaffari P, Zhang C, Larsson E, et al. 2015. The

gut microbiota modulates host amino acid and glutathione metabolism in mice. Mol Syst

Biol 11:1–15.

36. Medina DA, Pinto F, Ovalle A, Thomson P, Garrido D. 2017. Prebiotics mediate

microbial interactions in a consortium of the infant gut microbiome. Int J Mol Sci

18:2095.

37. Cheng J, Hu J, Geng F, Nie S. 2022. Bacteroides utilization for dietary polysaccharides

and their beneficial effects on gut health. Food Sci Hum Wellness 11:1101–10.

38. Rosser EC, Piper CJM, Matei DE, Blair PA, Rendeiro AF, Orford M, et al. 2020.

Microbiota-derived metabolites suppress arthritis by amplifying aryl-hydrocarbon

receptor activation in regulatory B cells. Cell Metab 31:837-851.

39. Ding X, Bin P, Wu W, Chang Y, Zhu G. 2020. Tryptophan metabolism, regulatory T cells,

and inflammatory bowel disease: a mini review. Mediators Inflamm 2020.

40. Ezeji JC, Sarikonda DK, Hopperton A, Erkkila HL, Cohen DE, Martinez SP, et al. 2021.

Parabacteroides distasonis: intriguing aerotolerant gut anaerobe with emerging

antimicrobial resistance and pathogenic and probiotic roles in human health. Gut

Microbes 13:1–27.

41. Cekanaviciute E, Yoo BB, Runia TF, Debelius JW, Singh S, Nelson CA, et al. 2017. Gut

bacteria from multiple sclerosis patients modulate human T cells and exacerbate

symptoms in mouse models. Proc Natl Acad Sci USA 114:10713–8.

42. Łaniewski P, Herbst-Kralovetz MM. 2021. Bacterial vaginosis and health-associated

bacteria modulate the immunometabolic landscape in 3D model of human cervix. NPJ

Biofilms Microbiomes 7:1–17.

43. Tomkovich S, Gharaibeh RZ, Dejea CM, Pope JL, Jiang J, Winglee K, et al. 2020.

Human colon mucosal biofilms and murine host communicate via altered mRNA and

microRNA expression during cancer. mSystems 5:e00451-19.

44. Cornick S, Tawiah A, Chadee K. 2015. Roles and regulation of the mucus barrier in the

gut. Tissue Barriers 3:1–2.

45. González-Morelo KJ, Vega-Sagardía M, Garrido D. Molecular insights into O-linked

glycan utilization by gut microbes. Frontiers in Microbiology. 2020 Nov 5;11:591568.

46. Png CW, Lindén SK, Gilshenan KS, Zoetendal EG, McSweeney CS, Sly LI, McGuckin

MA, Florin TH. Mucolytic bacteria with increased prevalence in IBD mucosa augmentin

vitroutilization of mucin by other bacteria. Official journal of the American College of

Gastroenterology| ACG. 2010 Nov 1;105(11):2420-8.

47. Zurkinden L, Sviridov D, Vogt B, Escher G. 2020. Downregulation of Cyp7a1 by cholic

acid and chenodeoxycholic acid in Cyp27a1/ApoE double knockout mice: differential

cardiovascular outcome. Front Endocrinol 11:1–13.

48. Biagioli M, Marchianò S, Carino A, Di Giorgio C, Santucci L, Distrutti E, et al. 2021.

Bile acids activated receptors in inflammatory bowel disease. Cells 10:1281.

49. Badi SA, Motahhary A, Bahramali G, Masoumi M, Khalili SFS, Ebrahimzadeh N, et al.

2020. The regulation of Niemann-Pick C1-Like 1 (NPC1L1) gene expression in opposite

direction by Bacteroides spp. and related outer membrane vesicles in Caco-2 cell line. J

Diabetes Metab Disord 19:415–22.

50. Wang J, Thingholm LB, Skiecevičie J, Rausch P, Kummen M, Hov JR, et al. 2016.

Genome-wide association analysis identifies variation in Vitamin D receptor and other

host factors influencing the gut microbiota. Nat Genet 48:1396–406.

51. Qin Y, Havulinna AS, Liu Y, Jousilahti P, Ritchie SC, Tokolyi A, et al. 2022. Combined

effects of host genetics and diet on human gut microbiota and incident disease in a single

population cohort. Nat Genet 54:134–42.

52. Zhang Q, Linke V, Overmyer KA, Traeger LL, Kasahara K, Miller IJ, et al. 2023. Genetic

mapping of microbial and host traits reveals production of immunomodulatory lipids by

Akkermansia muciniphila in the murine gut. Nat Microbiol 8:424–40.

53. Zeevi D, Korem T, Godneva A, Bar N, Kurilshikov A, Lotan-Pompan M, et al. 2019.

Structural variation in the gut microbiome associates with host health. Nature 568:43–8.

54. Wang Y, Baumdicker F, Schweiger P, Kuenzel S, Staubach F. 2021. Horizontal gene

transfer-mediated bacterial strain variation affects host fitness in Drosophila. BMC Biol

19:1-12.

55. Han B, Sivaramakrishnan P, Lin CCJ, Neve IAA, He J, Tay LWR, et al. 2017. Microbial

genetic composition tunes host longevity. Cell 169:1249-1262.

56. Zinnen AD, Vichich J, Metzger JM, Gambardella JC, Bondarenko V, Simmons HA, et al.

2022. Alpha-synuclein and tau are abundantly expressed in the ENS of the human

appendix and monkey cecum. PLoS One 17:e0269190.

57. Fisher RE. The primate appendix: A reassessment. Vol. 261, Anatomical Record. 2000. pp

228–36.

58. Debelius JW, Engstrand L, Matussek A, Brusselaers N, Morton JT, Stenmarker M,et al.

2023. The local tumor microbiome is associated with survival in late-stage colorectal

cancer patients. Microbiol Spectr. 2023 11:e05066-22.

59. Drewes JL, White JR, Dejea CM, Fathi P, Iyadorai T, Vadivelu J, et al. 2017. High-

resolution bacterial 16S rRNA gene profile meta-analysis and biofilm status reveal

common colorectal cancer consortia. NPJ Biofilms Microbiomes 3:34.

60. Čipčić Paljetak H, Barešić A, Panek M, Perić M, Matijašić M, Lojkić I, et al. 2022. Gut

microbiota in mucosa and feces of newly diagnosed, treatment-naïve adult inflammatory

bowel disease and irritable bowel syndrome patients. Gut Microbes 14:2083419.

61. Yu LC. 2018. Microbiota dysbiosis and barrier dysfunction in inflammatory bowel

disease and colorectal cancers: exploring a common ground hypothesis. J Biomed Sci.

25:1-4.

Figures

Fig. 1 Overview of the main analytical approach used in this study

This method involves the simultaneous acquisition of the host and microbiome transcriptome (metatranscriptome) from intestinal sites and conducts a gene co-expression network analysis of the host and microbiome community. The procedure is outlined from step A to E. (A) Perform dual RNA-seq of host and microbiome transcriptomes in internal site samples. (B) Separate host and microbiome RNA reads in silico using the host reference genome. (C) Compute the gene expression profiles of the host and microbiome, respectively. (D) Unify gene co-expression networks between host-host, host-microbiome, and microbiome-microbiome. The host-host and microbiome-microbiome gene co-expression networks are connected when there is a positive correlation between genes, while the host-microbiome gene co-expression network is connected when there is either a positive or negative correlation between genes. (E) Identify closely interacting gene modules from the gene co-expression network using a proposed clique-based algorithm. (E Step 1) This process begins with searching for cliques. (E Step 2) If overlaps exist between cliques, the cliqueness score is computed in three gene co-expression networks (host-host, host-microbiome, and microbiome-microbiome) when overlapping cliques are merged. (E Step 3) If the cliqueness score exceeds the merging threshold τ in all three networks, the cliques are merged into a single module. If the score does not exceed the merging threshold τ in at least one network, the original cliques are retained without merging. This process is repeated until no further modules can be merged.

Fig. 2 PCA analysis of gene expression data for host and microbiome

PCA was applied to the host gene expression profiles to visualize groupings according to site (A)

and individual (B). PCA was similarly applied to the microbiome gene expression profiles to

visualize groupings according to site (C) and individual (D).

Fig. 3 Variations in host gene expression and functional pathways among intestinal sites

(A) The top 20 functional pathways significantly upregulated in the cecum compared to the

transverse colon (normalized enrichment score > 1.5, FDR < 0.1), as revealed by gene set

enrichment analysis, are presented alongside their normalized enrichment score and FDR. (B)

Heatmap of differentially expressed genes belonging to the top 5 enriched functional pathways.

The heatmap on the left represents the z-scores of gene expression levels, with genes

significantly upregulated in the cecum compared to the transverse colon and rectum (FDR $<$

0.01) colored in pink on the middle heatmap. The heatmap on the right indicates the functional

pathways to which the genes belong, represented in green.

Fig. 4 Functional pathways characterized in modules extracted from the host-microbiome gene

co-expression network

This figure displays the identified modules along with the results of their functional pathway

enrichment analysis (Supplementary Section 8). To enhance clarity, the background of the bar

graphs alternates between gray and white for each module. Modules were characterized using

KEGG pathways, with those meeting an FDR < 0.05 as determined by Fisher's exact test. The

dashed line represents an FDR < 0.05 ($-\log 10(\text{FDR}) > 1.30$). All pathways with an FDR < 0.1

are represented, and modules with no identified function are denoted as "None".

Fig. 5 Metabolic pathway-enriched Module_07

A module showing a positive correlation between the host gene *FBP1*, which is highly expressed

in the transverse colon, and 40 microbiome genes, including those involved in fructose-mannose

metabolism and oxidative phosphorylation, was extracted. These genes were visualized

alongside bacterial species nodes that contribute to the expression of the microbiome genes. Note

that the bacterial species nodes were added after the module detection analysis. Edges among

genes indicate an absolute Spearman's correlation coefficient greater than 0.8. Edges connecting

microbiome genes to bacterial species are shown when the contribution rate of bacterial species

to microbiome gene expression exceeds 0.2. The thickness of the edge increases with higher

contribution rates.

Fig. 6 B cell receptor signaling pathway-enriched Module_01

Host genes involved in the immune response of B cells, including *CD19*, *CD22*, *CD79B* and

PTPN6, which were highly expressed in the cecum, co-expressed with the K01696 (*trpB*). These

genes were visualized alongside bacterial species nodes that contribute to the expression of the

microbiome genes. K01696 (*trpB*) expression was contributed to by *Parabacteroides distasonis*.

Edges among genes indicate an absolute Spearman's correlation coefficient greater than 0.8.

Edges connecting microbiome genes to bacterial species are shown when the contribution rate of

bacterial species to microbiome gene expression exceeds 0.2. Note that the bacterial species

nodes were added after the module detection analysis. The thickness of the edge increases with

higher contribution rates.

Fig. 7 Carbon metabolism pathway-enriched Module_21

Host genes, including *MMP10*, which is specifically highly expressed in the cecum, and 40

microbiome genes involved in carbon metabolism, were identified. These genes were visualized

alongside bacterial species nodes that contribute to the expression of the microbiome genes. Note

that the bacterial species nodes were added after the module detection analysis. Edges among

genes indicate an absolute Spearman's correlation coefficient greater than 0.8. Edges connecting

microbiome genes to bacterial species are shown when the contribution rate of bacterial species

to microbiome gene expression exceeds 0.2. The thickness of the edge increases with higher

contribution rates.

Fig. 8 Module_18 containing host genes and microbiome genes related to the production and

breakdown of bile acids

These genes were visualized alongside bacterial species nodes that contribute to the expression

of the microbiome genes. Note that the bacterial species nodes were added after the module

detection analysis. Edges among genes indicate an absolute Spearman's correlation coefficient

greater than 0.8. Edges connecting microbiome genes to bacterial species are shown when the

contribution rate of bacterial species to microbiome gene expression exceeds 0.2. The thickness

of the edge increases with higher contribution rates.

Supplemental Materials

Supplementary Notes. Supplementary information on the methods, consisting of: Section 1.

Animal experiments; Section 2. RNA extraction and dual RNA sequencing of host and

microbiome; Section 3. Partitioning of host RNA reads and microbiome RNA reads; Section 4.

Accuracy assessment of partitioning of RNA read data into host and microbiome reads; Section

5. Analysis of host gene expression; Section 6. Analysis of microbiome gene expression; Section

7. Network construction and determination of parameters for module extraction; Section 8.

Characterization and evaluation of identified modules; Section 9. Comparison with other module

extraction methods

Table S1. RIN score of total RNA samples

Table S2. Sequence statistics

Table S3. Number of RNA reads derived from the host and microbiome

Table S4. Results of DEG analysis with pairwise DEG analysis between three intestinal sites

Table S5. Statistics on the module extraction methods

Table S6. Evaluation of the module extraction methods through enrichment analysis

Table S7. Information on common marmoset samples

Table S8. Number of RNA reads at each preprocessing step

Fig. S1. Carbon metabolism-related microbiome genes detected in Module_21

The Carbon metabolism pathway of KEGG is shown with the genes in Module_21 highlighted in

red.

Fig. S2. Changes in the coefficient of determination for fitting to the scale-free topology model

in response to the soft threshold in WGCNA

The red line indicates a coefficient of determination = 0.8.

Fig. 1 Overview of the main analytical approach used in this study

This method involves the simultaneous acquisition of the host and microbiome transcriptome (metatranscriptome) from intestinal sites and conducts a gene co-expression network analysis of the host and microbiome community. The procedure is outlined from step A to E. (A) Perform dual RNA-seq of host and microbiome transcriptomes in internal site samples. (B) Separate host and microbiome RNA reads in silico using the host reference genome. (C) Compute the gene expression profiles of the host and microbiome, respectively. (D) Unify gene co-expression networks between host-host, host-microbiome, and microbiome-microbiome. The host-host and microbiome-microbiome gene co-expression networks are connected when there is a positive correlation between genes, while the host-microbiome gene co-expression network is connected when there is either a positive or negative correlation between genes. (E) Identify closely interacting gene modules from the gene co-expression network using a proposed clique-based algorithm. (E Step 1) This process begins with searching for cliques. (E Step 2) If overlaps exist between cliques, the cliqueness score is computed in three gene co-expression networks (host-host, host-microbiome, and microbiome-microbiome) when overlapping cliques are merged. (E Step 3) If the cliqueness score exceeds the merging threshold τ in all three networks, the cliques are merged into a single module. If the score does not exceed the merging threshold τ in at least one network, the original cliques are retained without merging. This process is repeated until no further modules can be merged.

Fig. 2 PCA analysis of gene expression data for host and microbiome

PCA was applied to the host gene expression profiles to visualize groupings according to site (A) and individual (B). PCA was similarly applied to the microbiome gene expression profiles to visualize groupings according to site (C) and individual (D).

Fig. 3 Variations in host gene expression and functional pathways among intestinal sites

(A) The top 20 functional pathways significantly upregulated in the cecum compared to the transverse colon (normalized enrichment score > 1.5, FDR < 0.1), as revealed by gene set enrichment analysis, are presented alongside their normalized enrichment score and FDR. (B) Heatmap of differentially expressed genes belonging to the top 5 enriched functional pathways. The heatmap on the left represents the z-scores of gene expression levels, with genes significantly upregulated in the cecum compared to the transverse colon and rectum (FDR < 0.01) colored in pink on the middle heatmap. The heatmap on the right indicates the functional pathways to which the genes belong, represented in green.

Fig. 4 Functional pathways characterized in modules extracted from the host-microbiome gene co-expression network

This figure displays the identified modules along with the results of their functional pathway enrichment analysis (Supplementary Section 8). To enhance clarity, the background of the bar graphs alternates between gray and white for each module. Modules were characterized using KEGG pathways, with those meeting an FDR $<$ 0.05 as determined by Fisher's exact test. The dashed line represents an FDR $<$ 0.05 (-log10(FDR) $>$ 1.30). All pathways with an FDR $<$ 0.1 are represented, and modules with no identified function are denoted as "None".

Fig. 5 Metabolic pathway-enriched Module_07

A module showing a positive correlation between the host gene *FBP1*, which is highly expressed in the transverse colon, and 40 microbiome genes, including those involved in fructose-mannose metabolism and oxidative phosphorylation, was extracted. These genes were visualized alongside bacterial species nodes that contribute to the expression of the microbiome genes. Note that the bacterial species nodes were added after the module detection analysis. Edges among genes indicate an absolute Spearman's correlation coefficient greater than 0.8. Edges connecting microbiome genes to bacterial species are shown when the contribution rate of bacterial species to microbiome gene expression exceeds 0.2. The thickness of the edge increases with higher contribution rates.

Fig. 6 B cell receptor signaling pathway-enriched Module_01

Host genes involved in the immune response of B cells, including *CD19*, *CD22*, *CD79B* and *PTPN6*, which were highly expressed in the cecum, co-expressed with the K01696 (*trpB*). These genes were visualized alongside bacterial species nodes that contribute to the expression of the microbiome genes. K01696 (*trpB*) expression was contributed to by *Parabacteroides distasonis*. Edges among genes indicate an absolute Spearman's correlation coefficient greater than 0.8. Edges connecting microbiome genes to bacterial species are shown when the contribution rate of bacterial species to microbiome gene expression exceeds 0.2. Note that the bacterial species nodes were added after the module detection analysis. The thickness of the edge increases with higher contribution rates.

Fig. 7 Carbon metabolism pathway-enriched Module_21

Host genes, including *MMP10*, which is specifically highly expressed in the cecum, and 40 microbiome genes involved in carbon metabolism, were identified. These genes were visualized alongside bacterial species nodes that contribute to the expression of the microbiome genes. Note that the bacterial species nodes were added after the module detection analysis. Edges among genes indicate an absolute Spearman's correlation coefficient greater than 0.8. Edges connecting microbiome genes to bacterial species are shown when the contribution rate of bacterial species to microbiome gene expression exceeds 0.2. The thickness of the edge increases with higher contribution rates.

Fig. 8 Module_18 containing host genes and microbiome genes related to the production and breakdown of bile acids

These genes were visualized alongside bacterial species nodes that contribute to the expression of the microbiome genes. Note that the bacterial species nodes were added after the module detection analysis. Edges among genes indicate an absolute Spearman's correlation coefficient greater than 0.8. Edges connecting microbiome genes to bacterial species are shown when the contribution rate of bacterial species to microbiome gene expression exceeds 0.2. The thickness of the edge increases with higher contribution rates.