Decoding host-microbiome interactions through co-expression network analysis within the

non-human primate intestine

Mika Uehara[1], Takashi Inoue[2], Sumitaka Hase[1], Erika Sasaki[2, 4], Atsushi Toyoda[3], Yasubumi

Sakakibara[1*]

1) Department of Biosciences and Informatics, Keio University, Yokohama, Kanagawa 223-

8522, Japan

2) Department of Marmoset Biology and Medicine, Central Institute for Experimental Animals,

Kawasaki, Kanagawa 210-0821, Japan

3) Department of Genomics and Evolutionary Biology, National Institute of Genetics, Mishima,

Shizuoka 411-8540, Japan

4) Laboratory for Marmoset Neural Architecture, RIKEN Center for Brain Science, 2-1

Hirosawa, Wako-shi, Saitama 351-0198, Japan.

*Corresponding Author

Yasubumi Sakakibara

3-14-1 Hiyoshi, Kohoku-ku, Yokohama, 223-8522, Japan

Phone/Fax: +81-45-566-1791

E-mail: yasu@bio.keio.ac.jp

The authors declare no competing interests.

ABSTRACT

The gut microbiome affects the health status of the host through complex interactions with the

host's intestinal wall. These host-microbiome interactions spatially vary along the physical and

chemical environment of the intestine, but these changes remain unknown. This study

investigated these intricate relationships through a gene co-expression network analysis based on

dual transcriptome profiling of different intestinal sites (cecum, transverse colon, rectum) of the

primate common marmoset. We proposed a gene module extraction algorithm based on graph

theory to find tightly interacting gene modules of the host and the microbiome from a vast co-

expression network. The 27 gene modules identified by this method, which include both host and

microbiome genes, not only produced results consistent with previous studies regarding the host-

microbiome relationships, but also provided new insights into microbiome genes acting as

potential mediators in host-microbiome interplays. Specifically, we discovered associations

between the host gene FBP1, a cancer marker, and polysaccharide degradation-related genes

(*pfkA*, *fucI*) coded by *Bacteroides vulgatus*, as well as relationships between host genes CD19,

CD22, CD79B, PTPN6 involved in B-cell immune responses and a tryptophan synthesis gene

(*trpB*) coded by *Parabacteroides distasonis*. Furthermore, our proposed module extraction

algorithm surpassed existing approaches by successfully defining more functionally related gene

modules, providing insights for understanding the complex relationship between the host and the

microbiome.

INTRODUCTION

The mutual interaction between the intestinal microbiome and the host constitutes a complex

biological process, with intertwined dependencies essential for survival and health of both. The

interactions between the intestinal microbiome and the host are formed by various factors such as

the effects of metabolites produced by bacteria on the host, the binding of bacterial-derived

proteins and host receptors, and the binding of bacterial cell surface structures and host receptors.

Studies on host-microbiome interaction through bacterial-derived proteins have discovered that a

novel protein P9 secreted by *Akkermansia muciniphila* binds to host ICAM-2, thereby increasing

the secretion of glucagon-like peptide 1 (GLP-1), suggesting that the interaction between P9 and

ICAM-2 could be a potential therapeutic target in type 2 diabetes patients [1]. It has also been

reported that bacterial lipopolysaccharides (LPS), cell wall components of gram-negative

bacteria, are structurally recognized by the Toll-like receptor 4 (TLR4) complex, inducing innate

immune responses [2]. Among these, the host-microbiome interaction mediated by metabolites is

particularly well-studied. For instance, primary bile acids, produced in the host's liver cells, are

converted into secondary bile acids by the microbiome in the colon. This microbiome-derived

secondary bile acid is known to regulate the homeostasis of regulatory T cells, immune

suppressive cells in the host [3]. In this way, the intestinal microbiome and host mutually interact

and establish a symbiotic relationship while forming a metabolic pathway or physiological

response.

To understand such complex interactions between the microbiome and the host, it is

beneficial to analyze combined profiles of both the microbiome and the host. A cohort study of

patients with colorectal cancer, irritable bowel syndrome, and inflammatory bowel disease

investigated host gene expression levels and microbiota composition [4]. While the relationship

between disease and microbiome, or disease and host genes, has been investigated individually,

these approach combining profiles of both the microbiome and the host have been a significant

clue to unravel the relationship among the microbiome, host genes, and diseases.

Recently, by profiling the gene expression of both the host and the microbiome, the functional

involvement of the microbiome in the host response has been revealed, extending beyond just the

relationship between microbiome members and the host [5, 6, 7]. Gene expression network

analysis is suitable for comprehensively deciphering the complex interactions between the host's

genes and the diverse microbiome's genes. Some studies have adopted network analysis to

highlight the characteristics of host-microbiome interactions, identifying functional gene

modules related to various physiological and pathological processes in both the host and

microbiome. For instance, a study [5]characterizing the host-microbiome interaction during

influenza virus infection identified modules from the host's gene co-expression network. It

discovered correlations between host gene modules involved in interferon signal transduction

and the sulfonamide antibiotic resistance genes of the microbiome. In another study [6] applying

network analysis to the gene expression profiles of the host and microbiome in the

nasopharyngeal airway during bronchiolitis, networks were built for both the host and

microbiome, then modules were identified, revealing host modules involved in host-T-cell

regulation and microbiome modules involved in microbial-BCAA metabolism and oxidative

stress response were associated with disease severity. Thus, considering network topology, such

as modularity, helps interpret mutual biological processes between the microbiome and host from

complex and large networks.

Despite such progress, there remain several limitations to the understanding of the

interactions between the host and the intestinal microbiome. First, there's a lack of understanding

about the spatial variation in host-microbiome interactions. As the function of the intestinal tract

varies by location and the physicochemical environment such as nutrients, oxygen, and pH also

differ [8, 9], the mechanisms of interaction between host gut cells and the microbiome could vary

as well. Indeed, studies using animal models have reported that the composition and functional

fluctuation of the microbiome varies due to environmental changes depending on intestinal

location (biogeographical location) [10, 11, 12, 13]. However, these studies focused solely on the

microbiome, and the coordinated spatial variation of both the microbiome and host gut cells

along the intestinal tract has yet to be clarified.

Second, there's a lack of computational methods for extracting significant modules

(subgraphs) from the host-microbiome gene expression network. Analyzing a massive gene

expression network integrating the gene expression profiles of both the host and microbiome

helps interpret the interactions between the genes of the host and the microbiome. Especially in a

huge gene expression network, it is crucial to extract densely connected subgraphs (modules) to

divide the network into understandable units; however, existing algorithms for extracting

modules from the network are insufficient for accurately capturing the interaction between the

host and microbiome because they were not designed with the purpose of understanding

interactions between organisms [14, 15].

Lastly, there are concerns regarding the applicability of the results of studies using animal

models to humans. Mouse models have been the basis of biological research for a long time due

to ethical constraints, experiments in controlled environments, and considerations of

reproducibility, but there are several issues and limitations to their use. The physiology, anatomy,

and pharmacology of mice differ from those of humans, and there are differences in the

composition and function of the intestinal microbiome between mice and humans [13, 16].

Particularly considering the complexity of host-microbiome interactions, it is challenging to

precisely replicate the relationship between humans and their intestinal microbiome. In contrast,

the common marmoset, a small New World primate, is considered a useful model in preclinical

trials because it is genetically close to humans and has common physiological and anatomical

characteristics [17]. Moreover, the common marmoset is the only non-human primate in which a

germ-free state has been successfully achieved [18], and it has a fecal microbiome similar to that

of humans compared to other animal models, including mice [13], potentially broadening the

scope of intestinal microbiome research.

To address these challenges, this study aimed to reveal the interactions between the host

and the microbiome along multiple sites of the common marmoset colon (cecum, transverse

colon, rectum) by developing a module extraction method applicable to the gene expression

network that represents the complex interactions between the host and the microbiome. Through

simultaneous profiling of host and microbiome transcripts in the intestinal sites of the common

marmoset, we constructed an integrated gene expression network, incorporating both host and

microbiome genes. From this network, we further identified densely covariant subgraphs (gene

modules) to interpret the interactions between genes. As conventional methods for extracting

gene modules from the network were insufficient for interpreting the interactions between

organisms examined in this study, we proposed a module extraction algorithm based on graph

theory. Our proposed method successfully extracted more modules containing functionally

related genes of both the host and microbiome compared to existing algorithms, underlining its

suitability for predicting host-microbiome interactions. This method extracted 27 gene modules

containing both host and microbiome, indicating that the gut microbiome is functionally closely

involved in the variation of host gene expression along the intestinal tract.

RESULTS

Overview of the analysis pipeline integrating host and microbiome transcriptomes:

Identification of closely interacting gene modules from gene expression networks

To investigate the host and microbiome interactions along the intestinal tract, we proposed an

analysis method integrating the host transcriptome and the microbiome metatranscriptome

(Figure 1). First, we computed gene expression profiles from the host transcriptome and

microbiome metatranscriptome obtained via dual RNA-seq and created three types of gene

expression networks: host-host, host-microbiome, and microbiome-microbiome. By unifying

these three networks, we constructed a host and microbiome gene co-expression network, with

the genes of the host and microbiome serving as nodes, and covariant genes interconnected.

Second, we unified gene modules containing closely interacting genes from both the host and

microbiome from the gene co-expression network. Conventional algorithms for extracting

subgraphs from the network, such as modularity maximization, may not be suitable for

accurately capturing interactions between the host and microbiome, as they were not designed

with the purpose of understanding inter-organism interactions. Therefore, we proposed a clique-

search-based module extraction method that takes into account the structure of three networks:

host-host, host-microbiome, and microbiome-microbiome. This method begins by exploring

cliques from the gene co-expression network. For overlapping cliques, it calculates the

cliqueness in each of the original three networks and merges them if the cliqueness of all

networks exceeds a certain threshold, thereby obtaining the final gene modules (refer to the

Methods section for details).

Computing gene expression profiles of the host and the microbiome from dual RNA-seq data

Firstly, we extracted high-quality RNA (RNA integrity number (RIN) scores ≥8.6;

Supplementary Table S1) from the intestinal sites (cecum, transverse colon, rectum) of five

common marmosets and carried out dual RNA-seq with NovaSeq 6000 v1.5 (100 bp PE). The

obtained RNA read data amounted to a total of 4,724 M reads (mean 157,450,033 paired-end

reads/sample; Supplementary Table S2).

Next, as the RNA read data obtained by dual RNA-seq contains a mix of reads derived

from both the host and the microbiome, it is necessary to separate these two read data in silico

(Supplementary Table S3). The pipeline established in this study separates the host-derived read

data and the microbiome-derived read data by aligning the RNA reads to the host reference

genome (Fig. 1B). To avoid cross-contamination of read data between organisms at this stage, we

adjusted the pipeline parameters using simulated RNA read data from the host and the

microbiome. As a result, we achieved a precision of 1.000 in separating the read data of the host

and the microbiome, and our established pipeline completely separated the reads derived from

the host and those from the microbiome (Supplementary Table S4, Supplementary Note Section

3).

After the above preprocessing, we calculated the gene expression levels for both the host

and the microbiome. As a result, we detected 16,578 host expressed genes and 8,533 microbiome

expressed genes in total (Supplementary Table S5 and S6, Supplementary Note Section 4 and 5).

To examine the expression patterns between different sites and between different individuals for

both host genes and microbiome genes, we applied Principal Component Analysis (PCA) to the

gene expression profiles for visualization. In the host profiles, distinct clusters were formed for

each site, while the clusters for each individual overlapped (Fig. 2A and B). In the microbiome

profiles, clusters were separated for the cecum and rectum, which are the beginning and end of

the colon, respectively, with the transverse colon positioned between the cecum and rectum

overlapping between the two clusters (Fig. 2C). The clusters for each individual also overlapped

in the microbiome profiles (Fig. 2D). These results suggest that both host and microbiome gene

expression profiles are more variable across different sites compared to those across individuals.

Spatial variations in host gene expression among intestinal sites

Of the 16,578 host genes detected in three regions of the common marmoset's intestinal tract

(cecum, transverse colon, and rectum), a total of 2,615 genes showed significant expression

variations among the sites (FDR<0.01). Furthermore, upon conducting functional pathway

analysis, 40 pathways were identified, all of which were significantly upregulated in the cecum

compared to the transverse colon (normalized enrichment score >1.5, FDR<0.1; Supplementary

Table S7). In particular, pathways with high normalized enrichment scores, such as Fc epsilon RI

signaling pathway, T cell receptor signaling pathway, leukocyte transendothelial migration,

Yersinia infection, and B cell receptor signaling pathway were detected, indicating that pathways

related to immune functions were upregulated in the cecum relative to the transverse colon (Fig.

3A). Additionally, a substantial number of genes involved in these functional pathways were

found to be significantly upregulated in the cecum (FDR<0.01). For example, among the genes

specifically upregulated in the cecum, we found genes related to B cell receptor signaling. These

include CD19, which acts as a coreceptor for the B cell antigen receptor complex and augments

B cell activation [19]; CD79A and CD79B, components of the B cell antigen receptor that

generate signals upon antigen recognition [20]; and CD22 and CD72, which inhibit B cell

antigen receptor signal transduction [21, 22]. In addition, genes associated with T cell receptor

signaling were also found as being highly expressed specifically in the cecum. These include

CD28, a co-stimulatory receptor for T cell receptors that activates T cells [23]; CD3D, a

component of CD3 that plays a role in signal transduction during T cell activation [24];

CD40LG, which is induced on the T cell surface following T cell receptor (TCR) activation [25];

and CTLA4, which attenuates T cell activation and fine-tunes immune responses [26] (Fig. 3B).

Interpreting host-microbiome interactions through spatial co-expression gene network

analysis within the intestine

We constructed the unified gene expression network composed of 3,219 gene nodes, which

embodies co-expression information on host-host, host-microbiome, and microbiome-

microbiome interactions. Subsequently, to identify densely co-expressed gene sets within this

network, we implemented the clique-based module extraction algorithm proposed in this study.

In graph theory, a "clique" refers to a subgraph within the network where all nodes are fully

connected. Our algorithm discerned modules of interacting genes in the gene co-expression

network as aggregates of overlapping cliques. As a result, we then identified 27 gene modules

containing both host and microbiome genes (Supplementary Table S8). Furthermore, upon

conducting a functional pathway enrichment analysis on these gene modules, we found that 19 of

them, representing 70.4%, could be functionally characterized (Supplementary Note Section 6).

The modules we identified were characterized by functional pathways including the B

cell receptor signaling pathway, fructose and mannose metabolism, oxidative phosphorylation,

and biosynthesis of amino acids (Fig. 4). In addition, we provided visualizations of the bacterial

species that contribute to microbiome gene expression within these identified modules, together

with host and microbiome genes (Fig.5, 6, 7 and S2). The results obtained from these identified

modules were not only in agreement with previous research examining host genes, microbiome

genes, and bacterial species but also revealed novel interactions between host genes and the

microbiome, including the identification of microbiome genes that serve as intermediaries

between host genes and the microbiome.

Module_06, as shown in Figure 5, was identified as a module with a positive correlation

between the host gene Fructose-1,6-bisphosphatase 1 (FBP1), which is highly expressed in the

transverse colon, and 40 microbiome genes. These 40 gut microbiome genes included those

associated with fructose-mannose metabolism, such as K01840 (manB), K00850 (pfkA), and

K01818 (fucI), as well as those related to oxidative phosphorylation, such as K00241 (sdhC,

frdC), K00426 (cydB), K02112 (atpD), and K02121 (ntpE, atpE). The most substantial

contributor to the expression of these microbiome genes was *Bacteroides vulgatus*.

FBP1 is known to have reduced expression in various tumor tissues, including the esophagus

[27], liver [28], lung [29], and ovarian cancer [30]. It has been confirmed that its expression

increases with fructose treatment in intestinal organoids [31]. Previous studies have identified

FBP1 as a differentially expressed gene in the small intestine of germ-free mice, suggesting that

the intestinal microbiome influences the expression of the host's FBP1 gene [32], but the

bacterial species and genes affecting the expression of the FBP1 gene have not been identified.

Bacteroides vulgatus, which contributed to the expression of fructose-mannose metabolism-

related microbiome genes observed in Module_06, has been reported to release fructose by

decomposing polysaccharides [33, 34]. In this module, the gene K00850 (*pfkA*) involved in the

fructo-oligosaccharide degradation and the gene K01818 (*fucI*) involved in the 2'-fucosyllactose

degradation were detected, revealing a novel association between polysaccharide degradation

genes encoded by *Bacteroides vulgatus* and the host gene FBP1.

In Module_01, the B cell receptor signaling pathway was enriched (Fig. 6). This module

featured covariation between host genes involved in B cell immune responses—including CD19,

CD22, CD79B, PTPN6, which were highly expressed in the cecum—and the tryptophan

synthesis gene K01696 (*trpB*). This tryptophan synthesis gene K01696 (*trpB*) expression was

contributed to by *Parabacteroides distasonis*. Tryptophan is an essential amino acid that, upon

uptake by intestinal cells and subsequent metabolism, can activate the Aryl hydrocarbon receptor

and influence B cell immune responses [35]. Tryptophan is also known to ameliorate

autoimmune diseases, such as inflammatory bowel disease [36]. *Parabacteroides distasonis* has

been suggested to be involved in various autoimmune conditions in several studies [37, 38]. For

instance, *Parabacteroides distasonis* levels have been shown to be lower in multiple sclerosis

(MS) patients compared to healthy controls, and transplanting the intestinal microbiome from

MS patients to germ-free mice resulted in increased severity of autoimmune encephalomyelitis

symptoms [37]. While the associations between tryptophan, B-cell immune responses, and

autoimmune diseases, as well as between *Parabacteroides distasonis* and autoimmune diseases,

are already reported, our study provides additional insight. We suggest that tryptophan

synthesized by *Parabacteroides distasonis* could exert an influence on the upregulation of B-cell

immune response-related genes in the cecum, including CD19, CD22, CD79B, and PTPN6.

In Module_19, a group consisting of three host genes and 40 microbiome genes was

identified. The three host genes included MMP10, two uncharacterized genes LOC103795407

and CCDC162P, all of which showed high expression in the cecum (Fig. 7). The microbiome

genes that showed a positive correlation with these genes included K01491 (folD), K00605

(gcvT), K01803 (*tpiA*), K00831 (*serC*), K15633 (*gpmI*), K00024 (*mdh*), all of which are

involved in carbon metabolism pathways (Supplementary Fig. S1). *Bacteroides vulgatus* was

identified as the bacteria contributing to the expression of these microbiome genes. The MMP10

gene plays a crucial role in maintaining epithelial barrier function and is necessary for resolving

colonic epithelial damage [39]. A study examining the host transcriptome and microbiota

composition of colorectal cancer tumors found high expression of MMP10 in tumor tissues and a

significant increase in the genus *Bacteroides*, indicating a link between the host gene MMP10

and the genus *Bacteroides* [40]. These findings are consistent with our observations.

Furthermore, in the identified Module_19, not only the relationship between the host gene

MMP10 and *Bacteroides vulgatus*, but also a group of microbiome genes related to carbon

metabolism pathways showing a positive correlation with the host gene MMP10 was identified.

One hypothesis that suggests a positive correlation between these carbon metabolism-related

genes coded by *Bacteroides vulgatus* and the host gene MMP10 is that the microbiome might

form colonies in the host's mucosal layer. Mucin, the main component of mucus generated by the

gastrointestinal mucosal epithelium, contains various sugars such as glucose and fructose [41]. In

addition to promoting bacterial settlement and adhesion, it also serves as a nutrient source for

bacteria. In this Module_19, it was observed that the microbiome gene group, including K01803

(*tpiA*), K00831 (*serC*), K15633 (*gpmI*) related to the glycolysis of glucose, fructose, and

galactose, positively correlated with the host gene MMP10, which is upregulated specifically in

the cecum. This suggests the possibility that in the cecum, the microbiome, including

Bacteroides vulgatus, forms colonies using mucin as a nutrient source, and the high expression

of the host's MMP10 gene protects the epithelium from the colonization of the microbiome. In

this Module_19, we observed that a group of peptide genes such as K01803 (*tpiA*), K00831

(*serC*), and K15633 (*gpmI*) related to glycolysis of glucose, fructose, and galactose were

positively correlated with the host gene MMP10, which was up-regulated specifically in the

cecum. This observation suggests that in the cecum, microbiome including *Bacteroides vulgatus*

colonize the mucin, the main mucus component, as a nutrient source, and that the host MMP10

gene is highly expressed to protect the epithelium from microbiome colony formation.

In Module_16, three host genes and 54 microbiome genes were identified, and

furthermore, the expression of these microbiome genes was influenced by five bacterial species.

The relationships between these host genes and microbiome genes, as well as between the host

genes and bacterial species, were consistent with existing knowledge about bile acid synthesis

and degradation (Fig. 8). This module included the host gene CYP27A1, which is involved in

bile acid production [42], and NR1H3, a host gene that codes for a bile acid receptor [43]. These

genes were upregulated in the cecum and transverse colon compared to the rectum. The

microbiome genes interacting with these host genes included K03664 (*smpB*), a gene essential

for bacterial survival when upregulated in the presence of secondary bile acid deoxycholic acid

(DCA), K02835 (*prfA*) which positively regulates the expression of a gene coding for bile acid

hydrolase, and K03527 (*ispH*, *lytB*), a gene involved in bile acid sensitivity. Additionally,

Bacteroides vulgatus contributed to the expression of all these microbiome genes, which aligns

with previous studies showing an increase in *Bacteroides* spp. correlating with the expression of

the host gene NR1H3 [44].

DISCUSSION

In this study, we investigated the interactions between the host and intestinal microbiome in the

common marmoset's intestinal sites (cecum, transverse colon, and rectum) by utilizing dual

transcriptome profiling and subsequent gene co-expression network analysis. To discern modules

of tightly interacting host and microbiome genes from a vast gene co-expression network with

nodes representing both host and microbiome genes, we proposed a clique-based algorithm that

considers the interconnectedness of nodes (cliqueness), in three different networks: host-host,

host-microbiome, and microbiome-microbiome. As a result, we identified 27 co-expression

modules composed of both host and microbiome genes, with 70.4% of them, or 19 modules,

being functionally characterized by enrichment analysis (Fisher's exact test, FDR<0.05;

Supplementary Table S9).

To compare our method with conventional module detection methods, we applied the

Newman [14] and Louvain [15] algorithms, both of which maximize modularity, to the same

gene co-expression network. Upon conducting pathway enrichment analyses on the modules

extracted by these conventional methods, the proportion of functionally characterized modules

was only 22.2% and 50.0% respectively, which is lower than what we achieved with our method

(Supplementary Table S10). These results underscore the success of our method in defining a

larger number of functionally interpretable gene modules compared to conventional approaches

and emphasize its suitability for predicting host-microbiome interactions, over conventional

modularity-maximizing approaches.

The gene modules extracted, which encompassed both host and microbiome genes,

frequently exhibited relationships where a small number of genes regulated a larger number of

genes between the host and microbiome genes. These observations are thought to reflect the

characteristics of the interactions between the host and the microbiome. Several studies have

reported the impact of host gene mutations on the intestinal microbiome. For instance, it has

been reported that the host gene VDR, which codes for the Vitamin D receptor, influences the

colonization of *Parabacteroides* [45], and that the host gene ABO is associated with

Faecalicatena lactaris [46]. In another study investigating the relationship between the intestinal

microbiome and host gene loci through quantitative trait locus analysis, it was suggested that

Akkermansia muciniphila regulates the expression of ATF3, which codes for a transcription

factor that plays a crucial role in metabolic and immune regulation, and this bacterium is related

to only four genes [47]. These studies highlight the fact that a single or a small number of host

genes are controlling, or are controlled by, the intestinal microbiome. Conversely, a small

number of microbiome genes can affect the host's metabolic [48], fitness [49], and longevity [50]

phenotypes. For example, structural mutations in the gene related to butyrate transport in

Anaerostipes hadrus are strongly associated with a reduction in the host's metabolic risk [48],

and the intestinal microbiome's thiamine biosynthesis genes, transferred by horizontal

transmission, influence the host's fitness [49]. Considering these previous studies and our results,

it is suggested that the interaction between the host and the microbiome involves widespread

influence of one's genes on the activity of the other.

The common marmoset used in this study bears anatomical and pharmacological

similarities to humans [17], and the fecal microbiome of the common marmoset is more similar

to that of humans compared to other animal models [13]. While the common marmoset's

cecum—one of the study's targeted intestinal sites—does not possess an appendage as

pronounced as the human appendix, it is known to share significant similarities with the human

appendix in terms of microanatomical structure and immune reactivity [51, 52]. The results of

this study using common marmosets could therefore hold higher applicability to humans

compared to other animal models.

In conclusion, we union host transcriptome and microbiome metatranscriptome profiles

of the common marmoset and performed gene co-expression network analysis. The algorithm we

proposed for extracting gene modules in the gene co-expression network enabled the

identification of dense, functionally related host-microbiome co-expression modules, which was

challenging using the conventional module maximization algorithms. The identified gene

modules provided new insights into relationships between host genes and intestinal microbiome

and microbial genes bridging interactions between the host and bacteria. Our findings underscore

that interactions between the host and microbiome are not uniform throughout the intestine but

are localized, indicating spatial variation in host-microbiome interactions along the gut. These

insights could be crucial in elucidating mechanisms of localized gut diseases, such as colorectal

cancer and inflammatory bowel disease, and in developing localized therapies.

MATERIALS AND METHODS

Animal experiments

Common marmosets were housed at the Central Institute for Experimental Animals (Kawasaki,

Japan). Five marmosets were euthanized with intravenous administration of pentobarbital

overdose, and the digestive tract was isolated (Supplementary Table S11, Supplementary Note

Section 1). Three gastrointestinal site samples (cecum, transverse colon and rectum), including

the mucosal layer, were used for dual RNA-seq. These three sites were targeted at the beginning,

middle and end of the colon, which has an abundant microbiome. The animal experiment

protocol was approved by the CIEA Institutional Animal Care and Use Committee (approval no.

17031, 18032, and 21084). The study was conducted in accordance with the guidelines of CIEA

that comply with the Guidelines for Proper Conduct of Animal Experiments published by the

Science Council of Japan. Animal care was conducted in accordance with the Guide for the Care

and Use of Laboratory Animals (Institute for Laboratory Animal Resources, 2011).

RNA extraction and dual RNA sequencing of host and microbiome

RNA libraries were prepared after removing rRNA from the extracted high-quality Total RNA

(RNA integrity number (RIN) scores ≥8.6; Supplementary Table S1). Illumina NovaSeq 6000

v1.5 sequencing yielded a total of 472 Gnt of paired-end reads (100 nt × 2). This dataset included

an average of 314.9 M reads per sample (Supplementary Table S2, Supplementary Note Section

2).

Construction of host-microbiome co-expression network and identification of gene modules

Network analysis

In this study, we utilized gene co-expression network analysis to identify interacting gene

modules of host and microbiome. We established three types of networks: host-host, host-

microbiome, and microbiome-microbiome, wherein genes were considered as nodes and co-

expressing genes were interconnected (Fig. 1D). Networks were defined by an adjacency matrix,

where the adjacency a_{ij} between gene i and gene j was defined as follows based on co-

expression similarity s_{ij}:

405
$$a_{ij} = \begin{cases} 1 \text{, if } s_{ij} > \alpha \\ 0 \text{, otherwise} \end{cases}$$

The co-expression similarity s_{ij} was computed using Spearman's correlation coefficients for

host-host and microbiome-microbiome networks, and absolute values of Spearman's correlation

coefficients for host-microbiome networks. The adjacency threshold α was set at 0.80. These

three types of networks were then integrated into one.

Identification of gene modules

We developed an algorithm to identify densely interacting gene modules from the host-

microbiome co-expression network. This algorithm, based on clique finding, progressively

merges cliques (Fig. 1E). In graph theory, a "clique" refers to a subgraph within the network

where all nodes are fully connected. In the context of a gene co-expression network, all genes

within a clique indicate a strong co-expression (correlation) amongst them. In this study, we

focus only on networks and cliques that include gene nodes from both the host and microbiome,

encompassing three types of networks: host-host, host-microbiome, and microbiome-

microbiome. Our algorithm identified modules of interacting genes in the gene co-expression

network as aggregates of overlapping cliques. It was implemented in three steps: (1) Cliques

within the network were found; (2) Beginning with the largest cliques, we examined their

overlap with any other cliques. If overlap occurred, we computed the cliqueness scores

independently for each of the three types of networks (host-host, host-microbiome, and

microbiome-microbiome networks) upon merging the overlapping cliques. The cliqueness scores

for the host-host, host-microbiome, and microbiome-microbiome networks were denoted as C_{hh},

C_{hm}, and C_{mm}, respectively; (3) Cliques were merged if they satisfied the condition: $C_{hh} > \tau$,

$C_{mm} > \tau$, and $C_{mm} > \tau$. The threshold τ for cliqueness was set at 0.60 based on parameter

determination (Supplementary Note Section 7). If there were multiple cliques with minimum

cliqueness scores exceeding the threshold, the clique with the highest cliqueness score was

selected. Steps (2) and (3) were repeated until no cliques could be merged.

For the host-host networks the cliqueness score C_{hh} between cliques A and B was defined as:

431
$$C_{hh} = \frac{E}{\frac{1}{2}N(N-1)}$$

Here, N is defined as the number of host gene nodes in the union of cliques A and B, represented

as $N = |N_A \cup N_B|$. N_A and N_B denote the number of host gene nodes within clique A and clique

B, respectively. E corresponds to the total number of edges between the host nodes included in

cliques A and/or B, formulated as $E = E_A + E_B + E_{AB}$. Here, E_A and E_B represent the number of

edges between host gene nodes within clique A and clique B, respectively, while E_{AB} denotes the

number of edges connecting across host gene nodes within clique A and clique B. The cliqueness

score for microbiome-microbiome networks C_{mm} was calculated in the same way.

For the host-microbiome network, the cliqueness score C_{hm}, was defined as:

440
$$C_{hm} = \frac{E}{N_{host} N_{microbiome}}$$

where N_{host} and $N_{microbiome}$ represent the number of host and microbiome gene nodes

respectively in the union of cliques A and B. E corresponds to the total number of edges between

the host or microbiome gene nodes included in cliques A and/or B.

Acknowledgements

We thank the scientists of the Central Institute for Experimental Animals for providing support for

animal experiments and the members of the Sakakibara Lab at Keio University for helpful

discussions.

This work was supported by grants from the Japan Agency for Medical Research and Development

(AMED PRIME JP19gm6010006). M. Uehara has received funding from JSPS KAKENHI Grant

Numbers JP20J21477.

Author contribution

M.U. performed the experiments, conducted the bioinformatics analysis and cowrote the paper;

T.I. and E.S. provided common marmoset samples; M.U. and S.H. performed RNA extraction; A.T.

performed deep sequencing with high-throughput sequencer; Y.S. designed and supervised the

research, analysed the data, and cowrote the paper. All authors have read and approved the

manuscript.

Competing Interests

The authors declare no competing interests.

Data and source code availability

All raw sequence data have been submitted to the DDBJ under project PSUB020951 from the

Ministry of Education, Culture, Sports, Science and Technology of Japan.

REFERENCES

1. Yoon HS, Cho CH, Yun MS, Jang SJ, You HJ, Kim JH, et al. Akkermansia muciniphila secretes a glucagon-like peptide-1-inducing protein that improves glucose homeostasis and ameliorates metabolic disease in mice. Nat Microbiol. 2021;6:563–73.

2. Vatanen T, Kostic AD, D'Hennezel E, Siljander H, Franzosa EA, Yassour M, et al. Variation in Microbiome LPS Immunogenicity Contributes to Autoimmunity in Humans. Cell. 2016;165:842–53.

3. Song X, Sun X, Oh SF, Wu M, Zhang Y, Zheng W, et al. Microbial bile acid metabolites modulate gut RORγ+ regulatory T cell homeostasis. Nature. 2020;577:410–5.

4. Priya S, Burns MB, Ward T, Mars RAT, Adamowicz B, Lock EF, et al. Identification of shared and disease-specific host gene–microbiome associations across human diseases using multi-omic integration. Nat Microbiol. 2022;7:780–95.

5. Zhang L, Forst CV, Gordon A, Gussin G, Geber AB, Fernandez PJ, et al. Characterization of antibiotic resistance and host-microbiome interactions in the human upper respiratory tract during influenza infection. Microbiome. 2020;8:1–12.

6. Fujiogi M, Raita Y, Pérez-Losada M, Freishtat RJ, Celedón JC, Mansbach JM, et al. Integrated relationship of nasopharyngeal airway host response and microbiome associates with bronchiolitis severity. Nat Commun. 2022;13:4970.

7. Lloyd-Price J, Arze C, Ananthakrishnan AN, Schirmer M, Avila-Pacheco J, Poon TW, et al. Multi-omics of the gut microbial ecosystem in inflammatory bowel diseases. Nature. 2019;569:655–62.

8. Vertzoni M, Augustijns P, Grimm M, Koziolek M, Lemmens G, Parrott N, et al. Impact of

regional differences along the gastrointestinal tract of healthy adults on oral drug

absorption: An UNGAP review. Eur J Pharm Sci. 2019;134:153–75.

9. Ilhan ZE, Marcus AK, Kang DW, Rittmann BE, Krajmalnik-Brown R. pH-mediated

microbial and metabolic interactions in fecal enrichment cultures. mSphere.

2017;2:e00047-17.

10. Yasuda K, Oh K, Ren B, Tickle TL, Franzosa EA, Wachtman LM, et al. Biogeography of

the intestinal mucosal and lumenal microbiome in the rhesus macaque. Cell Host

Microbe. 2015;17:385–91.

11. Kozik AJ, Nakatsu CH, Chun H, Jones-Hall YL. Comparison of the fecal, cecal, and

mucus microbiome in male and female mice after TNBS-induced colitis. PLoS One.

2019;14:1–14.

12. Zhang L, Wu W, Lee YK, Xie J, Zhang H. Spatial heterogeneity and co-occurrence of

mucosal and luminal microbiome across swine intestinal tract. Front Microbiol.

2018;9:48

13. Uehara M, Inoue T, Kominato M, Hase S, Sasaki E, Toyoda A, et al. Intraintestinal

Analysis of the Functional Activity of Microbiomes and Its Application to the Common

Marmoset Intestine. mSystems. 2022;7:e00520-22.

14. Newman ME. Detecting community structure in networks. Eur Phys J B. 2004;38:321–

30.

15. Blondel VD, Guillaume JL, Lambiotte R, Lefebvre E. Fast unfolding of communities in

large networks. J Stat Mech. 2008;10008.

16. Hugenholtz F, de Vos WM. Mouse models for human intestinal microbiota research: a

critical evaluation. Cell Mol Life Sci. 2018;75:149–60.

17. Kishi N, Sato K, Sasaki E, Okano H. Common marmoset as a new model animal for

neuroscience research and genome editing technology. Dev Growth Differ. 2014;56:53–

62.

18. Inoue T, Okahara N, Okahashi N, Sato K, Ueda M, Atarashi K, et al. 2021. Generation of

germ-free common marmosets. Research Square. Preprint.

https://assets.researchsquare.com/files/rs-428622/v1_covered.pdf?c=1631871455

19. Wang K, Wei G, Liu D. CD19: a biomarker for B cell development, lymphoma diagnosis

and therapy. Exp Hematol Oncol. 2012;1:1–7.

20. Huse K, Bai B, Hilden VI, Bollum LK, Våtsveen TK, Munthe LA, et al. Mechanism of

CD79A and CD79B support for IgM+ B cell fitness through B cell receptor surface

expression. J Immunol. 2022;209:2042–53.

21. Clark EA, Giltiay N V. CD22: a regulator of innate and adaptive B cell responses and

autoimmunity. Front Immunol. 2018;9:2235.

22. Pan C, Baumgarth N, Parnes JR. CD72-deficient mice reveal nonredundant roles of

CD72 in B cell development and activation. Immunity. 1999;11:495–506.

23. M.L. A, K.A. F, C.B. T. T-cell regulation by CD28 and CTLA-4. Nat Rev Immunol.

2001;1:220–8.

24. Yang Y, Zang Y, Zheng C, Li Z, Gu X, Zhou M, et al. CD3D is associated with immune

checkpoints and predicts favorable clinical outcome in colon cancer. Immunotherapy.

2020;12:25–35.

25. Hubbard N, Hagin D, Sommer K, Song Y, Khan I, Clough C, et al. Targeted gene editing

restores regulated CD40L function in X-linked hyper-IgM syndrome. Blood.

2016;127:2513–22.

26. Sobhani N, Tardiel-Cyril DR, Davtyan A, Generali D, Roudi R, Li Y. CTLA-4 in

regulatory T cells for cancer immunotherapy. Cancers. 2021;13:1–18.

27. He Y, Hua R, Li B, Gu H, Sun Y, Li Z. Loss of FBP1 promotes proliferation, migration,

and invasion by regulating fatty acid metabolism in esophageal squamous cell carcinoma.

Aging. 2021;13:4986–8.

28. Hirata H, Sugimachi K, Komatsu H, Ueda M, Masuda T, Uchi R, et al. Decreased

expression of fructose-1,6-bisphosphatase associates with glucose metabolism and tumor

progression in hepatocellular carcinoma. Cancer Res. 2016;76:3265–76.

29. Dong Y, Huaying S, Danying W, Chihong Z, Ruibin J, Xiaojiang S, et al. Significance of

methylation of FBP1 gene in non-small cell lung cancer. Biomed Res Int. 2018;2018:13–

9.

30. Li H, Qi Z, Niu Y, Yang Y, Li M, Pang Y, et al. FBP1 regulates proliferation, metastasis,

and chemoresistance by participating in C-MYC/STAT3 signaling axis in ovarian cancer.

Oncogene. 2021;40:5938–49.

31. Hou Y, Wei W, Guan X, Liu Y, Bian G, He D, et al. A diet-microbial metabolism

feedforward loop modulates intestinal stem cell renewal in the stressed gut. Nat

Commun. 2021;12:271.

32. Mardinoglu A, Shoaie S, Bergentall M, Ghaffari P, Zhang C, Larsson E, et al. The gut

microbiota modulates host amino acid and glutathione metabolism in mice. Mol Syst

Biol. 2015;11:1–15.

33. Medina DA, Pinto F, Ovalle A, Thomson P, Garrido D. Prebiotics mediate microbial

interactions in a consortium of the infant gut microbiome. Int J Mol Sci. 2017;18:2095.

34. Cheng J, Hu J, Geng F, Nie S. Bacteroides utilization for dietary polysaccharides and

their beneficial effects on gut health. Food Sci Hum Wellness. 2022;11:1101–10.

35. Rosser EC, Piper CJM, Matei DE, Blair PA, Rendeiro AF, Orford M, et al. Microbiota-

derived metabolites suppress arthritis by amplifying aryl-hydrocarbon receptor activation

in regulatory B cells. Cell Metab. 2020;31:837-851.

36. Ding X, Ding X, Ding X, Bin P, Bin P, Bin P, et al. Tryptophan metabolism, regulatory T

cells, and inflammatory bowel disease: a mini review. Mediators Inflamm. 2020;2020.

37. Ezeji JC, Sarikonda DK, Hopperton A, Erkkila HL, Cohen DE, Martinez SP, et al.

Parabacteroides distasonis: intriguing aerotolerant gut anaerobe with emerging

antimicrobial resistance and pathogenic and probiotic roles in human health. Gut

Microbes. 2021;13:1–27.

38. Cekanaviciute E, Yoo BB, Runia TF, Debelius JW, Singh S, Nelson CA, et al. Gut

bacteria from multiple sclerosis patients modulate human T cells and exacerbate

symptoms in mouse models. Proc Natl Acad Sci USA. 2017;114:10713–8.

39. Łaniewski P, Herbst-Kralovetz MM. Bacterial vaginosis and health-associated bacteria

modulate the immunometabolic landscape in 3D model of human cervix. NPJ Biofilms

Microbiomes. 2021;7:1–17.

40. Tomkovich S, Gharaibeh RZ, Dejea CM, Pope JL, Jiang J, Winglee K, et al. Human

colon mucosal biofilms and murine host communicate via altered mRNA and microRNA

expression during cancer. mSystems. 2020;5:e00451-19.

41. Cornick S, Tawiah A, Chadee K. Roles and regulation of the mucus barrier in the gut.

Tissue Barriers. 2015;3:1–2.

42. Zurkinden L, Sviridov D, Vogt B, Escher G. Downregulation of Cyp7a1 by cholic acid

and chenodeoxycholic acid in Cyp27a1/ApoE double knockout mice: differential

cardiovascular outcome. Front Endocrinol. 2020;11:1–13.

43. Biagioli M, Marchianò S, Carino A, Di Giorgio C, Santucci L, Distrutti E, et al. Bile

acids activated receptors in inflammatory bowel disease. Cells. 2021;10:1281.

44. Badi SA, Motahhary A, Bahramali G, Masoumi M, Khalili SFS, Ebrahimzadeh N, et al.

The regulation of Niemann-Pick C1-Like 1 (NPC1L1) gene expression in opposite

direction by Bacteroides spp. and related outer membrane vesicles in Caco-2 cell line. J

Diabetes Metab Disord. 2020;19:415–22.

45. Wang J, Thingholm LB, Skiecevičie J, Rausch P, Kummen M, Hov JR, et al. Genome-

wide association analysis identifies variation in Vitamin D receptor and other host factors

influencing the gut microbiota. Nat Genet. 2016;48:1396–406.

46. Qin Y, Havulinna AS, Liu Y, Jousilahti P, Ritchie SC, Tokolyi A, et al. Combined effects

of host genetics and diet on human gut microbiota and incident disease in a single

population cohort. Nat Genet. 2022;54:134–42.

47. Zhang Q, Linke V, Overmyer KA, Traeger LL, Kasahara K, Miller IJ, et al. Genetic

mapping of microbial and host traits reveals production of immunomodulatory lipids by

Akkermansia muciniphila in the murine gut. Nat Microbiol. 2023;8:424–40.

48. Zeevi D, Korem T, Godneva A, Bar N, Kurilshikov A, Lotan-Pompan M, et al. Structural

variation in the gut microbiome associates with host health. Nature. 2019;568:43–8.

49. Wang Y, Baumdicker F, Schweiger P, Kuenzel S, Staubach F. Horizontal gene transfer-

mediated bacterial strain variation affects host fitness in Drosophila. BMC Biol. 2021

Dec 1;19(1).

50. Han B, Sivaramakrishnan P, Lin CCJ, Neve IAA, He J, Tay LWR, et al. Microbial genetic

composition tunes host longevity. Cell. 2017;169:1249-1262.

51. Zinnen AD, Vichich J, Metzger JM, Gambardella JC, Bondarenko V, Simmons HA, et al.

Alpha-synuclein and tau are abundantly expressed in the ENS of the human appendix and

monkey cecum. PLoS One. 2022;17:e0269190.

52. Fisher RE. The primate appendix: A reassessment. Vol. 261, Anatomical Record. 2000. pp

228–36.

Fig. 1 Overview of the main analytical approach used in this study.

This method involves the simultaneous acquisition of the host and microbiome transcriptome

(metatranscriptome) from intestinal sites and conducts a gene co-expression network analysis of

the host and microbiome community. The procedure is outlined from step A to E. Perform dual

RNA-seq of host and microbiome transcriptomes in internal site samples. (A) Separate host and

microbiome RNA reads in silico using the host reference genome. (B) Compute the gene

expression profiles of the host and microbiome, respectively. (C) Unify gene expression

networks between host-host, host-microbiome, and microbiome-microbiome. The host-host and

microbiome-microbiome gene expression networks are connected when there is a positive

correlation between genes, while the host-microbiome gene expression network is connected

when there is either a positive or negative correlation between genes. (D) Identify closely

interacting gene modules from the gene expression network using a proposed clique-based

algorithm. (E) This process begins with searching for cliques (E Step1). If overlaps exist between

cliques, the cliqueness score is computed in three gene expression networks (host-host, host-

microbiome, and microbiome-microbiome) when overlapping cliques are merged (E Step2). If

the cliqueness score exceeds the merging threshold τ in all three networks, the cliques are

merged into a single module. If the score does not exceed the merging threshold τ in at least one

network, the original cliques are retained without merging. This process is repeated until no

further modules can be merged (E Step3).

Fig. 2 PCA analysis of gene expression data for host and microbiome

PCA was applied to the host gene expression profiles to visualize groupings according to site (A)

and individual (B). PCA was similarly applied to the microbiome gene expression profiles to

visualize groupings according to site (C) and individual (D).

Fig. 3 Variations in host gene expression and functional pathways among intestinal sites

(A) The top 20 functional pathways significantly upregulated in the cecum compared to the

transverse colon (FDR<0.1), as revealed by gene set enrichment analysis, are presented

alongside their Normalized Enrichment Score and FDR. (B) Heatmap of differentially expressed

genes belonging to the top 5 enriched functional pathways. The heatmap on the left represents

the z-scores of gene expression levels, with genes significantly upregulated in the cecum

compared to the transverse colon and rectum (FDR<0.01) colored in pink on the middle

heatmap. The heatmap on the right indicates the functional pathways to which the genes belong,

represented in green.

Fig. 4 Functional pathways characterized in modules extracted from host-microbiome gene

expression networks

This figure illustrates the functional characteristics of the modules determined by enrichment

analysis of functional pathways (FDR<0.1). The background of the bar graph alternates between

grey and white for each module for ease of differentiation.

Fig. 5 Metabolic pathway-enriched Module_06

A module showing a positive correlation between the host gene FBP1, which is highly expressed

in the transverse colon, and 40 microbiome genes, including those involved in fructose-mannose

metabolism and oxidative phosphorylation, was extracted. These genes were visualized

alongside bacterial species nodes that contribute to the expression of the microbiome genes.

Edges among genes indicate an absolute Spearman's correlation coefficient greater than 0.8.

Edges connecting microbiome genes to bacterial species are shown when the contribution rate of

bacterial species to microbiome gene expression exceeds 0.2. The thickness of the edge increases

with higher contribution rates.

Fig. 6 B cell receptor signaling pathway-enriched Module_01

Host genes involved in the immune response of B cells, including CD19, CD22, CD79B and

PTPN6, which were highly expressed in the cecum, co-expressed with the K01696 (*trpB*). These

genes were visualized alongside bacterial species nodes that contribute to the expression of the

microbiome genes. K01696 (*trpB*) expression was contributed to by *Parabacteroides distasonis*.

Edges among genes indicate an absolute Spearman's correlation coefficient greater than 0.8.

Edges connecting microbiome genes to bacterial species are shown when the contribution rate of

bacterial species to microbiome gene expression exceeds 0.2. The thickness of the edge increases

with higher contribution rates.

Fig. 7 Carbon metabolism pathway-enriched Module_19

Host genes, including MMP10, which is specifically highly expressed in the cecum, and 40

microbiome genes involved in carbon metabolism, were identified. These genes were visualized

alongside bacterial species nodes that contribute to the expression of the microbiome genes.

Edges among genes indicate an absolute Spearman's correlation coefficient greater than 0.8.

Edges connecting microbiome genes to bacterial species are shown when the contribution rate of

bacterial species to microbiome gene expression exceeds 0.2. The thickness of the edge increases

with higher contribution rates.

Fig. 8 Module_16 containing host genes and microbiome genes related to the production and

breakdown of bile acids. These genes were visualized alongside bacterial species nodes that

contribute to the expression of the microbiome genes. Edges among genes indicate an absolute

Spearman's correlation coefficient greater than 0.8. Edges connecting microbiome genes to

bacterial species are shown when the contribution rate of bacterial species to microbiome gene

expression exceeds 0.2. The thickness of the edge increases with higher contribution rates.

A) Dual RNA-seq of host and microbiome transctiptome

B) Separation of host and microbiome reads

C) Gene expression analysis

D) Construction of integrated gene expression network

Intestinal sample

Dual RNA-seq

Host RNA reads

Microbiome RNA reads

Host transcriptome profile

Metatranscriptome profile

Host-Host

Host-Microbiome

Microbiome-Microbiome

Host   Microbiome

E) Clique-based module extraction

Host   Microbiome

Host-Host

Host-Microbiome

Microbiome-Microbiome

✓ $Cliqueness > \tau$

✓ $Cliqueness > \tau$

✓ $Cliqueness > \tau$

Merge

Host   Microbiome

Host-Host

Host-Microbiome

Microbiome-Microbiome

✗ $Cliqueness \leq \tau$

✓ $Cliqueness > \tau$

✗ $Cliqueness \leq \tau$

Retain without merging

A Host gene expression — PC1 (24.7%) vs PC2 (15.4%)

B Host gene expression — PC1 (24.7%) vs PC2 (15.4%)

C Microbiome gene expression — PC1 (75.1%) vs PC2 (9.1%)

D Microbiome gene expression — PC1 (75.1%) vs PC2 (9.1%)

Individual:
- CJ01
- CJ02
- CJ03
- CJ04
- CJ05

Intestinal site:
- Cecum
- T.colon
- Rectum

Host gene	Microbiome gene	Microbiome

FBP1

K01818
K00241
K03631
K00174
K02004
K06168
K02121
K03654
K08325
K00876
K01000
K01814
K03269
K00721
K00677
K01736
K09815
K00764
K02337
K01840
K09825
K03585
K02519
K07566
K03701
K03595
K00426
K05515
K07588
K03771
K06973
K01647
K01284
K00133
K00850
K02112
K14652
K00661
K06213
K00655

Akkermansia_muciniphila

Bacteroides_dorei

Bacteroides_vulgatus

Bacteroides_uniformis

- ● Gene with the enriched function
- ● Gene without the enriched function
- ■ Species

— Positive correlation
— Negative correlation
— Microbiome gene expression contribution

Host gene   Microbiome gene   Microbiome

MMP10

CCDC162P

LOC103795407

K02563
K01972
K12267
K03977
K01893
K03664
K02564
K00024
K01928
K00605
K02528
K03474
K00566
K03797
K06287
K01610
K06949
K01770
K03979
K02551
K00791
K01939
K01803
K03046
K01924
K01258
K00831
K00939
K07407
K03551
K03650
K03572
K03527
K01491
K01881
K01735
K03526
K15633
K00215
K09903

Akkermansia_muciniphila

Parabacteroides_distasonis

Clostridioides_difficile

Flavonifractor_plautii

Bacteroides_vulgatus

● Gene with the enriched function
● Gene without the enriched function
■ Species

— Positive correlation
— Negative correlation
— Microbiome gene expression contribution

Host gene | Microbiome gene | Microbiome

K01662
K02563
K03979
K00147
K03590
K02860
K03797
K03650
K01918
K01735
K06949
K03977
K00604
K03664
K01866
K01939
K01887
K06168
K01869
K01876
K02876
K01000
K07407
K03551
K00930
K04077
K02528
K03217
K02885
K01868
K03527
K01881
K01889
K15633
K00215
K09903

NR1H3

CYP27A1

LOC103790783

Akkermansia_muciniphila
Streptococcus_lutetiensis
Parabacteroides_distasonis
Flavonifractor_plautii
Bacteroides_vulgatus

Legend:
● Gene
■ Species

— Positive correlation
— Negative correlation
— Microbiome gene expression contribution