Original Research:

Hippocampal-occipital connectivity reflects autobiographical memory deficits

in aphantasia

Merlin Monzel[1,2,*], Pitshaporn Leelaarporn[2,3,*], Teresa Lutz[2], Johannes Schultz[4,5], Sascha

Brunheim[3], Martin Reuter[1], and Cornelia McCormick[2,3]

*Both authors contributed equally.

1. Department of Psychology, University of Bonn, 53113 Bonn, Germany

2. German Center for Neurodegenerative Diseases, 53127 Bonn, Germany

3. Department of Neurodegenerative Diseases and Geriatric Psychiatry, University of Bonn

Medical Center, 53127 Bonn, Germany

4. Center for Economics and Neuroscience, University of Bonn, 53113 Bonn, Germany

5. Institute of Experimental Epileptology and Cognition Research, Medical Faculty, University

of Bonn, 53127 Bonn, Germany

Corresponding authors: Cornelia McCormick (cornelia.mccormick@dzne.de), Merlin Monzel

(merlin.monzel@uni-bonn-diff.de)

Conflict of interest statement:

The authors declare no competing financial interest.

Acknowledgments

C.M. was supported by the German Research Foundation (DFG, MC244/3-1) and the DZNE

Foundation (DZNE Stiftung-Forschung für ein Leben ohne Demenz, Parkinson & ALS). We thank

Anke Rühling and Jennifer Schlee for their technical support during scanning and Yilmaz Sagik for

serving as the secondary scorer for the Autobiographical Interview. In addition, we would like to

thank all participants for their time and effort to share their memories with us.

Abstract

Aphantasia prohibits people from experiencing visual imagery. While most of us can readily recall decade-old personal experiences (autobiographical memories, AM) with vivid mental images, there is a dearth of information about whether the loss of visual imagery in aphantasics affects their AM retrieval. The hippocampus is thought to be a crucial hub in a brain-wide network underlying AM. One important question is whether this network, especially the connectivity of the hippocampus, is altered in aphantasia. In the current study, we tested 14 congenital aphantasics and 16 demographically matched controls in an AM fMRI task to investigate how key brain regions (i.e., hippocampus and visual-perceptual cortices) interact with each other during AM re-experiencing. All participants were interviewed regarding their autobiographical memory to examine their episodic and semantic recall of specific events. Aphantasics reported more difficulties in recalling AM, were less confident about their memories, and described less internal and emotional details than controls. Neurally, aphantasics displayed decreased hippocampal and increased visual-perceptual cortex activation during AM retrieval compared to controls. In addition, controls showed strong negative functional connectivity between the hippocampus and the visual cortex during AM and resting-state functional connectivity between these two brain structures predicted better visualization skills. Our results indicate that visual mental imagery is essential for detail-rich, vivid AM, and that this type of cognitive function is supported by the functional connection between the hippocampus and the visual-perceptual cortex.

Keywords: autobiographical retrieval, episodic memory, visual cortex, neural networks, functional connectivity

Introduction

Our unique and personal memories are stored in autobiographical memories (AM) providing

stability and continuity of our self (Svoboda et al., 2006). For most of us, travelling mentally back

in time and re-visiting such unique personal events is associated with vivid, detail-rich mental

imagery (D'Argembeau & Van der Linden, 2006; Greenberg & Knowlton, 2014). This vivid mental

imagery during the re-experiencing of AMs has become a hallmark of autonoetic, episodic AM

retrieval. However, up to date, it remains unclear to what extent episodic AM retrieval depends

on visual mental imagery and what neural consequences a lack of mental imagery has on episodic

AM retrieval. This knowledge gap exists because separating AM retrieval from mental imagery is

a complex and challenging task.

One way to address this conundrum is to study people with aphantasia (Zeman et al., 2015).

Recent research defines aphantasia as a neuropsychological condition in which people

experience a marked reduction or complete lack of voluntary sensory imagery (Monzel et al.,

2022). This condition is associated with psychophysiological alterations, such as reduced

imagery-induced pupil contraction (Kay et al., 2022) and reduced imagery-induced priming

effects (Keogh & Pearson, 2018; Monzel, Keidel, et al., 2021). Thus, aphantasics offer the unique

opportunity to examine the consequences for episodic AM retrieval in the absence of voluntary

imagery. Indeed, a handful of previous studies report convergent evidence that aphantasics

report less sensory AM details than controls (Dawes et al., 2020, 2022; Milton et al., 2020; Zeman

et al., 2020), yet task-based functional activity has not been fully explored.

Neurally, the hippocampus has been established as a central brain structure to support the

detail-rich episodic AM retrieval in the healthy brain (Bauer et al., 2017; Brown et al., 2018;

Burianova et al., 2010; McCormick et al., 2020; Moscovitch et al., 2005), albeit some studies

differentiate between hippocampal support for remote and recent autobiographical memories

(see Bayley et al. 2016). In fact, hippocampal activity correlates with the vividness of AM

recollection (Addis et al., 2004; Sheldon & Levine, 2013) and patients with hippocampal damage

show marked deficits in detailed episodic AM retrieval (Miller et al., 2020; Rosenbaum et al.,

2008). In addition, neuroimaging studies illuminate that the hippocampus is almost always co-

activated with a wider set of brain regions, including the ventromedial prefrontal cortex (vmPFC),

lateral and medial parietal cortices, as well as visual-perceptual cortices (Svoboda et al., 2006).

Interestingly, especially during the elaboration phase of AM retrieval, when people engage in the

active retrieval of episodic details to a specific AM, the hippocampus exhibits a strong functional

connection to the visual-perceptual cortices, suggesting a crucial role of this connection for the

embedding of visual-perceptual details into AMs (McCormick et al., 2015).

Yet, not many studies have examined the neural correlates of aphantasia, and none during

AM retrieval. Of the little evidence there is, reports converge on a potential hyperactivity of the

visual-perceptual cortices in aphantasia (Fulford et al., 2018; Keogh et al., 2020). A prominent

theory posits that because of this hyperactivity, small signals elicited during the construction of

mental imagery may not be detected. If true, episodic AM retrieval deficits seen in aphantasia

may be due to altered connectivity between hippocampus and visual-perceptual cortices

(Pearson, 2019). In the same vein, Blomkvist (2023) proposes the extended constructive episodic

simulation hypotheses (CESH+) that suggests that imagination and memory rely on similar neural

structures, since both represent simulated recombinations of previous impressions. This

hypothesis has been supported by shared representations for memory and mental imagery in

early visual cortex (Albers et al., 2013; see also Zeidman & Maguire, 2016). Within this

framework, the hippocampus is supposed to initiate sensory retrieval processes (e.g., in the

visual-perceptual cortices; Danker & Anderson, 2010), comparable to its role in the hippocampal

memory indexing theory (Langille & Gallistel, 2020). Blomkvist (2023) speculates that in

aphantasics, either the hippocampal memory index or the retrieval processes may be impaired.

Thus, the main goal of our study was to examine the neural correlates of AM deficits associated

with aphantasia. We hypothesized that the deficits in AM seen in aphantasia rely on altered

involvement of the hippocampus, visual-perceptual cortices and their functional connectivity.

110   **Materials and methods**

Participants

In total, 31 healthy individuals with no previous psychiatric or neurological condition participated

in this study. 15 congenital aphantasics and 16 matched controls were recruited from the

database of the *Aphantasia Research Project Bonn* (Monzel, Keidel, et al., 2021; Monzel,

Vetterlein, et al., 2021). Due to technical issues during MRI scanning, one aphantasic had to be

excluded from the analyses. Groups were matched for basic demographic data, that is, sex, age,

and education, as well as intelligence assessed with a short intelligence screening (Baudson &

Preckel, 2015) (see Table 1). Oral and written informed consent was obtained from all

participants prior to the commencement of experimental procedure in accordance with the

Declaration of Helsinki (World Medical Association, 2013) and the local ethics board of the

University Hospital Bonn.

Table 1.

Demographic data for aphantasics, controls and the total sample.

	Total (n = 30)	Aphantasics (n = 14)	Controls (n = 16)	Test statistic	p	BF_{01}
Age				0.80^a	.431	2.30
M	29.77	31.47	28.19			
SD	11.36	10.45	12.27			
IQ						
M	93.77	91.73	95.69	0.81^a	.425	2.29
SD	13.53	16.61	10.02			
Sex				2.76^b	.097	0.69
Male (%)	32.3	53.3	81.3			
Female (%)	67.7	46.7	18.8			
Education				1.59^b	.662	7.90
Secondary school (%)	6.5	6.7	6.3			
A-levels (%)	35.5	40.0	31.3			
University degree	54.8	46.7	62.5			

(%)			
Doctoral		6.7	
degree	3.2		0.0
(%)			

Vividness of visual imagery questionnaire

Aphantasia is typically assessed with the Vividness of Visual Imagery Questionnaire (VVIQ; Marks, 1973, 1995), a subjective self-report questionnaire that measures how vivid mental scenes can be constructed by an individual. For example, individuals are asked to visualize a sunset with as much details as possible and rate their mental scene based on a 5-point Likert scale (ranging from 'no image at all, you only "know" that you are thinking of the object' to 'perfectly clear and as vivid as normal vision'). Since there are 16 items, the highest score of the VVIQ is 80 indicating the ability to visualize mental images with such vividness as if the event were happening right there and then. The minimum number of points is 16 indicating that an individual reported no mental image for any of the items at all. Aphantasia is at the lower end of the spectrum of imagery-abilities and usually diagnosed with a VVIQ-score between 16 and 32 (e.g., Dawes et al., 2020, 2022).

Binocular rivalry task

Since self-report questionnaires such as the VVIQ are associated with several drawbacks, such as their reliance on introspection (Schwitzgebel, 2002), we administered a mental imagery priming-based binocular rivalry task to assess mental imagery more objectively (for more details, see Keogh & Pearson, 2018; Pearson et al., 2008). In short, after imagining either red-horizontal or

blue-vertical Gabor patterns, participants were presented with a red-horizontal Gabor pattern to

one eye and a blue-vertical Gabor pattern to the other eye. Subsequently, participants were

asked to indicate which type of Gabor pattern they predominantly observed. Usually, successful

mental imagery leads subjects to select the Gabor pattern which they had just visualized. This

selection bias can be transferred into a priming score representing visual imagery strength. Mock

stimuli consisting of only red-horizontal or blue-vertical Gabor patterns were displayed in 12.5 %

of the trials to be able to detect decisional biases. Previous studies have shown that the binocular

rivalry task validly measures mental imagery strength (Pearson et al., 2011; Wagner & Monzel,

2023).

Autobiographical interview

Detailed behavioral AM measures were obtained in blinded semi-structured interviews either in-

person or online via Zoom (Zoom Video Communications Inc., 2016) using the Autobiographical

Interview (AI; Levine et al., 2002). During the AI, the interviewer asks the participant to recall five

episodic AMs from different life periods: early childhood (up to age 11), adolescent years (ages

11–17), early adulthood (ages 18–35), middle age (35–55), and the previous year. For participants

who were younger than 34 years, the middle age memory was replaced by another early

adulthood memory. Memories from the first four periods were considered remote, whereas the

memory from the previous year was considered recent. The interview is structured so that each

memory recollection consists of three parts: free recall, general probe, and specific probe. During

free recall, the participants were asked to recall as many details as possible for a memory of their

choice that is specific in time and place within the given time period. When the participant came

to a natural ending, the free recall was followed by the general and specific probes. During the

general probe, the interviewer asked the participant encouragingly to promote any additional

details. During the specific probe, specific questions were asked for details about the time, place,

perception, and emotion/thoughts of each memory. Then, participants were instructed to rate

their recall in terms of their ability to visualize the event on a 6-point Likert scale (ranging from

'not at all' to 'very much'). The interview was audiotaped, and afterwards transcribed and then

scored by two independent raters according to the standard protocol (Levine et al., 2002). The

interviews were scored after all data had been collected, in random order, and scorers were blind

to the group membership of the participant.

For scoring, the memory details were assigned to two broad categories, that is, internal and

external details. There were the following subcategories of internal details: internal events

(happenings, weather conditions, people present, actions), place (country, city, building, part of

room), time (year, month, day, time of the day), perceptual details (visual, auditory, gustatory,

tactile, smell, body position), and emotion/thought (emotional state, thoughts). The

subcategories for external details were semantic details (factual or general knowledge), external

events (other specific events in time and place but different to the main event), repetition

(repeated identical information without request), and other details (metacognitive statements,

editorializing). In addition, following the standard procedure, an episodic richness score was

given for each memory by the rater on a 7-point Likert scale (ranging from 'not at all' to 'perfect').

Furthermore, we added a novel rating score of confidence to the protocol since many

participants indicated very strong belief in the details they provided, while others were insecure

about the correctness of their own memories. Confidence scores were again rated on a 7-point

Likert scale (ranging from 'not at all' to 'perfect').

Debriefing questions

Following the AI, we asked participants three general questions:

1. How difficult is it typically for you to recall autobiographical memories?

2. How difficult is it typically for you to orient you spatially?

3. How difficult is it typically for you to use your imagination?

After a free explanation, participants were asked to rate the difficulty on a Likert-scale from 1

('very easy') to 6 ('very difficult').

Autobiographical memory fMRI task

The experimental fMRI task was adapted from a previous protocol by McCormick et al. (2015).

Two conditions, an AM retrieval task and a simple math task (MA), each consisting of 20

randomized trials, were included in this experiment. During AM trials, cue words, such as 'a

party', were presented on the screen for 12 s and participants were instructed to recall a personal

event related to the word cue which was specific in time and place (e.g., their 20th birthday party).

Participants were asked to press a response button once an AM was retrieved to indicate the

time point by which they would start to engage in the AM elaboration phase. For the rest of the

trial duration, participants were asked to re-experience the chosen AM and elaborate as many

details as possible in their mind's eye (elaboration period). After each AM trial, participants were

instructed to rate via button presses whether their retrieval had been vivid or faint. During MA

trials, simple addition or subtraction problems, for example, 47 + 19, were presented on the

screen for 12 s. Here, participants were instructed to press a response button once the problems

were solved and asked to engage in adding 3s to the solutions, for example, (47 + 19) + 3 + … +

3, until the trial ended. The MA trials were followed by a rating whether the MA problems had

been easy or difficult to solve. For both AM and MA, each trial lasted for 12 s, the maximum time

for rating of 3 s, and a jittered inter-stimulus interval (ISI) between 1 to 4 s. Since we were

especially interested in the elaboration phase of AM retrieval, for the fMRI analyses, we modelled

the last 8 s of each AM and MA trial just before the rating screen appeared.

MRI data acquisition

Anatomical and functional data were acquired at the German Center for Neurodegenerative

Diseases (DZNE), Bonn, Germany, using a 3 Tesla MAGNETOM Skyra MRI scanner (Siemens

Healthineers, Erlangen, Germany). A mirror was mounted on the 32 channel receiver head coil

and was placed in the scanner for the participants to view the stimuli shown on an MRI

conditional 30-inch TFT monitor (Medres medical research, Cologne, Germany) placed at the

scanner's rear end. The MRI protocol consisted of anatomical, resting-state, and AM task-based

fMRI scanning sessions. In addition, we acquired further experimental fMRI and DTI data which

are not part of this manuscript. Of note, the resting-state scans were acquired before participants

engaged in the AM task in order to prevent reminiscing about personal memories during the

resting-state. For the anatomical scans, an in-house developed 0.8 mm isotropic whole-brain T1-

weighted sagittal oriented multi-echo magnetization prepared rapid acquisition gradient echo

(MEMPRAGE; Brenner et al., 2014) was employed with the following parameters: TR = 2.56 s, TEs

= 1.72/3.44/5.16/6.88 ms, TA = 6:48, matrix = 320 x 320 x 224, readout pixel bandwidth = 680

Hz/Pixel, CAIPIRINHA mode. Resting-state (190 volumes, TA = 7 min) and AM task-based fMRI

scans (460 volumes, TA = 15 min) were acquired using an interleaved multi-slice 3.5 mm isotropic

echo planar imaging (EPI) sequence with TR = 2 s, TE = 30 ms, matrix = 64 x 64 x 39, readout pixel

bandwidth = 2112 Hz/Pixel (see Jessen et al., 2018). The images were obtained in an oblique-axial

slice orientation along the anterior-posterior commissure line. During resting-state, the

participants were asked to close their eyes and to think about nothing at all. The first 5 frames of

each functional session were excluded for the scanner to reach equilibrium. Before each

functional session an optimal B0 shim was determined and individually mapped by 2-echo

gradient echo (GRE) with same voxel resolution and positioning for later post-processing.

Manual segmentation of the hippocampus

Since our main goal was to assess hippocampal involvement during AM retrieval in aphantasia,

we sought to examine in depth whether there were any group differences in hippocampal

activation in respect to the hemispheric laterality or along its long-axis. For this reason, we

segmented the hippocampus based on the T1 structural images using ITKSnap (www.itksnap.org,

Version 3.8). Although we did not segment specific hippocampal subfields, our masks included

the dentate gyrus, CA1-4, subiculum and pre- and parasubiculum. Whole masks of the left and

right hippocampus were segmented manually in their respective native space and were divided

afterwards into anterior and posterior portions, using the location of the uncus as boundary.

fMRI preprocessing

SPM12 (Statistical Parametric Mapping 12) software package (www.fil.ion.ucl.ac.uk/spm/) on

MATLAB v19a (MathWorks) computing platform (https://matlab.mathworks.com/) was used to

perform resting-state and AM task-based fMRI data preprocessing. The anatomical T1w RMS of

all MEMPRAGE's echoes and functional 2D-EPI images were reoriented along the anterior-

posterior commissure axis. The phase and magnitude images within the field maps were applied

to calculate the voxel displacement maps (VDM) for geometrical correction of the distorted EPI

images. The echo times were set to 4.92 ms (short) and 7.38 ms (long). The total EPI readout time

was 34.56 ms. The calculated EPI and VDMs were applied to the functional scans for realignment

and unwarping. The functional scans were then co-registered to the segmented bias corrected

T1 scans.

Whole-brain differences between groups were evaluated. Thus, co-registered scans were

normalized to the Montreal Neurological Institute (MNI) space and a Gaussian smoothing kernel

of 8 mm FWHM was applied. In addition, for functional connectivity analyses, denoising was

applied using a linear regression model of potential confounding effects (global white matter

signal, CSF signal, and ART scrubbing parameters) in the BOLD signal using CONN software

package v20.b (www.nitrc.org/projects/conn/). Temporal band pass filter was set from 0.01 to

infinite to further minimize artifacts.

Statistical analyses

Behavioral analyses

Two samples t-tests were calculated to assess differences in the priming scores of aphantasics

and controls in the binocular rivalry task. One sample t-tests were used to distinguish the

performances of both groups from chance. To assess differences of Autobiographical Interview

scores between aphantasics and controls, a 2x2x2 ANOVA with post-hoc t-tests were calculated

with type of memory details (internal vs. external) and memory recency (remote vs. recent) as

within-subject factors and group (aphantasics vs. controls) as between-subject factor.

Afterwards, Bonferroni-corrected t-tests were conducted for specific internal (time, place,

internal event, perception, emotion) and external (external event, semantic, repetition, other)

memory details. Differences in memory ratings (confidence, episodic richness), self-reported

visualization scores, debriefing questions, and behavioral responses during fMRI scanning were

also assessed via two sample t-tests.

Hippocampal activity associated with autobiographical memory in native space

In order to examine hippocampal activity associated with autobiographical memory, we

extracted signal intensity values for both AM and MA trials for each participant for our manually

segmented anatomical masks of the left and right, anterior and posterior portions of the

hippocampus using the MATLAB-based Response Exploration toolbox (REX;

www.nitrc.org/projects/rex/). We then calculated for each participant for each anatomical mask

the difference between AM and MA signal intensities. Afterwards, group differences between

aphantasics and controls with respect to the laterality effects and effects between the anterior

and posterior hippocampus were assessed using a 2-way ANOVA with a post-hoc Tukey's multiple

comparison test, applying a significance threshold of α = .05.

Whole-brain fMRI activation analyses

After focusing on the hippocampus, we examined group differences of whole-brain activation

associated with AM and MA following the standard GLM procedure in SPM12. Owing to the

prominence of mental imagery during the elaboration phase of AM retrieval, we analyzed the

last 8 s of the AM and MA trials prior to the display of the vividness rating. These trials were

modelled as mini blocks in the GLM with motion correction regressors included as covariate of

no interest. We specified our main contrast of interest, i.e., AM versus MA on the first level, which

was then brought to the second group level using a one-sample t-test. Finally, the activation maps

of the two groups were compared using a two-sample t-test. For whole-brain analysis, we applied

a significance threshold of $p < .001$, and voxel cluster size of 10 adjacent voxels, uncorrected.

ROI-to-ROI functional connectivity analyses

One of our main a priori hypotheses stated that the hippocampus and the visual-perceptual

cortex show differential engagement during AM retrieval associated with aphantasia which was

confirmed by our whole-brain activation analyses. Based on these results, we sought to examine

the functional connectivity between those two areas. Towards this end, we created regions of

interest (ROIs, spheres with a diameter of 10 mm consisting of 536 voxels) around the three peaks

of the activation differences, following the whole-brain fMRI activation analyses, using the

MarsBaR HBM toolbox (Brett et al., 2002). The ROIs comprised (1) the right hippocampus, MNI:

x = 39, y = -31, z = -13, (2) the right visual cortex, MNI: x = 12, y = -79, z = 5, and (3) the left visual

cortex, MNI: x = -9, y = -76, z = 29. Using CONN, we examined functional connectivity (i.e.,

Generalized Psycho-Physiological Interactions, weighted general linear model with bivariate

correlation) between the hippocampal ROI and the ROIs situated in the visual-perceptual cortex

during AM task-based fMRI and during resting-state. Furthermore, in order to examine how well

functional connectivity between hippocampus and the visual cortex reflected an individuals'

ability to visualize mental events, we examined a regression model with the visualization scores

of the AI as criterion and resting state connectivity values, group allocation and the interaction

term of the connectivity values and group allocation as predictors. For these a priori driven

analyses, we applied a significance threshold of $p < .05$, small volume corrected, and a voxel

cluster threshold of 10 adjacent voxels.

329   **Results**

VVIQ and binocular rivalry task

Aphantasics (M = 16.57, SD = 1.02), scored significantly lower on the VVIQ than controls (M =

62.94, SD = 8.71), $t(15.47)$ = 21.12, $p < .001$, d = 7.23. Furthermore, aphantasics and controls

differed in the priming score of the binocular rivalry task, $t(18.04)$ = 2.41, p = .027, d = 0.87. While

controls were primed by their own mental imagery in 61.3 % (SD = 13.1 %) of the trials,

aphantasics were only primed 52.6 % (SD = 4.9 %) of the time. In fact, the performance of controls

differed significantly from chance, $t(14)$ = 3.34, p = .005, d = 0.86, whereas performance of

aphantasics did not, $t(13)$ = 1.96, p = .072. Moreover, the VVIQ scores correlated positively with

the performance on the binocular rivalry task, $r(28)$ = .43, p = .022. For the mock trials, no

significant differences in priming scores were found between groups, $t(28)$ = 0.86, p = .396, or

related to chance (aphantasics: $t(13)$ = 0.74, p = .475; controls: $t(15)$ = 0.42, p = .682). These

findings validate our groups by indicating that visual imagery strength was diminished in

aphantasics.

Autobiographical Interview

Regarding the AI, we found significant main effects of memory period, $F(1, 27) = 11.88$, $p = .002$,

$\eta_p^2 = .31$, type of memory details, $F(1, 27) = 189.03$, $p < .001$, $\eta_p^2 = .88$, and group, $F(1, 27) = 9.98$,

$p = .004$, $\eta_p^2 = .27$. On average, aphantasics ($M = 26.29$, $SD = 9.58$) described less memory details

than controls ($M = 38.36$, $SD = 10.99$), more details were reported for recent ($M = 35.17$, $SD =$

14.19) than remote memories ($M = 29.06$, $SD = 11.12$), and internal details ($M = 43.59$, $SD = 17.91$)

were reported more often than external details ($M = 20.64$, $SD = 8.94$). More importantly, a 2-

way interaction was found between type of memory details and group, $F(1, 27) = 54.09$, $p < .001$,

$\eta_p^2 = .67$, indicating that aphantasics reported significantly less internal memory details, $t(27) =$

5.07, $p < .001$, $d = 1.83$, but not significantly less external memory details, $t(27) = 0.13$, $p = .898$,

compared to controls (see Figure 1b). No 2-way interaction between memory period and group,

$F(1, 27) = 0.62$, $p = .439$, and no 3-way interaction between memory period, group and type of

memory details was found, $F(1, 27) = 3.87$, $p = .060$.

Based on the interaction effect between group and type of memory details, we compared

specific categories of internal and external memory details between the groups. For internal

details and in comparison to controls, aphantasics reported less internal events, $t(27) = 3.22$, $p =$

.016, $d = 1.17$, less emotional details, $t(27) = 4.40$, $p < .001$, $d = 1.59$, less perceptual details, $t(27)$

$= 4.95$, $p < .001$, $d = 1.79$, and less details regarding time, $t(27) = 5.27$, $p < .001$, $d = 1.90$, and

place, $t(27) = 3.31$, $p < .013$, $d = 1.20$ (see Figure 1c). On the other hand, no significant differences

were found for external details, including external events, $t(27) = 0.71$, $p > .999$, semantic details,

$t(27) = 0.02$, $p > .999$, repetition, $t(27) = 0.46$, $p > .999$, and other details, $t(27) = 0.45$, $p > .999$

(see Figure 1C). Regarding the rating scales, we found that aphantasics showed less episodic

richness, $t(27) = 7.50$, $p < .001$, $d = 2.71$, and less memory confidence, $t(27) = 5.85$, $p < .001$, $d = $

2.11 (see Figure 1a) as well as lower self-reported visualization scores, $t(27) = 11.92$, $p < .001$, d

$= 4.30$, than controls.

Figure 1. AM deficits associated with aphantasia. (A) Mean amount (± SEM) of episodic richness

and confidence in the Autobiographical Interview for controls and aphantasics. (B) Mean amount

(± SEM) of internal details and external details for recent and remote memories. (C) Mean

amount (± SEM) of specific internal and external memory details for aphantasics and controls.

* $p < .05$, ** $p < .01$, *** $p < .001$, **** $p < .0001$, n.s. = non-significant.

Debriefing questions

There were stark differences between the groups in how they answered our debriefing questions.

While aphantasics reported that they typically have great difficulty to recall autobiographical

memories, $t(27) = 6.20$, $p < .001$, $d = 2.31$, and to use their imagination in daily life, $t(24) = 10.18$,

$p < .001$, $d = 3.93$, they did not report difficulties in spatial orientation, $t(27) = 0.62$, $p = .541$, $d =$

0.23.

Behavioral results of the fMRI AM task

We found stark differences for the vividness response between groups, $t(28) = 5.29$, $p < .001$.

While controls reported in 86 % ($SD = 26$ %) of trials that their AM retrieval had been vivid,

aphantasics indicated only in 20 % ($SD = 20$ %) of trials that their AM retrieval had been vivid.

Moreover, aphantasics responded slower ($M = 1.34$ s, $SD = 0.38$ s) than controls ($M = 1.00$ s, SD

$= 0.29$ s) when they were asked if their retrieved memories were vivid or faint, $t(28) = 2.78$, $p =$

.009. In contrast, there were no significant differences between groups during the MA trials,

neither on the easy/hard response, $t(28) = 1.16$, $p = .255$, nor on the reaction times, $t(28) = 0.58$,

$p = .567$.

In addition, aphantasics and controls spent equal amounts of time to search for a memory

in AM trials, $t(19) = 1.03$, $p = .315$. On average, aphantasics spent 3.42 s ($SD = 0.74$ s) and controls

spent 3.15 s ($SD = 0.48$ s). Furthermore, both groups did not differ in their speed to solve the

math problems, $t(23) = 0.09$, $p = .926$. Aphantasics spent 3.87 s ($SD = 0.97$ s) to solve a math

problem and controls spent 3.90 s ($SD = 0.72$ s). For the button press, there were 9 % missing

values in AM trials and 7 % missing values in MA trials with no significant differences of missing

values between groups, neither for AM trials, $t(19.98) = 1.11$, $p = .281$, nor for MA trials, $t(18.13)$

$= 0.52$, $p = .609$.

Native space differences in hippocampal activation during AM retrieval

First, we sought to examine the hippocampal activation associated with AM in aphantasics and

controls (see Figure 2). The extracted fMRI signals illustrated stark group differences in

hippocampal AM-associated activation, $F(17, 252) = 3.03$, $p < .001$. Aphantasics showed reduced

activation of bilateral hippocampi, including the left anterior ($p = .033$), left posterior ($p = .027$),

right anterior ($p = .047$), and right posterior hippocampus ($p = .025$). There was no laterality effect

nor differences along the pattern of activation down the anterior-posterior axis between the

groups (all $p > .05$). These findings indicate that the behavioral AM deficit associated with

aphantasia is reflected neurally by a reduced bilateral hippocampal activation.

Figure 2. Reduced hippocampal activity during AM retrieval associated with aphantasia. The

signal intensities during AM and MA were extracted from anatomical hippocampal masks created

from each individual participant. (A) An example of a 3D reconstruction of the hippocampus,

separated into anterior and posterior portions for the left hippocampus. (B) The comparison

between the percentage of signal change during the AM and MA tasks in the hippocampus of

aphantasics and controls. Aphantasics show reduced differentiation between AM and MA than

controls in all portions of the hippocampus. * p < .05.

Activation patterns associated with AM retrieval

Second, we examined whole-brain activation during AM retrieval of both groups and the results

are displayed in Figure 3a and b. Additionally, the peak coordinates of AM and MA activation for

aphantasics and controls are shown in Table 1 and 2, respectively.

Overall, both groups showed greater activation in all areas typically associated with AM,

including bilateral hippocampus, vmPFC, and medial/lateral parietal regions, during AM retrieval.

When examining the group differences, aphantasics displayed greater activation in bilateral

visual-perceptual regions (maximum in lingual gyrus) in the occipital lobe than controls, $t(28) =$

4.41, $p < .001$ (MNI: right visual cortex: x = 12, y = -79, z = 5; left visual cortex: x = -9, y = -76, z =

29, see Figure 3c, d, and e). In contrast, controls showed greater activation in the right

hippocampus than aphantasics, $t(28) = 3.77$, $p < .001$ (MNI: x = 39, y = -31, z = -13). An additional

correlational analysis revealed that those participants with higher visual-perceptual cortex

activation had less hippocampal activation, $r(28) = -.39$, $p = .041$, indicating that there was a

trade-off between increased visual-perceptual cortex activation and decreased hippocampal

activation.

Figure 3. **Activation during AM retrieval task.** (A) Stronger activated cortical regions during AM

retrieval (in warm colors) in comparison to math problem solving (in cool colors) in aphantasics

and (B) controls. (C) Aphantasics showed greater activation in visual-perceptual cortices than

controls, and (D) controls showed stronger activation in the right posterior hippocampus than

aphantasics. Images are thresholded at $p < .001$, cluster size 10, uncorrected, except (D) which is

thresholded at $p < .01$, cluster size 10, for display purposes. (E) The percentage of signal change

for the contrast AM versus MA were extracted from the peaks of activated voxels, each with 1

mm sphere for display purposes. * $p < .05$

Table 1. Peak coordinates of the AM and MA activation for Aphantasia.

Region	Hemisphere	MNI Coordinates			Voxels	T-value
		X	Y	Z		
Activation AM > MA						
Posterior Cingulate Gyrus	Right	18	-57	11	4657	11.00
Parahippocampual Gyrus*	Left	-21	-31	-13		9.06
Hippocampus	Left	-27	-17	-19	205	8.40
Superior Frontal Gyrus	Left	-12	47	50	926	8.38
Angular Gyrus	Left	-42	-55	23	165	7.88
Lateral Orbitofrontal Cortex	Left	-42	38	-16	208	7.58
Hippocampus	Right	18	-37	-1	199	6.89
Cerebellum	Right	15	-79	-37	109	6.33
Brainstem	Right	3	-46	-52	43	6.03
Parahippocampual Gyrus*	Right	24	-31	-13		6.02
Middle Temporal Gyrus	Right	60	2	--19	76	5.27
Supramarginal Gyrus	Right	54	-58	32	26	4.99
Middle Frontal Gyrus	Left	-39	20	50	12	4.52
Activation MA > AM						
Precuneus	Left	-18	-58	41	594	-3.85
Inferior Temporal Gyrus	Right	51	-46	-13	123	-3.85
Precuneus	Right	24	-49	53	718	-3.85
Insula	Left	-30	23	11	48	-3.85
Inferior Temporal Gyrus	Left	-51	-49	-13	67	-3.86
Cerebellum	Right	30	-67	-52	27	-3.87
Middle Frontal Gyrus	Right	33	41	17	34	-3.87
Superior Frontal Gyrus	Right	30	5	59	52	-3.87
Inferior Frontal Gyrus	Right	54	14	29	35	-3.88
Insula	Right	39	11	8	51	-3.88
Inferior Frontal Gyrus	Left	-57	11	26	182	-3.88
Lateral Globus Pallidus	Right	23	-7	14	14	-3.92
Cerebellum	Left	-24	-64	-46	16	-3.93

451 *Sub-cluster level, Cluster size = 10 voxels, p-value = 0.001

454 **Table 2. Peak coordinates of the AM and MA activation for healthy controls.**

Region	Hemisphere	MNI Coordinates			Voxels	T-value
		X	Y	Z		
Activation AM > MA						
Parahippocampal Gyrus	Right	27	-28	-19	11319	12.41
Parahippocampal Gyrus*	Left	-24	-25	-16		9.01
Cerebellum	Left	-18	-76	-37	108	7.67
Anterior Cingulate	Right	9	35	11	13	7.01
Medial Frontal Gyrus	Right	18	32	29	233	6.93
Inferior Frontal Gyrus	Right	60	32	11	53	5.94
Hippocampus	Left	-36	-22	-16	252	5.64
Hippocampus	Right	27	-22	-16	233	5.28
Hypothalamus	Right	3	-4	-10	16	4.93
Activation MA > AM						
Post Central Gyrus	Left	-33	-43	62	643	-3.73
Precuneus	Right	21	-52	53	483	-3.74
Inferior Frontal Gyrus	Right	51	8	26	16	-3.74
Middle Occipital Gyrus	Right	33	-82	2	26	-3.75
Middle Temporal Gyrus	Left	-51	-58	-1	18	-3.76

455 *Sub-cluster level, Cluster size = 10 voxels, p-value = 0.001

Exploring functional connectivity of hippocampus and visual-perceptual cortices during AM

The whole-brain analyses strengthened our hypothesis that a core difference between

aphantasics and controls lies in the interplay between the visual-perceptual cortex and the

hippocampus. To test this interplay, in a third step, we examined functional connectivity between

the peak differences of the hippocampus and the visual-perceptual cortex during AM retrieval

(see Figure 4). We found a group difference in functional connectivity between the right

hippocampus and left visual-perceptual cortices, $t(28) = 2.65$, $p = .006$. Interestingly, while

aphantasics show almost no functional connectivity between those two ROIs, controls displayed

a strong negative connectivity between the two locations, $p = .013$.

Figure 4. Functional connectivity between the visual-perceptual cortex and hippocampus

during AM retrieval. (A) During AM retrieval, group differences in functional connectivity

amongst the ROIs were only found between the right hippocampus, and left visual-perceptual

cortices. (B) Controls displayed a stark negative correlation, whereas aphantasics did not.

Exploring functional connectivity of hippocampus and visual-perceptual cortices during resting-

state

Lastly, we examined resting-state connectivity between our identified ROIs in the right posterior

hippocampus and left and right visual-perceptual cortex. There was no group difference during

resting-state between these ROIs. In order to examine whether resting-state functional

connectivity carried information about one's ability to visualize AMs, we added the visualization

scores as a regressor of interest in our model. While connectivity alone did not predict the

visualization scores in the interview, $F(14, 40) = 1.651$, $p = .391$, $\beta = -.06$, we found a main effect

of group, $F(3, 40) = 353.2$, $p < .001$, $\beta = .92$, and an interaction between group and connectivity,

$F(3, 54) = 305.1$, $p < .001$, $\beta = .26$. Interestingly, for controls, we found a positive correlation

between the resting-state connectivity of the right hippocampus and the visual cortex and the

visualization scores from the interview, $r(13) = .65$, $p = .011$ (see Figure 5). On the other hand, for

aphantasics, we found a negative correlation between the resting-state connectivity of the right

hippocampus and the visual cortex and the visualization scores from the interview, *r*(14) = -.57,

p = .027.

In sum, our fMRI results indicate that the impaired AM retrieval associated with

aphantasia is reflected by functional alterations of the hippocampus and visual-perceptual

cortex, as well as the interaction between them.

Figure 5. Functional connectivity between the visual-perceptual cortex and hippocampus

during resting-state explains visualization abilities. Resting-state functional connectivity

between the right hippocampus and the right visual-perceptual cortex correlates with

visualization abilities*.* Fitted straight lines indicate a negative correlation for aphantasics (red)

and a positive correlation for controls (blue).

Discussion

In this study, we set out to examine the neural correlates of episodic AM retrieval in aphantasia

as a way to examine the influence of visual imagery on episodic AM. In line with previous reports,

we found that aphantasics reported less sensory details during AM retrieval regardless of the

recency of memory (cf. Dawes et al., 2020; 2022; Milton et al., 2020; Zeman et al., 2020).

Strikingly, the deficit in constructing visual imagery associated with aphantasia did not only lead

to a reduced retrieval of visual-perceptual details but to a broader impairment in retrieving

episodic AMs, including reduced emotions and confidence attached to the memories. Thus, in

agreement with a recent account of aphantasia (Blomkvist, 2023), our results support the idea

that a diminished construction of visual details during AM retrieval leads to a more general

episodic memory deficit. We expand the current knowledge by adding that this AM deficit is

reflected neurally by an altered activation and connectivity pattern between the hippocampus

and visual-perceptual cortices. Our findings provide novel insights into three current debates: 1)

the mechanisms of aphantasia-related AM deficits, 2) the similarities and differences between

aphantasics and individuals with hippocampal damage, and 3) the neural models of AM.

Potential mechanisms underlying aphantasia-related AM deficits

We report that aphantasics show increased activation of bilateral visual-perceptual cortices

as well as decreased hippocampal activation during AM retrieval in comparison to controls.

Increased activity in the visual-perceptual cortices in aphantasics has been reported previously,

albeit not associated with AM (Fulford et al., 2018; Keogh et al., 2020). A prominent hypothesis

states that this heightened activity of the visual-perceptual cortices hinders aphantasics to detect

small imagery-related signals (Pearson, 2019). Our findings support this hypothesis, indicating a

more generally heightened state of arousal rather than a memory-specific reaction.

We had the a priori hypothesis that hippocampal activation will be decreased in

aphantasics; a hypothesis which we could confirm in our native space hippocampal analysis. On

the whole-brain level, perhaps due to our small sample size, only the cluster of activation group

differences in the right posterior hippocampus survived the statistical threshold. Given the low

power, further studies are needed to confirm this effect; however, the right posterior

hippocampus interacts in the healthy brain heavily with the visual-perceptual cortex only during

the elaboration phase of AM retrieval (McCormick et al. 2015).

In addition, controls exhibited a strong functional connection between both brain

structures during AM retrieval and this functional connectivity predicted better visualization

skills. At first glance, it is surprising that this functional connectivity was negative. However,

negative visual-perceptual cortex activation during perceiving and imagining scenes has been

reported before (McCormick et al. 2021). One possible explanation might be that signals from

the hippocampus selectively inhibit imagery-irrelevant activation in the visual-perceptual

cortices (e.g., sensory noise) to carve out imagery-related signals (cf. Pace et al., 2023). This would

be in line with the hypothesis stated above, that a bad signal-to-noise ratio in the visual cortex

hiders aphantasics to create mental imagery. Either way, the described functional connectivity

between the hippocampus and the visual-perceptual cortex fits well to previous neuroimaging

studies pointing towards a central role of the dynamic interplay between these brain structures

during AM retrieval (McCormick, et al. 2015). This interplay seems to be especially important

during the elaboration stage of AM retrieval, a period when specific visual-perceptual details are

being actively brought back into the mind's eye. At this point, however, it remains unclear

whether the disruption of AM elaboration associated with aphantasia takes place during the

encoding, storage, or retrieval process.

From a theoretical point of view, the extended constructive episodic simulation hypothesis

proposes a top-down hierarchy during mental imagery (Blomkvist, 2023). In this model, the

hippocampus initiates retrieval processes in primary sensory brain regions, such as the visual-

perceptual cortex in order to retrieve visual-perceptual details associated with a specific AM.

Evidence for such top-down hierarchies during mental imagery have been observed in fronto-

parietal and occipital networks via effective connectivity analyses, such as Granger Causality and

Dynamic Causal Modelling (Dentico et al., 2014; Dijkstra et al., 2017; Mechelli et al., 2004). For

example, intracranial and high-density scalp electroencephalography (EEG) provided evidence of

high frequency hippocampal signaling during the recall of perceptual cues in patients with

epilepsy, indicating that the hippocampus drives the switch from perception to memory recalling

(Treder et al., 2021). In aphantasia, it is hypothesized that this top-down hierarchy is disrupted

and therefore, the hippocampus can no longer initiate the retrieval and incorporation of visual-

perceptual details in one coherent mental event. Because of the slow temporal resolution of our

fMRI sequence, our data cannot directly speak to the question of temporal directionality

between the hippocampus and visual-perceptual cortex. Nonetheless, our findings suggest that

the bidirectional connectivity between both brain structures is crucial for the re-experience of

episodic AMs. As such, hippocampal processes may be needed to retrieve specific details and if

these details are not provided by the visual-perceptual cortices, the entire episodic AM retrieval

seems to fail.

Similarities and differences between aphantasics and individuals with hippocampal lesions

At face value, the episodic AM deficits in aphantasia in our data and reported previously (Dawes

et al., 2020; 2022; Milton et al., 2020; Zeman et al., 2020), as well as the decreased hippocampal

activation during AM retrieval suggest that aphantasia is a selective episodic memory condition,

similar to AM amnesia known from individuals with hippocampal damage. In fact, previous and

the current study show that aphantasics and individuals with hippocampal damage report less

internal details across several memory detail subcategories (Rosenbaum et al., 2008; St-Laurent

et al., 2009; Steinvorth et al., 2005) and these deficits can be observed regardless of the recency

of the memory (Miller et al., 2020). These similarities suggest that aphantasics are not merely

missing the visual-perceptual details to specific AM, but they have a profound deficit associated

with the retrieval of AM.

Nonetheless, there are also stark differences between aphantasics and individuals with

hippocampal damage. Foremost, aphantasics seem not to have difficulties to retrieve spatial

information (Blomkvist, 2023), which is another inherent function of the hippocampus (Burgess

et al., 2002; O'Keefe, 1991). In the current study, we did not set out to examine spatial cognition

in aphantasics, however, parts of our data speak to this aspect. While in our study aphantasics

reported less amount of spatial details during the AI, this standard scoring procedure only counts

place details when the exact place is recalled and is not meant to assess the recall of spatial layout

(Levine et al. 2002). Thus, this place score may not represent spatial cognition per se. In fact,

when asking aphantasics about their experience, they point out difficulties in recalling AM and

using imagination in daily life, however they report no difficulties in spatial orientation. Indeed,

often during the interview, aphantasics would explain that they know how the space around

them felt, they just cannot see it in front of their mind's eye. One aphantasic put her finger on it,

describing it as: "I can put my consciousness in my kitchen at home and feel all around but there

is no visual image attached to this feeling." These observations support the idea that some

hippocampal processes, at least regarding spatial navigation, may be intact in people with

aphantasia. However, spatial cognition should be formally addressed in future studies. One way

to assess this hippocampal function would be to examine tasks, which rely on scene construction.

The scene construction theory states that the hippocampus is crucially needed for the

construction of spatially coherent mental models of scenes (Maguire & Mullally, 2013). For

example, patients with hippocampal damage cannot imagine the spatial layout of fictitious

scenes (Hassabis et al., 2007), they detect less errors in spatially-incoherent scenes than controls

(McCormick et al., 2017), and they show less scene-dependent mind-wandering episodes

(McCormick, Rosenthal, et al., 2018). In contrast, we would predict that aphantasics have

diminutive deficits in tasks that depend on hippocampal scene construction processes.

What could be impaired in aphantasics are all cognitive functions which rely on the

population of the constructed scenes with visual-perceptual details, such as episodic AM

retrieval, episodic future thinking, complex decision-making, and complex empathy tasks. This is

in line with work from Bergmann and Ortiz-Tudela (2023), who propose that 'the memory deficits

observed in aphantasia might be better understood as a difference in high-level feedback

processing along the 'what' stream, rather than an episodic memory impairment'.

Towards a novel neural model of autobiographical memory

While more research is required exploring the cognitive landscape associated with aphantasia,

such as spatial cognition and scene construction, our data contribute to an old debate of how AM

retrieval and visual imagery are intertwined. We propose that the hippocampus is embedded in

a brain-wide network, comprising the vmPFC and visual-perceptual cortices, in which each of

these nodes contributes specific processes to the re-construction of extended detail-rich mental

events (see also Ciaramelli et al., 2019; McCormick, Ciaramelli, et al., 2018). Within this model,

the vmPFC initiates and overseas the scene construction process which takes place in the

hippocampus. Further, the visual-perceptual cortex provides the visual details which are essential

to populate the hippocampally constructed scenes. This model is backed up by a previous MEG

study revealing that the vmPFC directs hippocampal activity during the initiation of AM retrieval

(McCormick et al., 2020). This finding has been replicated and extended by Chen et al. (2021),

showing that the vmPFC leads hippocampal involvement during scene construction and other

scene-based processes (Monk et al., 2021). Moreover, the connection between the hippocampus

and the visual-perceptual cortex seems equally crucial. There are a few case reports of damage

to the occipital cortex causing AM amnesia (Greenberg et al., 2005), potentially by preventing

the population of the hippocampally constructed scenes. Furthermore, our current study

suggests that a reliable connectivity between the hippocampus and the visual-perceptual cortices

is essential to provide the visual details necessary for successful vivid, detail-rich AM retrieval.

Conclusion

Aphantasia provides a natural knock-out model for the influence of visual imagery on different

cognitive functions. We here report a tight link between visual imagery and our ability to retrieve

vivid and detail-rich personal past events, as aphantasics do not only report fewer visual-

perceptual details during episodic AM retrieval but also show decreased confidence and

emotionality associated with these memories. In this context, we highlight the central role of the

functional connectivity between the hippocampus and occipital cortex to assemble visual-

perceptual details into one coherent extended mental event. Exciting novel research avenues will

be to examine hippocampal-dependent spatial cognition in aphantasics and to investigate

whether neuroscientific interventions can be used to enhance AM retrieval by enhancing visual

imagery.

References

Addis, D. R., Moscovitch, M., Crawley, A. P., & McAndrews, M. P. (2004). Recollective qualities

modulate hippocampal activation during autobiographical memory retrieval.

Hippocampus, *14*(6), 752–762.

Albers, A. M., Kok, P., Toni, I., Dijkerman, H. C., & de Lange, F. P. (2013). Shared representations

for working memory and mental imagery in early visual cortex. *Current Biology*, *23*(15),

1427–1431. https://doi.org/10.1016/j.cub.2013.05.065

Baudson, T. G., & Preckel, F. (2015). mini-q: Intelligenzscreening in drei Minuten. *Diagnostica*.

Bauer, P. J., Pathman, T., Inman, C., Campanella, C., & Hamann, S. (2017). Neural correlates of

autobiographical memory retrieval in children and adults. *Memory*, *25*(4), 450–466.

https://doi.org/10.1080/09658211.2016.1186699

Bergmann, J., & Ortiz-Tudela, J. (2023). Feedback signals in visual cortex during episodic and

schematic memory retrieval and their potential implications for aphantasia. *Neuroscience*

and Biobehavioral Reviews, *152*, 105335.

https://doi.org/10.1016/j.neubiorev.2023.105335

Blomkvist, A. (2023). Aphantasia: In search of a theory. *Mind and Language*, *38*(3), 866–888.

https://doi.org/10.1111/mila.12432

Brenner, D., Stirnberg, R., Pracht, E. D., & Stöcker, T. (2014). Two-dimensional accelerated MP-

RAGE imaging with flexible linear reordering. *Magnetic Resonance Materials in Physics,*

Biology and Medicine, *27*(5), 455–462. https://doi.org/10.1007/S10334-014-0430-

Y/TABLES/2

Brett, M., Anton, J.-L., Valabregue, R., Poline, J.-B., & others. (2002). Region of interest analysis

using an SPM toolbox. *8th International Conference on Functional Mapping of the Human*

Brain, *16*(2), 497.

Brown, T. I., Rissman, J., Chow, T. E., Uncapher, M. R., & Wagner, A. D. (2018). Differential

Medial Temporal Lobe and Parietal Cortical Contributions to Real-world Autobiographical

Episodic and Autobiographical Semantic Memory. *Scientific Reports*, *8*(1), 1–14.

https://doi.org/10.1038/s41598-018-24549-y

Burgess, N., Maguire, E. A., & O'Keefe, J. (2002). The human hippocampus and spatial and

episodic memory. *Neuron*, *35*(4), 625–641.

Burianova, H., McIntosh, A. R., & Grady, C. L. (2010). A common functional brain network for

autobiographical, episodic, and semantic memory retrieval. *NeuroImage*, *49*(1), 865–874.

https://doi.org/10.1016/j.neuroimage.2009.08.066

Chen, D., Kunz, L., Lv, P., Zhang, H., Zhou, W., Liang, S., Axmacher, N., & Wang, L. (2021). Theta

oscillations coordinate grid-like representations between ventromedial prefrontal and

entorhinal cortex. *Science Advances*, *7*(44), eabj0200.

Ciaramelli, E., De Luca, F., Monk, A. M., McCormick, C., & Maguire, E. A. (2019). What" wins" in

VMPFC: Scenes, situations, or schema? *Neuroscience & Biobehavioral Reviews*, *100*, 208–

210.

D'Argembeau, A., & Van der Linden, M. (2006). Individual differences in the phenomenology of

mental time travel: the effect of vivid visual imagery and emotion regulation strategies.

Consciousness and Cognition, *15*(2), 342–350.

https://doi.org/10.1016/j.concog.2005.09.001

Danker, J. F., & Anderson, J. R. (2010). The ghosts of brain states past: remembering reactivates

the brain regions engaged during encoding. *Psychological Bulletin*, *136*(1), 87.

Dawes, A. J., Keogh, R., Andrillon, T., & Pearson, J. (2020). A cognitive profile of multi-sensory

imagery, memory and dreaming in aphantasia. *Scientific Reports*, *10*(1), 10022.

https://doi.org/10.1038/s41598-020-65705-7

Dawes, A. J., Keogh, R., Robuck, S., & Pearson, J. (2022). Memories with a blind mind:

Remembering the past and imagining the future with aphantasia. *Cognition*, *227*, 105192.

https://doi.org/10.1016/J.COGNITION.2022.105192

Dentico, D., Cheung, B. L., Chang, J. Y., Guokas, J., Boly, M., Tononi, G., & Van Veen, B. (2014).

Reversal of cortical information flow during visual imagery as compared to visual

perception. *NeuroImage*, *100*, 237–243.

https://doi.org/10.1016/j.neuroimage.2014.05.081

Dijkstra, N., Zeidman, P., Ondobaka, S., Van Gerven, M. A. J., & Friston, K. (2017). Distinct Top-

down and Bottom-up Brain Connectivity during Visual Perception and Imagery. *Scientific*

Reports, *7*(1), 1–9. https://doi.org/10.1038/s41598-017-05888-8

Fulford, J., Milton, F., Salas, D., Smith, A., Simler, A., Winlove, C., & Zeman, A. (2018). The neural

correlates of visual imagery vividness – An fMRI study and literature review. *Cortex*, *105*,

26–40. https://doi.org/10.1016/j.cortex.2017.09.014

Greenberg, D. L., Eacott, M., Brechin, D., & Rubin, D. (2005). Visual memory loss and

autobiographical amnesia: a case study. *Neuropsychologia*, *43*(10), 1493–1502.

https://doi.org/10.1016/j.neuropsychologia.2004.12.009

Greenberg, D. L., & Knowlton, B. J. (2014). The role of visual imagery in autobiographical

memory. *Memory & Cognition*, *42*(6), 922–934. https://doi.org/10.3758/s13421-014-0402-

5

Hassabis, D., Kumaran, D., Vann, S. D., & Maguire, E. A. (2007). Patients with hippocampal

amnesia cannot imagine new experiences. *Proceedings of the National Academy of*

Sciences, *104*(5), 1726–1731.

Jessen, F., Spottke, A., Boecker, H., Brosseron, F., Buerger, K., Catak, C., Fliessbach, K., Franke,

C., Fuentes, M., Heneka, M. T., Janowitz, D., Kilimann, I., Laske, C., Menne, F., Nestor, P.,

Peters, O., Priller, J., Pross, V., Ramirez, A., … Düzel, E. (2018). Design and first baseline

data of the DZNE multicenter observational study on predementia Alzheimer's disease

(DELCODE). *Alzheimer's Research and Therapy*, *10*(1), 1–10.

https://doi.org/10.1186/S13195-017-0314-2/TABLES/1

Kay, L., Keogh, R., Andrillion, T., & Pearson, J. (2022). The pupillary light response as a

physiological index of aphantasia, sensory and phenomenological imagery strength. *ELife*,

11, e72484. https://doi.org/10.7554/eLife.72484

Keogh, R., Bergmann, J., & Pearson, J. (2020). Cortical excitability controls the strength of

mental imagery. *ELife*, *9*, e50232. https://doi.org/10.7554/eLife.50232

Keogh, R., & Pearson, J. (2018). The blind mind: No sensory visual imagery in aphantasia.

Cortex, *105*, 53–60. https://doi.org/10.1016/j.cortex.2017.10.012

Langille, J. J., & Gallistel, C. R. (2020). Locating the engram: Should we look for plastic synapses

or information-storing molecules? *Neurobiology of Learning and Memory*, *169*, 107164.

Levine, B., Svoboda, E., Hay, J. F., Winocur, G., & Moscovitch, M. (2002). Aging and

autobiographical memory: dissociating episodic from semantic retrieval. *Psychology and*

Aging, *17*(4), 677–689. https://doi.org/10.1037/0882-7974.17.4.677

Maguire, E. A., & Mullally, S. L. (2013). The hippocampus: a manifesto for change. *Journal of*

Experimental Psychology: General, *142*(4), 1180–1189. https://doi.org/10.1037/a0033650

Marks, D. F. (1973). Visual imagery differences in the recall of pictures. *British Journal of*

Psychology, *64*(1), 17–24. https://doi.org/10.1111/j.2044-8295.1973.tb01322.x

Marks, D. F. (1995). New directions for mental imagery research. *Journal of Mental Imagery*,

19(3–4), 153–167. https://psycnet.apa.org/record/1996-29150-001

McCormick, C., Barry, D. N., Jafarian, A., Barnes, G. R., & Maguire, E. A. (2020). VmPFC Drives

Hippocampal Processing during Autobiographical Memory Recall Regardless of

Remoteness. *Cerebral Cortex*, *30*(11), 5972–5987. https://doi.org/10.1093/cercor/bhaa172

McCormick, C., Ciaramelli, E., De Luca, F., & Maguire, E. A. (2018). Comparing and Contrasting

the Cognitive Effects of Hippocampal and Ventromedial Prefrontal Cortex Damage: A

Review of Human Lesion Studies. *Neuroscience*, *374*, 295–318.

https://doi.org/10.1016/j.neuroscience.2017.07.066

McCormick, C., Rosenthal, C. R., Miller, T. D., & Maguire, E. A. (2017). Deciding what is possible

and impossible following hippocampal damage in humans. *Hippocampus*, *27*(3), 303–314.

https://doi.org/10.1002/hipo.22694

McCormick, C., Rosenthal, C. R., Miller, T. D., & Maguire, E. A. (2018). Mind-wandering in

people with hippocampal damage. *Journal of Neuroscience*, *38*(11), 2745–2754.

https://doi.org/10.1523/JNEUROSCI.1812-17.2018

McCormick, C., St-Laurent, M., Ty, A., Valiante, T. A., & McAndrews, M. P. (2015). Functional

and effective hippocampal-neocortical connectivity during construction and elaboration of

autobiographical memory retrieval. *Cerebral Cortex*, *25*(5), 1297–1305.

https://doi.org/10.1093/cercor/bht324

Mechelli, A., Price, C. J., Friston, K. J., & Ishai, A. (2004). Where bottom-up meets top-down:

Neuronal interactions during perception and imagery. *Cerebral Cortex*, *14*(11), 1256–1265.

https://doi.org/10.1093/cercor/bhh087

Miller, T. D., Chong, T. T. J., Davies, A. M. A., Johnson, M. R., Irani, S. R., Husain, M., Ng, T. W. C.,

Jacob, S., Maddison, P., Kennard, C., & others. (2020). Human hippocampal CA3 damage

disrupts both recent and remote episodic memories. *Elife*, *9*, e41836.

Milton, F., Fulford, J., Dance, C., Gaddum, J., Heuerman-Williamson, B., Jones, K., Knight, K. F.,

MacKisack, M., Winlove, C., & Zeman, A. (2020). Behavioral and neural signatures of visual

imagery vividness extremes: aphantasia vs. hyperphantasia. *Cerebral Cortex*

Communications, *2*(2), tgab035. https://doi.org/10.31234/OSF.IO/J2ZPN

Monk, A. M., Dalton, M. A., Barnes, G. R., & Maguire, E. A. (2021). The role of hippocampal–

ventromedial prefrontal cortex neural dynamics in building mental representations.

Journal of Cognitive Neuroscience, *33*(1), 89–103.

Monzel, M., Keidel, K., & Reuter, M. (2021). Imagine, and you will find - Lack of attentional

guidance through visual imagery in aphantasics. *Attention, Perception {\&} Psychophysics*.

https://doi.org/10.3758/s13414-021-02307-z

Monzel, M., Mitchell, D., Macpherson, F., Pearson, J., & Zeman, A. (2022). Proposal for a

consistent definition of aphantasia and hyperphantasia: A response to Lambert and Sibley

(2022) and Simner and Dance (2022). *Cortex*, *152*, 74–76.

https://doi.org/10.1016/J.CORTEX.2022.04.003

Monzel, M., Vetterlein, A., & Reuter, M. (2021). Memory deficits in aphantasics are not

restricted to autobiographical memory – Perspectives from the Dual Coding Approach.

Journal of Neuropsychology, *16*(2), 444–461. https://doi.org/10.1111/jnp.12265

Moscovitch, M., Rosenbaum, R. S., Gilboa, A., Addis, D. R., Westmacott, R., Grady, C.,

McAndrews, M. P., Levine, B., Black, S., Winocur, G., & Nadel, L. (2005). Functional

neuroanatomy of remote episodic, semantic and spatial memory: A unified account based

on multiple trace theory. *Journal of Anatomy*, *207*(1), 35–66.

https://doi.org/10.1111/j.1469-7580.2005.00421.x

O'Keefe, J. (1991). *The hippocampal cognitive map and navigational strategies.*

Pace, T., Koenig-Robert, R., & Pearson, J. (2023). Different Mechanisms for Supporting Mental

Imagery and Perceptual Representations: Modulation Versus Excitation. *Psychological*

Science, *34*(11), 9567976231198436. https://doi.org/10.1177/09567976231198435

Pearson, J. (2019). The human imagination: the cognitive neuroscience of visual mental

imagery. *Nature Reviews Neuroscience*, *20*(10), 624–634. https://doi.org/10.1038/s41583-

019-0202-9

Pearson, J., Clifford, C. W. G., & Tong, F. (2008). The Functional Impact of Mental Imagery on

Conscious Perception. *Current Biology*, *18*(13), 982–986.

https://doi.org/10.1016/j.cub.2008.05.048

Pearson, J., Rademaker, R. L., & Tong, F. (2011). Evaluating the mind's eye: The metacognition

of visual imagery. *Psychological Science*, *22*(12), 1535–1542.

https://doi.org/10.1177/0956797611417134

Rosenbaum, R. S., Moscovitch, M., Foster, J. K., Schnyer, D. M., Gao, F., Kovacevic, N., Verfaellie,

M., Black, S. E., & Levine, B. (2008). Patterns of autobiographical memory loss in medial-

temporal lobe amnesic patients. *Journal of Cognitive Neuroscience*, *20*(8), 1490–1506.

Schwitzgebel, E. (2002). How well do we know our own conscious experience? The case of

visual imagery. *Journal of Consciousness Studies*, *9*(5–6), 35–53.

https://www.jstor.org/stable/43154688

Sheldon, S., & Levine, B. (2013). Same as it ever was: Vividness modulates the similarities and

differences between the neural networks that support retrieving remote and recent

autobiographical memories. *NeuroImage*, *83*, 880–891.

St-Laurent, M., Moscovitch, M., Levine, B., & McAndrews, M. P. (2009). Determinants of

autobiographical memory in patients with unilateral temporal lobe epilepsy or excisions.

Neuropsychologia, *47*(11), 2211–2221.

Steinvorth, S., Levine, B., & Corkin, S. (2005). Medial temporal lobe structures are needed to re-

experience remote autobiographical memories: evidence from HM and WR.

Neuropsychologia, *43*(4), 479–496.

Svoboda, E., McKinnon, M. C., & Levine, B. (2006). The functional neuroanatomy of

autobiographical memory: a meta-analysis. *Neuropsychologia*, *44*(12), 2189–2208.

https://doi.org/10.1016/j.neuropsychologia.2006.05.023

Teyler, T. J., & DiScenna, P. (1986). The Hippocampal Memory Indexing Theory. *Behavioral*

Neuroscience, *100*(2), 147–154. https://doi.org/10.1037/0735-7044.100.2.147

Wagner, S., & Monzel, M. (2023). Measuring imagery strength in schizophrenia: no evidence of

enhanced mental imagery priming. *Brain and Behavior*, *13*(9), e3146.

https://doi.org/10.1002/brb3.3146

World Medical Association. (2013). World Medical Association Declaration of Helsinki: Ehtical

principles for medical research involving human subjects. *JAMA*, *310*(20), 2191.

https://doi.org/10.1001/jama.2013.281053

Zeidman, P., & Maguire, E. A. (2016). Anterior hippocampus: the anatomy of perception,

imagination and episodic memory. *Nature Reviews Neuroscience*, *17*(3), 173–182.

Zeman, A., Dewar, M., & Della Sala, S. (2015). Lives without imagery – congenital aphantasia.

Cortex, *73*, 378–380. https://doi.org/10.1016/j.cortex.2015.05.019

Zeman, A., Milton, F., Della Sala, S., Dewar, M., Frayling, T., Gaddum, J., Hattersley, A.,

Heuerman-Williamson, B., Jones, K., MacKisack, M., & Winlove, C. (2020). Phantasia – the

psychological significance of lifelong visual imagery vividness extremes. *Cortex*, *130*, 426–

440. https://doi.org/10.1016/j.cortex.2020.04.003