Original Research:

**Hippocampal-occipital connectivity reflects autobiographical memory deficits**

**in aphantasia**

Merlin Monzel[1,2,*], Pitshaporn Leelaarporn[2,3,*], Teresa Lutz[2], Johannes Schultz[4,5], Sascha

Brunheim[3], Martin Reuter[1], and Cornelia McCormick[2,3]

*Both authors contributed equally.

1.  Department of Psychology, University of Bonn, 53113 Bonn, Germany

2.  German Center for Neurodegenerative Diseases, 53127 Bonn, Germany

3.  Department of Neurodegenerative Diseases and Geriatric Psychiatry, University of Bonn

Medical Center, 53127 Bonn, Germany

4.  Center for Economics and Neuroscience, University of Bonn, 53113 Bonn, Germany

5.  Institute of Experimental Epileptology and Cognition Research, Medical Faculty, University

of Bonn, 53127 Bonn, Germany

Corresponding authors: Cornelia McCormick (cornelia.mccormick@dzne.de), Merlin Monzel

(merlin.monzel@uni-bonn-diff.de)

**Conflict of interest statement:**

The authors declare no competing financial interest.

**Acknowledgments**

C.M. was supported by the German Research Foundation (DFG, MC244/3-1) and the DZNE

Foundation (DZNE Stiftung-Forschung für ein Leben ohne Demenz, Parkinson & ALS). We thank

Anke Rühling and Jennifer Schlee for their technical support during scanning and Yilmaz Sagik

for serving as the secondary scorer for the Autobiographical Interview. In addition, we would

like to thank all participants for their time and effort to share their memories with us.

**Abstract**

Aphantasia refers to reduced or absent visual imagery. While most of us can readily recall

decade-old personal experiences (autobiographical memories, AM) with vivid mental images,

there is a dearth of information about whether the loss of visual imagery in aphantasics affects

their AM retrieval. The hippocampus is thought to be a crucial hub in a brain-wide network

underlying AM. One important question is whether this network, especially the connectivity of

the hippocampus, is altered in aphantasia. In the current study, we tested 14 congenital

aphantasics and 16 demographically matched controls in an AM fMRI task to investigate how

key brain regions (i.e., hippocampus and visual-perceptual cortices) interact with each other

during AM re-experiencing. All participants were interviewed regarding their autobiographical

memory to examine their episodic and semantic recall of specific events. Aphantasics reported

more difficulties in recalling AM, were less confident about their memories, and described less

internal and emotional details than controls. Neurally, aphantasics displayed decreased

hippocampal and increased visual-perceptual cortex activation during AM retrieval compared

to controls. In addition, controls showed strong negative functional connectivity between the

hippocampus and the visual cortex during AM and resting-state functional connectivity

between these two brain structures predicted better visualization skills. Our results indicate

that visual mental imagery plays an important role in detail-rich vivid AM, and that this type of

cognitive function is supported by the functional connection between the hippocampus and the

visual-perceptual cortex.

**Keywords:** autobiographical retrieval, episodic memory, visual cortex, neural networks,

functional connectivity

## Introduction

Our unique and personal memories are stored in autobiographical memories (AM) providing

stability and continuity of our self (Svoboda et al., 2006). For most of us, travelling mentally

back in time and re-visiting such unique personal events is associated with vivid, detail-rich

mental imagery (D'Argembeau & Van der Linden, 2006; Greenberg & Knowlton, 2014). This

vivid mental imagery during the re-experiencing of AMs has become a hallmark of autonoetic,

episodic AM retrieval. However, up to date, it remains unclear to what extent episodic AM

retrieval depends on visual mental imagery and what neural consequences a lack of mental

imagery has on episodic AM retrieval. This knowledge gap exists because separating AM

retrieval from mental imagery is a complex and challenging task.

One way to address this conundrum is to study people with aphantasia (Zeman et al.,

2015). Recent research defines aphantasia as a neuropsychological difference in which people

experience a marked reduction or complete lack of voluntary sensory imagery (Monzel et al.,

2022). This state is associated with psychophysiological alterations, such as reduced imagery-

induced pupil contraction (Kay et al., 2022) and reduced imagery-induced priming effects

(Keogh & Pearson, 2018; Monzel, Keidel, et al., 2021). Thus, aphantasics offer the unique

opportunity to examine the consequences for episodic AM retrieval in the absence or marked

reduction of voluntary imagery. Indeed, a handful of previous studies report convergent

evidence that aphantasics report less sensory AM details than controls (Dawes et al., 2020,

2022; Milton et al., 2020; Zeman et al., 2020), which may also be less emotional (Monzel et al.,

2023; Wicken et al., 2021). Spatial accuracy, on the other hand, was not found to be impaired

(Bainbridge et al., 2021). Yet, task-based functional activity has not been fully explored.

Neurally, the hippocampus has been established as a central brain structure to support

the detail-rich episodic AM retrieval in the healthy brain (Bauer et al., 2017; Brown et al., 2018;

Burianova et al., 2010; McCormick et al., 2020; Moscovitch et al., 2005), albeit some studies

differentiate between hippocampal support for remote and recent autobiographical memories

(see Bayley et al. 2016). In fact, hippocampal activity correlates with the vividness of AM

recollection (Addis et al., 2004; Sheldon & Levine, 2013) and patients with hippocampal damage

show marked deficits in detailed episodic AM retrieval (Miller et al., 2020; Rosenbaum et al.,

2008). In addition, neuroimaging studies illuminate that the hippocampus is almost always co-

activated with a wider set of brain regions, including the ventromedial prefrontal cortex

(vmPFC), lateral and medial parietal cortices, as well as visual-perceptual cortices (Svoboda et

al., 2006). Interestingly, especially during the elaboration phase of AM retrieval, when people

engage in the active retrieval of episodic details to a specific AM, the hippocampus exhibits a

strong functional connection to the visual-perceptual cortices, suggesting a crucial role of this

connection for the embedding of visual-perceptual details into AMs (McCormick et al., 2015).

Yet, not many studies have examined the neural correlates of aphantasia, and none

during AM retrieval. Of the little evidence there is, reports converge on a potential

hyperactivity of the visual-perceptual cortices in aphantasia (Fulford et al., 2018; Keogh et al.,

2020). A prominent theory posits that because of this hyperactivity, small signals elicited during

the construction of mental imagery may not be detected (Pearson, 2019, Keogh et al., 2020).

Pearson further speculates that since spatial abilities seem to be spared, the hippocampus may

not be the underlying cause of aphantasia. In agreement, Bergmann and Ortiz-Tudela (2023)

speculate that individuals with aphantasia might lack the ability to reinstate visually precise

episodic elements from memory due to altered feedback from the visual cortex. In the same

vein, Blomkvist (2022) proposes the extended constructive episodic simulation hypotheses

(CESH+) that suggests that imagination and memory rely on similar neural structures, since both

represent simulated recombinations of previous impressions. This hypothesis has been

supported by shared representations for memory and mental imagery in early visual cortex

(Albers et al., 2013; see also Zeidman & Maguire, 2016). Within this framework, the

hippocampus is supposed to initiate sensory retrieval processes (e.g., in the visual-perceptual

cortices; Danker & Anderson, 2010), comparable to its role in the hippocampal memory

indexing theory (Langille & Gallistel, 2020). Blomkvist (2022) speculates that in aphantasics,

either the hippocampal memory index or the retrieval processes may be impaired.

Thus, the main goal of our study was to examine the neural correlates of AM deficits

associated with aphantasia. We hypothesized that the deficits in AM seen in aphantasia rely on

altered involvement of the hippocampus, visual-perceptual cortices and their functional

connectivity.

119                                **Materials and methods**

**Participants**

In total, 31 healthy individuals with no previous psychiatric or neurological condition

participated in this study. 15 congenital aphantasics and 16 matched controls were recruited

from the database of the *Aphantasia Research Project Bonn* (Monzel, Keidel, et al., 2021;

Monzel, Vetterlein, et al., 2021). Due to technical issues during MRI scanning, one participant

(with aphantasia) had to be excluded from the analyses. Groups were matched for basic

demographic data, that is, sex, age, and education, as well as intelligence assessed with a short

intelligence screening (Baudson & Preckel, 2015) (see Table 1). Oral and written informed

consent was obtained from all participants prior to the commencement of experimental

procedure in accordance with the Declaration of Helsinki (World Medical Association, 2013)

and the local ethics board of the University Hospital Bonn.

**Table 1.**

*Demographic data for aphantasics, controls and the total sample.*

| | Total (n = 30) | Aphantasics (n = 14) | Controls (n = 16) | Test statistic | $p$ | $BF_{01}$ |
|---|---|---|---|---|---|---|
| **Age** | | | | $0.80^a$ | .431 | 2.30 |
| $M$ | 29.77 | 31.47 | 28.19 | | | |
| $SD$ | 11.36 | 10.45 | 12.27 | | | |
| **IQ** | | | | | | |
| $M$ | 93.77 | 91.73 | 95.69 | $0.81^a$ | .425 | 2.29 |

| | | | | | | |
|---|---|---|---|---|---|---|
| *SD* | 13.53 | 16.61 | 10.02 | | | |
| **Sex** | | | | 2.76[b] | .097 | 0.69 |
| Male (%) | 32.3 | 53.3 | 81.3 | | | |
| Female (%) | 67.7 | 46.7 | 18.8 | | | |
| **Education** | | | | 1.59[b] | .662 | 7.90 |
| Secondary school (%) | 6.5 | 6.7 | 6.3 | | | |
| A-levels (%) | 35.5 | 40.0 | 31.3 | | | |
| University degree (%) | 54.8 | 46.7 | 62.5 | | | |
| Doctoral degree (%) | 3.2 | 6.7 | 0.0 | | | |

*Note.* $BF_{01}$ = Bayes Factor, indicates how much more likely H0 is compared to H1. [a] *t*-test, [b] $\chi^2$-

test.

***Vividness of visual imagery questionnaire***

Aphantasia is typically assessed with the Vividness of Visual Imagery Questionnaire (VVIQ;

Marks, 1973, 1995), a subjective self-report questionnaire that measures how vivid mental

scenes can be constructed by an individual. For example, individuals are asked to visualize a

sunset with as much details as possible and rate their mental scene based on a 5-point Likert

scale (ranging from 'no image at all, you only "know" that you are thinking of the object' to

'perfectly clear and as vivid as normal vision'). Since there are 16 items, the highest score of the

VVIQ is 80 indicating the ability to visualize mental images with such vividness as if the event

were happening right there and then. The minimum number of points is 16 indicating that an

individual reported no mental image for any of the items at all. Aphantasia is at the lower end

of the spectrum of imagery-abilities and usually identified with a VVIQ-score between 16 and

32 (e.g., Dawes et al., 2020, 2022).

***Binocular rivalry task***

Since self-report questionnaires such as the VVIQ are associated with several drawbacks, such

as their reliance on introspection (Schwitzgebel, 2002), we administered a mental imagery

priming-based binocular rivalry task to assess mental imagery more objectively (for more

details, see Keogh & Pearson, 2018; Pearson et al., 2008). In short, after imagining either red-

horizontal or blue-vertical Gabor patterns, participants were presented with a red-horizontal

Gabor pattern to one eye and a blue-vertical Gabor pattern to the other eye. Subsequently,

participants were asked to indicate which type of Gabor pattern they predominantly observed.

Usually, successful mental imagery leads subjects to select the Gabor pattern which they had

just visualized. This selection bias can be transferred into a priming score representing visual

imagery strength[1]. Mock stimuli consisting of only red-horizontal or blue-vertical Gabor

patterns were displayed in 12.5 % of the trials to be able to detect decisional biases. Previous

studies have shown that the binocular rivalry task validly correlated with mental imagery

strength (Pearson et al., 2011; Wagner & Monzel, 2023).

*Autobiographical interview*

Detailed behavioral AM measures were obtained in blinded semi-structured interviews either

in-person or online via Zoom (Zoom Video Communications Inc., 2016) using the

Autobiographical Interview (AI; Levine et al., 2002). All interviews were conducted in German.

During the AI, the interviewer asks the participant to recall five episodic AMs from different life

periods: early childhood (up to age 11), adolescent years (ages 11–17), early adulthood (ages

18–35), middle age (35–55), and the previous year. In order to acquire five AMs in every

participant, the middle age memory was replaced by another early adulthood memory for

participants who were younger than 34 years old (see Levine et al., 2002). Hence, all

participants provided the last time period with memories from their previous year. Memories

from the first four periods were considered remote, whereas the memory from the previous

year was considered recent. The interview is structured so that each memory recollection

consists of three parts: free recall, general probe, and specific probe. During free recall, the

---

[1] Imagery strength is used to describe the results of the Binocular Rivalry Task, whereas vividness of mental imagery is used to describe the results of the VVIQ. Although both tasks are correlated, the VVIQ measures vividness, whereas the dimension of the Binocular Rivalry Task is not clearly defined.

participants were asked to recall as many details as possible for a memory of their choice that is

specific in time and place within the given time period. When the participant came to a natural

ending, the free recall was followed by the general and specific probes. During the general

probe, the interviewer asked the participant encouragingly to provide any additional details.

During the specific probe, specific questions were asked for details about the time, place,

perception, and emotion/thoughts of each memory. Then, participants were instructed to rate

their recall in terms of their ability to visualize the event on a 6-point Likert scale (ranging from

'not at all' to 'very much'). The interview was audiotaped, and afterwards transcribed and then

scored by two independent raters according to the standard protocol (Levine et al., 2002). The

interviews were scored after all data had been collected, in random order, and scorers were

blind to the group membership of the participant.

For scoring, the memory details were assigned to two broad categories, that is, internal

and external details. There were the following subcategories of internal details: internal events

(happenings, weather conditions, people present, actions), place (country, city, building, part of

room), time (year, month, day, time of the day), perceptual details (visual, auditory, gustatory,

tactile, smell, body position), and emotion/thought (emotional state, thoughts). The

subcategories for external details were semantic details (factual or general knowledge),

external events (other specific events in time and place but different to the main event),

repetition (repeated identical information without request), and other details (metacognitive

statements, editorializing). In addition, following the standard procedure, an episodic richness

score was given for each memory by the rater on a 7-point Likert scale (ranging from 'not at all'

to 'perfect'). Furthermore, we added a novel rating score of confidence to the protocol since

many participants indicated very strong belief in the details they provided, while others were

insecure about the correctness of their own memories. Confidence scores were again rated on

a 7-point Likert scale (ranging from 'not at all' to 'perfect').

***Debriefing questions***

Following the AI, we asked participants three general questions. These were thought of as open

questions to get people to talk about their personal perspective on AM, spatial cognition, and

imagination.

1.  Typically, how difficult is it for you to recall autobiographical memories?

2.  Typically, how difficult is it for you to orient yourself spatially?

3.  Typically, how difficult is it for you to use your imagination?

After a free report, participants were asked to rate the difficulty on a Likert-scale from 1 ('very

easy') to 6 ('very difficult').

***Autobiographical memory fMRI task***

The experimental fMRI task was adapted from a previous protocol by McCormick et al. (2015).

Two conditions, an AM retrieval task and a simple math task (MA), each consisting of 20

randomized trials, were included in this experiment. During AM trials, cue words, such as 'a

party', were presented on the screen for 12 s and participants were instructed to recall a

personal event related to the word cue which was specific in time and place (e.g., their $20^{th}$

birthday party). Participants were asked to press a response button once an AM was retrieved

to indicate the time point by which they would start to engage in the AM elaboration phase. For

the rest of the trial duration, participants were asked to re-experience the chosen AM and try

to recall as many details as possible without speaking out loud. After each AM trial, participants

were instructed to rate via button presses whether their retrieval had been vivid or faint. We

chose a simple two-button response in order to keep the task as easy as possible. During MA

trials, simple addition or subtraction problems, for example, 47 + 19, were presented on the

screen for 12 s. Here, participants were instructed to press a response button once the

problems were solved and asked to engage in adding 3s to the solutions, for example, (47 + 19)

+ 3 + ... + 3, until the trial ended. The MA trials were followed by a rating whether the MA

problems had been easy or difficult to solve. For both AM and MA, each trial lasted for 12 s, the

maximum time for rating of 3 s, and a jittered inter-stimulus interval (ISI) between 1 to 4 s.

Since we were especially interested in the elaboration phase of AM retrieval, for the fMRI

analyses, we modelled the last 8 s of each AM and MA trial just before the rating screen

appeared.

*MRI data acquisition*

Anatomical and functional data were acquired at the German Center for Neurodegenerative

Diseases (DZNE), Bonn, Germany, using a 3 Tesla MAGNETOM Skyra MRI scanner (Siemens

Healthineers, Erlangen, Germany). A mirror was mounted on the 32 channel receiver head coil

and was placed in the scanner for the participants to view the stimuli shown on an MRI

conditional 30-inch TFT monitor (Medres medical research, Cologne, Germany) placed at the

scanner's rear end. The MRI protocol consisted of anatomical, resting-state, and AM task-based

fMRI scanning sessions. In addition, we acquired further experimental fMRI and DTI data which

are not part of this manuscript. Of note, the resting-state scans were acquired before

participants engaged in the AM task in order to prevent reminiscing about personal memories

during the resting-state. For the anatomical scans, an in-house developed 0.8 mm isotropic

whole-brain T1-weighted sagittal oriented multi-echo magnetization prepared rapid acquisition

gradient echo (MEMPRAGE; Brenner et al., 2014) was employed with the following parameters:

TR = 2.56 s, TEs = 1.72/3.44/5.16/6.88 ms, TA = 6:48, matrix = 320 x 320 x 224, readout pixel

bandwidth = 680 Hz/Pixel, CAIPIRINHA mode. Resting-state (190 volumes, TA = 7 min) and AM

task-based fMRI scans (460 volumes, TA = 15 min) were acquired using an interleaved multi-

slice 3.5 mm isotropic echo planar imaging (EPI) sequence with TR = 2 s, TE = 30 ms, matrix = 64

x 64 x 39, readout pixel bandwidth = 2112 Hz/Pixel (see Jessen et al., 2018). The images were

obtained in an oblique-axial slice orientation along the anterior-posterior commissure line.

During resting-state, the participants were asked to close their eyes and to think about nothing

at all. The first 5 frames of each functional session were excluded for the scanner to reach

equilibrium. Before each functional session an optimal B0 shim was determined and individually

mapped by 2-echo gradient echo (GRE) with same voxel resolution and positioning for later

post-processing.

***Manual segmentation of the hippocampus***

Since our main goal was to assess hippocampal involvement during AM retrieval in aphantasia,

we sought to examine in depth whether there were any group differences in hippocampal

activation in respect to the hemispheric laterality or along its long-axis. For this reason, we

segmented the hippocampus based on the T1 structural images using ITKSnap

(www.itksnap.org, Version 3.8). Although we did not segment specific hippocampal subfields,

our masks included the dentate gyrus, CA1-4, subiculum and pre- and parasubiculum. Whole

masks of the left and right hippocampus were segmented manually in their respective native

space and were divided afterwards into anterior and posterior portions, using the location of

the uncus as boundary.

***fMRI preprocessing***

SPM12 (Statistical Parametric Mapping 12) software package (www.fil.ion.ucl.ac.uk/spm/) on

MATLAB v19a (MathWorks) computing platform (https://matlab.mathworks.com/) was used to

perform resting-state and AM task-based fMRI data preprocessing. The anatomical T1w RMS of

all MEMPRAGE's echoes and functional 2D-EPI images were reoriented along the anterior-

posterior commissure axis. The phase and magnitude images within the field maps were

applied to calculate the voxel displacement maps (VDM) for geometrical correction of the

distorted EPI images. The echo times were set to 4.92 ms (short) and 7.38 ms (long). The total

EPI readout time was 34.56 ms. The calculated EPI and VDMs were applied to the functional

scans for realignment and unwarping. The functional scans were then co-registered to the

segmented bias corrected T1 scans.

Whole-brain differences between groups were evaluated. Thus, co-registered scans were

normalized to the Montreal Neurological Institute (MNI) space and a Gaussian smoothing

kernel of 8 mm FWHM was applied. In addition, for functional connectivity analyses, denoising

was applied using a linear regression model of potential confounding effects (global white

matter signal, CSF signal, and ART scrubbing parameters) in the BOLD signal using CONN

software package v20.b (www.nitrc.org/projects/conn/). Temporal band pass filter was set

from 0.01 to infinite to further minimize artifacts.

**Statistical analyses**

***Behavioral analyses***

Two samples t-tests were calculated to assess differences in the priming scores of aphantasics

and controls in the binocular rivalry task. One sample t-tests were used to distinguish the

performances of both groups from chance. To assess differences of Autobiographical Interview

scores between aphantasics and controls, a 2x2x2 ANOVA with post-hoc t-tests were calculated

with type of memory details (internal vs. external) and memory recency (remote vs. recent) as

within-subject factors and group (aphantasics vs. controls) as between-subject factor.

Afterwards, Bonferroni-corrected t-tests were conducted for specific internal (time, place,

internal event, perception, emotion) and external (external event, semantic, repetition, other)

memory details. Differences in memory ratings (confidence, episodic richness), self-reported

visualization scores, debriefing questions, and behavioral responses during fMRI scanning were

also assessed via two sample t-tests.

***Hippocampal activity associated with autobiographical memory in native space***

In order to examine hippocampal activity associated with autobiographical memory, we

extracted signal intensity values for both AM and MA trials for each participant for our

manually segmented anatomical masks of the left and right, anterior and posterior portions of

the hippocampus using the MATLAB-based Response Exploration toolbox (REX;

www.nitrc.org/projects/rex/). We then calculated for each participant for each anatomical

mask the difference between AM and MA signal intensities. Afterwards, group differences

between aphantasics and controls with respect to the laterality effects and effects between the

anterior and posterior hippocampus were assessed using a 2-way ANOVA with a post-hoc

Tukey's multiple comparison test, applying a significance threshold of $\alpha$ = .05.

### *Whole-brain fMRI activation analyses*

After focusing on the hippocampus, we examined group differences of whole-brain activation

associated with AM and MA following the standard GLM procedure in SPM12. Owing to the

prominence of mental imagery during the elaboration phase of AM retrieval, we analyzed the

last 8 s of the AM and MA trials prior to the display of the vividness rating. These trials were

modelled as mini blocks in the GLM with motion correction regressors included as covariate of

no interest. We specified our main contrast of interest, i.e., AM versus MA on the first level,

which was then brought to the second group level using a one-sample t-test. Finally, the

activation maps of the two groups were compared using a two-sample t-test. For whole-brain

analysis, we applied a significance threshold of $p$ < .001, and voxel cluster size of 10 adjacent

voxels, uncorrected.

### *ROI-to-ROI functional connectivity analyses*

One of our main a priori hypotheses stated that the hippocampus and the visual-perceptual

cortex show differential engagement during AM retrieval associated with aphantasia which was

confirmed by our whole-brain activation analyses. Based on these results, we sought to

examine the functional connectivity between those two areas. Towards this end, we created

regions of interest (ROIs, spheres with a diameter of 10 mm consisting of 536 voxels) around

the three peaks of the activation differences, following the whole-brain fMRI activation

analyses, using the MarsBaR HBM toolbox (Brett et al., 2002). The ROIs comprised (1) the right

hippocampus, MNI: x = 39, y = -31, z = -13, (2) the right visual cortex, MNI: x = 12, y = -79, z = 5,

and (3) the left visual cortex, MNI: x = -9, y = -76, z = 29. Using CONN, we examined functional

connectivity (i.e., Generalized Psycho-Physiological Interactions, weighted general linear model

with bivariate correlation) between the hippocampal ROI and the ROIs situated in the visual-

perceptual cortex during AM task-based fMRI and during resting-state. Furthermore, in order to

examine how well functional connectivity between hippocampus and the visual cortex reflected

an individuals' ability to visualize mental events, we examined a regression model with the

visualization scores of the AI as criterion and resting state connectivity values, group allocation

and the interaction term of the connectivity values and group allocation as predictors. For these

a priori driven analyses, we applied a significance threshold of $p < .05$, small volume corrected,

and a voxel cluster threshold of 10 adjacent voxels.

349                                      **Results**

**VVIQ and binocular rivalry task**

Aphantasics (*M* = 16.57, *SD* = 1.02), scored significantly lower on the VVIQ than controls (*M* =

62.94, *SD* = 8.71), *t*(15.47) = 21.12, *p* < .001, *d* = 7.23. Furthermore, aphantasics and controls

differed in the priming score of the binocular rivalry task, *t*(18.04) = 2.41, *p* = .027, *d* = 0.87.

While controls were primed by their own mental imagery in 61.3 % ($SD$ = 13.1 %) of the trials,

aphantasics were only primed 52.6 % ($SD$ = 4.9 %) of the time. In fact, the performance of

controls differed significantly from chance, $t(14)$ = 3.34, $p$ = .005, $d$ = 0.86, whereas

performance of aphantasics did not, $t(13)$ = 1.96, $p$ = .072. Moreover, the VVIQ scores

correlated positively with the performance on the binocular rivalry task, $r(28)$ = .43, $p$ = .022.

For the mock trials, no significant differences in priming scores were found between groups,

$t(28)$ = 0.86, $p$ = .396, or related to chance (aphantasics: $t(13)$ = 0.74, $p$ = .475; controls: $t(15)$ =

0.42, $p$ = .682). These findings validate our groups by indicating that visual imagery strength was

diminished in aphantasics.

**Autobiographical Interview**

Regarding the AI, we found significant main effects of memory period, $F(1, 27)$ = 11.88, $p$ = .002,

$\eta_p^2$ = .31, type of memory details, $F(1, 27)$ = 189.03, $p < .001$, $\eta_p^2$ = .88, and group, $F(1, 27)$ =

9.98, $p$ = .004, $\eta_p^2$ = .27. When the other conditions were collapsed, aphantasics (M = 26.29, SD

= 9.58) described less memory details than controls (M = 38.36, SD = 10.99). For aphantasics

and controls combined, more details were reported for recent (M = 35.17, SD = 14.19) than

remote memories (M = 29.06, SD = 11.12), and internal details (M = 43.59, SD = 17.91) were

reported more often than external details (M = 20.64, SD = 8.94). More importantly, a 2-way

interaction was found between type of memory details and group, $F(1, 27)$ = 54.09, $p < .001$, $\eta_p^2$

= .67, indicating that aphantasics reported significantly less internal memory details, $t(27)$ =

5.07, $p < .001$, $d$ = 1.83, but not significantly less external memory details, $t(27)$ = 0.13, $p$ = .898,

compared to controls (see Figure 1b). No 2-way interaction between memory period and group,

$F(1, 27) = 0.62$, $p = .439$, and no 3-way interaction between memory period, group and type of

memory details was found, $F(1, 27) = 3.87$, $p = .060$.

Based on the interaction effect between group and type of memory details, we compared

specific categories of internal and external memory details between the groups. For internal

details and in comparison to controls, aphantasics reported less internal events, $t(27) = 3.22$, $p$

$= .016$, $d = 1.17$, less emotional details, $t(27) = 4.40$, $p < .001$, $d = 1.59$, less perceptual details,

$t(27) = 4.95$, $p < .001$, $d = 1.79$, and less details regarding time, $t(27) = 5.27$, $p < .001$, $d = 1.90$,

and place, $t(27) = 3.31$, $p < .013$, $d = 1.20$ (see Figure 1c). On the other hand, no significant

differences were found for external details, including external events, $t(27) = 0.71$, $p > .999$,

semantic details, $t(27) = 0.02$, $p > .999$, repetition, $t(27) = 0.46$, $p > .999$, and other details, $t(27)$

$= 0.45$, $p > .999$ (see Figure 1C). Regarding the rating scales, we found that aphantasics showed

less episodic richness, $t(27) = 7.50$, $p < .001$, $d = 2.71$, and less memory confidence, $t(27) = 5.85$,

$p < .001$, $d = 2.11$ (see Figure 1a) as well as lower self-reported visualization scores, $t(27) =$

$11.92$, $p < .001$, $d = 4.30$, than controls.

**Figure 1. AM deficits associated with aphantasia.** (A) Mean amount (± SEM) of episodic

richness and confidence in the Autobiographical Interview for controls and aphantasics. (B)

Mean amount (± SEM) of internal details and external details for recent and remote memories.

(C) Mean amount (± SEM) of specific internal and external memory details for aphantasics and

controls.

* $p < .05$, ** $p < .01$, *** $p < .001$, **** $p < .0001$, n.s. = non-significant.

**Debriefing questions**

The debriefing questions were employed as a way for participants to reflect on their own cognitive abilities. Of note, these were not meant to represent or replace necessary future experiments. There were stark differences between the groups in how they answered our debriefing questions. While aphantasics reported that they typically have greater difficulty to recall autobiographical memories, $t(27) = 6.20$, $p < .001$, $d = 2.31$, and to use their imagination in daily life, $t(24) = 10.18$, $p < .001$, $d = 3.93$, they did not report difficulties in spatial orientation, $t(27) = 0.62$, $p = .541$, $d = 0.23$.

**Behavioral results of the fMRI AM task**

We found stark differences for the vividness response between groups, $t(28) = 5.29$, $p < .001$. While controls reported in 86 % ($SD$ = 26 %) of trials that their AM retrieval had been vivid, aphantasics indicated only in 20 % ($SD$ = 20 %) of trials that their AM retrieval had been vivid. Moreover, aphantasics responded slower ($M$ = 1.34 s, $SD$ = 0.38 s) than controls ($M$ = 1.00 s, $SD$ = 0.29 s) when they were asked whether their retrieved memories were vivid or faint, $t(28) = 2.78$, $p = .009$, possibly reflecting uncertainty in their response. In contrast, there were no significant differences between groups during the MA trials, neither on the easy/hard response, $t(28) = 1.16$, $p = .255$, nor on the reaction times, $t(28) = 0.58$, $p = .567$.

In addition, aphantasics and controls did not differ significantly in their time searching for a memory in AM trials, $t(19) = 1.03$, $p = .315$. On average, aphantasics spent 3.42 s ($SD$ =

0.74 s) and controls spent 3.15 s (*SD* = 0.48 s). Furthermore, both groups did not differ in their

speed to solve the math problems, $t(23) = 0.09$, $p = .926$. Aphantasics spent 3.87 s (*SD* = 0.97 s)

to solve a math problem and controls spent 3.90 s (*SD* = 0.72 s). For the button press, there

were 9 % missing values in AM trials and 7 % missing values in MA trials with no significant

differences of missing values between groups, neither for AM trials, $t(19.98) = 1.11$, $p = .281$,

nor for MA trials, $t(18.13) = 0.52$, $p = .609$.

**Native space differences in hippocampal activation during AM retrieval**

First, we sought to examine the hippocampal activation associated with AM in aphantasics and

controls (see Figure 2). The extracted fMRI signals illustrated stark group differences in

hippocampal AM-associated activation, $F(17, 252) = 3.03$, $p < .001$. Aphantasics showed

reduced activation of bilateral hippocampi, including the left anterior ($p = .033$), left posterior

($p = .027$), right anterior ($p = .047$), and right posterior hippocampus ($p = .025$). There was no

laterality effect nor differences along the pattern of activation down the anterior-posterior axis

between the groups (all $p > .05$). These findings indicate that the behavioral AM deficit

associated with aphantasia is reflected neurally by a reduced bilateral hippocampal activation.

439
***Figure 2. Reduced hippocampal activity during AM retrieval associated with aphantasia.*** The

signal intensities during AM and MA were extracted from anatomical hippocampal masks

created from each individual participant. (A) An example of a 3D reconstruction of the

hippocampus, separated into anterior and posterior portions for the left hippocampus. (B) The

comparison between the percentage of signal change during the AM and MA tasks in the

hippocampus of aphantasics and controls. Aphantasics show reduced differentiation between

AM and MA than controls in all portions of the hippocampus. * $p < .05$.

**Activation patterns associated with AM retrieval**

Second, we examined whole-brain activation during AM retrieval of both groups and the results

are displayed in Figure 3a and b. Additionally, the peak coordinates of AM and MA activation

for aphantasics and controls are shown in Table 1 and 2, respectively.

Overall, both groups showed greater activation in all areas typically associated with AM,

including bilateral hippocampus, vmPFC, and medial/lateral parietal regions, during AM

retrieval. When examining the group differences, aphantasics displayed greater activation in

bilateral visual-perceptual regions (maximum in lingual gyrus) in the occipital lobe than

controls, $t(28)$ = 4.41, $p$ < .001 (MNI: right visual cortex: x = 12, y = -79, z = 5; left visual cortex: x

= -9, y = -76, z = 29, see Figure 3c, d, and e). In contrast, controls showed greater activation in

the right hippocampus than aphantasics, $t(28)$ = 3.77, $p$ < .001 (MNI: x = 39, y = -31, z = -13). An

additional correlational analysis revealed that those participants with higher visual-perceptual

cortex activation had less hippocampal activation, $r(28)$ = -.39, $p$ = .041, indicating that there

was a trade-off between increased visual-perceptual cortex activation and decreased

hippocampal activation.

*Figure 3.*  **Activation during AM retrieval task.** (A) Stronger activated cortical regions during AM

retrieval (in warm colors) in comparison to math problem solving (in cool colors) in aphantasics

and (B) controls. (C) Aphantasics showed greater activation in visual-perceptual cortices than

controls, and (D) controls showed stronger activation in the right posterior hippocampus than

aphantasics. Images are thresholded at $p < .001$, cluster size 10, uncorrected, except (D) which

is thresholded at $p < .01$, cluster size 10, for display purposes only (i.e., the peak voxel and

adjacent 10 voxels also survived $p < 0,001$, uncorrected). (E) The percentage of signal change

for the contrast AM versus MA were extracted from the peaks of activated voxels, each with 1

mm sphere for display purposes.

**Table 1. Peak coordinates of the AM and MA activation for Aphantasia.**

| Region | Hemisphere | MNI Coordinates | | | Voxels | T-value |
|---|---|---|---|---|---|---|
| | | X | Y | Z | | |
| **Activation AM > MA** | | | | | | |
| Posterior Cingulate Gyrus | Right | 18 | -57 | 11 | 4657 | 11.00 |
| Parahippocampual Gyrus* | Left | -21 | -31 | -13 | | 9.06 |
| Hippocampus | Left | -27 | -17 | -19 | 205 | 8.40 |
| Superior Frontal Gyrus | Left | -12 | 47 | 50 | 926 | 8.38 |
| Angular Gyrus | Left | -42 | -55 | 23 | 165 | 7.88 |
| Lateral Orbitofrontal Cortex | Left | -42 | 38 | -16 | 208 | 7.58 |
| Hippocampus | Right | 18 | -37 | -1 | 199 | 6.89 |
| Cerebellum | Right | 15 | -79 | -37 | 109 | 6.33 |
| Brainstem | Right | 3 | -46 | -52 | 43 | 6.03 |
| Parahippocampual Gyrus* | Right | 24 | -31 | -13 | | 6.02 |
| Middle Temporal Gyrus | Right | 60 | 2 | --19 | 76 | 5.27 |
| Supramarginal Gyrus | Right | 54 | -58 | 32 | 26 | 4.99 |
| Middle Frontal Gyrus | Left | -39 | 20 | 50 | 12 | 4.52 |
| **Activation MA > AM** | | | | | | |
| Precuneus | Left | -18 | -58 | 41 | 594 | -3.85 |
| Inferior Temporal Gyrus | Right | 51 | -46 | -13 | 123 | -3.85 |
| Precuneus | Right | 24 | -49 | 53 | 718 | -3.85 |
| Insula | Left | -30 | 23 | 11 | 48 | -3.85 |
| Inferior Temporal Gyrus | Left | -51 | -49 | -13 | 67 | -3.86 |
| Cerebellum | Right | 30 | -67 | -52 | 27 | -3.87 |
| Middle Frontal Gyrus | Right | 33 | 41 | 17 | 34 | -3.87 |
| Superior Frontal Gyrus | Right | 30 | 5 | 59 | 52 | -3.87 |
| Inferior Frontal Gyrus | Right | 54 | 14 | 29 | 35 | -3.88 |
| Insula | Right | 39 | 11 | 8 | 51 | -3.88 |
| Inferior Frontal Gyrus | Left | -57 | 11 | 26 | 182 | -3.88 |
| Lateral Globus Pallidus | Right | 23 | -7 | 14 | 14 | -3.92 |
| Cerebellum | Left | -24 | -64 | -46 | 16 | -3.93 |

477 *Sub-cluster level, Cluster size = 10 voxels, p-value = 0.001

480    **Table 2. Peak coordinates of the AM and MA activation for healthy controls.**

| Region | Hemisphere | MNI Coordinates | | | Voxels | T-value |
|---|---|---|---|---|---|---|
| | | X | Y | Z | | |
| **Activation AM > MA** | | | | | | |
| Parahippocampal Gyrus | Right | 27 | -28 | -19 | 11319 | 12.41 |
| Parahippocampal Gyrus* | Left | -24 | -25 | -16 | | 9.01 |
| Cerebellum | Left | -18 | -76 | -37 | 108 | 7.67 |
| Anterior Cingulate | Right | 9 | 35 | 11 | 13 | 7.01 |
| Medial Frontal Gyrus | Right | 18 | 32 | 29 | 233 | 6.93 |
| Inferior Frontal Gyrus | Right | 60 | 32 | 11 | 53 | 5.94 |
| Hippocampus | Left | -36 | -22 | -16 | 252 | 5.64 |
| Hippocampus | Right | 27 | -22 | -16 | 233 | 5.28 |
| Hypothalamus | Right | 3 | -4 | -10 | 16 | 4.93 |
| **Activation MA > AM** | | | | | | |
| Post Central Gyrus | Left | -33 | -43 | 62 | 643 | -3.73 |
| Precuneus | Right | 21 | -52 | 53 | 483 | -3.74 |
| Inferior Frontal Gyrus | Right | 51 | 8 | 26 | 16 | -3.74 |
| Middle Occipital Gyrus | Right | 33 | -82 | 2 | 26 | -3.75 |
| Middle Temporal Gyrus | Left | -51 | -58 | -1 | 18 | -3.76 |

481    *Sub-cluster level, Cluster size = 10 voxels, p-value = 0.001

**Exploring functional connectivity of hippocampus and visual-perceptual cortices during AM**

The whole-brain analyses strengthened our hypothesis that a core difference between aphantasics and controls lies in the interplay between the visual-perceptual cortex and the hippocampus. To test this interplay, in a third step, we examined functional connectivity between the peak differences of the hippocampus and the visual-perceptual cortex during AM retrieval (see Figure 4). We found a group difference in functional connectivity between the right hippocampus and left visual-perceptual cortices, $t(28) = 2.65$, $p = .006$. Interestingly, while aphantasics show almost no functional connectivity between those two ROIs, controls displayed a strong negative connectivity between the two locations, $p = .013$.

*Figure 4. Functional connectivity between the visual-perceptual cortex and hippocampus during AM retrieval.* (A) During AM retrieval, group differences in functional connectivity amongst the ROIs were only found between the right hippocampus, and left visual-perceptual cortices. (B) Controls displayed a stark negative correlation, whereas aphantasics did not. Image is displayed at $p < .05$, small volume corrected, and a voxel cluster threshold of 10 adjacent voxels.

**Exploring functional connectivity of hippocampus and visual-perceptual cortices during resting-state**

Lastly, we examined resting-state connectivity between our identified ROIs in the right posterior hippocampus and left and right visual-perceptual cortex. There was no group difference during resting-state between these ROIs. In order to examine whether resting-state functional connectivity carried information about one's ability to visualize AMs, we added the visualization scores as a regressor of interest in our model. While connectivity alone did not predict the visualization scores in the interview, $F(14, 40) = 1.651$, $p = .391$, $\beta = -.06$, we found a main effect of group, $F(3, 40) = 353.2$, $p < .001$, $\beta = .92$, and an interaction between group and connectivity, $F(3, 54) = 305.1$, $p < .001$, $\beta = .26$. Interestingly, for controls, we found a positive correlation between the resting-state connectivity of the right hippocampus and the visual cortex and the visualization scores from the interview, $r(13) = .65$, $p = .011$ (see Figure 5). On the other hand, for aphantasics, we found a negative correlation between the resting-state connectivity of the right hippocampus and the visual cortex and the visualization scores from the interview, $r(14) = -.57$, $p = .027$.

In sum, our fMRI results indicate that the impaired AM retrieval associated with aphantasia is reflected by functional alterations of the hippocampus and visual-perceptual cortex, as well as the interaction between them.

Figure 5. Functional connectivity between the visual-perceptual cortex and hippocampus
during resting-state explains visualization abilities. Resting-state functional connectivity
between the right hippocampus and the right visual-perceptual cortex correlates with
visualization abilities. Fitted straight lines indicate a negative correlation for aphantasics (red)
and a positive correlation for controls (blue).

 **Discussion**

In this study, we set out to examine the neural correlates of episodic AM retrieval in aphantasia

as a way to examine the influence of visual imagery on episodic AM. In line with previous

reports, we found that aphantasics reported less sensory details during AM retrieval regardless

of the recency of memory (cf. Dawes et al., 2020; 2022; Milton et al., 2020; Zeman et al., 2020).

Strikingly, the deficit in constructing visual imagery associated with aphantasia did not only lead

to a reduced retrieval of visual-perceptual details but to a broader impairment in retrieving

episodic AMs, including reduced emotions and confidence attached to the memories. Thus, in

agreement with a recent account of aphantasia (Blomkvist, 2022), our results support the idea

that a diminished construction of visual details during AM retrieval leads to a more general

episodic memory deficit. We expand the current knowledge by adding that this AM deficit is

reflected neurally by an altered activation and connectivity pattern between the hippocampus

and visual-perceptual cortices. Our findings provide novel insights into three current debates: 1)

the mechanisms of aphantasia-related AM deficits, 2) the similarities and differences between

aphantasics and individuals with hippocampal damage, and 3) the neural models of AM.

**Potential mechanisms underlying aphantasia-related AM deficits**

We report that aphantasics show increased activation of bilateral visual-perceptual

cortices as well as decreased hippocampal activation during AM retrieval in comparison to

controls. Increased activity in the visual-perceptual cortices in aphantasics has been reported

previously, albeit not associated with AM (Fulford et al., 2018; Keogh et al., 2020). In a

prominent review, Pearson synthesizes evidence about the neural mechanism of imagery

strength (Pearson, 2019). Indeed, activity metrics in the visual cortex predict imagery strength

(Cui et al., 2007; Dijkstra et al., 2017). Interestingly, lower resting activity and excitability result

in stronger imagery, and reducing cortical activity in the visual cortex via transcranial direct

current stimulation (tDCS) increases visual imagery strength (Keogh et al., 2020). Thus, one

potential mechanism of aphantasia-related AM deficits is that the heightened activity of the

visual-perceptual cortices observed in our and previous work hinders aphantasics to detect

weaker imagery-related signals.

Further, we had the a priori hypothesis that hippocampal activation will be decreased in

aphantasics; a hypothesis which we could confirm in our native space hippocampal analysis. On

the whole-brain level, perhaps due to our small sample size, only the cluster of activation group

differences in the right posterior hippocampus survived the statistical threshold. Given the low

power, further studies are needed to confirm this effect; however, the right posterior

hippocampus interacts in the healthy brain heavily with the visual-perceptual cortex only during

the elaboration phase of AM retrieval (McCormick et al. 2015).

In addition, controls exhibited a strong functional connection between both brain

structures during AM retrieval and this functional connectivity predicted better visualization

skills. At first glance, it is surprising that this functional connectivity was negative. However,

negative visual-perceptual cortex activation during perceiving and imagining scenes has been

reported before (McCormick et al. 2021). One possible explanation might be that signals from

the hippocampus selectively inhibit imagery-irrelevant activation in the visual-perceptual

cortices (e.g., sensory noise) to carve out imagery-related signals (cf. Pace et al., 2023). This

would be in line with the hypothesis stated above, that a bad signal-to-noise ratio in the visual

cortex hiders aphantasics to create mental imagery. Either way, the described functional

connectivity between the hippocampus and the visual-perceptual cortex fits well to previous

neuroimaging studies pointing towards a central role of the dynamic interplay between these

brain structures during AM retrieval (McCormick, et al. 2015). This interplay seems to be

especially important during the elaboration stage of AM retrieval, a period when specific visual-

perceptual details are being actively brought back into the mind's eye. At this point, however, it

remains unclear whether the disruption of AM elaboration associated with aphantasia takes

place during the encoding, storage, or retrieval process.

From a theoretical point of view, the extended constructive episodic simulation

hypothesis proposes a top-down hierarchy during mental imagery (Blomkvist, 2022). In this

model, the hippocampus initiates retrieval processes in primary sensory brain regions, such as

the visual-perceptual cortex in order to retrieve visual-perceptual details associated with a

specific AM. Evidence for such top-down hierarchies during mental imagery have been

observed in fronto-parietal and occipital networks via effective connectivity analyses, such as

Granger Causality and Dynamic Causal Modelling (Dentico et al., 2014; Dijkstra et al., 2017;

Mechelli et al., 2004). For example, intracranial and high-density scalp electroencephalography

(EEG) provided evidence of high frequency hippocampal signaling during the recall of

perceptual cues in patients with epilepsy, indicating that the hippocampus drives the switch

from perception to memory recalling (Treder et al., 2021). In aphantasia, it is hypothesized that

this top-down hierarchy is disrupted and therefore, the hippocampus can no longer initiate the

retrieval and incorporation of visual-perceptual details in one coherent mental event. Because

of the slow temporal resolution of our fMRI sequence, our data cannot directly speak to the

question of temporal directionality between the hippocampus and visual-perceptual cortex.

Nonetheless, our findings suggest that the bidirectional connectivity between both brain

structures is crucial for the re-experience of episodic AMs. As such, hippocampal processes may

be needed to retrieve specific details and if these details are not provided by the visual-

perceptual cortices, the entire episodic AM retrieval seems to fail.

**Similarities and differences between aphantasics and individuals with hippocampal lesions**

At face value, the episodic AM deficits in aphantasia in our data and reported previously

(Dawes et al., 2020; 2022; Milton et al., 2020; Zeman et al., 2020), as well as the decreased

hippocampal activation during AM retrieval suggest that aphantasia is a selective episodic

memory condition (see Blomkvist, 2022), similar to AM amnesia known from individuals with

hippocampal damage. In fact, previous and the current study show that aphantasics and

individuals with hippocampal damage report less internal details across several memory detail

subcategories, such as emotional details and temporal details (Rosenbaum et al., 2008; St-

Laurent et al., 2009; Steinvorth et al., 2005), and these deficits can be observed regardless of

the recency of the memory (Miller et al., 2020). These similarities suggest that aphantasics are

not merely missing the visual-perceptual details to specific AM, but they have a profound

deficit associated with the retrieval of AM.

Nonetheless, there are also stark differences between aphantasics and individuals with

hippocampal damage. Foremost, aphantasics seem not to have difficulties to retrieve spatial

information (Bainbridge et al., 2021), which is another inherent function of the hippocampus

(Burgess et al., 2002; O'Keefe, 1991). In the current study, we did not set out to examine spatial

cognition in aphantasics, however, parts of our data speak to this aspect. While in our study

aphantasics reported less amount of spatial details during the AI, this standard scoring

procedure only counts place details when the exact place is recalled and is not meant to assess

the recall of spatial layout (Levine et al., 2002). Thus, this place score may not represent spatial

cognition per se. In fact, when asking aphantasics about their experience, they point out

difficulties in recalling AM and using imagination in daily life, however they report no difficulties

in spatial orientation. Indeed, often during the interview, aphantasics would explain that they

know how the space around them felt, they just cannot see it in front of their mind's eye. One

aphantasic put her finger on it, describing it as: "I can put my consciousness in my kitchen at

home and feel all around but there is no visual image attached to this feeling." These

observations support the idea that some hippocampal processes, at least regarding spatial

navigation, may be intact in people with aphantasia (Bainbridge et al., 2021). However, spatial

cognition should be formally addressed in future studies. One way to assess this hippocampal

function would be to examine tasks, which rely on scene construction. The scene construction

theory states that the hippocampus is crucially needed for the construction of spatially

coherent mental models of scenes (Maguire & Mullally, 2013). For example, patients with

hippocampal damage cannot imagine the spatial layout of fictitious scenes (Hassabis et al.,

2007), they detect less errors in spatially-incoherent scenes than controls (McCormick et al.,

2017), and they show less scene-dependent mind-wandering episodes (McCormick, Rosenthal,

et al., 2018). In contrast, we would predict that aphantasics have diminutive deficits in tasks

that depend on hippocampal scene construction processes.

What could be impaired in aphantasics are all cognitive functions which rely on the

population of the constructed scenes with visual-perceptual details, such as episodic AM

retrieval, episodic future thinking, complex decision-making, and complex empathy tasks.

**Towards a novel neural model of autobiographical memory**

While more research is required exploring the cognitive landscape associated with aphantasia,

such as spatial cognition and scene construction, our data contribute to an old debate of how

AM retrieval and visual imagery are intertwined. We propose that the hippocampus is

embedded in a brain-wide network, comprising the vmPFC and visual-perceptual cortices, in

which each of these nodes contributes specific processes to the re-construction of extended

detail-rich mental events (see also Ciaramelli et al., 2019; McCormick, Ciaramelli, et al., 2018).

Within this model, the vmPFC initiates and oversees the scene construction process which takes

place in the hippocampus. Further, the visual-perceptual cortex provides the visual details

which are essential to populate the hippocampally constructed scenes. This model is backed up

by a previous MEG study revealing that the vmPFC directs hippocampal activity during the

initiation of AM retrieval (McCormick et al., 2020). This finding has been replicated and

extended by Chen et al. (2021), showing that the vmPFC leads hippocampal involvement during

scene construction and other scene-based processes (Monk et al., 2021). Moreover, the

connection between the hippocampus and the visual-perceptual cortex seems equally crucial.

There are a few case reports of damage to the occipital cortex causing AM amnesia (Greenberg

et al., 2005), potentially by preventing the population of the hippocampally constructed scenes.

Furthermore, our current study suggests that a reliable connectivity between the hippocampus

and the visual-perceptual cortices is important to provide the visual details necessary for

successful vivid, detail-rich AM retrieval.

**Conclusion**

Aphantasia provides a natural knock-out model for the influence of visual imagery on different

cognitive functions. We here report a tight link between visual imagery and our ability to

retrieve vivid and detail-rich personal past events, as aphantasics do not only report fewer

visual-perceptual details during episodic AM retrieval but also show decreased confidence and

emotionality associated with these memories. In this context, we highlight the central role of

the functional connectivity between the hippocampus and occipital cortex to assemble visual-

perceptual details into one coherent extended mental event. Exciting novel research avenues

will be to examine hippocampal-dependent spatial cognition in aphantasics and to investigate

whether neuroscientific interventions can be used to enhance AM retrieval by enhancing visual

imagery.

**References**

Addis, D. R., Moscovitch, M., Crawley, A. P., & McAndrews, M. P. (2004). Recollective qualities

683           modulate hippocampal activation during autobiographical memory retrieval.

684           *Hippocampus*, *14*(6), 752–762.

685      Addis, D. R., Wong, A. T., & Schacter, D. L. (2007). Remembering the past and imagining the

686           future: Common and distinct neural substrates during event construction and elaboration.

687           *Neuropsychologia*, 45(7), 1363-1377.

688      Albers, A. M., Kok, P., Toni, I., Dijkerman, H. C., & de Lange, F. P. (2013). Shared representations

689           for working memory and mental imagery in early visual cortex. *Current Biology*, *23*(15),

690           1427–1431. https://doi.org/10.1016/j.cub.2013.05.065

691      Bainbridge, W. A., Pounder, Z., Eardley, A. F., & Baker, C. I. (2021). Quantifying aphantasia

692           through drawing: Those without visual imagery show deficits in object but not spatial

693           memory. Cortex, 135, 159-172.

694      Baudson, T. G., & Preckel, F. (2015). mini-q: Intelligenzscreening in drei Minuten. *Diagnostica*.

695      Bauer, P. J., Pathman, T., Inman, C., Campanella, C., & Hamann, S. (2017). Neural correlates of

696           autobiographical memory retrieval in children and adults. *Memory*, *25*(4), 450–466.

697           https://doi.org/10.1080/09658211.2016.1186699

698      Bergmann, J., & Ortiz-Tudela, J. (2023). Feedback signals in visual cortex during episodic and

699           schematic memory retrieval and their potential implications for aphantasia. *Neuroscience*

700           *and Biobehavioral Reviews*, *152*, 105335.

701           https://doi.org/10.1016/j.neubiorev.2023.105335

Blomkvist, A. (2022). Aphantasia: In search of a theory. *Mind and Language*, *38*(3), 866–888.

https://doi.org/10.1111/mila.12432

Brenner, D., Stirnberg, R., Pracht, E. D., & Stöcker, T. (2014). Two-dimensional accelerated MP-

RAGE imaging with flexible linear reordering. *Magnetic Resonance Materials in Physics,*

*Biology and Medicine*, *27*(5), 455–462. https://doi.org/10.1007/S10334-014-0430-

Y/TABLES/2

Brett, M., Anton, J.-L., Valabregue, R., Poline, J.-B., & others. (2002). Region of interest analysis

using an SPM toolbox. *8th International Conference on Functional Mapping of the Human*

*Brain*, *16*(2), 497.

Brown, T. I., Rissman, J., Chow, T. E., Uncapher, M. R., & Wagner, A. D. (2018). Differential

Medial Temporal Lobe and Parietal Cortical Contributions to Real-world Autobiographical

Episodic and Autobiographical Semantic Memory. *Scientific Reports*, *8*(1), 1–14.

https://doi.org/10.1038/s41598-018-24549-y

Burgess, N., Maguire, E. A., & O'Keefe, J. (2002). The human hippocampus and spatial and

episodic memory. *Neuron*, *35*(4), 625–641.

Burianova, H., McIntosh, A. R., & Grady, C. L. (2010). A common functional brain network for

autobiographical, episodic, and semantic memory retrieval. *NeuroImage*, *49*(1), 865–874.

https://doi.org/10.1016/j.neuroimage.2009.08.066

Chen, D., Kunz, L., Lv, P., Zhang, H., Zhou, W., Liang, S., Axmacher, N., & Wang, L. (2021). Theta

oscillations coordinate grid-like representations between ventromedial prefrontal and

entorhinal cortex. *Science Advances*, *7*(44), eabj0200.

Ciaramelli, E., De Luca, F., Monk, A. M., McCormick, C., & Maguire, E. A. (2019). What" wins" in

VMPFC: Scenes, situations, or schema? *Neuroscience & Biobehavioral Reviews*, *100*, 208–

210.

D'Argembeau, A., & Van der Linden, M. (2006). Individual differences in the phenomenology of

mental time travel: the effect of vivid visual imagery and emotion regulation strategies.

*Consciousness and Cognition*, *15*(2), 342–350.

https://doi.org/10.1016/j.concog.2005.09.001

Danker, J. F., & Anderson, J. R. (2010). The ghosts of brain states past: remembering reactivates

the brain regions engaged during encoding. *Psychological Bulletin*, *136*(1), 87.

Dawes, A. J., Keogh, R., Andrillon, T., & Pearson, J. (2020). A cognitive profile of multi-sensory

imagery, memory and dreaming in aphantasia. *Scientific Reports*, *10*(1), 10022.

https://doi.org/10.1038/s41598-020-65705-7

Dawes, A. J., Keogh, R., Robuck, S., & Pearson, J. (2022). Memories with a blind mind:

Remembering the past and imagining the future with aphantasia. *Cognition*, *227*, 105192.

https://doi.org/10.1016/J.COGNITION.2022.105192

Dentico, D., Cheung, B. L., Chang, J. Y., Guokas, J., Boly, M., Tononi, G., & Van Veen, B. (2014).

Reversal of cortical information flow during visual imagery as compared to visual

perception. *NeuroImage*, *100*, 237–243.

https://doi.org/10.1016/j.neuroimage.2014.05.081

Dijkstra, N., Zeidman, P., Ondobaka, S., Van Gerven, M. A. J., & Friston, K. (2017). Distinct Top-

down and Bottom-up Brain Connectivity during Visual Perception and Imagery. *Scientific*

*Reports*, *7*(1), 1–9. https://doi.org/10.1038/s41598-017-05888-8

Fulford, J., Milton, F., Salas, D., Smith, A., Simler, A., Winlove, C., & Zeman, A. (2018). The neural

correlates of visual imagery vividness – An fMRI study and literature review. *Cortex*, *105*,

747   26–40. https://doi.org/10.1016/j.cortex.2017.09.014

748 Greenberg, D. L., Eacott, M., Brechin, D., & Rubin, D. (2005). Visual memory loss and

749   autobiographical amnesia: a case study. *Neuropsychologia*, *43*(10), 1493–1502.

750   https://doi.org/10.1016/j.neuropsychologia.2004.12.009

751 Greenberg, D. L., & Knowlton, B. J. (2014). The role of visual imagery in autobiographical

752   memory. *Memory & Cognition*, *42*(6), 922–934. https://doi.org/10.3758/s13421-014-0402-

753   5

754 Hassabis, D., Kumaran, D., Vann, S. D., & Maguire, E. A. (2007). Patients with hippocampal

755   amnesia cannot imagine new experiences. *Proceedings of the National Academy of*

756   *Sciences*, *104*(5), 1726–1731.

757 Jessen, F., Spottke, A., Boecker, H., Brosseron, F., Buerger, K., Catak, C., Fliessbach, K., Franke,

758   C., Fuentes, M., Heneka, M. T., Janowitz, D., Kilimann, I., Laske, C., Menne, F., Nestor, P.,

759   Peters, O., Priller, J., Pross, V., Ramirez, A., … Düzel, E. (2018). Design and first baseline

760   data of the DZNE multicenter observational study on predementia Alzheimer's disease

761   (DELCODE). *Alzheimer's Research and Therapy*, *10*(1), 1–10.

762   https://doi.org/10.1186/S13195-017-0314-2/TABLES/1

763 Kay, L., Keogh, R., Andrillion, T., & Pearson, J. (2022). The pupillary light response as a

764   physiological index of aphantasia, sensory and phenomenological imagery strength. *ELife*,

765   *11*, e72484. https://doi.org/10.7554/eLife.72484

766 Keogh, R., Bergmann, J., & Pearson, J. (2020). Cortical excitability controls the strength of

767   mental imagery. *ELife*, *9*, e50232. https://doi.org/10.7554/eLife.50232

768 Keogh, R., & Pearson, J. (2018). The blind mind: No sensory visual imagery in aphantasia.

*Cortex*, *105*, 53–60. https://doi.org/10.1016/j.cortex.2017.10.012

Langille, J. J., & Gallistel, C. R. (2020). Locating the engram: Should we look for plastic synapses

or information-storing molecules? *Neurobiology of Learning and Memory*, *169*, 107164.

Leelaarporn, P., Dalton, M. A., Stirnberg, R., Stöcker, T., Spottke, A., Schneider, A., &

McCormick, C. (2024). Hippocampal subfields and their neocortical interactions during

autobiographical memory. *Imaging Neuroscience.*

Levine, B., Svoboda, E., Hay, J. F., Winocur, G., & Moscovitch, M. (2002). Aging and

autobiographical memory: dissociating episodic from semantic retrieval. *Psychology and*

*Aging*, *17*(4), 677–689. https://doi.org/10.1037/0882-7974.17.4.677

Maguire, E. A., & Mullally, S. L. (2013). The hippocampus: a manifesto for change. *Journal of*

*Experimental Psychology: General*, *142*(4), 1180–1189. https://doi.org/10.1037/a0033650

Marks, D. F. (1973). Visual imagery differences in the recall of pictures. *British Journal of*

*Psychology*, *64*(1), 17–24. https://doi.org/10.1111/j.2044-8295.1973.tb01322.x

Marks, D. F. (1995). New directions for mental imagery research. *Journal of Mental Imagery*,

*19*(3–4), 153–167. https://psycnet.apa.org/record/1996-29150-001

McCormick, C., Barry, D. N., Jafarian, A., Barnes, G. R., & Maguire, E. A. (2020). VmPFC Drives

Hippocampal Processing during Autobiographical Memory Recall Regardless of

Remoteness. *Cerebral Cortex*, *30*(11), 5972–5987. https://doi.org/10.1093/cercor/bhaa172

McCormick, C., Ciaramelli, E., De Luca, F., & Maguire, E. A. (2018). Comparing and Contrasting

the Cognitive Effects of Hippocampal and Ventromedial Prefrontal Cortex Damage: A

Review of Human Lesion Studies. *Neuroscience*, *374*, 295–318.

https://doi.org/10.1016/j.neuroscience.2017.07.066

McCormick, C., Rosenthal, C. R., Miller, T. D., & Maguire, E. A. (2017). Deciding what is possible

and impossible following hippocampal damage in humans. *Hippocampus, 27*(3), 303–314.

https://doi.org/10.1002/hipo.22694

McCormick, C., Rosenthal, C. R., Miller, T. D., & Maguire, E. A. (2018). Mind-wandering in

people with hippocampal damage. *Journal of Neuroscience, 38*(11), 2745–2754.

https://doi.org/10.1523/JNEUROSCI.1812-17.2018

McCormick, C., St-Laurent, M., Ty, A., Valiante, T. A., & McAndrews, M. P. (2015). Functional

and effective hippocampal-neocortical connectivity during construction and elaboration of

autobiographical memory retrieval. *Cerebral Cortex, 25*(5), 1297–1305.

https://doi.org/10.1093/cercor/bht324

Mechelli, A., Price, C. J., Friston, K. J., & Ishai, A. (2004). Where bottom-up meets top-down:

Neuronal interactions during perception and imagery. *Cerebral Cortex, 14*(11), 1256–1265.

https://doi.org/10.1093/cercor/bhh087

Miller, T. D., Chong, T. T. J., Davies, A. M. A., Johnson, M. R., Irani, S. R., Husain, M., Ng, T. W. C.,

Jacob, S., Maddison, P., Kennard, C., & others. (2020). Human hippocampal CA3 damage

disrupts both recent and remote episodic memories. *Elife, 9*, e41836.

Milton, F., Fulford, J., Dance, C., Gaddum, J., Heuerman-Williamson, B., Jones, K., Knight, K. F.,

MacKisack, M., Winlove, C., & Zeman, A. (2020). Behavioral and neural signatures of visual

imagery vividness extremes: aphantasia vs. hyperphantasia. *Cerebral Cortex*

*Communications, 2*(2), tgab035. https://doi.org/10.31234/OSF.IO/J2ZPN

Monk, A. M., Dalton, M. A., Barnes, G. R., & Maguire, E. A. (2021). The role of hippocampal–

ventromedial prefrontal cortex neural dynamics in building mental representations.

*Journal of Cognitive Neuroscience, 33*(1), 89–103.

Monzel, M., Keidel, K., & Reuter, M. (2021). Imagine, and you will find - Lack of attentional

guidance through visual imagery in aphantasics. *Attention, Perception {\&} Psychophysics*.

https://doi.org/10.3758/s13414-021-02307-z

Monzel, M., Keidel, K., & Reuter, M. (2023). Is it really empathy? The potential confounding role

of visual imagery in self-reports of empathy. *Journal of Research in Personality, e104354*.

https://doi.org/10.1016/j.jrp.2023.104354

Monzel, M., Mitchell, D., Macpherson, F., Pearson, J., & Zeman, A. (2022). Proposal for a

consistent definition of aphantasia and hyperphantasia: A response to Lambert and Sibley

(2022) and Simner and Dance (2022). *Cortex, 152*, 74–76.

https://doi.org/10.1016/J.CORTEX.2022.04.003

Monzel, M., Vetterlein, A., & Reuter, M. (2021). Memory deficits in aphantasics are not

restricted to autobiographical memory – Perspectives from the Dual Coding Approach.

*Journal of Neuropsychology, 16*(2), 444–461. https://doi.org/10.1111/jnp.12265

Moscovitch, M., Rosenbaum, R. S., Gilboa, A., Addis, D. R., Westmacott, R., Grady, C.,

McAndrews, M. P., Levine, B., Black, S., Winocur, G., & Nadel, L. (2005). Functional

neuroanatomy of remote episodic, semantic and spatial memory: A unified account based

on multiple trace theory. *Journal of Anatomy, 207*(1), 35–66.

https://doi.org/10.1111/j.1469-7580.2005.00421.x

O'Keefe, J. (1991). *The hippocampal cognitive map and navigational strategies.*

Pace, T., Koenig-Robert, R., & Pearson, J. (2023). Different Mechanisms for Supporting Mental

Imagery and Perceptual Representations: Modulation Versus Excitation. *Psychological*

*Science*, *34*(11), 9567976231198436. https://doi.org/10.1177/09567976231198435

Pearson, J. (2019). The human imagination: the cognitive neuroscience of visual mental

imagery. *Nature Reviews Neuroscience*, *20*(10), 624–634. https://doi.org/10.1038/s41583-

019-0202-9

Pearson, J., Clifford, C. W. G., & Tong, F. (2008). The Functional Impact of Mental Imagery on

Conscious Perception. *Current Biology*, *18*(13), 982–986.

https://doi.org/10.1016/j.cub.2008.05.048

Pearson, J., Rademaker, R. L., & Tong, F. (2011). Evaluating the mind's eye: The metacognition

of visual imagery. *Psychological Science*, *22*(12), 1535–1542.

https://doi.org/10.1177/0956797611417134

Rosenbaum, R. S., Moscovitch, M., Foster, J. K., Schnyer, D. M., Gao, F., Kovacevic, N., Verfaellie,

M., Black, S. E., & Levine, B. (2008). Patterns of autobiographical memory loss in medial-

temporal lobe amnesic patients. *Journal of Cognitive Neuroscience*, *20*(8), 1490–1506.

Schwitzgebel, E. (2002). How well do we know our own conscious experience? The case of

visual imagery. *Journal of Consciousness Studies*, *9*(5–6), 35–53.

https://www.jstor.org/stable/43154688

Sheldon, S., & Levine, B. (2013). Same as it ever was: Vividness modulates the similarities and

differences between the neural networks that support retrieving remote and recent

autobiographical memories. *NeuroImage*, *83*, 880–891.

St-Laurent, M., Moscovitch, M., Levine, B., & McAndrews, M. P. (2009). Determinants of

autobiographical memory in patients with unilateral temporal lobe epilepsy or excisions.

*Neuropsychologia*, *47*(11), 2211–2221.

Steinvorth, S., Levine, B., & Corkin, S. (2005). Medial temporal lobe structures are needed to re-

experience remote autobiographical memories: evidence from HM and WR.

*Neuropsychologia, 43*(4), 479–496.

Svoboda, E., McKinnon, M. C., & Levine, B. (2006). The functional neuroanatomy of

autobiographical memory: a meta-analysis. *Neuropsychologia, 44*(12), 2189–2208.

https://doi.org/10.1016/j.neuropsychologia.2006.05.023

Teyler, T. J., & DiScenna, P. (1986). The Hippocampal Memory Indexing Theory. *Behavioral*

*Neuroscience, 100*(2), 147–154. https://doi.org/10.1037/0735-7044.100.2.147

Wagner, S., & Monzel, M. (2023). Measuring imagery strength in schizophrenia: no evidence of

enhanced mental imagery priming. *Brain and Behavior, 13*(9), e3146.

https://doi.org/10.1002/brb3.3146

Wicken, M., Keogh, R., & Pearson, J. (2021). The critical role of mental imagery in human

emotion: insights from fear-based imagery and aphantasia. *Proceedings of the Royal*

*Society B, 288*(1946). https://doi.org/10.1098/RSPB.2021.0267

World Medical Association. (2013). World Medical Association Declaration of Helsinki: Ehtical

principles for medical research involving human subjects. *JAMA, 310*(20), 2191.

https://doi.org/10.1001/jama.2013.281053

Zeidman, P., & Maguire, E. A. (2016). Anterior hippocampus: the anatomy of perception,

imagination and episodic memory. *Nature Reviews Neuroscience, 17*(3), 173–182.

Zeman, A., Dewar, M., & Della Sala, S. (2015). Lives without imagery – congenital aphantasia.

*Cortex, 73*, 378–380. https://doi.org/10.1016/j.cortex.2015.05.019

Zeman, A., Milton, F., Della Sala, S., Dewar, M., Frayling, T., Gaddum, J., Hattersley, A.,

Heuerman-Williamson, B., Jones, K., MacKisack, M., & Winlove, C. (2020). Phantasia – the

psychological significance of lifelong visual imagery vividness extremes. *Cortex, 130*, 426–

440. https://doi.org/10.1016/j.cortex.2020.04.003