Successful regeneration of the adult zebrafish retina is dependent on inflammatory signaling

Oliver Bludau[1,2], Anke Weber[1], Viktoria Bosak[1], Veronika Kuscha[1], Kristin Dietrich[1],

Stefan Hans[1] & Michael Brand[1]*

1 - CRTD - Center for Regenerative Therapies, and PoL - Cluster of Excellence

Physics of Life, TU Dresden, Dresden, Germany.

2 – present address: Biomedical Center, LMU München, Munich, Germany.

*: author for correspondence

Email: michael.brand@tu-dresden.de

ORCID ID http://orcid.org/0000-0001-5711-6512

Word count: 8052 (excluding materials & methods, references)

Running title: Neuroinflammation as a regulator of adult retina regeneration

Key words: immune suppression/M-CSF/ microglia/Müller glia/NF-κB

pathway/regeneration/retina/zebrafish

Abstract

Inflammation can lead to persistent and irreversible loss of retinal neurons and photoreceptors in mammalian vertebrates. In contrast, in the adult zebrafish brain, acute neural inflammation is both necessary and sufficient to stimulate regeneration of neurons. Here, we report on the critical, positive role of the immune system to support retina regeneration in adult zebrafish. After sterile, ablation of photoreceptors by phototoxicity, we find rapid response of tissue-resident microglia and neutrophils, which returns to homeostatic levels within 14 days post lesion. Pharmacological or genetic impairment of immune cell reactivity results in a reduced Müller glia stem cell response, seen as decreased reactive proliferation, and a strikingly reduced number of regenerated cells from them, including photoreceptors. Conversely, injection of the immune stimulators flagellin, zymosan, or M-CSF into the vitreous of the eye, in spite of the absence of a retinal lesion, leads to a robust proliferation response and the up-regulation of regeneration-associated marker genes in Müller glia. Our results suggest that neuroinflammation is a necessary and sufficient driver for retinal regeneration in the adult zebrafish retina.

Introduction

Immune system activation is one of the first responses to tissue damage, e.g. by infection, disease or injury. Cells of the immune system (leukocytes) can recognize invading pathogens or factors that are secreted by damaged or dying cells (Ferrero-Miliani et al., 2007). Subsequently, leukocytes accumulate at the affected area, removing pathogens and clearing cellular debris, thus supporting the reestablishment of a physiological balance (Nathan & Ding, 2010). Conversely, if inflammation cannot be resolved, a detrimental chronic inflammation can occur, causing progressive tissue damage and pathology (Zhou et al., 2016).

In the lesioned mammalian central nervous system (CNS), accumulation of reactive astrocytes often results in a glial scar that acts as a barrier for successful regeneration (Buffo et al., 2008; Fitch & Silver, 2008; Sofroniew, 2009). Similarly, neurodegenerative diseases like Parkinson's or Alzheimer's show characteristics of chronic inflammation, causing subsequent neuronal death (Amor et al., 2010; Herrero et al., 2015; Kinney et al., 2018).

In contrast to mammals, lesioning of the zebrafish CNS results in a strong regenerative response despite the initial manifestation of inflammation (Kroehne et al., 2011; Kyritsis et al., 2012; Kizil et al., 2015; Bosak et al., 2018; Mitchell et al., 2018; Tsarouchas et al., 2018; White et al., 2017; Silva et al., 2020; Zhang et al., 2020). Whereas immune system activation in the mammalian CNS is typically detrimental for regeneration, studies in zebrafish demonstrated a strong beneficial link between an immune response and neural stem cell reactivity (Kyritsis et al., 2012; Aurora & Olson, 2014; Bosak et al., 2018; Tsarouchas et al., 2018). Remarkably, in the zebrafish adult telencephalon, inflammation is required to initiate

a successful regenerative response, and a lipid inflammatory cue, leukotriene-C4, is sufficient to stimulate proliferation of radial glia-type stem cells (Kyritsis et al., 2012; Kizil et al., 2015).

In the retina, zebrafish Müller glial cells similarly act as stem cells, and generate neuronal precursor cells (NPCs) in response to retinal lesion (Lenkowski & Raymond, 2014; Goldman, 2014; Gorsuch & Hyde, 2014). These cells then amplify, migrate to the lesion site, and differentiate into the lost neuronal subtypes and thus gradually restore vision (Raymond et al., 2006; Hammer et al., 2022). Dying neurons release the proinflammatory cytokine TNF-α, triggering the regenerative response of Müller glia (Conner et al., 2014; Nelson et al., 2013). Likewise, the inflammation-associated factors Interleukin-11 and TGF-ß stimulate Müller glia cell cycle re-entry and NPC generation (Lenkowski et al., 2013; Zhao et al., 2014). Furthermore, microglia – the CNS tissue resident macrophages – support this initial regenerative response by secreting proinflammatory factors (Kizil et al., 2015; Conedera et al., 2019; Iribarne & Hyde, 2022; Zhang et al., 2020). However, to date the molecular pathways involved in damage recognition, stem cell proliferation, neuronal precursor cell amplification and differentiation during retinal regeneration are poorly understood; in particular, the role of inflammation is unclear (Lahne et al., 2020; Lenkowski & Raymond, 2014; Mitchell et al., 2018, 2019; Iribarne & Hyde, 2022).

Here, we analyze the contribution of inflammation to regeneration using a non-invasive, sterile phototoxic ablation model of photoreceptor cells, as the key cell type affected by retinal disease, in the adult zebrafish retina (Mitchell et al., 2018; Silva et al., 2020; Weber et al., 2013; Zhang et al., 2020). Following light lesion, we observe strong convergence of tissue resident microglia to the lesion site. Blocking inflammation pharmacologically caused reduced reactive proliferation of Müller glia

stem cells and impaired photoreceptor regeneration. Similarly, in a genetic model of

microglia deficiency, we find that microglia are required to support Müller cell

proliferation. Conversely, when the immune stimulators flagellin, zymosan, or M-CSF

are injected into the vitreous of the eye, notably in the absence of retinal lesion,

Müller glial cells are triggered to undergo reactive proliferation and regeneration-

associated marker gene expression. Taken together, our results show that in the

regeneration-competent adult zebrafish retina, acute inflammation is an important

positive regulator of retina regeneration.

Results

Leukocytes react to sterile ablation of photoreceptor cells

The immune system of vertebrates rapidly responds to retinal damage. In larval zebrafish, *mpeg1*:mCherry positive monocytes react to photoreceptor ablation by rapid migration towards the site of lesion (White et al., 2017). Similarly, microglia respond by accumulation and phagocytosis of debris in response to neurotoxic ablation of inner retinal cells in adult zebrafish (Mitchell et al., 2018). To investigate if leukocytes are recruited in an injury model at adult stages, we used intense diffuse light that causes sterile ablation of all photoreceptor subtypes by phototoxicity (Weber et al., 2013). In this model, due to the refractive properties of the adult zebrafish visual system, the photoreceptors in a central stripe of the retina are ablated, whereas ventral and dorsal retina is much less affected and conveniently serves as an internal control area (Weber et al., 2013; Fig. 1 D). We analyzed the accumulation and appearance of leukocytes in the central retina on sections by immunohistochemistry for L-Plastin, a pan-leukocyte marker (Redd et al., 2006; Kroehne et al. 2011) in *Tg(mpeg1:mCherry)* reporter animals, labelling the monocyte lineage, including macrophages and microglia (Figure 1A). In comparison to unlesioned (sham) controls, L-Plastin positive cells accumulated in the central lesion zone already at 2 day post lesion (dpl; Figure 1A). Whereas leukocytes showed a ramified morphology in sham control retinae, their appearance changed to an amoeboid and swollen shape at 2 dpl (insets in Figure 1A, D), as described by Mitchell et al., 2018 upon neurotoxic lesion. At 2 dpl an increased number of L-Plastin positive leukocytes at the outer nuclear layer (ONL) in comparison to sham controls is revealed (Figure 1A, B). In flatmounts of *Tg(mpeg1:mCherry) x Tg(opn1sw1:GFP)* double transgenic zebrafish that express GFP as a marker in the

entire UV cone population, displaying the central lesion by the absence of GFP, the

accumulation of monocytes can nicely be observed (Figure 1D). While in sham,

microglia display an equal distribution and a ramified structure they accumulate in the

central lesion area. Interestingly they disappear from the unlesioned peripheral

regions. In a time course, the number of L-Plastin positive cells peaked at 2 dpl and

subsequently declined to sham levels at 14 dpl (Figure 1B). To further determine the

identity of retinal leukocytes, we analyzed immunoreactivity of L-Plastin in transgenic

Tg(mpeg1:mCherry) animals.In sham control retinae, *mpeg1*:mCherry positive cells

always co-expressed L-Plastin, indicating that all tissue resident homeostatic

leukocytes were of the monocyte lineage, namely microglia (Figure 1A, S1A). In

contrast, additional L-Plastin positive, but *mpeg1*:mCherry negative cells could be

detected upon injury at 2 dpl (insets in Figure 1A). Subsequent quantification of the

number of L-Plastin and *mpeg1*:mCherry double positive cells revealed a decrease

from 99% to approximately 83%, indicating the presence of another cell type of the

leukocyte lineage at the lesion site (Figure S1A). Neutrophils are known to rapidly

respond to tissue damage and are labeled by transgenic *Tg(mpo:GFP)* (Renshaw et

al., 2016; Kurimoto et al., 2013; Figure 1D, S1B). In contrast to sham controls, which

never showed any *mpo*:GFP positive cells in the homeostatic retina, we observed

rapid accumulation of neutrophils after light lesion. *mpo*:GFP positive cells first

appeared in the retina by 12 hpl, and a diffuse GFP positive pattern (referred to as

matrix), in the outer segment layer and the inner nuclear layer (INL) was present from

15 hpl onwards (Figure 1C), presumably reflecting neutrophil NETosis (Zhu et al.,

2021). Analyses of *Tg(mpeg1:mCherry)* and *Tg(mpo:GFP)* double transgenic animals

revealed that *mpeg1*:mCherry positive cells co-localized with the GFP positive matrix,

suggesting an uptake of the matrix material by monocytes (Figure S1B). T cells of the

lymphoid lineage were reported to stimulate Müller glia proliferation, and to augment

retina regeneration in a stab wound assay (Hui et al., 2017). To examine if T cells are

also recruited to the lesion site, we used transgenic *Tg(lck:NLS-DsRed)* animals, but

could not detect any *lck* positive T cells, neither in the homeostatic nor in the

regenerating retina up to 7 days post lesion (Figure S2).

Taken together, our data show a strong accumulation and activation of innate

immune cells following a sterile light lesion that is resolved by 14 dpl, consistent with

the mounting of an acute inflammatory response after retinal injury.

Müller glia activate NF-κB signaling in response to injury

In zebrafish, Müller glia have key functions in the regulation of retinal homeostasis as

well as during regeneration (Lenkowski & Raymond, 2014). To study how Müller glia

react to injury during regeneration and inflammation, we analyzed the activation of

the proinflammatory signaling pathway NF-κB using *Tg(NF-κB:GFP)* and

Tg(gfap:NLS-mCherry) transgenic animals. The *Tg(NF-κB:GFP)* line reports

canonical NF-κB activity via the expression of GFP under the control of six human

NF-κB binding motifs driving a c-fos minimal promoter (Kanther et al., 2011). In sham

control retinae, *NF-κB*:GFP expression was only found in the retinal vasculature,

microglia and, occasionally in Müller glia residing in the inner nuclear layer (Figure

2A). In sharp contrast, robust activation of *NF-κB*:GFP was present in numerous

gfap:NLS-mCherry positive Müller glia already at 1 dpl (Figure 2A). Moreover, *NF-*

κB:GFP expression was present in cells located in the ONL showing the

characteristic morphology of photoreceptor cells. Both cell types continued to express

NF-κB:GFP at 2 dpl. At 4 dpl, *NF-κB*:GFP expression was absent from the ONL but

remained detectable in Müller glia. To determine if NF-κB activation might be

functionally relevant, we investigated the expression of *matrix metallopeptidase 9*

(*mmp9*), a known downstream target of NF-κB with important functions in degrading

extracellular matrix and chemokines (Cheng et al., 2012; LeBert et al., 2015; Xu et

al., 2018; Yang et al., 2017). Thus, we performed *mmp9 in situ* hybridization in

combination with immunohistochemistry against glial fibrillary acidic protein

(GFAP/Zrf-1) labeling Müller glia and proliferating cell nuclear antigen (PCNA)

labeling cells in S-phase and shortly after. In contrast to sham controls that never

showed *mmp9* expression in the homeostatic retina, *mmp9* was strongly expressed

by Zrf-1 positive Müller glia at 1 dpl, indicating *de novo mmp9* expression in these

cells (Figure 2B). Moreover, numerous *mmp9* and Zrf-1 expressing cells were also

positive for PCNA, identifying them as reactively proliferating Müller glia in response

to injury. Expression of *mmp9* remained strong at 2 dpl, but dropped below detection

levels at 4 dpl, consistent with transcriptome data for *mmp9* (Kramer et al., 2021;

Silva 2020; Celotto et al., 2023). Taken together, our results showed that the NF-κB

signaling pathway is transiently activated in Müller glia in response to a sterile

ablation of photoreceptor cells, indicative of an inflammatory response by Müller glia.

Dexamethasone-mediated immunosuppression reduces retinal regeneration

To study the role of the immune system during retinal regeneration, we used

Dexamethasone (Dex), a potent immunosuppressant (Coutinho & Chapman, 2011;

Donika Gallina et al., 2015; Kyritsis et al., 2012; Silva et al., 2020; Zhang et al.,

2020). Experimental zebrafish were treated with Dex from 10 days prior to lesion until

the time point of analysis (Figure 3A). Vehicle control experiments were carried out

with the respective amount of the solvent methanol (MeOH). In contrast to MeOH-

treated controls, Dex-treatment for 10 consecutive days resulted in an overall

reduction of retinal microglia (Figure 3B). Similarly, leukocyte recruitment upon light

lesion was significantly reduced in Dex-treated animals at 2 dpl, compared to control

(Figure 3B). Quantification of L-Plastin positive cells corroborated a significant

reduction at all indicated time points (Figure 3C). In addition, Dex-treatment reduced

the number of resident retinal microglia during homeostasis, and no increase in

leukocyte number was noted after lesion. To address if the observed changes might

be caused indirectly by the known neuroprotective properties of Dex (Gallina et al.,

2015), we analyzed cell death using the TdT-mediated dUTP-digoxigenin nick end

labeling (TUNEL) assay; however, we did not observe any difference in the number

of TUNEL positive cells after light lesion in control MeOH- or Dex-treated animals

(Figure S3).

Reactive proliferation of Müller glia is a hallmark of retinal regeneration; we therefore

further analyzed if Dex-treatment influences proliferation after sterile light lesion using

immunolabeling of the proliferation marker PCNA (Figure 3D). In contrast to MeOH-

treated controls, Dex-treated animals showed a significant reduction in PCNA

positive cells at 4 dpl. Quantification of PCNA positive cells during the course of

regeneration revealed that reactive proliferation, driven by Müller cells and neuronal

progenitors, is significantly reduced in Dex-treated animals at all time-points

examined in comparison to MeOH controls. Furthermore, homeostatic proliferation

(most likely by cells of the ciliary margin) in unlesioned retinae is not affected (Figure

3E). To investigate if Dex-treatment specifically affects Müller glia proliferation, we

analyzed the number of PCNA positive cells in transgenic *Tg(gfap:NLS-GFP)*

animals, which express strong nuclear GFP in all Müller glia (with some leakage of

GFP to the cytoplasm, identifying the characteristic Müller glia cell shape, Figure 3F).

Indeed, in comparison to MeOH-treated animals, the number of PCNA and *gfap*:NLS-

GFP double positive cells was significantly reduced after Dex-treatment.

Quantification showed that the percentage of Müller glia co-localizing with PCNA is

reduced to less than 50% at all analyzed time points (Figure 3G). This suggests that

Müller glia reactivity is impaired upon Dex-treatment. Consistent with this possibility,

NF-κB activation is reduced in Müller glia after Dex-treatment, as seen in the *Tg(NF-*

κB:GFP) activation reporter line (Figure 3H). As expected, cells in the ONL and

Müller glia show a robust *NF-κB*:GFP expression in MeOH-treated control animals at

2 dpl, whereas a reduced number of cells in the ONL activated the *NF-κB*:GFP

transgene and no GFP positive Müller glia could be detected at 2 dpl after Dex-

treatment. Consistently, activation of the NF-κB downstream target *mmp9* was almost

absent in Dex-treated animals at 2 dpl, compared to MeOH controls (Figure 3I).

Taken together, these results show that Dex-mediated immunosuppression efficiently

reduces the accumulation of leukocytes and impairs Müller glia reactivity, as well as

reactive proliferation and regeneration-associated marker gene expression, during

retinal regeneration.

236 Immunosuppression reduces regeneration of photoreceptors

To examine if the reduced cell proliferation has consequences for the regenerative

outcome, we probed the effect of Dex-immunosuppression on restoration of

photoreceptors after light lesion. Thus, we performed Dex-treatment and light lesions

on *Tg(opn1sw1:GFP)* animals expressing GFP in all UV cones, followed by repeated

EdU injections to label newborn cells, and analyzed the animals at 28 dpl when

cellular regeneration is completed (Figure 4A). Consistent with the afore-mentioned

decrease of retinal leukocytes after Dex-treatment, we observed an almost complete

loss of retinal leukocytes after 28 days of continuous immune suppression (Figure

S4). In comparison to MeOH-treated control animals, the overall number of EdU

positive cells was significantly lower after Dex-treatment at 28 dpl (Figure 4B).

Similarly, the number of EdU positive UV-cones (Figure 4C) was severely decreased

in Dex-treated animals compared to MeOH-treated controls. Moreover, in addition to

a reduction in number, EdU positive UV cones appeared malformed with improperly

shaped outer segments in Dex-treated animals (insets Figure 4B). Quantifications

showed that the number of EdU positive nuclei was similarly reduced in all retinal

layers (Figure 4C). In conclusion, Dex-mediated immunosuppression interferes with

reactive proliferation and reactive neurogenesis, indicating that the injury-induced

immune response is essential for proper retina regeneration.

Retinal microglia support reactive Müller glia proliferation

Activated microglia and macrophages clear debris from dead cells, and they interact

with Müller glia and influence their cellular response (Aslanidis et al., 2015; Keightley

et al., 2014; Palazzo et al., 2020; Wang et al., 2011). Consistently, our results show

that Dex-mediated immunosuppression interferes with reactive proliferation and

neurogenesis, indicating that the injury-induced immune response is essential for

successful retina regeneration. To independently test this notion, we investigated

whether microglia positively contribute to reactive proliferation during retina

regeneration, using Interferon regulatory factor 8 (*irf8*) myeloid-defective mutants to

genetically deplete microglia in embryonic and juvenile fish (Shiau et al., 2015). To

verify leukocyte deficiency at adult stages, we analyzed L-Plastin immunoreactivity in

homozygous adult *irf8* mutants and heterozygous control siblings, and found that the

number of leukocytes in the homeostatic retina was significantly reduced in *irf8*

mutants (Figure 5A, B). Next, we asked whether the microglia deficiency also affects

the proliferation of Müller cells and NPCs during regeneration. We therefore combined the light lesion paradigm with EdU-labeling at 3 dpl prior to analysis at 4 dpl (Figure 5C). Compared to heterozygous control siblings, the number of accumulated L-Plastin positive cells was clearly reduced in *irf8* mutant animals at 4 dpl (Figure 5D). Similarly, the number of EdU positive cells is lower in *irf8* mutants at 4 dpl compared to heterozygous controls (Figure 5D). Quantification of both the number of L-Plastin positive cells as well as the number of EdU positive cells confirmed a statistically significant reduction (Figure 5E). Thus, consistent with our above findings using Dex-mediated immunosuppression, these results show that genetic depletion of leukocytes correlates with decreased reactive proliferation upon injury, supporting the notion of an important role of immune cells during zebrafish retina regeneration.

Inflammatory stimuli trigger a regeneration response in Müller glia in the absence of lesion

We hypothesized that inflammation might not only be necessary, but also sufficient, to trigger reactive proliferation of Müller glia and the generation of NPCs. To address this notion, we injected different immune activators into the vitreous of the eye - thus leaving the retina unlesioned (Fig 6A) - and analyzed the response of Müller glia at 2 and 4 days post injections (dpi; Figure 6). The toll-like receptor (TLR) agonist zymosan has previously been shown to stimulate an inflammatory response both in the zebrafish and mouse eye, and in the zebrafish telencephalon (Kyritsis et al., 2012; Kurimoto et al., 2013; Zhang et al., 2020). Next, we tested injections of flagellin, the principal structural protein of bacterial flagella, which is known to cause sepsis in zebrafish larvae (Barber et al., 2016). As a control, to ensure that injection

of both zymosan and flagellin did not cause retinal cell death and thereby trigger a

regenerative response indirectly, we performed TUNEL assays at 2 dpi. Neither

zymosan nor flagellin injection resulted in a significant increase of TUNEL positive

cells in comparison to control PBS-injected eyes (Figure 6B). Interestingly, the

number of L-Plastin positive cells also did not significantly increase after injecting

zymosan or flagellin (Figure 6C). In contrast, cell proliferation using PCNA

immunohistochemistry is clearly induced in both zymosan as well as flagellin-injected

animals in comparison to PBS-injected controls (Figure 6D). Quantification of PCNA

positive nuclei confirmed a significant increase in zymosan and flagellin-injected

animals at 4 dpi, but not at 2 dpi (Figure 6E). Furthermore, we analyzed activity of

NF-κB after injections of PBS, zymosan and flagellin using the *NF-κB*:GFP reporter

line. Consistent with the afore-mentioned expression in non-injected retinae, PBS-

injected controls show *NF-κB*:GFP expression in blood vessels, microglia and few

Müller glia residing in the INL (Figure 6F). In contrast, *NF-κB*:GFP is strongly

activated in additional cells of the inner and outer nuclear layer in zymosan and

flagellin-injected animals at 2 dpi. Again, GFP positive cells in the ONL showed

mostly the characteristic morphology of photoreceptor cells (see also above, Fig. 2B

& 3H), in particular in the zymosan-injected animals. Consistent with the activation of

the *NF-κB*:GFP reporter, retinal expression of *mmp9* is also found in zymosan and

flagellin-injected animals, but not in PBS-injected controls at 2 dpi (Figure 6G).

Finally, we analyzed the expression of the transcription factor *her4.1*, a downstream

target of the Notch signaling pathway, which is upregulated in NPCs generated in

response to photoreceptor damage (Wan et al., 2012). No expression of *her4.1* is

detected in PBS-injected controls, but *her4.1* expression is activated in zymosan and

flagellin-injected samples at 4 dpi (Figure 6H). These results show that activators of

the immune system can strongly stimulate Müller glia reactivity and proliferation,

even in the absence of tissue damage.

M-CSF injection triggers leukocyte accumulation and Müller glia reactivity in

the absence of a lesion

In order to identify the stimulatory potential of specific individual inflammatory

mediators, we focused on macrophage colony-stimulating-factor (M-CSF/CSF-1),

based on preliminary data from our recent single-cell RNAseq dataset of

regenerating retina (Celotto et al., 2023), and using the same injection paradigm as

before (Figure 7A). In mammals, M-CSF stimulate CSF-1 receptor signaling, which is

involved in monocyte colonization and stimulation (Chitu et al., 2016; Wu et al.,

2018). Injection of human M-CSF induced an increased cell proliferation at 4 dpi

compared to PBS controls (Figure 7B, C), despite the absence of unspecific damage

to the retina, as seen by lack of an increase in TUNEL positive cells at 2 dpi.

Quantification corroborated a significant increase of proliferating cells, scored as

PCNA positive nuclei, in M-CSF-injected retinae, compared to PBS-injected controls

(Figure 7D). To address if M-CSF causes an increase in the number of retinal

leukocytes, we performed immunolabeling of L-Plastin in M-CSF- and PBS-injected

specimens. We observed a significant accumulation of L-Plastin positive cells in M-

CSF-, but not in PBS-injected, animals at 4 dpi (Figure 7E & F). Further, we found

augmented NF-κB activity after M-CSF injection in *NF-κB*:GFP reporter animals.

PBS-injected controls showed the homeostatic *NF-κB*:GFP expression described

above (Figure 7G). In sharp contrast, additional *NF-κB*:GFP positive cells were found

in the M-CSF-injected animals at 2 dpi, located in the inner nuclear layer and

showing the characteristic morphology of Müller glia. Photoreceptor cells in the ONL

did not activate the *NF-κB*:GFP reporter following M-CSF injection whereas they do

so upon lesion (see above, Fig. 2B & 3H), which we tentatively suggest to reflect a

more selective role of M-CSF signaling for Müller glia. Consistent with the activation

of *NF-κB*:GFP, *mmp9* was also found to be expressed in M-CSF injected animals,

but not in PBS-injected controls at 2 dpi (Figure 7H). Taken together, these data

demonstrate that human recombinant M-CSF can induce leukocyte accumulation in

the neuronal retina of zebrafish, and can stimulate Müller glia reactive proliferation

and alter gene expression even in the absence of a lesion.

Discussion

In contrast to mammals, the zebrafish retina readily regenerates photoreceptor cells that are lost following a phototoxic lesion. Several signaling pathways have been implicated in the restoration of lost neurons (Lahne et al., 2020; Lenkowski & Raymond, 2014; Goldman, 2014; Hochmann et al., 2012). A role for the immune system in regeneration of the adult zebrafish retina has previously been suggested (Mitchell et al., 2018; Sifuentes et al., 2016; Silva et al., 2020; White et al., 2017; Zhang et al., 2020; Iribarne et al., 2022), but it is not understood in detail. Here, we confirm and extend these previous studies on the role of inflammation, and its impact on Müller glia reactivity, by modulating immune activity during regeneration of the adult zebrafish retina after a sterile phototoxic ablation of photoreceptor cells. We find that microglia and neutrophils infiltrate and accumulate at damaged sites of the retina. Importantly, Müller glia themselves react by activating the proinflammatory NF-κB signaling pathway in response to retinal damage. Our functional studies show that Dexamethasone-mediated immunosuppression reduces (i) leukocyte accumulation, and (ii) the reactive Müller glia response at the proliferative and gene expression level, and thus reduces the regeneration of photoreceptors. (iii) Conversely, in gain-of-function assays, injection of flagellin, zymosan or M-CSF as inflammatory factors that are thought to facilitate the immune system, some Müller glia are stimulated and reactive proliferation is induced (Gorsuch & Hyde, 2014; Lenkowski et al., 2013, Nelson et al., 2013; Figure 8). Overall, we suggest that cells of the innate immune system act as necessary and sufficient positive regulators for successful regeneration of the adult zebrafish retina. Regulators of the inflammatory state, such as Dexamethasone and the M-CSF that stimulates Müller glia-based photoreceptor regeneration in our injection assays, might therefore provide advanced

options for clinical treatments of retinal disease, as recently also suggested for

corticosteroid management of early phases of spinal cord injury (Nelson et al.,

2019).

Sterile ablation of photoreceptors recruits innate immune cells

Consistent with previous studies, we find that retinal *mpeg1*:mCherry positive tissue-

resident microglia display a ramified morphology in homeostasis, but change to an

activated amoeboid cell shape upon a sterile light lesion, similar to what is observed

after CNS lesion (Karlstetter et al., 2010; Kroehne et al., 2011; Kyritsis et al., 2012;

Mitchell et al., 2018). Along with activation, we find a strong accumulation of microglia

at the lesion site, which is resolved over a period of 14 days. The underlying dynamic

events indicate an acute inflammatory response of leukocytes that is tightly controlled

in a spatiotemporal manner. We assume that it is mostly tissue-resident microglia

that mediate the observed immune response; however, we cannot exclude that

additional peripheral macrophages are also recruited from the bloodstream and

contribute to the observed reaction, in particular since we also observe neutrophils

entering the retina. Consistent with our assumption, live imaging in larval zebrafish

revealed that peripheral macrophages do not enter the developing retina during drug-

mediated ablation of rod photoreceptors (Oosterhof et al., 2017; White et al., 2017).

In contrast, neurotoxic ablation of the ganglion cells and neurons of the inner nuclear

layer using ouabain triggers proliferation of adult retinal microglia, and it has been

suggested that the observed increase in cell number might also involve peripheral

macrophages entering the retina from circulation (Mitchell et al., 2018). The

observation of *mpo*:GFP positive neutrophils in the vasculature, as well as neutrophil

accumulation in the photoreceptor outer segment layer upon light lesion, indicates a

certain permeabilization of the blood-retina-barrier, which may also allow additional

leukocytes to invade the damaged retina (Eshaq et al., 2017; McMenamin et al.,

2019; Sim et al., 2015). Further studies of the cell-permeability of the blood-retina-

barrier during regeneration, e.g. by *in vivo* imaging approaches, and specific labeling

of either peripheral macrophage or microglia populations, will likely be interesting in

this regard.

Our observation that neutrophils arrive in the retina following non-invasive light lesion

suggests an interesting aspect of retina regeneration, with parallels to observations

after neurotoxic ablation of retinal ganglion cells (Mitchell et al., 2018). Neutrophils

have different functions at the site of injury, and they are among the first cells to react

to injury. It has been proposed that they participate in clearing cellular debris, by

secretion of factors influencing angiogenesis and regeneration, but they may also

support resolution of inflammation (Wang, 2018). In mice, neutrophils promote the

regeneration of the optic nerve by expression of oncomodulin, and can influence the

response of glial cells and axonal outgrowth after spinal cord injury (Kurimoto et al.,

2013; Stirling et al., 2009; Tsarouchas et al., 2018). Furthermore, neutrophils are

indicated to resolve inflammation by a localized O_2 depletion (Campbell et al., 2014).

Thus, neutrophils could be involved in the mediation of the immune response in adult

retina regeneration in zebrafish upon lesion. Further investigation of the observed

GFP positive matrices may therefore be interesting. Here, we speculate that these

structures might act as 'extracellular-traps' that could prevent cellular debris from

spreading further into the surrounding tissues, protecting the latter from secondary

cell death (Branzk & Papayannopoulos, 2013; Walker et al., 2007), which might help

suppress a systemic inflammatory/regenerative reaction in the retina. The

observation of GFP positive inclusions in mpeg1 positive macrophages furthermore

suggests degradation of the trap-like structures by macrophage uptake (Farrera &

Fadeel, 2013).

While microglia and neutrophils seem to play important roles in retina regeneration,

we did not observe any infiltration of *lck*:NLS-dsRed positive T cells into the retina

after lesion, arguing that T cells may be dispensable at least for the initial

regeneration response. A recent report suggests that T cells can positively influence

the regenerative response after an invasive mechanical lesion of the retina (Hui et al.,

2017); this difference might be due to differences in the lesion paradigm employed,

and/or timing of the regeneration events. As mentioned above, the non-invasive light

lesion that we employed results in a specific ablation of photoreceptors without

physically disrupting the blood-retina barrier. In contrast, the needle stab employed

by Hui et al., 2017 disrupts the blood-retina barrier, thereby allowing T cells to access

the lesion.

Müller glia react to photoreceptor cell death in an inflammatory manner

Upon retinal injury and progressive dystrophies, mammalian Müller glia undergo

physiological changes and show transcriptional alterations as a hallmark of their

gliotic response (Bringmann et al., 2009; Bringmann & Wiedemann, 2011). Despite

the regenerative response of zebrafish Müller glia, they exhibit initial signs of reactive

gliosis following retinal damage (Thomas et al., 2016). Here, we report that ablation

of photoreceptors induces an acute inflammatory response at the molecular level.

Importantly, we observed an inflammatory-like response directly in the lesion-

responding Müller glia. Using *NF-κB*:GFP reporter fish (Kanther et al., 2011), we find

that canonical NF-κB becomes activated in Müller glia upon damage, in keeping with transcriptomic changes of Müller glia upon light lesion (Sifuentes et al., 2016; Celotto et al., 2023). In the regenerating chick retina, NF-κB is active in Müller glia and acts as an important pathway controlled by factors that are secreted by microglia (Palazzo et al., 2020). A potential activator of NF-κB may be TNFα; indeed, several studies suggested that dying neurons secrete this factor and thus trigger Müller glia proliferation and the generation of NPCs (Kumar & Shamsuddin, 2012; Nelson et al., 2013).

Consistent with previous reports, we also found the NF-κB target gene, matrix metallopeptidase *mmp9,* to be expressed in reactive Müller glia, displaying a highly dynamic and strictly spatiotemporally controlled activation profile (Kaur et al., 2018; Kim et al., 2007; Silva et al., 2020; Takada et al., 2004). In *mmp9* knockout zebrafish, Müller glia-derived progenitors show a hyperproliferative response after injury, but reduced survival of regenerated photoreceptors (Silva et al., 2020). The exact function of *mmp9* is not completely clear yet; we speculate that *mmp9,* as a metalloproteinase, might be involved in the degradation of extracellular matrix (ECM), thus 'opening' the retina for incoming immune or neuronal precursor cells, and/or modulating ECM interacting cytokines (Gong et al., 2008; LeBert et al., 2015; Xu et al., 2018; Yoong et al., 2007). Interestingly *mmp9* is a key marker gene known to be differentially expressed between wet and dry forms of age-related macular degeneration, providing further evidence for a possibly critical role during inflammatory processes (Fritsche et al., 2016). Understanding the contribution and molecular control of *mmp9* in zebrafish retina regeneration may therefore provide further insights into the mechanisms and molecules that are required to restore lost neural tissues.

Immunosuppression alters the regenerative response of the zebrafish retina

Immunosuppression with the glucocorticoid dexamethasone drastically decreases the
amount of retinal microglia and blocks accumulation of reactive leukocytes in the
retina after lesion (Gallina et al., 2014; White et al., 2017). In addition, activation of
NF-κB signaling was also successfully blocked in Müller glia, as shown by the loss of
NF-κB:GFP and expression of its target gene. NF-κB activation has been described
in a plethora of cellular functions such as cell survival, cell death and mediation of
inflammation by the regulation of cytokine expression, as well as cell proliferation and
migration (Liu et al., 2017). In Müller glia, NF-κB signaling could mediate an
inflammatory-like response, consistent with their expression of pro- and anti-
inflammatory cytokines (Nelson et al., 2013; Polager & Ginsberg, 2002; Zhao et al.,
2014). Consistently, following Dex-treatment, we find that the reactive proliferation of
Müller glia was strongly reduced, indicating that inflammatory-like signaling promotes
Müller glia cell cycle reentry. Interestingly, homeostatic proliferation, which mainly
occurs in the ciliary marginal zone and only sporadically in the central retina, remains
unaffected, similar to our findings in the zebrafish telencephalon (Kyritsis et al.,
2012). Moreover, Dex-treatment is not only affecting reactive proliferation, as is also
confirmed by other research groups (Iribarne & Hyde, 2022; Silva et al., 2020; White
et al., 2017). Regenerated UV-cones show dysmorphic outer segments, indicating
that Dex-treatment also impairs the proper differentiation of photoreceptor cells. On
the other hand, in mmp9 mutants, initial Dex-treatment supports the restoration of the
photoreceptors and thus can act positively (Silva et al., 2020). Collectively, we
propose that inflammatory signaling is required for cell cycle reentry of Müller glia
during regeneration, but might also play additional roles during maturation of
regenerated photoreceptors (Figure 8). An incomplete regenerative outcome was

also detected in larval retinal pigment epithelium (RPE) regeneration in *irf8*-mutants

deficient in microglia. Upon ablation of RPE, the neural retina showed an increased

reactive proliferation in mutants, but the amount of regenerated RPE cells is lower

than in wild type siblings (Hanovice et al., 2019).

To further understand the impact of resident microglia on the regeneration process in

adult retina, we addressed the proliferative potential of retinae in response to injury in

irf8 mutant fish, as a genetic model for myeloid deficiency in larval and juvenile

zebrafish (Shiau et al., 2015). We confirm microglia deficiency at adult stages (>6

month), indicating that *irf8* is required throughout lifetime in primitive as well as a

definitive wave of monocyte populations of the eye (Ferrero et al., 2018; Shiau et al.,

2015). However, a reduced number of leukocytes is still found to populate the adult

retina. Upon lesion, we observed that the number of cells that are in cell cycle is

significantly reduced in *irf8* mutants in comparison to heterozygous siblings in

response to lesion. Interestingly, the number of L-Plastin+ leukocytes in *irf8* mutants

increases upon injury, but never reaches levels comparable to heterozygous control

siblings. In line with other reports, these findings indicate that microglia are

supportive for Müller cell proliferation and NPC amplification and thereby support

initial steps of regeneration (Conedera et al., 2019; Zhang et al., 2020; Iribarne &

Hyde, 2022). The necessity of (tissue-resident) macrophages is also observed in

other tissues, e.g. telencephalon, spinal cord, caudal fin and heart, showing the

contribution of the innate immune system to tissue regeneration (Kyritsis et al., 2012;

De Preux Charles et al., 2016; Petrie et al., 2015; Portillo et al., 2017; Roche et al.,

2018; Tsarouchas et al., 2018). This supportive role of macrophages in zebrafish is

unlike that of the rodent retina, where microglia appear to inhibit Müller-glia-mediated

tissue regeneration by suppression of *ascl1* expression. This interesting species

difference might be explained by distinct expression profiles of zebrafish vs. mouse

microglia during the time course of regeneration (Issaka Salia & Mitchell, 2020;

Mitchell et al., 2019; Todd et al., 2020).

Immune stimulation triggers Müller glia reactivity and proliferation

Retinal pathology is often coupled to neuronal loss, activation of the immune system

and Müller glia reactivity (Iribarne & Hyde, 2022). In particular, mammalian Müller glia

react to injury mostly by (proliferative) gliosis, which contributes to glial scarring, thus

enhancing loss of vision (Bringmann et al., 2006). In zebrafish, retinal injury also

results in an acute immune reaction; however, in contrast to mammals, this is

followed by a regenerative response.

Importantly, inflammatory signals can indeed initiate reactive proliferation, even in the

absence of injury, as seen by our vitreous injections of toll-like receptor (TLR)

agonists zymosan and flagellin to trigger an inflammatory response (Akira & Takeda,

2004; Barber et al., 2016; Zhang et al., 2020). Leukocyte number was not increased

following these injections, which might be explained by three possible scenarios: i)

Zymosan and flagellin do not stimulate leukocyte proliferation ii) recruitment from the

bloodstream is not induced; while in both scenarios a local leukocyte activation can

occur. iii) The accumulation could already be resolved within 48h, since the response

of immune cells is rapid and highly dynamic. Furthermore, the injected concentration

of zymosan in a rodent model, where leukocytes are seen to accumulate, is ten-fold

higher than the concentration used in our injections, and could reflect a dose

dependency (Kurimoto et al., 2013). In contrast to the more generic inflammatory

stimuli, M-CSF-injected retinae show an increased number of leukocytes. This may

be the consequence of both proliferation and recruitment from the bloodstream, since

M-CSF receptor (CSF-1R) is thought to regulate the population of microglia in neural

tissue during development (Wu et al., 2018). Following M-CSF injection, we find that

Müller glia activate NF-κB signaling, which fits well with studies reporting TLR and

CSF-1 receptor expression in Müller glia and microglia (Bsibsi et al., 2002; Kumar &

Shamsuddin, 2012; Olson & Miller, 2004; Oosterhof et al., 2017; Rocío Nieto-

Arellano, 2019). Most importantly, we found that in the absence of injury, stimulation

of the immune system activates the regenerative response program of Müller glia, as

shown by upregulation of *mmp9* and *her4.1,* as well as an increase in proliferation;

consistently, expression of *her4.1* also indicates the generation of NPCs from Müller

glia. The expression of the CSF-1 receptor in neural progenitor cells is also

associated with proliferation, survival and differentiation, supporting the theory that

M-CSF stimulates a proliferative/regenerative response in Müller glia (Chitu et al.,

2016). Nevertheless, whether Müller glia reactivity is a direct consequence of

CSF/TLR-agonist treatment, or an indirect consequence of potentially altered

microglia signaling, is so far not clear, because receptors for CSF/TLR signaling

appear to be expressed in both cell populations (Kochan et al., 2012; Kumar &

Shamsuddin, 2012; Letiembre et al., 2007; Lin et al., 2013; Mitchell et al., 2019;

Tsarouchas et al., 2018; Wu et al., 2018). Hence, it will be interesting to test whether

CSF-1R or TLR agonists alone are sufficient to drive increased Müller glia

proliferation, using a model that completely lacks microglial cells in the retina.

Moreover, it will be interesting to determine if the observed effects of the

CSF1R/TLR-agonists are mediated via NF-κB signaling, or via other pathways, such

as the MAP kinase pathway, or combinations thereof (Chen et al., 2018; Kawasaki &

Kawai, 2014; Wan et al., 2012).

Conclusion

Our work strongly supports the notion that acute inflammatory signaling provides a beneficial contribution during adult zebrafish retina regeneration, akin to findings in the telencephalon (Kyritsis et al., 2012; Kizil et al., 2015). Reactivity of retinal leukocytes appears to be tightly controlled after injury, and the presence of leukocytes positively influences reactive proliferation of Müller glial cells, as well as the downstream differentiation of Müller glia progeny into mature photoreceptor cells. Importantly, immune stimulation through TLRs, or via M-CSF injection, was sufficient, in the absence of injury, to trigger Müller glia proliferation and NPC formation. Hence, we conclude that inflammatory signaling plays a critical role during zebrafish retina regeneration.

Further investigations on possible interactions between leukocytes and Müller glia will help to understand the balance between the inflammatory reaction and its resolution during regeneration of neural tissue in zebrafish. In addition, it will be important to identify critical factors secreted by leukocytes in an acute reaction to lesion that are - directly or indirectly- involved in triggering Müller glia reactivity, and thus, regeneration. Finally, this study provides a framework for investigating potentially beneficial effects of an acute immune response, also in comparison with failed regeneration and chronic inflammatory outcome in mammalian models.

Materials and Methods

Ethics statement

The animal experiments were performed in strict accordance with the European Union and German law (Tierschutzgesetz). Experimental procedures were approved by the animal ethics committee of the TU Dresden and the Landesdirektion Sachsen (permits AZ: 24-9168.11-1/2013-5; AZ: TV A 1/2017; TVV 21/2018; AZ: TVV 55/2018).

Zebrafish Maintenance

Zebrafish (*Danio rerio*) were kept under standard housing conditions as previously described (Brand et al., 2002). All fish used in this study were adults, both males and females, and 6 to 12 months of age. Transgenic fish used are: *Tg(−5.5opn1sw1:EGFP)*[kj9] (Takechi et al., 2003); *Tg(mpeg1:mCherry)* (Ellett et al., 2011); *Tg(mpo:GFP)* (Renshaw et al., 2016); *Tg(gfap:NLS-GFP)* (described here); *Tg(gfap:NLS-mCherry)* (Lange et al., 2020), *Tg(lck:NLS-mCherry)*[sd31] (Butko et al., 2015), *Tg(NF-kB:GFP)* (Kanther et al., 2011), Δirf8[st95] (Shiau et al., 2015).

Generation of *Tg(gfap:nls-GFP)*

The *Tg(gfap:NLS-GFP)* reporter line with its NLS-tag shows highly nuclear localization of GFP, but also some leaky cytoplasmic expression in Müller glial cells, and thus serves as an ideal marker for MGCs in our experiments. To generate this line, the GFP reporter was PCR amplified and flanked by restriction sites using GFP-for (atatGGCCGGCCgccaccatggctccaaagaagaagcgtaaggt) and GFP-rev (ggtgtgcatgttttgacgttgatggc) primers. By PCR, the nuclear localization sequence (NLS) was added as a 5' overhang to the GFP. The PCR product was subcloned into

the p2.1 Topo Vector. Next, Topo vector with the reporters and the

pTol(gfap:mcherry-T2A-CreERT2) construct (SH and MB, unpublished) were

digested using the enzymes Asc and FseI and ligated to replace the mCherry-

CreERT2 cassette with the NLS-reporter. For germline transformation, either

linearized plasmid DNA or plasmid DNA with transposase mRNA were injected into

fertilized eggs (F0) in E3 medium (Brand et al., 2002), raised to adulthood and

crossed to AB wild-type fish as previously described (Kawakami et al., 2004).

Diffuse light lesion

To ablate photoreceptor cells, diffuse light lesion was performed as described in

Weber et al., 2013. Briefly, dark-adapted fish were transferred to a beaker containing

250 ml system water and exposed to the light of an EXFO X-Cite 120 W metal halide

lamp (~200,000 lux) for 30'. For recovery, fish were connected to the system and

kept under standard light conditions.

Drug treatment

Dexamethasone (Dex, Sigma-Aldrich, Germany) was diluted from a 25 mg/ml stock

solution in methanol to 15 mg/l in sterilized system water. For Dex treatment, adult

fish were kept individually in sterilized tanks. Control groups with the corresponding

amount of methanol (MeOH) without Dex, both groups did not show retinal cell death

(supplement S3). Solutions were renewed daily, and tanks were exchanged every 5

days. Fish were fed with brine shrimp 1h prior to solution exchange. Animals were

pretreated for 10 days and respective conditions were maintained until sacrificed.

Light lesions were performed in corresponding solutions.

Tissue preparation and sectioning

For retinal flat mounts, eyes were removed and a small incision in the cornea was

made using a sapphire blade scalpel (ZT215 W0.5mm A60°). Eyes were prefixed in

4% PFA in calcium-free ringer solution (PFAR) for 30' with slow agitation. Cornea,

sclera and the lens were removed in calcium-free ringer solution and the retinae were

cut at four sides. Flat mounts were fixed in 4% PFAR at 4 C overnight with gentle

agitation. For storage, samples were transferred to 100% methanol and stored at -

20°C.

For sections, the lens was removed and fish heads were fixed at 4 C in 4% PFA in

0.1 M phosphate buffer (PB). Following that, samples were decalcified and cryo-

protected with 20% sucrose/20% EDTA in 0.1 M PB. Tissues were embedded in

7.5% gelatine/20% sucrose in 0,1% PB, stored at -80 °C and sectioned into 14 µm

cryo-sections using a Microm HM560. Sections were stored at -20°C.

***In situ* Hybridization**

In situ hybridization on sections was performed as described in Ganz et al., 2015.

Hybridization was carried out overnight at 62 °C. Probes were generated using

mmp9 (primer forward: CTTGGAGTCCTGGCGTTTCT; primer reverse:

GCCCGTCCTTGAAGAAGTGA) and *her4.1* (Takke et al., 1999) as targets. For

detection, NBT/BCIP or SIGMA*FAST*™ Fast Red TR/Naphtanol AS-MX was used.

For combination with immunohistochemistry, the primary antibody was incubated with

anti-digoxigenin-AP. Secondary antibody was applied after staining development for

2h and washed with PBS with 0.3% Triton-X100 (PBSTx). Sections were mounted in

glycerol.

Immunohistochemistry & TUNEL

Sections were dried for 2 h at 50 °C and rehydrated with PBSTx. Following this,

sections were incubated with primary antibody: anti-Zrf-1 ZIRC mouse IgG, 1:200

(Fausett & Goldman, 2006), anti-Zpr-3 ZIRC IgG, 1:200 (Zou et al., 2008), anti-GFP

Abcam chicken 1:3000 (Leinninger et al., 2009), anti-DsRed Clontech rabbit 1:500

(Glass et al., 2005), anti-PCNA mouse IgG$_{2a}$ DACO PC10 1:500 (Grandel et al.,

2006), anti-L-Plastin rabbit 1:5000, (Redd et al., 2006) and incubated overnight at

4 °C. Excess antibody was washed off using PBSTx followed by incubation with

secondary antibody against the respective primary antibody host (Molecular Probes;

Alexa 488, Alexa 555, Alexa 635; 1:750) containing 1 µg/ml 4′,6-Diamidin-2-

phenylindol (DAPI). Antibody-solution was washed off and slides were mounted with

glycerol.

To retrieve the PCNA antigen, sections were incubated for 8' in 50 mM Tris buffer pH

8.0 at 99 °C followed by 10 min PBS prior to primary antibody incubation.

Modifications for retinal flat mounts are prolonged incubation times for primary and

secondary antibody solution for 48 h respectively at 4 °C with slow agitation and

extensive washing 10x 30' after each antibody incubation. Flat mounts were mounted

on slides in glycerol.

TUNEL (TdT-mediated dUTP-biotin nick end labeling) assays were performed on

sections using ApopTag® Red in Situ Apoptosis Detection Kit (Merk) according to

manufacturer's instructions.

EdU labeling & detection

To trace proliferating cells, EdU (5-ethynyl-2′-deoxyuridine) pulses were given

intraperitoneally. Fish were anesthetized in 0.024% Tricaine and EdU was injected

intraperitoneally (20 µl of 2.5 mg/ml EdU in PBS per pulse). Detection was performed

on sections using "Click-iT® Plus EdU Alexa Fluor® 555 Imaging Kit" (Thermo

Fischer Scientific) according to manufacturer's instructions.

Intravitreal injections

Fish were anesthetized in 0.024 % tricaine and covered with tricaine-moistened

paper towel, leaving the eye exposed. The outer cornea was removed.

Subsequently, a small incision in the cornea was made using a sapphire blade

scalpel (ZT215 W0.5mm A60°). A Hamilton syringe equipped with a 33 gauge, blunt-

end needle was inserted and 0,5 µl of solution (flagellin from *S. typhimurium*, 0,1

µg/µl, Sigma; zymosan from *S. cerevisiae*, 0,2 µg/µl BioParticles; Human M-CSF, 50

ng/µl, PreproTech; Human Il-34, 50 ng/µl, PreproTech; PBS) was injected into the

vitreous of the eye without touching the retina. The incision was covered with

Histoacryl® (Braun) to avoid leakage.

Image acquisition

Images were acquired using a Zeiss Axio Imager, equipped with ApoTome. EC Plan-

Neofluar 5x/0.16, Plan-Apochromat 20x/0.8 and LD C-Apochromat 40x/1.1 W Korr

UV VIS IR objectives were used for magnification. For detection either a

monochromatic Axiocam HR (1388x1040 pixels, 6.45*6.45) for fluorescence or a

polychromatic Axiocam MR Rev3 (1388x1040 pixels, 6.45*6.45) for bright-field

imaging was used. Sequential image acquisition was used in co-stained samples with

multiple fluorophores. Images were acquired in AxioVision Rel. 4.8 and processed

using Fiji (Schindelin et al., 2012). Figure panels were assembled in Adobe

Photoshop CS6.

Cell counting and statistical analysis

For cell quantification, retinal sections with a thickness of 14 µm were used. For

single fluorophore cell quantification, the first 3-5 consecutive retinal sections rostral

to the optic nerve were selected and manually counted. For normalization, the length

of the retina was determined for every section by measuring a snapshot of the retina

at the level of the inner nuclear layer. For quantification of multiple fluorophores, 20x

images were acquired. Cells were quantified using the cell counter tool in Fiji. Length

measurements were applied on the same images for normalization. For statistical

analyses, ≥3 fish were used for calculations. Counted cell numbers were normalized

to the respective parameter (reference cell population or length) and the counts per

fish were averaged. The average number for every fish was used as an analytical

unit. For statistical analysis, Graph Pad Prism was used to determine p-values with a

one-way ANOVA test with Tukey's post hoc analysis. For unpaired two-tailed

student's T-Test, Excel was used. Significance levels are displayed as: not significant

= p>0.05, * = p≤0.5, ** = p≤0.01, *** = p≤0.001. Graphs shown in bar charts were

created using Excel, error bars represent standard error of the mean.

Acknowledgments

We are very grateful to Marika Fischer, Jitka Michling, Daniela Mögel, and Claudia

Meyer who provided outstanding fish care, and Dr. Judith Konantz for managing the

fish facility. We also thank Michaela Geffarth and Nathalie Franke for excellent

technical support and John F Rawls for sharing the *Tg(NF-kB:GFP)* reporter line.

Further, we thank past and present members of the Brand lab for many interesting

discussions, and Dr. Thomas Quail, Dr. Judith Konantz, Dr. Christian Lange, Prof. Dr.

Thomas Becker, Prof. Dr. Catherina Becker, Prof. Dr. Marius Ader, Ecem Çayıroğlu

and Prof. Dr. Susanne Koch for comments on earlier versions of this manuscript.

Funding

742 This work was supported by project grants of the German Research Foundation

743 (DFG, project numbers BR 1746/3, BR 1746/11-1) and an ERC advanced grant (Zf-

744 BrainReg) to MB. We also gratefully acknowledge support of OB by a pre-doctoral

745 fellowship of the ProRetina foundation.

Competing interests

748 The authors declare that they have no conflict of interest.

749
750

References

Akhtar-Schäfer, I., Wang, L., Krohne, T. U., Xu, H., & Langmann, T. (2018). Modulation of three key innate immune pathways for the most common retinal degenerative diseases. *EMBO Molecular Medicine*, *10*(10), e8259. https://doi.org/10.15252/emmm.201708259

Akira, S., & Takeda, K. (2004). Toll-like receptor signalling. *Nature Reviews Immunology*, *4*(7), 499–511. https://doi.org/10.1038/nri1391

Amor, S., Puentes, F., Baker, D., & Van Der Valk, P. (2010). Inflammation in neurodegenerative diseases. *Immunology*, *129*(2), 154–169. https://doi.org/10.1111/j.1365-2567.2009.03225.x

Aslanidis, A., Karlstetter, M., Scholz, R., Fauser, S., Neumann, H., Fried, C., Pietsch, M., & Langmann, T. (2015). Activated microglia/macrophage whey acidic protein (AMWAP) inhibits NFKB signaling and induces a neuroprotective phenotype in microglia. *Journal of Neuroinflammation*, *12*(1), 1–14. https://doi.org/10.1186/s12974-015-0296-6

Aurora, A. B., & Olson, E. N. (2014). Immune modulation of stem cells and regeneration. *Cell Stem Cell*, *15*(1), 14–25. https://doi.org/10.1016/j.stem.2014.06.009

Barber, A. E., Fleming, B. A., & Mulvey, M. A. (2016). Similarly Lethal Strains of Extraintestinal Pathogenic Escherichia coli Trigger Markedly Diverse Host Responses in a Zebrafish Model of Sepsis. *MSphere*, *1*(2), 1–19. https://doi.org/10.1128/msphere.00062-16

Bosak, V., Murata, K., Bludau, O., & Brand, M. (2018). *Role of the immune response in initiating central nervous system regeneration in vertebrates : learning from the fish. 417*, 403–417.

Brand, M., Granato, M. and Nüsslein-Volhard, C. (2002). Keeping and raising zebrafish. In Zebrafish: A Practical Approach, (ed. C. Nüsslein-Volhard and R. Dahm). Oxford: Oxford University Press.

Branzk, N., & Papayannopoulos, V. (2013). Molecular mechanisms regulating NETosis in infection and disease. *Seminars in Immunopathology*, *35*(4), 513–530. https://doi.org/10.1007/s00281-013-0384-6

Bringmann, A, Pannike, T., Grosche, J., Francke, M., Wiedemann, P., Skatchkov, S., Osborne, N., & Reichenbach, A. (2006). Müller cells in the healthy and diseased retina. *Progress in Retinal and Eye Research*, *25*(4), 397–424. https://doi.org/10.1016/j.preteyeres.2006.05.003

Bringmann, Andreas, Pannicke, T., Biedermann, B., Francke, M., Iandiev, I., Grosche, J., Wiedemann, P., Albrecht, J., & Reichenbach, A. (2009). Role of retinal glial cells in neurotransmitter uptake and metabolism. *Neurochemistry International*, *54*(3–4), 143–160. https://doi.org/10.1016/J.NEUINT.2008.10.014

Bringmann, Andreas, & Wiedemann, P. (2011). Müller glial cells in retinal disease. *Ophthalmologica*, *227*(1), 1–19. https://doi.org/10.1159/000328979

Bsibsi, M., Ravid, R., Gveric, D., & Van Noort, J. M. (2002). Broad expression of Toll-like receptors in the human central nervous system. *Journal of Neuropathology and Experimental Neurology*, *61*(11), 1013–1021. https://doi.org/10.1093/jnen/61.11.1013

Buffo, A., Rite, I., Tripathi, P., Lepier, A., Colak, D., Horn, A.-P., Mori, T., & Götz, M. (2008). Origin and progeny of reactive gliosis: A source of multipotent cells in the injured brain. *Proceedings of the National Academy of Sciences*, *105*(9), 3581–3586. https://doi.org/10.1073/pnas.0709002105

Butko, E., Distel, M., Pouget, C., Weijts, B., Kobayashi, I., Ng, K., Mosimann, C., Poulain, F. E., McPherson, A., Ni, C.-W., Stachura, D. L., Del Cid, N., Espin-Palazon, R., Lawson, N. D., Dorsky, R., Clements, W. K., & Traver, D. (2015). Gata2b is a restricted early regulator of hemogenic endothelium in the zebrafish embryo. *Development*, *142*(6), 1050–1061. https://doi.org/10.1242/dev.119180

Campbell, E. L., Bruyninckx, W. J., Kelly, C. J., Glover, L. E., McNamee, E. N., Bowers, B. E., Bayless, A. J., Scully, M., Saeedi, B. J., Golden-Mason, L., Ehrentraut, S. F., Curtis, V. F., Burgess, A., Garvey, J. F., Sorensen, A., Nemenoff, R., Jedlicka, P., Taylor, C. T., Kominsky, D. J., & Colgan, S. P. (2014). Transmigrating neutrophils shape the mucosal microenvironment through localized oxygen depletion to influence resolution of inflammation. *Immunity*, *40*(1), 66–77. https://doi.org/10.1016/j.immuni.2013.11.020

Chen, L., Deng, H., Cui, H., Fang, J., Zuo, Z., Deng, J., Li, Y., Wang, X., & Zhao, L. (2018). Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget*, *9*(6), 7204–7218. https://doi.org/10.18632/oncotarget.23208

Cheng, C. Y., Hsieh, H. L., Hsiao, L. Der, & Yang, C. M. (2012). PI3-K/Akt/JNK/NF-κB is essential for MMP-9 expression and outgrowth in human limbal epithelial cells on intact amniotic membrane. *Stem Cell Research*, *9*(1), 9–23. https://doi.org/10.1016/j.scr.2012.02.005

Chitu, V., Gokhan, Ş., Nandi, S., Mehler, M. F., & Stanley, E. R. (2016). Emerging Roles for CSF-1 Receptor and its Ligands in the Nervous System. *Trends in Neurosciences*, *39*(6), 378–393. https://doi.org/10.1016/J.TINS.2016.03.005

Celotto, L., Rost, F., Machate, A., Bläsche, J., Dahl, A., Weber, A., Hans, S., Brand, M. (2023). Single cell RNA sequencing unravels the transcriptional network underlying zebrafish retina regeneration *eLife* 12:RP86507. https://doi.org/10.7554/eLife.86507.1

Conedera, F. M., Pousa, A. M. Q., Mercader, N., Tschopp, M., & Enzmann, V. (2019). Retinal microglia signaling affects Müller cell behavior in the zebrafish following laser injury induction. *Glia*, *6*(November 2018), 1–17. https://doi.org/10.1002/glia.23601

Conner, C., Ackerman, K. M., Lahne, M., Hobgood, J. S., & Hyde, D. R. (2014). Repressing Notch Signaling and Expressing TNFα Are Sufficient to Mimic Retinal Regeneration by Inducing Muller Glial Proliferation to Generate Committed Progenitor Cells. *Journal of Neuroscience*, *34*(43), 14403–14419. https://doi.org/10.1523/JNEUROSCI.0498-14.2014

Coutinho, A. E., & Chapman, K. E. (2011). The anti-inflammatory and immunosuppressive effects of glucocorticoids, recent developments and mechanistic insights. *Molecular and Cellular Endocrinology*, *335*(1), 2–13. https://doi.org/10.1016/j.mce.2010.04.005

De Preux Charles, A. S., Bise, T., Baier, F., Marro, J., & Jaźwińska, A. (2016). Distinct effects of inflammation on preconditioning and regeneration of the adult zebrafish heart. *Open Biology*, *6*(7). https://doi.org/10.1098/rsob.160102

Ellett, F., Pase, L., Hayman, J. W., Andrianopoulos, A., & Lieschke, G. J. (2011). Phagocytes, Granulocytes, and Myelopoiesis mpeg1 promoter transgenes direct macrophage-lineage expression in zebrafish. *Blood*, *27*(4), e49–e56. https://doi.org/10.1182/blood-2010-10-314120: 10.1182/blood-2010-10-314120

Eshaq, R. S., Aldalati, A. M. Z., Alexander, J. S., & Harris, N. R. (2017). Diabetic retinopathy: Breaking the barrier. *Pathophysiology*, *24*(4), 229–241. https://doi.org/10.1016/J.PATHOPHYS.2017.07.001

Farrera, C., & Fadeel, B. (2013). Macrophage Clearance of Neutrophil Extracellular Traps Is
a Silent Process. *The Journal of Immunology*, *191*(5), 2647–2656.
https://doi.org/10.4049/jimmunol.1300436

Fausett, B. V., & Goldman, D. (2006). A Role for ß1 Tubulin-Expressing Muller Glia in
Regeneration of the Injured Zebrafish Retina. *Journal of Neuroscience*, *26*(23), 6303–
6313. https://doi.org/10.1523/jneurosci.0332-06.2006

Ferrero-Miliani, L., Nielsen, O. H., Andersen, P. S., & Girardin, S. E. (2007). Chronic
inflammation: Importance of NOD2 and NALP3 in interleukin-1β generation. *Clinical and*
Experimental Immunology, *147*(2), 227–235. https://doi.org/10.1111/j.1365-
2249.2006.03261.x

Ferrero, G., Mahony, C. B., Dupuis, E., Yvernogeau, L., Di Ruggiero, E., Miserocchi, M.,
Caron, M., Robin, C., Traver, D., Bertrand, J. Y., & Wittamer, V. (2018). Embryonic
Microglia Derive from Primitive Macrophages and Are Replaced by cmyb-Dependent
Definitive Microglia in Zebrafish. *Cell Reports*, *24*(1), 130–141.
https://doi.org/10.1016/j.celrep.2018.05.066

Fitch, M. T., & Silver, J. (2008). CNS injury, glial scars, and inflammation: Inhibitory
extracellular matrices and regeneration failure. *Experimental Neurology*, *209*(2), 294–301.
https://doi.org/10.1016/j.expneurol.2007.05.014

Fritsche, L. G., Igl, W., Bailey, J. N. C., Grassmann, F., Sengupta, S., Bragg-Gresham, J. L.,
Burdon, K. P., Hebbring, S. J., Wen, C., Gorski, M., Kim, I. K., Cho, D., Zack, D., Souied,
E., Scholl, H. P. N., Bala, E., ELee, K., Hunter, D. J., Sardell, R. J., … Heid, I. M. (2016).
A large genome-wide association study of age-related macular degeneration highlights
contributions of rare and common variants. *Nature Genetics*, *48*(2), 134–143.
https://doi.org/10.1038/ng.3448

Gallina, D., Zelinka, C., & Fischer, A. J. (2014). Glucocorticoid receptors in the retina, Muller
glia and the formation of Muller glia-derived progenitors. *Development*, *141*(17), 3340–
3351. https://doi.org/10.1242/dev.109835

Gallina, D., Zelinka, C. P., Cebulla, C. M., & Fischer, A. J. (2015). Activation of
glucocorticoid receptors in Müller glia is protective to retinal neurons and suppresses
microglial reactivity. *Experimental Neurology*, *273*, 114–125.
https://doi.org/10.1016/j.expneurol.2015.08.007

Ganz, J., Kroehne, V., Freudenreich, D., Machate, A., Geffarth, M., Braasch, I., Kaslin, J., &
Brand, M. (2015). Subdivisions of the adult zebrafish pallium based on molecular marker
analysis. *F1000Research*, *3*(3), 308. https://doi.org/10.12688/f1000research.5595.2

Gasparini, S. J., Llonch, S., Borsch, O., & Ader, M. (2019). Transplantation of photoreceptors
into the degenerative retina: Current state and future perspectives. *Progress in Retinal and*
Eye Research, *69*(July 2018), 1–37. https://doi.org/10.1016/j.preteyeres.2018.11.001

Glass, R., Synowitz, M., Kronenberg, G., Walzlein, J., Markovic, D., Wang, L., Gast, D.,
Kiwit, J., Kempermann, G., & Kettenmann, H. (2005). Glioblastoma-Induced Attraction of
Endogenous Neural Precursor Cells Is Associated with Improved Survival. *Journal of*
Neuroscience, *25*(10), 2637–2646. https://doi.org/10.1523/jneurosci.5118-04.2005

Goldman, D. (2014). Müller glial cell reprogramming and retina regeneration. *Nature*
Reviews. Neuroscience, *15*(7), 431–442. https://doi.org/10.1038/nrn3723

Gong, Y., Hart, E., Shchurin, A., & Hoover-Plow, J. (2008). Inflammatory macrophage
migration requires MMP-9 activation by plasminogen in mice. *Journal of Clinical*
Investigation, *118*(9), 3012–3024. https://doi.org/10.1172/JCI32750

Gorsuch, R. A., & Hyde, D. R. (2014). Regulation of Müller glial dependent neuronal
regeneration in the damaged adult zebrafish retina. *Experimental Eye Research*, *123*, 131–
140. https://doi.org/10.1016/j.exer.2013.07.012

Grandel, H., Kaslin, J., Ganz, J., Wenzel, I., & Brand, M. (2006). Neural stem cells and
neurogenesis in the adult zebrafish brain: Origin, proliferation dynamics, migration and
cell fate. *Developmental Biology*, *295*(1), 263–277.
https://doi.org/10.1016/j.ydbio.2006.03.040

Hammer, J., Röppenack, P., Yousuf, S., Schnabel, C., Weber, A., Zöller, D., Koch, E., Hans,
S., & Brand, M. (2022). Visual function is gradually restored during retina regeneration in
adult zebrafish. *Frontiers in Cell and Developmental Biology*, *9*(February), 1–18.
https://doi.org/10.3389/fcell.2021.831322

Hanovice, N. J., Leach, L. L., Slater, K., Gabriel, A. E., Romanovicz, D., Shao, E., Collery,
R., Burton, E. A., Lathrop, K. L., Link, B. A., & Gross, J. M. (2019). Regeneration of the
zebrafish retinal pigment epithelium after widespread genetic ablation. In *PLoS Genetics*
(Vol. 15, Issue 1). https://doi.org/10.1371/journal.pgen.1007939

Herrero, M.-T., Estrada, C., Maatouk, L., & Vyas, S. (2015). Inflammation in Parkinsons
disease: role of glucocorticoids. *Frontiers in Neuroanatomy*, *9*(April), 1–12.
https://doi.org/10.3389/fnana.2015.00032

Hochmann, S., Kaslin, J., Hans, S., Weber, A., Machate, A., Geffarth, M., Funk, R. H., Brand,
M. (2012). Fgf signaling is required for photoreceptor maintenance in the adult zebrafish
retina. *PLoS One*.;7(1), https://doi.org/10.1371/journal.pone.0030365.

Hui, S. P., Sheng, D. Z., Sugimoto, K., Gonzalez-Rajal, A., Nakagawa, S., Hesselson, D., &
Kikuchi, K. (2017). Zebrafish Regulatory T Cells Mediate Organ-Specific Regenerative
Programs. *Developmental Cell*, *43*(6), 659-672.e5.
https://doi.org/10.1016/j.devcel.2017.11.010

Iribarne, M., & Hyde, D. R. (2022). Different inflammation responses modulate Müller glia
proliferation in the acute or chronically damaged zebrafish retina. *Frontiers in Cell and*
Developmental Biology, *10*(August), 1–19. https://doi.org/10.3389/fcell.2022.892271

Issaka Salia, O., & Mitchell, D. M. (2020). Bioinformatic analysis and functional predictions
of selected regeneration-associated transcripts expressed by zebrafish microglia. *BMC*
Genomics, *21*(1), 1–17. https://doi.org/10.1186/s12864-020-07273-8

Kanther, M., Sun, X., Mühlbauer, M., MacKey, L. C., Flynn, E. J., Bagnat, M., Jobin, C., &
Rawls, J. F. (2011). Microbial colonization induces dynamic temporal and spatial patterns
of NF-κB activation in the zebrafish digestive tract. *Gastroenterology*, *141*(1), 197–207.
https://doi.org/10.1053/j.gastro.2011.03.042

Karlstetter, M., Ebert, S., & Langmann, T. (2010). Microglia in the healthy and degenerating
retina: Insights from novel mouse models. *Immunobiology*, *215*(9–10), 685–691.
https://doi.org/10.1016/j.imbio.2010.05.010

Kaur, S., Gupta, S., Chaudhary, M., Khursheed, M. A., Mitra, S., Kurup, A. J., &
Ramachandran, R. (2018). let-7 MicroRNA-Mediated Regulation of Shh Signaling and the
Gene Regulatory Network Is Essential for Retina Regeneration. *Cell Reports*, *23*(5), 1409–
1423. https://doi.org/10.1016/j.celrep.2018.04.002

Kawakami, K., Kobayashi, M., Matsuda, N., Takeda, H., Mishina, M., & Kawakami, N.
(2004). A Transposon-Mediated Gene Trap Approach Identifies Developmentally
Regulated Genes in Zebrafish. *Developmental Cell*, *7*(1), 133–144.
https://doi.org/10.1016/j.devcel.2004.06.005

Kawasaki, T., & Kawai, T. (2014). Toll-like receptor signaling pathways. *Frontiers in*
Immunology, *300*(5625), 1524–1525. https://doi.org/10.1126/science.1085536

Keightley, M. C., Wang, C. H., Pazhakh, V., & Lieschke, G. J. (2014). Delineating the roles
of neutrophils and macrophages in zebrafish regeneration models. *International Journal of*
Biochemistry and Cell Biology, *56*, 92–106. https://doi.org/10.1016/j.biocel.2014.07.010

Kim, S. D., Yang, S.-I., Kim, H.-C., Shin, C. Y., & Ko, K. H. (2007). Inhibition of GSK-3β
mediates expression of MMP-9 through ERK1/2 activation and translocation of NF-κB in
rat primary astrocytes. *Brain Research*, *1186*, 12–20.
https://doi.org/10.1016/J.BRAINRES.2007.10.018

Kinney, J. W., Bemiller, S. M., Murtishaw, A. S., Leisgang, A. M., Salazar, A. M., & Lamb,
B. T. (2018). Inflammation as a central mechanism in Alzheimer's disease. *Alzheimer's*
and Dementia: Translational Research and Clinical Interventions, *4*, 575–590.
https://doi.org/10.1016/j.trci.2018.06.014

Kizil, C., Kyritsis, N., & Brand, M. (2015). Effects of inflammation on stem cells: together
they strive?. *EMBO reports*, *16*(4), 416–426. https://doi.org/10.15252/embr.201439702

Kochan, T., Singla, A., Tosi, J., & Kumar, A. (2012). Toll-Like Receptor 2 Ligand
Pretreatment Attenuates Retinal Microglial Inflammatory Response but Enhances
Phagocytic Activity toward Staphylococcus aureus. *Infection and Immunity*, *80*(6), 2076–
2088. https://doi.org/10.1128/iai.00149-12

Kramer, A. C., Gurdziel, K., & Thummel, R. (2021). A Comparative Analysis of Gene and
Protein Expression Throughout a Full 28-Day Retinal Regeneration Time-Course in Adult
Zebrafish. *Frontiers in Cell and Developmental Biology*, *9*.
https://doi.org/10.3389/fcell.2021.741514

Kumar, A., & Shamsuddin, N. (2012). Retinal muller glia initiate innate response to infectious
stimuli via toll-like receptor signaling. *PLoS ONE*, *7*(1), e29830.
https://doi.org/10.1371/journal.pone.0029830

Kurimoto, T., Yin, Y., Habboub, G., Gilbert, H.-Y., Li, Y., Nakao, S., Hafezi-Moghadam, A.,
& Benowitz, L. I. (2013). Neutrophils Express Oncomodulin and Promote Optic Nerve
Regeneration. *Journal of Neuroscience*, *33*(37), 14816–14824.
https://doi.org/10.1523/JNEUROSCI.5511-12.2013

Kyritsis, N., Kizil, C., Zocher, S., Kroehne, V., Kaslin, J., Freudenreich, D., Iltzsche, A., &
Brand, M. (2012). Acute inflammation initiates the regenerative response in the adult
zebrafish brain. *Science*, *338*(6112), 1353–1356. https://doi.org/10.1126/science.1228773

Lahne, M., Nagashima, M., Hyde, D. R., & Hitchcock, P. F. (2020). Reprogramming Müller
Glia to Regenerate Retinal Neurons. *Annual Review of Vision Science*, *6*, 171–193.
https://doi.org/10.1146/annurev-vision-121219-081808

Lange, C., Rost, F., MacHate, A., Reinhardt, S., Lesche, M., Weber, A., Kuscha, V., Dahl, A.,
Rulands, S., & Brand, M. (2020). Single cell sequencing of radial glia progeny reveals the
diversity of newborn neurons in the adult zebrafish brain. *Development (Cambridge)*,
147(1). https://doi.org/10.1242/dev.185595

LeBert, D. C., Squirrell, J. M., Rindy, J., Broadbridge, E., Lui, Y., Zakrzewska, A., Eliceiri,
K. W., Meijer, A. H., & Huttenlocher, A. (2015). Matrix metalloproteinase 9 modulates
collagen matrices and wound repair. *Development*, *142*(12), 2136–2146.
https://doi.org/10.1242/dev.121160

Leininger, G. M., Jo, Y. H., Leshan, R. L., Louis, G. W., Yang, H., Barrera, J. G., Wilson,
H., Opland, D. M., Faouzi, M. A., Gong, Y., Jones, J. C., Rhodes, C. J., Chua, S., Diano,
S., Horvath, T. L., Seeley, R. J., Becker, J. B., Münzberg, H., & Myers, M. G. (2009).
Leptin Acts via Leptin Receptor-Expressing Lateral Hypothalamic Neurons to Modulate
the Mesolimbic Dopamine System and Suppress Feeding. *Cell Metabolism*, *10*(2), 89–98.
https://doi.org/10.1016/j.cmet.2009.06.011

Lenkowski, J. R., Qin, Z., Sifuentes, C. J., Thummel, R., Soto, C. M., Moens, C. B., &
Raymond, P. A. (2013). Retinal regeneration in adult zebrafish requires regulation of
TGFβ signaling. *Glia*, *61*(10), 1687–1697. https://doi.org/10.1002/glia.22549

Lenkowski, J. R., & Raymond, P. A. (2014). Müller glia: Stem cells for generation and
regeneration of retinal neurons in teleost fish. *Progress in Retinal and Eye Research*, *40*,
94–123. https://doi.org/10.1016/j.preteyeres.2013.12.007

Letiembre, M., Hao, W., Liu, Y., Walter, S., Mihaljevic, I., Rivest, S., Hartmann, T., &
Fassbender, K. (2007). Innate immune receptor expression in normal brain aging.
Neuroscience, *146*(1), 248–254. https://doi.org/10.1016/J.NEUROSCIENCE.2007.01.004

Lin, X., Fang, D., Zhou, H., & Su, S. B. (2013). The expression of Toll-like receptors in
murine Müller cells, the glial cells in retina. *Neurological Sciences*, *34*(8), 1339–1346.
https://doi.org/10.1007/s10072-012-1236-1

Liu, T., Zhang, L., Joo, D., & Sun, S. C. (2017). NF-κB signaling in inflammation. *Signal*
Transduction and Targeted Therapy, *2*(March). https://doi.org/10.1038/sigtrans.2017.23

Madeira, M. H., Rashid, K., Ambrósio, A. F., Santiago, A. R., & Langmann, T. (2018).
Blockade of microglial adenosine A2A receptor impacts inflammatory mechanisms,
reduces ARPE-19 cell dysfunction and prevents photoreceptor loss in vitro. *Scientific*
Reports, *8*(1), 1–15. https://doi.org/10.1038/s41598-018-20733-2

McMenamin, P. G., Saban, D. R., & Dando, S. J. (2019). Immune cells in the retina and
choroid: Two different tissue environments that require different defenses and surveillance.
In *Progress in Retinal and Eye Research* (Vol. 70, pp. 85–98). Pergamon.
https://doi.org/10.1016/j.preteyeres.2018.12.002

Mitchell, D. M., Lovel, A. G., & Stenkamp, D. L. (2018). Dynamic changes in microglial and
macrophage characteristics during degeneration and regeneration of the zebrafish retina.
Journal of Neuroinflammation, *15*(1), 1–20. https://doi.org/10.1186/s12974-018-1185-6

Mitchell, D. M., Sun, C., Hunter, S. S., New, D. D., & Stenkamp, D. L. (2019). Regeneration
associated transcriptional signature of retinal microglia and macrophages. *Scientific*
Reports, *9*(1), 4768. https://doi.org/10.1038/s41598-019-41298-8

Nathan, C., & Ding, A. (2010). Nonresolving Inflammation. *Cell*, *140*(6), 871–882.
https://doi.org/10.1016/j.cell.2010.02.029

Nelson, C. M., Ackerman, K. M., O'Hayer, P., Bailey, T. J., Gorsuch, R. A., & Hyde, D. R.
(2013). Tumor Necrosis Factor-Alpha Is Produced by Dying Retinal Neurons and Is
Required for Muller Glia Proliferation during Zebrafish Retinal Regeneration. *Journal of*
Neuroscience, *33*(15), 6524–6539. https://doi.org/10.1523/JNEUROSCI.3838-12.2013

Olson, J. K., & Miller, S. D. (2004). Microglia Initiate Central Nervous System Innate and
Adaptive Immune Responses through Multiple TLRs. *The Journal of Immunology*, *173*(6),
3916–3924. https://doi.org/10.4049/jimmunol.173.6.3916

Oosterhof, N., Holtman, I. R., Kuil, L. E., van der Linde, H. C., Boddeke, E. W. G. M.,
Eggen, B. J. L., & van Ham, T. J. (2017). Identification of a conserved and acute

neurodegeneration-specific microglial transcriptome in the zebrafish. *Glia*, *65*(1), 138–149. https://doi.org/10.1002/glia.23083

Palazzo, I., Deistler, K., Hoang, T. V., Blackshaw, S., & Fischer, A. J. (2020). NF-κB signaling regulates the formation of proliferating Müller glia-derived progenitor cells in the avian retina. *Development*, *147*(10), dev183418. https://doi.org/10.1242/dev.183418

Petrie, T. A., Strand, N. S., Yang, C.-T., Rabinowitz, J. S., & Moon, R. T. (2015). Macrophages modulate adult zebrafish tail fin regeneration. *Development*, *142*(2), 406–406. https://doi.org/10.1242/dev.120642

Polager, S., & Ginsberg, D. (2002). NF-kappaB at the crossroads of life and death. *Trends in Cell Biology*, *3*(3), 528–535. http://www.ncbi.nlm.nih.gov/pubmed/18805009

Portillo, J. A. C., Corcino, Y. L., Miao, Y., Tang, J., Sheibani, N., Kern, T. S., Dubyak, G. R., & Subauste, C. S. (2017). CD40 in retinal müller cells induces P2X7-dependent cytokine expression in macrophages/microglia in diabetic mice and development of early experimental diabetic retinopathy. *Diabetes*, *66*(2), 483–493. https://doi.org/10.2337/db16-0051

Raymond, P. A., Barthel, L. K., Bernardos, R. L., & Perkowski, J. J. (2006). Molecular characterization of retinal stem cells and their niches in adult zebrafish. *BMC Developmental Biology*, *6*(36), 1–17. https://doi.org/10.1186/1471-213X-6-36

Redd, M. J., Kelly, G., Dunn, G., Way, M., & Martin, P. (2006). Imaging macrophage chemotaxis in vivo: Studies of microtubule function in zebrafish wound inflammation. *Cell Motility and the Cytoskeleton*, *63*(7), 415–422. https://doi.org/10.1002/cm.20133

Renshaw, S. A., Loynes, C. A., Trushell, D. M. I., Elworthy, S., Ingham, P. W., & Whyte, M. K. B. (2016). Plenary paper A transgenic zebrafish model of neutrophilic inflammation. *Blood Journal*, *108*(13), 3976–3979. https://doi.org/10.1182/blood-2006-05-024075.The

Roche, S. L., Ruiz-Lopez, A. M., Moloney, J. N., Byrne, A. M., & Cotter, T. G. (2018). Microglial-induced Müller cell gliosis is attenuated by progesterone in a mouse model of retinitis pigmentosa. *Glia*, *66*(2), 295–310. https://doi.org/10.1002/glia.23243

Rocío Nieto-Arellano, H. S.-I. (2019). zfRegeneration: a database for gene expression profiling during regeneration. *Bioinformatics*, *35*(4), 703–705. https://doi.org/10.1093/bioinformatics/bty659

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Schmidt, M., Pauels, H. G., Lügering, N., Lügering, A., Domschke, W., & Kucharzik, T. (1999). Glucocorticoids induce apoptosis in human monocytes: potential role of IL-1 beta. *Journal of Immunology (Baltimore, Md. : 1950)*, *163*(6), 3484–3490. http://www.ncbi.nlm.nih.gov/pubmed/10477621

Shiau, C. E., Kaufman, Z., Meireles, A. M., & Talbot, W. S. (2015). Differential Requirement for irf8 in Formation of Embryonic and Adult Macrophages in Zebrafish. *PLOS ONE*, *10*(1), e0117513. https://doi.org/10.1371/journal.pone.0117513

Sifuentes, C. J., Kim, J. W., Swaroop, A., & Raymond, P. A. (2016). Rapid, dynamic activation of Müller Glial stem cell responses in Zebrafish. *Investigative Ophthalmology and Visual Science*, *57*(13), 5148–5160. https://doi.org/10.1167/iovs.16-19973

Silva, N. J., Nagashima, M., Li, J., Kakuk-Atkins, L., Ashrafzadeh, M., Hyde, D. R., & Hitchcock, P. F. (2020). Inflammation and matrix metalloproteinase 9 (Mmp-9) regulate photoreceptor regeneration in adult zebrafish. *Glia*, *68*(7), 1445–1465. https://doi.org/10.1002/glia.23792

Sim, D. A., Chu, C. J., Selvam, S., Powner, M. B., Liyanage, S., Copland, D. A., Keane, P. A., Tufail, A., Egan, C. A., Bainbridge, J. W. B., Lee, R. W., Dick, A. D., & Fruttiger, M. (2015). A simple method for in vivo labelling of infiltrating leukocytes in the mouse retina using indocyanine green dye. *Disease Models & Mechanisms*, *8*(11), 1479–1487. https://doi.org/10.1242/dmm.019018

Sofroniew, M. V. (2009). Molecular dissection of reactive astrogliosis and glial scar formation. *Trends in Neurosciences*, *32*(12), 638–647. https://doi.org/10.1016/j.tins.2009.08.002

Stirling, D. P., Liu, S., Kubes, P., & Yong, V. W. (2009). Depletion of Ly6G/Gr-1 leukocytes after spinal cord injury in mice alters wound healing and worsens neurological outcome. *Journal of Neuroscience*, *29*(3), 753–764. https://doi.org/10.1523/JNEUROSCI.4918-08.2009

Takada, Y., Khuri, F. R., & Aggarwal, B. B. (2004). Protein farnesyltransferase inhibitor (SCH 66336) abolishes NF-κB activation induced by various carcinogens and inflammatory stimuli leading to suppression of NF-κB-regulated gene expression and up-regulation of apoptosis. *Journal of Biological Chemistry*, *279*(25), 26287–26299. https://doi.org/10.1074/jbc.M400963200

Takechi, M., Hamaoka, T., & Kawamura, S. (2003). Fluorescence visualization of ultraviolet-sensitive cone photoreceptor development in living zebrafish. *FEBS Letters*, *553*(1–2), 90–94. https://doi.org/10.1016/S0014-5793(03)00977-3

Takke, C., Dornseifer, P., v Weizsäcker, E., & Campos-Ortega, J. A. (1999). her4, a zebrafish homologue of the Drosophila neurogenic gene E(spl), is a target of NOTCH signalling. *Development (Cambridge, England)*, *126*(9), 1811–1821. http://www.ncbi.nlm.nih.gov/pubmed/10101116

Thomas, J. L., Ranski, A. H., Morgan, G. W., & Thummel, R. (2016). Reactive gliosis in the adult zebrafish retina. *Experimental Eye Research*, *143*, 98–109. https://doi.org/10.1016/j.exer.2015.09.017

Todd, L., Finkbeiner, C., Wong, C. K., Hooper, M. J., & Reh, T. A. (2020). Microglia Suppress Ascl1-Induced Retinal Regeneration in Mice. *Cell Reports*, *33*(11), 108507. https://doi.org/10.1016/j.celrep.2020.108507

Tsarouchas, T. M., Wehner, D., Cavone, L., Munir, T., Keatinge, M., Lambertus, M., Underhill, A., Barrett, T., Kassapis, E., Ogryzko, N., Feng, Y., van Ham, T. J., Becker, T., & Becker, C. G. (2018). Dynamic control of proinflammatory cytokines Il-1β and Tnf-α by macrophages in zebrafish spinal cord regeneration. *Nature Communications*, *9*(1), 4670. https://doi.org/10.1038/s41467-018-07036-w

Walker, M. J., Hollands, A., Sanderson-Smith, M. L., Cole, J. N., Kirk, J. K., Henningham, A., McArthur, J. D., Dinkla, K., Aziz, R. K., Kansal, R. G., Simpson, A. J., Buchanan, J. T., Chhatwal, G. S., Kotb, M., & Nizet, V. (2007). DNase Sda1 provides selection pressure for a switch to invasive group A streptococcal infection. *Nature Medicine*, *13*(8), 981–985. https://doi.org/10.1038/nm1612

Wan, J., Ramachandran, R., & Goldman, D. (2012). HB-EGF Is Necessary and Sufficient for Müller Glia Dedifferentiation and Retina Regeneration. *Developmental Cell*, *22*(2), 334–347. https://doi.org/10.1016/j.devcel.2011.11.020

Wang, J. (2018). Neutrophils in tissue injury and repair. *Cell and Tissue Research*, *371*(3), 531–539. https://doi.org/10.1007/s00441-017-2785-7

Wang, M., Ma, W., Zhao, L., Fariss, R. N., & Wong, W. T. (2011). Adaptive Müller cell responses to microglial activation mediate neuroprotection and coordinate inflammation in the retina. *Journal of Neuroinflammation*, *8*(1), 173. https://doi.org/10.1186/1742-2094-8-173

Weber, A., Hochmann, S., Cimalla, P., Gärtner, M., Kuscha, V., Hans, S., Geffarth, M., Kaslin, J., Koch, E., & Brand, M. (2013). Characterization of Light Lesion Paradigms and Optical Coherence Tomography as Tools to Study Adult Retina Regeneration in Zebrafish. *PLOS ONE*, *8*(11), e80483. https://doi.org/10.1371/journal.pone.0080483

Webster, J. I., Tonelli, L., & Sternberg, E. M. (2002). Neuroendocrine regulation of immunity. *Annual Review of Immunology*, *20*(1), 125–163. https://doi.org/10.1146/annurev.immunol.20.082401.104914

White, D. T., Sengupta, S., Saxena, M. T., Xu, Q., Hanes, J., Ding, D., Ji, H., & Mumm, J. S. (2017). Immunomodulation-accelerated neuronal regeneration following selective rod photoreceptor cell ablation in the zebrafish retina. *Proceedings of the National Academy of Sciences*, *114*(18), E3719–E3728. https://doi.org/10.1073/pnas.1617721114

Wu, S., Xue, R., Hassan, S., Nguyen, T. M. L., Wang, T., Pan, H., Xu, J., Liu, Q., Zhang, W., & Wen, Z. (2018). Il34-Csf1r Pathway Regulates the Migration and Colonization of Microglial Precursors. *Developmental Cell*, *46*(5), 552-563.e4. https://doi.org/10.1016/j.devcel.2018.08.005

Xu, S., Webb, S. E., Lau, T. C. K., & Cheng, S. H. (2018). Matrix metalloproteinases (MMPs) mediate leukocyte recruitment during the inflammatory phase of zebrafish heart regeneration. *Scientific Reports*, *8*(1), 1–14. https://doi.org/10.1038/s41598-018-25490-w

Yang, J.-S., Lin, C.-W., Hsieh, Y.-H., Chien, M.-H., Chuang, C.-Y., & Yang, S.-F. (2017). Overexpression of carbonic anhydrase IX induces cell motility by activating matrix metalloproteinase-9 in human oral squamous cell carcinoma cells. *Oncotarget*, *8*(47), 83088–83099. https://doi.org/10.18632/oncotarget.20236

Yoong, S., O'Connell, B., Soanes, A., Crowhurst, M. O., Lieschke, G. J., & Ward, A. C. (2007). Characterization of the zebrafish matrix metalloproteinase 9 gene and its developmental expression pattern. *Gene Expression Patterns*, *7*(1–2), 39–46. https://doi.org/10.1016/j.modgep.2006.05.005

Zhang, Z., Hou, H., Yu, S., Zhou, C., Zhang, X., Li, N., Zhang, S., Song, K., Lu, Y., Liu, D., Lu, H., & Xu, H. (2020). Inflammation-induced mammalian target of rapamycin signaling is essential for retina regeneration. *Glia*, *68*(1), 111–127. https://doi.org/10.1002/glia.23707

Zhao, X. F., Wan, J., Powell, C., Ramachandran, R., Myers, M. G., & Goldman, D. (2014). Leptin and IL-6 family cytokines synergize to stimulate Müller Glia reprogramming and retina regeneration. *Cell Reports*, *9*(1), 272–284. https://doi.org/10.1016/j.celrep.2014.08.047

Zhou, Y., Hong, Y., & Huang, H. (2016). Triptolide Attenuates Inflammatory Response in Membranous Glomerulo-Nephritis Rat via Downregulation of NF-κB Signaling Pathway. *Kidney and Blood Pressure Research*, *41*(6), 901–910. https://doi.org/10.1159/000452591

Zhu, S., Yu, Y., Ren, Y., Xu, L., Wang, H., Ling, X., Jin, L., Hu, Y., & Zhang, H. (2021). The emerging roles of neutrophil extracellular traps in wound healing. *Cell Death Dis*, *12*(October), 984. https://doi.org/10.1038/s41419-021-04294-3

1159   Zou, J., Lathrop, K. L., Sun, M., & Wei, X. (2008). Intact Retinal Pigment Epithelium
1160   Maintained by Nok Is Essential for Retinal Epithelial Polarity and Cellular Patterning in
1161   Zebrafish. *Journal of Neuroscience*, *28*(50), 13684–13695.
1162   https://doi.org/10.1523/jneurosci.4333-08.2008

1163

Figure 1: Sterile phototoxic ablation of photoreceptors triggers leukocyte accumulation. (A) Retinal sections show that the majority of L-Plastin+ cells are *Tg(mpeg1:mCherry)*+, display morphological changes after lesion, and accumulate at the outer nuclear layer (ONL) in response to lesion. However, L-Plastin+ *Tg(mpeg1:mCherry)* negative cells were also observed upon injury (arrowhead). (B) Quantification on sections of L-Plastin+ cells in sham and regenerating retinae at 1, 2, 4, 7 and 14 days post lesion (dpl) shows an increase of cells within 2dpl that is released within 14 dpl. (C) In contrast to sham, *Tg(mpo:GFP)*+ neutrophils are detected upon lesion. They gather at the lesion site (12 hpl) and a diffuse GFP positive pattern (matrix) is formed from 15 hpl onwards. (D) Retinal flat mounts of *Tg(opn1sw1:GFP)* x *Tg(mpeg1:mCherry)* show the ramified structure of leukocytes in a UV-cone-specific reporter line and their reaction to lesion. Upon light lesion, leukocytes increase in number at the lesion site and display an amoeboid, activated morphology and accumulate in the central part of the lesion while unharmed areas appear devoid of *mpeg1:*mCherry positive cells. Scale bars in A & C: 50 µm Scale Bars in D: 200µm, inlets 20µm, error bars indicate standard error; * = p≤0,05; ** = p≤0,01; *** = p<0,001; n≥4; ANOVA Tukey's post hoc test; INL= inner nuclear layer; GCL=ganglion cell layer.

Figure 2: Müller glia transiently activate the *NF-κB:*GFP reporter and express the *NF-κB* target metalloproteinase *mmp9* in response to injury. (A) In comparison to sham, the *NF-κB:*GFP reporter is activated at 1 and 2 days post lesion (dpl) in *gfap:*NLS-mCherry labeled Müller glia localized in the inner nuclear layer (INL) and photoreceptors of the outer nuclear layer (ONL) and is restricted to the INL at 4 dpl. (B) *In situ* hybridization of *mmp9* in combination with immunohistochemistry for proliferating cell nuclear antigen (PCNA) and glial fibrillary acidic protein (GFAP/Zrf-1) labelling Müller glia, shows that *mmp9* is not expressed in sham retinae. In contrast, *mmp9* is transiently expressed at 1 and 2 dpl and returns to undetectable levels at 4 dpl. Scale bar: 50µm. GCL=ganglion cell layer.

Figure 3: Immunosuppression interferes with leukocyte accumulation and Müller glia reactivity. (A) Scheme of experimental outline. Fish were treated with Dexamethasone (Dex) or vehicle (Methanol; MeOH) from 10 days prior to injury until the day of analysis (sham, 1, 2, 4, 7 and 14 days post lesion; dpl). (B) Dex-treatment reduces the number of L-Plastin+ leukocytes (arrowheads) in sham (left panel) and regenerating retinae at 2 dpl (right panel). (C) Quantification of L-Plastin+ cells in MeOH- and Dex-treated sham or lesioned animals at 1, 2, 4, 7 and 14 dpl. (D) Immunohistochemistry for proliferating cell nuclear antigen (PCNA) reveals impaired proliferation in Dex-treated animals at 4 dpl. (E) Quantification of PCNA+ cells in MeOH- and Dex-treated sham and regenerating retinae at 1, 2, 4, 7 and 14 dpl shows impaired proliferative response in the Dex-treated group. (F) Immunohistochemistry for PCNA in *gfap*:NLS-GFP labeled Müller glia shows reduced numbers of proliferating Müller glia in the Dex treated group at 2 dpl compared to MeOH controls (arrowheads indicating non proliferative Müller cells). (G) Quantification of proliferating Müller glia in vehicle and Dex-treated sham and regenerating retinae at 1, 2, 4, 7 and 14 dpl, indicating that Dex hinders Müller glia proliferation. (H) In comparison to vehicle, the *NFκB*:GFP reporter is not activated in the inner nuclear layer (INL) of Dex-treated retinae at 2 dpl. (I) Injury-induced expression of *mmp9* is strongly reduced in Dex-

treated retinae at 2 dpl. Scale bar: 50 µm, Error bars indicate standard error; ns = p>0,05, * = p≤0,05; ** = p≤0,01; *** = p<0,001; n≥6; two-tailed student's T-Test, GCL=ganglion cell layer, ONL=outer nuclear layer.

Figure 4: Long-term immune suppression impairs photoreceptor regeneration.
(A) Scheme of experimental outline. *opn1sw1*:GFP, labeling UV-cones, transgenic animals were treated with Dexamethasone (Dex) or vehicle (Methanol; MeOH) from 10 days prior to lesion until the day of analysis (28 days post lesion; dpl 38 days post treatment; dpt). EdU pulses were applied at 2, 4, 6, 8 and 10 dpl. (B) In comparison to MeOH-treated animals, the number of regenerated EdU+ UV-cones is significantly reduced after Dex-treatment (compare upper panel with lower panel). Furthermore, the morphology of photoreceptor cells is disrupted in comparison to MeOH controls (arrowheads). (C) Quantification of EdU+ cells with respect to retinal layers showing significant reduction of EdU positive nuclei in the Dex-treated group. Scale bars: 50 µm. Error bars indicate standard error; * = p≤0,05; ** = p≤0,01; *** = p<0,001; n≥6; two-tailed student's T-Test. ONL=outer nuclear layer; INL=inner nuclear layer; GCL=ganglion cell layer.

Figure 5: Genetic reduction of microglia inhibits retinal proliferation in response to injury. (A) In contrast to heterozygous control siblings (*irf8*[+/-]), interferon regulatory factor 8 (*irf8)* homozygous myeloid-deficient mutant retinae (*irf8*[-/-]) display a decreased number of L-Plastin[+] leukocytes in homeostasis. (B) Quantifications of L-Plastin[+] cells in *irf8*[-/-] and control heterozygous siblings *irf8*[+/-]. (C) Scheme of experimental outline. *irf8*[-/-] and *irf8*[+/-] animals received an EdU pulse at 3 days post lesion (dpl) and were analyzed at 4 dpl. (D) During regeneration, the accumulation of L-Plastin[+] cells at the outer nuclear layer (ONL) is impaired in *irf8*[-/-] but not in control *irf8*[+/-] siblings. Moreover, the number of EdU[+] cells is significantly decreased in *irf8*[-/-] retinae. (E) Quantification of L-Plastin[+] cells and EdU[+] nuclei in *irf8*[-/-] and control *irf8*[+/-] animals at 4 dpl, show

reduced amounts of positive cells respectively. Scale bar: 50 μm. Error bars indicate standard error; * = p≤0,05; *** = p<0,001; n=5; two-tailed student's T-Test; INL=inner nuclear layer; GCL ganglion cell layer.

Figure 6: Inflammatory stimuli trigger Müller glia reactivity. (A) Scheme of experimental outline. Wild type or *NF-κB*:GFP reporter animals were injected with flagellin or zymosan into the vitreous of the eye and analyzed at 2 and 4 days post injection (dpi). (B) Quantification of TUNEL⁺ nuclei in control and injected retinae at 2 dpi revealing no increase in cell death. (C) Quantification of L-Plastin⁺ cells in control (PBS) and flagellin or zymosan injected retinae at 2 and 4 dpi display no significant increase in leukocytes due to the injection of factors. (D) Immunohistochemistry for proliferating cell nuclear antigen (PCNA) reveals proliferation in flagellin or zymosan injected but not in control animals at 4 dpi. (E) Quantification of PCNA⁺ nuclei in injected retinae at 2 and 4 dpi. (F) In comparison to PBS injected animals, injection of flagellin or zymosan results in strong activation of the *NFκB*:GFP reporter at 2 dpi. (G & H) *In situ* hybridization of *mmp9* and *her4.1* show expression of both genes in flagellin or zymosan injected but not in PBS injected eyes at 2 and 4 dpi, respectively. Scale bar: 50 µm. Error bars indicate standard error; * = p≤0,05; ** = p≤0,01; *** = p<0,001; n=6; two-tailed student's T-Test; ONL=outer nuclear layer; INL=inner nuclear layer; GCL=ganglion cell layer.

A

Injection   Analysis   Analysis

0   2   4 dpi

B

PBS
M-CSF

TUNEL$^+$ [mm^{-1}]

0,4
0,3
0,2
0,1

2dpi

C   PCNA DAPI 4dpi

PBS   ONL INL GCL

M-CSF

D

PBS
M-CSF

PCNA$^+$ [mm^{-1}]

30
20
10

***

4dpi

E   L-Plastin DAPI 4dpi

PBS

M-CSF

F

PBS
M-CSF

L-Plastin$^+$ [mm^{-1}]

40
30
20
10

**

4dpi

G   *NF-κB*:GFP DAPI 2dpi

PBS

M-CSF

H   *mmp9* 2dpi

PBS   GCL INL ONL

M-CSF

Figure 7: M-CSF stimulates an inflammatory response and initiates Müller glia cell cycle re-entry in the absence of lesion. (A) Scheme of experimental outline. Wild type or *NFκB*:GFP reporter animals were injected with M-CSF or PBS into the vitreous of the eye and analyzed at 2 and 4 days post injection (dpi). (B) Quantification of TUNEL+ nuclei in control and M-CSF-injected retinae at 2 dpi reveals no chance in cell death. (C) Immunohistochemistry for proliferating cell nuclear antigen (PCNA) reveals proliferation in M-CSF injected but not in control injected animals at 4 dpi. (D) Quantification of PCNA+ cells in control (PBS) and M-CSF injected retinae at 4 dpi. (E) L-Plastin staining shows more positive cells in M-CSF injected retinae at 4 dpi compared to controls. (F) Quantification of L-Plastin+ cells in control and M-CSF injected retinae at 4 dpi, show accumulation of L-Plastin cells in M-CSF injected specimen. (G) In comparison to PBS, injection of M-CSF results in strong activation of the *NFκB*:GFP reporter at 2 dpi. (H) *In situ* hybridization of *mmp9* shows transcriptional activation of this gene in M-CSF injected but not in control-injected animals at 2 dpi. Scale bars = 50µm, Error bars indicate standard error; ** = p≤0,01; *** = p<0,001; n>5; two-tailed student's T-Test. ONL=outer nuclear layer; INL=inner nuclear layer; GCL=ganglion cell layer.

Figure 8: Schematic summary of the impact of inflammation in retinal regeneration. In response to injury, microglia undergo a phenotypic change from amoeboid to ramified shape and accumulate at the lesion site. This process is inhibited by immune suppression. The reactive proliferation of Müller glia during regeneration is also impaired upon immunosuppression. Leukocytes stimulate the proliferation of neuronal precursor cells. M-CSF as inflammatory mediator triggers Müller glia proliferation, leukocyte accumulation and the generation of neuronal precursor cells. This process may also act indirectly via the stimulation of immune cells. Overall, the innate immune system, via mediators such as M-CSF, supports the differentiation into photoreceptor cells and thereby acts as a regenerative cue for retina regeneration.