Machine learning of cellular metabolic rewiring

Joao B. Xavier

Program for Computational and Systems Biology

Sloan Kettering Institute for Cancer Research

Abstract

Metabolic rewiring allows cells to adapt their metabolism in response to evolving environmental conditions. Traditional metabolomics techniques, whether targeted or untargeted, often struggle to interpret these adaptive shifts. Here we introduce *MetaboLiteLearner*, a machine learning framework that harnesses the detailed fragmentation patterns from electron ionization (EI) collected in scan mode during gas chromatography/mass spectrometry (GC/MS) to predict abundance changes in metabolically adapted cells. When tested on breast cancer cells with different preferences to metastasize to specific organs, *MetaboLiteLearner* successfully predicted the impact of metabolic rewiring on metabolites withheld from the training dataset using only the EI spectra, without metabolite identification or pre-existing knowledge of metabolic networks. Our analysis highlights shared and unique metabolomic shifts between brain- and lung-homing metastatic lineages, suggesting potential organ-tailored cellular adaptations. *MetaboLiteLearner*'s integration of machine learning and metabolomics paves the way for new insights into complex cellular adaptations.

Significance

Metabolic rewiring—the cellular adaptation to shifts in environment and nutrients—plays key roles in many contexts, including cancer metastasis. Traditional metabolomics often falls short of capturing the nuances of these metabolic shifts. This work introduces *MetaboliLiteLearner*, a machine learning framework that harnesses the rich fragmentation patterns from electron ionization collected in scan mode during gas chromatography/mass spectrometry, paving the way for enhanced insights into metabolic adaptations. Demonstrating its robustness on a breast cancer model, we highlight *MetaboliLiteLearner*'s potential to reshape our understanding of metabolic rewiring, with implications in diagnostics, therapeutics, and basic cell biology.

Introduction

Cells dynamically rewire their intracellular metabolism in response to changing nutrients, signals, and environmental cues (1). These adjustments allow cells to maintain homeostasis, optimize energy production, and fulfill the biosynthetic demands for growth and repair. From unicellular organisms like yeast adapting to changing nutrient availability (2) to the cells of multicellular organisms during development (3), disease (4) , or environmental stress (5), metabolic rewiring is a universal feature of cellular adaptation and survival.

Metabolic rewiring shifts the biochemical composition of cells, and the changes observed in the intracellular metabolome provide a window into the underlying cellular adaptation (6). Traditional approaches to metabolomics are either targeted or untargeted. Targeted metabolomics allows for the accurate and sensitive quantification of a predefined set of metabolites but risks missing novel or unexpected ones (7, 8). On the other hand, untargeted metabolomics allows for the comprehensive survey of all detectable metabolites within a biological sample. Still, it has challenges regarding reproducibility, identification of unknown compounds, and handling complex data sets (9).

To address these limitations, we present *MetaboliLiteLearner.* This novel computational method
utilizes machine learning to investigate metabolic rewiring using metabolomic data without
relying on prior knowledge of the metabolic network or having to identify the compounds from
their spectra. *MetaboliLiteLearner* deploys the extensive molecular fragmentation achieved
through electron ionization (EI) in gas chromatography/mass spectrometry (GC/MS), treating the
fragmentation patterns as input features in a supervised learning model; it associates these
features with the changes in metabolite abundance observed in cells undergoing metabolic
rewiring, effectively using the information about the molecular structure of metabolites to predict
how their abundances change in cells adapted metabolically to diverse circumstances.

As proof of concept, we utilized clones derived from the MDA-MB-231 breast cancer cell line,
which home to the lung and brain (10). These derivatives, developed through *in vivo* selection in
mice (11, 12), show pronounced transcriptomic shifts in laboratory cultures. In a previous study
(13), we employed targeted metabolomics to analyze a comprehensive panel of 645 metabolites
from these cells. Our analyses unveiled distinct intracellular metabolic profiles between the
brain- and lung-homing cells and their parental counterparts. The lung-homing cells displayed a
pronounced Warburg effect, underscored by a high lactate dehydrogenase (LDH) to pyruvate
dehydrogenase (PDH) expression ratio, a potential biomarker for lung metastasis. We confirmed
these distinct metabolic rewiring patterns through direct metabolic exchange measurements and
indirect assessments via stable isotope tracing. Given this detailed characterization, the MDA-
MB-231 system serves as an exemplary model to benchmark the efficacy of
MetaboliLiteLearner.

MetaboliLiteLearner confirmed that brain- and lung-homing cells exhibit unique metabolic
features that may be adaptations to the specific challenges of their target organs. This insight
aligns with the findings from our earlier approach (13), which relied on a large panel of
metabolites validated and measured through LC/MS, a method considered more laborious and
expensive. Unlike the earlier approach, *MetaboliLiteLearner* achieves this using GC/MS without
relying on prior information such as spectral libraries or extensive validation to draw its insights.
Beyond streamlining the analytical process, *MetaboliLiteLearner* paves the way for in-depth
insights into metabolic rewiring patterns by finding associations between molecular structure and
metabolic adaptability.

Results

***MetaboliLiteLearner*: Theory and Application.** *MetaboliLiteLearner* capitalizes on GC/MS
data from intracellular metabolome extracts. The extracts are primed for GC/MS analysis using
trimethylsilyl (TMS) derivatization, which enhances metabolite volatility. EI fragmentation to
provide a detailed molecular structure spectrum (14). The TMS derivatization, combined with
GC/MS and EI fragmentation, produces reproducible spectra and retention times for each
metabolite. While targeted metabolomics often measures specific ion peak areas (**Fig. 1A**),
MetaboliLiteLearner harnesses the entirety of data available in scan mode (**Fig. 1B**). This spans
ion fragment abundances from 50 to 600 mass-to-charge ratio units (m/z) and all GC retention
times.

MetaboliLiteLearner integrates full spectra with the corresponding log two-fold changes in
abundance, reflecting their abundance changes due to metabolic rewiring. Each molecule's EI
fragmentation spectrum is encoded as a 550-dimensional vector (**Fig. 1C**), with corresponding
log2 fold changes (obtained by comparing baseline vs. rewired cell metabolite abundances) as

the training data labels. Following the supervised learning paradigm, the model's efficacy is
gauged by predicting log2 fold changes on a new compound set and comparing these predictions
with actual data (15).

MetaboliLiteLearner uses Partial Least Squares Regression (PLSR) to construct a linear model
by projecting the predictors and the response variables onto a new N-dimensional space (16).
PLSR generates a model that balances complexity and interpretability without sacrificing the
power of linear combinations of original variables. This approach is analogous to an Artificial
Neural Network (ANN) that employs a hidden layer with N neurons, except that PLSR employs
linear functions instead of non-linear activation functions networks (17).

To transform the high-dimensional input (a 550-dimensional EI spectrum) and output (log2 fold
changes) into a latent space, *MetaboliLiteLearner* uses two matrices: 'w' for the spectra and 'c'
for the log2 fold changes (**Fig. 1D**). The latent space dimensionality, N, dictates the model's
complexity. An optimal number of dimensions, N_{opt}, is determined through cross-validation to
prevent under- and overfitting (15). When applied to metabolomics data, the loadings derived
from PLSR—coefficients that describe the relationship between the original predictors and the
new latent factors—provide insights into the underlying cellular adaptations. These loadings are
directly tied to the EI-fragmentation spectra of metabolites and indicate molecular structural
features linked to their abundance changes in metabolically rewired cells. Therefore, once the
model is trained, the loadings provide insights into the relationships between metabolite structure
and metabolic rewiring, shedding light on cellular adaptation mechanisms

Breast Cancer Cell Data Integration into *MetaboliLiteLearner*. We collected the data for
MetaboLiteLearner from the MDA-MB-231 breast cancer cell line and its brain and lung-
targeted derivatives. The derivatives, originating from *in vivo* mouse selection (11, 12), exhibited
transcriptomic differences compared to the original MDA-MB-231 cells in lab cultures. Our
prior work identified marked intracellular metabolome variances between the brain- and lung-
homing cells and the parent cells, indicating metabolic rewiring (13).

We cultured all three cell variants, and the metabolites were harvested during balanced growth,
which is when cells divide at exponential growth and before any slowdown in growth due to
confluency (**Fig. 2A**). We then extracted soluble metabolites, dried all samples and controls, and
derivatized the polar metabolites using trimethylsilyl (TMS) derivatization. Post-processing with
GC/MS with EI in scan mode (18), we aligned all data to consistent retention (6-30 mins at 0.01
min intervals) and m/z (50 to 600 m/z at 1 m/z steps) ranges. The resulting individual data
matrices were joined into a singular "virtual bulk sample" matrix (**Fig. 2B**). Spectrum extraction
from this composite data enabled us to determine the log2 fold change in metabolite abundance
in the organ-homing cells relative to the parental cells through a linear mixed-effects model. Our
input for *MetaboliLiteLearner* consisted of spectra arrays from m/z intervals of 50 to 600,
normalized to their norms. The dataset, encompassing 153 unique spectra alongside their
respective log2 fold-changes, is now presented as the *MetaboLiteLearner* Open Dataset
(MLOD).

Training and Evaluating *MetaboLiteLearner*. The analysis began by determining the optimal
number of latent components, N_{opt}, using the MLOD. Through hold-out cross-validation, as we
increased N from 1 (simplest model) to 30 (most complex), the training error dropped
monotonically, suggesting a better fit of the training data with increased model complexity (**Fig.**
3A). The test error reached a minimum at $N=11$ components before rising, indicating potential
overfitting in the more complex models. Utilizing a conservative methodology from supervised

learning (15), we selected N_{opt}=5, which showcased the smallest test error within one standard
error. The N_{opt}=5 model predictions correlated robustly with the actual log2 fold changes
(ρ=0.39, P-value$\ll$0.01). We should note that other datasets may require a different value of N_{opt}.

The model with N_{opt}=5, when trained on the entire MLOD, optimally adjusted its loadings to
maximize covariance between the projections of both inputs (the spectra) and outputs (log2 fold
changes) onto the 5-dimensional latent space (**Fig. 3B**). Unlike unsupervised methods such as
principal component analysis (PCA), which primarily focuses on variance within individual
datasets, the PLSR emphasizes joint variance optimization (15). The resultant transformations
into the latent space accounted for 32% of the predictor variance and 68% of the response
variance. The variance explained by each latent factor for the predictor and response can be
analyzed separately (**Fig. 3B'** and **Fig. 3B''**, respectively).

Next, we performed a randomization test to determine our model's ability to capture biologically
relevant patterns, not statistical artifacts. By shuffling the log2 fold changes, we retained internal
correlations but disrupted correlations with input spectra. Notably, *MetaboLiteLearner*'s error,
when trained with these shuffled data, was consistently worse than with the original dataset (**Fig.**
3C). This reaffirms our model's capability to discern significant links between metabolite spectra
and abundance shifts in rewired cells.

MetaboLiteLearner **Reveals Metabolic Changes in Metastatic Breast Cancer Cells.** The
optimal model with 5 latent components (N_{opt}=5) trained on the entire MLOD can transform a
spectrum (a 500-dimensional vector) into log2 fold changes for brain- and lung-homing cells (a
2-dimensional vector). This transformation can be visualized using a biplot, with each m/z ionic
fragment represented by a vector (**Fig. 1A**). The biplot shows that specific fragments, such as
m/z=104 which arises from the EI fragmentation of TMS derivatives of amino acids (14),
correlate with increased levels in both cell types. In contrast, fragments like m/z=306, associated
with EI-fragmentation of certain TMS-derivatized sugars (14), relate to decreased levels.

To delve deeper, we analyzed a set of 263 EI-fragmentation spectra for TMS-derivatized
metabolites (18) which we grouped into seven categories from the KEGG database of
compounds with biological roles (19). We fed the spectra into our trained model to predict fold
changes. While most metabolites showed concurrent changes in both cell types, specific
carbohydrates, and nucleic acids increased only in lung-homing cells (**Fig. 1A**).

The 5-latent-component model explains 67.7% of the variance in log2 fold changes (**Fig. 3B''**).
PLSR's strength lies in decomposing the model's coefficients into latent components. These
components, organized based on their contribution to explain the joint variance, shed light on the
molecular shifts underpinning cellular adaptations. Components 1 and 3 showcase metabolites
with decreased levels in both cell types, indicating overlapping metabolic shifts (**Fig. 4B**).
Components 2 and 5 indicate metabolites with increased levels in both cell types. Component 4
highlights differences between the cell types—some metabolites decrease in brain-homing but
increase in lung-homing cells.

Component 1, which explains 27% of the response variance, captures a common trend—reduced
levels in both cell types. Evaluating spectra from various compound classes, most amino acids
follow this trend, whereas carbohydrates deviate from it (**Fig. 4C**). Analyzing all latent
components (**Fig. 4D-G**) revealed component 4's unique role. Despite accounting for just 4.9%
of the response variance, it distinctly captures the variation between brain- and lung-homing
cells. Carbohydrates and deoxyribonucleosides dominate this component, indicating potential
metabolic shifts between the two cell types. These observed metabolic changes could reflect

adaptations to the unique environments of the brain and lungs. With its rich blood supply, the
lung might favor carbohydrate-utilizing cells. Conversely, the elevated deoxyribonucleosides in
lung-homing cells could suggest more robust DNA repair mechanisms than in brain-homing
cells. These findings align with our previous work (13) and suggest potential pathways for
further exploration.

Discussion

Here, we presented the capabilities of *MetaboLiteLearner*, a supervised learning approach to
study the rewiring of intracellular metabolomes. Through computational experiments, we have
shown that *MetaboLiteLearner* can discern patterns even within the confines of a relatively small
dataset comprising 153 unidentified metabolites without prior knowledge of metabolic networks.

A defining feature of *MetaboLiteLearner*'s is its use of PLSR—a computationally lightweight
and robust statistical algorithm, especially useful for smaller datasets. PLSR can handle internal
correlations within predictors and responses, which in our case arise due to shared ion fragments
produced by naturally abundant isotopes and functional groups shared among certain metabolite
classes and the universally witnessed metabolic shifts in disseminated cells, as noted in previous
work (13). PLSR handles these internal correlations by generating a reduced-dimensional latent
space that maximizes the joint variance between predictors and responses. Here, the latent space
represents the leading associations between a metabolite's molecular features (the ions produced
by EI fragmentation) and its abundance change in rewired cells. While the current
implementation of PLSR in our model is efficient, there is potential for further refinement, such
as using regularization and Laplacian constraints (20). Incorporating regularization might
improve model interpretability. Meanwhile, Laplacian constraints could let the model
incorporate existing knowledge from comprehensive spectral libraries and metabolic network
models.

In this study, we also introduce the MLOD, an open dataset tailored for machine learning
research focusing on the metabolic reconfigurations in cancer cells. The MLOD, enriched with
meticulously captured spectra and corresponding abundance shifts, can be a standard for
benchmarking future supervised learning endeavors in this domain. Our decision to use GC/MS
spectra from TMS-derivatized samples within a specific interval was primarily driven by its
alignment with prevalent practices (18). However, we recognize the potential benefits of
integrating data from high-resolution instruments like time-of-flight (TOF) or Orbitrap. These
devices could capture finer details, thus enhancing model predictions. Furthermore, harnessing
data generated by tandem instruments, such as MS/MS or MS2, can open doors to more
sophisticated computational strategies, as witnessed in works external to cancer research (21–
23).

Our dataset is limited by its focus on log2 fold changes computed from unsynchronized
intracellular metabolite concentrations. Thus, our insights primarily pertain to consistent changes
across a generalized cell population. Synchronized data acquisition at distinct cellular growth
stages (24) or the integration of stable isotope tracing (25) can add dynamics to future datasets,
further enhancing the granularity of our models. Our breast cancer case study had a
bidimensional response variable, representing metabolite abundance changes in brain- and lung-
homing cells. As we move forward, incorporating additional dimensions and more expansive
datasets could pave the way for deploying advanced computational techniques, like deep neural
networks.

This study showcases the potential synergy between machine learning and metabolomics. With
evolving datasets and improving computational methods, we can make new strides in unraveling
the intricacies of metabolic rewiring—a fundamental aspect of cellular adaptation.

Methods

Cell Culture, Metabolite Extraction and Derivatization. Cell lines were cultured in DMEM
(Fisher 11965118) supplemented with 10% FBS, produced in the MSKCC media core facility,
and 1% penicillin/streptomycin (Fisher 15140122). The cultivation conditions included a 37°C
incubator with regulated humidity and a 5% CO2 atmosphere. Authenticated cell lines were
procured from the Massague lab and were developed as delineated in previous studies (11, 12).
Cells were subjected to metabolite extraction using 1 ml of ice-cold 80% methanol, followed by
overnight storage at -80°C. Subsequent to this, the extracts underwent a drying process using an
evaporator. Resuspension was achieved by incubation with shaking at 30°C for 2 hours in a
solution containing 50 μl of 40 mg ml−1 methoxyamine hydrochloride in pyridine.
Derivatization was performed by adding 80 μl of N-methyl-N-(trimethylsilyl) trifluoroacetamide
(with or without 1% TCMS from Thermo Fisher Scientific) and 70 μl of ethyl acetate (sourced
from Sigma-Aldrich). This mixture was then incubated at 37°C for 30 minutes.

Gas Chromatography/Mass Spectrometry (GC/MS) Analysis. Analytical procedures utilized
the Agilent 7890A gas chromatograph paired with an Agilent 5977C mass selective detector. The
gas chromatograph operated in splitless injection mode, maintaining a constant helium gas flow
at 1 ml/min. The injection involved introducing 1 μl of the derivatized metabolites onto an HP-
5ms column. The temperature of the gas chromatograph oven was systematically ramped from
60°C to 290°C over a 25-minute interval. Samples comprised four distinct types: blank media,
parental cells, lung-homing cells, and brain-homing cells. Each type was cultured in triplicate
groups over a span of three days, resulting in nine replicates for each sample category.

Data Processing: GC/MS raw data, stored in the Agilent .D format, underwent processing to
generate the MLOD dataset. These raw .D files were initially converted into CSV. By creating a
"virtual bulk sample" from this CSV data, we could detect and extract spectra from the total ion
chromatogram (TIC). With the aid of the Matlab function *mspeaks*, peaks associated with
different metabolites were identified within the TIC, followed by the extraction of their spectra.
An integration process then yielded a peak area table. To maintain the relevance of the dataset,
compounds that did not exhibit significant differences compared to blank media samples, as
determined using ANOVA, were removed. Following this filtering process, the refined data
formed the MLOD dataset, which has 153 unique spectra labeled with abundance alterations,
represented as log2 fold changes for both brain-homing and lung-homing cells.

Machine Learning: Supervised learning attempts to learn from training data containing inputs
matched to their correct outputs, a model $Y = f(X)$ that can generalize to a new dataset (15).
During the learning phase, the model is a "student" given a set X_{train}, Y_{train} of training. For each
example x_i in the training data set X_{train} the model presents the answer $f(x_i) = \hat{y}_i$. The
supervisor or "teacher" compares $\hat{y}_i$ with the correct answer, y_i, and gives an error associated
with the "student's" answer. The examples' errors are used to calculate a loss, $L(y_i, \hat{y}_i)$, and the
model learns by adjusting its parameters to minimize the loss. After the training, the model is
presented with a test dataset (X_{test}) containing a new set of examples. The model's accuracy in

predicting the output for the new inputs shown is evaluated by comparing these predictions $\hat{Y}_{test}$
with the actual data, Y_{test}.

MetaboLiteLearner learns a linear function f that takes a feature vector representation of a
metabolite as input. It predicts, as output, how the intracellular level of that metabolite is
impacted by metabolic rewiring. Each metabolite is represented by an input vector x_i, a p-
dimensional array of the features of metabolite i. In our case, x_i is 550-dimensional, representing
the abundances of ionic fragments of metabolite i of sizes between 50 and 600 mass-to-charge
ratio units (m/z) binned at unit intervals. This vector is obtained from the electron ionization (EI)
mass spectrum. The EI-spectrum is a mass-to-charge histogram of the ion fragments produced
when a metabolite undergoes breakdown via electron ionization. It offers a direct insight into the
metabolite's structure—and potentially its function—without needing to identify that metabolite
first. The output vector y_i is two-dimensional, representing the $\log_2$ fold-change of that
metabolite in brain-homing and lung-homing cells compared to the parental lineage.

MetaboLiteLearner's learning algorithm is the partial least squares regression (PLSR) (16). This
algorithm has similarities with artificial neural networks (17) but is more stable and less
computationally demanding to run. We used the MATLAB implementation of PLSR encoded in
function *plsregress*. Leave-one-out cross-validation was used to determine the optimal number of
latent components, N_{opt}. To ensure the robustness of the analytical outcomes, a shuffling test was
administered, wherein the order of observed data was randomly rearranged a thousand times.
Augmenting the biological context of the study, a comprehensive table detailing compounds with
biological roles was sourced from the Kyoto Encyclopedia of Genes and Genomes (KEGG) (19).
The spectra for TMS-derivatized compounds were obtained from the Fiehn Library (18). In
conjunction with the KEGG table, these spectra were input into the *MetaboLiteLearner* model to
compute their fold changes in brain- and lung-homing cells and the values for the five latent
components.

Data Availability: The raw GC/MS data supporting the findings of this study are available in
Zenodo (26).

Code Availability: The data processing and analysis code is available on GitHub (27).

Acknowledgments

This work was supported by the National Institutes of Health (NIH) grant R01 CA266068
"Mathematical modeling of metabolism rewiring in cancer eco-evolution and metastasis
tropism" to J.B.X. Acknowledgments to Chen Liao, Deepti Mathur, Heesoo Seong, and Carlos
Carmona Fontaine for their comments on the manuscript.

Bibliography

1.   Gomes AP, Blenis J (2015) A nexus for cellular homeostasis: the interplay between
metabolic and signal transduction pathways. *Curr Opin Biotechnol* 34:110–117.
2.   Moxley JF, et al. (2009) Linking high-resolution metabolic flux phenotypes and
transcriptional regulation in yeast modulated by the global regulator Gcn4p. *Proc Natl*
Acad Sci USA 106(16):6477–6482.
3.   Miyazawa H, Aulehla A (2018) Revisiting the role of metabolism during development.

Development 145(19).

4.   Chapman NM, Chi H (2022) Metabolic adaptation of lymphocytes in immunity and
disease. *Immunity* 55(1):14–30.

5.   Sengupta N, Rose ST, Morgan JA (2011) Metabolic flux analysis of CHO cell metabolism
in the late non-growth phase. *Biotechnol Bioeng* 108(1):82–92.

6.   Schmidt DR, et al. (2021) Metabolomics in cancer research and emerging applications in
clinical oncology. *CA Cancer J Clin* 71(4):333–358.

7.   Lu W, et al. (2017) Metabolite measurement: pitfalls to avoid and practices to follow.
Annu Rev Biochem 86:277–304.

8.   Wieder C, et al. (2021) Pathway analysis in metabolomics: Recommendations for the use
of over-representation analysis. *PLoS Comput Biol* 17(9):e1009105.

9.   Dunn WB, et al. (2013) Mass appeal: metabolite identification in mass spectrometry-
focused untargeted metabolomics. *Metabolomics* 9(S1):44–66.

10.   Cailleau R, Olivé M, Cruciger QV (1978) Long-term human breast carcinoma cell lines of
metastatic origin: preliminary characterization. *In Vitro* 14(11):911–915.

11.   Minn AJ, et al. (2005) Genes that mediate breast cancer metastasis to lung. *Nature*
436(7050):518–524.

12.   Bos PD, et al. (2009) Genes that mediate breast cancer metastasis to the brain. *Nature*
459(7249):1005–1009.

13.   Mathur D, et al. (2020) Divergent use of metabolic fluxes in breast cancer metastasis.
BioRxiv.

14.   Lai Z, Fiehn O (2016) Mass spectral fragmentation of trimethylsilylated small molecules.
Mass Spectrom Rev 37(3):245–257.

15.   Hastie T, Tibshirani R, Friedman J (2009) *The Elements of Statistical Learning* (Springer
New York, New York, NY). 2nd Ed.

16.   de Jong S (1993) SIMPLS: An alternative approach to partial least squares regression.
Chemometrics and Intelligent Laboratory Systems 18(3):251–263.

17.   Tzu-Chien Hsiao, Chii-Wann Lin, Mang-Ting Zeng, Chiang HHK (1998) in *Proceedings*
of the 20th Annual International Conference of the IEEE Engineering in Medicine and
Biology Society. Vol.20 Biomedical Engineering Towards the Year 2000 and Beyond (Cat.
No.98CH36286) (IEEE), pp 1341–1343.

18.   Kind T, et al. (2009) FiehnLib: mass spectral and retention index libraries for
metabolomics based on quadrupole and time-of-flight gas chromatography/mass
spectrometry. *Anal Chem* 81(24):10038–10048.

19.   Kanehisa M, Goto S (2000) KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic*
Acids Res 28(1):27–30.

20.   Li X, Panea C, Wiggins CH, Reinke V, Leslie C (2010) Learning "graph-mer" motifs that
predict gene expression trajectories in development. *PLoS Comput Biol* 6(4):e1000761.

21.   Watrous J, et al. (2012) Mass spectral molecular networking of living microbial colonies.
Proc Natl Acad Sci USA 109(26):E1743-52.

22.   Tripathi A, et al. (2021) Chemically informed analyses of metabolomics mass
spectrometry data with Qemistree. *Nat Chem Biol* 17(2):146–151.

23.   Stravs MA, Dührkop K, Böcker S, Zamboni N (2022) MSNovelist: de novo structure
generation from mass spectra. *Nat Methods* 19(7):865–870.

24.   Diehl FF, et al. (2022) Nucleotide imbalance decouples cell growth from cell proliferation.
Nat Cell Biol 24(8):1252–1264.

25.   Jang C, Chen L, Rabinowitz JD (2018) Metabolomics and isotope tracing. *Cell*
173(4):822–837.
26.   Xavier J (2023) Raw GC/MS data used for MLOD (MetaboLiteLearner). *Zenodo*.
27.   Xavier J (2023) joaobxavier/learn_metabolic_rewiring_matlab:
MetaboLiteLearner08032023. *Zenodo*.

Figures

Figure 1. Workflow and Functionality of *MetaboliLiteLearner*. (A) Traditional targeted
metabolomics uses specific ion peak areas. **(B)** *MetaboLiteLearner* uses GC/MS data acquired in
scan mode in *MetaboliLiteLearner*, which encompasses ion fragment abundances ranging from
50 to 600 m/z captured at all GC retention times. **(C)** Each molecule's electron impact (EI)
fragmentation spectrum is depicted as a 550-dimensional vector. These high-dimensional vectors
are paired with their corresponding log2 fold change values, which serve as training labels. **(D)**
The transformation matrices ('w' for spectra and 'c' for log2 fold changes) are the Partial Least
Squares Regression (PLSR) loadings used to map data into the N-dimensional latent space. This
enables *MetaboliLiteLearner* to learn the relationship between metabolite structure and
metabolic rewiring.

Figure 2. Data Acquisition and Processing for *MetaboLiteLearner* from Breast Cancer Cell
Derivatives. (A) Cell Culture and Metabolite Extraction: Breast cancer cell lines, including the
parental MDA-MB-231 cells and its brain- and lung-homing derivatives, were cultivated. These
derivatives were procured through *in vivo* selection using mice. Under consistent media
conditions *in vitro*, intracellular metabolites from these cells were extracted to ensure a reliable
data source for subsequent processing. **(B)** GC/MS Processing and Data Aggregation: Samples
underwent GC/MS analysis following the TMS derivatization protocol. The generated data
matrices, unique for each sample, were amalgamated to create a virtual "bulk" sample. Peaks
were identified, and their spectra were extracted from this consolidated matrix. The input (X) for
MetaboLiteLearner encompasses the mass spectra of each peak. The output data (Y) indicates
the comparative abundance shift of each peak in brain- and lung-homing cells relative to the
parental cells.

Figure 3. Optimization and Evaluation of MetaboLiteLearner's Predictive Performance.
(A) Model Complexity vs. Error: Through hold-out cross-validation, the training error
consistently decreased as latent components (N) increased from 1 to 30. Test error reached its
lowest at N=11 before it began to rise, highlighting potential overfitting with more complex
models. The chosen optimal model had N_{opt}=5 components, correlating strongly with true log2
fold changes (ρ=0.39, P-value$\ll$0.01). **(B)** Schematic of the model trained with 5 components.
(B' and B'') Variance Explained by Latent Factors: Using the Nopt=5 model, the
transformations into the 5-dimensional latent space covered 32% of the predictor variance and
68% of the response variance. The variance explained by each latent factor for the predictor and
response datasets can be viewed separately. **(C)** Randomization Test Results: After shuffling the
log2 fold changes, disrupting their correlations with input spectra, *MetaboLiteLearner*'s error
with shuffled data was consistently higher than with the original dataset, confirming its ability to
identify genuine relationships between metabolite spectra and abundance changes in rewired

cells.
404
Figure 4. Interpretation of the Model for Metastatic Breast Cancer Cells. (A) Biplot
representation of the m/z ionic fragment vectors. Specific fragments, such as m/z=104, are
associated with increased levels in both cell types in contrast to fragments like m/z=306 which
are associated with decreased levels. **(B)** The proportional contribution of the five latent
components to the variance in log2 fold changes, with components 1 and 3 indicating
overlapping metabolic shifts with decreased levels in both cell types, components 2 and 5
showcasing increases in both, and component 4 underscoring the divergence between the two
cell types. **(C)** In component 1, accounting for 27% of the response variance, most amino acids
follow the trend of reduced levels in both cell types, while carbohydrates differ. **(D-G)** Among
the latent components, we see the distinctive role of component 4 which is dominated by
carbohydrates and deoxyribonucleosides and highlights potential metabolic variances between
brain- and lung-homing cells.
417

A. Sampling and extraction

MDA-MB-231   Brain metastasis   Lung metastasis   Blank control

x 9   x 9   x 9   x 9

Extract intracellular metabolites

x 9   x 9   x 9   x 9

B. Data processing

 x 36

Derivatize, GC/MS analysis

Resample, integrate 236 peaks

236 peaks
x 36

Remove 83 background peaks
(153 peaks remain)

Extract average EI spectrum between
50 - 600 m/z, then normalize

$$\begin{matrix} x_{1,50} & x_{1,51} & x_{1,52} & \dots & \dots & \dots & \dots & x_{1,600} \\ x_{2,50} & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ x_{153,50} & \dots & \dots & \dots & \dots & \dots & \dots & x_{153,600} \end{matrix}$$

$= \quad$ 153 × 550
predictor variable
X

 vs.   Brain-homing vs. parental Log_2{Fold Change} × 153

 vs.   Lung-homing vs. parental Log_2{Fold Change} × 153

$= \quad$ 153 × 2
response
variable
Y