1 **TITLE:**
2 DIFFERENTIAL EFFECTS OF REACTIVE OXYGEN SPECIES ON AUTOPHAGY IN ARABIDOPSIS
3
4
5 **RUNNING TITLE:**
6 Redox regulation of autophagy
7
8

9 **AUTHORS:**
10 Alejandro Enet [1†], Germán Robert [1,2,3 † *], Laura Saavedra [1], Ramiro Lascano [1,2 *]
11
12 **†** Both authors contributed equally
13
14 *** Corresponding authors:**
15 robert.german@inta.gob.ar; ramiro.lascano@unc.edu.ar
16
17
AFFILIATIONS
[1] Plant Stress Biology Group, Unidad de Doble Dependencia INTA-CONICET (UDEA), Av. 11 de
Septiembre, Córdoba 4755-X5020ICA, Argentina.
[2] Cátedra de Fisiología Vegetal, Facultad de Ciencias Exactas Físicas y Naturales, Universidad
Nacional de Córdoba, Av. Velez Sarsfield 299, Córdoba, Argentina;
[3] Instituto Nacional de Tecnología Agropecuaria (INTA), Instituto de Fisiología y Recursos
Genéticos Vegetales (IFRGV), Av. 11 de Septiembre, Córdoba 4755-X5020ICA, Argentina.
25
26
27 Alejandro Enet: ale.403@hotmail.com
28 Germán Robert: robert.german@inta.gob.ar
29 Laura Saavedra: saavedra.laura@inta.gob.ar
30 Ramiro Lascano: ramiro.lascano@unc.edu.ar
31
32

33 **KEYWORDS**
34 Autophagy, stress, reactive oxygen species, singlet oxygen, superoxide radical, hydrogen
35 peroxide, redox state, Arabidopsis thaliana.

49　**HIGHLIGHT**
50　This research shows the differential effects of reactive oxygen species on autophagic flux,
51　highlighting their impact on the cytoplasmic redox state.
52
53

ABSTRACT
Autophagy plays a key role in the responses to different stress condition in plants. Reactive
oxygen species (ROS) are common modulators of stress responses, having both toxic and
signaling functions. In this context, the relationship between ROS and autophagy regulation
remains unclear, and in some aspects, contradictory. In this study, we employed
pharmacological and genetic approaches to investigate the effects of different ROS on the
cytoplastic redox state and autophagic flux in *Arabidopsis thaliana*. Ours results demonstrated
that oxidative treatments with H_2O_2 and MV, which drastically increased the oxidized state of
the cytoplasm, reduced the autophagic flux. Conversely, singlet oxygen, which did not have
significant effects on the cytoplasmic redox state, increased the autophagic flux. Additionally,
our findings indicated that after H_2O_2 and high light treatments and during the recovery period,
the cytoplasm returned to its reduced state, while autophagy was markedly induced. In
summary, our study unveils the differential effects of ROS on the autophagic flux, establishing a
correlation with the redox state of the cytoplasm. Moreover, it emphasizes the dynamic nature
of autophagy in response to oxidative stress and the subsequent recovery period.

69
70
71　**KEYWORDS**
72　Autophagy, stress, reactive oxygen species, singlet oxygen, superoxide radical, hydrogen
73　peroxide, redox state, Arabidopsis thaliana.
74
75
ABBREVIATIONS
ATG: autophagy genes
Conc A: Concanamicina A
GFP: green fluorescent protein
GSH: reduced glutathione
GSSG: oxidized glutathione
MV: methyl viologen
roGFP: redox-sensitive green fluorescent proteins
ROS: reactive oxygen species

INTRODUCTION

Plants as sessile organisms are constantly exposed to different biotic and abiotic stress conditions, so they have evolved integrated defense and acclimation mechanisms to cope with adverse conditions (Fichman and Mittler, 2020; Peláez-vico *et al.*, 2022). A common effect induced in plants by different stress conditions is the increase in the generation of reactive oxygen species (ROS), such as superoxide (O_2^-), hydrogen peroxide (H_2O_2), singlet oxygen ($^1O_2$), and the hydroxyl radical (-OH), at different subcellular compartments, such as chloroplasts, mitochondria, peroxisomes, cytosol and the apoplast (Apel and Hirt, 2004; Foyer and Noctor, 2009, 2011*a*). ROS are potent oxidants, that depending on their intensity, duration, and location site of production within the cell can modify different biomolecules, affecting the redox state in different subcellular compartments, provoking oxidative stress, inducing senescence symptoms and cell death. Furthermore, due to their high reactivity and toxicity, the maintenance of tightly regulated ROS levels becomes crucial for plant acclimation, adaptation, and survival, facilitated by the activity of a sophisticated plant antioxidative system (Apel and Hirt, 2004; Foyer and Noctor, 2009, 2011*b*; Trippi *et al.*, 2009). In this way, controlling the different types of ROS produced, along with their intensity, duration, and site of production, has diverse implications in redox signaling, influencing processes such as growth, development, and responses to changing environmental conditions (Melchiorre *et al.*, 2009; Miller *et al.*, 2009; Robert *et al.*, 2009, 2018; Muñoz *et al.*, 2012; Fernandez-göbel *et al.*, 2019; Fichman and Mittler, 2020; Zandalinas *et al.*, 2020; Peláez-vico *et al.*, 2022). Consequently, ROS are recognized as critical signaling molecules, and the cellular redox state is considered an intensive property of the living system that integrates information from metabolic, energetic and environmental cues controlling growth, development and acclimation responses, as well as cell suicide events (Breusegem and Dat, 2006; Trippi *et al.*, 2009; Fichman and Mittler, 2020; Peláez-vico *et al.*, 2022).

In plants, chloroplast, mitochondria and peroxisome, which are redox active organelles where carbon and energetic metabolisms take place, are major sources of ROS production (Apel and Hirt, 2004). In chloroplasts, the photosynthetic process involves physical, biophysical, and biochemical redox reactions with different reaction rates. In consequence, oxygen became a potent univalent electron or energy acceptor from highly activated intermediates under conditions of excess excitation energy (EEE) (Karpinski *et al.*, 1999; Foyer and Hanke, 2022). ROS produced in the chloroplasts are particularly important in the retrograde signaling of EEE conditions, affecting nuclear gene expression, acclimation responses as well as programmed cell death (Karpinski *et al.*, 1999; Breusegem *et al.*, 2001; Breusegem and Dat, 2006; Bechtold *et al.*, 2008; Fichman and Mittler, 2020; Foyer and Hanke, 2022).

Likewise, several stress conditions in plants have been also shown to be capable of triggering or increasing autophagy (Signorelli *et al.*, 2019; Wang *et al.*, 2021). Autophagy is an ancient and conserved process in eukaryotes by which different cellular components are delivered to vacuole/lysosome to be degraded, and is essential for plant energetic metabolisms, growth, development and response to different stress conditions (Su *et al.*, 2020; Wang *et al.*, 2021). Different types of autophagy have been identified in plants (Yoshimoto, 2012). Macroautophagy (hereafter termed autophagy) is the type of autophagy most thoroughly studied in all organisms. The autophagic process is initiated, and mainly modulated in the cytosol, by the formation of a double membrane cup, the phagophore, that expands to engulf macromolecules and cellular components to finally form a double-membrane vesicle called

autophagosomes. The autophagosomes are then transported and fused with the vacuole,
facilitating the delivery of their cargo for subsequent degradation (Yoshimoto, 2012; Yoshimoto
and Ohsumi, 2018). The several steps of the autophagic process are coordinated by ATG
(AuTophaGy) proteins as well as by some components of the endosome pathway, as recently
reported (Zhuang *et al.*, 2015; Yoshimoto and Ohsumi, 2018; Robert *et al.*, 2021). While early
evidence pointed to a strong transcriptionally regulation of autophagy in response to nutritional
starvation conditions and foliar senescence (Yoshimoto *et al.*, 2004; Breeze *et al.*, 2011), recent
studies have shed light on the widespread modulation of autophagy initiation and progression
by post-translational modifications of core autophagy proteins. These modifications primarily
occur at the cytoplasmic level in response to environmental cues (Feng *et al.*, 2015; Yang *et al.*,
2019; Qi *et al.*, 2021; Agbemafle *et al.*, 2023).
It has been suggested a positive correlation between ROS generation and the autophagic
process, where ROS can serve as both sources of damage and signals (Xiong *et al.*, 2007*a,b*;
Signorelli *et al.*, 2019). Autophagy promotes the degradation of ROS-overproducing damaged
organelles including chloroplasts, mitochondria, and peroxisomes, thus fulfilling antioxidant
roles (Xiong *et al.*, 2007*b*; Shibata *et al.*, 2013; Nakamura *et al.*, 2018; Oikawa *et al.*, 2022).
Supporting this notion, observations in different autophagy-mutant plants demonstrate the
accumulation of oxidized and ubiquitinated proteins (Xiong *et al.*, 2007*a*; Munch *et al.*, 2014),
high levels of H_2O_2 (Yoshimoto *et al.*, 2009), increased amounts of total glutathione (Masclaux-
Daubresse *et al.*, 2014), accelerated senescence (Doelling *et al.*, 2002; Hanaoka *et al.*, 2002;
Yoshimoto *et al.*, 2004), and hypersensitivity to different stress conditions including oxidative
stresses (Xiong *et al.*, 2007*b*; Liu *et al.*, 2009; Zhou *et al.*, 2013), indicating a significant redox
imbalance. On the other hand, oxidative treatments with H_2O_2 and methyl viologen (MV) have
been shown to induce autophagy (Xiong *et al.*, 2007*b*; Pérez-Pérez *et al.*, 2012*b*). Likewise,
autophagosome formation triggered by nutritional starvation and saline stress in Arabidopsis
roots depends on the NADPH-oxidase complex, a major source of ROS in the apoplast, which is
essential for local and systemic redox signal transduction (Liu *et al.*, 2009). However, the
induction of autophagy under osmotic stress was shown to be NADPH oxidase-independent,
suggesting that different signaling pathways regulate autophagy induction, and in certain
scenarios are dependent on ROS signaling (Liu *et al.*, 2009). More recently, it has been
hypothesized that H_2O_2 accumulation and membrane damage may mediate chlorophagy
induced by UV and high light treatments in Arabidopsis leaves (Izumi *et al.*, 2017; Woodson,
2019). Consequently, it is plausible that autophagy induction occurs as a response to the
accumulation of cellular components that have been damaged. Notably, recent studies
conducted in yeast, animals and Chlamydomonas have revealed that certain proteins involved
in the autophagic process could be inhibited by ROS in *in vitro* experiments or artificial oxidative
stress *in vivo*, indicating that ROS may inhibit the function of some autophagy-related proteins
(Pérez-Pérez *et al.*, 2014, 2016; Frudd *et al.*, 2018). These findings raise apparently opposite
evidence regarding the redox regulation of autophagy. Hence, the picture about the relationship
among different ROS, subcellular redox changes, and autophagy regulation in plants is still
unclear. Under this scenario, our study aimed to address the effects of different ROS on
autophagy in plants.
Herein, we provide evidence that oxidative treatments with H_2O_2 and MV, which
drastically increased the oxidized state of the cytoplasm, led to a reduction in the autophagic
flux in Arabidopsis seedlings. Likewise, after the oxidative treatments and during the subsequent

recovery period, the cytoplasm returned to its reduced state while autophagy was markedly induced. On the other hand, pharmacological and genetic approaches widely used to enhance the generation of $^1O_2$ resulted in an increased autophagic flux in Arabidopsis seedlings, with no significant effects observed on the cytoplasmic redox state. In summary, our findings unveil the distinct effects of ROS on autophagic flux, highlighting a strong correlation between autophagy and the physicochemical properties of the cytoplasm, with autophagy inhibition occurring under conditions where the redox state of the cytoplasm is highly oxidized.

MATERIALS AND METODS

Plant materials and growth conditions.

Arabidopsis thaliana ecotype Columbia (Col-0) was used in this study. 35S:GFP-ATG8a transgenic seeds (CS39996) (Thompson et al., 2005) were obtained from the Arabidopsis Biological Resource Center. The cytosol-targeted fluorescent redox sensors GFP2-Orp1 or GRX-roGFP2 were provided by Dr. Markus Schwarzländer. Fluorescent (*flu*) and ferrochelatase 2 (*fc2*) mutants were provided by Dr. Jesse Woodson. We introgressed the GFP-ATG8a, GFP2-Orp1 and GRX-roGFP2 transgenes into the *flu* and *fc2* mutant lines by genetic crosses. Homozygous offspring were then selected from the F3 backcross generations.

Seeds were surface-sterilized and sown on culture plates containing nutrient-rich Murashige and Skoog (MS) solid medium lacking sucrose and containing 1 % agar. After sowing, plates were kept in darkness at 4°C for 2 to 3 days, and then transferred to a growth chamber under a 16 h light/8 h dark photoperiod. Photosynthetic photon flux density was 110 µmol m^{-2} sec^{-1}. Average temperature was 22°C. Plants were grown for 7 days under these conditions and then passed to MS liquid medium to carry out the different treatments. For salt and osmotic stresses, NaCl (50 mM) and mannitol (100 mM) were used, respectively. Oxidative treatments were performed by the supplementation of the MS liquid medium with 1 µM methyl viologen (MV), 10 mM H_2O_2, 20 µM 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU), 20 µM methylene blue (MB). To scavenge superoxide and peroxide, 10 mM tiron and 10 mM benzoic acid (BA) were co-administrated, respectively. 10 mM dithiothreitol (DTT) was used as a reducing agent for protein disulfide bonds.

Confocal fluorescence microscopy and image analysis.

Confocal microscopy was performed on an Inverted microscope Nikon Eclipse CZ1. 35S:GFP-ATG8a expressing plants were observed with a 488-nm argon laser (BP 495–530 nm). To analyze autophagic bodies accumulation, 1 µM Concanamycin A was added to the different treatments. The number of autophagic bodies in epidermal cells was quantified after 16 h of treatments using ImageJ by counting their numbers in the central vacuole of multiple cells in sections of 50000 µm^2. A macro for automating autophagosome counting was developed in ImageJ. Briefly, the input images were converted into an 8-bit format, followed by background subtraction using a rolling ball algorithm with a 50-pixel radius. Subsequently, particle analysis was performed on the binary image, considering particle size, circularity, and displaying the resulting masks.

Cytosol-targeted fluorescent redox sensors GFP2-Orp1 or GRX-roGFP2 were excited at 488 nm and 405 nm and fluorescent emissions were collected at 505-535 nm. All imaging parameters were kept constant throughout all measurements. Subsequently, a single ratiometric image was generated using the ImageJ software, where the value of each pixel is the product of the ratio between the intensity at 405 nm with respect to that obtained at 488 nm according to (Schwarzlander *et al.*, 2008). Quantitative measurements were calculated as the ratio of the mean intensity from each channel. The dynamic range of these reporter lines was established

234 using 10 mM DTT and a curve of increasing concentrations of H_2O_2. For pseudocolor display, the
235 ratio was coded on a spectral color scale ranging from blue (fully reduced) to red (fully oxidized).
236

Immunoblot analyses.
For GFP-ATG8 cleavage assay, seven-day old 35S:GFP-ATG8a transgenic plants were transferred
to MS liquid medium to carry out the different treatments for 6 h. Then, total proteins were
extracted. The extracts were centrifuged at 10 000 g for 10 min, and equal amounts of proteins
from the supernatants were subjected to SDS-PAGE and immunoblotted with anti-GFP
antibodies (Roche). Comparisons of absolute integrated density values for each line (GFP-ATG8
and free GFP) were made. Corresponding background values were subtracted from integrated
density values, then integrated density values for bands corresponding to free GFP were
expressed as relative values of GFP-ATG8 present in the same sample, and relativized to a
control condition.

Physiological parameters.
Seven-day old Arabidopsis seedlings were exposed to the different treatments for 8 h and then
the production of H_2O_2 and O_2^-, quantum efficiency of PSII photochemistry (ΦPSII), as well as the
content of chlorophyll, soluble sugars and malondialdehyde (MDA) were analyzed. To assess
H_2O_2 and O_2^- accumulation, leaves were stained with DAB and NBT, respectively. The staining
was quantified by measuring the intensity per area using ImageJ. Quantum efficiency of PSII
photochemistry (ΦPSII) was measured from the upper surface using a pulse amplitude
modulated fluorometer (FMS2, Hansatech Instruments, Pentney King's Lynn, UK).
Measurements were performed 6 h after the growth chamber lights were turned on. For
chlorophyll, soluble sugars, and MDA analysis an alcoholic extract (1:2 w/v) was obtained from
the samples. After centrifugation for 15 minutes at 12,000 g, the supernatant was diluted in
ethanol, and the absorbance was measured at 654 nm using a spectrophotometer. Total
chlorophylls were calculated following the method reported by (Wintermans M. and De Mots
A., 1965). For soluble sugars determination, an aliquot of the sample was dried, and
resuspended in a same volume of water. Then, the concentration of sugars was measured with
anthrone (Fales, 1951) evaluating the absorbance at 620 nm using a glucose standard curve. For
MDA determination, a volume of the supernatant was mixed with a same volume of a mixture
of 0,5 % TBA and 20 % TCA, and incubated 20 minutes at 90 °C. The mixture was centrifuged at
12,000 g for 5 minutes, and the absorbance was measured at 532 and 600 nm (Heath and Packer,
1968).

RNA extraction and RT-qPCR
Samples were homogenized in a cold mortar with TRIzol Reagent, mixed for 1 min and incubated
at room temperature for 5 min. Then, 0.2 ml chloroform per ml of TRIzol Reagent was added and
incubated at room temperature for 3 min. After incubation, the samples were centrifuged at 12
000 g at 4 °C for 15 min and the aqueous phases were transferred to clean tubes. RNA was
precipitated by adding 1 vol of isopropanol, incubated at room temperature for 10 min and
centrifuged at 12 000 g at 4 °C for 15 min. The precipitate was washed with 70 % ethanol and
the samples were centrifuged again at 12 000 g at 4 °C for 15 min. The precipitates were dried
and resuspended in diethyl pyrocarbonate-treated water and its concentration was quantified
using a NanoDrop 3300 spectrometer (Thermo Scientific). Purified RNA was treated with DNase
I (Invitrogen) to remove genomic DNA, according to the manufacturer's instructions. DNA-free
RNA (1 to 2.5 μg) was used with oligo(dT) for first strand cDNA synthesis using the Moloney
murine leukemia virus reverse transcriptase (M-MLV RT, Promega) according to the
manufacturer's instructions. For each primer pair, the presence of a unique product of the

expected size was checked on ethidium bromide-stained agarose gels after PCR reactions.
Absence of contaminant genomic DNA was confirmed in reactions with DNase-treated RNA as
template. The qPCR reaction was performed using iQ Universal SYBR Green Supermix (Bio-Rad).
Amplification of actin with the forward primer 5'-TTCAGATGCCCAGAAGTCTTG-3' and the
reverse primer 5-GCGATACCTGAGAACATAGTGG-3 was used to normalize the amount of
template cDNA. The specific primers pair employed for the detection of *Atg6* was forward primer
5'-AAAACTGGGAGAGCGATGAG-3' and reverse primer 5'-GATAGATGAGCCCTGCGTAG-3', for
Atg4a was 5'-CCTTTGGAATCACTCGACCCATCT-3' and 5'-ATCCGCAAACCCGTAGTTGCTT-3', for
Atg7 was 5'-GCGGCTTTAGGTTTTGACAG-3' and 5'-GTTGGTCTAGAGTTCGATCAGTC -3', and for
Atg5 was 5'-TCCTGTTCGGTTGTATGTTCG-3' and 5'-CGCTCTGTCTCCCATAAACTC-3'. qPCR was
performed in thermocycler iQ5 (Bio-Rad) at 59 °C with iQ SYBR Green Supermix (Bio-Rad),
according to the manufacturer's instruction. Relative transcript abundance was calculated (Livak
and Schmittgen, 2001).
296
Microarray analysis.
Publicly available microarray data were utilized to compare the expression of *Atg* genes under
various conditions, including nitrogen deficiency (Rubin *et al.*, 2009), treatment with 10 mM
H_2O_2 for 24 h (Willems *et al.*, 2016), treatment with 10 µM MV for 2 h (Scarpeci *et al.*, 2008),
and the examination of *flu* mutant plants subjected to an 8-hour cycle of darkness followed by
2 h of light (Op Den Camp *et al.*, 2003). Expression values were normalized to the corresponding
control conditions employed in each study and were expressed as $\log_2$ values.
304
305 **Statistical analyses.**
306 Data were analysed by analysis of variance (ANOVA) followed by the Di Rienzo–Guzman–
307 Casanoves (DGC) test model, a cluster-based method for identifying groups of non-
308 homogeneous means, using InfoStat software (Di Rienzo *et al.*, 2002).
309
RESULTS
H_2O_2 and methyl viologen treatments oxidized the cytoplasmic redox state and inhibited
autophagy in Arabidopsis.
Initially, to gain insights into the interplay between ROS, redox state changes and
autophagy, *Arabidopsis thaliana* seedlings were subjected to well-established autophagy-
inducing conditions, namely nutritional starvation and different abiotic stress treatments (Wang
et al., 2021), and both autophagy and the cytoplasmic redox state were analyzed (Fig. 1). To
monitor autophagy, 35S:GFP-ATG8a (GFP-ATG8) transgenic plants were grown 7 days in MS solid
medium, and then transferred to the different treatments in MS liquid medium supplemented
with 1 µM Concanamycin A (Conc A) for a 16 hour-period. Conc A is a V-ATPase inhibitor widely
used to prevent vacuole acidification, thereby effectively inhibiting the lytic activity of vacuoles,
thus allowing the detection of autophagic bodies accumulation into the vacuoles through
confocal microscopy (Fig. 1A and B). Analysis of autophagic flux using Conc A better defines
autophagic activity since the increase of autophagosome number in the cytoplasm does not
necessarily corresponds to an enhanced autophagic activity (Robert *et al.*, 2021). When
subjected to carbon and nitrogen starvation treatments, epidermal cells showed a high
accumulation of autophagic bodies in the vacuole compared to control conditions, indicating an
activated autophagic flux (Fig. 1A and B). Similarly, saline and osmotic stress treatments resulted
in comparable outcomes, with a significant increase in autophagic bodies accumulating within
the vacuoles (Fig. 1A and B). Besides, to evaluate the cytoplasmic redox state *in vivo*, 7 days old
transgenic plants expressing the cytosol-targeted fluorescent redox sensors GFP2-Orp1 or GRX-

roGFP2 were subjected to the different treatments (Fig. 1). These transgenic plants express
redox-sensitive green fluorescent proteins (roGFPs) genetically fused to different redox-active
proteins sensing H_2O_2 production and GSH:GSSG, respectively (Schwarzländer *et al.*, 2009, 2015;
Aller *et al.*, 2013; Nietzel *et al.*, 2019). The relative intensity of the two fluorescence excitation
maxima depends on the redox state of the modified GFP, enabling ratiometric measurements,
independent of the probe concentration (Schwarzländer *et al.*, 2016). The dynamic range of
these reporter lines was previously monitored using DTT and H_2O_2 (Fig. S1). Notably, in
epidermal cells of seedlings expressing either GFP2-Orp1 or GRX-roGFP2, no changes in the
cytoplasmic redox state were detected after 16 h of nutritional starvation, neither under saline
or osmotic stress treatments compared to control conditions (Fig. 1A and C). These findings align
with earlier assessments of the redox state in cells of GRX-roGFP2 and roGFP2-ORP1 plants
under different stress conditions (Schwarzländer *et al.*, 2009).
To further investigate the interplay between ROS, redox state changes and autophagy,
oxidative stress treatments using H_2O_2 and methyl viologen (MV) were conducted. We evaluated
different concentrations of H_2O_2 and observed a dose-dependent response for both the
oxidation of the cytoplasmic redox state and autophagy inhibition (Fig. S1). A similar approach
was taken with MV, an herbicide widely used for the induction of photooxidative stress as it
takes electrons from photosystem I, generating O_2^- and H_2O_2 inside chloroplasts in the presence
of light (Mehler, 1951; Farrington *et al.*, 1973; Lascano *et al.*, 2012; Nazish *et al.*, 2022). Based
on these results and with the intention of employing moderate oxidative stress conditions that
do not induce cell death, we used 10 mM H_2O_2 and 1 µM MV for subsequent experiments.
Although oxidative treatments have been reported as autophagy-inducing conditions in
Arabidopsis and Chlamydomonas (Xiong *et al.*, 2007*b*; Pérez-Pérez *et al.*, 2012*b*), no significant
differences were observed in the accumulation of autophagic bodies in the epidermal cells of
seedlings treated with MV or H_2O_2 compared to the control condition (Fig. 1A and B). In contrast
to observations under autophagy-inducing conditions such as nutritional starvation, saline, or
osmotic stresses, these oxidative treatments significantly oxidized the cytoplasmic redox state,
as indicated by the reduced GSSG:GSH ratio in the cytoplasm (GRX-roGFP2) and increased the
accumulation of H_2O_2, as revealed by analyzing roGFP2-ORP1 plants and DAB staining (Fig. 1A
and C; Fig. 2; Fig. S1; Fig. S2).
To further characterize the effects of oxidative stress on autophagy, treatments with MV
and H_2O_2 were performed on 35S:GFP-ATG8a plants under autophagic inducing conditions,
specifically nitrogen and carbon starvation (Fig. 2). Interestingly, treatments with MV and H_2O_2
reduced the accumulation of autophagic bodies after 16 h of starvation conditions (Fig. 2A and
B). Furthermore, the inhibitory effects of the oxidative treatments on the autophagic flux were
also confirmed by analyzing the degradation of the GFP-ATG8a fusion protein by western blot
(Fig. 2D). The evaluation of autophagic flux through the analysis of the degradation of the GFP-
ATG8 fusion protein by immunoblotting allowed us to examine the effects of oxidative stress on
autophagy in shorter time periods (6 h, Fig. 2D).. Next, tiron and benzoic acid were co-
administered with the MV treatment to scavenge superoxide and hydrogen peroxide
(Navabpour *et al.*, 2003; Mansouri *et al.*, 2005; Velika and Kron, 2012; Liubimovskii *et al.*, 2021).
The effects of MV on autophagic flux, H_2O_2 accumulation, and cytoplasmic redox state oxidation
were blocked in the presence of ROS scavengers (Fig. 2, Fig. S2). Similarly, photooxidative
treatments induced by high light treatments showed an inhibitory effect on the autophagy flux,
which was reversed in the presence of the superoxide radical scavenger, tiron (Fig. 2D).
Moreover, both tiron and benzoic acid prevented photosynthetic damage caused by MV
treatments, as measured by the quantum yield of photosystem II (ΦPSII), and prevented the
decrease in soluble sugar levels (Fig. S2). Furthermore, the inhibitory effects of MV on

autophagic flux and the oxidation of the GSH:GSSG sensor probe (GRX-roGFP2) were reversed when 10 mM dithiothreitol (DTT), a reducing agent for protein disulfide bonds, was added (Fig. S2). These results emphasize that the effects of MV in plants are associated with its capacity to induce ROS generation, contributing to a more oxidized cellular redox state and oxidative damage (Lascano *et al.*, 2012). Notably, the restoration of autophagic flux in MV-treated plants by tiron, benzoic acid, and DTT suggests that H_2O_2 may inhibit autophagy, potentially by oxidizing disulfide bonds in crucial ATG proteins (Fig. S3 and S4) as demonstrated in recent studies in yeast, Chlamydomonas, and animals (Pérez-Pérez *et al.*, 2014, 2016; Frudd *et al.*, 2018; Zheng *et al.*, 2020). In this regard, the inhibitory effect of H_2O_2 treatments on autophagy was also observed in plants exposed to carbon or nitrogen starvation (Fig. 2A, B and D; Fig. S1).

Autophagy is induced in the recovery phase after oxidative stress in Arabidopsis.

Previous reports indicated that autophagy is involved in degrading oxidized proteins and organelles in response to oxidative stress (Xiong *et al.*, 2007*b*; Shibata *et al.*, 2013; Izumi *et al.*, 2017; Oikawa *et al.*, 2022). However, our experimental results demonstrate that oxidative stress treatments inhibit autophagy, as illustrated in Figure 2. In light of these findings, we aimed to investigate autophagic activity following oxidative stress treatments by assessing the autophagic flux in Arabidopsis plants during the recovery phase. With this purpose, 7-day-old Arabidopsis seedlings were subjected to a brief exposure of 10 mM H_2O_2 or high light intensity ($\approx$1000 μmoles m^{-2} s^{-1}) for 4 hours, followed by transfer to H_2O_2-free MS liquid medium under control conditions ($\approx$100 μmoles m^{-2} s^{-1}) for a 20-hour recovery period. Additionally, a set of plants were incubated in 10 mM H_2O_2 or to control conditions (H_2O_2-free MS liquid medium) for 24 h. This experimental design aimed to elucidate the dynamic changes in autophagy following oxidative stress and determine whether autophagic activity is modulated during the recovery phase. The results showed that autophagic flux was inhibited in plants incubated 24 h with H_2O_2 compared to control conditions (Fig. 3A, B and D). However, autophagic flux was not only recovered but also induced during the recovery period after the stress treatments surpassing the levels observed in plants subjected to control conditions (Fig. 3A, B and D). Likewise, after the 20-hour recovery period, the redox biosensors returned to their reduced state, indicating both a successful cytoplasmic redox recovery, and a well-defined correlation between reduced cytoplasmic redox state and autophagic flux (Fig. 3A and C).

Singlet oxygen induced autophagic flux but did not affect the cytoplasmic redox state.

Singlet oxygen ($^1O_2$) is mainly produced in the chloroplast, and its targets are predominantly confined inside this organelle due to the specific physicochemical characteristic of this ROS (Dmitrieva *et al.*, 2020). To investigate the effects of $^1O_2$ on the autophagic activity, GFP-ATG8a seedlings were treated with 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU) (Fig. 4), an herbicide that specifically induces $^1O_2$ generation by blocking plastoquinone reduction. This process promotes $^1O_2$ generation without affecting superoxide anion and hydroxyl radicals at PSII (Fufezan *et al.*, 2002; Flors *et al.*, 2006; Dmitrieva *et al.*, 2020). The analysis of autophagic body accumulation and cytoplasmic redox state was conducted after a 16-hour treatment (Fig. 4A-C). Additionally, treatments with methylene blue (MB), a type-II photosensitizer able to generate $^1O_2$ in leaves under visible light (Shao *et al.*, 2007, 2013), with minor effects on photosynthesis (Fig. 4E), were also performed. Treatments with these chemicals increased the accumulation of autophagic bodies but did not show statistically significant effects on the redox state of the cytoplasm and H_2O_2 production in Arabidopsis cells compared to control treatment (Fig. 4A and B). The autophagic flux-inducing effect of DCMU and MB was also evidenced after

6 hours of treatments through the analysis of GFP-ATG8 degradation by immunoblotting (Fig.
4D).

To further investigate the effects of $^1O_2$ on the autophagic process and the cytoplasmic
redox state, autophagy reporter- and redox sensor-transgenic plants were crossed with widely
used endogenously chloroplast $^1O_2$-hyperproducing *Arabidopsis* mutant plants: fluorescent (*flu*)
and two different lines of ferrochelatase 2 (*fc2*): *fc2.1* which presents reduced expression levels
of *Fc2*, and the null mutant *fc2.2*. These mutant plants accumulate different photosensitizing
intermediates of chlorophyll biosynthesis in the dark, which, upon light exposure lead to an
exacerbated production of $^1O_2$ (Op Den Camp *et al.*, 2003). The *flu* mutant plants accumulate
protochlorophyllide (Meskauskiene *et al.*, 2001), while mutations in plastid ferrochelatase 2 lead
to the accumulation of protoporphyrin IX (Scharfenberg *et al.*, 2015). Initially, wild type, *flu* and
fc2 mutant plants were grown 7 days under continuous light conditions (control conditions), and
several growth and physiological parameters were evaluated (Fig. S5). Interestingly, the *fc2*
mutants showed arrested root growth, reduced chlorophyll levels, increased levels of
malondialdehyde (MDA) an intermediary metabolite of lipid peroxidation used as oxidative
stress marker (Robert *et al.*, 2014), and reduced levels of soluble sugars compared to wild type,
while no significant differences were observed between wild type and *flu* mutant (Fig. S5).
Likewise, the cytoplasmic redox state in *fc2* plants was significantly more oxidized than that in
the wild type and *flu* backgrounds (Fig. S5, Fig. 5). Furthermore, and consistent with a previous
study (Lemke *et al.*, 2021), the autophagic flux was induced in *fc2* mutants plants compared to
wild type and *flu* mutant under continuous light conditions (Fig. S5). Hence, considering the
altered phenotypes observed in *fc2* mutants even under control conditions, we used the *flu*
mutants for subsequent analysis.

flu/GFP-ATG8a plants were grown under continuous light conditions for 7 days,
transferred to darkness for 8 h, and then re-exposed to light in the presence of Conc A for
another 8 h. After the light/dark/light cycling conditions, a significantly increased autophagic
flux was observed in *flu*/GFP-ATG8a compared to Col-0/GFP-ATG8 plants. However, no
statistically significant differences were observed among these genotypes under continuous
light conditions (Fig. 5A, B and D). Besides, the accumulation of H_2O_2 was monitored under
control conditions as well as after a period of darkness by analyzing the fluorescent redox sensor
roGFP2-ORP1 expressed in the wild type and *flu* backgrounds. Although a slight tendency
towards a more oxidized state was observed in the cytoplasm of *flu* mutant plants once re-
exposed to light, the differences were not statistically significant (Fig. 5C). Thus, in contrast to
H_2O_2 and MV treatments, $^1O_2$ increased the autophagic flux and did not oxidize the cytoplasmic
redox state, indicating differential effects on the autophagic activity depending on the ROS
nature and their effects on the cytoplasmic redox state. Moreover, the transcript levels of *Atg4*,
Atg5, *Atg6* and *Atg7* were analyzed by RTq-PCR in *flu* mutant after 8 h of light re-exposure. No
statistically significant differences were observed in the expression level of the analyzed genes
between wild type and *flu* mutant neither under control conditions nor after cycling conditions
(Fig. S6).

DISCUSSION

Plants, as aerobic and sessile organisms, respond to environmental stresses by
increasing the production of reactive oxygen species (ROS), which, due to their high reactivity,
fulfill dual roles by acting both as agents of damage and as signaling molecules. Consequently,
ROS are universally implicated in plant stress symptomatology, and also play a pivotal role in
modulating plant growth, development, and cell death (Apel and Hirt, 2004; Trippi *et al.*, 2009;
Baxter *et al.*, 2014; Choudhury *et al.*, 2017; Fichman and Mittler, 2020). Similarly, a multitude of

stress conditions in plants have been demonstrated to initiate or enhance autophagy (Bassham *et al.*, 2006; Liu *et al.*, 2009; Signorelli *et al.*, 2019; Su *et al.*, 2020). Moreover, defects in antioxidant proteins in Arabidopsis and Chlamydomonas contribute to uncontrolled ROS bursts and autophagy induction (Pérez-Pérez *et al.*, 2012*a*; Shibata *et al.*, 2013). While the concurrent induction of ROS production and autophagy activation suggests a positive correlation between these cellular processes, certain molecular mechanisms have been reported in animal cells, yeast, and Chlamydomonas, indicating that ROS could potentially exert a negative influence on autophagy (Pérez-Pérez *et al.*, 2014, 2016; Frudd *et al.*, 2018). However, direct evidence of ROS-mediated inhibition of autophagy remains scarce in plants. One of the fundamental unanswered queries is whether a particular type of ROS functions as the primary messenger/modulator, if any, to regulate autophagy. Our findings point in that direction, highlighting the differential effects of ROS on autophagy and suggesting that, under specific stress conditions leading to a highly oxidized cytoplasmic state, the autophagic process may be suppressed.

In this study, MV was used to specifically generate O_2^- and H_2O_2 in the chloroplast (*revised in* Asada, 2006; Lascano *et al.*, 2012). Previous studies have reported the formation of autophagosomes in root cells following incubation with MV (Xiong *et al.*, 2007*b*). Nevertheless, our findings differ from these outcomes. It is important to acknowledge that certain variations in experimental and methodological approaches between our research and the prior studies may have a considerable impact on the interpretation of the outcomes. In the earlier studies, MV treatments were administered on solid MS media, subjecting plants to 10 µM MV for a duration of two days, and the observations were made in roots (heterotrophic tissues where MV produces O_2^- and H_2O_2 in mitochondria). Conversely, in our study, MS liquid media was supplemented with 1 µM MV, and leaves were analyzed (tissues where MV generates O_2^- and H_2O_2 in chloroplast). Furthermore, a critical distinction lies in the method employed for quantifying autophagosomes. Previous investigations employed monodansylcadaverine (MDC) staining to identify acidic compartments. However, experiments involving autophagy-deficient *atg* mutants, as well as co-localization analysis in GFP-ATG8 plants, indicated that the MDC-stained structures do not clearly manifest as autophagosomes and/or autolysosomes (Merkulova et al., 2014). In contrast, we used GFP-ATG8 plants for autophagy analysis through confocal microscopy and western blot analysis, which are more reliable and widely accepted method for identifying autophagosomes/autophagic bodies and monitoring autophagic activity in general.

The relationship between autophagy and ROS is intricate, and our findings revealed that different types of ROS, even when generated in the same organelle, can have different effects on the autophagic process. We observed that under conditions of nutritional starvation, which are canonical inducers of autophagy, oxidative treatments with MV and H_2O_2 had an inhibitory effect on the autophagic flux (Fig. 2). This implies that the comparatively limited autophagosome biogenesis rate under optimal growth conditions could potentially account for the absence of observable inhibitory effects of ROS on autophagy in prior *in vivo* plant studies (Fig. 1). Opposite to the inhibitory effects of H_2O_2 on the autophagic flux, the specific generation of $^1O_2$ in the *flu* mutant plants, as well as treatments with DCMU and MB (Fufezan *et al.*, 2002; Op Den Camp *et al.*, 2003; Flors *et al.*, 2006; Shao *et al.*, 2007, 2013; Dmitrieva *et al.*, 2020), increased the autophagic activity (Fig. 4 and Fig. 5). These findings are consistent with previous research demonstrating that treatments with norflurazon, an inhibitor of carotenoid biosynthesis, triggers autophagy in *Chlamydomonas reinhardtii* (Pérez-Pérez *et al.*, 2016). Similarly to what is observed in *flu* mutants (Op Den Camp *et al.*, 2003), norflurazon treatments lead to perturbations in plastid homeostasis that increase the photosensitizing activity of tetrapyrroles, which can transfer the excitation energy to the oxygen molecule, thereby generating $^1O_2$ (Kim

and Apel, 2013). Singlet oxygen, carotenoid derivatives, and tetrapyrroles are principal
components of the retrograde signaling pathway, a key process that regulates not only
chloroplastic processes but also intersects with other cellular signaling events playing broader
roles in plant function (Chan *et al.*, 2016).

With a shorter half-life and inability to cross membranes, the activity of $^1O_2$ is confined
to the thylakoid membrane of chloroplasts (Dmitrieva *et al.*, 2020). These characteristics provide
potential explanations for the absence of observable effects on the cytoplasmic redox state in
treatments involving DCMU, MB, and *flu* mutant plants (Fig. 4 and Fig. 5). Conversely, treatments
with MV and H_2O_2 highly oxidized the cytoplasmic redox state (Fig. 1 and Fig. 2). These results
establish a well-defined correlation between the cytoplasmic redox state of the cytoplasm and
autophagic activity.

The observation that the inhibition of autophagic flux and the oxidation of the
cytoplasmic redox state in MV treatments can be prevented by ROS scavengers such as tiron or
benzoic acid, as well as by treatment with the disulfide bonds reductant DTT, strengthens the
notion that H_2O_2 inhibits autophagy, but also suggests that protein oxidation in the cytoplasm
may be involved in this response. Notably, it has been proposed that proteins are the primary
targets of reactive oxidants and other radicals *in vivo* (Davies, 2005). At the molecular level, the
activity of certain ATG proteins has been shown to be regulated by redox changes. For instance,
ATG4, a protein involved in the cleavage of ATG8 for its lipidation and de-lipidation, is inhibited
by H_2O_2 in yeast and the green alga *Chlamydomonas reinhardtii* (Pérez-Pérez *et al.*, 2014, 2016).
Conversely, reducing agents such as DTT can restore the activity of ATG4 (Pérez-Pérez *et al.*,
2014, 2016). Furthermore, it has been reported in animal cells that oxidation of ATG3 and ATG7
leads to the inhibition of autophagy (Frudd *et al.*, 2018). The conservation of amino acid
sequence between the isoforms from yeast, human cells, and Arabidopsis (Fig. S3 and S4)
suggest that this type of regulation could also occur in plants.

The complexity of the relationship between autophagy and ROS becomes even more
complex when considering that this catabolic process is involved in the degradation of proteins
and oxidized organelles, which are primary sources of ROS production. The increased autophagic
flux observed during the recovery period after exposure to oxidative stress, accompanied by the
restoration of a reduced redox state in the cytoplasm, indicated that the inhibition of autophagy
by ROS is transient (Fig. 3). Through the assessment of autophagic flux, our objective was to
illuminate the role of autophagy in cellular adaptation to oxidative stress and its potential
involvement in stress recovery mechanisms. Our findings imply that inducing autophagy once
the oxidative stress has subsided is linked to the intracellular clearance of cytoplasmic
components damaged during oxidative stress. This is supported by experiments with the
antioxidants tiron and benzoic acid, which regulated ROS production and thwarted cell injury
(Fig. S2). These observations underscore the dynamic nature of autophagy in response to
oxidative stress and its subsequent recovery, pointing towards a complex regulatory mechanism
in Arabidopsis.

In summary, our work provides a novel perspective on the relationship between ROS
and autophagy. The fact that the generation of H_2O_2 and $^1O_2$ have different effects on autophagy
imply that the term "ROS" encompasses molecules with the ability to modulate this process in
distinct ways. Further studies are needed to unravel the molecular mechanisms underlying the
differential modulation of autophagy by ROS, being ATG4, ATG3 and ATG7 possible targets to
be oxidatively regulated.

567
568
569

SUPPLEMENTARY DATA

Supplementary Figure 1: Optimization of methyl viologen (MV) and H_2O_2 concentrations, and calibration of the redox sensors. Supplementary Figure 2: DTT, tiron and benzoic acid reversed the effects of MV treatments.

Supplementary Figure 3: Sequence alignment of ATG4 proteases from different organisms.

Supplementary Figure 4: Sequence alignment of ATG3 and ATG7 proteins from different organisms.

Supplementary Figure 5: Characterization of *fc2* and *flu* mutant plants under continuous light conditions.

Supplementary Figure 6: Transcript level of *Atg* genes in Arabidopsis plants subjected to different oxidative treatments.

ACKNOWLEDGEMENTS

We thank Dr. Leandro Ortega and Dr. Virginia Lobatto (Unidad de Doble Dependencia INTA-CONICET (UDEA)) for helpful support with the confocal microscopy and for technical support, respectively. We also thank Sistema Nacional de Microscopia of the Ministerio de Ciencia, tecnología e innovación Productiva (MINCyT), Argentina.

AUTHOR CONTRIBUTION:

GR and RL: conceptualization; GR and AE: investigation; GR, AE, LS and RL: formal analysis; GR, LS, and RL: resources; GR and AE: writing original draft and incorporated feedback from all authors; GR and AE: visualization; GR and RL: supervision; GR, LS and RL: funding acquisition.

CONFLICT OF INTEREST

No conflict of interest declared.

FUNDING

This work was supported by grants from Agencia Nacional de Promoción Científica y Tecnológica, Argentina: FONCYT-PICT-2019-0533 (RL), FONCyT PICT-2017-2863 (GR), and FONCyT PICT-2020-2482 (LS); Consejo Nacional de Investigaciones Científicas y Técnicas CONICET-PIP2015-PIP2019 (RL), Proyecto INTA 2019-PD-E6-I116-001 (RL and GR), and Proyectos Consolidar, Secretaría de Ciencia y Técnica de la Universidad Nacional de Córdoba (UNC) (RL).

REFERENCES

Agbemafle W, Wong MM, Bassham DC. 2023. Transcriptional and post-translational regulation of plant autophagy. Journal of Experimental Botany **74**, 6006–6022.

Aller I, Rouhier N, Meyer AJ. 2013. Development of roGFP2-derived redox probes for measurement of the glutathione redox potential in the cytosol of severely glutathione-deficient rml1 seedlings. Frontiers in Plant Science **4**, 1–12.

Apel K, Hirt H. 2004. Reactive oxygen species: metabolism, oxidative stress, and signal transduction. Annual review of plant biology **55**, 373–99.

Asada K. 2006. Production and Scavenging of Reactive Oxygen Species in Chloroplasts and Their Functions. Plant Physiology **141**, 391–396.

Bassham DC, Laporte M, Marty F, Moriyasu Y, Ohsumi Y, Olsen LJ, Yoshimoto K. 2006. Autophagy in Development and Stress Responses of Plants. Autophagy **2**, 2–11.

Baxter A, Mittler R, Suzuki N. 2014. ROS as key players in plant stress signalling. Journal of experimental botany **65**, 1229–1240.

Bechtold U, Richard O, Zamboni A, Gapper C, Geisler M, Pogson B, Karpinski S, Mullineaux

PM. 2008. Impact of chloroplastic- and extracellular-sourced ROS on high light-responsive gene
expression in Arabidopsis. Journal of experimental botany **59**, 121–33.

Breeze E, Harrison E, McHattie S, et al. 2011. High-resolution temporal profiling of transcripts
during Arabidopsis leaf senescence reveals a distinct chronology of processes and regulation.
Plant Cell **23**, 873–894.

Breusegem F Van, Dat JF. 2006. Reactive Oxygen Species in Plant Cell Death. Plant physiology
141, 384–390.

Breusegem F Van, Vranova E, Dat JF, Inze D. 2001. The role of active oxygen species in plant
signal transduction. Plant Science **161**, 405–414.

Chan KX, Phua SY, Crisp P, McQuinn R, Pogson BJ. 2016. Learning the Languages of the
Chloroplast: Retrograde Signaling and beyond. Annual Review of Plant Biology **67**, 25–53.

Choudhury FK, Rivero RM, Blumwald E, Mittler R. 2017. Reactive oxygen species, abiotic
stress and stress combination. Plant Journal **90**, 856–867.

Davies MJ. 2005. The oxidative environment and protein damage. Biochimica et Biophysica
Acta - Proteins and Proteomics **1703**, 93–109.

Dmitrieva VA, Tyutereva E V., Voitsekhovskaja O V. 2020. Singlet oxygen in plants:
Generation, detection, and signaling roles. International Journal of Molecular Sciences **21**.

Doelling JH, Walker JM, Friedman EM, Thompson AR, Vierstra RD. 2002. The APG8/12-
activating enzyme APG7 is required for proper nutrient recycling and senescence in
Arabidopsis thaliana. Journal of Biological Chemistry **277**, 33105–33114.

Fales FW. 1951. The assimilation and degradation of carbohydrates by yeast cells. The Journal
of biological chemistry **193**, 113–124.

Farrington JA, Ebert M, Land EJ, Fletcher K. 1973. Bipyridylium quaternary salts and related
compounds. V. Pulse radiolysis studies of the reaction of paraquat radical with oxygen.
Implications for the mode of action of bipyridyl herbicides. Biochimica et Biophysica Acta **314**,
372–381.

Feng Y, Yao Z, Klionsky DJ. 2015. How to control self-digestion: transcriptional, post-
transcriptional, and post-translational regulation of autophagy. Trends in cell biology **25**, 354–
363.

Fernandez-göbel TF, Deanna R, Muñoz NB, Robert G, Asurmendi S, Lascano R. 2019. Redox
Systemic Signaling and Induced Tolerance Responses During Soybean – Bradyrhizobium
japonicum Interaction : Involvement of Nod Factor Receptor and Autoregulation of
Nodulation. Frontiers in Plant Science **10**, 1–15.

Fichman Y, Mittler R. 2020. Rapid systemic signaling during abiotic and biotic stresses: is the
ROS wave master of all trades? Plant Journal **102**, 887–896.

Flors C, Fryer MJ, Waring J, Reeder B, Bechtold U, Mullineaux PM, Nonell S, Wilson MT,
Baker NR. 2006. Imaging the production of singlet oxygen in vivo using a new fluorescent
sensor, Singlet Oxygen Sensor Green. Journal of experimental botany **57**, 1725–34.

Foyer CH, Hanke G. 2022. ROS production and signalling in chloroplasts: cornerstones and
evolving concepts. Plant Journal **111**, 642–661.

Foyer CH, Noctor G. 2009. Redox Regulation in Photosynthetic Organisms: signaling,
acclimation, and practical implications. Antioxidants & Redox Signaling **11**, 861–905.

Foyer CH, Noctor G. 2011*a*. Ascorbate and glutathione: the heart of the redox hub. Plant
physiology **155**, 2–18.

Foyer CH, Noctor G. 2011*b*. Ascorbate and glutathione: the heart of the redox hub. Plant
physiology **155**, 2–18.

Frudd K, Burgoyne T, Burgoyne JR. 2018. Oxidation of Atg3 and Atg7 mediates inhibition of
autophagy. Nature Communications **9**, 1–15.

Fufezan C, Rutherford AW, Krieger-liszkay A. 2002. Singlet oxygen production in herbicide-
treated photosystem II. FEBS **532**, 407–410.

Hanaoka H, Noda T, Shirano Y, Kato T, Hayashi H, Shibata D, Tabata S, Ohsumi Y. 2002. Leaf
senescence and starvation-induced chlorosis are accelerated by the disruption of an

Arabidopsis autophagy gene. Plant Physiol **129**, 1181–1193.

Heath RL, Packer L. 1968. Photoperoxidation in isolated chloroplasts. Archives of Biochemistry
and Biophysics **125**, 189–198.

Izumi M, Ishida H, Nakamura S, Hidema J. 2017. Entire Photodamaged Chloroplasts Are
Transported to the Central Vacuole by Autophagy. The Plant Cell **29**, 377.

Karpinski S, Reynolds H, Karpinska B, Wingsle G, Creissen G, Mullineaux P. 1999. Systemic
signaling and acclimation in response to excess excitation energy in Arabidopsis. Science **284**,
654–7.

Kim C, Apel K. 2013. 1O2-mediated and EXECUTER-dependent retrograde plastid-to-nucleus
signaling in norflurazon-treated seedlings of arabidopsis thaliana. Molecular Plant **6**, 1580–
1591.

Lascano R, Munoz N, Robert G, Rodriguez M, Melchiorre M, Trippi V, Quero G. 2012.
Paraquat: An Oxidative Stress Inducer. Herbicides - Properties, Synthesis and Control of
Weeds. London: IntechOpen, 135–148.

Lemke MD, Fisher KE, Kozlowska MA, Tano DW, Woodson JD. 2021. The core autophagy
machinery is not required for chloroplast singlet oxygen - mediated cell death in the
Arabidopsis thaliana plastid ferrochelatase two mutant. BMC plant biology **21**, 1–20.

Liu Y, Xiong Y, Bassham DC. 2009. Autophagy is required for tolerance of drought and salt
stress in plants. Autophagy **5**, 954–63.

Liubimovskii SO, Ustynyuk LY, Tikhonov AN. 2021. Superoxide radical scavenging by sodium
4,5-dihydroxybenzene-1,3-disulfonate dissolved in water: Experimental and quantum chemical
studies. Journal of Molecular Liquids **333**, 20–23.

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta CT) Method. Methods **25**, 402–408.

Mansouri A, Makris DP, Kefalas P. 2005. Determination of hydrogen peroxide scavenging
activity of cinnamic and benzoic acids employing a highly sensitive peroxyoxalate
chemiluminescence-based assay: Structure-activity relationships. Journal of Pharmaceutical
and Biomedical Analysis **39**, 22–26.

Masclaux-Daubresse C, Clement G, Anne P, Routaboul J-M, Guiboileau A, Soulay F, Shirasu K,
Yoshimoto K. 2014. Stitching together the Multiple Dimensions of Autophagy Using
Metabolomics and Transcriptomics Reveals Impacts on Metabolism, Development, and Plant
Responses to the Environment in Arabidopsis. The Plant Cell **26**, 1857–1877.

Mehler AH. 1951. Studies on reactions of illuminated chloroplasts. II. Stimulation and
inhibition of the reaction with molecular oxygen. Archives of Biochemistry and Biophysics **34**,
339–351.

Melchiorre M, Robert G, Trippi V, Racca R, Lascano HR. 2009. Superoxide dismutase and
glutathione reductase overexpression in wheat protoplast: photooxidative stress tolerance and
changes in cellular redox state. Plant Growth Regulation **57**, 57–68.

Meskauskiene R, Nater M, Goslings D, Kessler F, Op den Camp R, Apel K. 2001. FLU: A
negative regulator of chlorophyll biosynthesis in arabidopsis thaliana. Proceedings of the
National Academy of Sciences of the United States of America **98**, 12826–12831.

Miller G, Schlauch K, Tam R, Cortes D, Torres MA, Shulaev V, Dangl JL, Mittler R. 2009. The
plant NADPH oxidase RBOHD mediates rapid systemic signaling in response to diverse stimuli.
Science Signaling **2**.

Munch D, Rodriguez E, Bressendorff S, Park OK, Hofius D, Petersen M. 2014. Autophagy
deficiency leads to accumulation of ubiquitinated proteins, ER stress, and cell death in
Arabidopsis. Autophagy **10**, 1579–1587.

Muñoz N, Robert G, Melchiorre M, Racca R, Lascano R. 2012. Saline and osmotic stress
differentially affects apoplastic and intracellular reactive oxygen species production, curling
and death of root hair during Glycine max L.-Bradyrhizobium japonicum interaction.
Environmental and Experimental Botany **78**, 76–83.

Nakamura S, Hidema J, Sakamoto W, Ishida H, Izumi M. 2018. Selective Elimination of

Membrane-Damaged Chloroplasts via microautophagy. Plant Physiology **177**, 1007–1026.

Navabpour S, Morris K, Allen R, Harrison E, A-H-Mackerness S, Buchanan-Wollaston V. 2003. Expression of senescence-enhanced genes in response to oxidative stress. Journal of experimental botany **54**, 2285–92.

Nazish T, Huang YJ, Zhang J, Xia JQ, Alfatih A, Luo C, Cai XT, Xi J, Xu P, Xiang C Bin. 2022. Understanding paraquat resistance mechanisms in Arabidopsis thaliana to facilitate the development of paraquat-resistant crops. Plant Communications **3**, 1–11.

Nietzel T, Elsässer M, Ruberti C, *et al.* 2019. The fluorescent protein sensor roGFP2-Orp1 monitors in vivo H_2O_2 and thiol redox integration and elucidates intracellular H_2O_2 dynamics during elicitor-induced oxidative burst in Arabidopsis. New Phytologist **221**, 1649–1664.

Oikawa K, Goto-yamada S, Hayashi Y, *et al.* 2022. Pexophagy suppresses ROS-induced damage in leaf cells under high-intensity light.

Op Den Camp RGL, Przybyla D, Ochsenbein C, *et al.* 2003. Rapid Induction of Distinct Stress Responses after the Release of Singlet Oxygen in Arabidopsis. Plant Cell **15**, 2320–2332.

Peláez-vico MÁ, Fichman Y, Zandalinas SI, Van F, Karpiński SM, Mittler R. 2022. ROS and redox regulation of cell-to-cell and systemic signaling in plants during stress. Free Radical Biology and Medicine.

Pérez-Pérez ME, Couso I, Crespo JL. 2012*a*. Carotenoid deficiency triggers autophagy in the model green alga Chlamydomonas reinhardtii. Autophagy **8**, 376–388.

Pérez-Pérez ME, Lemaire SD, Crespo JL. 2012*b*. Reactive oxygen species and autophagy in plants and algae. Plant physiology **160**, 156–64.

Pérez-Pérez ME, Lemaire SD, Crespo JL. 2016. Control of Autophagy in *Chlamydomonas* Is Mediated through Redox-Dependent Inactivation of the ATG4 Protease. Plant Physiology **172**, 2219–2234.

Pérez-Pérez ME, Zaffagnini M, Marchand CH, Crespo JL, Lemaire SD. 2014. The yeast autophagy protease Atg4 is regulated by thioredoxin. Autophagy **10**, 1–12.

Qi H, Xia FN, Xiao S. 2021. Autophagy in plants: Physiological roles and post-translational regulation. Journal of Integrative Plant Biology **63**, 161–179.

Di Rienzo JA, Guzmán AW, Casanoves F. 2002. A multiple-comparisons method based on the distribution of the root node distance of a binary tree. Journal of Agricultural, Biological, and Environmental Statistics **7**, 129–142.

Robert G, Melchiorre M, Racca R, Trippi V, Lascano HR. 2009. Apoplastic superoxide level in wheat protoplast under photooxidative stress is regulated by chloroplast redox signals : Effects on the antioxidant system. Plant Science **177**, 168–174.

Robert G, Muñoz N, Alvarado-Affantranger X, Saavedra L, Davidenco V, Rodríguez-Kessler M, Estrada-Navarrete G, Sánchez F, Lascano R. 2018. Phosphatidylinositol 3-kinase function at very early symbiont perception: A local nodulation control under stress conditions? Journal of Experimental Botany **69**, 2037–2048.

Robert G, Muñoz N, Melchiorre M, Sánchez F, Lascano R. 2014. Expression of animal anti-apoptotic gene Ced-9 enhances tolerance during Glycine max L.-Bradyrhizobium japonicum interaction under saline stress but reduces nodule formation. PLoS One **9**, e101747.

Robert G, Yagyu M, Koizumi T, Naya L, Masclaux-Daubresse C, Yoshimoto K. 2021. Ammonium stress increases microautophagic activity while impairing macroautophagic flux in Arabidopsis roots. Plant Journal **105**, 1083–1097.

Rubin G, Tohge T, Matsuda F, Saito K, Scheible W-R. 2009. Members of the LBD Family of Transcription Factors Repress Anthocyanin Synthesis and Affect Additional Nitrogen Responses in Arabidopsis. The Plant Cell **21**, 3567–3584.

Scarpeci TE, Zanor MI, Carrillo N, Mueller-Roeber B, Valle EM. 2008. Generation of superoxide anion in chloroplasts of Arabidopsis thaliana during active photosynthesis : a focus on rapidly induced genes. Plant Mol Biol **66**, 361–378.

Scharfenberg M, Mittermayr L, Von Roepenack-Lahaye E, Schlicke H, Grimm B, Leister D,

Kleine T. 2015. Functional characterization of the two ferrochelatases in Arabidopsis thaliana.
Plant Cell and Environment **38**, 280–298.

Schwarzländer M, Dick TP, Meye AJ, Morgan B. 2015. Dissecting Redox Biology using
Fluorescent Protein Sensors. Antioxidants & Redox Signaling **00**, 150413084017007.

Schwarzländer M, Dick TP, Meyer AJ, Morgan B. 2016. Dissecting redox biology using
fluorescent protein sensors. Antioxidants and Redox Signaling **24**, 680–712.

Schwarzlander M, Fricker MD, Muller C, Marty L, Brach T, Novak J, Sweetlove LJ, Hell R,
Meyer AJ. 2008. Confocal imaging of glutathione redox potential in living plant cells. Journal of
Microscopy **231**, 299–316.

Schwarzländer M, Fricker MD, Sweetlove LJ. 2009. Monitoring the in vivo redox state of plant
mitochondria: effect of respiratory inhibitors, abiotic stress and assessment of recovery from
oxidative challenge. Biochimica et biophysica acta **1787**, 468–75.

Shao N, Duan GY, Bock R, Molekulare M. 2013. A Mediator of Singlet Oxygen Responses in
Chlamydomonas reinhardtii and Arabidopsis Identi fi ed by a Luciferase-Based Genetic Screen
in Algal Cells. The Plant Cell **25**, 4209–4226.

Shao N, Krieger-liszkay A, Schroda M, Beck CF. 2007. A reporter system for the individual
detection of hydrogen peroxide and singlet oxygen : its use for the assay of reactive oxygen
species produced in vivo. The Plant journal **50**, 475–487.

Shibata M, Oikawa K, Yoshimoto K, Kondo M, Mano S, Yamada K, Hayashi M, Sakamoto W,
Ohsumi Y, Nishimura M. 2013. Highly Oxidized Peroxisomes Are Selectively Degraded via
Autophagy in Arabidopsis. The Plant cell **25**, 4967–83.

Signorelli S, Tarkowski ŁP, Van den Ende W, Bassham DC. 2019. Linking Autophagy to Abiotic
and Biotic Stress Responses. Trends in Plant Science **24**, 413–430.

Su T, Li X, Yang M, Shao Q, Zhao Y, Ma C, Wang P. 2020. Autophagy: An Intracellular
Degradation Pathway Regulating Plant Survival and Stress Response. Frontiers in Plant Science
11, 1–16.

Trippi VS, Lascano HR, Melchiorre M. 2009. Plant senescence: A self-induced process. Current
topics in plant biology **10**, 1–15.

Velika B, Kron I. 2012. Antioxidant properties of benzoic acid derivatives against Superoxide
radical. Free Radicals and Antioxidants **2**, 62–67.

Wang P, Wang T, Han J, Li M, Zhao Y, Su T, Ma C. 2021. Plant Autophagy : An Intricate Process
Controlled by Various Signaling Pathways. Frontiers in Plant Science **12**, 754982.

Willems P, Mhamdi A, Stael S, Storme V, Kerchev P, Noctor G, Gevaert K, Van Breusegem F.
2016. The ROS wheel: Refining ROS transcriptional footprints. Plant Physiology **171**, 1720–
1733.

Wintermans M. j. FG, De Mots A. 1965. SPECTROPHOTOMETRIC CHARACTERISTICS OF
CHLOROPHYLLS a AND b AND THEIR PHEOPHYTINS IN ETHANOL. Biochimica et biophysica acta
109, 448–453.

Woodson JD. 2019. Chloroplast stress signals: regulation of cellular degradation and
chloroplast turnover. Current Opinion in Plant Biology **52**, 30–37.

Xiong Y, Contento AL, Bassham DC. 2007*a*. Disruption of Autophagy Results in Constitutive
Oxidative Stress in Arabidopsis. Autophagy **3**, 257–258.

Xiong Y, Contento AL, Nguyen PQ, Bassham DC. 2007*b*. Degradation of oxidized proteins by
autophagy during oxidative stress in Arabidopsis. Plant physiology **143**, 291–9.

Yang M, Bu F, Huang W, Chen L. 2019. Multiple regulatory levels shape autophagy activity in
plants. Frontiers in Plant Science **10**, 1–7.

Yoshimoto K. 2012. Beginning to understand autophagy, an intracellular self-degradation
system in plants. Plant & cell physiology **53**, 1355–65.

Yoshimoto K, Hanaoka H, Sato S, Kato T, Tabata S, Noda T. 2004. Processing of ATG8s ,
Ubiquitin-Like Proteins , and Their Deconjugation by ATG4s Are Essential for Plant Autophagy.
The Plant Cell **16**, 2967–2983.

Yoshimoto K, Jikumaru Y, Kamiya Y, Kusano M, Consonni C, Panstruga R, Ohsumi Y, Shirasu

K. 2009. Autophagy negatively regulates cell death by controlling NPR1-dependent salicylic
acid signaling during senescence and the innate immune response in Arabidopsis. The Plant
cell **21**, 2914–27.
Yoshimoto K, Ohsumi Y. 2018. Unveiling the molecular mechanisms of plant autophagy-from
autophagosomes to vacuoles in plants. Plant and Cell Physiology **59**, 1337–1344.
Zandalinas SI, Fichman Y, Devireddy AR, Sengupta S, Azad RK, Mittler R. 2020. Systemic
signaling during abiotic stress combination in plants. Proceedings of the National Academy of
Sciences **117**, 13810–13820.
Zheng X, Yang Z, Gu Q, Xia F, Fu Y, Liu P, Yin XM, Li M. 2020. The protease activity of human
ATG4B is regulated by reversible oxidative modification. Autophagy **16**, 1838–1850.
Zhou J, Wang J, Cheng Y, Chi YJ, Fan B, Yu JQ, Chen Z. 2013. NBR1-Mediated Selective
Autophagy Targets Insoluble Ubiquitinated Protein Aggregates in Plant Stress Responses. PLoS
Genetics **9**.
Zhuang X, Cui Y, Gao C, Jiang L. 2015. Endocytic and autophagic pathways crosstalk in plants.
Current Opinion in Plant Biology **28**, 39–47.

842
Figure legends

Figure 1: Autophagic flux and cytoplasmic redox state in *Arabidopsis thaliana* plants exposed to
different abiotic stress conditions. (A-C) Seven-day old 35S:GFP-ATG8a transgenic plants and
cytosol-targeted fluorescent redox sensors GFP2-Orp1 and GRX-roGFP2 expressing plants were
exposed to different stress conditions in MS liquid medium. To induce autophagic bodies
accumulation, MS liquid medium was supplemented with 1 µM Concanamycin A. (A)
Representative images showing autophagic bodies accumulation and cytoplasmic redox state.
The stress treatments were performed for 16 h, followed by the evaluation of autophagic bodies
accumulation (GFP-ATG8 columns) and the cytoplasmic redox state (GRX-roGFP and GFP2-Orp1
columns) by confocal microscopy. Arrowheads indicate some of the autophagic bodies
observed. Scale bars: 30 µm. GRX-roGFP and GFP2-Orp1 columns are ratiometric images. For
pseudocolor display, the ratio was coded on a spectral color scale ranging from blue (fully
reduced) to red (fully oxidized). The treatments included: MS with nitrogen (control), MS with
nitrogen in darkness (-C), MS without nitrogen (-N), MS with nitrogen supplemented with 100
mM mannitol (Manitol), 50 mM NaCl (NaCl), 1 µM methyl viologen (MV) and 10 mM H_2O_2. (B)
Average number of autophagic bodies accumulation per section. The average number of
autophagic bodies was calculated from single-plane confocal images. Two sections from at least
7 plants were analyzed. (C) Average of cytoplasmic redox state expressed as the ratio of the
fluorescence intensities measured at 488 nm and 405 nm. At least 6 plants were analyzed.
Statistically significant differences are indicated by asterisks (*$P < 0.05$; error bars: means ± SEM,
DGC test).

864
865

Figure 2: Treatments with MV and H_2O_2 oxidized the cytoplasmic redox state and prevented
autophagy induction in Arabidopsis. (A-C) Seven-day old 35S:GFP-ATG8a transgenic plants and
cytosol-targeted fluorescent redox sensors GFP2-Orp1 and GRX-roGFP2 expressing plants were
exposed to autophagy-inducing conditions and the effect of methyl viologen (MV) and H_2O_2 on
autophagic flux and cytoplasmic redox state were analyzed. (A) Representative images showing
autophagic bodies accumulation and cytoplasmic redox state. The stress treatments were
performed for 16 h, followed by the evaluation of autophagic bodies accumulation (GFP-ATG8
columns) and the cytoplasmic redox state (GRX-roGFP and GFP2-Orp1 columns) by confocal
microscopy. To induced autophagic bodies accumulation, MS liquid medium was supplemented
with 1 µM Concanamycin A. Arrowheads indicate some of the autophagic bodies observed. Scale
bars: 30 µm. GRX-roGFP and GFP2-Orp1 columns are ratiometric images. The treatments
included: MS with nitrogen (control), MS with nitrogen in darkness (-C) supplemented with 10

mM H_2O_2 (-C + H_2O_2), MS without nitrogen (-N) supplemented with 1 µM methyl viologen (-N + MV) or 10 mM H_2O_2 (-N + H_2O_2). 10 mM Tiron and 10 mM Benzoic acid (BA) were used for ROS scavenging (-N + MV + Tiron and -N + MV + BA). (B) Average of autophagic bodies accumulation per section. The average of autophagic bodies number was calculated from single-plane confocal images. Two sections from at least 6 plants were analyzed. (C) Average of cytoplasmic redox state expressed as the ratio of the fluorescence intensities measured at 488 nm and 405 nm from single-plane confocal images. At least 6 plants were analyzed. (D) Effects of oxidative treatments on the autophagic flux evaluated through the degradation of the GFP-ATG8a fusion protein by western blot. Seven-day old 35S:GFP-ATG8a transgenic Arabidopsis seedlings were exposed to the different treatments for 6 h. The GFP-ATG8a fusion has a predicted molecular weight of approximately 40 kDa (closed arrowheads); free GFP has a predicted molecular weight of 27 kDa (open arrowheads). Representative images are shown. Ratio: the numbers in brackets denote the GFP/GFP-ATG8 ratio relativized to the same condition without oxidative treatment. Three replicates were performed (more than fifty plants per replicate). HL: high light treatment (≈1000 µmoles m^{-2} s^{-1}). Asterisks indicate statistically significant differences (*P < 0.05; error bars: means ± SEM, DGC test).

Figure 3: Autophagy is induced after oxidative stress treatments in Arabidopsis. (A-D) Seven-day old transgenic plants expressing 35S:GFP-ATG8a and cytosol-targeted fluorescent redox sensors GFP2-Orp1 and GRX-roGFP2 were subjected to a brief exposure of oxidative stress, either 10 mM H_2O_2 or high light intensity (≈1000 µmoles m^{-2} s^{-1}), for 4 h. Subsequently they were transferred to MS liquid medium for a 20-hour recovery period under continuous light (120 µmoles m^{-2} s^{-1}) (recovery). Control plants were incubated under continuous light (120 µmoles m^{-2} s^{-1}) in H_2O_2-free MS liquid medium for 24 h. Stressed plants were incubated in MS supplemented with 10 mM H_2O_2 or high light intensity (≈1000 µmoles m^{-2} s^{-1}) for 24 h. (A) Representative images showing autophagic bodies accumulation and cytoplasmic redox state. To induce accumulation of autophagic bodies, MS liquid medium was supplemented with 1 µM Concanamycin A. Arrowheads indicate some of the autophagic bodies observed. Scale bars: 30 µm. GRX-roGFP and GFP2-Orp1 columns represent ratiometric images, with pseudocolor display indicating the ratio on a spectral color scale ranging from blue (fully reduced) to red (fully oxidized). (B) Average number of autophagic bodies accumulation per section. The average number of autophagic bodies was calculated from single-plane confocal images. Two sections from at least 6 plants were analyzed. (C) Average cytoplasmic redox state expressed as the ratio of the fluorescence intensities measured at 488 nm and 405 nm from single-plane confocal images. At least 6 plants were analyzed. Asterisks indicate statistically significant differences (*P < 0.05; error bars: means ± SEM, DGC test). (D) Autophagic flux evaluated through the degradation of the GFP-ATG8a fusion protein by western blot. Total protein (40 µg) was subjected to immunoblot analysis with anti-GFP antibodies. The GFP-ATG8a fusion has a predicted molecular weight of approximately 40 kDa (closed arrowheads), while free GFP has a predicted molecular weight of 27 kDa (open arrowheads). Representative images are shown. The numbers in brackets denote the GFP/GFP-ATG8 ratio relativized to the control condition. Three replicates were performed (n > 50 plants per replicate). Asterisks indicate statistically significant differences (*P < 0.05; error bars: means ± SEM, DGC test).

Figure 4: DCMU and methylene blue treatments induced autophagic flux in Arabidopsis. (A-D) Seven-day old transgenic plants expressing 35S:GFP-ATG8a and cytosol-targeted fluorescent redox sensors GFP2-Orp1 and GRX-roGFP2 were incubated in MS medium supplemented with either 20 µM 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU) or 20 µM methylene blue (MB), and the autophagic flux and cytoplasmic redox state were evaluated. (A) Representative images

showing autophagic bodies accumulation and cytoplasmic redox state after 16 h of treatment. To induce autophagic bodies accumulation, MS liquid medium was supplemented with 1 µM Concanamycin A. Some of the autophagic bodies are indicated by arrowheads. Scale bars: 30 µm. The GRX-roGFP and GFP2-Orp1 columns represent ratiometric images. For pseudo color display, the ratio was coded using a spectral color scale ranging from blue (fully reduced) to red (fully oxidized). (B) Average number of autophagic bodies per section. The average number of autophagic bodies was calculated from single-plane confocal images. Two sections from at least 5 plants were analyzed. (C) Average cytoplasmic redox state expressed as the ratio of the fluorescence intensities measured at 488 nm and 405 nm from single-plane confocal images. Analysis was performed on at least 4 plants. Asterisks indicate statistically significant differences (*$P < 0.05$; error bars: means ± SEM, DGC test). (D) Evaluation of the effects of DCMU and MB on autophagic flux through degradation of the GFP-ATG8a fusion protein by western blot. Seven-day old transgenic Arabidopsis seedlings expressing 35S:GFP-ATG8a were exposed to the different treatments for 6 h. Total protein (40 µg) was subjected to immunoblot analysis with anti-GFP antibodies. The GFP-ATG8a fusion has a predicted molecular weight of approximately 40 kDa (closed arrowheads), while free GFP has a predicted molecular weight of 27 kDa (open arrowheads). Representative images are shown. The numbers in brackets denote the GFP/GFP-ATG8 ratio relative to the control condition. Four replicates were performed with more than fifty plants per replicate. (E) Seven-day old Arabidopsis seedlings were exposed to the different treatments for 6 h and soluble sugar content was analyzed. FW: fresh weight. Asterisks indicate statistically significant differences (*$P < 0.05$; error bars: means ± SEM, DGC test).

Figure 5: Autophagic flux is induced in conditionally chloroplast 1O2-hyperproducing *flu* mutant Arabidopsis plants. (A-C) Seven-day old wildtype (Col-0) and *flu* mutant plants expressing 35S:GFP-ATG8a or GFP2-Orp1 were transferred to darkness for 8 h, followed by re-exposure to light. Control plants were maintained under continuous light conditions. After 8 h, the autophagic flux and cytoplasmic redox state were analyzed. (A) Representative images showing autophagic bodies accumulation and cytoplasmic redox state. To induce autophagic bodies accumulation, the MS liquid medium was supplemented with 1 µM Concanamycin A. The GFP2-Orp1 columns represent ratiometric images. For pseudocolor display, the ratio was coded on a spectral color scale ranging from blue (fully reduced) to red (fully oxidized). Arrowheads indicate some of the autophagic bodies observed. Scale bars: 30 µm. (B) Average number of autophagic bodies accumulation per section. The average number of autophagic bodies was calculated from single-plane confocal images. Two different sections from at least 6 plants were analyzed. (C) Average cytoplasmic redox state expressed as the ratio of the fluorescence intensities measured at 405 nm and 488 nm from single-plane confocal images. At least 4 plants were analyzed. Asterisks indicate statistically significant differences (*$P < 0.05$; error bars: means ± SEM, DGC test). (D) Evaluation of autophagic flux through the degradation of the GFP-ATG8a fusion protein by western blot. Seven-day old wildtype (Col-0) and *flu* mutant plants expressing 35S:GFP-ATG8a were transferred to darkness for 8 h and then re-exposed to light for 6 h (after darkness). Continuous light was used as a control (light). Total protein (40 µg) was subjected to immunoblot analysis with anti-GFP antibodies. The GFP-ATG8a fusion has a predicted molecular weight of approximately 40 kDa (closed arrowheads); while free GFP has a predicted molecular weight of 27 kDa (open arrowheads). Representative images are shown. The numbers in brackets denote the GFP/GFP-ATG8 ratio relative to *flu* mutans under control condition. Three replicates were performed (more than fifty plants per replicate). Asterisks indicate statistically significant differences.

978   **FIGURE 1**

992   **FIGURE 3**

 FIGURE 4