# Antigen presentation by tumor-associated macrophages mediates progenitor to terminal exhaustion transition in GBM and other solid tumors

## AUTHORS

Jessica Waibl Polania[1], Alexandra Hoyt-Miggelbrink[1], William H. Tomaszewski[2], Lucas P. Wachsmuth[1,3], Selena J. Lorrey[2], Daniel S. Wilkinson[4], Emily Lerner[3,5], Karolina Woroniecka[1], John B. Finlay[3], and Peter E. Fecci[4,6,*]

## AFFILIATIONS

[1]Department of Pathology, Duke University School of Medicine, Durham, NC 27710, USA

[2]Department of Immunology, Duke University, Durham, NC 27710, USA

[3]Duke Medical Scientist Training Program, Duke University School of Medicine, Durham, NC 27710, USA

[4]Department of Neurosurgery, Duke University School of Medicine, Durham, NC 27710, USA

[5]Department of Biomedical Engineering, Duke University, Durham, NC 27710, USA

[6]Lead contact

*Correspondence: peter.fecci@duke.edu

19 GRAPHICAL ABSTRACT

# SUMMARY

While terminally exhausted T cells (Tex_term) retain important anti-tumor cytotoxic function, it is the relative preservation of renewable, stem-like progenitor exhaustion (Tex_prog) that better indicates immunotherapeutic responsivity. Although restraining the progression from Tex_prog to Tex_term thus takes on clinical significance, the cellular interactions in a tumor microenvironment (TME) governing such progression remain less established. Employing glioblastoma (GBM) and other solid tumors as models of severe exhaustion, we provide a detailed characterization of the progression from Tex_prog to Tex_term within the TME, where we observe a striking and disproportionate loss of Tex_prog over time, leading to a low progenitor exhaustion to terminal exhaustion ratio (PETER). We find exhaustion concentrated within tumor-specific T cell subsets, with cognate antigenic exposure requisite for acquisition of the Tex_term phenotype. However, we implicate tumor-associated macrophages (TAM), and not tumor cells, as the source of antigenic exposure governing the Tex_prog to Tex_term transition. Using cell – cell interaction analysis, we additionally highlight candidate receptor–ligand communications that may be specifically mediating the progression to Tex_term and resultant decline in PETER within the TME.

# KEYWORDS

T cell exhaustion, tumor-associated macrophage, tumor microenvironment, tumor immunology, immunotherapy, brain tumor, brain metastasis, glioblastoma

# INTRODUCTION

Brain tumors, both primary and metastatic, continue to herald poor survival[1,2]. Immunotherapies, such as immune checkpoint blockade (ICB) and chimeric antigen receptor (CAR) T cells, continue to hold promise for improving outcomes, but remain only marginally effective within the intracranial confines[3,4]. This has been especially true in the case of primary malignant brain tumors, such as glioblastoma (GBM)[5]. An established

contributor to immunotherapeutic failures in GBM is tumor-imposed T cell dysfunction,

particularly exhaustion[6-13]. T cell exhaustion is a differentiation state characterized by

progressive and hierarchical loss of effector functions following chronic antigen

exposure[14-17]. As we have previously shown, it is particularly severe in GBM,[18] making

GBM an important model tumor for studying the dynamics and contributors to T cell

exhaustion. Two subsets of T cell exhaustion have now been described by several

groups: progenitor exhaustion (Tex_prog) and terminal exhaustion (Tex_term)[19-21].

Tex_prog describes a stem-like state, characterized by the expression of PD1 and TCF1

(surrogate Ly108), which retains T cell polyfunctionality and is immunotherapy-

responsive[21]. Tex_prog may subsequently differentiate toward Tex_term, which, in turn,

is characterized by high expression of TOX, PD1 and TIM3. Terminally exhausted T cells

have greater cytotoxic capability, but are short-lived and non-replenishing[15,16,21]. Various

studies have suggested that promoting the maintenance of Tex_prog is an important

consideration for generating an immunotherapy-responsive tumor microenvironment

(TME)[19-22]. Details of the tumor-associated factors governing progression from Tex_prog

to Tex_term, however, are less established.

Tumor cells have long been regarded as direct sources of T cell dysfunction within the

TME, and have likewise been a presumed source of the chronic antigen exposure

promoting T cell exhaustion[23,24]. Here, we provide a detailed characterization of the

progression from Tex_prog to Tex_term within the TME, where we observed a striking

and disproportionate loss of Tex_prog over time in GBM, leading to a notably low

progenitor exhaustion to terminal exhaustion ratio (PETER). We reveal that exhaustion is

indeed concentrated within tumor-specific T cell subsets, with cognate antigenic exposure

being requisite for acquisition of the Tex_term phenotype . However, we implicate tumor-

associated macrophages (TAM), and not tumor cells, as the relevant source of antigenic

exposure governing the Tex_prog to Tex_term transition. Using cell – cell interaction

analsyis, we additionally highlight candidate TAM – T cell receptor/ligand communications

that may be specifically mediating the progression to Tex_term and resultant decline in

PETER within the TME.

# RESULTS

**GBM tumor-infiltrating lymphocytes (TIL) demonstrate a severe decline in the Progenitor Exhaustion to Terminal Exhaustion (PETE) Ratio over time**

To better examine the dynamics of T cell exhaustion in GBM and other solid tumor models, but to eliminate any variables that might be imposed by anatomic compartment, we implanted murine GBM, melanoma, and lung (CT2A, B16F0, and LLC, respectively) tumor cells intracranially into syngeneic C57BL/6 mice, harvesting tumors when mice became moribund. We began by evaluating the frequency of both progenitor (Tex_prog) (PD1$^+$Ly108$^+$TIM3$^-$) and terminal (Tex_term) (PD1$^+$Ly108$^-$TIM3+) exhaustion in each model (**Figure 1A**). Consistent with our previous findings[6], GBM tumors demonstrated a higher frequency of Tex_term and correspondingly lower frequency of Tex_prog than the brain metastasis models (**Figure 1B and Supplemental Figure 1A**).

Having found that GBM possessed a particularly low progenitor exhaustion to terminal exhaustion ratio (which we have coined the PETE ratio, or PETER), we took this to indicate that GBM-infiltrating T cells may transition from Tex_prog to Tex_term at an especially high rate. We therefore selected GBM as an archetype tumor model to further examine the processes by which this transition occurs. To begin, we validated  the expected transcriptional and functional differences across tumor-infiltrating Tex_prog and Tex_term from CT2A-bearing mice. We observed higher TOX expression in Tex_term compared to Tex_prog, with both subsets expressing higher expression than non-exhausted PD1$^-$ CD8$^+$ T cells, confirming the typical transcriptional landscape (**Figure 1C**). Likewise, upon PMA/Ionomycin stimulation *ex-vivo*, Tex_prog retained the ability to secrete IL2, IFN$\gamma$, and TNF$\alpha$, whereas Tex_term predominantly secreted IFN$\gamma$ alone (**Figure 1D**).

We next examined the evolution of T cell exhaustion within the tumor over time. We harvested CT2A tumors every 3 days from day 9 (D9) through D21 post-implantation, where mice were moribund at D21 (**Figure 1E**). T cell infiltration was observed within the tumor at D9, with substantial increases seen by D15 (**Figure 1F**). Furthermore, CD8$^+$ PD1$^+$CD44$^+$ T cells significantly increased in frequency from <20% on day 9 to >75% by D21 (**Figure 1G**). Within the PD1$^+$CD44$^+$ subset, we identified Tex_prog and Tex_term

as above, with transcriptional exhaustion confirmed via TOX expression (**Figure 1H**).

Higher TOX levels were consistently seen in Tex_term than in Tex_prog at all timepoints

in which both subsets were present (D12-21), confirming the anticipated pattern of

expression (**Figure 1H**).

Importantly, Tex_prog was readily identifiable by D9 and accounted for ~60% of the

PD1$^+$CD8$^+$ T cells found at the tumor, while Tex_term was notably absent at this early

timepoint. The frequency of Tex_prog subsequently decreased over the course of tumor

progression, accounting for less than 20% of PD1$^+$CD8$^+$ T cells by D21. (**Figures 1I and**

**1J**). Tex_term was identifiable starting D12 at low frequencies (~10%). By D21, however,

Tex_term accounted for ~70% of PD1$^+$CD44$^+$ CD8$^+$ T cells (**Figures 1I and 1J**), further

revealing a drop in PETER over time.

**scRNAseq defines transcriptional heterogeneity amongst GBM-infiltrating**
**Tex_prog and Tex_term with an enrichment for cell-cell communication**
**pathways in Tex_prog**

To better understand the requirements for the transition from Tex_prog to Tex_term, we

first performed scRNAseq on sorted CD8$^+$ T cells from glioma-bearing mice. Informed by

our timecourse, we chose an earlier (D13) and a late (D19) timepoint for our analyses

(**Figure 2A**). We identified six clusters of CD8$^+$ T cells, including two separate terminal

exhaustion populations, which we termed Tex_term_early and Tex_term_late. While 5 of

the clusters were present at both timepoints (D13 and 19), Tex_term_early was present

only at D13 (**Figures 2B and 2C**). Clusters were annotated based on their top

differentially expressed genes and their expression of known T cell exhaustion genes,

namely *Tcf7*, *Tox*, and *Pdcd1* (**Figure 2D and Supplemental Figures 2A and 2B**).

Xcl1+_Tcm and Naïve-like clusters expressed memory/naïve T cell markers (*Sell, Il7r,*

*Id3, Tcf7*), while Xcl1+_Tcm additionally expressed genes associated with activation and

TCR signaling (*Xcl1, Klrk1, Nr4a2*) (**Figure 2D)**. The expression of *Pdcd1* and *Tox* on

Tex_prog, Tex_int, Tex_term_early, and Tex_term suggested that these were T cell

exhaustion subsets (**Figure 2D and Supplemental Figure 2B**). Tex_prog expressed

effector (*Ifng, Tbx21*) and stem-like genes (*Cd28, Il7r, Tcf7, Tbx21*) and had a lower T

cell exhaustion signature score (described by Im et al[19]) than Tex_int, Tex_term_early,

and Tex_term_late (**Figures 2D and 2E**). This could suggest that the exhaustion program may not be fully established in Tex_prog. Next, Tex_int had increased expression of exhaustion-associated genes (*Pdcd1, Tox, Entpd1*) and a decrease in effector (*Klrg1*) and memory associated genes (*Tcf7, Il2*) (**Figure 2D**). Cytotoxicity-associated genes, such as *Gzmb, Gzma*, and *Gzmk* were expressed at the highest levels in Tex_term_early and Tex_term_late, consistent with previous characterization of terminal exhaustion[21] (**Figure 2D**). Finally, Tex_term_late, predominantly found at the late timepoint and consistent with our time-course characterizations, expressed the highest levels of exhaustion-associated genes (*Pdcd1, Tox, Havcr2, Entpd1*) and lost *Tcf7* expression (**Figure 2D and Supplemental Figure 2B**).

Once confident in our annotations, we employed pathway enrichment analysis for both Tex_prog and Tex_term_late, which represented the two extreme ends of the exhaustion pathway. This analysis revealed an abundance of pathways related to T cell migration, differentiation, cell-cell communication and response to stimuli (cytokine, chemokines) in Tex_prog (**Figures 2F and 2G**). This is in keeping with the current understanding of the Tex_prog state and its prevalence among T cells entering or recently entering tumor [25,26]. In contrast, Tex_term were enriched for cytotoxic pathways, almost exclusively (**Figures 2F and 2G**). Importantly, this highlights that Tex_prog, unlike Tex_term, likely remain in a signal-responsive state within the TME.

Lasty, we applied pseudotime analysis to our dataset to infer the temporal progression amongst the T cell exhaustion subsets. This analysis spatially separated the Naïve_like and Xcl1+_Tcm clusters from the other four clusters, supporting their identification as separate, non-exhausted lineages (**Figure 2H**). The pseudotime plot ultimately validated the progression from progenitor to terminal exhaustion through an intermediate population (Tex_int). Interestingly, however, two trajectories were determined, both beginning with Tex_prog (**Figure 2I**). While one ended with Tex_term_late, the other ended in Tex_term_early. This finding is likely simply highlighting differences in phenotypes for the predominant ending node at the early and late timepoints.

To further examine lineage relationships between the T cell exhaustion clusters, we utilized our paired T cell receptor sequencing (TCRseq) dataset. To this end, we observed

TCR clonal overlap between predicted Tex clusters (**Figure 2J**). In combination with our
pseudotime results, this analysis suggests a trajectory of shared TCR clones, beginning
with Tex_prog, progressing through Tex_int and eventually ending with Tex_term. We
next compared clonal expansion between clusters, where we observed the greatest
frequency of "Large" expanded clones, (defined as between 20 and 100 cells with the
same clone), in Tex_term_late, and the lowest such frequency in Tex_prog (**Figure 2K**
**and 2L**). Correspondingly, Tex_prog had the greatest diversity index score, as measured
by the Shannon and Inverse Simpson indices, which has often been associated with
response to immunotherapy (**Supplemental Figure 2C**)[27].  Moreover, when we plotted
the 10 most expanded clones on a UMAP and divided the analysis by timepoint, we
observed that the expanded clones overwhelmingly resided in Tex_int, Tex_term_early
and Tex_term_late, but not Tex_prog (**Supplemental Figure 2D**). This suggests that
clonal expansion in response to TCR stimulation closely reflects T cell exhaustion
progression.

**Tumor cells are not a necessary source of the TCR stimulation mediating**
**Tex_prog to Tex_term transition**

In light of the above observations, we hypothesized that antigen-specific TCR activation
would be required for progression of Tex_prog to Tex_term within the TME, and thus that
Tex_term would be concentrated amongst tumor-specific T cells. To test this directly, we
employed a model in which CT2A gliomas were engineered to express the antigen TRP2
(See STAR Methods). We then adoptively transferred either tumor-specific TRP2 TCR-
transgenic CD8$^+$ T cells (TRP2-TCR) or irrelevant OVA antigen-specific (OT1) CD8$^+$ T
cells into CT2A-TRP2-bearing mice (**Figure 3A**). All T cells were activated *ex vivo* prior
to transfer. Two weeks later, tumors were harvested, and adoptively transferred T cells
were identified as TRP2 tetramer$^+$CD45.1$^+$ (TRP2-TCR) or OVA tetramer$^+$ (OT1) and
evaluated for exhaustion phenotypes. Unsurprisingly, recovery of TRP2-TCR T cells from
tumor was significantly better than recovery of OT1 T cells (**Figure 3B**). Recovered TRP2-
TCR T cells also demonstrated significantly higher acquisition of a Tex_term phenotype
than OT1 T cells, which only minimally acquired PD1 expression (**Figures 3C and 3D**
**and Supplemental Figure 3A**). Unlike OT1 T cells, transferred TRP2-TCR T cells

developed into predominantly Tex_int and Tex_term, similar to endogenous tumor-specific T cells (TRP2 tetramer[+]CD45.2[+]) (**Supplemental Figure 3B)**. Taken together, these data strongly suggest that T cells in the TME must encounter their cognate antigen to progress along the exhaustion pathway and complete the transition to Tex_term.

As tumor antigen-specific TCR stimulation appeared to be necessary for exhaustion progression, we first supposed that the relevant antigen-cognate interaction would be between T cells and tumor cells. To evaluate this, we compared the progression of T cell exhaustion *in vivo* within CT2A-TRP2 and CT2A-TRP2-β2mKO tumors, where the latter constitutes a cell line where β2m is knocked out using CRISPR-Cas9 (See STAR Methods). Knockout of β2m leaves the tumor without the capacity to build functional MHC I molecules, and thus without the capacity to present antigen to T cells (**Figure 3E and Supplemental Figure 3C**). Surprisingly, we observed no significant difference in the frequency of PD1[+] TRP2-TCR T cells or in the frequency of T cell exhaustion subsets (Tex_prog or Tex_term) in the absence of tumor MHC I expression and antigen presentation (**Figures 3F, 3G, and 3H**). Of note, the extent of T cell infiltration was comparable between MHC I expressing and non-expressing tumors (**Supplemental Figure 3D**).

## TAM are the most frequent and most potent cross-presenting antigen-presenting cells (APC) within the intracranial TME

With tumor cells seemingly eliminated as the source of exhaustion-relevant antigen-cognate interactions with T cells, we next sought to narrow down additional candidate antigen presenting populations. We therefore evaluated other sources of potentially determinant antigen-cognate interactions within the TME. Such relevant APC might include monocytes, TAM, microglia, and/or dendritic cells (DC)[28]. Using a scRNAseq dataset of CD45[+] cells sorted from CT2A tumors previously published by our group[29], we identified predicted sources of MHC I cell-cell interactions in the TME. We isolated a component of the dataset that included the above APC types and CD8[+] T cells, then conducted cell-cell interaction analysis using the CellChat pipeline[29,30] (Schematic in **Figure 4A**). Importantly, MHC I signaling was a highly predicted interaction between T cells and all potential APC (**Figure 4B**). This analysis revealed the greatest *number* of

predicted interactions between CD8$^+$ T cells and Apoe$^+$ TAM, while the *strength* of

predicted interactions was similar across APC (**Figure 4C and Supplemental Figure**

**4A**). Notably, TAM greatly outnumber other APC in the CT2A TME, comprising ~40% of

tumor-infiltrating immune cells (**Figure 4D**). Combined, these results suggested that the

relevant antigen-cognate interactions promoting exhaustion progression within the TME

are occurring between CD8$^+$ T cells and APC (and not tumor), with TAM being the the

most abundant source of antigen.

To investigate the relevant antigen-presenting functions of the various APC subtypes

within CT2A, we directly evaluated their relative *in vivo* phagocytic activity within the TME.

To this end, we implanted GFP-expressing CT2A tumors and evaluated the frequency of

GFP positivity within intratumoral APC when mice were moribund (schematic in **Figure**

**4E**). We focused on monocytes (CD11b$^+$F4/80$^+$Ly6C$^{hi}$), TAM (CD11b$^+$F4/80$^+$MHC II$^+$),

DC (F4/80$^-$CD11c$^+$MHC II$^+$), and microglia (CD45$^{int}$CD11b$^+$F4/80$^+$) (**Supplemental**

**Figure 4B**). We found that monocytes, TAM, and DC all took up GFP (**Figure 4F**). After

phagocytosing exogenous antigen, APC must load and cross-present the antigen via

MHC I to interact with CD8 T cells[31]. We therefore next probed for differential MHC I

expression as a surrogate for subsequent tumor antigen cross-presentation. TAM and

DC both had significantly higher MHC I on their surface than other APC populations

(**Figures 4G and 4H**).

To more directly evaluate the cross-presentation capacity of intratumoral APC, we

implanted OVA-expressing CT2A tumors and subsequently assessed the various APC

populations by flow cytometry for the presence of SIINFEKL-loaded MHC I (Schematic in

**Figure 4I**). Among APC, we observed the greatest frequency of SIINFEKL-loaded MHC

I on TAM and their monocyte precursors (**Figure 4J**). Combining our data, then, TAM

appeared to exhibit the greatest potency for both phagocytosing and cross-presenting

tumor antigen within the TME.

**TAM mediate progenitor to terminal exhaustion transition in GBM and other**
**solid tumor models**

To better establish the requirement for TAM in the progression from Tex_prog to

Tex_term, we implanted CT2A tumors into CCR2KO mice. CCR2KO mice exhibit

impaired monocyte trafficking toward sites of inflammation[32], therefore establishing a

TME that is relatively deficient in TAM, which are monocyte-derived. To further the extent

of TAM depletion in this model, however, we also administered tumor-bearing CCR2KO

mice an anti-CSF1R antibody, which eliminates differentiated TAM (schematic in **Figure**

**5A**)[33]. We observed efficient depletion of TAM with this approach (**Supplementary**

**Figure 5A**). Importantly, there was a significant increase in PETER observed in the

CCR2KO setting (**Figures 5B and 5C**), suggesting a relative failure to progress from

Tex_prog to Tex_term. Notably, we likewise observed a strong positive correlation

between the frequency of Tex_term and the frequency of TAM remaining within the TME,

as Tex_term proportions dropped substantially with improved TAM depletion (**Figure 5D**).

Importantly, we did not observe similar alterations to PETER when we depleted DC

(**Figure 5E and 5F**) or microglia (**Figure 5G and 5H) (Supplemental Figures 5B-5E**),

further suggesting TAM as the pertinent APC population for driving the progression of T

cell exhaustion in this model.

With these observations, we next evaluated the role of MHC I presentation by TAM in

driving progression of T cell exhaustion. To this end, we generated $\beta$2m bone marrow

chimeras ($\beta$2mKO donors into WT recipients), wherein hematopoietic cells (including

hemaptopoietically-derived APC) lacked $\beta$2m and antigen presenting capacity (see STAR

Methods). While we continued to observe low levels of MHC I expression on TAM in this

model, likely as a result of TAM trogocytosis of MHC I from surrounding $\beta$2m-replete

cells[34] (**Supplemental Figures 5F and 5G**), we observed a positive correlation between

persisting MHC I[+] on TAM and the frequency of Tex_term (**Figure 5I**).

These findings in CT2A raised the question as to whether the role of TAM in driving

exhaustion progression was exclusive to GBM and/or to the intracranial environment. To

assess this, we employed the same TAM depletion approach in both orthotopic and

intracranial murine models of melanoma (schematic in **Figure 5J**). As before, we saw

efficient depletion of TAM (**Supplementary Figures 5H and 5I**). Importantly, in both

peripheral and intracranial melanoma models, we observed a significant increase in

PETER within the TME in the setting of TAM depletion (**Figures 5K-5N**). These findings

implicate TAM as a primary source of tumor-antigen presentation mediating T cell

exhaustion progression across multiple solid tumor types and locations. Taken together,

these data suggest Tex_prog progresses to Tex_term independently of tumor-derived

antigen presentation, and instead requires antigen-cognate interactions between T cells

and antigen cross-presenting APC, with TAM being the most relevant of these.

**297 CellChat predicts secondary interactions between TAM and CD8+ T cells in**
**298 mice and humans that may contribute to exhaustion progression**

Given the above role uncovered for TAM, we returned to the cell – cell interaction analysis

from Figure 3 to further interrogate interactions between TAM and Tex_prog that might

be contributors to Tex_prog progression to Tex_term (schematic in **Figure 6A**). Several

TAM ligands or surface molecules were highlighted in our scRNAseq dataset, with

pathways ranging from inhibitory (PD-L1, SPP1, GALECTIN) to stimulatory (CD86), to

cell-cell adhesion (ICAM, JAM) to migration (CXCL), among others (**Figure 6B**). We

further refined these possibilities by selecting receptors/ligands that were expressed in

our scRNAseq dataset and specifically present in the Tex_prog cluster (**Figure 6C**). From

these analyses, we observed that cell – cell adhesion (*Itgb1, Itgb2, Itga4*), migration

(*Cxcr3*) and stimulatory molecules (*Cd28*) were particularly enriched in Tex_prog

compared to Tex_term, consistent with the pathways identified in Figure 2F. Tex_term,

by comparison, had greater expression of inhibitory receptors (*Pdcd1, Ctla4, Tigit*) and

the chemokine receptor, *Cxcr6*.

To determine which of the above predicted interactions might also be prevalent in

patients, we turned to publicly available human GBM and metastatic melanoma datasets.

We integrated 15 newly diagnosed GBM (ndGBM) and 5 treatement-naïve metastatic

brain melanoma (MBM) samples and focused on TAM and CD8+ T cells for further

analysis[35,36]. Once integrated, we derived 8 TAM clusters and 3 CD8 clusters that were

shared between tumor types (**Figure 6D**). As expected, the abundance of TAM was

greater in ndGBM than in MBM (**Figure 6E**). We then performed cell – cell communication

analysis with the human samples using CellChat as previously described[30]. Broadly, we

noted that CD8+ T cells possessed high "incoming interaction" levels but low "outgoing

interaction" strength for both tumor types, whereas TAM had comparable

incoming/outgoing strengths, highlighting their interconnectedness within the TME

(**Figure 6F**). The total interaction strength was much higher in MBM than ndGBM, suggesting the MBM TME, as a whole, exhibits receptor – ligand pairs at a higher level (**Figure 6F**).

Given the higher strength of incoming interactions amongst CD8[+] T cells, we evaluated which predicted interactions were prevalent between TAM and CD8[+] T cells across tumor types. Among those predicted, several overlapped with our CellChat analysis of murine GBM, including MHC-I, SPP1, CD86, ICAM, FN1, JAM, CXCL, and GALECTIN (**Figures 6B and 6G**). Interestingly, ndGBM was primarily enriched for cell – cell adhesion interactions, whereas MBM had exhibited a greater representation of inflammatory/migration pathways. These results align with perceptions of GBM as an "immunologically cold" tumor compared to MBM[37,38]. When comparing the intensity of these pathways between tumor types, we again observed an increased strength and number of interactions in the MBM group (**Figure 6H**). Notably, the SPP1 and FN1 pathways were significantly upregulated in the ndGBM samples compared to MBM. These findings highlight enriched interactions between TAM and CD8[+] T cells that may be implicated in T cell exhaustion progression across tumor types, tumor locations, and species, thus warranting further investigation.

# DISCUSSION

The abundance of Tex_prog within a TME is currently believed to reflect immunotherapeutic responsivity [15,21,22,38,39]. Initially identified as the T cell differentiation state remaining reactive to anti-PD1 treatment in chronic infection, Tex_prog has now been recognized as a consistent correlate of ICB response[15,19,20]. Some studies have further identified the abundance of Tex_prog relative to Tex_term as critical[22]. Here, we detail the severe and disproportionate loss of Tex_prog over time in GBM and other solid tumors and implicate TAM, but not tumor cells, as the mediators of the Tex_prog to Tex_term transition. Recognizing the importance of this shift from Tex_prog to Tex_term, we, in turn, coin and identify the PETER as an important indicator of the dynamics of T cell exhaustion within the TME. As GBM possesses a particularly low PETER and is likewise unresponsive to immune checkpoint blockade, we suggest that a therapeutic goal for licensing improved immunotherapeutic responses may ultimately be finding ways

to optimize PETER within a given TME .

Further characterizing Tex_prog and Tex_term, we observe an enrichment for
chemotactic and cell-cell adhesion pathways in Tex_prog, and we implicate TAM as
driving a subsequent loss of T cell polyfunctionality and stem-like properties as T cells
increasingly differentiate toward Tex_term in tumor. In the absence of TAM, we instead
observe retention of the Tex_prog population and a resultant increase in PETER, with the
retained Tex_prog possessing established implications for immunotherapeutic
responsivity[16,19-22].

Until recently, most defining studies for T cell exhaustion have been conducted in a
chronic infection model of lymphocytic choriomeningitis virus (LCMV)[14,17,40]. In the LCMV
model, it is unfortunately impossible to determine whether T cell chronic antigen exposure
occurs on the surface of infected target cells or APC, as they are typically one and the
same. As a result, a common presumption is that chronic exposure to antigen presented
by target cells is a driver of T cell exhaustion progression. In contrast, in tumors, T cell
antigen exposure may occur on the surface of tumor cells presenting endogenous
antigen, or in the context of TME APC cross-presenting exogenous antigen. Looking to
assess whether one antigenic context might be more crucial to the derivation of
exhaustion in the TME, we removed the capacity for MHC I presentation from tumor cells
and  observed no change in the frequency of T cell exhaustion subsets. This suggested
that encountering antigen on tumor cells is not required for the differentiation of Tex_prog
to Tex_term, leaving intratumoral APC as our next-investiagted source of instigatory
antigen-cognate interactions.

In our intial querying of intratumoral APC, DC were a leading logical candidate for
exhaustion-eliciting interactions with Tex_prog. Surprisingly, however, interactions with
DC, although present, proved to be neither necessary nor sufficient for the differentiation
of Tex_prog to Tex_term. Rather, we found that both TAM and TAM MHC I are necessary
components. Importantly, in addition to being the most abundant APC, TAM also
demonstrated the greatest propensity for phagocytosis and cross-presentation in the
TME. The sheer number of  predicted TAM – T cell interactions are important as they are
generally only weakly activating and often even immunosuppressive compared to DC –

T cell interactions[28,41-43]. Our findings, however, highlight a previously unappreciated

immunoregulatory role for TAM and provide a new focus on determining relevant  T cell

– TAM interactions for targeting in an effort to better license immunotherapeutic

responses.

While antigen cognate interactions with TAM may be a necessary first step in driving T

cell exhaustion within the TME, our data suggest that there are additional secondary

interactions that follow that may more proximally mediate the Tex_prog to Tex_term

transition. Other groups have proposed co-stimulatory signals through the CD86/CD28

axis as contributors[26]. Our CellChat findings suggest that such interactions may indeed

play a role in more inflammatory tumors, such as MBM, but this axis was not prevalent in

our human ndGBM dataset. Interestingly, we identified SPP1, which binds to integrins

(CD44 or CD29) on CD8$^+$ T cells, as a highly predicted component of interaction between

TAM and CD8$^+$ T cells in in both murine and human tumors. Importantly, SPP1 has

recently been ascribed a T cell exhaustive role in the setting of GBM. Kilian *et al.* identified

a pathway in which the absence of CD4$^+$ T cell – TAM MHC II interactions led to increased

secretion of SPP1 by TAM, which suppressed surrounding CD8$^+$ T cells and increased

their expression of exhaustion-associated factors[44]. Further studies are ultimately needed

to better address the nature and role of specific interactions between TAM and CD8$^+$ T

cells.

Taken together, we have highlighted TAM – T cell interactions as critical for driving CD8$^+$

T cell exhaustion progression in GBM and other solid tumors. Future work will focus on

defining specific TAM – T cell interactions and their relative contributions to the

differentiation of Tex_prog to Tex_term. The goal remains to promote a TME with a

heightened PETER and renewable Tex_prog populations in an effort to better license

immunotherapeutic responses.

## ACKNOWLEDGMENTS

Dr. Fecci, and the research contained herein, is supported by a Cancer Research Institute Lloyd J Old STAR award. We would like to thank the Duke Molecular Genomics Core, Duke Human Vaccine Institute Flow Cytometry Core, and the Duke Cancer Institute Flow Cytometry Core.

## AUTHOR CONTRIBUTIONS

Conceptualization, J.WP and P.E.F.; Methodology, J.WP., W.H.T., A.M.H-M., L.P.W., D.S.W., E.L., and K.W.; Investigation, J.WP.; Resources, J.WP., W.H.T., A.M.H-M., and J.B.F.; Writing – Original Draft, J.WP.; Writing – Review & Editing, J.WP., A.M.H-M., W.H.T., L.P.W., S.J.L. and P.E.F.; Supervision, P.E.F.; Funding Acquisition, J.WP., and P.E.F.

## DECLARATIONS OF INTEREST

The authors declare no competing interests.

## FIGURE TITLES and LEGENDS

Figure 1: GBM tumor-infiltrating lymphocytes (TIL) demonstrate a severe decline in the Progenitor Exhaustion to Terminal Exhaustion (PETE) Ratio over time.

(A) Representative contour plots of Tex_prog and Tex_term across tumor types.

(B) PETER across CT2A, B16F0 and LLC intracranial tumors, harvested when mice were moribund. n = 8-10 mice/group.

(C and D) Geometric mean fluorescence intensity (gMFI) of intracellular TOX expression (C) and quantification of TNFa/IFNy/IL2 of Tex_prog and Tex_term in 10K CT2A at D18 (D). n = 5 mice. Representative of two independent experiments.

(E) Experimental layout. 10K CT2A cells implanted D0, tumors were harvested every 3 days starting D9 until mice were moribund on D21. Created with Biorender.

(F) number of CD8+ T cells per harvested tumor. n = 5 mice/timepoint.

(G) Frequency of PD1$^+$CD44$^+$ of CD8$^+$ T cells. n = 5 mice/timepoint.

(H) Geometric mean fluorescence intensity (gMFI) of TOX of Tex_prog and Tex_term across timepoints with >100 total PD1$^+$CD44$^+$ CD8$^+$ T cells. n = 5 mice/timepoint.

(I and J) Representative plots showing Ly108 vs. TIM3 progression from D9 to D21, Tex_prog = top left, Tex_int = bottom left, Tex_term = bottom right (I) and quantification of Tex subsets of PD1+CD44+ CD8+ T cells over time (J). n = 5 mice/timepoint.

All of the data are means ± SEMs; p-values are shown on individual plots as determined by 1-way ANOVA and post-hoc Tukey test (A, F, G), two-sided paired t-test (B) or RM 2-way ANOVA and post-hoc Tukey test (D and H). See also Figure S1.

Figure 2: scRNAseq defines transcriptional heterogeneity amongst GBM-infiltrating Tex_prog and Tex_term with an enrichment for cell-cell communication pathways int Tex_prog.

(A) Experimental layout for scRNAseq experiment. tumors were harvested at an early (D13) and late (D19) timepoint. Created with Biorender.

(B) UMAP of 6 CD8 T cell clusters following QC.

(C) Frequency of each cluster per timepoint. (D) Heatmap of exhaustion associated transcription factors, cytokines/secreted molecules, and surface markers.

(E) AddModuleScore of exhaustion signature from Im et al., 2016, violin plot and density plot representations.

(F) Significantly upregulated pathways in Tex_prog and Tex_term from Hallmark, Reactome, Gene Ontology databases.

(G) Relevant genes from upregulated pathways in F.

(H) Trajectory analysis overlaid onto clusters from B.

(I) Trajectory analysis overlaid onto pseudotime.

(J) Circle plot of overlapping clones across Tex_clusters. Tex_prog, Tex_int have overlapping clones with all other clusters. Tex_term and Tex_int_2 do not have overlapping clones.

(K) UMAP of cloneTypes across clusters.

(L) Frequency of cloneTypes across clusters.

See also Figure S2.

Figure 3: Tumor cells are not a necessary source of the TCR stimulation mediating Tex_prog to Tex_term transition

(A) Experimental layout. 250K *ex-vivo* activated and transduced OT1 and TRP2-TCR T cells were intracranially implanted into D7 CT2A-TRP2-bearing mice. Tumors were harvested 14 days later.

(B) Frequency of implanted cells recovered on D21. n = 7 mice/group.

(C and D) Frequency of PD1+ CD8+ T cells in OT1 vs. TRP2-TCR (C) and representative contour plot PD1 and TIM3 expression of OT1 and TRP2-TCR T cells 14 days post-transfer (D). n = 7 mice/group

(E) Experimental layout. 50K CT2A-TRP2 or CT2A-TRP2-B2mKO were implanted

intracranially and harvested when mice were moribund. Created with Biorender.

(F) Frequency of PD1+ TRP2-TCR T cells between groups. n = 10 mice/group.

(G and H) Frequency of Tex subset of PD1+TRP2-Tetramer+ CD8+ T cells and

representative contour plot (G) and representative contour plot of Ly108 and TIM3

expression between groups (H). n = 10 mice/group.

All of the data are means ± SEMs; p-values are shown on individual plots as measured

by two tailed t-test (B, C, F) or RM 2-way ANOVA (G). See also Figure S3.

Figure 4: TAM are the most frequent and most potent cross-presenting antigen-

presenting cells (APC) within the intracranial TME.

(A) Experimental layout. CD45+ sorted scRNAseq from Tomaszewski et al., 2022 was

analyzed with CellChat to determine which APC had predicted MHC I interactions with

CD8 T cells. Created with Biorender.

(B) Heatmap showing communication probability of MHC I interactions in the TME. All

APC were predicted to interact with CD8 TIL through MHC I.

(C) Circle plot displaying number of interactions (thickness of arrow/number) across APC

populations.

(D) Frequency of infiltrating immune cells from Tomaszewski et al., 2022 dataset.

(E) Experimental layout. 25K CT2A-GFP was implanted intracranially and harvested

when mice were moribund. GFP+ APC were evaluated. Created with Biorender.

(F) Frequency of GFP+ T cells (CD3$^+$), TAM (CD45$^{hi}$CD11b$^+$F4/80$^+$), DC (CD11b$^+$MHC

II$^+$F4/80$^-$) and Microglia (CD45$^{hi}$CD11b$^+$F4/80$^+$). n = 5 mice.

(G and H) gMFI of MHC I of APC populations from F (G) and representative histogram

(H). n = 5 mice.

(I) Experimental layout. 100K CT2A-OVA was implanted intracranially and harvested

when mice were moribund.

(J) Frequency of SIINFEKL-loaded MHC I of T cells, TAM, Monocytes

(CD11b+F480$^{int}$Ly6C$^{hi}$), DC, and Microglia. n = 9 mice.

p-values are plotted on individual graphs. Statistical significance was determined by 1-
way ANOVA (F, G, J) with post-hoc Tukey's test for multiple comparisons. All data are
plotted as mean ± SEM. See also Figure S4.

Figure 5: TAM mediate progenitor to terminal exhaustion transition in GBM and other
intracranial solid tumor models.

(A) Experimental layout. TAM and monocytes were depleted in CCR2KO mice with
aCSF1R depletion antibody. 10K CT2A was implanted intracranially and harvested when
mice were moribund. Created with Biorender.

(B and C) Frequency of Tex_subsets of PD1+CD44+ (B) and PETER between WT and
TAM depleted groups (C). Data pooled from 2 independent experiments. n = 10-11
mice/group.

(D) Simple linear correlation of %Tex_term of PD1+CD44+ CD8+ T cells vs. % TAM of
CD45+. Data pooled from 2 independent experiments. n = 11 mice/group.

(E and F) CD11c+ DC were depleted with 5ng/g DTR injections on D8, D11, D14, D17
post CT2A-implantation. Frequency of T cell exhaustion subsets (E) and and PETER
between WT and DC depleted groups (F). n = 5 mice/group.

(G and H) Microglia were depleted with 290mg/kg PLX3397 free-fed chow starting 1 week
prior to CT2A implantation. Frequency of T cell exhaustion subsets (G) and PETER
between WT and microglia depleted groups (H). n = 10 mice/group.

(I) Simple linear correlation of frequency of Tex_term vs. Frequency of MHC I+ TAM in
B2mKO chimera mice implanted with CT2A.

(J) Experimental layout. TAM and monocytes were depleted in CCR2KO mice with
aCSF1R depletion antibody. 1K B16F0 was implanted intracranially or 100K B16F0 was
implanted subcutaneously and harvested when mice were moribund. Created with
Biorender.

(K and L) Frequency of Tex_subsets of PD1+CD44+ B16F0 IC (K) and PETER between

WT and TAM depleted groups (L). n = 9-10 mice/group.

(M and N) Frequency of Tex_subsets of PD1+CD44+ B16F0 SQ (M) and PETER
between WT and TAM depleted groups (N). Data pooled from 2 independent
experiments. n = 6-7 mice/group.

p-values are plotted on individual graphs. Statistical significance was determined by two-
sided unpaired t test (C, F, H, L, N) or RM-2-way ANOVA with post-hoc Tukey's test for
multiple comparisons (B, E, G, K, M). All data are plotted as mean ± SEM. See also Figure
S5.

Figure 6: CellChat predicts secondary interactions between TAM and CD8[+] T cells in mice
and humans that may contribute to exhaustion progression

(A) Experimental layout. CD45+ sorted scRNAseq from murine GBM, ndGBM and MBC
was analyzed with CellChat to determine additional TAM-T cell interactions. Created with
Biorender.

(B) Enriched predicted interactions with TAM (orange) as the source and CD8 TIL (blue)
as the target from murine GBM.

(C) Receptors expression in Tex_prog vs. Tex_term from CD3+ sorted dataset from
Figure 2 of predicted interactions from 6B.

(D) UMAP of human ndGBM and MBM datasets.

(E) Frequency of TAM vs. CD8s in ndGBM vs. MBM.

(F) Incoming vs. outgoing interaction strength across cell types in ndGBM vs. MBM.

(G) Heatmap of all incoming signaling patterns in ndGBM vs. MBM.

(H) Significantly upregulated predicted pathways in ndGBM (maroon) vs. MBM (grey).

 TABLES with TITLES and LEGENDS

 **Table S1: Exhaustion signature (Im et al. 2016)**

| |
| --- |
| *Pdcd1* |
| *Tox* |
| *Il21r* |
| *Sh2d2a* |
| *Cd8a* |
| *Nkg7* |
| *Fgl2* |
| *Entpd1* |
| *Fasl* |
| *Ccl5* |
| *Apol7e* |
| *Arhgap9* |
| *Nedd9* |
| *Cd38* |
| *Pvrig* |
| *Trbc2* |
| *Chn2* |
| *2900026A02Rik* |
| *Pstpip1* |
| *Chst12* |
| *Cd3e* |
| *Cxcr6* |
| *Cd3g* |
| *Adgrg1* |
| *Prkch* |
| *Tap2* |
| *Irf1* |
| *Tapbpl* |
| *AW112010* |
| *Arl6ip1* |
| *Gng2* |
| *Adam19* |
| *Gimap7* |
| *Gzmk* |

## STAR METHODS

561 **Key Resources Table**

| Reagent or Resource | Source | Identifier |
|---|---|---|
| Antibodies | | |
| anti-mouse anti-GFP - Alexa Fluor® 488 (clone FM264G) | Biolegend | Cat# 338008; RRID:AB_2563288 |
| anti-mouse CD11b - APCR700 (clone M1/70) | BD Biosciences | Cat# 564985; RRID:AB_2739033 |
| anti-mouse CD11c - BV421 (clone N418) | BD Biosciences | Cat# 565452; RRID:AB_2744278 |
| anti-mouse CD16/32TruStain FcX™ (clone 93) | Biolegend | Cat# 101320; RRID:AB_1574975 |
| anti-mouse CD19 - Alexa Fluor® 488 (clone 6D5) | Biolegend | Cat# 115521; RRID:AB_389307 |
| anti-mouse CD24 - Brilliant Violet 605™ (clone M1/69) | Biolegend | Cat# 101827; RRID:AB_2563464 |
| anti-mouse CD3 - APC (clone 145-2C11) | Biolegend | Cat# 100312; RRID:AB_312677 |
| anti-mouse CD3 - PECy5 (clone 145-2C11) | Biolegend | Cat# 552774; RRID:AB_394460 |
| anti-mouse CD3 - PECy7 (clone 145-2C11) | Biolegend | Cat# 100310; RRID:AB_312675 |
| anti-mouse CD39 - PE-eFluor610 (clone 24DMS1) | Invitrogen | Cat# 61-0391-82; RRID:AB_2848486 |
| anti-mouse CD39 - PECy7 (clone 24DMS1) | Invitrogen | Cat# 61-0391-82; RRID:AB_2848486 |
| anti-mouse CD44 - Brilliant Violet 785™ (clone IM7) | Biolegend | Cat# 103059; RRID:AB_2571953 |
| anti-mouse CD44 - PE-CF594 (clone IM7) | BD Biosciences | Cat# 562464; RRID:AB_11153123 |
| anti-mouse CD45 - APCCy7 (clone 30-F11) | Biolegend | Cat# 103116; RRID:AB_312981 |

| anti-mouse CD45 - BV605 (clone 30-F11) | Biolegend | Cat# 103155; RRID:AB_2650656 |
|---|---|---|
| anti-mouse CD45.1 - Alexa Fluor® 488 (clone A20) | Biolegend | Cat# 110718; RRID:AB_492862 |
| anti-mouse CD45.1 - PECy7 (clone A20) | BD Biosciences | Cat# 110730; RRID:AB_1134168 |
| anti-mouse CD45.2 - APCCy7 (clone 104) | Biolegend | Cat# 109824; RRID:AB_830789 |
| anti-mouse CD86 - Brilliant Violet 650™ (clone GL-1) | Biolegend | Cat# 109824; RRID:AB_830789 |
| anti-mouse CD8a - APCR700 (clone 53-6.7 | BD Biosciences | Cat# 564983; RRID:AB_2739032 |
| anti-mouse CX3CR1 - APC (clone SA011F11) | Biolegend | Cat# 149007; RRID:AB_256449 |
| anti-mouse CX3CR1 - APC (clone SA011F11) | Biolegend | Cat# 149006; RRID:AB_2564315 |
| anti-mouse F4/80 - PECy5 (clone BM8) | Invitrogen | Cat# 15-4801-82; RRID:AB_468798 |
| anti-mouse H2Kb bound to SIINFEKL - PE (clone 25-D1.16) | Biolegend | Cat# 141604; RRID:AB_10895905 |
| anti-mouse H2Kb/H2Kd - PECy7 (clone 28-8-6) | Biolegend | Cat# 114616; RRID:AB_2750196 |
| anti-mouse IA/IE - Brilliant Violet 785™ (clone M5/114.15.2 | Biolegend | Cat# 107645; RRID:AB_2565977 |
| anti-mouse IFNy - PE/Dazzle 594 (clone XMG1.2) | Biolegend | Cat# 505846; RRID:AB_2563980 |
| anti-mouse IFNý - Alexa Fluor® (clone XMG1.2) | Invitrogen | Cat# 53-7311-82; RRID:AB_469932 |
| anti-mouse IL2 - Brilliant Violet 785™ (clone JES6-5H4) | Biolegend | Cat# 503843; RRID:AB_2832801 |
| anti-mouse Ki67 - BV785 (clone B56) | BD Biosciences | Cat# 563756; RRID:AB_2732007 |
| anti-mouse Ki67 - PECy7 (clone B56) | BD Biosciences | Cat# 561283; RRID:AB_10716060 |
| anti-mouse KLRG1 - PECy5 (clone 2F1) | Invitrogen | Cat# 15-5893-80; RRID:AB_2811751 |

| anti-mouse Ly108 - BV650 (clone 13G3) | BD Biosciences | Cat# 740628; RRID:AB_2740323 |
|---|---|---|
| anti-mouse Ly108 - BV786 (clone 13G3) | BD Biosciences | Cat# 741030; RRID:AB_2740648 |
| anti-mouse Ly6C - PE (clone HK1.4) | Biolegend | Cat# 128007; RRID:AB_1186133 |
| anti-mouse MHC I Tetramer H2Kb OVA - PE | MBL | Cat# TB-5001-1 |
| anti-mouse MHC I Tetramer H2Kb TRP2 - PE | MBL | Cat# TB-5004-1 |
| anti-mouse PD1 - Brilliant Violet 711™ (clone 29F.1.A12) | Biolegend | Cat# 135231; RRID:AB_2566158 |
| anti-mouse TCF1 - PE (clone S33-966) | BD Biosciences | Cat# 564217; RRID:AB_2687845 |
| anti-mouse TCR Vβ3 - BV605 (clone KJ25) | BD Biosciences | Cat# 743414; RRID:AB_2741487 |
| anti-mouse TCR Vβ3 - PE (clone KJ25) | BD Biosciences | Cat# 553209; RRID:AB_394709 |
| anti-mouse TIM3 - Brilliant Violet 421™ (clone RMT3-23) | Biolegend | Cat# 119723; RRID:AB_2616908 |
| anti-mouse TNFα - APC (clone MP6-XT22) | Biolegend | Cat# 506308; RRID:AB_315429 |
| anti-mouse TNFα - Brilliant Violet 785™ (clone MP6-XT22) | Biolegend | Cat# 506341; RRID:AB_2565951 |
| anti-mouse TOX - eFluor660 (clone TXRX10) | Invitrogen | Cat# 50-6502-82; RRID:AB_2574265 |
| Biologic Samples | | |
| Mouse tissue samples (brain tumor, spleen) | Duke University | IACUC: A163-21-08 |
| | | |
| Buffers, chemicals, and recombinant proteins | | |
| 0.05% Trypsin-EDTA | GIBCO | Cat# 25300-054 |
| 2-Mercaptoethanol | GIBCO | Cat# 21985023 |
| BD Golgiplug™ (Brefeldin A) | BD Biosciences | Cat# 555029 |
| BD Golgiplug™ (Brefeldin A) | BD Biosciences | Cat# 554724 |
| Concanavalin A (Con A) Solution (500X) | eBioscience | Cat# 00-4978-93 |

| | | |
|---|---|---|
| CountBright Absolute Counting Beads | Invitrogen | Cat# C36950 |
| DMEM | GIBCO | Cat# 11965118 |
| DnaseI | Sigma-Aldrich | Cat# 10104159001 |
| Fetal Bovine Serum (FBS) | Sigma | Cat# F4135 - 500mL |
| Foxp3/Transcription factor staining buffer set | eBioscience | Cat# 00-5523-00 |
| Intracellular Fixation and Permeabilization Set | eBioscience | Cat# 88-8824-00 |
| Ionomycin calcium salt from Streptomyces | Sigma | Cat# I0634-1MG |
| L-glutamine 200mM | GIBCO | Cat# 25030-081 |
| Liberase™ DL Research Grade | Sigma | Cat# 5401160001 |
| Liberase™ TL Research | Sigma | Cat# 5466202001 |
| Non-essential amino acids | GIBCO | Cat# 11140-035 |
| PBS | GIBCO | Cat# 20012-068 |
| Percoll® | Sigma | Cat# P1644-100ML |
| PMA (Phorbol 12-myristate 13-acetate) | Sigma | Cat# P8139-1MG |
| RBC Lysis Buffer | eBioscience | Cat# 00-4300-54 |
| RPMI 1640 | GIBCO | Cat# 21875034 |
| UltraComp eBeads Plus Compensation Beads | Invitrogen | Cat# 01-3333-42 |
| Zombie Aqua™ Viability Stain | Biolegend | Cat# 423101 |
| | | |
| Deposited Data | | |
| Single cell RNA/TCRseq data of T cells from CT2A tumor | This paper | GEO: *in progress, raw and processed data files available upon request* |
| Single cell RNAseq of CD45 cells from CT2A tumor | (Tomaszewski et al. 2022) | GEO: GSE197879 |
| Single cell RNAseq from human newly diagnosed GBM | (Abdelfattah et al., 2022) | GEO: GSE160189 |
| Single cell RNAseq from human metastatic brain melanoma | (Biermann et al., 2022) | GEO: GSE200218 |

| | | |
|---|---|---|
| **Experimental Models: Cell lines** | | |
| CT2A | Massachusetts General Hospital | (Martinez-Murillo et al., 2007) |
| CT2A-TRP2 | Duke University | N/A |
| CT2A-TRP2-β2mKO | Duke University | N/A |
| CT2A-TRP2-OVA | Duke University | N/A |
| CT2A-GFP-LUC | Duke University | (Swan et al., 2023) |
| B16F0 | ATCC | CRL-6322 |
| LLC | ATCC | CRL-1642 |
| E0771 | ATCC | CRL-3461 |
| | | |
| **Experimental Models: Organisms/Strains** | | |
| Mouse: C57BL/6J | Jackson | Strain# 000664; RRID: IMSR_JAX:000664 |
| Mouse: B6.SJL-Ptprc[a] Pepc[b]/BoyJ | Jackson | Strain# 002014; RRID: IMSR_JAX:002014 |
| Mouse: B6.129S4-Ccr2tm1Ifc/J | Jackson | Strain# 004999; RRID: IMSR_JAX:004999 |
| Mouse: B6.129P2-B2mtm1Unc/DcrJ | Jackson | Strain# 002087; RRID: IMSR_JAX:002087 |
| Mouse: B6.FVB-1700016L21RikTg(Itgax-DTR/EGFP)57Lan/J | Jackson | Strain# 004509; RRID: IMSR_JAX:004509 |
| Mouse: C57BL/6-Tg(TcraTcrb)1100Mjb/J | Jackson | Strain# 003831; RRID: IMSR_JAX:003831 |
| | | |
| **Software and Algorithms** | | |
| Flowjo | FlowJo, LLC | https://www.flowjo.com/ |
| R 3.6.1 | CRAN | https://cran.r-project.org/bin/macosx/ |
| RStudio 2022.07.1 | RStudio | https://posit.co/download/rstudio-desktop/ |
| Seurat V4.0.3 | (Stuart et al., 2019) | https://satijalab.org/seurat/ |
| CellChat | (Jin et al., 2021) | https://www.cellchat.org/ |
| CellRanger 4.0.0 | 10X Genomics | https://www.10xgenomics.com/ |
| scRepertoire | (Borcherding et al., 2020) | https://github.com/ncborcherding/scRepertoire |
| fgsea | (Korotkevich et al., 2021) | https://github.com/ctlab/fgsea |

| FIJI (v.2.13.1) | ImageJ | https://imagej.net/software/fiji/; RRID:SCR_002285 |
|---|---|---|
|  |  |  |
| Other Resources |  |  |
| 70 µM cell strainers | Corning Falcon | Cat# 352350 |
| Dounce Tissue Grinders, Wheaton | VWR | Cat# 62400-620 |
|  |  |  |

# RESOURCE AVAILABILITY

**Lead contact**

Further information and requests should be directed to Peter Fecci.

**Materials availability**

Materials generated by this study are available upon request.

**Data and code availability**

Raw FASTQ files and associated single cell RNA and TCR sequencing processed files have been uploaded to the NCBI Gene Expression Omnibus (GEO) database and are awaiting an accession number (raw and processed data files are available upon request). Custom R scripts for TCR, and RNA analysis are also available upon request.

# EXPERIMENTAL MODEL AND SUBJECT DETAILS

**Mice**

All mice were used in accordance with National Institutes of Health and the Duke University Institutional Animal Care and Use Committee guidelines. C57BL/6J (cat #: 664), CCR2KO (cat #: 4999), CD11c-DTR (cat #: 4509), CD45.1 (cat #: 2014), and β2mKO (cat #: 2087) were purchased from Jackson Laboratory at 7-8 weeks.

**Tumor cell lines**

C57BL/6 syngeneic CT2A glioma was provided by Robert L. Martuza (Massachusetts General Hospital)[45]. B16F0 melanoma, E0771 breast cancer, and Lewis lung cancer (LLC) were purchased from ATCC. CT2A-TRP2 was generated by overexpressing a TRP2-containing lentiviral plasmid. CT2A-TRP2-OVA was generated through lentiviral

transduction of parental CT2A-TRP2 with an OVA-containing lentiviral plasmid. CT2A-
GFP-LUC was kindly provided by Alexa Bramall and Nalini Mehta (Duke University)[46].
CT2A-TRP2-β2mKO was made through β2m knockout via CRISPR/Cas9. sgRNA
targeting murine β2m (gRNA sequence: AGUCGUCAGCAUGGCUCGCU) and Cas9
nuclease were purchased from STEMCELL technologies (Cat#: 200-0013, 76002,
respectively). sgRNA (5.8 µg) was incubated at room temperature with 10.8 µg Cas9
protein (STEMCELL Technologies, Cat#: 76002) in Opti-MEM™ + GlutaMAX™
(ThermoFisher Scientific Cat#: 51985034) for 20 min prior to electroporation.
sgRNA/Cas9 was then added to $2.5 \times 10^5$ CT2A-Trp2 cells in Opti-MEM™ + GlutaMAX™.
Cells were electroporated in a 0.2 cm cuvette using a BioRad Gene Pulser XCell™ in
100ul total volume (exponential decay, 1000 uF, 155 V). Cells were FACS isolated after
96 h on $H-2K^b/H-2D^b$ negative cells. CT2A-TRP2-β2MKO cell line was established from
a single cell clone, cultured as described below. CT2A, CT2A-TRP2, CT2A-TRP2-OVA,
CT2A-TRP2- β2mKO, CT2A-GFP, E0771, LLC were cultured in complete DMEM (Gibco
11995-065, 10% FBS). B16F0 was cultured in complete DMEM + 1mM L-glutamine. All
cell lines were authenticated and tested negative for mycoplasma, and interspecies
contamination by IDEXX Laboratories.

# METHOD DETAILS

**Tumor cell line preparation and injection**
All cell lines were cultured at 37˚C in 5% $CO_2$, in their respective media. For tumor
inoculation, all cell lines were taken out of culture using 0.05% Trypsin, washed once with
media and once with PBS. Cells were resuspended in PBS for subcutaneous injections
(200 µl injection volume) and a 1:1 mix of PBS and methylcellulose for intracranial
injections (5 µl injection volume). $1\times10^4$ CT2A, $5\times10^4$ CT2A-TRP2, $5\times10^4$ CT2A-TRP2-
β2mKO, $5\times10^4$ CT2A-TRP2-OVA, $2.5\times10^4$ CT2A-GFP-LUC, $1\times10^3$ B16F0, $1\times10^3$
E0771, and $1\times10^3$ LLC were injected intracranially. $5\times10^5$ CT2A, $2.5\times10^5$ E0771, $1\times10^5$
B16F0 were injected subcutaneously. Mice were monitored and sacrificed in accordance
with Duke University Institutional Animal Care and Use Committee guidelines.

**In vivo antibody administration (aCD8, aCSF1R)**

For macrophage depletion experiments, 400µg aCSF1R (AFS98, BioXCell) was administered 3 times weekly starting 2 days before tumor implantation into CCR2KO mice, as described previously[33].

For CD8[+] T cell depletion, mice received 200ug anti-CD8$\alpha$ (clone 2.43; BioXCell) daily for 3 days prior to tumor implantation, then every 5 days.

**Dendritic cell depletion**

CD11c-DTR or control C57Bl/6J mice were injected intraperitoneally with 5ng/g diptheria toxin (Sigma-Aldrich, D0643) every 3 days, starting D8 after tumor implantation for 4 treatments as previously described[2].

**Microglia depletion**

C57Bl/6J mice were put on PLX3397 diet (290 mg/kg) or a control diet formulated in AIN-76A standard chow as previously described[47].

**Mouse brain tumor tissue processing**

For processing intracranial tumors, brains were harvested as before[29]. In brief, mice were transcardially perfused with 10ml PBS + 1% heparin. The tumor-bearing hemisphere was mechanically dissociated with a dounce tissure homgenizer, then enzymatically dissociated (0.05mg ml$^{-1}$ Liberase DL (Roche), 0.05 mg ml$^{-1}$ Liberase TL (Roche), 0.2 mg ml$^{-1}$ Dnase I (Roche) in HBSS with Ca$^{2+}$ and Mg$^{2+}$) for 15 min at 37 ˚C. The single cell suspension was then washed with PBS and resuspended in RBC lysis buffer (Invitrogen) for 3 min. After quenching the lysis reaction with PBS, the cell suspension was passed through a 70 µm filter. Myelin was removed from the sample with Percoll density gradient centrifugation. Samples were resuspended in 30% Percoll (Sigma-Aldrich) and centrifuged at 500 g for 20 min at 18 ˚C with low brake. The myelin layer was aspirated, and cells were washed with PBS and counted for subsequent flow cytometry staining.

**Flow cytometry**

For cytokine re-stimulation, samples were resuspended in RPMI + 10% FBS at 0.5 - 1 $\times$
$10^7$ cells/ml in a 48 well plate after being counted. PMA (50 ng/ml) + Ionomycin (1000
ng/ml) + GolgiStop™ (BD Biosciences) + GolgiPlug™ (BD Biosciences) were added and
then samples were incubated for 4 h at 37 ˚C.

If no re-stimulation was necessary, samples were resuspended at 1-2 $\times$ $10^7$ cells/ml in
100µl PBS and transferred to a 96-well plate. Cells were resuspended in Zombie Aqua
Viability Dye (1:500, Biolegend) and incubated for 25 min at RT.

For extracellular staining, samples were resuspended in 50 µl TruStain FcX Plus (1:500,
Biolegend) in FACS (PBS + 2% FBS) for 15 min at RT. After blocking, 50 µl of a 2X
antibody mastermix were added and samples were incubated for 25 min at 4 ˚C.  Stained
samples were washed with FACS, then fixed with 2% formaldehyde in PBS for 20 min at
4 ˚C.

For intracellular staining, cells were instead fixed and permeabilized with a 1X FoxP3
Fixation/Perm buffer (eBioscience FoxP3/Transcription Factor Staining Buffer Set,
Thermo Fisher Scientific) for 30 min at 4 ˚C. Intracellular antigens were stained for 30 min
in 1X perm buffer at RT.

Before sample acquisition, 10 µl of Count Bright counting beads (Thermo Fisher
Scientific) were added to each sample. Samples were acquired on an LSRII or Fortessa
(BD Biosciences) using FACS Diva software v.9 and analyzed using FlowJo v.10 (Tree
Star).

**Immunofluorescence**

Brains from PLX3397 treated tumor-bearing mice were harvested after intracardial
perfusion with 10ml 1% PFA and 10ml PBS. Brains were left in 1% PFA for 6 h, then
washed and placed in 10% sucrose for 48h. Once brains had sunk in the solution, they
were washed and snap frozen in tissue freezing media (General Data Company) then
kept at -80 ˚C. Frozen brains were sectioned at 25 µm and mounted onto slides. Sections

were rehydrated with PBS, then permeabilized with Triton-X and washed with PBS.

Permeabilzed sections were blocked using PBS and 2% BSA. Primary and secondary

antibodies were subsequently stained in PBS for 1 h at RT each. Slides were imaged on

a Leica STED confocal microscope (Leica). FIJI was used to process and analyze

images.

*Ex vivo* **TRP2-TCR transgenic T cell generation and adoptive transfer**

Spleens were harvested from CD45.1 mice. Spleens were processed through 70µm filters

then red blood cells were lysed with RBC lysis (Invitrogen). Splenocytes were plated at 2

$\times 10^6$ cells/ml in RPMI + 10% FBS, 50 U/ml IL-2, 2.5 µg/ml Concavalin A, L-glutamine

(Gibco), Pen/Strep (Gibco), 50 µM βME and kept in culture for 48h. Cells were then

retrovirally transduced with the pMX-TRP2- TCRβ-2A-α vector (a kind gift from Dr.

Schumacher, Netherlands Cancer Institute) as described previously[48]. This TCR

recognizes the $TRP2_{180-188}$ epitope in the context of $H-2K^b$. Briefly, HEK293T cells were

transfected with pCL-Eco and the TRP2 vector using Lipofectamine 2000 (Invitrogen).

Transduction was then performed in non-tissue culture treated 24-well plates, previously

coated with Retronectin (25 µg/ml) (Takara) by combining viral supernatant and activated

splenocytes and spinning for 1.5 h at 2000 RPM. Transgenic (Tg) TRP2-TCR T cells were

cultured for another 3 days before use, splitting daily. $10 \times 10^6$ or $2.5 \times 10^5$ Tg TRP2-

TCR T cells were adoptively transferred through tail vein or intracranial injection in 100 µl

PBS or 24 µl PBS, respectively.

OT1 T cells were isolated from OT1 mice by culturing OT1 splenocytes as above for 48

h.

**Chimera Generation**

Bone marrow was harvested from 7 week old donor mice (either C57BL/6J or β2mKO).

Recipient mice (CD45.1) received whole-body irradiation with a 9 Gy dose from a Cesium

irradiator (Mark I-68A [137]Cs irradiator, JL Shepherd and Associates). Recipient mice then

received an intravenous infusion of $10 \times 10^6$ donor bone marrow cells in 100 µl PBS.

Recipient mice were then put on antibiotic-treated water for 2 weeks post bone marrow

transfer.

After 10 weeks of reconstitution, chimerism was validated in the blood, with >95%
chimerism in all queried mice. Then, CT2A-TRP2 or CT2A-TRP2-β2mKO was implanted
intracranially into chimeras. On D7 post-implantation, $10 \times 10^6$ TRP2-TCR T cells were
adoptively transferred intravenously.

**scRNAseq library generation**

$CD3^+CD4^+$ or $CD3^+CD8^+$ were sorted from D13 (n = 16) and D19 (n = 4) intracranial
tumors. Four mice were pooled for each D13 replicate to achieve desired number of cells.
10,000 total cells from each timepoint were targeted with an average read depth of 36,000
reads per cell for the gene expression library. Reads were sequenced on a NovaSeq S1
flow cell. For the TCR library, 9,000 reads per cell were targeted and sequenced on a
NovaSeq SP.

*10x Transcriptome library prep:* GEMs are generated by combining barcoded Single Cell
5' Gel Beads, Master Mix, Partitioning Oil and single cells on a Chromium microfluidic
Chip. Immediately following GEM generation, the Gel Bead is dissolved and any co-
partitioned cell is lysed. Oligonucleotides containing an Illumina R1 sequence (read 1
sequencing primer), a 16 nt 10x Barcode, a 10 nt unique molecular identifier (UMI), and
13 nt template switch oligo (TSO) are released and mixed with the cell lysate and a Master
Mix containing reverse transcription (RT) reagents and poly(dT) RT primers. Incubation
of the GEMs produces 10x Barcoded, full-length cDNA from polyadenylated mRNA.

GEMs are then broken and pooled and the 10x Barcoded first-strand cDNA is purified
from the post GEM-RT reaction mixture and amplified. Full-length V(D)J segments (10x
Barcoded) are enriched from amplified cDNA via PCR amplification with universal primer
for 5' adapter and successive nested primers specific to the TCR and/or BCR constant
regions. Enrichment and library preparation of the TCR and BCR constant regions takes
place in two separate workflows using the same amplified cDNA stock.

Enzymatic fragmentation and size selection are used to generate variable length
fragments that collectively span the V(D)J segments of the enriched TCR and/or BCR

transcripts prior to library construction. P5, P7, a dual sample index, and an Illumina R2

sequence (read 2 primer sequence) are added via End Repair, A-tailing, Adaptor Ligation,

and Sample Index PCR. The final libraries contain the P5 and P7 priming sites used in

Illumina sequencing. Sequence is generated using Illumina short read paired end

sequencing.

**scRNAseq data processing and analysis**

The Cell Ranger pipeline (10X Genomics) was used to perform sample demultiplexing

and to generate FASTQ files for the gene expression and ADT libraries. FASTQ files were

demultiplexed from the raw sequencing reads (bcl2fastq, v2.20), aligned to the mouse

mm10 reference genome (cellranger, v4.0.0), and raw gene count matrices were

generated using STAR (v2.7.5c).

Downstream    analysis    was    performed    using    the    R    software    Seurat

package[49] (v4.0.3, http://satijalab.org/seurat/). Hashtag oligos, which corresponded to

biological replicates, were demultiplexed using the HTODemux function and appended to

the meta-data for each sample. Low-quality cells, expressing less than 200 genes, and

genes expressed by fewer than three cells were removed. The gene expression matrix

for each genotype was then concatenated using the merge function in Seurat. The

percentage    of    mitochondrial    gene    content    was    calculated    using    the

Mouse.MitoCarta3.0[50] gene set and the PercentageFeatureSet function in Seurat. The

Seurat object was converted to a SingleCellExperiment object, and outlier exclusion was

performed in scater (v.1.18.6). Using the isOutlier function in scater, cells were discarded

if their percentage of mitochondrial gene content, number of expressed genes, or number

of reads for a given cell was considered an outlier. A total of 6,742 CD8$^+$ cells passed QC

and the SingleCellExperiment object was then converted back into a Seurat object.

Normalization and regression of cell cycle scoring and percent mitochondrial gene

content were performed using the SCTransform function in Seurat. Principal component

analysis (PCA) was performed on all genes, and the number of principal components to

be utilized in further analysis was determined heuristically using the elbow method.

Eleven principal components were used for clustering and dimensionality reduction using

FindNeighbors, RunUMAP, and a resolution of 0.5 was used for FindClusters in Seurat.

This approach identified six distinct cell clusters. All subsequent analyses were conducted

with this cleaned object.

For TCRSeq analysis, scRepertoire standard workflow was utilized[51]. In brief filtered

contig outputs from the 10X Genomics Cell Ranger Pipeline were loaded into R (v4.1.2)

and combined into a list using combineTCR. Next combineExpression was used to attach

clonotypic information to the Seurat object. Visualizations were made using the

scRepertoire vignette.

For exhaustion signature score generation, the built in AddModuleScore function from

Seurat was utilized with a gene list from Im et al. 2016[19]. A detailed list can be found in

Table S1. For pseudotime analysis, the object was passed into Monocle v3 without any

changes to the UMAP dimensional reduction.

**Publicly available scRNA-seq data analysis**

Human newly diagnosed GBM and metastatic brain melanoma datasets were

downloaded from the GEO (accession # :GSE160189, GSE200218). Datasets were QC'd

as above, removing cells with high mitochondrial percentage and those that lacked

expression of PTPRC1 (non-immune cells). Then datasets were integrated using the

standard Seurat protocol[52].

**Cellchat Analysis Mouse/Human**

For murine CellChat analysis utilizing the CD45+ sorted dataset generated by

Tomaszewski et al[29], we subsetted the object to keep CD8 TIL, Apoe+ TAM, Nos2+ TAM,

DC1, DC2, DC3, pDC and microglia from the WT sample. We then fed this cleaned object

into the CellChat pipeline with built in murine ligand-receptor database. Following the

recommended vignette, we visualized the enrichment of the MHC-I pathway across cell

types. Additionaly we visualized the strength and number of interactions with CD8 TIL as

the target cells.

Human CellChat analysis followed a similar workflow, except the human ligand-receptor

database was utilized. Additionally, integrated ndGBM and MBM was subsetted so samples of the same tumor type were analyzed separately, as suggested by the CellChat vignette.

# QUANTIFICATION AND STATISTICAL ANALYSIS

**Statistical Analysis**

Unless specifically noted, all data are representative of ≥2 separate experiments. Experimental group assignment was determined by random designation. Statistical analyses were performed using GraphPad Prism software. Error bars represent ± Standard Error of the mean (SEM) calculated using Prism. Specific statistical tests used were paired t-tests, unpaired t-tests, and one/two-way ANOVA unless otherwise noted. P-values <0.05 were considered statistically significant. Investigators were not blinded to group assignment during experimental procedures or analysis.

# SUPPLEMENTAL INFORMATION TITLES AND METHODS

Supplemental Figure 1:

(A) Gating strategy for Tex_prog and Tex_term.

(B) Frequency of Tex subsets of PD1+CD44+ CD8+ T cells in intracranial B16F0, LLC or CT2A. n = 8-10 mice/group.

All of the data are means ± SEMs; p-values are shown on individual plots as determined by 1-way ANOVA and post-hoc Tukey test (B), two-sided paired t-test (B).

Supplemental Figure 2:

(A) Dot plot of top 3 genes for each cluster.

(B) Violin plot of *Tcf7, Pdcd1, Tox,* and *Havcr2* expression across clusters.

(C) UMAP of cloneType across biological replicates (n=4 per timepoint).

(D) Shannon and Inverse Simpson diversity indices across clusters, calculated with the scRepertoire package.

(E) 10 most expanded TCR clones across timepoints are concentrated to Tex_int and
Tex_term clusters.

Supplemental Figure 3:

(A) PD1 expression on transferred OT1 vs. TRP2-TCR Tg T cells on day of ALT (D7) and
day of tumor harvest (D21) and representative histogram. n = 7 mice/group.

(B) Frequency of Tex subsets of PD1+ endogenous or transgenic TRP2-TCR T cells and
representative contour plot. n = 7 mice.

(C) gMFI of MHC I in CT2A-TRP2 vs. CT2A-TRP2-B2mKO validating knockout of MHC
I. n = 10 mice/group.

(D) Number of TRP2-TCR T cells/g of tumor in CT2A-TRP2 vs. CT2A-TRP2-B2mKO. n
= 10 mice/group.

p-values are plotted on individual graphs. Statistical significance was determined by two-
sided unpaired t-test (A, C, D) or 2-way RM ANOVA with post-hoc Tukey's test for multiple
comparisons (B). All data are plotted as mean ± SEM. See also Figure S4.

Supplemental Figure 4:

(A) Interaction strength across APC with CD8 TIL as the target.

(B) Gating strategy for intratumoral APC.

Supplemental Figure 5:

(A) Frequency of TAM of CD45$^+$ cells. Data pooled from 2 independent experiments. n =
11 mice/group.

(B) Experimental layout. DC were depleted in CD11c-DTR mice with 5ng/g DT, every 3
days for 4 total treatments, starting D8. Created with Biorender.com.

(C) Frequency of CD11c+CD11b- DC in WT vs. CD11c-DTR mice.

(D) Experimental layout. Microglia were depleted in WT mice with 270mg/kg PLX3397

chow. Chow was provided starting D(-)14. Created with Biorender.com.

(E) Representative immunofluorescence image of WT vs. PLX3397 treated non-tumor

brain. Scale bar = 100um.

(F) Chimera validation. Frequency of MHC I+ in WT $\rightarrow$ WT (WT) or B2mKO $\rightarrow$ WT

(B2mKO) chimeras across T cells, TAM and cDC. n = 8-10 mice/group.

(G) Representative histogram of MHC I expression on TAM in the setting of CT2A-TRP2

(Teal) vs. CT2A-TRP2-B2mKO (Dark Teal).

(H) Frequency of TAM of CD45$^+$ cells in B16F0 IC. n = 9-10 mice/group.

(I) Frequency of TAM of CD45$^+$ cells in B16F0 SQ. Data pooled from 2 independent

experiments. n = 6-7 mice/group.

p-values are plotted on individual graphs. Statistical significance was determined by two-

sided unpaired t test (A, H, I) or RM-2-way ANOVA with post-hoc Tukey's test for multiple

comparisons (F). All data are plotted as mean ± SEM.

References:

1. Nolan, C., and Deangelis, L.M. (2018). Overview of metastatic disease of the central nervous system. Handb Clin Neurol *149*, 3-23. 10.1016/B978-0-12-811161-1.00001-3.

2. Jung, S., Unutmaz, D., Wong, P., Sano, G., De los Santos, K., Sparwasser, T., Wu, S., Vuthoori, S., Ko, K., Zavala, F., et al. (2002). In vivo depletion of CD11c+ dendritic cells abrogates priming of CD8+ T cells by exogenous cell-associated antigens. Immunity *17*, 211-220. 10.1016/s1074-7613(02)00365-5.

3. Amin, S., Baine, M.J., Meza, J.L., and Lin, C. (2020). Association of Immunotherapy With Survival Among Patients With Brain Metastases Whose Cancer Was Managed With Definitive Surgery of the Primary Tumor. JAMA Netw Open *3*, e2015444. 10.1001/jamanetworkopen.2020.15444.

4. Brastianos, P.K., Curry, W.T., and Oh, K.S. (2013). Clinical discussion and review of the management of brain metastases. J Natl Compr Canc Netw *11*, 1153-1164. 10.6004/jnccn.2013.0133.

5. Reardon, D.A., Brandes, A.A., Omuro, A., Mulholland, P., Lim, M., Wick, A., Baehring, J., Ahluwalia, M.S., Roth, P., Bahr, O., et al. (2020). Effect of Nivolumab vs Bevacizumab in Patients With Recurrent Glioblastoma: The CheckMate 143 Phase 3 Randomized Clinical Trial. JAMA Oncol *6*, 1003-1010. 10.1001/jamaoncol.2020.1024.

6. Woroniecka, K., Chongsathidkiet, P., Rhodin, K., Kemeny, H., Dechant, C., Farber, S.H., Elsamadicy, A.A., Cui, X., Koyama, S., Jackson, C., et al. (2018). T-Cell Exhaustion Signatures Vary with Tumor Type and Are Severe in Glioblastoma. Clin Cancer Res *24*, 4175-4186. 10.1158/1078-0432.CCR-17-1846.

7. Woroniecka, K.I., Rhodin, K.E., Chongsathidkiet, P., Keith, K.A., and Fecci, P.E. (2018). T-cell Dysfunction in Glioblastoma: Applying a New Framework. Clin Cancer Res *24*, 3792-3802. 10.1158/1078-0432.ccr-18-0047.

8. Akbay, E.A., Koyama, S., Carretero, J., Altabef, A., Tchaicha, J.H., Christensen, C.L., Mikse, O.R., Cherniack, A.D., Beauchamp, E.M., Pugh, T.J., et al. (2013). Activation of the PD-1 pathway contributes to immune escape in EGFR-driven lung tumors. Cancer discovery *3*, 1355-1363. 10.1158/2159-8290.CD-13-0310.

9. Koyama, S., Akbay, E.A., Li, Y.Y., Herter-Sprie, G.S., Buczkowski, K.A., Richards, W.G., Gandhi, L., Redig, A.J., Rodig, S.J., Asahina, H., et al. (2016). Adaptive resistance to therapeutic PD-1 blockade is associated with upregulation of alternative immune checkpoints. Nat Commun *7*, 10501. 10.1038/ncomms10501.

10. Huang, R.Y., Francois, A., McGray, A.R., Miliotto, A., and Odunsi, K. (2017). Compensatory upregulation of PD-1, LAG-3, and CTLA-4 limits the efficacy of single-agent checkpoint blockade in metastatic ovarian cancer. Oncoimmunology *6*, e1249561. 10.1080/2162402X.2016.1249561.

11. Shayan, G., Srivastava, R., Li, J., Schmitt, N., Kane, L.P., and Ferris, R.L. (2017). Adaptive resistance to anti-PD1 therapy by Tim-3 upregulation is mediated by the PI3K-Akt pathway in head and neck cancer. Oncoimmunology *6*, e1261779. 10.1080/2162402X.2016.1261779.

12. Thommen, D.S., Schreiner, J., Muller, P., Herzig, P., Roller, A., Belousov, A., Umana, P., Pisa, P., Klein, C., Bacac, M., et al. (2015). Progression of Lung Cancer Is Associated with

Increased Dysfunction of T Cells Defined by Coexpression of Multiple Inhibitory
Receptors. Cancer Immunol Res *3*, 1344-1355. 10.1158/2326-6066.CIR-15-0097.

13.   Dunn, G.P., Fecci, P.E., and Curry, W.T. (2012). Cancer immunoediting in malignant
glioma. Neurosurgery *71*, 201-222; discussion 222-203.
10.1227/NEU.0b013e31824f840d.

14.   Wherry, E.J., Ha, S.J., Kaech, S.M., Haining, W.N., Sarkar, S., Kalia, V., Subramaniam, S.,
Blattman, J.N., Barber, D.L., and Ahmed, R. (2007). Molecular signature of CD8+ T cell
exhaustion during chronic viral infection. Immunity *27*, 670-684.
10.1016/j.immuni.2007.09.006.

15.   Kallies, A., Zehn, D., and Utzschneider, D.T. (2020). Precursor exhausted T cells: key to
successful immunotherapy? Nat Rev Immunol *20*, 128-136. 10.1038/s41577-019-0223-7.

16.   Utzschneider, D.T., Gabriel, S.S., Chisanga, D., Gloury, R., Gubser, P.M., Vasanthakumar,
A., Shi, W., and Kallies, A. (2020). Early precursor T cells establish and propagate T cell
exhaustion in chronic infection. Nat Immunol *21*, 1256-1266. 10.1038/s41590-020-0760-
z.

17.   Angelosanto, J.M., Blackburn, S.D., Crawford, A., and Wherry, E.J. (2012). Progressive loss
of memory T cell potential and commitment to exhaustion during chronic viral infection.
J Virol *86*, 8161-8170. 10.1128/JVI.00889-12.

18.   Woroniecka, K., Chongsathidkiet, P., Rhodin, K.E., Kemeny, H.R., Dechant, C.A., Farber,
S.H., Elsamadicy, A.A., Cui, X., Koyama, S., Jackson, C.C., et al. (2018). T Cell Exhaustion
Signatures Vary with Tumor Type and are Severe in Glioblastoma. Clin Cancer Res.
10.1158/1078-0432.ccr-17-1846.

19.   Im, S.J., Hashimoto, M., Gerner, M.Y., Lee, J., Kissick, H.T., Burger, M.C., Shan, Q., Hale,
J.S., Lee, J., Nasti, T.H., et al. (2016). Defining CD8+ T cells that provide the proliferative
burst after PD-1 therapy. Nature *537*, 417-421. 10.1038/nature19330.

20.   Utzschneider, D.T., Charmoy, M., Chennupati, V., Pousse, L., Ferreira, D.P., Calderon-
Copete, S., Danilo, M., Alfei, F., Hofmann, M., Wieland, D., et al. (2016). T Cell Factor 1-
Expressing Memory-like CD8(+) T Cells Sustain the Immune Response to Chronic Viral
Infections. Immunity *45*, 415-427. 10.1016/j.immuni.2016.07.021.

21.   Miller, B.C., Sen, D.R., Al Abosy, R., Bi, K., Virkud, Y.V., LaFleur, M.W., Yates, K.B., Lako, A.,
Felt, K., Naik, G.S., et al. (2019). Author Correction: Subsets of exhausted CD8(+) T cells
differentially mediate tumor control and respond to checkpoint blockade. Nat Immunol
*20*, 1556. 10.1038/s41590-019-0528-5.

22.   Sade-Feldman, M., Yizhak, K., Bjorgaard, S.L., Ray, J.P., de Boer, C.G., Jenkins, R.W., Lieb,
D.J., Chen, J.H., Frederick, D.T., Barzily-Rokni, M., et al. (2018). Defining T Cell States
Associated with Response to Checkpoint Immunotherapy in Melanoma. Cell *175*, 998-
1013 e1020. 10.1016/j.cell.2018.10.038.

23.   Schietinger, A., Philip, M., Krisnawan, V.E., Chiu, E.Y., Delrow, J.J., Basom, R.S., Lauer, P.,
Brockstedt, D.G., Knoblaugh, S.E., Hammerling, G.J., et al. (2016). Tumor-Specific T Cell
Dysfunction Is a Dynamic Antigen-Driven Differentiation Program Initiated Early during
Tumorigenesis. Immunity *45*, 389-401. 10.1016/j.immuni.2016.07.011.

24.   McLane, L.M., Abdel-Hakeem, M.S., and Wherry, E.J. (2019). CD8 T Cell Exhaustion During
Chronic Viral Infection and Cancer. Annu Rev Immunol *37*, 457-495. 10.1146/annurev-
immunol-041015-055318.

25. Connolly, K.A., Kuchroo, M., Venkat, A., Khatun, A., Wang, J., William, I., Hornick, N.I.,
Fitzgerald, B.L., Damo, M., Kasmani, M.Y., et al. (2021). A reservoir of stem-like CD8(+) T
cells in the tumor-draining lymph node preserves the ongoing antitumor immune
response. Sci Immunol *6*, eabg7836. 10.1126/sciimmunol.abg7836.

26. Prokhnevska, N., Cardenas, M.A., Valanparambil, R.M., Sobierajska, E., Barwick, B.G.,
Jansen, C., Reyes Moon, A., Gregorova, P., delBalzo, L., Greenwald, R., et al. (2023). CD8(+)
T cell activation in cancer comprises an initial activation phase in lymph nodes followed
by effector differentiation within the tumor. Immunity *56*, 107-124 e105.
10.1016/j.immuni.2022.12.002.

27. Postow, M.A., Manuel, M., Wong, P., Yuan, J., Dong, Z., Liu, C., Perez, S., Tanneau, I., Noel,
M., Courtier, A., et al. (2015). Peripheral T cell receptor diversity is associated with clinical
outcomes following ipilimumab treatment in metastatic melanoma. J Immunother Cancer
*3*, 23. 10.1186/s40425-015-0070-4.

28. Klemm, F., Maas, R.R., Bowman, R.L., Kornete, M., Soukup, K., Nassiri, S., Brouland, J.P.,
Iacobuzio-Donahue, C.A., Brennan, C., Tabar, V., et al. (2020). Interrogation of the
Microenvironmental Landscape in Brain Tumors Reveals Disease-Specific Alterations of
Immune Cells. Cell *181*, 1643-1660 e1617. 10.1016/j.cell.2020.05.007.

29. Tomaszewski, W.H., Waibl-Polania, J., Chakraborty, M., Perera, J., Ratiu, J., Miggelbrink,
A., McDonnell, D.P., Khasraw, M., Ashley, D.M., Fecci, P.E., et al. (2022). Neuronal CaMKK2
promotes immunosuppression and checkpoint blockade resistance in glioblastoma. Nat
Commun *13*, 6483. 10.1038/s41467-022-34175-y.

30. Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.H., Myung, P., Plikus,
M.V., and Nie, Q. (2021). Inference and analysis of cell-cell communication using CellChat.
Nat Commun *12*, 1088. 10.1038/s41467-021-21246-9.

31. Joffre, O.P., Segura, E., Savina, A., and Amigorena, S. (2012). Cross-presentation by
dendritic cells. Nat Rev Immunol *12*, 557-569. 10.1038/nri3254.

32. Boring, L., Gosling, J., Chensue, S.W., Kunkel, S.L., Farese, R.V., Jr., Broxmeyer, H.E., and
Charo, I.F. (1997). Impaired monocyte migration and reduced type 1 (Th1) cytokine
responses in C-C chemokine receptor 2 knockout mice. J Clin Invest *100*, 2552-2561.
10.1172/JCI119798.

33. MacDonald, K.P., Palmer, J.S., Cronau, S., Seppanen, E., Olver, S., Raffelt, N.C., Kuns, R.,
Pettit, A.R., Clouston, A., Wainwright, B., et al. (2010). An antibody against the colony-
stimulating factor 1 receptor depletes the resident subset of monocytes and tissue- and
tumor-associated macrophages but does not inhibit inflammation. Blood *116*, 3955-3963.
10.1182/blood-2010-02-266296.

34. Miyake, K., and Karasuyama, H. (2021). The Role of Trogocytosis in the Modulation of
Immune Cell Functions. Cells *10*. 10.3390/cells10051255.

35. Abdelfattah, N., Kumar, P., Wang, C., Leu, J.S., Flynn, W.F., Gao, R., Baskin, D.S.,
Pichumani, K., Ijare, O.B., Wood, S.L., et al. (2022). Single-cell analysis of human glioma
and immune cells identifies S100A4 as an immunotherapy target. Nat Commun *13*, 767.
10.1038/s41467-022-28372-y.

36. Biermann, J., Melms, J.C., Amin, A.D., Wang, Y., Caprio, L.A., Karz, A., Tagore, S., Barrera,
I., Ibarra-Arellano, M.A., Andreatta, M., et al. (2022). Dissecting the treatment-naive

ecosystem of human melanoma brain metastasis. Cell *185*, 2591-2608 e2530.
10.1016/j.cell.2022.06.007.

37.  Salmon, H., Remark, R., Gnjatic, S., and Merad, M. (2019). Host tissue determinants of
tumour immunity. Nat Rev Cancer *19*, 215-227. 10.1038/s41568-019-0125-9.

38.  Karimi, E., Yu, M.W., Maritan, S.M., Perus, L.J.M., Rezanejad, M., Sorin, M., Dankner, M.,
Fallah, P., Dore, S., Zuo, D., et al. (2023). Single-cell spatial immune landscapes of primary
and metastatic brain tumours. Nature *614*, 555-563. 10.1038/s41586-022-05680-3.

39.  Jansen, C.S., Prokhnevska, N., Master, V.A., Sanda, M.G., Carlisle, J.W., Bilen, M.A.,
Cardenas, M., Wilkinson, S., Lake, R., Sowalsky, A.G., et al. (2019). An intra-tumoral niche
maintains and differentiates stem-like CD8 T cells. Nature *576*, 465-470. 10.1038/s41586-
019-1836-5.

40.  Blackburn, S.D., Shin, H., Haining, W.N., Zou, T., Workman, C.J., Polley, A., Betts, M.R.,
Freeman, G.J., Vignali, D.A., and Wherry, E.J. (2009). Coregulation of CD8+ T cell
exhaustion by multiple inhibitory receptors during chronic viral infection. Nat Immunol
*10*, 29-37. 10.1038/ni.1679.

41.  Engelhardt, J.J., Boldajipour, B., Beemiller, P., Pandurangi, P., Sorensen, C., Werb, Z.,
Egeblad, M., and Krummel, M.F. (2012). Marginating dendritic cells of the tumor
microenvironment cross-present tumor antigens and stably engage tumor-specific T cells.
Cancer Cell *21*, 402-417. 10.1016/j.ccr.2012.01.008.

42.  Broz, M.L., Binnewies, M., Boldajipour, B., Nelson, A.E., Pollack, J.L., Erle, D.J., Barczak, A.,
Rosenblum, M.D., Daud, A., Barber, D.L., et al. (2014). Dissecting the Tumor Myeloid
Compartment Reveals Rare Activating Antigen-Presenting Cells Critical for T Cell
Immunity. Cancer Cell *26*, 938. 10.1016/j.ccell.2014.11.010.

43.  Kersten, K., Hu, K.H., Combes, A.J., Samad, B., Harwin, T., Ray, A., Rao, A.A., Cai, E.,
Marchuk, K., Artichoker, J., et al. (2022). Spatiotemporal co-dependency between
macrophages and exhausted CD8(+) T cells in cancer. Cancer Cell *40*, 624-638 e629.
10.1016/j.ccell.2022.05.004.

44.  Kilian, M., Sheinin, R., Tan, C.L., Friedrich, M., Kramer, C., Kaminitz, A., Sanghvi, K., Lindner,
K., Chih, Y.C., Cichon, F., et al. (2023). MHC class II-restricted antigen presentation is
required to prevent dysfunction of cytotoxic T cells by blood-borne myeloids in brain
tumors. Cancer Cell *41*, 235-251 e239. 10.1016/j.ccell.2022.12.007.

45.  Martinez-Murillo, R., and Martinez, A. (2007). Standardization of an orthotopic mouse
brain tumor model following transplantation of CT-2A astrocytoma cells. Histol
Histopathol *22*, 1309-1326. 10.14670/HH-22.1309.

46.  Swan, S.L., Mehta, N., Ilich, E., Shen, S.H., Wilkinson, D.S., Anderson, A.R., Segura, T.,
Sanchez-Perez, L., Sampson, J.H., and Bellamkonda, R.V. (2023). IL7 and IL7 Flt3L co-
expressing CAR T cells improve therapeutic efficacy in mouse EGFRvIII heterogeneous
glioblastoma. Front Immunol *14*, 1085547. 10.3389/fimmu.2023.1085547.

47.  Elmore, M.R., Najafi, A.R., Koike, M.A., Dagher, N.N., Spangenberg, E.E., Rice, R.A.,
Kitazawa, M., Matusow, B., Nguyen, H., West, B.L., and Green, K.N. (2014). Colony-
stimulating factor 1 receptor signaling is necessary for microglia viability, unmasking a
microglia progenitor cell in the adult brain. Neuron *82*, 380-397.
10.1016/j.neuron.2014.02.040.

48.    Bendle, G.M., Linnemann, C., Hooijkaas, A.I., Bies, L., de Witte, M.A., Jorritsma, A., Kaiser,
1028            A.D., Pouw, N., Debets, R., Kieback, E., et al. (2010). Lethal graft-versus-host disease in
1029            mouse models of T cell receptor gene therapy. Nat Med *16*, 565-570, 561p following 570.
1030            10.1038/nm.2128.

1031    49.    Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-
1032            cell transcriptomic data across different conditions, technologies, and species. Nat
1033            Biotechnol *36*, 411-420. 10.1038/nbt.4096.

1034    50.    Rath, S., Sharma, R., Gupta, R., Ast, T., Chan, C., Durham, T.J., Goodman, R.P., Grabarek,
1035            Z., Haas, M.E., Hung, W.H.W., et al. (2021). MitoCarta3.0: an updated mitochondrial
1036            proteome now with sub-organelle localization and pathway annotations. Nucleic Acids
1037            Res *49*, D1541-D1547. 10.1093/nar/gkaa1011.

1038    51.    Borcherding, N., Bormann, N.L., and Kraus, G. (2020). scRepertoire: An R-based toolkit for
1039            single-cell immune receptor analysis. F1000Res *9*, 47. 10.12688/f1000research.22139.2.

1040    52.    Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao,
1041            Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-
1042            Cell Data. Cell *177*, 1888-1902 e1821. 10.1016/j.cell.2019.05.031.

1043

# Main Figures:

Antigen-presentation by tumor-associated macrophages mediates progenitor to terminal exhaustion transition in GBM and other solid tumors

**Figure 1: GBM TIL demonstrate a severe decline in the Progenitor Exhaustion to Terminal Exhaustion (PETE) Ratio over time**

(A) Representative contour plots of Tex_prog and Tex_term across tumor types.

(B) PETER across CT2A, B16F0 and LLC intracranial tumors, harvested when mice were moribund. n = 8-10 mice/group.

(C and D) Geometric mean fluorescence intensity (gMFI) of intracellular TOX expression (C) and quantification of TNFa/IFNy/IL2 of Tex_prog and Tex_term in 10K CT2A at D18 (D). n = 5 mice. Representative of two independent experiments.

(E) Experimental layout. 10K CT2A cells implanted D0, tumors were harvested every 3 days starting D9 until mice were moribund on D21. Created with Biorender.

(F) number of CD8+ T cells per harvested tumor. n = 5 mice/timepoint.

(G) Frequency of PD1+CD44+ of CD8+ T cells. n = 5 mice/timepoint

(H) Geometric mean fluorescence intensity (gMFI) of TOX of Tex_prog and Tex_term across timepoints with >100 total PD1+CD44+ CD8+ T cells. n = 5 mice/timepoint

(I and J) Representative plots showing Ly108 vs. TIM3 progression from D9 to D21, Tex_prog = top left, Tex_int = bottom left, Tex_term = bottom right (I) and quantification of Tex subsets of PD1+CD44+ CD8+ T cells over time (J). n = 5 mice/timepoint

All of the data are means ± SEMs; p-values are shown on individual plots as determined by 1-way ANOVA and post-hoc Tukey test (B. F. G), two-sided paired t-test (C) or RM 2-way ANOVA and post-hoc Tukey test (D and H). See also Figure S1.

**Figure 2: scRNAseq defines transcriptional heterogeneity amongst tumor-infiltrating Tex_prog and Tex_term with an enrichment for cell-cell communication pathways in Tex_prog**

**Figure 2: scRNAseq defines transcriptional heterogeneity amongst tumor-infiltrating Tex_prog and Tex_term with an enrichment for cell-cell communication pathways in Tex_prog**

(A) Experimental layout for scRNAseq experiment. tumors were harvested at an early (D13) and late (D19) timepoint. Created with Biorender.

(B) UMAP of 6 CD8 T cell clusters following QC.

(C) Frequency of each cluster per timepoint.

(D) Heatmap of exhaustion associated transcription factors, cytokines/secreted molecules, and surface markers.

(E) AddModuleScore of exhaustion signature from Im et al., 2016, violin plot and density plot representations.

(F) Significantly upregulated pathways in Tex_prog and Tex_term from Hallmark, Reactome, Gene Ontology databases.

(G) Relevant genes from upregulated pathways in H.

(H) Trajectory analysis overlaid onto clusters from B.

(I) Trajectory analysis overlaid onto pseudotime.

(J) Circle plot of overlapping clones across Tex_clusters. Tex_prog, Tex_int have overlapping clones with all other clusters. Tex_term and Tex_int_2 do not have overlapping clones.

(K) UMAP of cloneTypes across clusters.

(L) Frequency of cloneTypes across clusters.

See also Figure S2.

**Figure 3: Tumor cells are not a necessary source of the TCR stimulation mediating Tex_prog to Tex_term transition**

(A) Experimental layout. 250K *ex-vivo* activated and transduced OT1 and TRP2-TCR T cells were intracranially implanted into D7 CT2A-TRP2-bearing mice. Tumors were harvested 14 days later.

(B) Frequency of implanted cells recovered on D21. n = 7 mice/group.

(C and D) Frequency of PD1+ CD8+ T cells in OT1 vs. TRP2-TCR (C) and representative contour plot PD1 and TIM3 expression of OT1 and TRP2-TCR T cells 14 days post-transfer (D). n = 7 mice/group

(E) Experimental layout. 50K CT2A-TRP2 or CT2A-TRP2-B2mKO were implanted intracranially and harvested when mice were moribund. Created with Biorender.

(F) Frequency of PD1+ TRP2-TCR T cells between groups. n = 10 mice/group.

(G and H) Frequency of Tex subset of PD1+TRP2-Tetramer+ CD8+ T cells and representative contour plot (G) and representative contour plot of Ly108 and TIM3 expression between groups (H). n = 10 mice/group.

p-values are shown on individual plots as measured by two tailed t-test (B, C, F) or RM 2-way ANOVA (G). See also Figure S3.

**Figure 4: TAM are the most frequent and most potent cross-presenting antigen-presenting cells (APC) within the intracranial TME.**

(A) Experimental layout. CD45+ sorted scRNAseq from Tomaszweski et al., 2022 was analyzed with CellChat to determine which APC had predicted MHC I interactions with CD8 T cells. Created with Biorender. ## = reference

(B) Heatmap showing communication probability of MHC I interactions in the TME. All APC were predicted to interact with CD8 TIL through MHC I.

(C) Circle plot displaying number of interactions (thickness of arrow/number) across APC populations.

(D) Frequency of infiltrating immune cells from Tomaszewski et al., 2022 dataset.

(E) Experimental layout. 25K CT2A-GFP was implanted intracranially and harvested when mice were moribund. GFP+ APC were evaluated. Created with Biorender.

(F) Frequency of GFP+ T cells (CD3+), TAM (CD45hiCD11b+F4/80+), DC (CD11b+MHC II+F4/80-) and Microglia (CD45hiCD11b+F4/80+). n = 5 mice.

(G and H) gMFI of MHC I of APC populations from F (G) and representative histogram (H). n = 5 mice.

(I) Experimental layout. 100K CT2A-OVA was implanted intracranially and harvested when mice were moribund.

(J) Frequency of SIINFEKL-loaded MHC I of T cells, TAM, Monoctyes (CD11b+F480intLy6Chi), DC, and Microglia. n = 9 mice.

p-values are plotted on individual graphs. Statistical significance was determined by 1-way ANOVA (F, G, J) with post-hoc Tukey's test for multiple comparisons. All data are plotted as mean ± SEM. See also Figure S4.

**Figure 5: TAM mediate progenitor to terminal exhaustion transition in GBM and other intracranial solid tumor models**

(A) Experimental layout. TAM and monocytes were depleted in CCR2KO mice with aCSF1R depletion antibody. 10K CT2A was implanted intracranially and harvested when mice were moribund. Created with Biorender.

(B and C) Frequency of Tex_subsets of PD1+CD44+ (B) and PETER between WT and TAM depleted groups (C). Data pooled from 2 independent experiments. n = 10-11 mice/group.

(D) Simple linear correlation of %Tex_term of PD1+CD44+ CD8+ T cells vs. % TAM of CD45+. Data pooled from 2 independent experiments. n = 11 mice/group.

(E and F) CD11c+ DC were depleted with 5ng/g DTR injections on D8, D11, D14, D17 post CT2A-implantation. Frequency of T cell exhaustion subsets (E) and and PETER between WT and DC depleted groups (F). n = 5 mice/group.

(G and H) Microglia were depleted with 290mg/kg PLX3397 free-fed chow starting 1 week prior to CT2A implantation. Frequency of T cell exhaustion subsets (G) and PETER between WT and microglia depleted groups (H). n = 10 mice/group.

(I) Simple linear correlation of frequency of Tex_term vs. Frequency of MHC I+ TAM in B2mKO chimera mice implanted with CT2A.

(J) Experimental layout. TAM and monocytes were depleted in CCR2KO mice with aCSF1R depletion antibody. 1K B16F0 was implanted intracranially or 100K B16F0 was implanted subcutaneously and harvested when mice were moribund. Created with Biorender.

(K and L) Frequency of Tex_subsets of PD1+CD44+ B16F0 IC (K) and PETER between WT and TAM depleted groups (L). n = 9-10 mice/group.

(M and N) Frequency of Tex_subsets of PD1+CD44+ B16F0 SQ (M) and PETER between WT and TAM depleted groups (N). Data pooled from 2 independent experiments. n = 6-7 mice/group.

p-values are plotted on individual graphs. Statistical significance was determined by two-sided unpaired t test (C, F, H, L, N) or RM-2-way ANOVA with post-hoc Tukey's test for multiple comparisons (B, E, G, K, M). All data are plotted as mean ± SEM. See also Figure S5.

**Figure 6: CellChat predicts secondary interactions between TAM and CD8+ T cells in mice and humans that may contribute to exhaustion progression**

**Figure 6: CellChat predicts secondary interactions between TAM and CD8+ T cells in mice and humans that may contribute to exhaustion progression**

(A) Experimental layout. CD45+ sorted scRNAseq from murine GBM, ndGBM and MBC was analyzed with CellChat to determine additional TAM-T cell interactions. Created with Biorender.

(B) Enriched predicted interactions with TAM (orange) as the source and CD8 TIL (blue) as the target from murine GBM.

(C) Receptors expression in Tex_prog vs. Tex_term from CD3+ sorted dataset from Figure 2 of predicted interactions from 6B.

(D) UMAP of human ndGBM and MBM datasets.

(E) Frequency of TAM vs. CD8s in ndGBM vs. MBM.

(F) Incoming vs. outgoing interaction strength across cell types in ndGBM vs. MBM.

(G) Heatmap of all incoming signaling patterns in ndGBM vs. MBM.

(H) Significantly upregulated predicted pathways in ndGBM (maroon) vs. MBM (grey).

# Supplemental Figures

**Supplemental Figure 1:**
(A) Gating strategy for Tex_prog and Tex_term.
(B) Frequency of Tex subsets of PD1+CD44+ CD8+ T cells in intracranial B16F0, LLC or CT2A. n = 8-10 mice/group.
All of the data are means ± SEMs; p-values are shown on individual plots as determined by 1-way ANOVA and post-hoc Tukey test (B).

**Supplemental Figure 2**

(A) Dot plot of top 3 genes for each cluster.

(B) Violin plot of *Tcf7, Pdcd1, Tox,* and *Havcr2* expression across clusters.

(C) UMAP of cloneType across biological replicates (n=4 per timepoint).

(D) Shannon and Inverse Simpson diversity indices across clusters.

(E) 10 most expanded TCR clones across timepoints are concentrated to Tex_int and Tex_term clusters.

**Supplemental Figure 3:**
(A) PD1 expression on transferred OT1 vs. TRP2-TCR Tg T cells on day of ALT (D7) and day of tumor harvest (D21) and representative histogram. n = 7 mice/group.
(B) Frequency of Tex subsets of PD1+ endogenous or transgenic TRP2-TCR T cells and representative contour plot. n = 7 mice.
(C) gMFI of MHC I in CT2A-TRP2 vs. CT2A-TRP2-B2mKO validating knockdown of MHC I. n = 10 mice/group.
(D) Number of TRP2-TCR T cells/g of tumor in CT2A-TRP2 vs. CT2A-TRP2-B2mKO. n = 10 mice/group.
p-values are plotted on individual graphs. Statistical significance was determined by two-sided unpaired t-test (A, C, D) or 2-way RM ANOVA with post-hoc Tukey's test for multiple comparisons (B). All data are plotted as mean ± SEM.

**A**

**B**

**Supplemental Figure 4:**

(A) Interaction strength across APC with CD8 TIL as the target.

(B) Gating strategy for intratumoral APC.

**Supplemental Figure 5:**

(A) Frequency of TAM of CD45+ cells. Data pooled from 2 independent experiments. n = 11 mice/group.

(B) Experimental layout. DC were depleted in CD11c-DTR mice with 5ng/g DT, every 3 days for 4 total treatments, starting D8. Created with Biorender.com.

(C) Frequency of CD11c+CD11b- DC in WT vs. CD11c-DTR mice.

(D) Experimental layout. Microglia were depleted in WT mice with 270mg/kg PLX3397 chow. Chow was provided starting D(-)14. Created with Biorender.com.

(E) Representative immunofluorescence image of WT vs. PLX3397 treated non-tumor brain. Scale bar = 100um.

(F) Chimera validation. Frequency of MHC I+ in WT → WT (WT) or B2mKO → WT (B2mKO) chimeras across T cells, TAM and cDC. n = 8-10 mice/group.

(G) Representative histogram of MHC I expression on TAM in the setting of CT2A-TRP2 (Teal) vs. CT2A-TRP2-B2mKO (Dark Teal).

(H) Frequency of TAM of CD45+ cells in B16F0 IC. n = 9-10 mice/group.

(I) Frequency of TAM of CD45+ cells in B16F0 SQ. Data pooled from 2 independent experiments. n = 6-7 mice/group.

p-values are plotted on individual graphs. Statistical significance was determined by two-sided unpaired t test (A, H, I) or RM-2-way ANOVA with post-hoc Tukey's test for multiple comparisons (F). All data are plotted as mean ± SEM.