1
2
3

# Sexually dimorphic leading-edge serrations evolved in silent swallows

Running title: **leading-edge serrations in silent swallows**

**Masaru Hasegawa[1],***

[1]*Department of Environmental Science, Ishikawa Prefectural University, Nonoichi, Ishikawa 921-8836, Japan*

Number of table 1

Number of figures 2

Number of supplementary material 1

*Author for corresponding

Dr  Masaru  Hasegawa

E-mail: perorobomusadiobe@gmail.com

**Abstract**

Leading-edge serrations are specialized feather structures, mitigating acoustic noise

during foraging flight in owls, and have been extensively studied and applied to

man-made noise-reducing structures. Similar structures occur in other avian species,

such as swallows, although the ecological functions of the serrations in these species

remain unclear. I conducted macroevolutionary analyses of hirundines (subfamily:

Hirundininae), in which leading-edge serrations evolved multiple times, to examine

their evolution in relation to vocal activity. I found that silent hirundines showed a

higher probability of possessing leading-edge serrations, indicating that leading-edge

serrations of swallows serve some functions associated with acoustics as in owls. I also

found that silent vocal activity accompanied loss of song, a well-known acoustic sexual

signal, but not with sexual plumage dimorphism, indicating that not sexual selection in

general but sexual selection on acoustic traits matters. These findings explain sexually

dimorphic leading-edge serrations, a unique characteristic in these diurnal

visually-foraging birds: stealth males rather than noisy males would be favored.

Although leading-edge serrations in hirundines would serve acoustic functions like

nocturnal birds such as owls, they might serve a unique sexual function.

**Keywords:** aeroacoustics, leading-edge comb, sexual plumage dimorphism,

vocalization

Leading-edge serrations, also known as leading-edge comb, are curved hooklets on the

leading edge of the primary feathers in owls (reviewed in Wagner et al. 2017; Jaworski

& Peake 2020; Terrill & Schultz 2023). This structure has intensively been studied and

shown to reduce noise in flight and thus can be adaptive to improve their hearing during

foraging flights and to avoid detection by prey (e.g., Rao et al. 2017; reviewed in

Wagner et al. 2017; Clark et al. 2020), which is now applied to man-made structure to

reduce acoustic noise (e.g., pantograph of Japanese Shinkansen trains: reviewed in

Wagner et al. 2017). However, owls are not the only animals that possess leading-edge

serrations (Clark et al. 2020); similar structures, albeit somewhat smaller and less

curved, can be found in other avian groups, including frogmouths (family: Podargidae)

and swallows (family: Hirundinidae), and the function of the serrations in these birds

remains unknown (reviewed in Clark et al. 2020).

In swallows (family: Hirundinidae), leading-edge serrations are known to have

evolved independently at least twice, once in rough-winged swallows (genus:

*Stelgidopteryx*) and the other in saw-wings (genus: *Psalidoprocne*), evident from their

English names (Turner & Rose 1994). Similar to other species with leading-edge

serrations (see above), these structures might serve to mitigate flight noise. However,

different from the majority of owls and frogmouths, swallows are diurnal,

visually-foraging birds, and, furthermore, they possess sexually dimorphic leading-edge

serrations, in which males, but not females, possess serrations (reviewed in Turner &

Rose 1994). This suggests that they might serve somewhat different functions than

sexually monomorphic leading-edge serrations found in mainly nocturnal, acoustic

predators (i.e., owls and frogmouths; e.g., see Roulin et al. 2013). In fact, although its

function has not been shown, several hypotheses are proposed to explain the evolution

of this sexually dimorphic structure (reviewed in Lunk 1962); for example, a clasping

structure for copulation, to prevent slipping from females (Steinbacher 1931) and a

means for generating non-vocal sound production during courtship or territorial chases

(Lunk 1962; de Jong 2020). It is also unclear, if leading-edge serrations are adaptive,

why only a few swallows possess leading-edge serrations (Steinbacher 1931; Lunk

1962), which sharply contrasts with owls in which virtually all owl species possess

some forms of leading-edge serrations (e.g., Le Piane & Clark 2022).

Here, I studied ecological correlates of leading-edge serrations in swallows and

martins (subfamily: Hirundininae). Because leading-edge serrations evolved at least two

times (see above), phylogenetic comparative approaches can be used to find out

ecological factors promoting the evolution of leading-edge serrations. Of several

hypotheses proposed, the most promising hypothesis is that, as in owls, leading-edge

serrations serve a function to reduce noise in flight (see above). Then, I can predict that

vocal activity would be negatively linked to the evolution of leading-edge serrations.

This is because reducing acoustic noise cannot make sense when birds continue to

vocalize. I also studied whether leading-edge serrations and its ecological correlates are

associated with proxies of sexual selection to clarify the possible role of sexual selection

on the evolution of sexually dimorphic leading-edge serrations in hirundines.

**Methods**

**Data collection**

As before (e.g., Hasegawa & Arai 2020a,b), information on morphological traits (i.e.,

wing length) and migratory habits (i.e., migrant or not) was obtained from Hasegawa &

Arai (2020b), which was derived from Turner & Rose (1994). I also collected data

concerning the presence/absence of song, which is usually sung by males, based on the

descriptions in the "voice" category of each species (i.e., whether or not "song" is noted

in the species) in Turner & Rose (1994). Further details are available elsewhere

(Hasegawa 2023). Likewise, I collected data concerning the vocal activities (silent or

not). Species with "silent", "quiet" or similar expression noted in the "voice" category

was regarded as silent species and other species were regarded as non-silent species

(unless data was unavailable).

As before (e.g., Hasegawa & Arai 2022), sexual dimorphism in overall

plumage characteristics as a proxy of sexual selection was obtained from Turner & Rose

(1994). Concerning a proxy of extrapair mating opportunities (frequent/rare), I used

incubation type (female-only versus biparental incubation) from a previous study

(Hasegawa & Arai 2020b), sourced mainly from Turner & Rose (1994). Incubation type

was tightly linked to extrapair paternity both within and between species in this clade

(Magrath & Elgar 1997; Hasegawa & Arai 2020b). The dataset used can be found in

Table S1.

**Statistics**

A Bayesian phylogenetic mixed model using a binomial error distribution was used to

examine the presence/absence of leading-edge serrations. As in previous studies (e.g.,

Hasegawa & Arai 2020a,b; Hasegawa 2023), potential confounding factors, body size

[measured via log(wing length)] and migratory habit, were included alongside vocal

activity in the model. To account for phylogenetic uncertainty, I fit the models to each

tree and applied a multimodel inference (e.g., Garamszegi & Mundry 2014), using 1000

alternative trees for swallows from birdtree.org (see Jetz et al. 2012, 2014; also see

Sheldon et al. 2005 for an original phylogeny on swallows). 1000 trees are sufficient to

control for phylogenetic uncertainty (Rubolini et al. 2015). The residual variance was

set at 1, in accordance with recommendations for categorical dependent variables (de

Villemereuil et al. 2013). I derived the mean coefficients, 95% credible intervals (CIs)

based on the highest posterior density, and Markov chain Monte Carlo (MCMC)-based

P-values, together with the phylogenetic signal (de Villemereuil & Nakagawa 2014). All

analyses were conducted in R ver. 4.1.0 (R Core Team 2021) using the "MCMCglmm"

function in the "MCMCglmm" package (ver. 2.29; Hadfield 2010). I ran the analysis for

140,000 iterations, with a burn-in period of 60,000 and a thinning interval of 80 for each

tree. The reproducibility of the MCMC simulation was assessed by calculating the

Brooks–Gelman–Rubin statistic (Rhat), which must be <1.2 for all parameters (Kass et

al. 1998), by repeating the analyses three times.

A discrete module in BayesTraits (Pagel & Meade 2006) was used to examine

evolutionary transitions among states with vocal activities (silent or not) and the

presence/absence of leading-edge serrations. I ran for 1,010,000 iterations with a

burn-in period of 10,000 and a thinning interval of 1000 (note that the phylogenetic tree

was randomly chosen from 1000 trees in each iteration). I denoted means as the

representatives of each transition rate. The reproducibility of the MCMC simulation was

assessed using the Brooks–Gelman–Rubin statistic (Rhat), which must be <1.2 for all

parameters (Kass et al. 1998), by repeating the analyses three times. Bayes factor was

calculated via the stepping stone sampler in BayesTraits, comparing the marginal

likelihood of a dependent model that assumed correlated evolution of the two traits (i.e.,

leading-edge serrations and vocal activity) to that of an independent model assuming

independent evolution. Bayes factors of >2, 5–10, and >10 indicate positive, strong, and

very strong evidence of correlated evolution, respectively (Kass & Raftery 1995;

BayesTraits Manual Ver. 3).

**Results**

**Leading-edge serrations and silent vocal activity**

Probability of possessing leading-edge serrations was significantly explained by the

vocal activity of swallows (Table 1): after controlling for phylogeny, hirundines with

relatively silent vocal activities showed a higher probability of possessing leading-edge

serrations (Fig. 1). Two covariates, body size measured as log(wing length) and

migratory habits (migrant or not) were non-significant (Table 1).

When examining evolutionary transitions among states (i.e., presence/absence

of leading-edge serrations and silent/non-silent vocal activities), I found strong support

for correlated evolution between these two state variables (n = 69, Bayes Factor = 8.49).

Leading-edge serrations evolved from the state with silent vocal activities, but virtually

did not evolve from the state with non-silent vocal activities (Fig. 2), though the results

should be carefully treated as leading-edge serrations evolved only a few times (Fig. 1).

**Silent vocal activity and proxies of sexual selection**

Silent species were less likely to have song (Bayesian phylogenetic mixed model with a

binomial error distribution: Coefficient = -3.09, 95% CI = -5.23, -1.08, $P_{MCMC}$ = 0.001,

phylogenetic signal = 0.24). In contrast, silent vocal activity had no detectable

relationship with sexual plumage dimorphism/monomorphism, a well-known proxy of

sexual selection (Bayesian phylogenetic mixed model with a binomial error distribution:

Coefficient = -0.04, 95% CI = -2.17, 2.08, $P_{MCMC}$ = 0.97, phylogenetic signal = 0.38).

Likewise, silent vocal activity had no detectable relationship with another proxy of

sexual selection in this clade, opportunity for extrapair matings measured as incubation

type, although the sample size for this analysis was reduced to 41 (Bayesian

phylogenetic mixed model with a binomial error distribution: Coefficient = -0.62, 95%

CI = -3.47, 2.15, $P_{MCMC}$ = 0.66, phylogenetic signal = 0.40). When silent vocal activity

was replaced by each of these variables, the presence of leading-edge serrations was not

significantly explained by these variables (i.e., song: $P_{MCMC}$ = 0.10; sexual plumage

dimorphism: $P_{MCMC}$ = 0.87; incubation type: $P_{MCMC}$ = 0.08, though this time migratory

species had higher probability of possessing leading-edge serrations than others, $P_{MCMC}$

< 0.01, perhaps because two non-migratory *Psalidoprocne* species with leading-edge

serrations were excluded from dataset due to the missing data, n = 41; details not

shown).

**Discussion**

The main finding of the current study is that leading-edge serrations evolved from

relatively silent hirundines, supporting that leading-edge serrations serve a function

associated with acoustics. Silent vocal activity evolved in a limited number of branches,

mostly in relatively young ones (Fig. 1), explaining the limited occurrence of

leading-edge serrations in hirundines. Although the current study does not clarify the

exact acoustic function, they may reduce noise in flight as found in owls and in artificial

wing models (e.g., Rao et al. 2017; see Introduction).

I also found that silent vocal activity had no detectable association with proxies

of sexual selection, i.e., sexual plumage dimorphism and opportunity for extra-pair

matings, and thus silent vocal activities would not result from weak (or intense) sexual

selection in general. Rather, silent vocal activity was associated with the absence of

male song, a conspicuous acoustic sexual signal (see Hasegawa 2023 for ancestral

character reconstruction of song in hirundines), indicating selection favors silent males

in these species. As selection sometimes favors muting in acoustic species accompanied

with deformed sound-producing structures in males (e.g., Hawaiian crickets; Pascoal et

al. 2014), selection for silent males might favor deformed leading-edges in males, which

engage in courtship in flight and territorial chases. Although this explanation assumes

viability cost of conspicuous sounds, possibly due to predation and other interspecific

interactions (e.g., competition over nest sites), silent flight activity might be directly

sexually selected. For example, it could aid in avoiding male aggression or female

evasions to succeed within-pair reproduction or extra-pair mating (e.g., see Hasselquist

& Bensch 1991 for an example of such "silent sneaking" in birds). In fact, this

explanation is more likely than the predator avoidance explanation (see above), because

even without leading-edge serrations, flight noise is inaudible from a distance (Wagner

et al. 2017). Leading-edge serrations would be functional only in a close-range, as in

intraspecific interaction. Of course, I cannot exclude the possibility that, as in owls (see

Introduction), hirundines use leading-edge serrations to reduce noise in foraging flight,

but this hypothesis alone cannot explain sexual dimorphism in leading-edge serrations,

and thus insufficient to explain the overall pattern of macroevolution of leading-edge

serrations in hirundines. Serrations are thought to mitigate flight noise just before

landing, which is corresponding to the final portion of hunting in owls (Wagner et al

2017; Clark et al. 2017) but this does not align with aerial foraging in hirundines

(Turner & Rose 1994).

An alternative explanation of the observed pattern is that leading-edge

serrations are used for non-vocal sound production during courtship or territorial chases

(Lunk 1962; de Jong 2020). Silent vocal activity might be a pre-condition for the

evolution of non-vocal acoustic communication. Although this is a fascinating idea,

there is no evidence that leading-edge serrations can produce non-vocal sounds and thus

is an unlikely explanation on the basis of an enormous amount of evidence that

serrations reduce, rather than increase, acoustic noise (see Introduction). It is, however,

possible that leading-edge serrations are a device to reduce noise to increase the

detectability of seemingly non-vocal, subtle sounds reported in swallows with

leading-edge serrations (i.e., genus: *Stelgidopteryx* and *Psalidoprocne*: Turner & Rose

1994; de Jong 2020), though we still need to identify the mechanisms of the possible

non-vocal sounds.

In summary, I demonstrated here that leading-edge serrations co-evolved with

silent vocal activities. The current study is the first macroevolutionary analyses of

sexually dimorphic leading-edge serrations. Since silent vocal activities are linked to the

loss of song, a well-known acoustic sexual signal, changing direction or target of sexual

selection (and its relative importance to viability selection: see below) would explain

sexually dimorphic expression of leading-edge serrations, which remains to be

demonstrated using manipulative experiments (e.g., removal of serrations). Thus, in

hirundines, a diurnal visually-oriented foragers, sexual selection might be involved in

the evolution of leading-edge serrations, different from acoustic foragers like owls and

other nocturnal birds, in which sexually monomorphic leading-edge serrations would

provide viability advantage for capturing prey. For swallows that are hyperaerial birds,

flight costs associated with leading-edge serrations (sensu Le Piane & Clark 2022; also

see Rao et al. 2017 for reduced lift:drag ratio in serrated wings) might prevent females

from evolutionarily maintaining male-like serrations, which is likely the case but should

be tested in future.

**Acknowledgments**

I thank Dr Emi Arai, Dr Shumpei Kitamura and his lab members at Ishikawa Prefectural

University for their kindest advices. I am grateful to Dr Angela Turner for her kindly

support on the valuable information on swallows.

**Author contributions:** I performed data analysis and wrote the manuscript alone.

**Availability of data and materials:** The data sets supporting this article are uploaded

as Table S1, which is also available from osf.io.

**Conflict of interest:** I have no competing interests.

**Animal welfare note:** This comparative study does not include any treatments of

animals, as all the information was gathered from bodies of literature.

**References**

Clark CJ, LePaine K, Liu L (2020) Evolution and ecology of silent flight in owls and

other flying vertebrates. Int Org Biol 2:obaa001

de Jong M. J. (2020). Northern Rough-winged Swallow (*Stelgidopteryx serripennis*),

version 1.0. In Birds of the World (A. F. Poole and F. B. Gill, Editors). Cornell

Lab of Ornithology, Ithaca, NY, USA. https://doi.org/10.2173/bow.nrwswa.01

de Villemereuil P, Nakagawa S (2014) General quantitative genetic methods for

comparative biology. In Garamszegi LZ (Ed), Modern phylogenetic comparative

methods and their application in evolutionary biology: Concepts and practice (pp.

287–301). Springer, New York.

de Villemereuil P, Gimenez O, Doligez B (2013) Comparing parent offspring regression

with frequentist and Bayesian animal models to estimate heritability in wild

populations: A simulation study for Gaussian and binary traits. Methods Ecol Evol

4:260–275.

Garamszegi LZ, Mundry R (2014) Multimodel-inference in comparative analyses. In

Garamszegi LZ (Ed), Modern phylogenetic comparative methods and their

application in evolutionary biology: concepts and practice (pp. 305–331). Springer,

New York.

Hadfield JD (2010) MCMC methods for multi-response generalized linear mixed

models: The MCMCglmm R package. J Stat Soft 33:1–22.

Hasegawa M (2023) Coevolutionary maintenance of forked tails and song in hirundines

(Aves: Hirundininae). J Avian Biol 2023:e03090.

Hasegawa M, Arai E (2020a) Fork tails evolved differently in swallows and swifts. J

Evol Biol 33:911–919.

Hasegawa M, Arai E (2020b) Correlated evolution of biparental incubation and sexual

tail monomorphism in swallows and martins (Aves: Hirundinidae). Evol Ecol

34:777–788.

Hasegawa M, Arai E (2022) Extrapair mating opportunity explains the evolutionary

transition between the presence and absence of tail ornamentation in swallows.

Evol Ecol 36:899–913.

Hasselquist D, Bensch S (1991) Trade-off between mate guarding and mate attraction in

the polygynous great reed warbler. Behav Ecol Sociobiol 28:187–193.

Jaworski JW, Peake N (2020) Aeroacoustics of silent owl flight. Ann Rev Fluid

Mechanics 52:395–420.

Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO (2012) The global diversity of

birds in space and time. Nature 491:444–448.

Jetz W, Thomas GH, Joy JB, Hartmann K, Redding D, Mooers AO (2014) Distribution

and conservation of global evolutionary distinctness in birds. Current Biology

24:919–930.

Kass RE, Raftery A (1995) Bayes Factors. J Am Stat Assoc 90:773–795.

Kass RE, Carlin BP, Gelman A, Neal R (1998) Markov chain Monte Carlo in practice: a

round table discussion. Am Stat 52:93–100.

Le Piane K, Clark CJ (2022) Quiet flight, the leading edge comb, and their ecological

correlates in owls (Strigiformes). Biol J Linn Soc 135:84–97.

Lunk WA (1962) The Rough-winged Swallow: a study based on its breeding biology in

Michigan. Publications of the Nuttall Ornithological Club 4.

Magrath MJL, Elgar MA (1997) Paternal care declines with increased opportunity for

extra-pair matings in fairy martins. Proc R Soc Lond B 264:1731–1736

Pagel M, Meade A (2006) Bayesian analysis of correlated evolution of discrete

characters by reversible-jump Markov chain Monte Carlo. Am Nat 167:808–825.

Pascoal S, Cezard T, Eik-Nes A, Gharbi K, Majewska J, Payne E, Ritchie MG, Zuk M,

Bailey NW (2014) Rapid convergent evolution in wild crickets. Curr Biol

24:1369–1374.

R Core Team (2021) R: A language and environment for statistical computing. R

Foundation for Statistical Computing.

Rao C, Ikeda T, Nakata T, Liu H (2017) Owl-inspired leading-edge serrations play a

crucial role in aerodynamic force production and sound suppression. Bioinspir

Biomim 12:046008.

Roulin A, Mangels J, Wakamatsu K, Bachmann T (2013) Sexually dimorphic

melanin-based colour polymorphism, feather melanin content, and wing feather

structure in the barn owl (*Tyto alba*). Biol J Linn Soc 109:562–573.

Rubolini D, Liker A, Garamszegi LZ, Møller AP, Saino N (2015) Using the

BirdTree.org website to obtain robust phylogenies for avian comparative studies:

A primer. Curr Zool 61:959–965.

Sheldon FH, Whittingham LA, Moyle RG, Slikas B, Winkler DW (2005) Phylogeny of

swallows (Aves: Hirundinidae) estimated from nuclear and mitochondrial DNA

sequences. Molecular Phylogenetics and Evolution 35:254–270.

Steinbacher G (1931) Ueber die biologische Bedeutung des Geschleehtsdimorphismus

bei den Sagelfliigelschwalben *Psalidoprocne* und *Stelgidopteryx*. Ornith. Monatsb.

39: 71–73.

Terrill RS, Schultz AJ (2023) Feather function and evolution f birds. Biol Rev 98:540–

566.

Turner AK, Rose C (1994) A Handbook to the Swallows and Martins of the World.

Helm.

Wagner H, Weger M, Klaas M, Schroder W (2017) Features of owl wings that promote

silent flight. Interface Focus 7:20160078.

**Table 1** Multivariable Bayesian phylogenetic mixed model with a binary error
distribution predicting the presence/absence of leading-edge serrations in relation to
vocal activities and covariates (n = 69) in swallows and martins (Aves: Hirundininae).

| Variables | Coefficient [95% CI] | $P_{MCMC}$ |
|---|---|---|
| Intercept | 36.35 [-60.36, 138.33] | 0.45 |
| log(wing length) | -9.44 [-31.45, 11.49] | 0.36 |
| Migration (non-migrant = 0, migrant = 1) | 1.40 [-2.67, 5.60] | 0.48 |
| Vocal activity (non-silent = 0, silent =1) | **4.79 [0.88, 8.96]** | **<0.01** |
| | Mean phylogenetic signal = 0.51 | |

339 Model-averaged coefficients and 95% credible intervals (CIs) are shown.
340 Bold indicates significant variables (i.e., $P_{MCMC} < 0.05$).
341

 Figure legends

Fig. 1 An example of ancestral character reconstruction of the leading-edge serrations (left) and of silent vocal activity (right) in swallows and martins (Aves: Hirundininae). Left side: presence and absence of leading-edge serrations are indicated by white and black circles, respectively; silent and non-silent vocal activity are indicated by white and black circles, respectively. The proportions of white and black at the nodes indicate the probability of the ancestral state. Here, I used the functions "ace" in the R package "ape" (with model = "ER", i.e., an equal-rates model; Paradis et al. 2005) and "plotTree" in the R package "phytools" (Revell 2012) for ancestral character reconstruction.

**Fig. 2** The most likely evolutionary transitions between states with the presence/absence of leading-edge serrations and the silent/non-silent vocal activity (see illustrations beside each state) in swallows and martins (Aves: Hirundininae). Numbers next to arrows, represented by arrow size, indicate transition rates between pairs of states. Leading-edge serrations of swallows are illustrated by referring to fig. 6 in de Jong (2020).

393    **Table S1**    Dataset of the current study (n = 69).

| Scientific Name | Serrations | Wing length | Migratory habit | Silent/non-silent vocalization | Presence/absence of song | Sexual plumage dimorphism | Incubation type† |
|---|---|---|---|---|---|---|---|
| *Neochelidon tibialis* | 0 | 85 | 0 | 0 | 0 | 0 | NA |
| *Alopochelidon fucata* | 0 | 100 | 1 | 1 | 0 | 0 | NA |
| *Stelgidopteryx ruficollis* | 1 | 109 | 1 | 1 | 0 | 0 | 1 |
| *Stelgidopteryx serripennis* | 1 | 110 | 1 | 1 | 0 | 0 | 1 |
| *Tachycineta bicolor* | 0 | 119 | 1 | 0 | 1 | 1 | 1 |
| *Tachycineta albilinea* | 0 | 97 | 0 | 0 | 1 | 0 | NA |
| *Tachycineta albiventer* | 0 | 104 | 1 | 0 | 1 | 0 | NA |
| *Tachycineta thalassina* | 0 | 122 | 1 | 1 | 0 | 1 | 1 |
| *Tachycineta leucorrhoa* | 0 | 115 | 1 | 0 | 1 | 0 | 1 |
| *Tachycineta meyeni* | 0 | 110 | 1 | 0 | 1 | 0 | 2 |
| *Tachycineta cyaneoviridis* | 0 | 115 | 0 | 0 | 0 | 1 | 1 |
| *Tachycineta euchrysea* | 0 | 106 | 0 | 0 | 0 | 1 | NA |
| *Notiochelidon murina* | 0 | 111 | 0 | 1 | 1 | 0 | NA |
| *Notiochelidon flavipes* | 0 | 90 | 0 | 0 | 0 | 0 | 2 |
| *Notiochelidon pileata* | 0 | 95 | 0 | 0 | 1 | 0 | NA |
| *Pygochelidon cyanoleuca* | 0 | 94 | 1 | 0 | 1 | 0 | 2 |
| *Atticora fasciata* | 0 | 101 | 0 | 0 | 1 | 0 | NA |
| *Atticora melanoleuca* | 0 | 93 | 0 | 0 | 0 | 0 | NA |
| *Progne tapera* | 0 | 130 | 1 | 0 | 1 | 0 | 1 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| *Progne subis* | 0 | 145 | 1 | 0 | 1 | 1 | 1 |
| *Progne chalybea* | 0 | 131 | 1 | 0 | 1 | 1 | 1 |
| *Progne dominicensis* | 0 | 143 | 1 | 0 | 1 | 1 | NA |
| *Progne modesta* | 0 | 125 | 1 | 0 | 1 | 1 | NA |
| *Riparia paludicola* | 0 | 104 | 0 | 0 | 1 | 0 | 2 |
| *Riparia riparia* | 0 | 107 | 1 | 0 | 1 | 0 | 2 |
| *Riparia cincta* | 0 | 130 | 1 | 0 | 1 | 0 | NA |
| *Psalidoprocne fuliginosa* | 1 | 104 | 0 | 0 | 1 | 0 | NA |
| *Psalidoprocne albiceps* | 1 | 102 | 1 | 1 | 0 | 1 | 2 |
| *Psalidoprocne pristoptera* | 1 | 106 | 1 | 1 | 1 | 0 | 1 |
| *Psalidoprocne obscura* | 1 | 96 | 0 | 1 | 1 | 1 | NA |
| *Psalidoprocne nitens* | 1 | 93 | 0 | 1 | 0 | 0 | NA |
| *Cheramoeca leucosterna* | 0 | 102 | 0 | 0 | 1 | 0 | NA |
| *Pseudhirundo griseopyga* | 0 | 97 | 0 | 1 | 1 | 0 | NA |
| *Phedina borbonica* | 0 | 116 | 0 | 1 | 1 | 0 | 1 |
| *Hirundo rupestris* | 0 | 130 | 1 | 0 | 1 | 0 | 1 |
| *Hirundo fuligula* | 0 | 129 | 0 | 1 | 1 | 0 | 2 |
| *Hirundo concolor* | 0 | 106 | 0 | 0 | 1 | 0 | 2 |
| *Hirundo rustica* | 0 | 124 | 1 | 0 | 1 | 1 | 1 |
| *Hirundo lucida* | 0 | 111 | 0 | 0 | 1 | 0 | NA |
| *Hirundo angolensis* | 0 | 119 | 0 | 0 | 1 | 0 | NA |
| *Hirundo tahitica* | 0 | 105 | 0 | 0 | 1 | 0 | 1 |
| *Hirundo neoxena* | 0 | 112 | 0 | 0 | 1 | 0 | 1 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| *Hirundo albigularis* | 0 | 128 | 1 | 0 | 1 | 1 | NA |
| *Hirundo aethiopica* | 0 | 106 | 0 | 0 | 1 | 0 | 1 |
| *Hirundo smithii* | 0 | 110 | 1 | 0 | 1 | 1 | 1 |
| *Hirundo nigrita* | 0 | 106 | 0 | 0 | 1 | 0 | 1 |
| *Hirundo leucosoma* | 0 | 99 | 0 | 1 | 0 | 0 | NA |
| *Hirundo dimidiata* | 0 | 102 | 1 | 0 | 1 | 1 | 1 |
| *Hirundo atrocaerulea* | 0 | 113 | 1 | 0 | 1 | 1 | 1 |
| *Hirundo nigrorufa* | 0 | 112 | 1 | 1 | 1 | 1 | NA |
| *Hirundo cucullata* | 0 | 125 | 1 | 0 | 1 | 1 | 1 |
| *Hirundo abyssinica* | 0 | 106 | 1 | 0 | 1 | 1 | 1 |
| *Hirundo semirufa* | 0 | 132 | 1 | 0 | 1 | 0 | 1 |
| *Hirundo senegalensis* | 0 | 144 | 0 | 0 | 1 | 0 | NA |
| *Hirundo daurica* | 0 | 124 | 1 | 0 | 1 | 0 | 2 |
| *Hirundo striolata* | 0 | 124 | 0 | 0 | 1 | 0 | NA |
| *Hirundo preussi* | 0 | 95 | 0 | 0 | 1 | 0 | NA |
| *Hirundo rufigula* | 0 | 97 | 1 | 0 | 1 | 0 | 2 |
| *Hirundo spilodera* | 0 | 111 | 1 | 0 | 0 | 0 | 2 |
| *Hirundo fuliginosa* | 0 | 88 | 0 | 0 | 1 | 1 | NA |
| *Petrochelidon pyrrhonota* | 0 | 109 | 1 | 0 | 1 | 0 | 2 |
| *Haplochelidon andecola* | 0 | 115 | 0 | 0 | 1 | 0 | NA |
| *Petrochelidon fulva* | 0 | 102 | 1 | 0 | 1 | 0 | 2 |
| *Hirundo fluvicola* | 0 | 91 | 1 | 0 | 1 | 0 | 2 |
| *Hirundo ariel* | 0 | 91 | 1 | 0 | 1 | 0 | 2 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| *Hirundo nigricans* | 0 | 107 | 1 | 0 | 1 | 0 | NA |
| *Delichon urbicum* | 0 | 110 | 1 | 0 | 1 | 0 | 2 |
| *Delichon dasypus* | 0 | 108 | 1 | 0 | 1 | 0 | 2 |
| Delichon nipalense | 0 | 95 | 0 | 1 | 0 | 0 | 2 |

394 †Female-only incubation is denoted as 1 and biparental incubation is denoted as 2.

395 NA indicates data "Not Available."

396 See text for detailed information.