Stress granule dysfunction via chromophore-associated light inactivation

Takumi Koizumi[1], Haruka Kawaguchi[1], Tsumugi Kurosaki[1], Akira Kitamura[2,3*]

[1]Laboratory of Molecular Cell Dynamics, Graduate School of Life Science, Hokkaido University,
Sapporo, Japan

[2]Laboratory of Molecular Cell Dynamics, Faculty of Advanced Life Science, Hokkaido University,
Sapporo, Japan

[3]PRIME, Japan Agency for Medical Research and Development, Chiyoda-ku, Tokyo, Japan

* Correspondence:
Akira Kitamura
akita@sci.hokudai.ac.jp
ORCiD ID: https://orcid.org/0000-0001-8357-8593

Keywords: stress granule, G3BP1, TDP-43, cell viability, chromophore-associated light
inactivation

Abstract

Stress granules (SGs) are membraneless organelles composed of various proteins and RNAs that
protect translation-associated machinery from harmful conditions during stress. However, the
functional differences among SG compartments remain unclear. Here, we show that the spatial and
temporal control of SG subcompartments by chromophore-associated light inactivation (CALI) using
a genetically encoded red fluorescence protein (SuperNova-Red) as a photosensitizer leads to
differences in cell viability during recovery from hyperosmotic stress and that the inactivation of the
core phase in the SGs could be more critical than the shell phase. CALI delayed the disassembly
kinetics of SGs during recovery from hyperosmotic stress. Consequently, CALI could inactivate the
subcompartments of the SGs, and the cellular fate due to SGs could be analyzed. Furthermore, CALI
is an effective spatiotemporal knockdown method for intracellular condensates/aggregates and
contributes to the elucidation of membraneless organelles.

1   Introduction

Stress granules (SGs) are membraneless organelles; they are biomolecular condensates composed of
proteins and RNAs that emerge in the cytoplasm under cellular stresses such as hyperosmotic, heat,
and oxidative stress (1, 2). After being released from stress, the SGs are reversibly disassembled (1).
Although the function of SGs remains elusive, they have long been proposed to play a role in
protecting translation-associated RNAs from harmful conditions under stress because translation-
associated proteins and RNAs, such as ribosomes, pre-translation complexes with mRNA, and
various translation-associated RNA-binding proteins (RNPs), are widely sequestered into the SGs (3-
5). Therefore, SGs are believed to play a cytoprotective role under stress conditions. SGs contain a
biphasic structure with a stable "core" phase surrounded by a less concentrated "shell" phase (6).
Ras-GTPase-activating protein SH3 domain-binding protein 1 (G3BP1) is an RNA-binding protein

essential for SG formation, which can primarily be found in the core phase (1, 2). Loss of G3BP1
increases the number of ubiquitinated proteins and toxic aggregates (7, 8) and inhibits the
proliferation of cancer cells (9). Transactivation response DNA/RNA-binding protein 43 kDa (TDP-
43) is an amyotrophic lateral sclerosis-associated aggregation-prone protein and a component of SGs.
One starting point for the pathological aggregation of TDP-43 is its localization to and interaction
with SG components in the shell phase (10). As the functional structure of SGs, the core phase is an
important scaffold for SG formation. The genetic depletion of endogenous G3BP1 reduces SG
formation efficiency (11). However, it is still unknown whether the loss of SG proteins in the shell
phase causes severe defects in the SGs. To elucidate the mechanisms underlying this issue, the
spatiotemporal inactivation of the shell phase, rather than the genetic knockout/knockdown of genes
of interest, is required because compartment-specific protein knockdown is not feasible with
traditional genetic methods.

Chromophore-assisted light inactivation (CALI) is used for the spatiotemporal inactivation of
specific compartments of live cells (12, 13). Absorption of light by the chromophore in phototoxic
photosensitizers produces reactive oxygen species (ROS), which, in turn, inactivates photosensitizer-
proximate biomolecules. Although various photosensitizers have been used, fluorescent protein-
based photosensitizers can be labeled at a defined molar ratio (e.g., 1:1) with the target protein,
expressed in cells via gene transfection, and localized to specific subcellular compartments (14).
SuperNova-Red (SNR) is a monomeric photosensitizer and a red fluorescent protein with high ROS
production (15). Therefore, we hypothesized that the spatiotemporal inactivation of SGs can be
achieved by tagging the SNR to specific proteins within specific compartments in the SGs under
stress.

In this report, we show how CALI changes the cell viability and kinetics of SG disassembly using
SNR to G3BP1 and TDP-43 in the SGs as targets during the recovery process after hyperosmotic
stress.

2　　Materials and methods

Plasmids: To create the expression plasmids of proteins that SNR was tagged at the C-terminus of
G3BP1 and TDP-43, SNR DNA fragment (gift from Prof. T. Nagai) was inserted into the *Bam*HI and
*Not*I sites of pcDNA3.1(+) (Thermo Fisher Scientific, Waltham, MA, USA) (pcDNA-SNR). Next,
the DNA fragments of human TDP-43 obtained from the TDP-43 expression plasmid as previously
reported (16) and G3BP1 from Addgene (Clone#129339, Watertown, MA, USA) were inserted into
the *Hind*III and *Bam*HI sites of pcDNA-SNR (pTDP43-SNR and pG3BP1-SNR, respectively). To
create monomeric enhanced green fluorescent protein (meGFP)-tagged TDP-43 or G3BP1 expression
vectors, the DNA fragment of the SNR of pTDP43-SNR and pG3BP1-SNR was cut and substituted
with that of meGFP digested from pmeGFP-N1 (16) using the *Bam*HI and *Not*I sites (pTDP43-GFP
and pG3BP1-GFP, respectively). The expression plasmid for meGFP-tagged histone H2B (pBOS-
H2B-GFP) was the same as that used previously (16, 17).

Cell Culture and Transfection: Neuro-2a murine neuroblastoma (N2a) cells were obtained from the
American Type Culture Collection (ATCC; Manassas, VA, USA) and maintained in Dulbecco's
Modified Eagle Medium (DMEM) (D5796, Sigma-Aldrich, St. Louis, MO, USA) supplemented with
10% heat-inactivated fetal bovine serum (FBS) (12676029, Thermo Fisher Scientific), 100 U/mL
penicillin G (Sigma-Aldrich), and 0.1 mg/mL streptomycin (Sigma-Aldrich), as previously described
(16). One day before transfection, 2.0×10^5 N2a cells were transferred to a glass bottom dish (3910-
035, IWAKI-AGC Technoglass, Shizuoka, Japan). The mixture of expression plasmids was

transfected using Lipofectamine 2000 (Thermo Fisher Scientific) according to the manufacturer's
protocol. Details on the amounts of the plasmids are described in the Supplementary Table. After 24
h of incubation, the spent medium was replaced with Opti-MEM I (Thermo Fisher Scientific) or
Opti-MEM I containing 290 mM NaCl (hyperosmotic medium).

Stress granule formation, light irradiation, and time-lapse observation: To generate SGs under
hyperosmotic stress, N2a cells were incubated in a hyperosmotic medium for 3 h. After the
hyperosmotic medium was removed, the cells were washed thrice in Hank's balanced salt solution,
and then the medium was exchanged with Opti-MEM I (recovery medium) containing 0.3 mM
DRAQ7 (BioStatus, Leicestershire, UK) as an indicator for dead cells. After medium exchange, the
glass-based dish was immediately placed on a microscope stage within 2 min. Light irradiation and
time-lapse observations were performed using an Axioobserver Z1 inverted microscope (Carl Zeiss,
Jena, Germany) equipped with an LSM510 META laser scanning confocal unit (Carl Zeiss), a C-
Apochromat 40×/1.2NA W Korr. UV-VIS-IR water immersion objective (Carl Zeiss), laser sources,
a mercury lamp, and a heat stage incubator (at 37°C supplemented with 5% CO_2). The field stop was
set to a minimum. For CALI, cells 1–5 expressing SNR-tagged proteins were simultaneously
irradiated with light from a mercury lamp through a 540–580 nm bandpass filter (ET560/40x;
Chroma Technology Corporation, Bellows Falls, VT, USA) and a 585 nm dichroic mirror (T585lpxr;
Chroma Technology Corporation) for 2 min. The aperture diaphragm was adjusted so that the light
intensity was 0.9 mW just above the objective lens. After the CALI of SNR, bright-field and DRAQ7
fluorescence images of the irradiated cells were acquired at 30-minute intervals for up to 600 min
(Supplementary Figure 1). The DRAQ7-stained cells were excited at 633 nm. The excitation and
emission beams were split using a beam splitter (HFT405/488/543/633). Fluorescence was detected
using photomultiplier tubes through a 650 nm long-pass filter (LP650). The diameter of the pinhole
for DRAQ7 was 92 μm. The image size was 1024 × 1024 pixels and the zoom factor was 3. DRAQ7-
positive cells were considered dead. To observe SG disassembly, GFP was excited at 488 nm. The
excitation and emission beams were split using a beam splitter (HFT488/594). Fluorescence was
detected using photomultiplier tubes through a 505–570 nm bandpass filter (BP505–570). The
pinhole diameter was 72 μm, the image size was 1024 × 1024 pixels, and the zoom factor was 4.
After light irradiation for CALI, bright-field and GFP fluorescence images of the irradiated cells were
acquired at 30-minute intervals for up to 600 min.

Confocal microscopy: Image acquisition was performed using an inverted fluorescence microscope,
Axioobserver Z1 (Carl Zeiss, Jena, Germany), combined with an LSM 510META system (Carl
Zeiss), a C-Apochromat 40×/1.2NA W Korr. UV-VIS-IR water immersion objective (Carl Zeiss),
laser sources, a mercury lamp, and a heat stage incubator (at 37°C supplemented with 5% CO_2). GFP
and SNR were excited at 488 nm and 594 nm, respectively. The excitation and emission beams were
split using a beam splitter (NFT488/594). Fluorescence was detected using photomultiplier tubes
through a 615 nm long-pass filter (LP615). The diameter of the pinhole was 72 μm for GFP and 208
μm for SNR.

Statistical analyses: Student's t-tests were performed using Microsoft Excel. The generalized
Wilcoxon test (Gehan-Breslow method) was performed using the Bell Curve for Excel software
(Social Survey Research Information Co., Ltd., Tokyo, Japan).

3   Results

3.1   After CALI, SG-containing cells showed low viability during recovery from hyperosmotic
stress

To demonstrate whether SGs play a cytoprotective role, we established a procedure to observe the
viability of cells expressing a protein target in CALI during the recovery period after hyperosmotic
stress caused by 290 mM NaCl (Supplementary Figure 1). Under this condition, SGs form in the
cytoplasm because their assembly and disassembly have previously been clarified under
hyperosmotic stress and after the recovery process (18). First, we tested whether the SNR monomers
induce cell death in the recovery medium after hyperosmotic stress. After light irradiation, no SNR
fluorescence was detected, indicating that SNR was efficiently photobleached (Supplemental Figure
2A). Photobleaching is important for ROS production from fluorophores (19). The viability of cells
expressing SNR monomers decreased to 86% at 600 min after the start of the time-lapse observation
(Supplemental Figure 2B), indicating that neither ROS from SNR monomers nor the cellular
recovery process from hyperosmotic stress caused cell death.

Next, as G3BP1 localizes in the core phase of SGs, we chose G3BP1 as a target for the
inactivation of SGs in CALI. Cytoplasmic SGs of G3BP1-SNR were observed in some of the cells
exposed to hyperosmotic stress; however, cells without SGs of G3BP1-SNR were also observed
(Figure 1A). Hence, we compared the proportion of surviving cells harboring SGs with that of cells
not harboring SGs during recovery from hyperosmotic stress. As negative controls, almost all cells
without light irradiation were alive regardless of G3BP1-SNR SG formation (green and orange lines,
Figure 1B), indicating that neither the recovery process from hyperosmotic stress nor time-lapse
observation affected the survival of the cells expressing G3BP1-SNR. In contrast, cells exposed to
hyperosmotic stress with light irradiation gradually died in both the SG-positive and SG-negative cell
groups during recovery from hyperosmotic stress. However, cells harboring SGs of G3BP1-SNR
died within a shorter time than those that did not (blue and magenta lines, Figure 1B). Therefore, we
successfully confirmed the reduction in cell viability induced by CALI in cells expressing G3BP1-
SNR, and that SG-positive cells were more likely to be affected by CALI-induced cell death than
SG-negative cells.

As TDP-43 localizes in the shell phase of SGs (10), we chose it as a target for the inactivation of
SGs in CALI. We also tracked cell viability after light irradiation using the same procedure as
G3BP1-SNR, where cells with and without TDP43-SNR SGs were observed (Figure 2A). Similar to
G3BP1-SNR-expressing cells, cell death was not observed in TDP43-SNR-expressing cells,
regardless of SGs, when no light was used as a negative control (green and orange lines, Figure 2B).
In contrast, significant cell death was observed in cells harboring SGs of TDP43-SNR; most cells that
did not harbor TDP43-SNR SGs remained alive (blue and magenta lines, Figure 2B). Furthermore, a
higher proportion of CALI-induced cell death was observed in cells expressing G3BP1-SNR than in
those expressing TDP43-SNR (Figure 3).

3.2   CALI inhibits SG disassembly

As SGs are gradually disassembled during recovery from hyperosmotic stress (20), we investigated
whether CALI-induced G3BP1 or TDP-43 inactivation in SGs affected the disassembly of SGs. To
observe the subcellular localization of G3BP1 or TDP-43 after the photobleaching of SNR for CALI,
N2a cells co-expressing meGFP-tagged and SNR-tagged G3BP1/TDP-43 were subjected to
hyperosmotic stress to form SGs in the cytoplasm, and then irradiated with light immediately after
replacement with the recovery medium. Immediately after light irradiation in CALI, the SNR
fluorescence completely disappeared; however, the subcellular localization of GFP-tagged G3BP1
and TDP-43 was not affected (Supplementary Figure 3). Next, the disassembly of G3BP1- and TDP-
43-containing SGs was observed using time-lapse fluorescence microscopy (Figure 4A). G3BP1-
containing SGs were gradually disassembled with and without light irradiation in the recovery

medium. However, in the light-irradiated cells, the SG disassembly speed was delayed up to ~270
min in the recovery medium, and the population of cells harboring SGs decreased to the same extent
with and without light irradiation (>300 min) (Figure 4B). The TDP-43-containing SGs with and
without light irradiation gradually disassembled at a similar speed in the recovery medium (Figure
4B). Although the disassembly speed of TDP-43-containing SGs in the recovery medium did not
differ with or without light irradiation up to ~270 min, its disassembly in light-irradiated cells did not
progress after its abundance was reduced to approximately 40% of the original; the remaining
proportion of SGs in light-irradiated cells was greater than that in non-irradiated cells (>350 min)
(Figure 4C). In summary, CALI inhibited SG disassembly.

4   Discussion

Here, we showed that CALI using SNR to target G3BP1 and TDP-43 in SGs increased cell death
(Figures 1–3) and inhibited SG disassembly (Figure 4). First, we determined the possible
mechanisms by which CALI inhibits SG disassembly. Light-irradiated SNR produces both
superoxide and singlet oxygen as ROS; however, the quantum yield of SNR to generate singlet
oxygen is approximately 15-fold higher than that to generate superoxide (19). The higher efficiency
of SNR in generating singlet oxygen compared to superoxide is similar to other red fluorescent
proteins, such as mCherry (21) and KillerRed (22) but the highest among these proteins (15, 19). The
lifetime of singlet oxygen in water was reported at 3.7 μs (23). However, the travel distance of singlet
oxygen in cells is believed to be less than 2 nm because singlet oxygen reacts with water and other
biomolecules (24); hence, singlet oxygen can spread only within the local environment (25). Because
the short diameter of the fluorescent protein-conserved β-barrel of SNR is approximately 1.3 nm
(15), the singlet oxygen produced from SNR is consumed by biomolecules close to the outside of the
β-barrel.

Singlet oxygen damages proteins, peptides, and nucleic acids in various ways (26-30). In
proteins and peptides, singlet oxygen can react with the side chains of amino acids (e.g., Trp, Tyr,
His, Met, and Cys), resulting in the chemical modification of these amino acids, backbone
fragmentation, cross-linking, and subsequent aggregation (26, 31). In nucleic acids, guanosine is a
major oxidization target of singlet oxygen (29, 30). Because SGs are densely packed with various
proteins and RNAs (1, 2), singlet oxygen generated from G3BP1-SNR or TDP43-SNR may act on
the proteins and RNAs in the SGs, leading to side chain modifications, backbone fragmentations, and
cross-linking of SG components in the phase where SNR-tagged proteins are localized. These effects
do not produce an immediate change or instantaneous disappearance of SGs (Supplementary Figure
3); however, they inhibit SG disassembly (Figure 4B). As previously reported, the most easily
interpreted cause of difficulty in SG disassembly is intermolecular cross-linking (32-34). Another
possibility is that molecular chaperones such as HSP70 are incorporated in the SGs to protect and
possibly repair proteins that unfold during various types of stress and are required for the
disassembly of SGs (3). Inactivation of such molecular chaperones by CALI would cause defects in
SG disassembly. As the abundance, conformations, modifications, and interactions of RNA may
control the assembly and disassembly of SGs (1, 35), singlet oxygen-induced guanosine oxygenation
in RNA may retard the SG disassembly process.

Because cross-linking is a covalent bond, the dissociation of biomolecules is less likely to occur.
Interestingly, the disassembly kinetics differed between G3BP1 and TDP-43 (Figure 4B). As G3BP1
and TDP-43 localize to different phases in the SG (core and shell, respectively), CALI may affect the
disassembly process in each phase. As TDP-43 is localized in the shell, the outer phase of the SGs,
TDP-43 may be released not as a single molecule but as an oligomeric or multimeric unit, likely with

other SG components. However, the population of TDP-43 cross-linked with molecules that are static
in the SG may remain inside the SGs. In contrast, because G3BP1 localizes to the core phase of the
SGs, when the SG itself collapses, it is likely to dissociate less if it is cross-linked. However, light-
irradiated G3BP1-harboring SGs disassembled even after a delay, so there may be a repair system for
cross-linked species during SG disassembly. Although singlet oxygen was artificially generated from
SNR in this experiment, singlet oxygen and other ROS can be generated by intracellular redox
reactions in physiological environments (36). Singlet oxygen and other ROS generated in
physiological states form intermolecular cross-links in SGs. In other words, this study attempted to
mimic the effects of singlet oxygen-promoted inactivation of intracellular structures such as SGs.

Next, we discuss the influence of CALI outside the SGs (i.e., in the cytoplasm and nucleus). The
G3BP1-SNR and TDP-43-SNR used in this study were localized not only in the SGs but also in the
cytoplasm and nucleus. Because whole cells were irradiated with light, ROS may have been
produced in the cytoplasm and nucleus. Our results showed that cell death was more pronounced
after CALI in SG-positive cells for both G3BP1-SNR and TDP-43-SNR (Figures 1–3), suggesting
that SG inactivation was more strongly associated with cell death than defects in these proteins
localized in the cytoplasm or nucleus. However, CALI in cells not harboring SGs of G3BP1-SNR
also induced cell death (Figure 1), suggesting that the inactivation of G3BP1 in the cytoplasm and
nucleus may be involved in cell death. Loss of G3BP1 increases protein ubiquitination and the
aggregation of toxic huntingtin (7, 8) and diminishes the proliferation of breast cancer cells (9); the
inactivation of G3BP1 by CALI likely leads to proteostasis dysregulation and eventual cell death. In
contrast, the loss of TDP-43 leads to various cellular dysfunctions, such as the misregulation of RNA
splicing and its quality control, mitochondrial dysfunction, and cell death (37-40). However, cell
death during TDP-43 knockdown was observed at approximately 120 h after siRNA introduction
(40). This suggests that cell death by TDP-43 depletion may not be an acute response but rather via
the accumulation of various dysfunctions. CALI of TDP43-SNR in cells not harboring SGs did not
result in a dramatic increase in cell death (Figure 2B), consistent with the slow rate of cell death
observed in TDP43-depleted cells. This supports the specificity of CALI for the protein of interest
and its localization.

Finally, we discuss the possible mechanisms of cell death induced by CALI in SGs. SGs
attenuate apoptotic cell death via p53 inhibition and the recruitment of ROCK1 and RACK1 to SGs
(41-43). SG formation regulates several canonical signaling pathways (such as NF-κB, mTORC1,
and PKR) and pro-survival functions during the acute stress response (1, 4). The dysfunction of SGs
induced by CALI may reactivate transiently inhibited apoptotic signaling pathways. The CALI-
induced aberrant RNA in the SGs may disturb cellular functions. mRNA translation is stalled as a
pre-initiation complex in SGs, as translation is traditionally considered silent during SG formation
(1). Although some mRNAs are translationally silent, a host of RNA molecules undergo translation
in SGs (44). Therefore, the translation dysfunction required for stress recovery is delayed by aberrant
RNA induced by CALI in the SGs, leading to cell death.

Consequently, here, we showed that CALI of two SG components (G3BP1 and TDP-43) using
SNR retarded SG disassembly and led to cell death. These results suggest that CALI can inactivate
various intracellular condensates composed of proteins and RNA in a spatiotemporally controlled
manner, and the subsequent cellular fate can be analyzed. Furthermore, we believe that CALI is an
effective method that contributes to identifying intercellular condensates/aggregates as hubs
controlling signal transduction. However, the molecular mechanism of inactivation by CALI remains
unknown, and its identification remains an important open question.

5   Conflict of Interest

The authors declare no conflicts of interest.

6   Author Contributions

TKo: Methodology, Formal Analysis; HK: Methodology, Formal Analysis; TKu: Methodology; AK: Writing – original draft, review, editing, Conceptualization, Methodology, Project administration.

7   Funding

This research was funded by the Japan Agency for Medical Research and Development (AMED), grant numbers JP23gm6410028, JP22ym0126814, and JP23ym0126801; the Japan Society for the Promotion of Science (JSPS), grant number 22H04826, 22H02578, and 22K19886; the Nakatani Foundation for Advancement of Measuring Technologies in Biomedical Engineering; the Hagiwara Foundation; the Hoansha Foundation; the Northern Advancement Center for Science & Technology (NOASTEC) for AK; and the Hokkaido University Office for Developing Future Research Leaders (L-Station).

8   Acknowledgments

We would like to thank Ms. Asuka Murata for technical support and Dr. Takeharu Nagai for generously providing us with a plasmid vector for SuperNova-Red.

9   Figure legends

Figure 1. Cell viability after chromophore-associated light inactivation (CALI) targeting G3BP1 in cells recovering from hyperosmotic stress. (A) Fluorescence images of SG-positive and SG-negative cells expressing SuperNova-Red (SNR)-tagged G3BP1 (G3BP1-SNR; magenta) and Histon H2B-GFP (green) before light irradiation for CALI. (B) Time-course cell viability plot (600 min) in the recovery medium after hyperosmotic stress with or without light irradiation for CALI (20 cells). P-values were obtained using the generalized Wilcoxon test (Gehan-Breslow method). ###$p < 0.001$ (compared to the group without light irradiation); **$p < 0.01$ between the lines; NS, not significant ($p \geq 0.05$).

Figure 2. Cell viability after chromophore-associated light inactivation (CALI) targeting TDP-43 in cells recovering from hyperosmotic stress. (A) Fluorescence images of SG-positive and SG-negative cells expressing SuperNova-Red (SNR)-tagged TDP-43 (TDP43-SNR; magenta) and Histon H2B-GFP (green) before light irradiation for CALI. (B) Time-course cell viability plot (600 min) in the recovery medium after hyperosmotic stress with or without light irradiation for CALI (20 cells). P-values were obtained using the generalized Wilcoxon test (Gehan-Breslow method). ##$p < 0.01$ (compared to the group without light irradiation); *$p < 0.05$ between the lines; NS, not significant ($p \geq 0.05$).

Figure 3. Comparison of cell viability after chromophore-associated light inactivation (CALI) between G3BP1 and TDP-43 in cells recovering from hyperosmotic stress. The plot lines are derived from the data shown in Figures 1B and 2B (20 cells). P-values were obtained using the generalized Wilcoxon test (Gehan-Breslow method). **$p < 0.01$ between the lines.

Figure 4. Disassembly of SGs containing G3BP1-GFP or TDP43-GFP after chromophore-associated light inactivation (CALI) targeting G3BP1-SNR or TDP43-SNR, respectively. (A) Time-lapse observation of cells harboring SGs of G3BP1-GFP or TDP43-GFP with (+) or without (-) light irradiation. (B) Plot showing the decrease in the normalized number of SGs containing G3BP1-GFP with (magenta) or without (green) light irradiation (mean ± 95%CI; 15 cells). *P*-values were obtained using Student's t-test. *p < 0.05 at the time indicated on the lines. (C) Plot showing the decrease in the normalized number of SGs containing TDP43-GFP with (magenta) or without (green) light irradiation (mean ± 95%CI; 15 cells). *P*-values were obtained using Student's t-test. *p < 0.05 at the indicated time points.

Figure 5. The possible mechanism of internal damage in SGs caused by light irradiation.

10   References

1.   D. Campos-Melo, Z. C. E. Hawley, C. A. Droppelmann and M. J. Strong: The Integral Role of RNA in Stress Granule Formation and Function. *Front Cell Dev Biol*, 9, 621779 (2021) doi:10.3389/fcell.2021.621779

2.   D. S. W. Protter and R. Parker: Principles and Properties of Stress Granules. *Trends Cell Biol*, 26(9), 668-679 (2016) doi:10.1016/j.tcb.2016.05.004

3.   D. Tauber, G. Tauber and R. Parker: Mechanisms and Regulation of RNA Condensation in RNP Granule Formation. *Trends Biochem Sci*, 45(9), 764-778 (2020) doi:10.1016/j.tibs.2020.05.002

4.   H. Zhou, J. Luo, K. Mou, L. Peng, X. Li, Y. Lei, J. Wang, S. Lin, Y. Luo and L. Xiang: Stress granules: functions and mechanisms in cancer. *Cell Biosci*, 13(1), 86 (2023) doi:10.1186/s13578-023-01030-6

5.   S. Das, L. Santos, A. V. Failla and Z. Ignatova: mRNAs sequestered in stress granules recover nearly completely for translation. *RNA Biol*, 19(1), 877-884 (2022) doi:10.1080/15476286.2022.2094137

6.   J. R. Wheeler, T. Matheny, S. Jain, R. Abrisch and R. Parker: Distinct stages in stress granule assembly and disassembly. *Elife*, 5 (2016) doi:10.7554/eLife.18413

7.   S. Anisimov, M. Takahashi, T. Kakihana, Y. Katsuragi, H. Kitaura, L. Zhang, A. Kakita and M. Fujii: G3BP1 inhibits ubiquitinated protein aggregations induced by p62 and USP10. *Sci Rep*, 9(1), 12896 (2019) doi:10.1038/s41598-019-46237-1

8.   R. Gutierrez-Garcia, S. Koyuncu, F. Hommen, S. Bilican, H. J. Lee, A. Fatima and D. Vilchez: G3BP1-dependent mechanism suppressing protein aggregation in Huntington's models and its demise upon stress granule assembly. *Hum Mol Genet*, 32(10), 1607-1621 (2023) doi:10.1093/hmg/ddac304

9.   S. Liu, S. Tian, T. Lin, X. He, J. Eze Ideozu, R. Wang, Y. Wang, D. Yue and H. Geng: G3BP1 regulates breast cancer cell proliferation and metastasis by modulating PKCzeta. *Front Genet*, 13, 1034889 (2022) doi:10.3389/fgene.2022.1034889

10.   B. Wolozin and P. Ivanov: Stress granules and neurodegeneration. *Nat Rev Neurosci*, 20(11), 649-666 (2019) doi:10.1038/s41583-019-0222-5

11.   P. Yang, C. Mathieu, R. M. Kolaitis, P. Zhang, J. Messing, U. Yurtsever, Z. Yang, J. Wu, Y. Li, Q. Pan, J. Yu, E. W. Martin, T. Mittag, H. J. Kim and J. P. Taylor: G3BP1 Is a Tunable Switch that Triggers Phase Separation to Assemble Stress Granules. *Cell*, 181(2), 325-345 e28 (2020) doi:10.1016/j.cell.2020.03.046

12.   T. Surrey, M. B. Elowitz, P. E. Wolf, F. Yang, F. Nedelec, K. Shokat and S. Leibler:
Chromophore-assisted light inactivation and self-organization of microtubules and motors. *Proc Natl*
Acad Sci U S A, 95(8), 4293-8 (1998) doi:10.1073/pnas.95.8.4293

13.   M. E. Bulina, D. M. Chudakov, O. V. Britanova, Y. G. Yanushevich, D. B. Staroverov, T. V.
Chepurnykh, E. M. Merzlyak, M. A. Shkrob, S. Lukyanov and K. A. Lukyanov: A genetically
encoded photosensitizer. *Nat Biotechnol*, 24(1), 95-9 (2006) doi:10.1038/nbt1175

14.   K. Jacobson, Z. Rajfur, E. Vitriol and K. Hahn: Chromophore-assisted laser inactivation in
cell biology. *Trends Cell Biol*, 18(9), 443-50 (2008) doi:10.1016/j.tcb.2008.07.001

15.   K. Takemoto, T. Matsuda, N. Sakai, D. Fu, M. Noda, S. Uchiyama, I. Kotera, Y. Arai, M.
Horiuchi, K. Fukui, T. Ayabe, F. Inagaki, H. Suzuki and T. Nagai: SuperNova, a monomeric
photosensitizing fluorescent protein for chromophore-assisted light inactivation. *Sci Rep*, 3, 2629
(2013) doi:10.1038/srep02629

16.   A. Kitamura, Y. Nakayama, A. Shibasaki, A. Taki, S. Yuno, K. Takeda, M. Yahara, N.
Tanabe and M. Kinjo: Interaction of RNA with a C-terminal fragment of the amyotrophic lateral
sclerosis-associated TDP43 reduces cytotoxicity. *Sci Rep*, 6, 19230 (2016) doi:10.1038/srep19230

17.   T. Kanda, K. F. Sullivan and G. M. Wahl: Histone-GFP fusion protein enables sensitive
analysis of chromosome dynamics in living mammalian cells. *Curr Biol*, 8(7), 377-85 (1998)
doi:10.1016/s0960-9822(98)70156-3

18.   A. Kitamura, S. Oasa, H. Kawaguchi, M. Osaka, V. Vukojevic and M. Kinjo: Increased
intracellular crowding during hyperosmotic stress. *Sci Rep*, 13(1), 11834 (2023) doi:10.1038/s41598-
023-39090-w

19.   J. O. Onukwufor, A. J. Trewin, T. M. Baran, A. Almast, T. H. Foster and A. P. Wojtovich:
Quantification of reactive oxygen species production by the red fluorescent proteins KillerRed,
SuperNova and mCherry. *Free Radic Biol Med*, 147, 1-7 (2020)
doi:10.1016/j.freeradbiomed.2019.12.008

20.   E. M. Hock, Z. Maniecka, M. Hruska-Plochan, S. Reber, F. Laferriere, M. K. S. Sahadevan,
H. Ederle, L. Gittings, L. Pelkmans, L. Dupuis, T. Lashley, M. D. Ruepp, D. Dormann and M.
Polymenidou: Hypertonic Stress Causes Cytoplasmic Translocation of Neuronal, but Not Astrocytic,
FUS due to Impaired Transportin Function. *Cell Rep*, 24(4), 987-1000 e7 (2018)
doi:10.1016/j.celrep.2018.06.094

21.   N. C. Shaner, R. E. Campbell, P. A. Steinbach, B. N. Giepmans, A. E. Palmer and R. Y.
Tsien: Improved monomeric red, orange and yellow fluorescent proteins derived from Discosoma sp.
red fluorescent protein. *Nat Biotechnol*, 22(12), 1567-72 (2004) doi:10.1038/nbt1037

22.   M. E. Bulina, K. A. Lukyanov, O. V. Britanova, D. Onichtchouk, S. Lukyanov and D. M.
Chudakov: Chromophore-assisted light inactivation (CALI) using the phototoxic fluorescent protein
KillerRed. *Nat Protoc*, 1(2), 947-53 (2006) doi:10.1038/nprot.2006.89

23.   M. Bregnhoj, M. Westberg, F. Jensen and P. R. Ogilby: Solvent-dependent singlet oxygen
lifetimes: temperature effects implicate tunneling and charge-transfer interactions. *Phys Chem Chem*
Phys, 18(33), 22946-61 (2016) doi:10.1039/c6cp01635a

24.   M. Klaper, W. Fudickar and T. Linker: Role of Distance in Singlet Oxygen Applications: A
Model System. *J Am Chem Soc*, 138(22), 7024-9 (2016) doi:10.1021/jacs.6b01555

25.   M. K. Kuimova, G. Yahioglu and P. R. Ogilby: Singlet oxygen in a cell: spatially dependent
lifetimes and quenching rate constants. *J Am Chem Soc*, 131(1), 332-40 (2009)
doi:10.1021/ja807484b

26.   D. G. Jay: Selective destruction of protein function by chromophore-assisted laser
inactivation. *Proc Natl Acad Sci U S A*, 85(15), 5454-8 (1988) doi:10.1073/pnas.85.15.5454

27.   Y. Sano, W. Watanabe and S. Matsunaga: Chromophore-assisted laser inactivation--towards a
spatiotemporal-functional analysis of proteins, and the ablation of chromatin, organelle and cell
function. *J Cell Sci*, 127(Pt 8), 1621-9 (2014) doi:10.1242/jcs.144527

28.   K. Takemoto, H. Iwanari, H. Tada, K. Suyama, A. Sano, T. Nagai, T. Hamakubo and T.
Takahashi: Optical inactivation of synaptic AMPA receptors erases fear memory. *Nat Biotechnol*,
35(1), 38-47 (2017) doi:10.1038/nbt.3710

29.   J. L. Ravanat and E. Dumont: Reactivity of Singlet Oxygen with DNA, an Update.
Photochem Photobiol, 98(3), 564-571 (2022) doi:10.1111/php.13581

30.   L. F. Agnez-Lima, J. T. Melo, A. E. Silva, A. H. Oliveira, A. R. Timoteo, K. M. Lima-Bessa,
G. R. Martinez, M. H. Medeiros, P. Di Mascio, R. S. Galhardo and C. F. Menck: DNA damage by
singlet oxygen and cellular protective mechanisms. *Mutat Res Rev Mutat Res*, 751(1), 15-28 (2012)
doi:10.1016/j.mrrev.2011.12.005

31.   M. J. Davies: Singlet oxygen-mediated damage to proteins and its consequences. *Biochem
Biophys Res Commun*, 305(3), 761-70 (2003) doi:10.1016/s0006-291x(03)00817-9

32.   E. Fuentes-Lemus, P. Hagglund, C. Lopez-Alarcon and M. J. Davies: Oxidative Crosslinking
of Peptides and Proteins: Mechanisms of Formation, Detection, Characterization and Quantification.
Molecules, 27(1) (2021) doi:10.3390/molecules27010015

33.   Y. Furukawa, R. Fu, H. X. Deng, T. Siddique and T. V. O'Halloran: Disulfide cross-linked
protein represents a significant fraction of ALS-associated Cu, Zn-superoxide dismutase aggregates
in spinal cords of model mice. *Proc Natl Acad Sci U S A*, 103(18), 7148-53 (2006)
doi:10.1073/pnas.0602048103

34.   A. Kitamura, N. Inada, H. Kubota, G. Matsumoto, M. Kinjo, R. I. Morimoto and K. Nagata:
Dysregulation of the proteasome increases the toxicity of ALS-linked mutant SOD1. *Genes Cells*,
19(3), 209-24 (2014) doi:10.1111/gtc.12125

35.   M. Garcia-Jove Navarro, S. Kashida, R. Chouaib, S. Souquere, G. Pierron, D. Weil and Z.
Gueroui: RNA is a critical element for the sizing and the composition of phase-separated RNA-
protein condensates. *Nat Commun*, 10(1), 3230 (2019) doi:10.1038/s41467-019-11241-6

36.   M. Herb, A. Gluschko and M. Schramm: Reactive Oxygen Species: Not Omnipresent but
Important in Many Locations. *Front Cell Dev Biol*, 9, 716406 (2021) doi:10.3389/fcell.2021.716406

37.   M. Briese, L. Saal-Bauernschubert, P. Luningschror, M. Moradi, B. Dombert, V. Surrey, S.
Appenzeller, C. Deng, S. Jablonka and M. Sendtner: Loss of Tdp-43 disrupts the axonal
transcriptome of motoneurons accompanied by impaired axonal translation and mitochondria
function. *Acta Neuropathol Commun*, 8(1), 116 (2020) doi:10.1186/s40478-020-00987-6

38.   F. C. Fiesel, C. Schurr, S. S. Weber and P. J. Kahle: TDP-43 knockdown impairs neurite
outgrowth dependent on its target histone deacetylase 6. *Mol Neurodegener*, 6, 64 (2011)
doi:10.1186/1750-1326-6-64

39.   M. Polymenidou, C. Lagier-Tourenne, K. R. Hutt, S. C. Huelga, J. Moran, T. Y. Liang, S. C.
Ling, E. Sun, E. Wancewicz, C. Mazur, H. Kordasiewicz, Y. Sedaghat, J. P. Donohue, L. Shiue, C. F.

Bennett, G. W. Yeo and D. W. Cleveland: Long pre-mRNA depletion and RNA missplicing
contribute to neuronal vulnerability from loss of TDP-43. *Nat Neurosci*, 14(4), 459-68 (2011)
doi:10.1038/nn.2779

40. M. Yahara, A. Kitamura and M. Kinjo: U6 snRNA expression prevents toxicity in TDP-43-
knockdown cells. *PLoS One*, 12(11), e0187813 (2017) doi:10.1371/journal.pone.0187813

41. N. P. Tsai and L. N. Wei: RhoA/ROCK1 signaling regulates stress granule formation and
apoptosis. *Cell Signal*, 22(4), 668-75 (2010) doi:10.1016/j.cellsig.2009.12.001

42. Y. J. Park, D. W. Choi, S. W. Cho, J. Han, S. Yang and C. Y. Choi: Stress Granule Formation
Attenuates RACK1-Mediated Apoptotic Cell Death Induced by Morusin. *Int J Mol Sci*, 21(15)
(2020) doi:10.3390/ijms21155360

43. C. Mao, X. Wang, Y. Liu, M. Wang, B. Yan, Y. Jiang, Y. Shi, Y. Shen, X. Liu, W. Lai, R.
Yang, D. Xiao, Y. Cheng, S. Liu, H. Zhou, Y. Cao, W. Yu, K. Muegge, H. Yu and Y. Tao: A
G3BP1-Interacting lncRNA Promotes Ferroptosis and Apoptosis in Cancer via Nuclear Sequestration
of p53. *Cancer Res*, 78(13), 3484-3496 (2018) doi:10.1158/0008-5472.CAN-17-3454

44. D. Mateju, B. Eichenberger, F. Voigt, J. Eglinger, G. Roth and J. A. Chao: Single-Molecule
Imaging Reveals Translation of mRNAs Localized to Stress Granules. *Cell*, 183(7), 1801-1812 e13
(2020) doi:10.1016/j.cell.2020.11.010

Figure 1

Figure 2

(A)

(B)

Figure 3

Figure 4

Figure 5

Not disassemble

Toxic/dysfunction

Slowly disassemble

CALI

Toxic/dysfunction

= Cross-linking

★ Modification

poly(A)+ mRNAs | G3BP1 | TDP-43 | Core proteins | Shell proteins | 40S ribosome | Other RNPs | Shell phase | Core phase