Overexpression of tomato *SlBBX16* and *SlBBX17* impacts fruit development and GA biosynthesis

Valentina Dusi[1][§], Federica Pennisi[1][§], Daniela Fortini[1], Alejandro Atarès[2], Stephan Wenkel[3,4], Barbara Molesini[1], and Tiziana Pandolfini[1*]

[1]Department of Biotechnology, University of Verona, Strada Le Grazie, 15 - 37134, Verona, Italy

[2]Instituto de Biología Molecular y Celular de Plantas (UPV-CSIC), Universitat Politècnica de València, Ingeniero Fausto Elio s/n, 46011, Valencia, Spain.

[3]Department of Plant Physiology, Plant Science Centre, University of Umeå, Linnaeus väg 6, 907 36 Umeå, Sweden

[4]NovoCrops Center, University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg, Denmark

[§]These authors contributed equally to the article.

*Corresponding Author: tiziana.pandolfini@univr.it

Keywords: BBX, microProteins, flowering time, fruit development, *Solanum lycopersicum*, *Arabidopsis thaliana*, gibberellins

Abstract

BBX proteins are B-Box zinc finger proteins that can act as transcription factors and regulators of protein complexes. Several BBX proteins play important roles in plant development. Two *Arabidopsis thaliana* microProteins belonging to the BBX family, named miP1a and miP1b, homotypically interact with and modulate the activity of other BBX proteins, including CONSTANS (*At*CO), which transcriptionally activates the florigen gene, *FLOWERING LOCUS T* (*FT*). In tomato, the closest homologs of miP1a and miP1b are the microProteins *Sl*BBX16 and *Sl*BBX17. To deepen our understanding of the role of tomato microProteins in flowering, we constitutively expressed *Sl*BBX16/17 in *Arabidopsis* and tomato (MicroTom) and examined possible interacting partners. Overexpression of the two tomato microProteins in *Arabidopsis* caused a delay in the flowering transition; however, the effect was weaker than that observed in *Arabidopsis* plants overexpressing the native miP1a/b microProteins. In tomato, overexpression of *SlBBX17* prolonged the flowering period; this effect was accompanied by downregulation of the flowering inhibitor *Self Pruning*. Yeast-Two Hybrid analysis did not show a direct interaction between *Sl*BBX16 and *Sl*BBX17 and *At*CO and its closest homolog in tomato, *Sl*COL1. On the other hand, *Sl*BBX16 and *Sl*BBX17 were able to hetero-oligomerize with TCMP-2, a cystine-knot peptide involved in flowering pattern and fruit development in tomato. Increasing the expression of both tomato microProteins also caused alterations in fruit development: overexpression of *SlBBX17* resulted in a diminished number and size of ripe fruits as compared to WT plants, while overexpression of *SlBBX16* caused delayed fruit production up to the breaker stage. These effects were associated with changes in the expression of genes regulating gibberellin content.

Introduction

Crop productivity is the result of the interaction between the plant genetic background, the environmental cues and the agronomic practices. In the face of climate change, it becomes increasingly important to match the phenology of crop cultivars with the environmental conditions to maintain high crop yields. Therefore, the availability of cultivars with different phenological requirements could be advantageous for optimising plant reproduction and productivity. In horticultural plants, flowering, fruit set and the onset of ripening are processes markedly affected by temperature and photoperiod, and regulated by internal signals such as hormones, transcription factors and adaptor molecules. Among the different regulatory proteins known to control reproductive development, members of the B-Box (BBX) protein family have emerged as important molecular players that integrate environmental cues, for instance light and temperature, with endogenous signalling pathways (Gangappa and Botto 2014; Silvestre Lira et al., 2020; Yadav et al., 2020).

The BBX family represents a group of zinc (Zn)-finger proteins that are involved not only in reproductive development, but also in many other physiological processes, such as photomorphogenesis (Crocco et al.,2010; Holtan et al., 2011; Fan et al., 2012), anthocyanin accumulation, seed germination, carotenoid biosynthesis, and responses to biotic and abiotic stresses (Wang et al., 2013; Kiełbowicz-Matuk et al., 2014; Vaishak et al., 2019; Xu et al., 2022). The BBX family is characterized by the presence of one or two Zn-finger-containing BBX domain(s) in the N-terminal region. Depending on their consensus sequence and the distance between the Zn binding residues, two types of BBX domains have been described, B1 and B2 (Khanna et al., 2009; Gangappa and Botto, 2014; Graeff et al.,2016). Previous studies suggested that the B-Box domain plays a crucial role in protein-protein interactions. Some BBX proteins may possess a CCT domain, which is associated with a role in transcriptional regulation and nuclear transport (Gendron et al., 2012; Crocco and Botto, 2013). CCT-containing BBXs are likely transcription factors (TFs), while BBXs lacking

CCT domain might have other functions, linked to their protein-protein interaction capacity. *Arabidopsis* BBX proteins have been classified into five subgroups according to different combinations of the above-mentioned domains. Members of group I are characterized by the presence of B1, B2, and CCT domains, as are members of group II; however, some differences have been observed between the two groups in the B2 consensus sequences (Chang et al., 2008; Crocco and Botto, 2013). The presence of B1 and CCT, B1 and B2, and a single B1 domain characterizes members of group III, IV, and V, respectively (Khanna et al., 2009; Gangappa and Botto, 2014; Graeff et al., 2016). Following the same criteria, BBX proteins from other species such as tomato, potato, rice, and maize have been grouped into five subfamilies (Huang et al., 2012; Chu et al., 2016; Li et al., 2017; Talar et al., 2017). *A. thaliana* CONSTANS (*At*CO), one of the first BBX proteins to be identified and characterized, belongs to subgroup I. *At*CO plays a crucial role in photoperiodic control of flowering time, enabling the transcriptional activation of the *FLOWERING LOCUS T* (*FT*) in response to long day (LD) conditions. Proteins homologous to *At*CO have been shown to contribute to flowering regulation in other plants species with different photoperiodic requirements (Doi et al., 2004; Miller et al., 2008; Meng et al., 2011; Campoli et al., 2012; Kikuchi et al., 2012; Yang et al., 2014; Ping et al., 2019; Wang et al., 2019).

Other members of the *Arabidopsis* BBX family were successively discovered to be implicated in flowering control (Cheng and Wang, 2005; Park et al., 2011; Li et al., 2014; Tripathi et al., 2017). Recently, two *Arabidopsis* BBX proteins of the group V, microProtein (miP) miP1a and miP1b, (also referred as *At*BBX31 and *At*BBX30, respectively) were shown to modulate *At*CO activity (Graeff et al., 2016; Rodrigues et al., 2021). MiP1a and miP1b can interact with both *At*CO and TOPLESS (TPL), leading to the formation of a trimeric complex that limits *At*CO-mediated induction of *FT* expression. Consistently, *Arabidopsis* plants overexpressing miP1a and miP1b, grown under LD conditions, showed delayed flowering (Graeff et al., 2016). In tomato, the microProteins *Sl*BBX16 and *Sl*BBX17 are the

closest homologs of miP1a and miP1b, respectively, but their role in tomato reproductive development is largely elusive. The features of group V BBXs - a single B-Box domain and the lack of CCT - suggest that they act as microProteins regulating other larger multidomain protein complexes at the post-translational level (Eguen et al., 2015).

In this regard, *Sl*BBX16 was found to interact with the tomato cystine-knot peptide 2 (TCMP-2) (Molesini et al., 2020). TCMP-2 is specifically expressed in reproductive organs, its expression is low in pre-anthesis flower buds and gradually increases after fertilization, reaching a maximum in green and ripe fruits (Pear et al., 1989; Cavallini et al., 2011; Treggiari et al., 2015). Increased TCMP-2 expression in pre-anthesis flower buds has been demonstrated to lead to alterations in flowering pattern as well as early fruit production and a slight delay in the initiation of ripening (Molesini et al., 2018; Molesini et al., 2020).

In tomato, which is a day-neutral species, the flowering transition is principally regulated by the balance between the activity of the florigen *Single Flower Truss* (*SFT*), which is the ortholog of *FT*, and that of *Self Pruning* (*SP*), which maintains vegetative growth. Recently, it has been shown in tomato that *Sl*COL1, the ortholog of *At*CO, is able to bind the promoter region of *SFT* to repress its expression (Cui et al., 2022). Accordingly, RNA silencing of *SlCOL1* led to promotion of flowering and increased fruit yield. Our work was aimed at elucidating the functional role of *Sl*BBX16 and *Sl*BBX17 in tomato reproductive development. We demonstrated that *SlBBX17* overexpressing (*Sl*BBX17OE) plants show a prolonged period of flowering associated with a reduced expression of the *SP* flowering inhibitor. Furthermore, we observed in *Sl*BBX16OE plants a delay in the early phases of fruit growth and in *Sl*BBX17OE plants changes in ripening. The overexpression of both microProteins induced modifications in the expression pattern of genes regulating gibberellin (GA) metabolism, such that they can lead to reduced GA level in the fruit.

Results

Ectopic overexpression of *Sl*BBX16 and *Sl*BBX17 influences the flowering transition in *Arabidopsis*

The microProteins *Sl*BBX16 and *Sl*BBX17 show high sequence similarity to miP1a and miP1b (Figure S1A), which control flowering in *Arabidopsis* and, when overexpressed, delay flowering (Graeff et al., 2016). To test whether *Sl*BBX16 and *Sl*BBX17 can exert a similar effect, we analysed the flowering behaviour of *Arabidopsis Sl*BBX16OE and *Sl*BBX17OE plants compared to Col-0 WT (Figure 1 and Figure S1B, C). In the *Sl*BBX16OE plants, the number of rosette leaves at bolting did not differ from the control plants (Figure 1A). Also, the days from sowing to flowering did not vary (Figure 1B). On the other hand, both *Sl*BBX17OE lines displayed an increased number of rosette leaves at bolting and a longer time to reach flowering under LD photoperiodic conditions (Figure 1C, D). The observed inhibitory effect of *Sl*BBX17 on flowering was less pronounced than that produced by overexpressing miP1a and miP1b in *Arabidopsis* (Graeff et al., 2016). To examine whether *Sl*BBX16 and *Sl*BBX17 can substitute the function of miP1a/b, we overexpressed them also in *Arabidopsis* miP1a/b double KO mutant, which displays earlier flowering than WT (Heng et al., 2019). In both *Sl*BBX16OE lines and in the *Sl*BBX17OE line #5, the number of leaves at bolting (Figure 1E, G) increased significantly (*Sl*BBX16OE #2 13.3 ± 1.5; #3 13.6 ±1.9; *Sl*BBX17OE #5 12.7 ± 1.6) compared to miP1a/b double KO mutant (11.7± 1.2) and was slightly lower than or similar to Col-0 WT (13.6 ±0.3). The number of days to bolting was increased significantly in *Sl*BBX16OE line #2 (31.7± 1.6) and in both *Sl*BBX17OE lines (33.2 ± 3.3 and 34.4 ± 3.1 days for lines #1 and #5, respectively) reaching values similar to those observed in Col-0 WT (35 ± 0.3) (Figure 1F, H). These results suggest that *Sl*BBX16 and *Sl*BBX17 may impact the flowering transition in *Arabidopsis* and partially rescue the function of endogenous miP1a/b. Since the flowering delay exhibited by *Sl*BBX16OE and *Sl*BBX17OE plants resembles, albeit in an attenuated form, the phenotype shown by

miP1a/b-overexpressing plants (Graeff et al.,2016), we examined via Y2H whether *Sl*BBX16

and *Sl*BBX17 interact with *At*CO as already demonstrated for miP1a/b (Graeff et al., 2016

and Figure S3). We did not observe a direct interaction between *Sl*BBX16 and *Sl*BBX17 and

*At*CO (Figure S2).

Overexpression of *Sl*BBX16 alters fruit set and delays fruit growth in MicroTom

We recently demonstrated that *Sl*BBX16 interacts with TCMP-2, a cystine-knot

metallocarboxypeptidase inhibitor (Molesini et al., 2020). TCMP-2 is specifically expressed

in reproductive organs (Cavallini et al., 2011; Molesini et al., 2018) and, when overexpressed

in flower buds, caused alteration in flowering pattern and early fruit setting in the tomato

cultivar UC82 (Molesini et al., 2018; Molesini et al. 2020). When *TCMP-2* is globally

overexpressed (i.e., using the *CaMV 35S* promoter) in the cultivar MicroTom, we have

observed an anticipated formation of the primary inflorescence and a reduction in plant

height, suggesting an accentuated determinate habit (Figure S3). The fruit set calculated on

the first three inflorescences did not change significantly (data not shown). We then

analyzed the role of *Sl*BBX16 microProtein in reproductive development by overexpression

in MicroTom. The three overexpressing lines #3, #4, and #20 (Figure S4A) showed no

changes in the number of leaves before the first inflorescence compared to the WT plants

(Figure 2A). The shoot height at the first inflorescence was significantly reduced only in the

*Sl*BBX16OE line #3 (Figure 2B). These data suggest that flowering transition was not greatly

affected by the *Sl*BBX16 overexpression. In addition, the number of flowers measured in the

first three inflorescences was comparable in the WT and *Sl*BBX16OE lines (Figure 2C).

However, fruit set, calculated as a percentage of number of fruits over number of flowers in

the first three inflorescences, was significantly lower in the *Sl*BBX16OE lines than in the WT

plants (Figure 2D). We have also observed a delay in the initial fruit formation and the onset

of maturation assessed as number of fruits at the green stage (Figure 3A) and number of

fruits at the breaker stage or in maturation (Figure 3B) overtime, respectively. Considering

the total fruit production harvested at 110 days after sowing, in the *Sl*BBX16OE lines there

was a partial compensation of the fruit set impairment in the first three inflorescences (Figure

3C). The weight of WT and transgenic fruits was similar (Figure 3D), as was the percentage

of red and green fruits (data not shown).

Overexpression of *SlBBX17* prolongs flowering and reduces ripe fruit production in

MicroTom

The microProteins *Sl*BBX16 and *Sl*BBX17 are highly homologous, with 46% identity (Figure

S1A). The two genes have similar patterns of expression in flowers and fruits and are co-

expressed with *TCMP-2* (Figures S4B and S5A, Cavallini et al., 2011, Molesini et al., 2020).

Furthermore, Y2H analysis suggests that *Sl*BBX17 and *Sl*BBX16 share a common

interacting partner, TCMP-2 (Figure S5B and Molesini et al. 2020).

We investigated the phenotypic consequences of *SlBBX17* overexpression in MicroTom by

comparing two independent *Sl*BBX17OE transgenic lines (#15, and #18) to WT plants

(Figure 5C). Differently from what was previously reported by overexpressing *SlBBX17* in

the indeterminate cultivar Ailsa Craig (Xu et al., 2022), we did not observe retardation in

vegetative growth and a reduction in leaf size (data not shown). However, because of the

stunted growth of MicroTom it is likely that these growth effects were masked by other

mutations. In agreement with Xu and collaborators (2022), we did observe moderate

resistance to high temperatures (Figure S5D). When we monitored the transition from

vegetative to reproductive development in *Sl*BBX17OE plants, we observed a reduction in

the height of the shoot bearing the primary inflorescence compared to WT plants, but no

changes in the number of leaves before the first inflorescence (Figure 4A, B). The number

of flowers in the first three inflorescences was similar in the WT and transgenic lines (Figure 4C), but when recording the number of flowers at anthesis over time, we noticed a different trend in the WT and transgenic plants (Figure 4D, E). In the WT plants, the number of open flowers increased from 39 to 47 days after sowing (das) but decreased sharply thereafter. In the transgenic plants, we observed a tendency to persist in producing flowers over time: in line #15 the number of flowers was higher than in WT from 61 das onwards and in line #18 this effect was detected from 75 das onwards. Given the differences in the duration of flowering between WT and *Sl*BBX17OE plants (Figure 4D, E), we analysed the expression of *SFT*, the ortholog of *FT*, and *SP*, a flowering inhibitor. The dynamic ratio between SFT and SP regulates shoot termination in tomato, which is associated with flowering (Lifschitz et al., 2014). The expression of *SFT* did not vary, whereas the expression of *SP* was reduced in the transgenic plants compared to WT (Figure 4F, G). This suggests that the prolonged flowering observed in *Sl*BBX17OE might be associated with a weaker antagonistic effect of SP on SFT. *Solanum lycopersicum* is a day-neutral plant and the mechanism controlling the flowering process in this species is not fully elucidated. In this regard, the biological functions of CO-like (COL) genes in tomato remain elusive. Recently, *Sl*COL1 (Solyc02g089540) was shown to interact with the *SFT* promoter repressing its transcription and consequently inhibiting the flowering transition (Cui et al., 2022). Considering that miP1a/b interacts with *At*CO regulating its activity, we tested by Y2H the interaction between *Sl*BBX16 and *Sl*BBX17 and *Sl*COL1. The absence of interaction between tomato BBX proteins and *Sl*COL1, would suggest that either *Sl*BBX16 and *Sl*BBX17 do not bind tightly to *Sl*COL1 under our experimental conditions or that their interaction requires additional factors that are absent in yeast (Figure S2).

We then evaluated the fruiting process in the *Sl*BBX17OE lines assessing fruit set and fruit productivity. The fruit set recorded in the first three inflorescences and the total number of fruits per plant evaluated 4 months after sowing were similar to WT plants (Figure 5A, B).

However, the proportion of ripe fruits out of the total number of fruits in the transgenic lines was reduced (Figure 5C, F). In addition, ripe fruits were smaller and contained fewer seeds (Figure 5C, E). We have also observed a different weight distribution of green fruits with a decreasing trend in *Sl*BBX17OE fruits (Figure S6).

Overexpression of *SlBBX16* and *SlBBX17* affects the expression of GA biosynthetic genes

To elucidate if *SlBBX16* and *SlBBX17* affect the hormone homeostasis that controls fruit development, we decided to assay the expression of the hormone biosynthesis genes in immature green fruits, a phase of active growth and preparation for ripening. We examined in *Sl*BBX16OE and *Sl*BBX17OE fruits the expression of a set of genes implicated in ethylene, abscisic acid (ABA), and gibberellin (GA) metabolism. *Aminocyclopropane-1-carboxylic acid synthase 1A* (*ACS1A*) and *ACS6* encode enzymes of the ethylene biosynthetic system 1 that triggers the transition to breaker stage in green fruit (Barry et al., 2000; Alexander and Grierson, 2002). Their expression did not vary in *Sl*BBX16OE and *Sl*BBX17OE fruits (Figure 6A, B, G, H), suggesting that ethylene biosynthesis at this stage is similar to that in WT fruits. To test a possible alteration of ABA metabolism, we analysed the expression of *9-cis-epoxycarotenoid dioxygenase* (*SlNCED-1*)*,* the key determinant of ABA synthesis, and *SlZFP2,* a zinc finger transcription factor involved in the crosstalk between ABA and ethylene in the regulation of fruit ripening in tomato (Weng et al., 2015). The expression of *SlNCED-1* increased in two (#3 and #4) out of the three *Sl*BBX16OE lines, while it remained unaltered in *Sl*BBX17OE fruits compared to WT (Figure 6D, J). The transcript level of *SlZFP2* was unchanged in *Sl*BBX16OE and *Sl*BBX17OE fruits (Figure 6C, I). A reduced expression of the GA biosynthetic gene, *GA20 oxidase 2* (*SlGA20ox2*), was recorded in the three *Sl*BBX16OE lines and in the *Sl*BBX17OE line#15 (Figure 6F, L). On

the other hand, in the *Sl*BBX17OE line#18, *SlGA2ox4*, a gene encoding an enzyme responsible for GA catabolism, was induced (Figure 6K). These data suggest that the phenotypes observed in *Sl*BBX16OE and *Sl*BBX17OE fruits might be due to altered GA metabolism. Interestingly, treatment with exogenous GA induced the expression of both *SlBBX16* and *SlBBX17* (Figure S7).

Discussion

Plant microProteins are small polypeptides characterized by a single domain involved in protein-protein interaction. From an evolutionary point of view, miPs can originate from larger proteins by duplication and domain loss (*trans*-miPs) or derive from alternative splicing or alternative transcription start and stop sites (*cis*-miPs) (Eguen et al., 2015; Kushwaha et al., 2022). Their capacity to interact with homologous proteins of larger size (homotypic interaction) is at the basis of their activity as posttranslational regulators of protein complexes. Besides homotypic, plant miPs can engage also in heterotypic interactions with evolutionary unrelated proteins, widening the regulatory role of these proteins (Bhati et al., 2021).

Arabidopsis miP1a and miP1b are microProteins belonging to subclass V of the BBX family. They play a role in photomorphogenic development and control of flowering time. MiP1a and miP1b participate in seedling growth arrest after germination under stress conditions by interacting heterotypically with ABA-insensitive 5 (ABI5) and stabilizing it (Singh and Datta, 2023). During the transition from dark to light conditions, miP1a and miP1b inhibit the oligomerization of PHYTOCHROME-INTERACTING FACTORs (PIFs) and ETHYLENE-INSENSITIVE 3 (EIN3) (Wu et al., 2020). A homotypic interaction between miP1a/b and *At*CO regulates florigen *FT* transcription; the strong delay in flowering observed by

overexpressing miP1a/b in *Arabidopsis* is caused by the recruitment of TOPLESS to miP1a/b-CO complex resulting in inhibition of *FT* transcription (Graeff et al., 2016).

In tomato, *Sl*BBX16 and *Sl*BBX17 microProteins are the closest homologs of miP1a/b. A recently published study by Xu and collaborators (2022), reported that *SlBBX17* overexpression in an indeterminate tomato cultivar (i.e., Ailsa Craig), resulted in increased heat tolerance and growth retardation (Xu et al., 2022). The effects of *SlBBX17* overexpression on reproductive development were not investigated in that study. The first hints on the implication of *Sl*BBX17 and *Sl*BBX16 in reproductive development derive from the observation that *Sl*BBX16 and *Sl*BBX17 undergo heterotypic interaction with a tomato cystine-knot peptide, TCMP-2 (this work and Molesini et al., 2020), which plays a role in both flowering pattern and fruit development. In the tomato cultivar UC82, overexpression of *TCMP-2* in flower buds promoted the termination of sympodial units (Molesini et al., 2020) and global overexpression in MicroTom resulted in the early appearance of the first inflorescence (this work). In addition, ectopic overexpression of *TCMP-2* anticipated flowering in *Arabidopsis* (Molesini et al., 2020) and in both tomato and *Arabidopsis*, TCMP-2 induced the transcription of florigen. To investigate if TCMP-2 partners play a role in flowering, *SlBBX16* and *SlBBX17* were overexpressed in *Arabidopsis* WT Col-0 and miP1a/b double KO mutant and in tomato. The *Arabidopsis Sl*BBX16OE and *Sl*BBX17OE plants displayed a weak delay in flowering; while in tomato, overexpression of *SlBBX16* caused no alterations in flowering, and overexpression of *SlBBX17* resulted in protracted flowering associated with a reduced expression of the antiflorigen *SP*. Thus, overexpression of *TCMP-2* and its partners produced contrasting effects on flowering, suggesting an antagonistic interaction. The evidence that both TCMP-2 and *Sl*BBX16/17 showed no direct interaction by Y2H with *At*CO and *Sl*COL1 (this work and Molesini et al., 2020) suggests that a system similar to the miP1a/b-based flowering control of *Arabidopsis* is not operating in tomato. In this regard, *Sl*COL1 has been shown to bind the *SFT* promoter in tomato and,

when overexpressed, altered flower development (Cui et al., 2022). Based on our results, the activity of TCMP-2 and *Sl*BBX17 appeared to be independent of the *Sl*COL1/SFT regulatory module. We cannot exclude that *Sl*BBX16 and *Sl*BBX17 can interact with other members of the *Sl*COL family.

Our results demonstrate that *Sl*BBX16 and *Sl*BBX17 also participate in the regulation of fruit growth and maturation. In a recent paper, Lira and collaborators evaluated the transcript profiles of *SlBBX16* and *SlBBX17* in tomato fruits at different stages of development, starting from immature green (i.e., IG3 corresponding to 8 days post-anthesis) up to 5 days after the breaker stage (Lira et al., 2020). During the stages of green fruit growth (i.e., IG phases), the expression of *SlBBX16* progressively decreased (Lira et al., 2020). The expression pattern of *SlBBX16* might indicate that during the period of fruit enlargement its activity must be kept low. This observation is consistent with the phenotype of *Sl*BBX16OE plants, in which fruit growth is delayed from the early green phases (comparable to IG3) until the breaker stage. It is plausible that *Sl*BBX16 is one of the factors that restricts ovary expansion; therefore, its progressively reduced expression would be necessary for optimal fruit growth. The reduced fruit set observed in the first three inflorescences of *Sl*BBX16OE plants would support this hypothesis. The expression of *SlBBX17* is not strongly modulated during fruit development as for *SlBBX16* (Lira et al., 2020); its transcript level increases weakly from IG3 to the breaker stage and then declines shortly after (Lira et al., 2020). In the present work, the major effect of *SlBBX17* overexpression was observed at the ripening stage. The percentage of ripe to total fruits was markedly reduced, suggesting that the excess of this microProtein at the maturation phase hinders this process. The reduced weight of ripe *Sl*BBX17OE fruits and the similar tendency observed for green fruits, indicate that *Sl*BBX17 is also involved in fruit enlargement. Although the two microProteins seem to contribute to regulate fruit development at different stages, they both show negative regulatory activities which is consistent with the inhibitory capacity of microProteins on their

target partners. In this regard, the overexpression of their partner *TCMP-2* in the flower buds before anthesis, accelerated the formation of fruits in cultivar UC82 (Molesini et al., 2018).

Gibberellin (GA) appears to play an important role in the mechanisms of action of both *Sl*BBX16 and *Sl*BBX17 because the expression of genes regulating active GA levels is altered in transgenic immature green fruits. Remarkably, both *SlBBX16* and *SlBBX17* are light and gibberellin responsive (this work and Chu et al., 2016; Lira et., 2020).

In tomato, GA presents a bimodal accumulation pattern (Gillaspy et al., 1993) controlling different phases of fruit development. A high GA level promotes fruit set and induces fruit growth through cell expansion, while a low level is detected from the mature green stage to the breaker (Li et al., 2019) due to downregulation of GA-biosynthetic genes, such as *GA3ox1* and *GA20ox2*, and upregulation of GA deactivation genes, such as *GA2ox* (Zouine et al., 2017; Li et al., 2019). Consistently, exogenous application of GA$_3$ to mature green fruit delayed fruit ripening (Li et al., 2019). In *Sl*BBX16OE and *Sl*BBX17OE fruits at the immature green stage (IG3), the modulation of genes regulating GA levels suggests a decline in GA content incongruous with this developmental stage. The effect of this hormonal change would on one hand counteract cell expansion, delaying fruit growth, and on the other hand, would affect the transition to ripening (Figure 7). In *Sl*BBX16OE and *Sl*BBX17OE fruits, the expression of the ethylene biosynthetic genes (*ACS1A* and *ACS6*) remained unaltered indicating that the induction of ethylene biosynthesis system1 has not jet induced at this stage (IG3). Furthermore, the activity of *Sl*BBX16, but not that of *Sl*BBX17, seems also associated with an increased biosynthesis of ABA.

It will be critical for future research to decipher the regulatory functions of these tomato microProteins in the reproductive process, investigating whether they may be part of a multiprotein complex together with TCMP-2. Ultimately, a better understanding of the mechanism of action by which these microProteins alter GA homeostasis will provide

opportunities for their potential application in tomato breeding to match the reproductive

phases with changes in climate conditions.

Materials and Methods

Plant materials

Solanum lycopersicum MicroTom WT seeds (ID:TOMJPF00001) were obtained from the

TOMATOMA mutant archive (Saito et al., 2011).

Arabidopsis thaliana WT (ecotype Col-0) and *At*miP1a/b double KO mutant plants were

employed for flowering time assessment.

Plant genetic transformation

The tomato DNA sequences corresponding to the coding regions of *SlBBX16*

(Solyc12g005750) and *SlBBX17* (Solyc07g052620) were amplified by PCR from cDNAs

using the primers reported in Table S1. The DNA fragments were subcloned into the

pGEM®-T Easy Vector (Promega) and checked by sequencing. The coding regions of

SlBBX16 and *SlBBX17* were then cloned into a derivative of the pBin19 vector, under the

control of the *CaMV35S* promoter and the terminator sequence of the *Agrobacterium*

tumefaciens nopaline synthase gene.

For the overexpression of *TCMP-2* (Solyc07g049140) in MicroTom, a sequence

corresponding to the coding region was amplified using the Gateway System (Invitrogen)

(Table S1). After subcloning in the pDONR221, the resulting pENTRY vector was checked

by sequencing and used for recombination in the destination vector pK7WG2D.1 (Karimi et

al., 2002). The recombinant vectors obtained were introduced into *Agrobacterium*

tumefaciens cells (strain GV2260). The genetic transformation of MicroTom was obtained

from cotyledon explants of 8 days-old seedlings. *A. tumefaciens* cells were grown at 28°C

for 24 hours and used at OD_{600} of 0.1 for explant infection. After 48 hours of co-cultivation,

the explants were transferred to culture medium containing NAA (0.01 mgL^{-1}), zeatin

riboside (2 mgL^{-1}) and kanamycin (100 mgL^{-1}). The regenerated shoots were transferred to

rooting medium supplemented with kanamycin (75 mgL^{-1}). After 3-4 weeks, the rooted plants

were acclimatized in the greenhouse. Monitoring the ploidy level of putative transformants

according to Atares et al., 2011, we demonstrated that with this transformation method, the

vast majority (about 90%) of MicroTom transformed lines retained the diploid state.

Arabidopsis plants were transformed using the floral dip method (Zhang et al., 2006).

Phenotypic analysis

Tomato plants were grown in the greenhouse during springtime. For phenotypic

assessment, plants of the T1 generation were grown in pots and transgenic state was

confirmed by spraying with kanamycin (400 mgL^{-1}). Various flowering and fruiting

parameters were recorded, and the fruit yield was evaluated at about 110 days after sowing.

Arabidopsis plants were grown in a climatic chamber at a constant temperature of 25°C

under LD conditions (16/8 hours light/dark cycle, photosynthetic photon fluence rate of 150

µmol m^{-2} s^{-1}). Homozygous plants of the T3 generation were used for flowering time

analysis.

Yeast Two-Hybrid analysis

To examine protein-protein interactions, the Matchmaker Gold Yeast Two-Hybrid System

(Clontech) was used, following the manufacturer's instructions with minor modifications. To

test the interaction between TCMP-2 and *Sl*BBX17, the DNA sequence of the mature portion

of the TCMP-2 protein (Solyc07g049140; from amino acid 53 to 96) was expressed as a

fusion with the DNA-binding domain of GAL4 in the pGBKT7-BD vector, while the entire

coding region of *Sl*BBX17 (Solyc07g052620) was cloned in frame into pGADT7-AD. For the

interactions between *Sl*BBX16 (Solyc12g005750), *Sl*BBX17, *At*CO (*At*5g15840) and

*Sl*COL1 (Solyc02g089540), the entire coding regions of the BBX genes were cloned in frame

into the pGBKT7-BD vector and *At*CO and *Sl*COL1 were cloned in the pGADT7-AD vector.

The interaction between miP1a (At3g21890) and *At*CO represents the positive control. For

negative controls, pGBKT7 without insert (BD alone; Empty) and pGADT7 without insert (AD

alone; Empty) were used. The primers employed for the genetic constructs preparation are

reported in Table S1.

RT-qPCR analysis

Total RNA extraction was performed from leaves, floral organs and fruits using the

"NucleoSpin RNA Plant" kit (Macherey-Nagel). After DNase I treatment, first-strand cDNA

was synthesized using the ImProm-II Reverse Transcriptase (Promega). For each mRNA

level determination, three cDNA samples derived from three independent RNA extractions

were synthesized. The cDNA was amplified using Luna®Universal qPCR Master Mix (New

England Biolabs, Ipswich, MA, USA) on a QuantStudio 3 Real-Time PCR System (Thermo

Fisher Scientific).   The data were normalized using *a*ctin (Solyc11g005330) or *SAND*

(Solyc03g115810) as references for leaf and fruit tomato samples, respectively, and actin

(At3g18780) for genes expressed in *Arabidopsis*. Data analysis was performed using the $2^{-\Delta\Delta Ct}$

method (Livak and Schmittgen, 2001).

GA treatment

To test the responsiveness of *SlBBX16* and *SlBBX17* to the exogenous application of GA

(GA$_3$), tomato seedlings were grown *in vitro* for one month on half- strength MS agar medium

(pH 5.9). The plants were placed in half-strength MS liquid solution (pH 5.9) overnight. The

next day, GA$_3$ was added with a final concentration of 5 μM. The treatment was conducted

for 24 h, then the shoots were collected for expression analysis by RT-qPCR.

Statistical analysis

Statistical analyses were performed using GraphPad Prism version 5.0 software (GraphPad

Software, San Diego California USA). Data were compared using Student's t-test.

Acknowledgements

This research was supported by a MIUR grant (PRIN2017, number 20173LBZM2) given to

T.P. University of Verona, Italy. The laboratory of SW receives funding through NovoCrops

Centre (Novo Nordisk Foundation project number 2019OC53580), the Independent

Research Fund Denmark (0136- 00015B and 0135-00014B) and the Novo Nordisk

Foundation (NNF18OC0034226 and NNF20OC0061440).

Conflict of interest

The authors declare that they have no conflict of interest.

Author contributions

B.M., and T.P. conceived and designed this work; V.D., F.P., and D.F carried out the

molecular analyses and the genetic transformation of *Arabidopsis* and tomato plants; A.A

contributed to the analysis of the transformed plants; S.W. provided *Arabidopsis* mutant;

B.M., S.W., and T.P. discussed the results, B.M. and T.P. wrote the manuscript. All authors

read and approved the final manuscript.

References

Alexander L, Grierson D (2002) Ethylene biosynthesis and action in tomato: a model for
climacteric fruit ripening. J Exp Bot **53:** 2039-2055

Atarés A, Moyano E, Morales B, Schleicher P, García-Abellán JO, Antón T, García-Sogo B,
Perez-Martin F, Lozano R, Flores FB, Moreno V, Bolarin Mdel C, Pineda B. (2011) An
insertional mutagenesis programme with an enhancer trap for the identification and
tagging of genes involved in abiotic stress tolerance in the tomato wild-related species
Solanum pennellii. Plant Cell Rep. 2011 **30**:1865-79.

Barry CS, Immaculada Llop-Tous M, Grierson D (2000) The regulation of 1-
aminocyclopropane-1-carboxylic acid synthase gene expression during the transition
from system-1 to system-2 ethylene synthesis in tomato. Plant Physiol **123:** 979–986

Bhati KK, Dolde U, Wenkel S (2021) MicroProteins: Expanding functions and novel modes
of regulation. Molecular Plant **14:** 705–707

Campoli C, Drosse B, Searle I, Coupland G, Von Korff M (2012) Functional characterisation
of HvCO1, the barley (Hordeum vulgare) flowering time ortholog of CONSTANS. Plant
J **69:** 868–880

Cavallini C, Trettene M, Degan M, Delva P, Molesini B, Minuz P, Pandolfini T (2011) Anti-
angiogenic effects of two cystine-knot miniproteins from tomato fruit. Brit J Pharmacol
162: 1261-1273

Chang CSJ, Li YH, Chen LT, Chen WC, Hsieh WP, Shin J, Jane WN, Chou SJ, Choi G, Hu
JM, Somerville S, Wu SH (2008) LZF1, a HY5-regulated transcriptional factor, functions
in *Arabidopsis* de-etiolation. Plant J **54:** 205–219

Cheng XF, Wang ZY (2005) Overexpression of COL9, a CONSTANS-LIKE gene, delays
flowering by reducing expression of CO and FT in *Arabidopsis thaliana*. Plant J **43:**
758–768

Chu Z, Wang X, Li Y, Yu H, Li J, Lu Y, Li, H., & Ouyang, B. (2016). Genomic organization,
phylogenetic and expression analysis of the B-Box gene family in tomato. Front. Plant
Sci **7:** 1552

Crocco CD, Botto JF (2013) BBX proteins in green plants: insights into their evolution,
structure, feature and functional diversification. Gene **531:** 44–52

Crocco CD, Holm M, Yanovsky MJ, Botto JF (2010) AtBBX21 and COP1 genetically interact
in the regulation of shade avoidance. Plant J **64:** 551–562

Cui L, Zheng F, Wang J, Zhang C, Zhang D, Gao S, Zhang C, Ye J, Zhang Y, Ouyang B,
Wang T, Hong Z, Ye Z, Zhang J (2022) The tomato CONSTANS-LIKE protein SlCOL1
regulates fruit yield by repressing SFT gene expression. BMC Plant Biol **22:** 429

Doi K, Izawa T, Fuse T, Yamanouchi U, Kubo T, Shimatani Z, Yano M, Yoshimura A (2004)
Ehd1, a B-type response regulator in rice, confers short-day promotion of flowering and
controls FT-like gene expression independently of Hd1. Genes Dev. **18:** 926-936

Eguen T, Straub D, Graeff M, Wenkel S (2015). MicroProteins: small size-big impact. Trends
Plant Sci **20:** 477–482

Fan XY, Sun Y, Cao DM, Bai MY, Luo XM, Yang HJ, Wei CQ, Zhu SW, Sun Y, Chong K, Wang ZY (2012) BZS1, a B-box protein, promotes photomorphogenesis downstream of both brassinosteroid and light signaling pathways. Molecular Plant **5:** 591–600

Gangappa SN, Botto JF (2014) The BBX family of plant transcription factors. Trends Plant Sci **19:** 460–470

Gendron JM, Pruneda-Paz JL, Doherty CJ, Gross AM, Kang SE, Kay SA (2012) *Arabidopsis* circadian clock protein, TOC1, is a DNA-binding transcription factor. PNAS **109:** 3167–3172

Gillaspy G, Ben-David H, Gruissem W (1993) Fruits: a developmental perspective. Plant Cell **5:** 1439–1451

Graeff M, Straub D, Eguen T, Dolde U, Rodrigues V, Brandt R, Wenkel S (2016) MicroProtein-mediated recruitment of CONSTANS into a TOPLESS trimeric complex represses flowering in *Arabidopsis*. PLoS Genet **12:** e1005959

Heng Y, Lin F, Jiang Y, Ding M, Yan T, Lan H, Zhou H, Zhao X, Xu D, Deng XW (2019) B-Box containing proteins BBX30 and BBX31, acting downstream of HY5, negatively regulate photomorphogenesis in *Arabidopsis*. Plant Physiol **180:** 497–508

Holtan HE, Bandong S, Marion CM, Adam L, Tiwari S, Shen Y, Maloof JN, Maszle DR, Ohto MS, Preuss S, Meister R, Petracek M, Repetti PP, Reuber TL, Ratcliffe OJ, Khanna R (2011) BBX32, an *Arabidopsis* B-Box Protein, functions in light signaling by suppressing HY5-regulated gene expression and interacting with STH2/BBX21. Plant Physiol **156:** 2109-2123

Huang J, Zhao X, Weng X, Wang L, Xie W (2012) The rice B-Box zinc finger gene family: genomic identification, characterization, expression profiling and diurnal analysis. PLoS ONE **7:** e48242

Karimi M, Inzé D, Depicker A (2002) GATEWAY vectors for *Agrobacterium*-mediated plant transformation. Trends Plant Sci **7:** 193–195

Khanna R, Kronmiller B, Maszle DR, Coupland G, Holm M, Mizuno T, Wu SH (2009) The *Arabidopsis* B-Box zinc finger family. Plant Cell **21:** 3416-3420

Kiełbowicz-Matuk A, Rey P, Rorat T (2014) Interplay between circadian rhythm, time of the day and osmotic stress constraints in the regulation of the expression of a *Solanum* Double B-box gene. Ann Bot **113:** 831–842

Kikuchi R, Kawahigashi H, Oshima M, Ando T, Handa H (2012) The differential expression of HvCO9, a member of the CONSTANS-like gene family, contributes to the control of flowering under short-day conditions in barley. J Exp Bot **63:** 773–784

Kushwaha AK, Dwivedi S, Mukherjee A, Lingwan M, Dar MA, Bhagavatula L, Datta S (2022) Plant microProteins: small but powerful modulators of plant development. iScience **25:** 1-15

Li F, Sun J, Wang D, Bai S, Clarke AK, Holm M (2014) The B-box family gene STO (BBX24) in *Arabidopsis thaliana* regulates flowering time in different pathways. PloS One **9:** e87544

Li H, Wu H, Qi Q, Li H, Li Z, Chen S, Ding Q, Wang Q, Yan Z, Gai Y, Jiang X, Ding J, Gu T, Hou X, Richard M, Zhao Y, Li Y (2019) Gibberellins play a role in regulating tomato fruit ripening. Plant Cell Physiol **60:** 1619–1629

Li W, Wang J, Sun Q, Li W, Yu Y, Zhao M, Meng Z (2017) Expression analysis of genes encoding double B-box zinc finger proteins in maize. Funct Integr Genomics **17:** 653–666

Lifschitz E, Ayre BG, Eshed Y, Coupland G, Mount SM, Sharma R (2014) Florigen and anti-florigen – a systemic mechanism for coordinating growth and termination in flowering plants. Front Plant Sci **5:** 465

Lira BS, Oliveira MJ, Shiose L, Wu RTA, Rosado D, Lupi ACD, Freschi L, Rossi M (2020) Light and ripening-regulated BBX protein-encoding genes in *Solanum lycopersicum.* Sci Rep **10:** 19235

Livak KJ, Schmittgen TD (2001) Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method. Methods **25:** 402–408

Meng X, Muszynski MG, Danilevskaya ON (2011) The FT-Like ZCN8 gene functions as a floral activator and is involved in photoperiod sensitivity in maize. Plant Cell **23:** 942–960

Miller TA, Muslin EH, Dorweiler JE (2008) A maize CONSTANS-like gene, conz1, exhibits distinct diurnal expression patterns in varied photoperiods. Planta **227:** 1377–1388

Molesini B, Dusi V, Pennisi F, Di Sansebastiano GP, Zanzoni S, Manara A, Furini A, Martini F, Rotino GL, Pandolfini T (2020) TCMP-2 affects tomato flowering and interacts with BBX16, a homolog of the *Arabidopsi*s B-box MiP1b. Plant Direct **4:** e00283

Molesini B, Rotino GL, Dusi V, Chignola R, Sala T, Mennella G, Francese G, Pandolfini T (2018) Two metallocarboxypeptidase inhibitors are implicated in tomato fruit development and regulated by the Inner No Outer transcription factor. Plant Sci **266:** 19–26

Park HY, Lee SY, Seok HY, Kim SH, Sung ZR, Moon YH (2011) EMF1 interacts with EIP1, EIP6 or EIP9 involved in the regulation of flowering time in *Arabidopsis*. Plant Cell Physiol **52:** 1376–1388

Pear JR, Ridge N, Rasmussen R, Rose RE, Houck CM (1989) Isolation and characterization of a fruit-specific cDNA and the corresponding genomic clone from tomato. Plant Mol Biol **13:** 639-651

Ping Q, Cheng P, Huang F, Ren L, Cheng H, Guan Z, Fang W, Chen S, Chen F, Jiang J (2019) The heterologous expression in *Arabidopsis thaliana* of a chrysanthemum gene encoding the BBX family transcription factor CmBBX13 delays flowering. Plant Physiol Biochem **144:** 480–487

Rodrigues VL, Dolde U, SunB, Blaakmeer A, Straub D, Eguen T, Botterweg-Paredes E, Hong S, Graeff M, Li MW, Gendron JM, Wenkel S (2021) A microProtein repressor complex in the shoot meristem controls the transition to flowering. Plant Physiol **187:** 187–202

Saito T, Ariizumi T, Okabe Y, AsamizuE, Hiwasa-Tanase K, Fukuda N, Mizoguchi T,
Yamazaki Y, Aoki K, Ezura H (2011) TOMATOMA: a novel tomato   mutant
database distributing Micro-Tom mutant collections. Plant Cell Physiol **52:** 283-296

Singh D, Datta S (2023) BBX30/miP1b and BBX31/miP1a form a positive feedback loop
with ABI5 to regulate ABA-mediated post germination seedling growth arrest. New
Phytol **238:** 1908–1923

Talar U, Kieøbowicz-Matuk A, Czarnecka J, Rorat T (2017) Genome-wide survey of B-box
proteins in potato (*Solanum tuberosum*)—Identification, characterization and
expression patterns during diurnal cycle, etiolation and de-etiolation. PloS One **12:**
e0177471

Treggiari D, Zoccatelli G, Molesini B, Degan M, Rotino GL, Sala T, Cavallini C, Macrae CA,
Minuz P, Pandolfini T (2015) A cystine-knot miniprotein from tomato fruit inhibits
endothelial cell migration and angiogenesis by affecting vascular endothelial growth
factor receptor (VEGFR) activation and nitric oxide production. Mol Nutr Food Res **59:**
2255–2266

Tripathi P, Carvallo M, Hamilton EE, Preuss S, Kay SA (2017) *Arabidopsis* B-BOX32
interacts with CONSTANS-LIKE3 to regulate flowering. PNAS **114:** 172–177

Vaishak KP, Yadukrishnan P, Bakshi S, Kushwaha AK, Ramachandran H, Job N, Babu D,
Datta S (2019) The B-box bridge between light and hormones in plants. J Photochem
191: 164–174

Wang L, Sun J, Ren L, Zhou M, Han X, Ding L, Zhang F, Guan Z, Fang W, Chen S, Chen
F, Jiang J (2019) CmBBX8 accelerates flowering by targeting CmFTL1 directly in
summer chrysanthemum. Plant Biotechnol J **18:**1562–1572

Wang Q, Tu X, Zhang J, Chen X, Rao L (2013) Heat stress-induced BBX18 negatively
regulates the thermotolerance in *Arabidopsis*. Mol Biol Rep **40:** 2679–2688

Wang X, Wang Q, Nguyen P, Lin C (2014) Cryptochrome-mediated light responses in
plants. Enzymes **35:** 167–189

Weng L, Zhao F, Li R, Xiao H, Zhao F (2015) Cross-talk modulation between ABA and
ethylene by transcription factor SlZFP2 during fruit development and ripening in tomato.
Plant Signal Behav **10:** e1107691

Wu Q, Kuang K, Lyu M, Zhao Y, Li Y, Li J, Pan Y, Shi H, Zhong S (2020) Allosteric
deactivation of PIFs and EIN3 by microproteins in light control of plant development.
PNAS **117:** 18858–18868

Xu X, Wang Q, Li W, Hu T, Wang Q, Yin Y, Liu X, He S, Zhang M, Liang Y, Zhu J, Zhan X
(2022) Overexpression of SlBBX17 affects plant growth and enhances heat tolerance in
tomato. Int J Biol Macromol **206**: 799–811

Yadav A, Ravindran N, Singh D, Rahul PV, Datta S (2020) Role of *Arabidopsis* BBX proteins
in light signaling. J Plant Biochem Biotechnol **29:** 623–635

Yang S, Weers BD, Morishige DT, Mullet JE (2014) CONSTANS is a photoperiod regulated
activator of flowering in sorghum. BMC Plant Biol **14: 148**

Zhang X, Henriques R, Lin SS, Niu QW, Chua NH (2006) *Agrobacterium*-mediated
transformation of *Arabidopsis thaliana* using the floral dip method. Nat Protoc **1:** 641–
646

Zouine M, Maza E, Djari A, Lauvernier M, Frasse P, Smouni A, Pirrello J, Bouzayen M
(2017) TomExpress, a unified tomato RNA-Seq platform for visualization of expression
data, clustering and correlation networks. Plant J **92:** 727–735

Figure 1. Overexpression of *SlBBX16* and *SlBBX17* in *Arabidopsis* WT (Col-0) and miP1a/b KO mutant. The transition from vegetative to reproductive development was determined by counting the number of rosette leaves at the bolting stage and the number of days to bolting. **(A, B)** Three lines overexpressing *SlBBX16* (*Sl*BBX16OE #3, #4, and #5) and **(C, D)** two lines overexpressing *SlBBX17* (*Sl*BBX17OE #1, and #4) were compared with WT. Values are mean ± SE (n= 19-27, for panels A and B; n=16-26, for panels C and D). **(E, F)** Two *Sl*BBX16OE lines (#2, and #3) and **(G, H)** two *Sl*BBX17OE lines (#1, and #5) were compared with miP1a/b double KO mutant. Values are mean ± SE (n= 13-26, for panels E and F; n= 18-26 for panels G and H). *P<0.05 and ***P < 0.001 versus the respective control (Student's t-test).

Figure 2. Transition to flowering and fruit set parameters of SlBBX16OE plants. (A) Number of leaves before the first inflorescence **(B)** Shoot height at the first inflorescence **(C)** number of flowers in the first three inflorescences **(D)** Fruit set of the first three inflorescences calculated as the percentage of the number of fruits over the number of flowers. The values reported are means ± SE (n= 10-13). Student's t-test was used for the statistical analysis (*P < 0.05; **P < 0.01).

664 **Figure 3. Fruit growth and production in *Sl*BBX16OE plants. (A)** Number of green fruits
665 recorded from 58 to 81 days after sowing **(B)** Number of fruits in the stage of maturation **(C)** Total
666 fruit number per plant recorded 110 days after sowing **(D)** Average fruit weight. The values reported
667 are means ± SE (n= 10-13). Student's t-test was used for the statistical analysis (*P < 0.05).

Figure 4. Fruit growth and production in *Sl*BBX17OE plants. **(A)** Shoot height at the first inflorescence **(B)** number of leaves before the first inflorescence; **(C)** number of flowers in the first three inflorescences **(D** and **E)** number of flowers at anthesis per plant from 39 to 90 days after sowing. The values reported are means ± SE (n= 5-8). (**F** and **G**) Expression level of *SFT* and *SP* of WT and *Sl*BBX17OE plants. Values are means ± SE of three biological replicates. Student's t-test was used for the statistical analysis (*P < 0.05; **P < 0.01).

Figure 5. Fruit growth and production in *Sl*BBX17OE plants. **(A)** Fruit set percentage of the first three inflorescences calculated as the percentage of number of fruits over number of flowers. **(B)** Total fruit number per plant, **(C)** percentage of ripe over total fruits collected 110 days after sowing **(D)** weight of ripe fruits and **(E)** number of seeds collected from ripe fruits. **(F)** Representative picture of fruit production. The values reported are means ± SE (n 8-14 for panel A, B and C; n=18-34 for panel D and E). Student's t-test was used for the statistical analysis (*P < 0.05; **P < 0.01).

Figure 6. Expression of genes implicated in ethylene, abscisic acid and GA metabolism.
Transcript levels of *ACS1A*, *ACS6*, *ZFP2*, *NCED-1*, *GA2ox4* and *GA20ox2* in WT and *Sl*BBX17OE
(panels **A-F**) and *Sl*BBX17OE (**G-L**) fruits. Values are means ± SE of three biological
replicates. Student's t-test was used for the statistical analysis (*P < 0.05; **P < 0.01).

Figure 7. Speculative model of the microProteins *Sl*BBX16 and *Sl*BBX17 action during fruit development. (A) Trend of GA content and transcript levels of Sl*BBX16*, Sl*BBX17* and *TCMP-2* at different stages of fruit growth and ripening, as deduced from previously published articles. IG, immature green; MG, mature green; Br, breaker. **(B)** GA induces the expression of *SlBBX16* (left) and *SlBBX17* (right), which directly or indirectly (dotted lines/arrow) would cause alterations in genes involved in GA metabolism. These transcriptional changes would result in feedback regulation of active GA concentration. Overexpression of *SlBBX16,* by forming a regulatory complex with TCMP-2 and possibly other unknown factors (?), would act by delaying fruit expansion (Right). Overexpression of *SlBBX17*, through the formation of a similar complex, would negatively control fruit growth and misregulate ripening (Left).

SUPPORTING INFORMATION

Figure S1. *Sl*BBX16 and *Sl*BBX17 overexpressing lines in *Arabidopsis*. (A) Sequence alignment of *Sl*BBX16 and *Sl*BBX17 with the closest homologs in *Arabidopsis*, miP1a and miP1b. The amino acid sequences were aligned using the ClustalW2 (http://www.ebi.ac.uk/Tools/msa/clustalw2) program, choosing default alignment parameters and selecting BLOSUM for the protein matrix. Consensus symbols: '*' identical residues; ':' residues with strongly similar properties; '.' residues with weakly similar properties. **(B, C)** RT-PCR analysis carried out on *Sl*BBX16OE lines and *Sl*BBX17OE lines both in *Arabidopsis* WT and miP1a/b double KO mutant. N= no template control; M= DNA ladder.

Figure S2. Y2H interaction assay of *Sl*BBX16 and *Sl*BBX17 with *At*CO and *Sl*COL1. The empty bait and prey vectors (BD and AD) were used to detect self-activation. For each protein-protein interaction, two increasing dilutions of the mated cultures (10^{-2} and 10^{-3}) were spotted on control medium (SD/-Leu/-Trp) and selection medium plates (SD/-Leu/-Trp/X-Gal/Aureobasidin A). The interaction of *At*miP1a with AtCO (Graeff et al., 2016) and p53 with SV40 were used as positive mating controls.

Figure S3. Overexpression of *TCMP-2* in the cultivar MicroTom. (A) Expression level of *TCMP-2* evaluated by RT-qPCR in tomato leaves of WT and two overexpressing lines (#6 and #20). Values are mean ± SE of three biological replicates; n.d. means no detectable. Line #20 was used as calibrator. **(B)** Number of leaves before the first inflorescence, **(C)** total plant height. Values are mean ± SE (n=9-10 for panels B and C). **(D)** Phenotypical aspect of WT and TCMP-2OE #20 plants at the same developmental stage.

Figure S4. Tomato *SlBBX16* overexpressing lines and expression of *SlBBX16* in different organs. (A) *SlBBX16* expression in three transgenic lines (#3, #4, and #20). WT was used as calibrator. **(B)** Expression level of *SlBBX16* in ovary detached from flowers at anthesis, young fruits (0.5-1.0 cm diameter) and leaves. Ovary was used as calibrator. Values are mean ± SE of three biological replicates.

Figure S5. (A) Expression level of *SlBBX17* in ovary detached from flowers at anthesis, young fruits (0.5-1.0 cm diameter) and leaves. Ovary was used as calibrator. Values reported are means ± SE of three biological replicates. **(B)** Y2H interaction assay between TCMP-2 and *Sl*BBX17. For negative controls, pGBKT7 without insert (BD alone; Empty), pGADT7 without insert (AD alone; Empty) were used. For each protein-protein interaction, two increasing dilutions of the mated cultures (10^{-2} and 10^{-3}) were spotted on control medium (SD/-Leu/-Trp) and selection medium plates (SD/-Leu/-Trp/-His/-Ade/X-Gal/Aureobasidin A/10 mM 3-AT). The interaction of p53 with SV40 was used as a positive mating control. **(C)** Expression of *SlBBX17* in tomato transgenic lines (#15 and #18). WT was used as calibrator. Values reported are means ± SE of three biological replicates. **(D)** Heat stress response of one-month-old WT and line #18 plants. Damage index was calculated as the percentage of dead or severely damaged plants after imposing heat stress at 42°C for 24 h. Values are mean ± SE of three independent experiments. *P < 0.05, ***P < 0.001 versus the respective control (Student's t-test).

Figure S6. Green fruit weight of SIBBX17OE. Comparison between distribution of green fruits' weights in WT and *Sl*BBX17OE lines. Dashed lines represent median and dotted lines quartiles.

Figure S7. Response of *SlBBX16* and *SlBBX17* genes to GA. Relative expression of *SlBBX16* **(A)** and *SlBBX17* **(B)** in shoots of seedling treated for 24 h with 5 μM GA$_3$ in comparison with Mock-treated ones. Values are mean ± SE of three biological replicates. Student's t-test was used for the statistical analysis (*$P < 0.05$; **$P < 0.01$).

Name	Experiment	Forward sequence (5'->3')	Reverse sequence (5'->3')
SlBBX16 CDS	Cloning	GCAGGTACCATGTGCAATGGAAGAAGAGAAA	TGCGGATCCTTAGAGAAACAAAAAAGGTTCTTT
SlBBX17 CDS	Cloning	GCAGGTACCATGTGTAGTGGAAGAAGAGAG	TTAGGATCCTCAAATAAACAGAAAAAGCTCTT
TCMP-2 CDS	Cloning	GGGGACAAGTTTGTACAAAAAAGCAGGCTCCAACATGGCAGCAAAGAACAGTGAA	GGGGACCACTTTGTACAAGAAAGCTGGGTCCTATTATGGAAGTAAATTACAAGTTCT
SlBBX16 (Solyc12g005750)	qRT-PCR	AGGTATTTGATTGGAGTTTCACA	CTTTTGTTGAACAATTTTGCTCA
SlBBX17 (Solyc07g052620)	qRT-PCR	AATGTGATAGATCGGTTCACGGA	GTTCCAAACGGGCAATCGTCG
SlTCMP-2 (Solyc07g049140)	qRT-PCR	ACGAACCTTGCAGCTCAAAC	GCAACAGGTTGCATGTACGG
SlACTIN (Solyc11g005330)	qRT-PCR	TTCAAAGGGCGAGTACGACGAG	CAGCAGACCCGAGTTCACTTTT
SlSAND (Solyc03g115810.2)	qRT-PCR	TTTGCTTGGAGGAACAGACG	GCAAACAGAACCCCTGAATC
AtActin (At3g18780)	qRT-PCR	TGTTCTCTCCTTGTACGCCAGT	CAGCAAGGTCAAGACGGAGGA
SlSFT (Solyc03g063100)	qRT-PCR	GACAATTAGGTCGGCAAACA	AGCAGCAACAGGTAAACCAA
SlSP (Solyc06g074350)	qRT-PCR	TGAACTTGGCTCACCAGTTGCTGC	AGTAGTAGTCCGTCCACGCACTAA
SlCOL1 (Solyc02g089540)	qRT-PCR	AAGATCAACCACAAGCCCCTAA	TCCTCTGTGGAAAGCTGCTGTAG
SlGA20ox2 (Solyc06g035530.2)	qRT-PCR	CATCCTCCATCAAGACAACGTC	TTACTACAGCTCTGTGCAAACAACT
SlGA2ox4 (Solyc07g061720.2)	qRT-PCR	AGAAAATCGGACAAAATGGCGA	ACTCAAAACATTCGTCGGTTGA
SlACS1A (Solyc08g081550.2)	qRT-PCR	ACCAGCTCACTCTCCGATGA	TCACTCATCACTACAACAAAAGGAA
SlACS6 (Solyc08g008100.2)	qRT-PCR	ATTGCTCAGAGCCAGGTTGG	TGTTGCAACTTCAACTCCCTT
SlNCED-1 (Solyc07g056570.1.1)	qRT-PCR	AGGCAACTGTGAAGCTTCCATCA	TCCATTAAAGAGGAATATTCCGGG
SlZFP2 (Solyc07g006880.1)	qRT-PCR	TGAGCCCTAGTGGTCCGAGC	TGTCCAAAATCCTCTTGAACACTCT
BD-*SlBBX16* (Solyc12g005750)	Y2H	CATGGAGGCCGAATTCTGCAATGGAAGAAGAGAAATTGA	GCAGGTCGACGGATCCGAGAAACAAAAAAGGTTCTTTCACT
BD-*SlBBX17* (Solyc07g052620)	Y2H	CATGGAGGCCGAATTCTGTAGTGGAAGAAGAGAGGGTGAC	GCAGGTCGACGGATCCAATAAACAGAAAAAGCTCTTGCGGC

AD-*SlBBX17* (Solyc07g052620)	Y2H	GGAGGCCAGTGAATTCTGTAGTGGAAGAAGAGAGGGTGAC	CGAGCTCGATGGATCCAATAAACAGAAAAAGCTCTTGCGGC
BD-*SlTCMP-2* (Solyc07g049140)	Y2H	CATGGAGGCCGAATTCACAAATATTTTGGGACTTTGTAACG	GCAGGTCGACGGATCCAGGCAACAGGTTGCATGTACG
BD-*Atmip1A* (At3g21890)	Y2H	CATGGAGGCCGAATTCTGTAGAGGCTTGAATAATGAAGAG	GCAGGTCGACGGATCCGAGAAAAACAAACGGAACCTCATG
AD-*SlCOL1* (Solyc02g089540)	Y2H	GGAGGCCAGTGAATTCTTGAAAAAAGAGAACAGTAACAATTGGG	CGAGCTCGATGGATCCGAATGAAGGGACAATTCCATAATTGC
AD-*AtCO1* (At5g15840)	Y2H	GGAGGCCAGTGAATTCTTGAAACAAGAGAGTAACGACA	CGAGCTCGATGGATCCGAATGAAGGAACAATCCCATATCCT

841 **Table S1. List of primers**