A spatially constrained independent component analysis jointly informed by structural and functional network connectivity.

Mahshid Fouladivanda[1,2*], Armin Iraji[1,2], Lei Wu[1], Theodorus G.M. van Erp[3], Aysenil Belger[4], Faris Hawamdeh[6], Godfrey D. Pearlson[5], Vince D. Calhoun[1,2*]

[1] Tri-institute Translational Research in Neuroimaging and Data Science (TReNDS Center), Georgia State University, Georgia Institute of Technology, Emory University, Atlanta, GA, USA.
[2] Georgia State University, Atlanta, GA, USA.
[3] Clinical Translational Neuroscience Laboratory, Department of Psychiatry and Human Behavior School of Medicine, University of California, Irvine, CA, USA.
[4] Department of Psychiatry Director, Neuroimaging Research in Psychiatry Director, Clinical Translational Core, UNC Intellectual and Developmental Disabilities Research Center, University of North Carolina, Chapel Hill, NC, USA.
[5] Olin Neuropsychiatry Research Center, Department of Psychiatry and Neuroscience, Yale University, School of Medicine, New Haven, CT, USA.
[6] Center for Disaster Informatics and Computational Epidemiology (DICE), Georgia State University, Atlanta, GA, USA.

Abstract

There are a growing number of neuroimaging studies motivating joint structural and functional brain connectivity. Brain connectivity of different modalities provides insight into brain functional organization by leveraging complementary information, especially for brain disorders such as schizophrenia. In this paper, we propose a multi-modal independent component analysis (ICA) model that utilizes information from both structural and functional brain connectivity guided by spatial maps to estimate intrinsic connectivity networks (ICNs). Structural connectivity is estimated through whole-brain tractography on diffusion-weighted MRI (dMRI), while functional connectivity is derived from resting-state functional MRI (rs-fMRI). The proposed structural-functional connectivity and spatially constrained ICA (sfCICA) model estimates ICNs at the subject level using a multi-objective optimization framework. We evaluated our model using synthetic and real datasets (including dMRI and rs-fMRI from 149 schizophrenia patients and 162 controls). Multi-modal ICNs revealed enhanced functional coupling between ICNs with higher

structural connectivity, improved modularity, and network distinction, particularly in

schizophrenia. Statistical analysis of group differences showed more significant differences in the

proposed model compared to the unimodal model. In summary, the sfCICA model showed benefits

from being jointly informed by structural and functional connectivity. These findings suggest

advantages in simultaneously learning effectively and enhancing connectivity estimates using

structural connectivity.

Keywords: multi-modal independent components analysis, multi-objective model, diffusion MRI,

resting state fMRI, spatial constraint, schizophrenia

Corresponding authors: Mahshid Fouladivanda (mfouladivanda@gsu.edu),
47   Vince. D. Calhoun (vcalhoun@gsu.edu)

1. Introduction

The human brain can be modeled as a distinct functional unit that exhibits a temporally synchronized pattern known as intrinsic connectivity networks (ICNs) (Genon et al., 2018; Iraji et al., 2020). These ICNs interact with each other, collectively representing brain function. Consequently, accurate estimation of ICNs is a critical step in studying brain networks because it minimizes the impact of inaccuracies in functional activities (Iraji et al., 2020). One approach for identifying ICNs and gaining insights into brain networks, including brain disorders like schizophrenia (Iraji et al., 2019; Iraji, Faghiri, Fu, Kochunov, et al., 2022; Iraji, Faghiri, Fu, Rachakonda, et al., 2022), is data-driven independent component analysis (ICA) (Calhoun & Adalı, 2012; Calhoun & de Lacy, 2017). This method assumes spatial independence for the temporally synchronized functional units and recovers a set of maximally independent sources from a mixture of unknown source signals without any prior information (Calhoun et al., 2001), using single modality neuroimaging, particularly resting-state functional magnetic resonance imaging (rs-fMRI). Rs-fMRI images dynamically measure the hemodynamic response associated with neural activity in the brain during rest.

ICA has become widely used to partition the brain into spatially overlapping and distinct ICNs at the group level. Group ICNs are estimated using fMRI images from all subjects, then the corresponding subject-level ICNs are obtained. The most common approach is to apply ICA to the data, then use back reconstruction to estimate subject-specific maps (Allen et al., 2011; Calhoun et al., 2001; Erhardt et al., 2011). These approaches are fully data-driven; however, more recently, spatially constrained ICA approaches (Du & Fan, 2013; Lin et al., 2010) have been proposed to estimate ICNs using a set of templates for ICNs as priors. They can be used to build a fully

automated approach (e.g., the NeuroMark pipeline (Du et al., 2020)) that does not require post hoc

ICN selection or network matching (Du et al., 2020), and also automatically provides ICN

ordering, providing modular functional network connectivity. These approaches have been used

in many prior studies and offer a fully automated framework that can be integrated within a larger

framework (e.g., a containerized version is available from http://trendscenter.org/software/gift, as

well as a BIDSapp (Kim et al., 2023)), making them more easily comparable across studies.

Current constrained ICA approaches are unimodal and do not leverage information from other

modalities, particularly dMRI images. Although MRI images, such as T1 and T2, include structural

information about the brain, dMRI images provide information about the physical paths in the

human brain, captured as structural connectivity which drives functional activities of different

units. In this study, we investigate a model that estimates functional brain patterns (ICNs) informed

by multi-modal data to provide a more complete view of the connectome by employing both

structural and functional connectivity. This has many advantages, for example, a connectivity

domain ICA approach (Iraji et al., 2016) called joint connectivity matrix ICA has been used to

jointly parcellate structural and functional connectivity data, yielding a data-driven

structure/function parcellation (Wu & Calhoun, 2023).

One longstanding aspiration in brain research is to link brain function to its underlying architecture,

making the interplay between structural connectivity and functional network connectivity (FNC)

a fundamental in network neuroscience (Park & Friston, 2013). There have been studies, such as

(Calhoun & Sui, 2016) using computational models on the connectivity domain to estimate the

human brain functional activities propounding from different modalities, mainly single modality

information. Consequently, a fundamental question arises regarding how to model human brain

functional activity changes with a deeper understanding of structural and functional information

(Batista-García-Ramó & Fernández-Verdecia, 2018; Skudlarski et al., 2010). Several studies have

been proposed to fill this gap ((Batista-García-Ramó & Fernández-Verdecia, 2018; Suárez et al.,

2020)) by developing different models predicting FNC from structural connectivity using

statistical models (Messé et al., 2014; Mišić et al., 2016; Suárez et al., 2020), or communication

models based on structural-functional network connectivity (Goñi et al., 2014; Zamani Esfahlani

et al., 2022).

In addition, jointly analyzing models is a growing methodology used to investigate the human

brain and how structural and functional relate to each other (Calhoun & Sui, 2016; Puxeddu et al.,

2022; Zhu et al., 2021). One may provide variations in brain functions at different regions by

multilayer network models (Puxeddu et al., 2020; Shine et al., 2016). Multilayer network models

have emerged to analyze the human brain employing the concepts of graph theory with multiple

viewings of the human brain (Puxeddu et al., 2020; Puxeddu et al., 2022). In parallel, other studies

have investigated brain network modules, uncovering intrinsic brain network activations using

data-driven models (Calhoun & Adalı, 2012; Yeo et al., 2011), and community detection (Power

et al., 2011). Several studies have attempted to develop multi-modal fusion models (Calhoun &

Sui, 2016; Sui et al., 2014) incorporating complementary information across modalities into the

analysis of ICNs (Batista-García-Ramó & Fernández-Verdecia, 2018; Suárez et al., 2020). Multi-

modal fusion can provide additional insights into brain structure and function that are impacted by

psychopathology, including identifying which structural or functional aspects of pathology might

be linked to human behavior or cognition (Sui et al., 2014), particularly in disorders related to both

brain function and structures such as schizophrenia (Calhoun, 2018).

Moreover, information from different modalities can be jointly analyzed to investigate ICNs using

ICA. This has been done mostly by focusing on linking group-level spatial features, including joint

ICA (jICA) (Calhoun et al., 2006), parallel ICA (pICA) (Liu et al., 2009), parallel ICA with

reference (pICAR) (Chen et al., 2013), multi-modal canonical correlation analysis (mCCA)+jICA

(Sui et al., 2011), multisite canonical correlation analysis with reference (mCCAR)+jICA (Qi et

al., 2018; Shile et al., 2016), and linked ICA (Groves et al., 2011), or jointly optimizing for fMRI

networks and covarying networks from structural MRI as in parallel group ICA+ICA.

However, these approaches have not directly integrated structural and functional network

connectivity information (Qi et al., 2022; S. L. Qi et al., 2019). To our knowledge, fewer prior

works have used structural connectivity to estimate ICNs, as in (Wu & Calhoun, 2023), which

performed joint structure/functional parcellation. Specifically, there is a lack of methodologies to

guide adaptive ICN estimation by incorporating both structural and functional network

connectivity. A common practice in neuroscience is to compress Network connectivity

information into nodes and edges to understand functional interactions in the brain

(Babaeeghazvini et al., 2021). Thus, using network connectivity helps to drive the model to learn

from both structural connectivity (real connections among different functional networks) and FNC

(their functional interactions).

In this paper, we introduce a novel adaptive ICA model designed to directly investigate the ICNs.

This is achieved by imposing constraints using two different modalities at the subject level. Our

approach involves constructing a new multi-modal ICA model that jointly embeds structural and

functional network connectivity. This model is supported by previous studies that indicate that

structural connectivity forms the foundation of functional connectivity (Honey et al., 2009;

Litwińczuk et al., 2022; Stam et al., 2016; Zhu et al., 2021). In addition, we incorporated prior

spatial information into our multi-modal ICA model to increase the generalizability of the

estimated ICNs. Our model is an iterative pipeline that simultaneously learns from both dMRI and

fMRI, along with ICN spatial maps, to identify joint structural-functional independent

components.

While prior work (Wu & Calhoun, 2023) applied a connectivity-based model to perform joint data-

driven parcellation of dMRI and fMRI, no approaches have attempted to directly link the full

spatiotemporal fMRI data with dMRI data in the context of a connectivity constraint at the subject

level within a spatially constrained model, thus providing a fully automated approach. Our

proposed structural-functional network connectivity and spatially constrained ICA model

(sfCICA) identifies ICNs that are simultaneously optimized to be maximally spatially independent,

influenced by both structural and functional connectivity. Thus, the learning procedure to estimate

subject-level ICNs is informed by the structural and functional network connectivity information

of each subject. We first validated the approach employing synthetic data and then applied the

proposed model to a real dataset, including controls and patients with schizophrenia. The results

demonstrate that the proposed method can effectively impose prior information on ICNs while

jointly learning from structural-functional network connectivity in both synthetic and real human

brain data. Furthermore, the identified ICNs exhibit an increased ability to distinguish between

healthy controls and individuals with schizophrenia.

2. Material and Methods

2.1. Structural-functional network connectivity and spatially constrained ICA (sfCICA).

Model definition.

A multi-modal ICA (sfCICA) model, called *structural and functional network connectivity and spatially constrained ICA* was introduced. Briefly, the sfCICA framework estimates subject-level ICNs by imposing structural network connectivity weights on time course distances (functional network connectivity). The main advantage of the constraints in the sfCICA framework is driving the model to jointly learn from both structural and functional network connectivity at the subject level. Using the sfCICA model, subject-specific ICNs were estimated through multi-modal information, incorporating prior spatial maps for each ICN derived from a standard template.

In summary, we introduced a multi-objective function as described in **Eq. 1**, consisting of three terms. The first and second terms find independent ICNs which are learned from the spatial template. The third term constrains the learning procedure to account for both structural and functional network connectivity.

$$\max \quad (J(S_l^k), F(S_l^k), -C(w_l^k)) \tag{1}$$

$$\text{s. t.} \ \left\| w_l^k \right\|^{\square} = 1$$

The sfCICA model was proposed in a multi-objective framework, aiming to estimate ICNs by optimizing their independence, through maximizing their non-Gaussianity measured by negentropy or kurtosis ($J(S_l^k)$) (Hyvarinen, 1999) (**Eq. 2**).

$$\max \quad J(S_l^k) = (E[G(S_l^k)] - E[G(v)])^2 \tag{2}$$

$$\text{s. t.} \ \left\| w_l^k \right\|^{\square} = 1$$

Where, G is any quadratic function, E is the expectation maximization operator, and v is a random

variable. S_l^k and w_l^k are respectively the spatial map and time course of the l^{th} ICN (here we use

Neuromark_fMRI_1.0 including 53 ICNs) for k^{th} subject. These are estimated by maximizing the

square differences between the expectation maximization of two vectors, as defined in **Eq. 2**.

In addition, the similarity of the spatial maps with the prior spatial information was maximized by

another cost function $(F(S_l^k))$ in **Eq. 3**. Thus, independence and similarity to the prior spatial

information (here a standard template was used) were jointly maximized during optimization.

$$\max \quad F(S_l^k) = E[S_l S_l^k] \tag{3}$$

$$\text{s. t.} \ \left\| w_l^k \right\|^{\square} = 1$$

where the S_l denoted the spatial map of the l^{th} components in the template.

Furthermore, we integrate structural connectivity information, measured as the number of

streamlines connecting node pairs, into the model as a constraint on the FNC of the estimated time

course. To this end, the model is guided by minimizing the weighted distances of the time courses

based on structural connectivity. The structural connectivity weights captured from dMRI can

reflect existing physical connectivity between functionally related gray matter regions, this means,

more fiber bundles between two brain regions (here are ICNs) may result in stronger functional

correlations and shorter distances between time courses (Zhu et al., 2021).

Thus, we assumed regions with higher structural connectivity (more fiber tracts) would exhibit

closer functional signal activities because of their shorter distance. Prior works (He & Niyogi,

2003; Zhu et al., 2021), led us to reconstruct an objective function $(\min \sum_{jl}^{\square} M_{lj} \| FC_i^{\square} - FC_j^{\square} \|^2)$

to adjust the functional activity of an ICN by considering other structurally connected ICNs. Here,

M_{lj} represents structural connectivity weights and FC denotes the correlation between pairs of

197 ICNs (l, j). It is important to note that maximizing the FC between ICNs involves minimizing the

198 distance of their time courses. Therefore, we propose the following cost function (**Eq. 4**), in

199 addition to the previous cost functions, to be simultaneously optimized. This is aimed at

200 incorporating information from both structure and function as constraints.

$$\min \quad L\left(w_i^{\square}\right) = \sum_{j=1}^{L} M_{lj}\left\|w_l^k - w_j^k\right\|^2 \tag{4}$$

$$\text{s. t. } \left\|w_l^k\right\|^{\square} = 1$$

202 To solve a multi-objective optimization problem, one usually needs to find the Pareto optimal set

203 or its subset and critically evaluate which specific trade-off is more appropriate to the problem

204 under study. In addition, optimizing a linear weighted sum of cost functions is a commonly used

205 method for solving multi-objective optimization problems, which simplifies the problem

206 (Klamroth & Jørgen, 2007). In our study, we utilize a linear weighted approach to address the

207 multi-objective problem described in **Eq. 1.** This allows us to explore various points along the

208 Pareto front by adjusting the weighting values in the linear weighted sum objective function.

$$C = \alpha J\left(S_l^k\right) + \beta F\left(S_l^k\right) - \gamma L\left(w_l^k\right) \tag{5}$$

$$\text{s. t. } \left\|w_l^k\right\|^{\square} = 1$$

210 Where, α, β , and γ are constant weight values. To ensure a fair comparison and prevent the

211 optimization from being biased towards a cost function with a larger magnitude, we normalize the

212 cost functions in **Eq. 2-4** as described in (Du & Fan, 2013), and weights (α, β, and γ) are chosen

213 to be equal (0.33). Then a gradient ascent method was used to iteratively converge on an optimal

214 solution. The iterative algorithm for optimizing the cost (C) is derived as follows. Initially, $S_l^k =$

215 $\left(w_l^k\right)^T * X^k, S_l^k$, is substituted into **Eq. 6**.

$$C(w_l^k) = \alpha J(w_l^k) + \beta F(w_l^k) - \gamma L(w_l^k) \qquad (6)$$

217 Then, according to the derivatives of each term, we have:

$$\nabla C(w_l^k) = \alpha \nabla J(w_l^k) + \beta \nabla F(w_l^k) - \gamma \nabla L(w_l^k) \qquad (7)$$

219 More details regarding the derivative of each term are described in (Du & Fan, 2013). Once the

220 gradient of the objective function of **Eq. 7** is available, we employ the steepest ascent iterative

221 formula:

$$[w_l^k]^{t+1} = [w_l^k]^t + \mu^t . d^t \qquad (8)$$

Where $[w_l^k]^{t+1}$ represents the value of w_l^k after t+1 iteration, d^t is the normalized $\nabla C(w_l^k)$ and,

μ^t stands for the step-length. To ensure a sufficient increase in the objective function in an inexact

line search, μ^t was estimated using the Armijo condition (Jorge Nocedal, 2006).

2.2. Evaluation analysis.

To evaluate the benefit of leveraging multi-modal information, we compare our model with one

of the spatial constrained ICA (CICA) models, known as a multi-objective optimization ICA

model (Du et al., 2020), which is a simplified version of the proposed framework when the multi-

modal information is not utilized. To this end, first, we used synthetic data to confirm that the

model is working in the intended objective defined for the optimization of the time series. Then,

we cope to evaluate our hypothesis using real data, capturing better group differences by

leveraging multi-modal information.

2.2.1. Synthetic data

Simulation. The synthetic dMRI was from FiberCup phantom data (Fillard et al., 2011; Poupon

et al., 2008) (https://tractometer.org/fibercup/data/), including 3mm isotropic images in 64

uniformly distributed over a sphere with b-value = 2000 m/s. It also provided 16 brain regions and

fiber pathways among the regions to reconstruct structural connectivity, (shown in **Figure 1**). To

keep consistency between the simulated dMRI and fMRI data, we used the same sixteen predefined

FiberCup regions as a reference to design synthetic fMRI images, similar to (Chu et al., 2018). We

simulated synthetic fMRI images using the simTB toolbox (Erhardt et al., 2012),

(https://trendscenter.org/software/simtb/).

(a)   (b)

Figure 1. In panel (a) and (b), we present simulated fiber pathways and predefined regions from FiberCup, respectively,

for the reconstruction of synthetic structural connectivity.

The simTB toolbox employs a data generation model that assumes spatiotemporal separability,

allowing the simulated fMRI data to be represented as the multiplication of spatial maps and time

courses. The generated data exhibit realistic dimensions, spatiotemporal activations, and noise

characteristics similar to those observed in typical fMRI datasets (Erhardt et al., 2012). Our fMRI

simulation consists of 16 ICNs for each of the 30 subjects, evenly divided into 3 distinct groups

(10 subjects per group) with varying design parameters (Group A, B and C). Generally, all spatial

maps have $V = 64 \times 64$ voxels (where V is the total number of voxels) and time courses are $T =$

3000 time points in length with a repetition time (TR) of 2 seconds. Spatial maps of the ICNs

represented in **Figure 2** correspond to the regions in the FiberCup data shown in **Figure 1**.

Groups A, B, and C differ in four ways, as outlined below. Firstly, the noise level for each group

was specifically selected to determine the robustness of the proposed model. For Group A, the

mean was 1.7, with a standard deviation of 0.2; for Group B, the mean was 1.5 with a standard

deviation of 0.4; and for Group C, the mean was 1.1 with a standard deviation of 0.35. Secondly,

to introduce diversity among the ICNs across different Groups, we implemented additional

adjustments to both the amplitudes (g_{ic}), and shapes of the reference ICNs. For ICN 6 in Group A,

we selected a weaker amplitude compared to Group B and C, while Group C had a weaker

amplitude for ICNs 8 and 11 compared to Groups A and B. The amplitudes of the ICNs (g_{ic}) were

modeled on a normal distribution. For instance, the distribution of ICN 6 (g_{ic6}) is normal with a

mean of 2.5 and a standard deviation of 0.3 for Group A, and a mean of 3.5 with a standard

deviation of 0.3 for Group B. Similarly, ICN 8 and 11 in Group C follow a normal distribution

with a mean of 2.5 and a standard deviation of 0.3, whereas Groups A and B have a mean of 3.5

and a standard deviation of 0.3. Additionally, each ICN underwent distinct transitions in the x and

y directions, as well as rotations, introducing unique changes. Moreover, the ICNs in Groups A

and C were expanded by factors of 1.2 and 1.3, respectively, while the ICNs in Group B were

contracted by a factor of 0.9.

Then, we constructed a synthetic ICN template to be used as a prior spatial map in the constrained

ICA model. The synthetic template was determined by averaging each ICN over all groups and

introducing deformation and noise.

(a)   (b)

Figure 2. (a) It represents the 16 spatial maps of the ICNs, derived from the FiberCup regions, inserted into the

simTB toolbox as a reference to generate synthetic fMRI images. Using the generated synthetic data, we determined

an average template, shown in the first column from the left (b), to be used as a prior spatial map. The second and

third columns from the left in (b) illustrate examples of estimated ICNs by the sfCICA and CICA model using

synthetic fMRI.

295   **Evaluation criteria.** Our model works on decreasing the weighted distance between time courses

296   of the structurally connected ICNs. To ensure the objective is preserved, we used a distance metric

297   and sparsity to measure the effect of the structural connectivity constraint on TCs. For this purpose,

298   the sum of the distances between an ICN with all other ICs was determined utilizing the Euclidean

299   distance as outlined in **Eq 9.**

$$\text{Dis} = \sum_{j=1}^{L} \|w_l - w_j\|^2 \tag{9}$$

where w_l is the time course of ICN number l and w_j is the time courses of all other remaining ICNs.

This measure was estimated for both the sfCICA and CICA models.

Sparsity was measured as the number of weak connections to all possible connections in the FNC.

To identify weak connections, we first determined an optimal threshold by maximizing the global

cost-efficiency of the brain networks (Bassett et al., 2009), (here a threshold of 0.45 was

determined). The remaining connections after this threshold were considered strong connections

(ST). Then, all FNC networks were thresholded to retain strong connections and sparsity as

follows.

$$SP = \frac{N - ST}{N} \tag{10}$$

Where ST is the number of strong connections and N is the total number of possible connections.

Moreover, the robustness of the model was assessed by varying noise levels across the three

distinct simulated groups.

2.2.2.   Real data

In this study, the real rs-fMRI and dMRI images were used from the Function Biomedical

Informatics Research Network (FBIRN) phase III datasets (Keator et al., 2016). Data collection

was at multiple sites of healthy control (HC) subjects or with schizophrenia (SZ) disorder. FBIRN

dataset consists of rs-fMRI and dMRI images from 311 age-gender-matched adults (age ranging

from 18 to 65 years old). It is 162 healthy controls (HC) including 115 male and 45 female (Avg

$\pm$ SD age: 37.0 ± 10.9) and 149 schizophrenia (SZ) patients including 115 male and 36 female

(Avg $\pm$ SD age: 38.7 ± 11.6).

Acquisition. Images were scanned at six 3T Siemens Tim Trio System and one 3T GE Discovery

MR750 scanner at multiple sites. The acquisition parameters of the rs-fMRI imaging were as

follows: FOV of 220×220 mm (64×64 matrix), TR = 2 s, TE = 30 ms, flip angle (FA) = 77°,

162 volumes, 32 sequential ascending axial slices of 4 mm thickness and 1 mm skip. Subjects had

their eyes closed during the resting state scan. For more details see (Keator et al., 2016). For the

dMRI images, scanning protocols were settled virtually to ensure equivalent acquisitions with 30

directions of diffusion gradient at b = 800 s/mm^2 and five measurements of b = 0 (b0) s/mm^2

(Keator et al., 2016).

rs-fMRI preprocessing. The rs-fMRI images were preprocessed via the statistical parametric

mapping (SPM12, http://www.fil.ion.ucl.ac.uk/spm/) toolbox in MATLAB 2020b, following the

procedure outlined in (Iraji, Fu, et al., 2022). The main steps of the preprocessing included

removing the first five fMRI timepoints, motion correction, slice timing correction, image

registering to the standard Montreal Neurological Institute (MNI) space, spatial resampling (to 3

$\times$ 3 $\times$3 mm^3 isotropic voxels), spatial smoothing using Gaussian kernel with a full width at half

maximum (FWHM) = 6 mm.

dMRI preprocessing. For preprocessing of the dMRI images, first, all images were corrected for

eddy current and motion distortions using the eddy package (FSL 6.0 (Jenkinson et al., 2012)) by

motion induced signal dropout detection based on b0 volumes and replacement method

(Andersson & Sotiropoulos, 2016). Then, a data quality check was performed by an in-house

developed algorithm and visual inspection to exclude images with extreme head motion, signal

dropout, or noise level (Caprihan et al., 2011; Wu et al., 2015).

An additional step was carried out on preprocessed dMRI images to determine white matter fiber

tracts for constructing structural connectivity. First, we used linear regression under the dtifit in

the FSL toolbox (Jenkinson et al., 2012) to model a voxel-wise diffusion tensor. The estimated

diffusion tensors were employed in deterministic tractography using the CAMINO toolbox (Cook

et al., 2006). This procedure involved three stopping criteria: a fixed step size of 0.5mm, an

anisotropy threshold of 0.2, and an angular threshold of 60° (Cook et al., 2006). All fiber tracts

were extracted in native space. Next, we transformed the native fractional anisotropy (FA) maps

into the standard MNI space using advanced normalizations tools (ANTs) (Avants et al., 2009).

Then we applied inverted spatial normalization to the Neuromark_fMRI_1.0 atlas (including N =

53 ICNs) (Du et al., 2020) to obtain the corresponding atlas in native space.

The ICNs of the normalized Neuromark_fMRI_1.0 atlas, which divides the brain into distinct

regions, were used as nodes in the network to build the structural connectivity network. For each

subject, we reconstructed an *N×N* structural connectivity network (M), counting the number of

streamlines connecting pairs of nodes.

Evaluation criteria. First, to determine the effect of the defined objective function in our

multimodal model, we used the same criteria adopted for the synthetic data. Consequently, for the

ICNs of each model (sfCICA and CICA), the distance was estimated using **Eq. 9**. Then, we

computed the sparsity of the estimated FNCs as in **Eq. 10**.

To enhance distance representation, we determined the difference of the measured distances, as

shown in **Eq. 11.**

$$Diff = \text{Dis}_{sfCICA} - \text{Dis}_{CICA} \tag{11}$$

In addition, we evaluated the proposed model with real data, using a set of statistical analysis. A

Paired t-test analysis was employed to determine if the examined ICNs from the sfCICA model,

constrained by structural connectivity, differ significantly from those of the CICA model. The

paired t-test analysis assesses whether the mean difference between paired observations is

significantly different from zero. Here, we used the time course of the ICNs and their FNC matrices

as observations for the paired t-test. FNC matrices were assessed using Pearson's correlation

among the time courses of the ICNs at the subject level for both the sfCICA and CICA models.

Moreover, for group comparison between subjects with schizophrenia (SZ) and healthy control

(HC), we applied a Generalized Linear Regression model (GLM) utilizing FNC matrices and

spatial maps. GLM analysis is a statistical technique that provides a flexible framework for

modeling the relationship between a dependent variable and one or more independent variables.

In this study, we used GLM to regress out effects of the age, sex, diagnosis, motion, and imaging

site effect as covariates for the FNC matrices and spatial maps. Additionally, false discovery rate

(FDR) correction was performed to adjust the estimated p-values.

Finally, to show that structural connectivity helps tune the FNCs we used network graph

parameters. To this end, graph theory analysis, utilizing FNC features was employed. We

estimated the FNCs as Pearson's correlation between TC related to each network (ICN) to

construct the FNC network at the individual level for both the CICA and sfCICA. Then, we

determined four common brain network parameters including modularity (Newman, 2006), local

and global efficiency (Latora & Marchiori, 2001), and small worldness (Amaral et al., 2000) to

characterize the graph properties of the resulting FNC. In addition, randomness (Vergara et al.,

2018), and sparsity were also examined.

In a brain network, modularity indicates the presence of functional modules involved in specific

cognitive processes (Newman, 2006). Local efficiency represents the efficiency of the

communications between regions (here ICNs) and global efficiency is the capacity of the network

for transferring information across the whole brain (Latora & Marchiori, 2001). Additionally,

small-worldness is the ratio of clustering to shortest path length, which summarizes the balance

between the efficiency and clustering in the brain network (Amaral et al., 2000).

3. Results

To determine multimodal ICNs through an iterative optimization, we applied the proposed sfCICA

model to the diffusion MRI (dMRI) and the resting-state fMRI (rs-fMRI) data obtained from 30

synthetic samples and 311 real subjects (146 HC and 162 SZ). The number of the ICNs was

determined based on a predefined standard ICN template, used as prior spatial maps. In this study,

the NeuroMark_1.0 template was utilized as prior maps for the real data. This template consists of

seven distinct functional domains, including subcortical (SC), sensorimotor (SM), auditory

(AUD), visual (VS), cognitive control (CC), default mode (DM), and cerebellum (CB). Our results

represent multimodal ICNs that are more informative and enhanced by incorporating multi-modal

information. They provide supporting evidence indicating the impact of the structural connectivity

information on neural activities (Litwińczuk et al., 2022).

3.1. Estimated ICNs using synthetic datasets show a structural and functional learning

pattern.

To determine the applicability and the robustness of the proposed model, we investigate the time

course distances of the ICNs, as a measure of the correlation, for both the sfCICA and the CICA

model at the subject level for different synthetic groups.

Distance analysis on the synthetic dataset predominantly revealed smaller distances after

performing structural connectivity constraints using the sfCICA model compared to the CICA

model, shown in **Figure 3.** This is reflected by average paired distances for the sfCICA (red box)

and the CICA (green box) model. **Figure 3,** column **(a)** displays average paired distances at the

subject level for each ICN. In all three groups of synthetic datasets, the average distances of the

sfCICA were primarily constrained by structural connectivity and decreased compared to the

CICA model. The corresponding group structural connectivity is shown in column **(b)**.

Statistical analysis, employing a paired t-test, revealed significant differences (p-value <0.05, FDR

correction: q< 0.05) in the average group distances between the sfCICA and the CICA model for

all the ICNs, except for ICN 14 in Group B, and ICN 1, 3, 7, 8, 11, 13, 14, 15 and, 16 in Group C.

Figure 3. (**a**) Shows the average time course distance of the three groups of synthetic data, comparing the sfCICA

(red box) to CICA (green box) at the subject level for each ICN. (**b**) Displays the corresponding group-level structural

connectivity.

Figure 4. represents the time course distance of an ICN (here ICNs 2 and 15) from other connected

ICNs at the subject level across three different groups using both the sfCICA and CICA models.

We specifically selected ICNs with varying connection weights: one exhibiting a high structural

connection (ICN number 2) and another with a low structural connection (ICN number 15) to other

ICNs. The results in **Figure 4.** illustrate a greater decrease in time course distance for the connected

ICNs in the sfCICA model (red box) compared to the CICA (green box). As expected, in the

subject-level analysis, the distance between ICNs in the sfCICA model decreased more

significantly than those in the CICA model.

(a)   (b)

Figure 4. It depicts the estimated time course distance of ICN 2 and 15 with other connected ICNs at the subject

level, shown in columns (**a**) and (**b**) respectively. The time course distance of the ICNs in sfCICA model was

lower compared to CICA. The corresponding structural connectivity of ICN 2 and 15 with other ICNs comes at

top of each column (**a**) and (**b**).

Moreover, we examined changes in distance specifically between two pairs of ICNs: one with a

connection (between ICNs 2- 4) and another without a connection (between ICNs 2- 13), across

three different synthetic datasets in **Figure 5 (a-c)**. The time course distance between ICN 2 - 4

(with strong structural connectivity, M = 0.66) and ICN 2 -13 (without structural connectivity) was

investigated at the subject level. Results show that the proposed model exhibits a lower distance

compared to CICA for ICN 2-4 consistently across all subjects and various groups with different

noise levels. However, for ICNs 2 - 13 (without structural connectivity)**,** the time course distance

for both sfCICA and CICA models was similar. The distance for the connected ICNs (e.g. 2-4) is

higher than that for the unconnected ICNs (e.g. 2-13) when comparing the distance results between

ICN 2- 4 and ICN 2- 13 at the subject level. The decrease in time course distance observed for

connected ICNs in the sfCICA model is greater than in the CICA model. This reflects the influence

of the structural connection and the adaptivity of our model in fluctuating the ICN estimation. For

the non-connected ICNs (without structural connectivity), only functional information was

effective, and the observed changes were consistently at a similar level for both models.

Figure 5. The time course distance of ICN 2 with a connected ICN (e.g 4) and a non-connected ICN (e.g. 13) is represented

for synthetic data with three different noise levels **(a-c)**. The distance decreased for connected ICNs (e.g. ICN 2- 4), while

the disconnected ICNs had similar distances across all subjects.

Statistical analysis using a two-sample t-test reveals an increase in various graph metrics,

(modularity: 0.352 ± 0.01, global efficiency: 0.157 ± 0.01, local efficiency: 0.199 ± 0.009, small-

worldness: 1.415 ± 0.055, non-randomness: 5.0 ± 1.15) for sfCICA compared to CICA (modularity:

0.351 ± 0.01, global efficiency: 0.155 ± 0.01, local efficiency: 0.198 ± 0.009, small-worldness:

1.412 ± 0.006, non-randomness: 4.18 ± 1.15). A slight, but insignificant decrease (p-value>0.05) was

observed between the sparsity measurements of the CICA (Mean$\pm$Std: 0.28 ± 0.01) and sfCICA

(Mean$\pm$Std: 0.27 ± 0.01) models.

3.2. FNC is informed by structural connectivity weights in synthetic data.

To emphasize the influence of the structural connectivity information on the FNC of ICNs in

synthetic data, we assess the similarity of the FNC with structural connectivity, as well as the

residual FNC for both the proposed sfCICA and the CICA models. The residual FNC is determined

as the difference between the FNC matrices from the sfCICA and CICA models (FNC_{diff} =

FNC_{sfCICA} - FNC_{CICA}).

We perform two different evaluations by computing the similarity of the FNC matrix to determine

the degree to which the structural connectivity impacted the resulting model weights, as shown in

Figure. 6. For this purpose, the FNC matrix of the estimated ICNs from the sfCICA model was

determined by computing Pearson's correlation between paired ICNs for each subject. The

similarity of the FNCs was then computed: (1) with structural connectivity and, (2) with FNC_{diff},

at the subject level. A similar procedure was performed on the ICNs from the CICA model, and

the results from both the sfCICA and CICA models were compared. From both evaluations,

FNC_{sfCICA} demonstrated higher similarity with FNC_{diff} and structural connectivity than FNC_{CICA}.

These results suggest that the proposed sfCICA model is informed by structural information and

remains robust to noise. Even with an increase in the noise level from Group A to C, the proposed

model showed improvement compared to the CICA model. However, FNC_{sfCICA} and FNC_{CICA}

exhibited the same level of similarity with structural connectivity.

Figure 6. Similarity (Pearson correlation) of the FNC matrix with structural connectivity and with FNC_{diff} (FNC_{diff} = FNC_{sfCICA} - FNC_{CICA}) was estimated for both the sfCICA and the CICA model across three distinct groups of synthetic data (columns). The results indicate that the sfCICA model exhibits a higher correlation with both structural connectivity and FNC_{diff} compared to CICA, as expected. Notably, greater differences were observed for FC_{diff}. Furthermore, the findings demonstrate the robustness of the proposed model to noise, as evidenced by its consistent performance even as noise increases from Group A to C.

3.3. The subject-level structural-functional ICNs show more modular and integrated networks for the FBIRN data.

Using the sfCICA model, the ICNs are guided by structural connectivity and FNC for each subject.

In this paper, we used the NeuroMark template (Du et al., 2020) as prior spatial information, and

the included information was evaluated by estimating the spatial similarity between the estimated

ICNs and the corresponding NeuroMark template networks using Pearson's correlation for both

the sfCICA and the CICA models. To analyze the effectiveness of the proposed model for both

healthy (HC) and disease (SZ) cases, we presented separate results for each group. In **Figure 7a,**

the results showed an unimodal Gaussian distribution for HC and a bimodal Gaussian-like

distribution in SZ, which is more clearly highlighted in the proposed model compared to the CICA

model. It illustrates more significant spatial similarity for all subjects in the sfCICA model (Mean:

0.52) compared to the CICA model (Mean: 0.47) across both HC and SZ cohorts from the FBIRN

dataset. On average, the sfCICA model exhibited a 10% increase in similarity compared to the

spatial similarity estimated by the CICA model. The enhanced spatial similarity observed with

standard and reproducible templates across multiple subjects suggests that informing the

optimization process using structural connectivity could improve the results.

Moreover, in **Figure 7b**, we computed and depicted the time course distance between paired ICNs

at the subject level to assess the impact of structural connectivity. As we expected, the estimated

distance showed a dependence on the structural connectivity weights, being consistently smaller

for all the ICNs estimated by sfCICA in comparison to the CICA model. Overall, the distances, as

well as the average distance of the time courses, were smaller in sfCICA than in CICA for both

HC and SZ subjects, indicating greater functional integration (paired t-test: p-value<0.05).

However, the increased spatial similarity resulted in more modular networks compared to the

CICA model. In this context, distance refers to the mean square distances between paired time

courses.

(a) (b)

Figure 7. Distribution of average spatial similarity and time course distance over the ICNs at the subject level were represented for each model using FBIRN data. In (a), the distribution of the spatial similarity (Pearson correlation) with NeuroMark_01 template, used for prior spatial information, is illustrated for both the CICA and the sfCICA models in two groups: HC in green and SZ in red. Mean square distances of each time course from remaining time courses are shown in (b) within each HC (green) and SZ (red) group using the sfCICA and the CICA models. The sfCICA model showed higher spatial similarity and reduced time course distances.

3.4. The subject-level structural-functional ICNs show a structural and functional learning pattern for the FBIRN data.

Figure. 8 illustrates the differences in the estimated time course distances (a measure of the correlation) between the sfCICA and the CICA model for each ICN at the subject level. Negative distance values (below the pink line) indicate lower distances among ICNs in the sfCICA model compared to CICA. Interestingly, we observed predominantly negative correlations in the FBIRN dataset. In addition, ICN 18, 29, 37, 40, 4, 47, 48, 51, 52, and 53 show, on average, positive differences across all subjects. In general, we expect that ICNs with weak mean structural

connectivity weight (as shown in **Figure. 8**) will show higher distances in the sfCICA model than

in the CICA model, resulting in a positive average pattern for distance differences. **Figure. 8**

reveals that the weights of the structural connectivity are associated with the distance of the time

courses, i.e., where the structural connectivity is weak, the distance is large, and where the

structural connectivity is strong, the distance is small among the sfCICA relative to the CICA time

courses. These results suggest that both structural weights and functional spatial maps contribute

complementary information to the estimated ICNs.

Figure 8. Illustrates that the differences in time course distances for most of the Intrinsic Connectivity Networks (ICNs) estimated by the sfCICA model, as compared to the CICA model in the FBIRN dataset, are smaller. The average distance differences of the time courses are predominantly negative, signifying smaller distances in the sfCICA compared to the CICA model, and this difference is influenced by Structural Connectivity.

We also estimated the time course distance for six randomly selected paired ICNs including 13-

22, 48-47, 13-52, 12-48, 2-37, and 3-5, with different structural connectivity weights from weak

to strong in the FBIRN data. As depicted in **Figure 9**, and similarly to the synthetic dataset, we

observed distinct time course distances between sfCICA and CICA, with a decrease in sfCICA as

the structural connectivity weights increased. On average over all subjects, our model (red box)

has a higher distance for ICN 13 with 22 and 52 (both weak structural connections) compared to

the CICA model (green box). For ICN 48, we consider two moderate (structural connectivity

weight, M = 0.2) and weak connections (0.002) respectively with ICN 12 and 47. Interestingly,

the distance between ICN 48 and 47 was increased compared to the CICA model, similar to the

distance between ICN 13 and 52 (structural connectivity weight, M = 0.037). Conversely, the

distance between ICN 48 and 12 was lower in our model. We evaluated the distance for the ICNs

with strong structural connection, ICN 37 with 2 (M = 0.66) and ICN 3 with 5 (M = 0.91). As

expected, for both connections, the distance decreased compared to the CICA model.

638 **Figure 9.** Depicts the distribution of time course distances for six paired ICNs estimated by both sfCICA and CICA

639 models in the FBIRN dataset. These, paired ICNs were selected with varying structural connectivity weights ranging

640 from weak to strong (0 - 0.91) connections. The estimated distance demonstrates an increase in structural connectivity

641 weights, resulting in a decreased time course distance for the sfCICA model (red box) compared to the CICA model

(green box). This highlights the influence of structural connectivity on the estimated ICNs.

3.5. Distinguished spatial patterns for the ICNs in the sfCICA compared to the CICA model using the FBIRN data.

Significantly, we observed enhanced modularity and well-defined spatial maps in certain functional networks represented by ICNs IC 2, 22, 42, 7, 11, 21, 13,14, 34, 37, 50, 3, 27, 39, and 42, across most subjects, especially in those diagnosed with schizophrenia. These observations indicate that the structural connectivity constraints assist in better distinguishing between healthy and patient subjects within these specific ICNs. These functional networks mostly include subcortical, sensory networks (sensorimotor and, visual), default mode, and the cerebellum. In **Figure 10,** three different ICNs for subjects 29, 30, 37, and 74 are depicted in each row. All ICNs were converted to z-scores and thresholded at z-score>3, p-value<0.05. Additionally, a qualitative examination reveals more modular networks compared to the CICA model, **(**refer to **Figure 11).**

Moreover, we performed paired t-test analysis between sfCICA and CICA using the spatial maps within each group (HC / SZ) separately. In healthy subjects, significant differences (p-value<0.05) were observed in most of the ICNs, except for the ICNs 45 and 51. ICNs 9, 27, and 51 in schizophrenia showed no significant differences between sfCICA and CICA.

In addition, we computed graph-based network features such as modularity and randomness. Utilizing FNC matrices derived from the estimated Intrinsic Connectivity Networks (ICNs), our analysis revealed that the modularity coefficient for ICNs estimated with the sfCICA model was statistically significant (paired t-test: p-value < 0.05, Mean±std = 0.24±0.03) compared to the CICA model (Mean±std = 0.21±0.02). A randomness analysis was also conducted, indicating significant differences in non-randomness. Specifically, the sfCICA model exhibited a lower randomness coefficient (i.e., higher non-randomness characteristic) in the FNC patterns (paired t-test: p-value < 0.05, Mean±std = 13.3±3.5) compared to the CICA model (Mean±std = 11.5±2.05).

Figure 10. Each column displays spatial maps for selected ICNs (z-score>3) derived from different subjects using sfCICA, CICA, and the NeuroMark template (NM). The sfCICA consistently exhibits more modular characteristics (average modularity for sfCICA: 0.24, for the CICA: 0.21) across different subjects compared to the CICA model.

682
683
684
685 **3.6. FNC of the Structural-functional connectivity informed ICNs revealed significantly**

686 **constrained/ learned from structural connectivity using the FBIRN data.**

687
To determine the differences between the CICA and sfCICA models, we applied a paired t-test

analysis on FNC matrices. Interestingly the paired t-tests indicated rejection of the null hypothesis,

signifying significant differences between sfCICA and CICA across all ICNs. The estimated t-

values (**Figure. 11a**) represented significant differences in ICNs related to the SC, SM, VS, CC,

and DM networks. Paired t-test analysis was also performed regarding the absolute value for the

FNCs, as illustrated in **Figure. 11b,** it shows reduced t-values for certain ICNs within the AUD,

SM, VS, CC, DM, and CB networks. It replicates a decrease in the FNC association between the

two models and can be explained based on the hypothesis that the impact of the relationship

between structural connectivity and FNC on cognitive performance may depend on the functional

domain.

Moreover, regarding **Figure. 11 (a-d)**, and the comparison of structural connectivity patterns

(**Figure 11(c-d)**) with paired t-test analysis (**Figure 11(a-b)**), it is apparent that the structural

connectivity weights optimize the distances, thereby enhancing the estimation of the ICNs.

Further, analysis of the paired t-test between ICNs of both models reveals similarity for ICNs 1,

6, 11, 12, 14, 15, 18, 20, 22, 29, 39, 40, 43, 45, 50, and 52.

Figure 11. Illustrates the results for the paired t-test analysis between the FNC matrices of the sfCICA and CICA models, along with their corresponding structural connectivity, using FBIRN dataset. FNC represents the Pearson correlation between the estimated ICNs of each model. In (**a**). T-value maps for the paired t-test using FNC are shown, while (**b**) displays the t-map for the absolute FNCs. sfCICA model exhibits higher values in most ICNs compared to the CICA model. The group average structural connectivity, and its logarithmic scale connection weights (*log₁₀(connection weights)*), are represented in (**b**) and (**c**). The weights of the structural connectivity indicate the number of tracts connecting two ICNs.

The determined results were jointly estimated by incorporating information from the structural

connectivity weights into the correlation between the time courses overall time (i.e., the FNC

matrix). A similar analysis on synthetic data was conducted on real data as well to assess the

similarity of the FNC with structural connectivity and FNC with FNC_{diff}, determining the extent

to which structural connectivity impacts the resulting model weights, as illustrated in **Figure. 12**.

For this purpose, an FNC matrix was computed for each subject using Pearson's correlation among

the time courses of the ICNs determined by the sfCICA model. A similar procedure was performed

to estimate FNC_{CICA} using the ICNs of the CICA model. The comparison of results from both the

sfCICA and CICA models indicated that in real datasets (FBIRN), FNC_{sfCICA} exhibits greater

similarity with structural connectivity and FNC_{diff} to FNC_{CICA}. It indicates the proposed sfCICA

model is informed by structural information, showing the multi-modal sfCICA model provides

additional information that could capture variability between subjects and between groups.

(a) (b)

799 **Figure 12.** Similarity (Pearson correlation) between FNC matrix of estimated ICNs from the sfCICA model with

800 structural connectivity and, residual (diff) FNC matrix is represented for FBIRN dataset respectively in (**a**) and (**b**).

801 Results show similarity (correlation) with both structural connectivity and residual FNC (FNC_{diff}) is higher for the

802 sfCICA model than the CICA, representing more informatic ICNs.

3.7. Structural-functional connectivity-informed ICNs reveal significant differences between controls and schizophrenia subjects using the FBIRN data.

Figure. 13 (**a-b**) show the average FNC across all subjects for the sfCICA and CICA models. Additionally, group differences analysis using FNC for the HC vs SZ was performed and the results are represented in **Figure. 13 (c-d).**

Furthermore, to investigate the ability of the proposed method to find group differences (healthy/ patient) in FNC patterns using both the sfCICA and the CICA models, a generalized linear regression model (GLM) was performed with multiple covariates (**Figure. 13** (**c-d**)). The results indicated a more significant correlation (p-value<0.05, FDR: q<0.03) across all networks (displayed with red stars) comparing HC vs SZ group for the sfCICA model. We detected 503 significant cells in the upper triangle for the sfCICA model (**Figure 13 c**), a notable increase compared to the 421 significant cells observed for the CICA model (**Figure 13 d**). Correlations of the sensory networks were less significant in the sfCICA model compared to the CICA. In contrast, SC-related regions showed more significance. The findings suggested the involvement of structural connectivity in the SC domain, a factor not observed with the CICA model. ICNs for the SC, DMN, and CB networks exhibited significantly higher FNC in HC compared to SZ for the sfCICA model (**Figure. 13 c**), while sensory-related networks (AUD, VS, and SM) and CC network showed significantly higher FNC in HC compared to SZ for both models (**Figure. 13 (c-d)**). This suggests that considering structural connectivity plays an important role in identifying the FNC of the SC and that the sfCICA which jointly analyzes structural and functional ICA output provides a more sensitive model. The values illustrated in the higher triangle of matrices in **Figure 13 (c-d)**, are -log10(p_value) × $sign$ (t_value) and t-values are represented in the lower triangle.

Figure 13. Average FNC estimated across all subjects and statistical group difference maps (HC/ SZ) for the FBIRN

dataset. In **(a)** and **(b),** the average FNC of all subjects is presented for the sfCICA and CICA models respectively. For

the group (HC/ SZ) difference comparison, GLM analysis was performed to regress out covariates including age, sex,

motion, imaging site, and diagnosis from the FNC matrices. More significant differences (p-value<0.05 and FDR<0.03),

represented by red stars, were observed between HC and SZ in the proposed sfCICA compared to the CICA model.

Results were corrected for multiple comparisons using false discovery rate (FDR) with q<0.03; and represented in **(c-**

d), which upper triangle each cell of the matrices are *-log₁₀(p-value) *sign(t-value)* and lower triangle is t-value. ICNs

were categorized in seven different domains including subcortical network (SC), auditory network (AUD), sensorimotor

network (SM), visual network (VS), cognitive control network (CC), default mode network (DM) and cerebellum (CB).

Results show that FNC in HC are higher than SZ in most of the ICNs, in the sfCICA compared to the CICA model.

3.8. Multimodal ICNs represent more significant spatial maps between groups (controls and schizophrenia) using the FBIRN data.

To investigate the hypothesis of the proposed model, we conducted a group differences analysis between HC and SZ on the ICN spatial maps. For this purpose, a GLM analysis was adopted to regress out the effect of age, sex, motion, site, and diagnosis per ICN. Table 1 displays the percentage of significant voxels in each ICN of different networks. Results indicate that out of fifty-three ICNs, thirty-two showed a higher number of significant voxels (p-value<0.05) in the sfCICA model (dark blue rows). Conversely, in the remaining twenty-one ICNs, five (light blue row) demonstrated a similar percentage of significant voxels, and sixteens (white rows) represented a lower percentage with a deviation for the sfCICA model.

In addition, network properties including modularity, efficiency, small-worldness, randomness, and sparsity were estimated using FNC features. Results (Table 2) showed significantly higher (p-value<0.05) efficiency, modularity, and randomness for the sfCICA model using the FNC feature. Small-worldness was also greater for sfCICA but the difference between the two models was not significant.

Table 1. Shows the percentage of the significant voxels in each ICN using group differences analysis. ICNs (rows) with higher values for the sfCICA are represented in dark blue, while light blue and white indicate equal or lower values for the sfCICA compared to the CICA.

Networks	ICNs	Number of significant voxels using CICA (%)	Number of significant voxels using sfCICA (%)	Differences between significant voxels, sfCICA – CICA (%)
SC	IC1	4.9	5.6	0.7
	IC2	4.1	6	1.9
	IC3	4.2	4.9	0.7
	IC4	3	3.5	0.5
	IC5	8.3	9.6	1.3
AUD	IC1	2.8	2.9	0.1
	IC2	4.7	5.4	0.7
SM	IC1	2.1	2.9	0.7
	IC2	2.8	3.4	0.6
	IC3	7.1	5.7	-1.4
	IC4	8.3	9.1	0.8
	IC5	5.5	5.8	0.3
	IC6	4.4	4.2	-0.2
	IC7	3.9	3.9	0
	IC8	6.6	7	0.4
	IC9	5	5.1	0.1
VS	IC1	3.2	4.1	0.9
	IC2	10	10	0
	IC3	5.1	5.2	0.1
	IC4	8	6.8	-1.2
	IC5	3.5	4.2	0.7
	IC6	3.1	3.4	0.3
	IC7	3.8	4.5	0.7
	IC8	2.4	2.4	0
	IC9	4.1	4.8	0.7
CC	IC1	5.1	5.1	0
	IC2	8.2	7.3	-0.9
	IC3	5.5	5.9	0.4
	IC4	5.3	5.2	-0.1
	IC5	6.6	6.7	0.1
	IC6	6.7	6.2	-0.5
	IC7	5.2	6	0.8
	IC8	8.9	6.3	-2.6
	IC9	8	8.3	0.3
	IC10	7	6.4	-0.6
	IC11	9.2	8	-0.8
	IC12	3.3	3.7	0.4
	IC13	8	8.3	0.3
	IC14	4.3	5.5	1.2
	IC15	7	5.4	-1.6
	IC16	4.2	4.2	0
	IC17	5.6	5.3	-0.3
DM	IC1	3.1	3.4	0.3
	IC2	3.5	3.2	-0.3
	IC3	2.5	3.2	0.7
	IC4	8.2	8.1	-0.1
	IC5	4.5	3.6	-0.9
	IC6	5.1	5.2	0.1
	IC7	7.6	6	-1.6
CB	IC1	2.1	4.8	2.7
	IC2	9.1	6.4	-2.7
	IC3	2.1	5.9	3.8
	IC4	4.6	5.6	1

Graph parameters	Single-modality Model (**Mean±std**)	Multi-modality Model (**Mean±std**)
Modularity	**0.21±0.02**	**0.24±0.03**
Local efficiency	**0.32±0.01**	**0.37±0.01**
Global efficiency	**0.30±0.03**	**0.33±0.03**
Small-worldness	**1.15±0.08**	**1.17±0.06**
Non- Randomness	**11.30±3.50**	**13.50±2.05**
Sparsity	**0.28±0.01**	**0.27±0.01**

4. Discussion

This study proposes a new, joint structural-functional, data-driven model to estimate intrinsic

functional connectivity networks (ICNs) using multi-modal brain images. While most of the

studies typically rely on fixed regions weighted by diffusion MRI (dMRI), our proposed approach

is a multi-modal ICA model, named sfCICA, which is guided by the structural and functional

connectivity network domain. Using network science, structural and functional information is

compressed in nodes and edges which has become a common practice in neuroscience to

understand functional interactions in the brain (Babaeeghazvini et al., 2021). In the sfCICA model,

the functional connectivity from resting-state magnetic resonance imaging is constrained by

structural connectivity weights (normalized number of streams). We applied the proposed model

to datasets from subjects with schizophrenia and healthy controls, as well as simulated data.

Through this new model, we identified improved spatial maps, individual subject variability, and

modularity of the FNC networks. Furthermore, the sfCICA model exhibits less randomness and

greater sensitivity to group differences in healthy controls and subjects with schizophrenia (HC/SZ

subjects).

Most previous ICA models are focused on functional information or functional information with

prior spatial constraints (Du & Fan, 2013; Du et al., 2020). In recent years, there has been a shift

towards multi-modal models, incorporating features estimated from different modalities through

joint analysis (Wu & Calhoun, 2023). Joint-ICA, parallel ICA, link-ICA, and cmICA models

utilize features from fMRI, dMRI, or sMRI, such as connectivity, fractional anisotropy, and

structural measures to identify functional networks. The proposed model in this study diverges by

simultaneously learning from both functional and structural connectivity. This approach aims to

constrain functional changes over time based on estimates of structural connectivity.

Less work has been given to dMRI connectivity, as in the cmICA model (Wu & Calhoun, 2023).

cmICA (Wu & Calhoun, 2023), uses multi-modal fMRI and dMRI, integrating features derived

from each modality to identify ICNs shared between structural connectivity and FNC,

acknowledging that there might be a mismatch between features from each modality. Although

cmICA attempts to address this, the contributions of features from each modality are asymmetrical

(Wu & Calhoun, 2023). In contrast, our proposed model is based on both direct and indirect

features and the contribution of each modality enhances the other. Thus, when structural

connectivity weights are strong, functional data exhibits a higher correlation, while weaker

structural connectivity weights result in less correlated functional data. Notably, the sfCICA model

relies on two main concepts of an ICA analysis, maximizing independence and similarity, and tries

to estimate intrinsic brain functional networks which are optimized based on both structural and

functional connectivity information. Therefore, we posit that the proposed joint multi-modal

information integration allows for better characterization of healthy and disordered brain

connectivity (Iraji et al., 2016; Wu & Calhoun, 2023). Characterization of the intrinsic brain

activity, known as functional network modeling, has been widely used in human brain studies,

particularly because of its relevance to brain disorder research (Wang et al., 2021). One of the

greatest advancements in this area was uncovered by rs-fMRI images using ICA analysis (Calhoun

et al., 2001) and continues to advance for the field by the inclusion of additional imaging

modalities.

Including two different modalities of information in our model, the replicability and noise

robustness of the proposed model were assessed through three different synthetic datasets (**Figures**

3, 4, and 5). Each group has different levels of signal-to-noise ratio (SNR), and for each group, a

similar pattern, constrained by structural connectivity, was observed. Our findings highlighted the

effect of the structural connectivity constraints on the time courses, as their distances were

significantly smaller when structural connectivity was stronger. Additionally, by changing the

functional information and keeping the structural connectivity constant, the learning procedure is

dependent on information from both modalities during the estimation of ICNs.

Empirical results from our proposed multi-modal model on real data show that our model (the

sfCICA) is more sensitive in detecting functional connectivity in both healthy and schizophrenia

groups, compared to the CICA model. Estimated ICNs were significantly more spatially similar

(**Figure 7a**) to the reproducible template (NeuroMark template (Du et al., 2020)) for both healthy

and schizophrenia subjects. This may enhance the replicability of the components using different

datasets (Duda et al., 2023).

Moreover, the proposed sfCICA model aims to estimate a unified and integrated brain network

using regional functional information instead of FNC as in (Zhu et al., 2021) which is the first time

this has been performed in an ICA analysis. Overall, the results in **Figure 7b** showed that the

distance of the time courses was smaller, which can be interpreted as more functionally correlated,

and hence more uniform ICNs. While considering each ICN specifically (**Figures 8 and 9**), ICNs

related to cognitive control (CC), and cerebellum (CB) networks were less constrained. In other

words, distance minimization of them was less compared to other ICNs. It is highlighted that

overall, they are less structurally connected to other ICNs. For instance, ICNs number 37 and 41

in the CC network, represent the weakest connection with CC, SM, and CB networks. Also,

consistent with prior findings (Glasser et al., 2016; Van Essen et al., 2019; Wu & Calhoun, 2023),

the DM has the largest distances within its network ICNs and the weakest with intra-network such

as SC network. These findings suggest the advantages of multi-modal approaches, using a

combination of structural and functional features, in facilitating the interpretability of the results

which is consistent with previous findings (Glasser et al., 2016; Van Essen et al., 2019).

Alternatively, there is a hypothesis that structural connectivity may directly influence FNC, as

higher structural connectivity leads to higher FNC (Litwińczuk et al., 2022; Zhao et al., 2023).

Statistical analysis on FNC of the estimated multi-modal (sfCICA) ICNs and single model ICNs

(CICA) in **Figure 11 (a-b)**, represented no association between the FNCs of the two models in

specific domains including SM, AUD, VS, CC, and DM. Interestingly results were following

previous findings (Li et al., 2020; Tsai et al., 2019). There is evidence that structural connectivity

and FNC have no coupling in SM and VS, which means either structural connectivity is not dense,

or it may come from dysfunction characteristics of the SZ subjects accompanied by weaker

structural connectivity in SM, anterior cingulum cortex (ACC, related to DM domain) (Li et al.,

2020). So, our results revealed it in the estimated ICNs features (FNC) using the sfCICA model.

Regarding the VS domain, little is known about it, however, it has been shown that the connectivity

of the visual system in SZ would be reduced (Reavis et al., 2020) which is consistent with our

findings. Overall, findings suggested that integrating multi-modal information to estimate ICNs

could help to find more accurate intrinsic functional domains in comparison to single-modality

models such as the CICA.

Our proposed model (sfCICA) was more sensitive to detecting differences between individuals

with schizophrenia patients and healthy controls. In schizophrenia both structural and functional

networks are affected (Li et al., 2017; S. Qi et al., 2019), our results showed a broad range of

impacted networks. Regarding the FNC matrices (**Figure 13 a-b**), findings are consistent with, and

extend, those of prior studies. For example, prior studies have reported dysfunction in the CC

network, which was also observed in our statistical results using the single modality CICA model.

However, the proposed sfCICA model showed enhanced functional connectivity within the SC

network with most of the domains, the CC network with other domains (e.g., visual), and the DM

network as well as with SM, AUD, and VS. Overall, these findings suggest that the contribution

of structural connectivity information could enhance our ability to identify dysfunctional

connections. Also, ICNs related to memory and higher-order cognitive control functions related to

DM networks have been reported as affected regions in individuals with SZ (Aleman et al., 1999;

Minzenberg et al., 2009; Potkin et al., 2009; S. Qi et al., 2019). Consistent with prior findings our

results show that functional correlations of the DM network are lower for the positive connections

and higher for the negative connections (S. Qi et al., 2019), representing the effectiveness of our

multi-modal model. However, in the proposed model, mostly FNC is higher for healthy compared

to schizophrenia but there exist some ICNs that do not show the same pattern. For example, in

ICNs 8 (of the SM network) and 42 (of the CC network), FNC is higher for schizophrenia.

Compared to previous studies, mostly dysconnectivity occurs in DM and SM networks (Rong et

al., 2023), and FNC increment is observed in these domains.

There exist some limitations in this study which would be interesting to consider in future works.

We selected the model order using a standard ICN template including 53 ICNs (NeuroMark_1.0).

The identified ICNs are restricted to prior spatial maps and are modified spatially and temporally

during the optimization process. However, it might be variable over subjects, which should be

considered in future works. Moreover, the standard template is functionally defined. Our multi-

modal sfCICA model uncovers differences in functional connectivity between regions and their

interactions are affected by structural information. To include differences in spatial maps and

enable the comparability of results, a multi-modal (structural/functional) template would be

needed. In parallel, increasing structural and functional studies highlights the need for a multi-

modal simulator. Our synthetic data are independent and structural information has no impact on

fMRI simulation. It could be beneficial to develop a simulator for generating structurally informed

fMRI data (e.g. perhaps via a copula framework (Silva et al., 2014)). In addition, our current model

starts from fixed networks for the structural connectivity, it would be useful to extend the model

to also allow for adaptive updates to the structural connectivity at the single subject level, similar

to what is occurring with the fMRI data. Existing methods have used structural connectivity as a

deterministic constraint to functional connectivity (Seguin et al., 2022). In contrast, in our work,

we propose a data-driven multi-objective model using structural connectivity constraints to

incorporate these into an ICA model resulting in spatial maps, time courses, and FNC. Our

expectation is that the results would increase our sensitivity to patient vs control differences. Our

suggested model is one of the first tools for directly linking structural connectivity/FNC in a data-

driven analysis. Furthermore, future extensions can be developed that allow the structural

connectivity model to adapt to the functional data in the context of a supervised model, optimizing

structural connectivity/FNC to maximize group differences and then using this in a new dataset to

classify. Other future work could be to use our model with an alternative transformation of the

structural connectivity network. Other future work could be to use our model with an alternative

transformation of the structural connectivity network. Most of the not directly connected regions

have polysynaptic communication (Seguin et al., 2022). One way to guide polysynaptic signals in

structural connectivity/FNC models is by using communication network models (Seguin et al.,

2022). Communication network models are derived from structural connectivity and converse to

the structural connectivity, communication network models are dense (Seguin et al., 2022). One

limitation of data fusion is that the structural connectivity informed FNC may be more complex to

interpret than the FNC estimated from fMRI alone. However, there are also advantages as we point

out, especially in enhancing sensitivity to group differences. One way to address this is to perform

both analyses as we have done. An advantage of the use of the NeuroMark spatially constrained

ICA framework is that the data-driven networks maintain their correspondence with one another

due to the similar spatial priors in both. On the other hand, regions-based structural connectivity

was identified before optimization, and it remains constant during resting-state data acquisition. In

addition, for future works, it might be more informatic to use diffusion spectrum imaging (DSI).

Although time courses and spatial maps are updated, the model can be extended to modify

structural weights throughout the optimization process. Beyond this, we can also incorporate the

dynamics of brain function.

1051

5. Conclusion

In this work, we proposed a novel, structural-functional constrained independent component analysis (sfCICA), which incorporates both structural and functional connectivity information, guided by spatial maps to estimate functional networks in the human brain. The main motivation for the proposed model is to allow for structural and functional information to jointly influence the model estimation. The resulting ICNs were more spatially focal, and more synchronized in the sfCICA model compared to the CICA model. In addition, we observed that integrating structural connectivity information in the sfCICA enhances the sensitivity to group differences. The results were consistent with those based on synthetic structural and functional imaging data, with different functional information and noise levels. In sum, our findings demonstrate the effectiveness of the proposed model in both synthetic and real data. The approach applies to the study of a wide range of areas including brain development, aging, healthy control, and mental disorders.

6. Acknowledgment

This work is funded in part by the NSF grant 2112455 and the NIH grants R01MH118695 and R01MH123610.

7. Authors contribution

Mahshid Fouladivanda: Conceptualization; Formal analysis; Investigation; Methodology; Software; Visualization; Writing – original draft; Writing – review & editing. Armin Iraji: Conceptualization; Formal analysis; Writing – review & editing. Lei Wu: Software. Theodorus G.M. van Erp: Resources; Writing – review & editing. Aysenil Belger: Resources. Faris

Hawamdeh: Writing – review & editing. Godfrey Pearlson: Writing – review & editing. Vince Calhoun: Conceptualization; Funding acquisition; Investigation; Methodology; Project administration; Resources; Software; Writing – review & editing.

References

Aleman, A., Hijman, R., De Haan, E. H., & Kahn, R. S. (1999). Memory impairment in schizophrenia: a meta-analysis. *American Journal of Psychiatry*, *156*(9), 1358-1366.

Allen, E. A., Erhardt, E. B., Damaraju, E., Gruner, W., Segall, J. M., Silva, R. F., Havlicek, M., Rachakonda, S., Fries, J., Kalyanam, R., Michael, A. M., Caprihan, A., Turner, J. A., Eichele, T., Adelsheim, S., Bryan, A. D., Bustillo, J., Clark, V. P., Feldstein Ewing, S. W., . . . Calhoun, V. D. (2011). A baseline for the multivariate comparison of resting-state networks. *Front Syst Neurosci*, *5*, 2. https://doi.org/10.3389/fnsys.2011.00002

Amaral, L. A. N., Scala, A., Barthélémy, M., & Stanley, H. E. (2000). Classes of small-world networks. *Proceedings of the National Academy of Sciences*, *97*(21), 11149-11152. https://doi.org/doi:10.1073/pnas.200327197

Andersson, J. L. R., & Sotiropoulos, S. N. (2016). An integrated approach to correction for off-resonance effects and subject movement in diffusion MR imaging. *Neuroimage*, *125*, 1063-1078. https://doi.org/10.1016/j.neuroimage.2015.10.019

Avants, B. B., Tustison, N., & Song, G. (2009). Advanced normalization tools (ANTS). *Insight j*, *2*(365), 1-35.

Babaeeghazvini, P., Rueda-Delgado, L. M., Gooijers, J., Swinnen, S. P., & Daffertshofer, A. (2021). Brain Structural and Functional Connectivity: A Review of Combined Works of Diffusion Magnetic Resonance Imaging and Electro-Encephalography. *Front Hum Neurosci*, *15*, 721206. https://doi.org/10.3389/fnhum.2021.721206

Bassett, D. S., Bullmore, E. T., Meyer-Lindenberg, A., Apud, J. A., Weinberger, D. R., & Coppola, R. (2009). Cognitive fitness of cost-efficient brain functional networks. *Proceedings of the National Academy of Sciences*, *106*(28), 11747-11752. https://doi.org/doi:10.1073/pnas.0903641106

Batista-García-Ramó, K., & Fernández-Verdecia, C. I. (2018). What We Know About the Brain Structure-Function Relationship. *Behav Sci (Basel)*, *8*(4). https://doi.org/10.3390/bs8040039

Calhoun, V. (2018). Data-driven approaches for identifying links between brain structure and function in health and disease. *Dialogues in Clinical Neuroscience*, *20*(2), 87-99. <Go to ISI>://WOS:000442461300002

Calhoun, V. D., & Adalı, T. (2012). Multisubject independent component analysis of fMRI: a decade of intrinsic networks, default mode, and neurodiagnostic discovery. *IEEE Rev Biomed Eng*, *5*, 60-73. https://doi.org/10.1109/rbme.2012.2211076

Calhoun, V. D., Adali, T., Kiehl, K. A., Astur, R., Pekar, J. J., & Pearlson, G. D. (2006). A method for multitask fMRI data fusion applied to schizophrenia. *Human Brain Mapping*, *27*(7), 598-610. https://doi.org/10.1002/hbm.20204

Calhoun, V. D., Adali, T., Pearlson, G. D., & Pekar, J. J. (2001). A method for making group inferences from functional MRI data using independent component analysis. *Hum Brain Mapp*, *14*(3), 140-151. https://doi.org/10.1002/hbm.1048

Calhoun, V. D., & de Lacy, N. (2017). Ten Key Observations on the Analysis of Resting-state Functional MR Imaging Data Using Independent Component Analysis. *Neuroimaging Clin N Am*, *27*(4), 561-579. https://doi.org/10.1016/j.nic.2017.06.012

1119 Calhoun, V. D., & Sui, J. (2016). Multimodal fusion of brain imaging data: A key to finding the

1120   missing link(s) in complex mental illness. *Biol Psychiatry Cogn Neurosci Neuroimaging*,

1121   *1*(3), 230-244. https://doi.org/10.1016/j.bpsc.2015.12.005

1122 Caprihan, A., Abbott, C., Yamamoto, J., Pearlson, G., Perrone-Bizzozero, N., Sui, J., & Calhoun,

1123   V. D. (2011). Source-based morphometry analysis of group differences in fractional

1124   anisotropy in schizophrenia. *Brain Connect*, *1*(2), 133-145.

1125   https://doi.org/10.1089/brain.2011.0015

1126 Chen, J., Calhoun, V. D., Pearlson, G. D., Perrone-Bizzozero, N., Sui, J., Turner, J. A., Bustillo, J.

1127   R., Ehrlich, S., Sponheim, S. R., Canive, J. M., Ho, B. C., & Liu, J. (2013). Guided

1128   exploration of genomic risk for gray matter abnormalities in schizophrenia using parallel

1129   independent component analysis with reference. *Neuroimage*, *83*, 384-396.

1130   https://doi.org/10.1016/j.neuroimage.2013.05.073

1131 Chu, S.-H., Parhi, K. K., & Lenglet, C. (2018). Function-specific and Enhanced Brain Structural

1132   Connectivity Mapping via Joint Modeling of Diffusion and Functional MRI. *Scientific*

1133   *Reports*, *8*(1), 4741. https://doi.org/10.1038/s41598-018-23051-9

1134 Cook, P. A., Bai, Y., Nedjati-Gilani, S., Seunarine, K. K., Hall, M. D., Parker, G. J. M., &

1135   Alexander, D. C. (2006). Camino: Open-Source Diffusion-MRI Reconstruction and

1136   Processing.

1137 Du, Y., & Fan, Y. (2013). Group information guided ICA for fMRI data analysis. *Neuroimage*,

1138   *69*, 157-197. https://doi.org/10.1016/j.neuroimage.2012.11.008

1139 Du, Y. H., Fu, Z. N., Sui, J., Gao, S., Xing, Y., Lin, D. D., Salman, M., Abrol, A., Rahaman, M.

1140   A., Chen, J. Y., Hong, L. E., Kochunov, P., Osuch, E. A., Calhoun, V. D., & Neuroimaging,

1141   A. D. (2020). NeuroMark: An automated and adaptive ICA based pipeline to identify

reproducible fMRI markers of brain disorders. *Neuroimage-Clinical*, *28*.

https://doi.org/ARTN 102375

10.1016/j.nicl.2020.102375

Duda, M., Iraji, A., Ford, J. M., Lim, K. O., Mathalon, D. H., Mueller, B. A., Potkin, S. G., Preda,

A., Van Erp, T. G., & Calhoun, V. D. (2023). Reliability and clinical utility of spatially

constrained estimates of intrinsic functional networks from very short fMRI scans. *Human*

Brain Mapping, *44*(6), 2620-2635.

Erhardt, E. B., Allen, E. A., Wei, Y., Eichele, T., & Calhoun, V. D. (2012). SimTB, a simulation

toolbox for fMRI data under a model of spatiotemporal separability. *Neuroimage*, *59*(4),

4160-4167. https://doi.org/10.1016/j.neuroimage.2011.11.088

Erhardt, E. B., Rachakonda, S., Bedrick, E. J., Allen, E. A., Adali, T., & Calhoun, V. D. (2011).

Comparison of multi-subject ICA methods for analysis of fMRI data. *Hum Brain Mapp*,

32(12), 2075-2095. https://doi.org/10.1002/hbm.21170

Fillard, P., Descoteaux, M., Goh, A., Gouttard, S., Jeurissen, B., Malcolm, J., Ramirez-

Manzanares, A., Reisert, M., Sakaie, K., Tensaouti, F., Yo, T., Mangin, J. F., & Poupon,

C. (2011). Quantitative evaluation of 10 tractography algorithms on a realistic diffusion

MR phantom. *Neuroimage*, *56*(1), 220-234.

https://doi.org/10.1016/j.neuroimage.2011.01.032

Genon, S., Reid, A., Langner, R., Amunts, K., & Eickhoff, S. B. (2018). How to Characterize the

Function of a Brain Region. *Trends in Cognitive Sciences*, *22*(4), 350-364.

https://doi.org/10.1016/j.tics.2018.01.010

Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E., Ugurbil,

K., Andersson, J., Beckmann, C. F., & Jenkinson, M. (2016). A multi-modal parcellation

of human cerebral cortex. *Nature*, *536*(7615), 171-178.

Goñi, J., van den Heuvel, M. P., Avena-Koenigsberger, A., Velez de Mendizabal, N., Betzel, R.

F., Griffa, A., Hagmann, P., Corominas-Murtra, B., Thiran, J. P., & Sporns, O. (2014).

Resting-brain functional connectivity predicted by analytic measures of network

communication. *Proc Natl Acad Sci U S A*, *111*(2), 833-838.

https://doi.org/10.1073/pnas.1315529111

Groves, A. R., Beckmann, C. F., Smith, S. M., & Woolrich, M. W. (2011). Linked independent

component analysis for multimodal data fusion. *Neuroimage*, *54*(3), 2198-2217.

https://doi.org/10.1016/j.neuroimage.2010.09.073

He, X., & Niyogi, P. (2003). Locality preserving projections. *Advances in neural information*

processing systems, *16*.

Honey, C. J., Sporns, O., Cammoun, L., Gigandet, X., Thiran, J. P., Meuli, R., & Hagmann, P.

(2009). Predicting human resting-state functional connectivity from structural

connectivity. *Proceedings of the National Academy of Sciences*, *106*(6), 2035-2040.

https://doi.org/doi:10.1073/pnas.0811168106

Hyvarinen, A. (1999). Fast and robust fixed-point algorithms for independent component analysis.

IEEE Trans Neural Netw, *10*(3), 626-634. https://doi.org/10.1109/72.761722

Iraji, A., Calhoun, V. D., Wiseman, N. M., Davoodi-Bojd, E., Avanaki, M. R. N., Haacke, E. M.,

& Kou, Z. (2016). The connectivity domain: Analyzing resting state fMRI data using

feature-based data-driven and model-based methods. *Neuroimage*, *134*, 494-507.

https://doi.org/10.1016/j.neuroimage.2016.04.006

Iraji, A., Deramus, T. P., Lewis, N., Yaesoubi, M., Stephen, J. M., Erhardt, E., Belger, A., Ford, J.

M., McEwen, S., Mathalon, D. H., Mueller, B. A., Pearlson, G. D., Potkin, S. G., Preda,

A., Turner, J. A., Vaidya, J. G., van Erp, T. G. M., & Calhoun, V. D. (2019). The spatial

chronnectome reveals a dynamic interplay between functional segregation and integration.

Hum Brain Mapp, *40*(10), 3058-3077. https://doi.org/10.1002/hbm.24580

Iraji, A., Faghiri, A., Fu, Z., Kochunov, P., Adhikari, B. M., Belger, A., Ford, J. M., McEwen, S.,

Mathalon, D. H., Pearlson, G. D., Potkin, S. G., Preda, A., Turner, J. A., Van Erp, T. G.

M., Chang, C., & Calhoun, V. D. (2022). Moving beyond the 'CAP' of the Iceberg: Intrinsic

connectivity networks in fMRI are continuously engaging and overlapping. *Neuroimage*,

251. https://doi.org/ARTN 119013

10.1016/j.neuroimage.2022.119013

Iraji, A., Faghiri, A., Fu, Z., Rachakonda, S., Kochunov, P., Belger, A., Ford, J. M., McEwen, S.,

Mathalon, D. H., Mueller, B. A., Pearlson, G. D., Potkin, S. G., Preda, A., Turner, J. A.,

van Erp, T. G. M., & Calhoun, V. D. (2022). Multi-spatial-scale dynamic interactions

between functional sources reveal sex-specific changes in schizophrenia. *Netw Neurosci*,

6(2), 357-381. https://doi.org/10.1162/netn_a_00196

Iraji, A., Fu, Z., Faghiri, A., Duda, M., Chen, J., Rachakonda, S., DeRamus, T., Kochunov, P.,

Adhikari, B. M., Belger, A., Ford, J. M., Mathalon, D. H., Pearlson, G. D., Potkin, S. G.,

Preda, A., Turner, J. A., Erp, T. G. M. v., Bustillo, J. R., Yang, K., . . . Calhoun, V. D.

(2022). Canonical and Replicable Multi-Scale Intrinsic Connectivity Networks in 100k+

Resting-State   fMRI   Datasets.   *bioRxiv*,   2022.2009.2003.506487.

https://doi.org/10.1101/2022.09.03.506487

Iraji, A., Miller, R., Adali, T., & Calhoun, V. D. (2020). Space: A Missing Piece of the Dynamic

Puzzle. *Trends in Cognitive Sciences*, *24*(2), 135-149.

https://doi.org/10.1016/j.tics.2019.12.004

Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W., & Smith, S. M. (2012). FSL.

Neuroimage, *62*(2), 782-790. https://doi.org/10.1016/j.neuroimage.2011.09.015

Jorge Nocedal, S. J. W. (2006). *Numerical Optimization*. Springer New York, NY.

https://doi.org/https://doi.org/10.1007/978-0-387-40065-5

Keator, D. B., van Erp, T. G. M., Turner, J. A., Glover, G. H., Mueller, B. A., Liu, T. T., Voyvodic,

J. T., Rasmussen, J., Calhoun, V. D., Lee, H. J., Toga, A. W., McEwen, S., Ford, J. M.,

Mathalon, D. H., Diaz, M., O'Leary, D. S., Jeremy Bockholt, H., Gadde, S., Preda, A., . . .

Fbirn. (2016). The Function Biomedical Informatics Research Network Data Repository.

Neuroimage, *124*(Pt B), 1074-1079. https://doi.org/10.1016/j.neuroimage.2015.09.003

Kim, Y., Joshi, A. A., Choi, S., Joshi, S. H., Bhushan, C., Varadarajan, D., Haldar, J. P., Leahy,

R. M., & Shattuck, D. W. (2023). BrainSuite BIDS App: Containerized Workflows for

MRI Analysis. *bioRxiv*, 2023.2003.2014.532686.

https://doi.org/10.1101/2023.03.14.532686

Klamroth, K., & Jørgen, T. (2007). Constrained optimization using multiple objective

programming. *Journal of Global Optimization*, *37*, 325-355.

Latora, V., & Marchiori, M. (2001). Efficient Behavior of Small-World Networks. *Physical

Review Letters*, *87*(19), 198701. https://doi.org/10.1103/PhysRevLett.87.198701

Li, P., Fan, T.-T., Zhao, R.-J., Han, Y., Shi, L., Sun, H.-Q., Chen, S.-J., Shi, J., Lin, X., & Lu, L.

(2017). Altered brain network connectivity as a potential endophenotype of schizophrenia.

Scientific reports, *7*(1), 5483.

1231 Li, P., Jing, R. X., Zhao, R. J., Shi, L., Sun, H. Q., Ding, Z., Lin, X., Lu, L., & Fan, Y. (2020).

1232  Association between functional and structural connectivity of the corticostriatal network

1233  in people with schizophrenia and unaffected first-degree relatives. *J Psychiatry Neurosci*,

1234  *45*(6), 395-405. https://doi.org/10.1503/jpn.190015

1235 Lin, Q. H., Liu, J., Zheng, Y. R., Liang, H., & Calhoun, V. D. (2010). Semiblind spatial ICA of

1236  fMRI using spatial constraints. *Hum Brain Mapp*, *31*(7), 1076-1088.

1237  https://doi.org/10.1002/hbm.20919

1238 Litwińczuk, M. C., Muhlert, N., Cloutman, L., Trujillo-Barreto, N., & Woollams, A. (2022).

1239  Combination of structural and functional connectivity explains unique variation in specific

1240  domains of cognitive function. *Neuroimage*, *262*, 119531.

1241  https://doi.org/10.1016/j.neuroimage.2022.119531

1242 Liu, J., Pearlson, G., Windemuth, A., Ruano, G., Perrone-Bizzozero, N. I., & Calhoun, V. (2009).

1243  Combining fMRI and SNP data to investigate connections between brain function and

1244  genetics using parallel ICA. *Hum Brain Mapp*, *30*(1), 241-255.

1245  https://doi.org/10.1002/hbm.20508

1246 Messé, A., Rudrauf, D., Benali, H., & Marrelec, G. (2014). Relating structure and function in the

1247  human brain: relative contributions of anatomy, stationary dynamics, and non-

1248  stationarities. *PLoS Comput Biol*, *10*(3), e1003530.

1249  https://doi.org/10.1371/journal.pcbi.1003530

1250 Minzenberg, M. J., Laird, A. R., Thelen, S., Carter, C. S., & Glahn, D. C. (2009). Meta-analysis

1251  of 41 functional neuroimaging studies of executive function in schizophrenia. *Archives of*

1252  *general psychiatry*, *66*(8), 811-822.

Mišić, B., Betzel, R. F., de Reus, M. A., van den Heuvel, M. P., Berman, M. G., McIntosh, A. R.,

& Sporns, O. (2016). Network-Level Structure-Function Relationships in Human

Neocortex. *Cereb Cortex*, *26*(7), 3285-3296. https://doi.org/10.1093/cercor/bhw089

Newman, M. E. J. (2006). Modularity and community structure in networks. *Proceedings of the*

National   *Academy*   *of*   *Sciences*,   *103*(23),   8577-8582.

https://doi.org/doi:10.1073/pnas.0601602103

Park, H. J., & Friston, K. (2013). Structural and functional brain networks: from connections to

cognition. *Science*, *342*(6158), 1238411. https://doi.org/10.1126/science.1238411

Potkin, S., Turner, J., Brown, G., McCarthy, G., Greve, D., Glover, G., Manoach, D., Belger, A.,

Diaz, M., & Wible, C. (2009). Working memory and DLPFC inefficiency in schizophrenia:

the FBIRN study. *Schizophrenia bulletin*, *35*(1), 19-31.

Poupon, C., Rieul, B., Kezele, I., Perrin, M., Poupon, F., & Mangin, J. F. (2008). New diffusion

phantoms dedicated to the study and validation of high-angular-resolution diffusion

imaging   (HARDI)   models.   *Magn*   *Reson*   *Med*,   *60*(6),   1276-1283.

https://doi.org/10.1002/mrm.21789

Power, J. D., Cohen, A. L., Nelson, S. M., Wig, G. S., Barnes, K. A., Church, J. A., Vogel, A. C.,

Laumann, T. O., Miezin, F. M., Schlaggar, B. L., & Petersen, S. E. (2011). Functional

network   organization   of   the   human   brain.   *Neuron*,   *72*(4),   665-678.

https://doi.org/10.1016/j.neuron.2011.09.006

Puxeddu, M. G., Faskowitz, J., Betzel, R. F., Petti, M., Astolfi, L., & Sporns, O. (2020). The

modular organization of brain cortical connectivity across the human lifespan.

Neuroimage,   *218*,   116974.

https://doi.org/https://doi.org/10.1016/j.neuroimage.2020.116974

Puxeddu, M. G., Faskowitz, J., Sporns, O., Astolfi, L., & Betzel, R. F. (2022). Multi-modal and multi-subject modular organization of human brain networks. *Neuroimage*, *264*, 119673. https://doi.org/https://doi.org/10.1016/j.neuroimage.2022.119673

Qi, S., Sui, J., Chen, J., Liu, J., Jiang, R., Silva, R., Iraji, A., Damaraju, E., Salman, M., & Lin, D. (2019). Parallel group ICA+ ICA: Joint estimation of linked functional network variability and structural covariation with application to schizophrenia. *Human Brain Mapping*, *40*(13), 3795-3809.

Qi, S., Yang, X., Zhao, L., Calhoun, V. D., Perrone-Bizzozero, N., Liu, S., Jiang, R., Jiang, T., Sui, J., & Ma, X. (2018). MicroRNA132 associated multimodal neuroimaging patterns in unmedicated major depressive disorder. *Brain*, *141*(3), 916-926. https://doi.org/10.1093/brain/awx366

Qi, S. L., Silva, R. F., Zhang, D. Q., Plis, S. M., Miller, R., Vergara, V. M., Jiang, R. T., Zhi, D. M., Sui, J., & Calhoun, V. D. (2022). Three-way parallel group independent component analysis: Fusion of spatial and spatiotemporal magnetic resonance imaging data. *Human Brain Mapping*, *43*(4), 1280-1294. https://doi.org/10.1002/hbm.25720

Qi, S. L., Sui, J., Chen, J. Y., Liu, J. Y., Jiang, R. T., Silva, R., Iraji, A., Damaraju, E., Salman, M., Lin, D. D., Fu, Z. N., Zhi, D. M., Turner, J. A., Bustillo, J., Ford, J. M., Mathalon, D. H., Voyvodic, J., McEwen, S., Preda, A., . . . Calhoun, V. D. (2019). Parallel group ICA plus ICA: Joint estimation of linked functional network variability and structural covariation with application to schizophrenia. *Human Brain Mapping*, *40*(13), 3795-3809. https://doi.org/10.1002/hbm.24632

Reavis, E. A., Lee, J., Altshuler, L. L., Cohen, M. S., Engel, S. A., Glahn, D. C., Jimenez, A. M., Narr, K. L., Nuechterlein, K. H., Riedel, P., Wynn, J. K., & Green, M. F. (2020). Structural

and Functional Connectivity of Visual Cortex in Schizophrenia and Bipolar Disorder: A

Graph-Theoretic   Analysis.   *Schizophrenia   Bulletin   Open*,   *1*(1).

https://doi.org/10.1093/schizbullopen/sgaa056

Rong, B., Huang, H., Gao, G., Sun, L., Zhou, Y., Xiao, L., Wang, H., & Wang, G. (2023).

Widespread Intra- and Inter-Network Dysconnectivity among Large-Scale Resting State

Networks in Schizophrenia. *J Clin Med*, *12*(9). https://doi.org/10.3390/jcm12093176

Seguin, C., Mansour L, S., Sporns, O., Zalesky, A., & Calamante, F. (2022). Network

communication models narrow the gap between the modular organization of structural and

functional   brain   networks.   *Neuroimage*,   *257*,   119323.

https://doi.org/https://doi.org/10.1016/j.neuroimage.2022.119323

Shile, Q., Calhoun, V. D., van Erp, T. G., Damaraju, E., Bustillo, J., Yuhui, D., Turner, J. A.,

Mathalon, D. H., Ford, J. M., Voyvodic, J., Mueller, B. A., Belger, A., McEwen, S., Potkin,

S. G., Preda, A., Birn, F., Tianzi, J., & Jing, S. (2016). Supervised multimodal fusion and

its application in searching joint neuromarkers of working memory deficits in

schizophrenia. *Annu Int Conf IEEE Eng Med Biol Soc*, *2016*, 4021-4026.

https://doi.org/10.1109/embc.2016.7591609

Shine, James M., Bissett, Patrick G., Bell, Peter T., Koyejo, O., Balsters, Joshua H., Gorgolewski,

Krzysztof J., Moodie, Craig A., & Poldrack, Russell A. (2016). The Dynamics of

Functional Brain Networks: Integrated Network States during Cognitive Task

Performance.   *Neuron*,   *92*(2),   544-554.

https://doi.org/https://doi.org/10.1016/j.neuron.2016.09.018

Silva, R. F., Plis, S. M., Adalı, T., & Calhoun, V. D. (2014). A statistically motivated framework for simulation of stochastic data fusion models applied to multimodal neuroimaging. *Neuroimage*, *102*, 92-117.

Skudlarski, P., Jagannathan, K., Anderson, K., Stevens, M. C., Calhoun, V. D., Skudlarska, B. A., & Pearlson, G. (2010). Brain connectivity is not only lower but different in schizophrenia: a combined anatomical and functional approach. *Biol Psychiatry*, *68*(1), 61-69. https://doi.org/10.1016/j.biopsych.2010.03.035

Stam, C. J., van Straaten, E. C. W., Van Dellen, E., Tewarie, P., Gong, G., Hillebrand, A., Meier, J., & Van Mieghem, P. (2016). The relation between structural and functional connectivity patterns in complex brain networks. *International Journal of Psychophysiology*, *103*, 149-160. https://doi.org/https://doi.org/10.1016/j.ijpsycho.2015.02.011

Suárez, L. E., Markello, R. D., Betzel, R. F., & Misic, B. (2020). Linking Structure and Function in Macroscale Brain Networks. *Trends in Cognitive Sciences*, *24*(4), 302-315. https://doi.org/10.1016/j.tics.2020.01.008

Sui, J., Huster, R., Yu, Q., Segall, J. M., & Calhoun, V. D. (2014). Function-structure associations of the brain: evidence from multimodal connectivity and covariance studies. *Neuroimage*, *102 Pt 1*, 11-23. https://doi.org/10.1016/j.neuroimage.2013.09.044

Sui, J., Pearlson, G., Caprihan, A., Adali, T., Kiehl, K. A., Liu, J., Yamamoto, J., & Calhoun, V. D. (2011). Discriminating schizophrenia and bipolar disorder by fusing fMRI and DTI in a multimodal CCA+ joint ICA model. *Neuroimage*, *57*(3), 839-855. https://doi.org/10.1016/j.neuroimage.2011.05.055

Tsai, Y. H., Liang, X., Yang, J. T., & Hsu, L. M. (2019). Modular organization of brain resting

state networks in patients with classical trigeminal neuralgia. *Neuroimage Clin*, *24*,

102027. https://doi.org/10.1016/j.nicl.2019.102027

Van Essen, D. C., Donahue, C. J., Coalson, T. S., Kennedy, H., Hayashi, T., & Glasser, M. F.

(2019). Cerebral cortical folding, parcellation, and connectivity in humans, nonhuman

primates, and mice. *Proceedings of the National Academy of Sciences*, *116*(52), 26173-

26180.

Vergara, V. M., Yu, Q., & Calhoun, V. D. (2018). A method to assess randomness of functional

connectivity matrices. *Journal of Neuroscience Methods*, *303*, 146-158.

https://doi.org/https://doi.org/10.1016/j.jneumeth.2018.03.015

Wang, Z., Xin, J., Wang, Z., Yao, Y., Zhao, Y., & Qian, W. (2021). Brain functional network

modeling and analysis based on fMRI: a systematic review. *Cogn Neurodyn*, *15*(3), 389-

403. https://doi.org/10.1007/s11571-020-09630-5

Wu, L., & Calhoun, V. (2023). Joint connectivity matrix independent component analysis: Auto-

linking of structural and functional connectivities. *Hum Brain Mapp*, *44*(4), 1533-1547.

https://doi.org/10.1002/hbm.26155

Wu, L., Calhoun, V. D., Jung, R. E., & Caprihan, A. (2015). Connectivity-based whole brain dual

parcellation by group ICA reveals tract structures and decreased connectivity in

schizophrenia. *Hum Brain Mapp*, *36*(11), 4681-4701. https://doi.org/10.1002/hbm.22945

Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., Roffman,

J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fischl, B., Liu, H., & Buckner, R. L. (2011).

The organization of the human cerebral cortex estimated by intrinsic functional

connectivity. *J Neurophysiol*, *106*(3), 1125-1165. https://doi.org/10.1152/jn.00338.2011

Zamani Esfahlani, F., Faskowitz, J., Slack, J., Mišić, B., & Betzel, R. F. (2022). Local structure-

function relationships in human brain networks across the lifespan. *Nature*

Communications, *13*(1), 2053. https://doi.org/10.1038/s41467-022-29770-y

Zhao, J., Huang, C.-C., Zhang, Y., Liu, Y., Tsai, S.-J., Lin, C.-P., & Lo, C.-Y. Z. (2023). Structure-

function coupling in white matter uncovers the abnormal brain connectivity in

Schizophrenia. *Translational Psychiatry*, *13*(1), 214. https://doi.org/10.1038/s41398-023-

02520-4

Zhu, Q., Yang, J., Xu, B., Hou, Z., Sun, L., & Zhang, D. (2021). Multimodal Brain Network Jointly

Construction and Fusion for Diagnosis of Epilepsy. *Front Neurosci*, *15*, 734711.

https://doi.org/10.3389/fnins.2021.734711

1374